1   **Transport Pathways and Kinetics of Cerebrospinal Fluid Tracers in Mouse Brain**

2   **Observed by Dynamic Contrast-Enhanced MRI**

Yuran Zhu[1], Guanhua Wang[2], Chaitanya Kolluru[1], Yuning Gu[1], Huiyun Gao[3,4], Jing Zhang[5],

Yunmei Wang[3,4], David L. Wilson[1,6], Xiaofeng Zhu[5], Chris A. Flask[1,6,7], Xin Yu[1,6,8]

[1]Department of Biomedical Engineering, Case Western Reserve University, Cleveland, Ohio, USA

[2]Department of Biomedical Engineering, University of Michigan, Ann Arbor, Michigan, USA

[3]Cardiovascular Research Institute, Case Western Reserve University, Cleveland, Ohio, USA

[4]Department of Medicine, Case Western Reserve University, Cleveland, Ohio, USA

[5]Department of Biostatistics, Population and Quantitative Health Sciences, Case Western

Reserve University, Cleveland, Ohio, USA

[6]Department of Radiology, Case Western Reserve University, Cleveland, Ohio, USA

[7]Department of Pediatrics, Case Western Reserve University, Cleveland, Ohio, USA

[8]Department of Physiology and Biophysics, Case Western Reserve University, Cleveland, Ohio,

USA

Running headline: **Transport of CSF tracers in mouse brain**

18   Address correspondence to:
19   Xin Yu, Sc.D.,
20   Wickenden 430
21   10900 Euclid Avenue
22   Cleveland, OH 44106, USA
23   Tel: 216-368-3918
24   Email: xin.yu@case.edu

Abstract

Recent studies have suggested the glymphatic system as a key mechanism of waste removal in

the brain. Dynamic contrast-enhanced MRI (DCE-MRI) using intracisternally administered

contrast agents is a promising tool for assessing glymphatic function in the whole brain. In this

study, we evaluated the transport kinetics and distribution of three MRI contrast agents with

vastly different molecular sizes in mice. Our results demonstrate that oxygen-17 enriched water

($H_2^{17}O$), which has direct access to parenchymal tissues via aquaporin-4 water channels,

exhibited significantly faster and more extensive transport compared to the two gadolinium-

based contrast agents (Gd-DTPA and GadoSpin). Time-lagged correlation and clustering

analyses also revealed different transport pathways for Gd-DTPA and $H_2^{17}O$. Furthermore, there

were significant differences in transport kinetics of the three contrast agents to the lateral

ventricles, reflecting the differences in forces that drive solute transport in the brain. These

findings suggest the size-dependent transport pathways and kinetics of intracisternal contrast

agents and the potential of DCE-MRI for assessing multiple aspects of solute transport in the

glymphatic system.

Key words

Cerebrospinal fluid, interstitial fluid, glymphatic pathway, CSF transport, subarachnoid space

Introduction

Recent evidence suggests that the exchange of the cerebrospinal fluid (CSF) with the

parenchymal interstitial fluid (ISF) occurs via a highly regulated, brain-wide pathway[1]. The

glymphatic model proposes that CSF in the subarachnoid space is driven by cerebral arterial

pulsation along the perivascular space surrounding penetrating arteries[2], and its influx into the

parenchyma is facilitated by the astroglial water channel aquaporin-4 (AQP4) located on the

vascular endfeet[3]. The bulk flow from the influx of CSF into the parenchymal interstitium

provides an efficient clearance route for metabolic by-products and other toxic wastes from the

brain[4]. The transport of various CSF tracers has been studied extensively in rodent models to

evaluate many pathophysiological factors that may impact CSF transport and CSF-ISF

exchange, including the sleep-wake cycle, anesthesia, body postures, and cardiac function[5–10].

Further, impaired glymphatic function has been indicated in various disease conditions such as

stroke, diabetes, traumatic brain injury, Alzheimer's disease, and other dementias[11–16].

Glymphatic flow and its dependence on AQP4 were first characterized in vivo by two-

photon microscopy using fluorescent tracers with different molecular weights (MWs)[1].

Subsequent studies on four different lines of AQP4-knockout mice further confirmed the critical

role of AQP4 in solute transport in the glymphatic system[3]. Dynamic contrast-enhanced MRI

(DCE-MRI) provides the opportunity to assess both the kinetics and spatial distribution of CSF

tracers in the whole brain[17,18]. Iliff and colleagues were the first to use DCE-MRI to evaluate the

transport of MRI contrast agents in the glymphatic system in rat brains[19]. By comparing the

transport of two gadolinium-based contrast agents (GBCAs) with different molecular sizes (Gd-

DTPA, MW=938 Da; GadoSpin, MW=200 kDa), they showed that GadoSpin transport was

confined to the subarachnoid space and CSF conduits, while Gd-DTPA was able to participate

in CSF-ISF exchange. However, a limitation of using Gd-DTPA as a CSF tracer is that Gd-

DTPA has limited penetration to the parenchyma because of its large molecular size, which may

lead to an underestimation of the CSF-ISF exchange. Indeed, a recent study by Alshuhri et al

compared the transport of Gd-DTPA and oxygen-17 (^{17}O) enriched water ($H_2^{17}O$, MW=19 Da) in

rats and reported significantly faster transport kinetics of $H_2^{17}O$, which is exchanged into the

parenchyma directly via AQP4[20]. These studies suggest that the assessment of glymphatic

function by DCE-MRI is dependent on the molecular size of the contrast agent, and using $H_2^{17}O$

as a CSF tracer provides a unique opportunity to directly evaluate CSF-ISF exchange via

AQP4.

Following these foundational DCE-MRI studies in rats, interest in evaluating the

glymphatic function in mice grows rapidly due to the availability of genetically manipulated

mouse models[3,6,11,21–25]. Assessing mouse glymphatic function by DCE-MRI has been

challenged by the small size of a mouse brain and the limited volume (<20 µL) of fluids that can

be delivered intracisternally without significantly altering the intracranial pressure (ICP)[15,26].

Early studies on mice used the protocol of administering contrast agents on the bench to allow

close monitoring of the infusion process and to visually ensure proper sealing of the infusion

site. While subsequent MRI scanning enabled assessing tracer transport at a later stage, such

an approach inevitably missed the initial phase of contrast agent transport. More recently,

Stanton et al and Gomolka et al successfully performed mouse DCE-MRI studies with in-

scanner delivery of GBCA[9,27]. By monitoring the dynamics of GBCA transport for an hour, their

results show that contrast enhancement peaked within 20 min of infusion in most of the regions

characterized, highlighting the importance of delineating the kinetics of contrast agent transport

at the early phase of infusion in mouse studies.

To build upon these prior studies, the goal of the current study was to evaluate the size-

dependent transport kinetics and distribution of MRI contrast agents in the mouse glymphatic

system. We first established and validated an intracisternal infusion protocol in mice that

allowed the measurements of the entire time course of contrast agent transport for 2 hours.

DCE-MRI studies were performed to compare the transport kinetics of Gd-DTPA, GadoSpin,

and $H_2^{17}O$. Atlas-registered image analysis confirmed significantly faster transport of $H_2^{17}O$

compared to Gd-DTPA, as well as drastically different transport kinetics of the three contrast

agents from cisterna magna to the lateral ventricles. Furthermore, clustering analysis comparing

the kinetic profiles of contrast agent induced signal changes showed a remarkable difference

between $H_2^{17}O$ and Gd-DTPA transport, suggesting different transport pathways for these two

contrast agents.

Results

Transport of contrast agents in the whole brain

Fig. 1 shows representative sagittal slices of signal changes over the 2-hour DCE-MRI

scans for each contrast agent. The signal changes showed clear differences in the transport

dynamics of the three contrast agents (Figs. 1a-c). The time maximum intensity projection maps

(tMIPs) further demonstrated the differences in contrast agent distributions during the entire

time course of DCE-MRI scans (Figs. 1d-f). Following the infusion of the contrast agents at

cisterna magna, contrast enhancement in the cerebellum region proximal to the infusion site can

be observed within 5 min of infusion (CM in Fig. 1e). Subsequently, the transport of all three

contrast agents to the fourth ventricle (V4), as well as along the subarachnoid space on the

ventral surface of the brain, can be appreciated. Gd-DTPA was also transported into the

parenchyma towards the dorsal direction after 15 min (Figs. 1a and 1d). In contrast, the

transport of GadoSpin was primarily confined in the ventricles and the subarachnoid space

(Figs. 1b and 1e). Compared to the two GBCAs, the transport of $H_2^{17}O$ was more extensive and

its distribution at the dorsal side of the brain was more pronounced (Figs. 3c and 3f). Finally, the

downstream flow to the spinal cord (SC) was detectable for all three contrast agents.

Fig. 1 Contrast agent distribution. **a-c:** Representative sagittal views of group-averaged images, overlaid with signal changes from the baseline at selected time points, of mice infused with Gd-DTPA (**a**), GadoSpin (**b**), and $H_2^{17}O$ (**c**), respectively. **d-f:** Time maximum intensity projection maps of representative sagittal and axial views of mice infused with Gd-DTPA (**d**), GadoSpin (**e**), and $H_2^{17}O$ (**f**), respectively. CM, V4, LV, and SC in (**e**) indicate the contrast infusion site at cisterna magna, the fourth ventricle, the lateral ventricle, and the spinal cord, respectively.

Dynamics of contrast agent transport

Shown in Figs. 2 to 4 and Fig. S1 are the time courses of signal changes in selected regions of interest (ROIs) covering the cerebellum and the ventral brain surface (Fig. 2), the deep brain (Fig. 3 and Fig. S1), the dorsal brain and the ventricular regions (Fig. 4). The dynamics of signal changes of the three contrast agents differ drastically in both the magnitude

and the rate of change over time. In cerebellum (Fig. 2b), an ROI proximal to the site of contrast

agent infusion, all three contrast agents showed rapid uptake, with $H_2^{17}O$ being the fastest.

While the accumulation of the two GBCAs remained high, $H_2^{17}O$ exhibited very rapid clearance

in the cerebellum, suggesting fast transport of $H_2^{17}O$ from the infusion site. The transport of

$H_2^{17}O$ was also faster than that of Gd-DTPA in all the ROIs. In the deep brain and dorsal regions

(Figs. 3 and 4), the transport of Gd-DTPA was further delayed compared to that in the ventral

brain regions, with significantly reduced magnitude of signal enhancement that did not reach

steady-state over a time period of ~2 hrs. In contrast, $H_2^{17}O$ transport in these regions was

significantly faster and of a higher magnitude.

Fig. 2 Contrast agent transport in cerebellum and ventral regions. **a:** Segmentation of selected
ROIs. **b-g:** Time courses of signal changes in the selected ROIs. Gray bands indicate the time
period of contrast agent infusion. Red, blue, and green lines represent the mean time courses of
signal changes induced by $H_2^{17}O$, GadoSpin, and Gd-DTPA, respectively. Shaded areas
represent standard errors.

Fig. 3 Contrast agent transport in deep brain regions. **a:** Segmentation of selected ROIs. **b-f:** Time courses of signal changes in the selected ROIs. Gray bands indicate the time period of contrast agent infusion. Red, blue, and green lines represent the mean time courses of signal changes induced by $H_2{}^{17}O$, GadoSpin, and Gd-DTPA, respectively. Shaded areas represent standard errors. CC & EC: Corpus callosum and external capsule.

Compared to Gd-DTPA and $H_2{}^{17}O$, GadoSpin uptake only occurred in the ventricles and a few regions directly adjacent to the subarachnoid or large perivascular spaces, such as brainstem, inferior colliculi, and central gray. In the lateral ventricles (Fig. 4e), GadoSpin showed a prominent rapid uptake that peaked within 15 min after its infusion, followed immediately by a rapid clearance. Interestingly, a delayed but progressive accumulation of $H_2{}^{17}O$ was observed, and the transport of Gd-DTPA to the lateral ventricles was negligible.

Fig. 4 Contrast agent transport in dorsal brain, lateral ventricles and central gray regions. **a:** Segmentation of selected ROIs. **b-f:** Time courses of signal changes in the selected ROIs. Gray bands indicate the time period of contrast agent infusion. Red, blue, and green lines represent the mean time courses of signal changes induced by $H_2^{17}O$, GadoSpin, and Gd-DTPA, respectively. Shaded areas represent standard errors.

Clustering of ROIs

Figs 5 and 6 show results of correlation-matrix-based hierarchical clustering analysis of Gd-DTPA and $H_2^{17}O$ transport, respectively. The maximal cross-correlation coefficient (mCC) and the lag time corresponding to mCC between each pair of ROIs are shown in Figs. 5a and 6a. A large mCC value between two ROIs suggests a high similarity in the dynamic profiles of signal enhancement and the lag time indicates a relative delay in contrast agent transport between the two ROIs. Using 1-mCC as a measure of "dissimilarity" between two ROIs, hierarchical clustering analysis identified four clusters with a dissimilarity value of <0.4 were identified for both Gd-DTPA and $H_2^{17}O$. Bootstrap analysis with 1000 replications returned an approximately unbiased (AU) p-value of 0.92 to 1. These clusters are indicated by the rectangles in the correlation matrices in Figs. 5a and 6a, and their anatomical locations are outlined in Figs. 5b and 6b.

Fig. 5 Correlation-matrix-based clustering analysis of Gd-DTPA transport. **a:** Matrix of maximal cross-correlation (lower left) and lag time (upper right), and cluster dendrogram with bootstrap analysis (top). Values at nodes are AU p-values (red, left) and BP values (green, right), respectively. Clusters with a dissimilarity value of <0.4 are indicated by the rectangles in (**a**). **b:** Outlines of ROIs in each cluster. CC & EC: Corpus callosum and external capsule.

For Gd-DTPA, all five ROIs along the ventral brain surface were classified into the same

cluster (C1, cyan rectangle in Fig. 5a). The mCC values for this cluster were >93% for the four

ROIs in the posterior to midbrain regions and decreased slightly to 89% in the olfactory bulbs in

the anterior region. Further, the lag time was 0 among all these five ROIs, suggesting fast Gd-

DTPA transport along the ventral surface of the brain. The remaining three clusters were

grouped by their proximity to the subarachnoid space or ventral brain. Specifically, C2 (red) was

anatomically adjacent to these regions, while C3 (purple) and C4 (blue) were located distally in

the deep brain and dorsal regions. Correlation analysis of the ROIs in C3 and C4 with those in

the ventral surface (C1) also showed the lowest mCC, as well as a longer lag time ranging from

25 to 55 min (Fig. 5a). In contrast, the lag time between ROIs in C1 and C2 was 0 except for

anterior commissure (5 min).

Fig. 6 Correlation-matrix-based clustering analysis of $H_2^{17}O$ transport. **a:** Matrix of maximal
cross-correlation (lower left) and lag time (upper right), and cluster dendrogram with bootstrap
analysis (top). Values at nodes are AU p-values (red, left) and BP values (green, right),
respectively. Clusters with a dissimilarity value of <0.4 are indicated by the rectangles in (**a**). **b:**
Outlines of ROIs in each cluster.   CC & EC: Corpus callosum and external capsule.

9            $H_2^{17}O$ transport showed distinctly different patterns in correlation matrix, lag time, and

clustering from those of Gd-DTPA (Fig. 6a). The five ROIs along the ventral surface were

classified into three different clusters (C2-C4) instead of one. The clusters were primarily

grouped by their proximity to the infusion site, with two clusters (C1 and C2) located in the

posterior to midbrain regions and the other two clusters (C3 and C4) located in the midbrain to

anterior regions. Lag time between ROIs in the same cluster was 0 except for fimbria, the most

anterior ROI in C2, which showed a lag time of 10-15 min from other ROIs in C2. Further, the

lag time between two ROIs in different clusters was typically <15 min. Only a few ROIs showed

a longer lag time compared to other ROIs, such as the cerebellum, lateral ventricles, amygdala,

olfactory bulbs, corpus callosum and external capsule. The longer lag time between these ROIs

can be attributed to the differences in the kinetic profiles rather than a delay in $H_2^{17}O$ transport,

as indicated by the very low mCC values (⬚40%) between these ROIs.

Impact of $H_2^{17}O$ exchange with blood

To evaluate the impact of the exchange of intracisternal $H_2^{17}O$ with systemic circulation,

blood samples were collected from mice at 20 min (n=3) and 125 min (n=4) after $H_2^{17}O$ infusion,

as well as from mice without $H_2^{17}O$ infusion (n=3). T_2 measurements of the plasma samples did

not show significant differences among these three groups (Fig. S2), suggesting that $H_2^{17}O$

exchange across the blood-brain barrier (BBB) did not significantly alter the T_2 of circulating

blood.

Discussion

This study investigated the size-dependent transport of intracisternally administered CSF

tracers in mouse brain using DCE-MRI. Consistent with previous observations in rats[19], Gd-

DTPA showed relatively fast transport kinetics along the ventral surface of the brain but delayed

and reduced transport into the deep brain and dorsal regions. GadoSpin transport showed

limited penetration to all parenchymal regions of the brain. Compared to the two GBCAs, the

transport of $H_2^{17}O$ was both faster and more extensive, which is also in agreement with a

previous study in rats[20]. These findings were further supported by time-lagged cross-correlation

analysis in which Gd-DTPA showed a much longer lag time between ROIs in the ventral surface

versus those in the deep brain and dorsal regions as compared to $H_2^{17}O$. In addition, clustering

analysis also showed different patterns of transport between Gd-DTPA and $H_2^{17}O$. Further, the

three tracers also showed drastically different transport kinetics from cisterna magna to the

lateral ventricles.

Previous studies of rat glymphatic function have employed pixel-based clustering

analysis to identify regions with similar contrast enhancement features[19,28]. Using the area-

under-the-curve in k-means clustering algorithm, Iliff et al demonstrated deeper penetration of

Gd-DTPA to parenchymal tissues than GadoSpin[19]. In an effort to model the glymphatic system

using local input function, Davoodi-Bojd et al used the derivatives of the time signal curve to

identify clusters with similar profiles of contrast enhancement[28]. In the current study, we

performed ROI-based hierarchical clustering analysis using mCC-derived parameters as a

measure of "dissimilarity" between two ROIs or clusters. Such an approach allowed the

identification of ROIs with similar kinetic profiles over the entire time course of imaging (2

hours). While lag time was not used as a parameter in clustering analysis, most of the ROIs

within a cluster showed minimal lag time for both Gd-DTPA and $H_2^{17}O$, suggesting that this

approach is robust in identifying ROIs along the transport pathway.

The clustering of Gd-DTPA transport was highly consistent with that by Iliff et al using a

pixel-based clustering approach. In particular, all five ROIs along the ventral surface were

identified as a single cluster while the remaining ROIs in the parenchymal tissues were

clustered by their proximity to the subarachnoid space and ventral surface of the brain. More

importantly, ROIs in these clusters also showed a prolonged lag time from the ROIs in the

ventral cluster. These analyses suggest that Gd-DTPA transport occurred initially via the

perivascular network, followed by its penetration into the parenchyma. In contrast, $H_2^{17}O$

transport showed dramatically different clustering patterns compared to Gd-DTPA, with the five

ventral ROIs distributed in three different clusters. In general, clustering of $H_2^{17}O$ transport

followed a pattern of posterior to anterior while Gd-DTPA was along the ventral surface towards

the dorsal direction. These different clustering patterns reflect different transport pathways for

$H_2^{17}O$ because of its direct access to parenchymal tissues via AQP4.

A potential confounding factor in $H_2^{17}O$ transport is the exchange of water across the

BBB. Since systemic circulation is much faster than CSF flow[29,30], its participation in $H_2^{17}O$

transport may significantly accelerate the distribution of $H_2^{17}O$ to the whole brain. To evaluate

the impact of $H_2^{17}O$ exchange across BBB, we measured the T_2 of plasma collected at two time

points, one at the time when $H_2^{17}O$-induced T_2 change reached peak in parenchyma (20 min)

and one at the end of the scan protocol (125 min). Our measurements showed no significant

difference in plasma T_2 at both time points compared to samples collected from mice without

$H_2^{17}O$ infusion. This lack of change in plasma T_2 suggests that the intracisternal administration

of a small volume of $H_2^{17}O$ (10 µL) had a negligible effect on plasma $H_2^{17}O$ concentration.

Hence, it is unlikely that systemic circulation played a significant role in the observed fast $H_2^{17}O$

transport in the current study. The participation of cerebrovasculature in $H_2^{17}O$ transport needs

more careful evaluation, as the possibility of $H_2^{17}O$ entering the bloodstream near the infusion

site and impacting signals in the cerebellum cannot be ruled out completely. However,

considering that arteries in the cerebellum are not the feeding arteries to other regions of the

brain and the very short transit time of cerebral blood in a mouse brain[31], CSF-blood exchange

would have to be extremely fast for cerebrovasculature to play a significant role in $H_2^{17}O$

transport beyond cerebellum. Furthermore, Alshuhri et al reported that rats pretreated with an

AQP4 inhibitor exhibited ~80% reduction in $H_2^{17}O$ transport into the parenchyma, providing

strong evidence that AQP4 and the glymphatic system play a major role in $H_2^{17}O$ transport to

the parenchymal tissue[20].

A recent study using confocal microscopy and fluorescein-labeled dextran (MW=500

kDa) as a CSF tracer has suggested that tracers administered via cisterna magna can be

transported to the lateral ventricles via the interpeduncular cistern[32]. However, such observation

was not unequivocally supported by a DCE-MRI study using Gd-DOTA (MW=661.8 Da) as the

tracer, in which only one rat with an abnormally enlarged third ventricle exhibited Gd-DOTA

uptake in the lateral ventricles[33]. In the current study, mice also showed negligible uptake of Gd-

DTPA into the lateral ventricles. However, there was a rapid uptake of GadoSpin into the lateral

ventricles that peaked at the end of its infusion, and it was immediately followed by a rapid

clearance. Interestingly, despite its fast transport to the parenchymal tissues, $H_2^{17}O$ showed a

slow but progressive accumulation of $H_2^{17}O$ in the lateral ventricles (Fig. 4e). Considering that

cisterna magna is at the downstream of CSF circulation from the fourth ventricle[34], direct

transport of CSF tracers from cisterna magna to the lateral ventricles would require a reversal of

the pressure gradient. The transport of GadoSpin into the lateral ventricles suggests that

intracisternal infusion may have caused such a transient reversal of the pressure gradient

between the cisterna magna and the lateral ventricles, and the rapid clearance of GadoSpin

appears to be consistent with the restoration of the normal pressure gradient immediately after

the infusion. On the other hand, the delayed transport of $H_2^{17}O$ and the negligible transport of

Gd-DTPA into the lateral ventricles suggest that the pressure gradient and the transport

impedance for these two contrast agents may have favored their transport from the

subarachnoid space into the parenchymal tissues. As such, the delayed $H_2^{17}O$ uptake into the

lateral ventricles may have occurred at the parenchymal-ventricular interface following its

transport to the parenchyma. However, due to the slow and reduced penetration of Gd-DTPA

into the parenchyma, Gd-DTPA did not reach the parenchymal-ventricular interface in a

significant amount, leading to its negligible uptake into the lateral ventricles. Further

investigations rooted in the fluid dynamics of CSF circulation are warranted to gain more

quantitative insight of these observations.

There are several limitations in the current study. First, the use of T_1- or T_2-weighted

images in DCE-MRI is a semi-quantitative approach such that signal changes from the baseline

can only serve as a proxy of contrast agent concentration. While the concentrations of

gadolinium ions (Gd^{2+}) were matched for the two GBCAs, a direct comparison of contrast agent

concentrations is not possible because of the nonlinear relationship between signal changes

and contrast agent concentrations. As such, the current study focused on evaluating the kinetics

of contrast agent induced signal changes. To account for inter-subject variations, we normalized

the signal changes by using the maximal signal surrounding the infusion site as the common

denominator. While this approach has enabled the comparison of relative signal changes in

different ROIs, flux quantification is challenged by the nonlinear relationship between signal

change and contrast concentration. Second, due to hardware limitations, a lower spatial

resolution was used in the current study to achieve adequate temporal resolution, which has led

to the pronounced partial volume effect in a few ROIs. For example, the large signal increase in

the central gray from mice infused with GadoSpin (Fig. 4e) is likely caused by GadoSpin uptake

in the fourth ventricle. Improving the spatial resolution will reduce the partial volume effect and

enable the segmentation of smaller ROIs in future studies. Finally, previous studies have shown

that general anesthesia with isoflurane alone can impair the circulation of CSF through the brain

while low-dose isoflurane with supplementary dexmedetomidine or ketamine/xylazine can

enhance glymphatic transport[5,6,9]. Hence, an anesthetic regime that enhances glymphatic

function can be more desirable in studying solute transport in the brain, especially when using

tracers with a large molecular size such as Gd-DTPA.

In summary, the current study established an experimental protocol that enabled DCE-

MRI measurements of the entire time course of contrast agent transport in the mouse

glymphatic system. Comparison of the transport kinetics and distribution of three MRI contrast

agents with different molecular sizes showed drastically different transport profiles and

clustering patterns, suggesting that the transport pathways and kinetics in the glymphatic

system are size-dependent.

Materials and Methods

Animals

The animal protocol was approved by the Institutional Animal Care and Use Committee

of Case Western Reserve University. The experiments were performed on 13- to 15-week-old

male C57BL/6J mice (Jackson Laboratories, Bar Harbor, ME, US). The average body weight at

the time of MRI scan was 29.4 g. The animals were housed in a temperature- and humidity-

controlled environment with ad libitum access to food and water and a 12-hour light-dark cycle.

Cisterna magna cannulation

The animal was anesthetized with 3% isoflurane mixed with 100% O_2 in an induction

chamber and transferred to a stereotaxic frame with the head secured by ear bars and a tooth

bar. Anesthesia was maintained with 1.5% isoflurane in 100% O_2 delivered via a nose cone.

The body temperature was maintained at ~37°C with a heating tape attached to the surface of

the stereotaxis frame. A midline dorsal neck incision was made to expose the dura mater. A

small durotomy was made using a 30-gauge needle to expose the cisterna magna. A

polyethylene micro-catheter (0.13 mm ID × 0.25 mm OD, Scientific Commodities, Lake Havasu

City, AZ, US) was inserted into the intrathecal space and secured with cyanoacrylate glue.

Tracer delivery via cisterna magna was first confirmed with bright-field microscopy of Evans

blue and cryoimaging of CF594 hydrazide and FITC-dextran (Fig. S3).

MRI protocol

MRI studies were performed on a horizontal bore 9.4T preclinical scanner (Bruker

Biospin, Billerica, MA, US) using a 35-mm volume coil. Mice were randomly assigned to three

groups with intracisternal infusion of 1) 12.5 mM Gd-DTPA (n=7; MW=661.8 Da; Mallinckrodt, St

Louis, MO, US); 2) 4.17 mM GadoSpin-P (n=6; MW=200 kDa; Miltenyi Biotec, Bergisch

Gladbach, Germany); and 3) 90% enriched $H_2^{17}O$ (n=7; MW=19 Da; NUKEM Isotopes, Alzenau,

Germany). After the cannulation, the mouse was transferred to an MRI-compatible cradle and

placed in a prone position. Anesthesia was maintained with 1 to 1.5% isoflurane delivered via a

nose cone. Respiration rate and body temperature were monitored during the MRI scan. The

respiration rate was maintained within 90 to 110 breaths per minute (bpm) by adjusting the

anesthesia level. The body temperature was maintained at ~37°C by blowing warm air into the

scanner through a feedback control system (SA Instruments, Stony Brook, NY, US). After the

initial setup, an anatomic scan was acquired at baseline using a 3D spin-echo sequence with

the following parameters: TR/TE, 50/8.1 ms; FOV, 20×16×14 mm^3; matrix size, 100x80x70;

NAV, 2. Total scan time was ~9 min. Subsequently, DCE-MRI acquisition was started, and 10

μL of contrast agent was infused at a rate of 1 μL/min (10 min total infusion time).

The transport of Gd-DTPA and GadoSpin was tracked dynamically using a T_1-weighted

3D FLASH sequence with the following parameters: TR/TE, 50/2.5 ms; flip angle, 15°; FOV,

20×16×14 mm^3; matrix size, 100x80x70; NAV, 1, leading to an isotropic resolution of 200 μm

and a temporal resolution of 5 min. A single baseline scan and 25 dynamic scans were acquired

before, during, and after contrast agent infusion for 130 min.

$H_2^{17}O$ transport was imaged with a T_2-weighted multi-slice RARE sequence. A total of 28

slices were acquired with TR, 2500 ms; FOV, 20x16 mm^2; matrix size, 100x80; slice thickness,

0.5 mm. 8 echoes were acquired in each TR, leading to an effective TE of 31 ms. Acquisition

time for a single-average dataset was 25 s. A total of 312 images were acquired continuously

for 130 min with the first 12 images acquired as baseline.

To evaluate the significance of $H_2^{17}O$ exchange across the blood-brain barrier (BBB),

blood was collected from a subset of mice infused with $H_2^{17}O$ at the end of DCE-MRI scan

(n=4). Additional experiments were performed on the bench to collect blood from mice 10 min

after the completion of $H_2^{17}O$ infusion (i.e., 20 min from the start of $H_2^{17}O$ infusion, n=3), as well

as from mice without $H_2^{17}O$ infusion (n=3). The blood samples were centrifuged at 2400 RPM

for 20 min and the plasma was extracted. The T_2 values of the plasma samples were measured

using a single-slice Car-Purcell-Meiboom-Gill (CPMG) sequence with the following acquisition

parameters: TR, 10 s; FOV, 20x20 mm^2; matrix size, 128x128; slice thickness, 1 mm. A total of

64 echoes were acquired with an evenly spaced echo time of 8 ms.

MRI image analysis

All images and data analyses were performed using either in-house developed or open-

source software in MATLAB (MathWorks, Natick, MA, US) or Python (Python Software

Foundation, v.3.0). Single-average T_1-weighted images were reconstructed to delineate the

kinetics and distribution of Gd-DTPA and GadoSpin transport. For the analysis of $H_2^{17}O$

transport, T_2-weighted images were reconstructed using 6 averages, resulting in a temporal

resolution of 2.5 min. Motion correction was performed by registering the dynamic images to the

anatomic image acquired at baseline via affine transformation using the open-source toolkit,

Advanced Normalization Tools[35–37].

Following motion correction, image segmentation was performed by co-registering

images to an MRI mouse brain atlas[38]. Specifically, a representative animal was selected from

each contrast agent group, and the anatomic images of the representative animal were

registered to the atlas through affine and deformable transformation. Subsequently, images

from the remaining animals in each contrast agent group were registered to that of the

representative animal, followed by the generation of averaged dynamic images and tMIPs.

A total of 20 ROIs were generated from the co-registered brain atlas, covering the

intracisternal infusion and transport pathways, the brain parenchyma, and the lateral ventricles.

Mean signal intensity in each ROI was calculated, followed by the subtraction of the baseline

signal for the entire dynamic series. Subsequently, the maximal signal from a small region

surrounding the infusion catheter in the cisterna magna was used as the "input function" to

normalize the time course of signal changes in each ROI.

Clustering analysis

A correlation-matrix-based hierarchical clustering method was used to analyze the time

courses of Gd-DTPA and $H_2^{17}O$ transport among different ROIs[39]. Specifically, time-lagged

cross-correlation analysis was performed to determine the similarity of the kinetics of contrast

agent transport between two ROIs, as well as the lag time corresponding to the maximal cross-

correlation coefficient (mCC)[40]. Subsequently, the "dissimilarity", defined as 1-mCC, was used

to quantify the distance between two ROIs. A dendrogram representing the hierarchical

structure of the mCC matrix with complete linkage was generated using the open-source library

SciPy[41]. Bootstrap analysis using the Pvclust package was performed to assess the uncertainty

in clustering analysis[42]. A bootstrap replication of 1000 was used to calculate the approximately

unbiased (AU) p-values and the bootstrap probability (BP) values. Visualizations of ROIs in

each cluster were created using the Allen mouse brain atlas and brainrender[43,44].

References

1. Iliff, J. J. *et al.* A Paravascular Pathway Facilitates CSF Flow Through the Brain Parenchyma and the Clearance of Interstitial Solutes, Including Amyloid β. *Sci. Transl. Med.* **4**, (2012).

2. Iliff, J. J. *et al.* Cerebral Arterial Pulsation Drives Paravascular CSF-Interstitial Fluid Exchange in the Murine Brain. *J. Neurosci.* **33**, 18190–18199 (2013).

3. Mestre, H. *et al.* Aquaporin-4-dependent glymphatic solute transport in the rodent brain. *eLife* **7**, e40070 (2018).

4. Nedergaard, M. Garbage Truck of the Brain. *Science* **340**, 1529–1530 (2013).

5. Benveniste, H. *et al.* Anesthesia with Dexmedetomidine and Low-dose Isoflurane Increases Solute Transport *via* the Glymphatic Pathway in Rat Brain When Compared with High-dose Isoflurane. *Anesthesiology* **127**, 976–988 (2017).

6. Gakuba, C. *et al.* General Anesthesia Inhibits the Activity of the "Glymphatic System". *Theranostics* **8**, 710–722 (2018).

7. Hablitz, L. M. *et al.* Increased glymphatic influx is correlated with high EEG delta power and low heart rate in mice under anesthesia. *Sci. Adv.* **5**, eaav5447 (2019).

8. Lee, H. *et al.* The Effect of Body Posture on Brain Glymphatic Transport. *J. Neurosci.* **35**, 11034–11044 (2015).

9. Stanton, E. H. *et al.* Mapping of CSF transport using high spatiotemporal resolution dynamic contrast-enhanced MRI in mice: Effect of anesthesia. *Magn. Reson. Med.* **85**, 3326–3342 (2021).

10. Xie, L. *et al.* Sleep Drives Metabolite Clearance from the Adult Brain. *Science* **342**, 373–377 (2013).

11. Gaberel, T. *et al.* Impaired Glymphatic Perfusion After Strokes Revealed by Contrast-Enhanced MRI: A New Target for Fibrinolysis? *Stroke* **45**, 3092–3096 (2014).

12. Goulay, R. *et al.* Subarachnoid Hemorrhage Severely Impairs Brain Parenchymal

Cerebrospinal Fluid Circulation in Nonhuman Primate. *Stroke* **48**, 2301–2305 (2017).

13. Iliff, J. J. *et al.* Impairment of Glymphatic Pathway Function Promotes Tau Pathology after

Traumatic Brain Injury. *J. Neurosci.* **34**, 16180–16193 (2014).

14. Jiang, Q. *et al.* Impairment of the glymphatic system after diabetes. *J. Cereb. Blood Flow*

Metab. **37**, 1326–1337 (2017).

15. Kress, B. T. *et al.* Impairment of paravascular clearance pathways in the aging brain:

Paravascular Clearance. *Ann. Neurol.* **76**, 845–861 (2014).

16. Rasmussen, M. K., Mestre, H. & Nedergaard, M. The glymphatic pathway in neurological

disorders. *Lancet Neurol.* **17**, 1016–1024 (2018).

17. Jiang, Q. MRI and glymphatic system. *Stroke Vasc. Neurol.* **4**, 75–77 (2019).

18. Taoka, T. & Naganawa, S. Glymphatic imaging using MRI. *J. Magn. Reson. Imaging* **51**,

11–24 (2020).

19. Iliff, J. J. *et al.* Brain-wide pathway for waste clearance captured by contrast-enhanced MRI.

J. Clin. Invest. **123**, 1299–1309 (2013).

20. Alshuhri, M. S., Gallagher, L., Work, L. M. & Holmes, W. M. Direct imaging of glymphatic

transport using H217O MRI. *JCI Insight* **6**, e141159 (2021).

21. Ma, Q. *et al.* Rapid lymphatic efflux limits cerebrospinal fluid flow to the brain. *Acta*

Neuropathol. (Berl.) **137**, 151–165 (2019).

22. Xue, Y. *et al.* Sustained glymphatic transport and impaired drainage to the nasal cavity

observed in multiciliated cell ciliopathies with hydrocephalus. *Fluids Barriers CNS* **19**, 20

(2022).

23. Xue, Y. *et al.* In vivo T1 mapping for quantifying glymphatic system transport and cervical

lymph node drainage. *Sci. Rep.* **10**, 14592 (2020).

24. Zamani, A. *et al.* Impaired glymphatic function in the early stages of disease in a TDP-43

mouse model of amyotrophic lateral sclerosis. *Transl. Neurodegener.* **11**, 17 (2022).

25. Zhang, J. *et al.* Impaired Glymphatic Transport Kinetics Following Induced Acute Ischemic Brain Edema in a Mouse pMCAO Model. *Front. Neurol.* **13**, 860255 (2022).

26. Benveniste, H. *et al.* Glymphatic Cerebrospinal Fluid and Solute Transport Quantified by MRI and PET Imaging. *Neuroscience* **474**, 63–79 (2021).

27. Gomolka, R. S. *et al.* Loss of aquaporin-4 results in glymphatic system dysfunction via brain-wide interstitial fluid stagnation. *eLife* **12**, e82232 (2023).

28. Davoodi-Bojd, E. *et al.* Modeling glymphatic system of the brain using MRI. *NeuroImage* **188**, 616–627 (2019).

29. Li, J. *et al.* Whole-brain mapping of mouse CSF flow via HEAP-METRIC phase-contrast MRI. *Magn. Reson. Med.* **87**, 2851–2861 (2022).

30. Pardridge, W. M. Drug transport in brain via the cerebrospinal fluid. *Fluids Barriers CNS* **8**, 7 (2011).

31. Hirschler, L. *et al.* Transit time mapping in the mouse brain using time-encoded pCASL. *NMR Biomed.* **31**, e3855 (2018).

32. Bedussi, B. *et al.* Paravascular channels, cisterns, and the subarachnoid space in the rat brain: A single compartment with preferential pathways. *J. Cereb. Blood Flow Metab.* **37**, 1374–1385 (2017).

33. Lee, H. *et al.* Quantitative Gd-DOTA uptake from cerebrospinal fluid into rat brain using 3D VFA-SPGR at 9.4T: Quantitative Gd-DOTA Uptake from CSF into Rat Brain. *Magn. Reson. Med.* **79**, 1568–1578 (2018).

34. Simon, M. J. & Iliff, J. J. Regulation of cerebrospinal fluid (CSF) flow in neurodegenerative, neurovascular and neuroinflammatory disease. *Biochim. Biophys. Acta BBA - Mol. Basis Dis.* **1862**, 442–451 (2016).

35. Avants, B. B. *et al.* A reproducible evaluation of ANTs similarity metric performance in brain image registration. *NeuroImage* **54**, 2033–2044 (2011).

36. Hübner, N. S. *et al.* The connectomics of brain demyelination: Functional and structural

patterns in the cuprizone mouse model. *NeuroImage* **146**, 1–18 (2017).

37. Koch, S. *et al.* Atlas registration for edema-corrected MRI lesion volume in mouse stroke

models. *J. Cereb. Blood Flow Metab.* **39**, 313–323 (2019).

38. Ma, Y. *et al.* A three-dimensional digital atlas database of the adult C57BL/6J mouse brain

by magnetic resonance microscopy. *Neuroscience* **135**, 1203–1215 (2005).

39. Liu, X., Zhu, X.-H., Qiu, P. & Chen, W. A correlation-matrix-based hierarchical clustering

method for functional connectivity analysis. *J. Neurosci. Methods* **211**, 94–102 (2012).

40. Siegel, J. S., Snyder, A. Z., Ramsey, L., Shulman, G. L. & Corbetta, M. The effects of

hemodynamic lag on functional connectivity and behavior after stroke. *J. Cereb. Blood Flow*

Metab. **36**, 2162–2176 (2016).

41. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat.*

Methods **17**, 261–272 (2020).

42. Suzuki, R. & Shimodaira, H. Pvclust: an R package for assessing the uncertainty in

hierarchical clustering. *Bioinformatics* **22**, 1540–1542 (2006).

43. Lein, E. S. *et al.* Genome-wide atlas of gene expression in the adult mouse brain. *Nature*

445, 168–176 (2007).

44. Claudi, F. *et al.* Visualizing anatomically registered data with brainrender. *eLife* **10**, e65751

(2021).

Declarations

Acknowledgments

The author(s) would like to thank Drs. Junqing Zhu and Chunying Wu for their
assistance in establishing the intracisternal infusion protocol in mice.

Authors' contributions

YZ, experimental design, technical development and implementation, data collection and
analysis, manuscript development and editing; GW, data analysis; CK, data collection and
analysis; YG, technical development and implementation; HG, technical development; JZ, data
analysis; XZ, data analysis; YW, technical development; DW, technical development; CF,
experimental design, technical development; XY, experimental design, technical development,
data collection and analysis, manuscript development and editing.

Availability of data

Experimental data, images, and code from this study are available upon request to the
corresponding author.

Declaration of conflicting interests

There were no potential conflicts of interest with respect to the research, authorship,
and/or publication of this article.

Funding

This work was supported by grants from the National Institute of Health awards R01
NS124206 and R01 EB023704 to X. Y., and predoctoral fellowship award from American Heart
Association 23PRE1017924 to Y. Z.

Figure legends

Fig. 1 Contrast agent distribution. **a-c:** Representative sagittal views of group-averaged images,

overlaid with signal changes from the baseline at selected time points, of mice infused with Gd-

DTPA (**a**), GadoSpin (**b**), and $H_2^{17}O$ (**c**), respectively. **d-f:** Time maximum intensity projection

maps of representative sagittal and axial views of mice infused with Gd-DTPA (**d**), GadoSpin

(**e**), and $H_2^{17}O$ (**f**), respectively. CM, V4, LV, and SC in (**e**) indicate the contrast infusion site at

cisterna magna, the fourth ventricle, the lateral ventricle, and the spinal cord, respectively.

Fig. 2 Contrast agent transport in cerebellum and ventral regions. **a:** Segmentation of selected

ROIs. **b-g:** Time courses of signal changes in the selected ROIs. Gray bands indicate the time

period of contrast agent infusion. Red, blue, and green lines represent the mean time courses of

signal changes induced by $H_2^{17}O$, GadoSpin, and Gd-DTPA, respectively. Shaded areas

represent standard errors.

Fig. 3 Contrast agent transport in deep brain regions. **a:** Segmentation of selected ROIs. **b-f:**

Time courses of signal changes in the selected ROIs. Gray bands indicate the time period of

contrast agent infusion. Red, blue, and green lines represent the mean time courses of signal

changes induced by $H_2^{17}O$, GadoSpin, and Gd-DTPA, respectively. Shaded areas represent

standard errors. CC & EC: Corpus callosum and external capsule.

Fig. 4 Contrast agent transport in dorsal brain, lateral ventricles and central gray regions. **a:**

Segmentation of selected ROIs. **b-f:** Time courses of signal changes in the selected ROIs. Gray

bands indicate the time period of contrast agent infusion. Red, blue, and green lines represent

the mean time courses of signal changes induced by $H_2^{17}O$, GadoSpin, and Gd-DTPA,

respectively. Shaded areas represent standard errors.

Fig. 5 Correlation-matrix-based clustering analysis of Gd-DTPA transport. **a:** Matrix of maximal

cross-correlation (lower left) and lag time (upper right), and cluster dendrogram with bootstrap

analysis (top). Values at nodes are AU p-values (red, left) and BP values (green, right),

respectively. Clusters with a dissimilarity value of <0.4 are indicated by the rectangles in (**a**). **b:**

Outlines of ROIs in each cluster. CC & EC: Corpus callosum and external capsule.

Fig. 6 Correlation-matrix-based clustering analysis of $H_2^{17}O$ transport. **a:** Matrix of maximal

cross-correlation (lower left) and lag time (upper right), and cluster dendrogram with bootstrap

analysis (top). Values at nodes are AU p-values (red, left) and BP values (green, right),

respectively. Clusters with a dissimilarity value of <0.4 are indicated by the rectangles in (**a**). **b:**

Outlines of ROIs in each cluster.   CC & EC: Corpus callosum and external capsule.