The role of recent evolutionary history in resilience to environmental

change: social evolution effects versus founder effects

Abstract

Principles of social evolution have long been used retrospectively to interpret social

interactions, but have less commonly been applied predictively to inform conservation and

animal husbandry strategies. We investigate whether past social selection can predict resilience

to environmental change. Upon exposure to harsh novel environments, populations that

previously experienced more benign social environments are predicted either to suffer fitness

losses (the "mutation load hypothesis" and "selection filter hypothesis") or maintain fitness

(the "beneficial mutation hypothesis"). We tested these contrasting predictions using

populations of burying beetles Nicrophorus vespilloides evolved experimentally for 48

generations under harsh versus benign social environments through manipulation of parental

care. We exposed sexually immature adults from each population to varying heat stress and

measured the effect on survival and reproduction. The more benign the social environment

previously experienced by a population, the better its survival under heat stress during sexual

maturation. Although consistent with the "beneficial mutation hypothesis", it is also possible

that populations evolved under the harsher social environment were more prone to dying during

maturation, regardless of their thermal environment. Overall, stochastic genetic variation,

probably due to founder effects, was a better predictor of resilience. We discuss the

implications for translocation and captive breeding programmes.

Introduction

In a world changing rapidly through the effects of climate change (Pörtner et al., 2022), habitat degradation (Millenium Ecosystem Assessment, 2005) and excessive use of antibiotics (Graham et al., 2019) and pesticides (Hawkins et al., 2019), our ability to predict how effectively organisms will adapt to new environments is more important than ever. The principles of social evolution have long been used retrospectively to understand social interactions, however they have been more rarely used predictively to inform the development of practical strategies to address global problems (Carroll et al., 2014). Yet an understanding of a population's evolutionary history, by which we mean the type of selective environment it has experienced, could predict its resilience to future environmental perturbation (Lumley et al., 2015). This has practical value in its potential to address several major economic and societal problems, including food security (pest/crop interactions, crop development), human and animal medicine (antibiotic development, anti-parasite interventions) and wild animal conservation (choice of populations for translocation schemes, identification of populations vulnerable to disease and environmental change) (Carroll et al., 2014).

Few studies have tested experimentally how past exposure to different social selective environments affects a population's future response to environmental change (but see Lumley et al., 2015). In theory, the effect on resilience depends on whether the social environment experienced by the population is benign or harsh - because this influences the extent of standing genetic variation available.

Benign social interactions, which are usually due to some form of cooperative interaction, are known to buffer individuals from environmental stress and therefore to reduce the extent to

which mildly deleterious alleles are exposed to natural selection (Linksvayer & Wade, 2009; Pascoal et al., 2022). Elevated mutation loads resulting from relaxed selection have been shown to reduce reproductive success in animals kept in captive breeding programmes (Araki et al., 2007; McPhee, 2004; Robert, 2009) and have been linked to a higher prevalence of genetic disorders in human societies where selection is buffered by widely available healthcare (Lynch, 2016; Saniotis et al., 2021; You & Henneberg, 2018). The "mutation load hypothesis" therefore predicts that individuals previously exposed to more benign social environments should be less resilient to new environmental stresses. However, whilst most of the mutations that accumulate under relaxed selection are likely to be (mildly) harmful, some could potentially be beneficial and might even facilitate greater resilience to environmental change at the population level. The "beneficial mutation hypothesis" therefore conversely predicts that individuals previously exposed to more benign social environments could be more resilient to new environmental stresses, if by chance they have accumulated cryptic genetic variation that is beneficial when expressed in these new environments (Barrett & Schluter, 2008; Paaby & Rockman, 2014). At the other end of the spectrum, a harsh social environment potentially imposes a strong selection filter, which is likely to reduce the population's mutation load, causing individuals generally to be fitter (Balmford, 1996). The "selection filter hypothesis" predicts that a history of exposure to harsher social environments means that populations are more likely to be resilient to environmental change.

Species in which offspring are facultatively dependent on parental care (Clutton-Brock, 1991) offer a way to test these contrasting predictions. Offspring that receive care are exposed to a benign social environment during development which, if sustained over many generations relaxes selection (Snell-Rood et al., 2016). Conversely, offspring that receive no care over many generations experience a harsher environment during development and consequently

experience stronger selection against deleterious mutations. We exploited this natural history

to test whether, and how, past social interactions influence future resilience to environmental

change. Our experiments focus on the burying beetle *Nicrophorus vespilloides*, which exhibits

elaborate biparental care upon which offspring are facultatively dependent (Capodeanu-Nägler

et al., 2016).

Burying beetle adults locate the carcasses of small vertebrates using their highly sensitive

antennae (Kalinová et al., 2009; Trumbo & Steiger, 2020) and then fight intrasexually for

ownership (Otronen, 1988). The winning pair then tears the fur or feathers from the carcass,

covers the flesh with antimicrobial oral and anal exudates, rolls it into a ball and buries it

underground (Cotter & Kilner, 2010). The female then lays eggs in the soil surrounding the

carcass nest. When the larvae hatch, 2-3 days later, they crawl to the nest (Müller & Eggert,

1989), where their parents defend them from predators and intra- and interspecific competitors

and feed them trophallactically (Eggert et al., 1998). The duration of post-hatching care is

highly variable and in some cases is non-existent due to the death or desertion by parents after

preparation of the carcass nest (Schrader, Jarrett, et al., 2015a; Smiseth et al., 2003).

We evolved two replicate populations of burying beetles in the laboratory under contrasting

regimes of parental care for 48 generations. In the Full Care populations, beetles were allowed

to stay with their offspring throughout development. Individuals in these populations

experienced a benign environment during development and relaxed selection (Pascoal et al.,

2022). By contrast, in the No Care populations, parents were removed just prior to larval

hatching and so were unable to supply any post-hatching care at all. Therefore individuals in

these populations experienced a harsher social environment and stronger selection against

deleterious mutations (Pascoal et al., 2022).

After 47 generations of experimental evolution, we tested whether there were differences in

the resilience of beetles from these two types of experimental population when they were

exposed to higher temperatures during their juvenile adult stage. We define resilience as the

ability of a population to maintain normal processes, in particular reproductive success, in

response to a harsh novel environment (Lowerre-Barbieri et al., 2017; Martin & Wiebe, 2004;

Wingfield et al., 2011). Temperature was chosen as the environmental stressor because of its

ecological relevance in the context of climate change. Insects, being poikilothermic, are

particularly vulnerable to heat stress, and research on burying beetles specifically has shown

that they have reduced breeding success at higher temperatures (Moss & Moore, 2021; Quinby

et al., 2020; Sun & Kilner, 2020). We chose to expose the beetles to the temperature treatments

during sexual maturation alone because this is the time when an adult beetle's gonads are

developing (e.g. David et al., 2005; Gibbs et al., 2010; Sales et al., 2018; Zizzari and Ellers,

2011). Furthermore, Sales et al. (2021) show that this is the developmental stage that is most

vulnerable to effects of thermal stress on fertility – perhaps because in nature, sexually

immature individuals cannot stay underground to buffer themselves from heat stress because

they must emerge to feed.

We exposed sexually immature adults to different degrees of heat stress and measured the

effect on the following correlates of fitness: survival to sexual maturity, anal exudate lytic

activity, carcass sphericity, fecundity, and hatching success (Cotter and Kilner, 2010, Arce et

al., 2012; Cotter and Kilner, 2010, Duarte et al., 2021). We predicted that the contrasting

regimes of selection imposed by these different social environments for the 47 preceding

generations should yield a contrasting response to heat stress during sexual maturation. The

"mutation load hypothesis" and "selection filter hypothesis" each predict that individuals

derived from the No Care populations should be more resilient to heat stress, whereas the

"beneficial mutation hypothesis" predicts that individuals derived from the No Care

populations should be less resilient to heat stress.

Methods

Burying beetle husbandry in the laboratory: general methods

Burying beetles were bred by placing them in a plastic breeding box (17 x 12 x 6 cm), partially

filled with damp soil (Miracle-Gro Compost), furnished with a 10-15 g mouse carcass for

breeding. Boxes were placed in a cupboard after pairing to simulate underground conditions.

Larvae were counted and weighed 8 days after pairing and each one placed in its own cell (2 x

2 x 2 cm) within a plastic pupation box (10 x 10 x 2 cm), filled with damp peat. Sexually

immature adults were eclosed around 21 days later and each was placed in its own individual

box (12 x 8 x 2 cm). Adults were fed twice weekly with beef mince until breeding, which took

place 15 days post-eclosion. At all life stages, individuals were kept at 21°C. During breeding,

pairs were kept in a constantly dark environment, but for all other life stages they were kept on

a 16L: 8D hour light cycle.

Experimental Evolution

The *N. vespilloides* populations used in this study were part of a long-term experimental

evolution project that investigated how populations of burying beetles adapt to the loss of

parental care. There were four experimental populations in all: Full Care (FC) (replicated

twice) and No Care (NC) (replicated twice). Their establishment and husbandry have been

described in detail before (Duarte et al., 2021; Jarrett, Evans, et al., 2018; Jarrett, Rebar, et al.,

2018; Rebar et al., 2020; Schrader et al., 2017; Schrader, Jarrett, et al., 2015a, 2015b). Briefly,

these populations were established in 2014 with wild-caught beetles from four woodland sites

across Cambridgeshire, UK (Byron's Pool, Gamlingay Woods, Waresley Woods and Overhall

Grove). FC populations were left undisturbed (as described in general methods above) and

allowed to stay with their offspring until dispersal. However, in the NC population, at 53 h

post-pairing, when the carrion nest was complete but before the larvae had hatched, both

parents were removed to prevent post-hatching care. This procedure was repeated at every

generation. The experimental populations were bred in two separate blocks (FC1/NC1 and

FC2/NC2), and the timing of the lifecycle was staggered between blocks to be 7 days apart.

Testing for resilience to heat stress

Experimental generation 1: Partitioning current care environment from evolutionary

history of care

Beetles were taken for analysis after 47 generations of experimental evolution. We partitioned

the effects of current care experienced by individuals (No Care or Full Care), from an

evolutionary history of care experienced by the source population (No Care or Full Care), by

breeding beetles for one generation using the following treatments:

1) Adults from Full Care populations allowed to provide full parental care (Full Care

parents with a Full Care regime) = FC_{FC}

2) Adults from Full Care populations removed at 53 hours after pairing (Full Care parents

with a No Care regime) = FC_{NC}

3) Adults from No Care populations removed at 53 hours after pairing (No Care parents

with a No Care regime) = NC_{NC}

4) Adults from No Care populations allowed to provide full parental care (No Care parents

with a Full Care regime) = NC_{FC}

When larval development was complete, and larvae were starting to disperse away from the

remains of the carrion nest (8 days after pairing), larvae were put in pupation boxes to mature

and reach eclosion, as described above.

Experimental generation 2: Exposure to heat stress

Immediately after eclosion, each experimental individual was placed alone in a small plastic

box (12 x 8 x 6 cm) containing a fine layer of soil. The boxes were placed in incubators

(Panasonic MLR-352H-PE) set to either 22 °C, 25 °C or 28 °C, 0% humidity and a 16L: 8D

hour light cycle. The boxes were removed briefly once weekly so that the beetles could be fed

with beef mince. We then measured the following correlates of fitness for this generation of

beetles:

Survival

To measure the effect of heat stress on survival, after 14 days (at sexual maturity) all individuals

were removed from their incubators and classified as alive or dead.

The surviving individuals were then assigned to one of the following treatments for breeding:

5) 22 °C male x 22 °C female

6) 25 °C male x 25 °C female

7) 28 °C male x 28 °C female

Individuals were only bred within their populations and temperatures, e.g. 22 °C males from

FC_{FC} were only bred with 22 °C females from FC_{FC}, resulting in 12 treatments (Figure S1).

Pairs were placed in breeding boxes containing 350 ml of moist soil and a 10-13 g mouse

carcass, and the box was sealed and kept in a dark environment, as described under 'general

methods' above. All pairs were kept at the standard laboratory temperature of 22 °C, regardless

of their temperature treatment, to isolate the effects of the thermal environment during sexual

maturation on subsequent reproductive success.

Lytic activity

At 53 h after pairing, we collected anal exudates from parents from a subsample of broods from

each treatment (n = 27 FC_{FC}, 27 FC_{NC}, 22 NC_{NC}, 19 NC_{FC}) by holding each beetle by the

abdomen and pressing a 5 ul glass microcapillary tube (Drummond) to its anus. Each sample

was pipetted immediately into a 1.5 ml Eppendorf tube on dry ice and subsequently stored at -

80 °C until analysis. We quantified the lytic activity of each sample in an automated microplate

reader (BioTek ELx808) by running turbidity assays that measured the degradation of bacterial

cell walls (see Duarte et al., 2021 for full details). Briefly, 1 ul of each exudate sample was

diluted in 24 ul of 6.4 pH 0.02 M potassium phosphate buffer and 10 ul of this solution was

added to a well of a 96-well flat-bottomed microplate (Thermo Scientific Nunc MicroWell)

containing 100 ul of a 1.3 mg ml^{-1} solution of lyophilized *Micrococcus lysodeikticus* cells in

potassium phosphate buffer. The microplate reader was set to 25 °C and took an initial reading

at 450 nm. Plates then went through a cycle of being shaken at a medium setting every 10

minutes, followed immediately by a 450 nm reading, for 60 minutes.

Carcass sphericity

At 53 h, we also briefly removed the carcass from a subsample of broods (n = 33 FC$_{FC}$, 31

FC$_{NC}$, 33 NC$_{NC}$, 33 NC$_{FC}$) where the parents had experienced the 22 °C and 28 °C degree

treatments. Each carcass was placed on a white background, alongside a scale for measurement,

and photographed from perpendicular angles using two cameras (Canon DSLR, with 18-55mm

lens) at 30 cm away from the carcass (Figure S2).

At 53 h after pairing, we removed the parents from all broods and counted the number of eggs

laid, through inspection of the underside of the box (following Jarrett et al., 2017; Schrader et

al., 2015c). At 70, 80, 94 and 104 h after pairing we checked the carcass and the surface of the

soil for any first instar larvae, and counted them before discarding them.

Statistical analyses

All statistical tests were conducted in R version 3.6.1 (R Core Team, 2019) and a 0.05

significance threshold was used throughout. Data handling and visualisation were carried out

using base R and the 'tidyverse' suite of R packages (Wickham et al., 2019). A stepwise

deletion method using F and chi-squared tests (as appropriate), implemented in the base

'statistics' package in R, was used to determine significance of each term and remove non-

significant terms sequentially (Crawley, 2007). For all analyses, temperature was modelled

first as a continuous variable and then as a factor and the model fit examined. Model fit was

best when temperature was treated as a continuous variable so it was treated as such in all the

models presented here.

Survival

We ran a Gaussian linear model with survival status as the dependent variable ("0" = died in

the incubator, "1" = survived to sexual maturity), and population evolutionary history (Full

Care or No Care), generation 1 care regime (Full Care or No Care), temperature, sex, block

and the two-way interactions of these terms as the independent variables.

Lytic activity

In some of the broods only one of the parents produced enough exudates to allow us to conduct

the lytic activity analysis on their sample. The total useable sample size for exudates was 150

individuals, and for 110 of these we had a sufficient sample from their partner (55 pairs) and

in the other 40 only one partner in the pair could be analysed (we could not collect enough

exudate from the partners of 20 females and 20 males used in the analysis).

We used the percentage change between the 450 nm absorption at the first reading (0 minutes)

and at 60 minutes as the dependent variable (as per Duarte et al., 2021). We first ran a paired

t-test to determine whether there was a consistent sex difference within pairs in lytic activity.

As the lytic activity of one individual in a pair may influence its partner's levels, we then ran

separate binomial generalised linear models for males and females, with proportion of cells

degraded (450 nm reading at 60 minutes/450 nm reading at 0 minutes) in either males' or

females' anal exudate as the dependent variable, and population evolutionary history (Full Care

or No Care), generation 1 care regime (Full Care or No Care), temperature, partner's lytic

activity (if data were available), block, and the two-way interactions between temperature and

evolutionary history, and sex and temperature as independent variables.

Carcass sphericity

Carcass sphericity was calculated digitally in ImageJ (Schneider et al., 2012), using a macro

script adapted from De Gasperin et al (2016) (Script S1 and Figure S3). The ImageJ script

assumes that if a carcass were perfectly spherical, the two-dimensional images in the carcass

from the top and side would be exact circles. The script compares the actual images to a perfect

circle of the same perimeter and then calculates the difference between the two – such that a

perfectly spherical carcass would have a score of 1 to denote that it fills 100% of the space of

a perfect sphere (Figure S4).

We ran a binomial GLM with carcass sphericity as the dependent variable and population

evolutionary history (Full Care or No Care), generation 1 care regime (Full Care or No Care),

temperature, sex, block, and the two-way interactions of these terms as the independent

variables.

Failure to lay, clutch size, hatching failure and brood size

To investigate predictors of fecundity, we first ran a binomial GLM with clutch failure ("0" =

no eggs laid and "1" = at least one egg laid) as the dependent variable and population

evolutionary history (Full Care or No Care), generation 1 care regime (Full Care or No Care),

temperature, carcass mass, block, and two-way interactions as listed in Table 4A as

independent variables.

Then to examine predictors of clutch size in the broods that did lay eggs, we ran a Gaussian

GLM with clutch size (the number of eggs counted from the underside of the breeding box) as

the dependent variable, and temperature, carcass mass, block, carcass roundness, and two-way

interactions as listed in all two-way interactions as listed in Table 4B as independent variables.

We then ran two tests to investigate predictors of fertility. Firstly, we ran a binomial GLM with

brood success ("0" = no larvae, "1" = at least one larva alive at dispersal) as the dependent

variable and the same independent variables that were used in the binomial model analysing

clutch failure vs success. Next, we ran a quasi-Poisson GLM with brood size (number of larvae

counted on the carcass or soil) as the dependent variable and population evolutionary history

(Full Care or No Care), generation 1 care regime (Full Care or No Care), temperature, carcass

mass, block, clutch size, and two-way interactions as listed in all two-way interactions as listed

in Table 5B as independent variables.

Results

Survival

Beetles from populations with an evolutionary history of No Care were more likely to die in

the incubators ($\chi^2_{1,\,N=1218}$ = 20.244, p = <0.001, Table 1), regardless of the temperature they

experienced in the incubator or the care they received in generation 1 of the heat stress

experiment. Mean survival of Full Care individuals across blocks was 94%, but 87% for No

Care individuals. Beetles exposed to greater heat stress as they matured sexually were more

likely to die, but only in block 2. Temperature had no equivalent effect on mortality in block 1

(block x temperature interaction: $\chi^2_{1,\,N=1218}$ = 17.284, p = <0.001, Table 1; Figure 1).

Table 1 – Results of binomial model investigating the effects on likelihood of a beetle surviving from

eclosion to sexual maturity in incubators set to 22 °C, 25 °C or 28 °C.

Independent variable	Estimate	SE	χ^2	df	*p*
Intercept	7.628	2.716			
Evolutionary History (No Care)	**-1.008**	**0.231**	**20.244**	**1**	**<0.001**
Generation 1 Parental Care Regime (none)	-0.223	0.223	1.008	1	0.316
Temperature (°C)	**-0.137**	**0.104**	**109.86**	**1**	**<0.001**
Block (2)	**14.322**	**3.733**	**74.003**	**1**	**<0.001**
Sex (male)	0.076	0.222	0.117	1	0.733
Evolutionary History (No Care): Temperature (°C)	0.111	0.161	0.492	1	0.483
Gen 1 Parental Care Regime (none): Temperature (°C)	0.141	0.137	1.089	1	0.297
Temperature (°C):Sex (male)	0.113	0.133	0.735	1	0.391
Evolutionary History:Sex	-0.513	0.461	1.245	1	0.265
Gen 1 Parental Care Regime (none):Sex (male)	0.389	0.452	0.744	1	0.388
Gen 1 Parental Care Regime (none):Block (2)	0.160	0.589	0.074	1	0.785
Evolutionary History (No Care):Block (2)	-0.377	0.571	0.514	1	0.426

Temperature (°C):Block (2)	-0.607	0.140	17.284	1	<0.001
Sex (male):Block (2)	0.243	0.581	0.175	1	0.676

Figure 1 – The survival of beetles kept in incubators at different levels of heat stress throughout sexual

maturation (1 = survived, 0 = died) in relation to block. Points represent data for individual pairs, lines

represent the reaction norms for each block (1 = red, 2 = blue) and the shaded areas indicate the 95

% confidence intervals. N = 1,281 individuals.

To further investigate how block affected survival at higher temperatures we split the dataset

by block and ran generalised linear models with survival probability as the dependent variable

and temperature as the independent variable. In block 1 temperature did not significantly affect

survival, but in block 2 there was a significant negative effect ($\chi^2_{1, N=600}$ = 122.19, p = <0.001)

Lytic activity

There was no difference between males and females in lytic activity strength: $t(54) = 0.446$, *p*

$= 0.658$). Lytic activity in both sexes was only significantly affected by block, with individuals

having stronger lytic activity in block 2 (Figure S5, Table 2).

Table 2 – Results of binomial GLMs investigating the effects on A) male and B) female exudate lytic activity at 53h after pairing. We have defined lytic activity

as the proportion of cell degradation of Micrococcus lysodeikticus *as recorded at 450 nm after 60 minutes*

Independent variable	A) Male exudate					B) Female exudate				
	Estimate	SE	χ^2	df	p	Estimate	SE	χ^2	df	p
Intercept	0.065	0.026				0.067	0.028			
Female Exudate Lytic Activity	-0.003	0.002	1.405	1	0.236	-0.003	0.003	1.391	1	0.238
Evolutionary History (No Care)	0.033	0.041	0.648	1	0.421	-0.023	0.041	0.300	1	0.584
Generation 1 Parental Care Regime (none)	-0.048	0.041	1.063	1	0.303	0.022	0.042	0.290	1	0.590
Temperature	-0.006	0.007	0.793	1	0.373	-0.006	0.007	0.633	1	0.426
Carcass mass (g)	-0.020	0.029	0.484	1	0.487	-0.008	0.029	0.075	1	0.785
Block (2)	**0.164**	**0.034**	**22.906**	**1**	**<0.001**	**0.165**	**0.036**	**21.588**	**1**	**<0.001**

Evolutionary History (No Care): Generation 1 Parental Care Regime (none)	-0.003	0.083	0.001	1	0.970	0.049	0.082	0.358	1	0.550
Temperature: Evolutionary History (No Care)	-0.010	0.014	0.568	1	0.451	-0.005	0.014	0.146	1	0.703
Temperature: Generation 1 Parental Care Regime (none)	0.010	0.014	00.537	1	0.464	-0.008	0.014	0.350	1	0.554
Evolutionary History (No Care): Block (2)	0.023	0.082	0.082	1	0.774	-0.048	0.082	0.339	1	0.561

352   ***Carcass sphericity***

354   No variables tested explained a significant amount of variation in carcass sphericity (all *p*

355   values = >0.05; Table 3).

358   *Table 2 – Results of binomial GLM investigating variation in carcass nest roundness as a proxy for*

359   *carcass preparation success*

Independent variable	Estimate	S.E.	χ^2	d.f.	*p*
Intercept	-0.075	0.176			
Female Exudate Lytic Activity	-0.008	0.02	0.083	1	0.773
Male Exudate Lytic Activity	0.012	0.037	0.095	1	0.758
Evolutionary History (No Care)	0.026	0.610	0.002	1	0.966
Generation 1 Parental Care Regime (none)	-0.208	0.692	0.091	1	0.763
Temperature	-0.084	0.059	2.029	1	0.154
Carcass mass (g)	-0.094	0.490	0.037	1	0.848
Block (2)	0.227	1.205	0.036	1	0.851
Evolutionary History (No Care): Generation	0.333	1.379	0.058	1	0.809

1 Parental Care Regime (none)					
Temperature: Evolutionary History (No Care)	-0.058	0.245	0.056	1	0.814
Temperature: Generation 1 Parental Care Regime (none)	0.018	0.243	0.006	1	0.941
Carcass mass: Evolutionary History (No Care)	0.310	0.966	0.103	1	0.749
Carcass mass: Generation 1 Parental Care Regime (none)	-0.056	1.106	0.003	1	0.960

Failure to lay eggs and clutch size

The population's evolutionary history of care (Full Care or No Care) did not explain significant

variation in clutch size, and nor did the care regime experienced in the first generation of the

heat stress experiment (Table 4A). Exposure to higher temperatures during sexual maturation

were more likely to cause complete failure to lay, and there was a greater chance of complete

laying failure in block 2 (Table 4A, Figure 2). In broods where some laying occurred, higher

temperatures led to fewer eggs laid per clutch ($F_{1,304} = 166.8$, p=<0.001, Table 4B, Figure S6).

Table 3 – Results of GLMs looking at variables affecting A) failure to lay: no eggs produced versus at

least one egg laid and B) clutch size for pairs that laid at least one egg

A) Failure to lay					
Independent variable	**Estimate**	**S.E.**	**χ^2**	**d.f.**	***p***
Intercept	39.167	4.828			
Evolutionary History (No Care)	0.457	0.369	1.545	1	0.214
Generation 1 Parental Care Regime (none)	-0.041	0.364	0.013	1	0.910
Temperature	**-1.383**	**0.173**	**183.99**	**1**	**<0.001**
Carcass mass (g)	-0.140	0.269	0.270	1	0.603
Block (2)	**-2.082**	**0.422**	**30.439**	**1**	**<0.001**
Evolutionary History (No Care): Temperature	0.464	0.321	2.219	1	0.136
Gen 1 Parental Care Regime (none): Temperature	0.641	0.392	3.450	1	0.063
Evolutionary History (No Care): Gen 1 Parental Care Regime (none)	0.630	0.783	0.653	1	0.419
Block (2):Temperature	4.967	347.908	1.760	1	0.185
B) Clutch size					
Independent variable	**Estimate**	**S.E.**	**F**	**d.f.**	**p**
Intercept	86.468	5.153			

Evolutionary History (No Care)	2.234	2.734	0.668	1	0.421
Generation 1 Parental Care Regime (none)	-1.243	0.903	1.898	1	0.169
Temperature	**-2.717**	**0.210**	**166.8**	**1**	**<0.001**
Carcass mass (g)	-0.422	2.511	0.028	1	0.868
Carcass roundness	1.722	4.733	0.132	1	0.717
Block (2)	1.534	2.606	2.393	1	0.125
Male exudate lytic activity	0.205	0.154	1.774	1	0.193
Female exudate lytic activity	0.067	0.254	0.069	1	0.796
Evolutionary History (No Care): Temperature	-1.817	0.990	3.369	1	0.076
Gen 1 Parental Care Regime (none): Temperature	0.161	1.890	0.007	1	0.934
Evolutionary History (No Care): Gen 1 Parental Care Regime (none)	5.602	5.423	1.067	1	0.311
Evolutionary History (No Care): Carcass roundness	-12.378	27.842	0.001	1	0.972
Gen 1 Parental Care Regime (none): Carcass roundness	14.780	9.143	2.613	1	0.110
Block (2):Temperature	-0.776	0.999	0.604	1	0.444

Figure 2 – The effect of temperature and block on laying status of pairs of beetles kept in incubators of

22 °C, 25 °C and 28 °C during sexual maturation (1 = at least 1 egg laid, 0 = no eggs laid). Each point

represents a different clutch, lines represent the reaction norms for each block (1 = red, 2 = blue) and

the shaded areas represent the 95 % confidence intervals. N = 383 clutches.

Hatching failure and brood size

There was no significant difference in hatching failure mediated by parents' evolutionary

history (Table 5A). Parents kept at higher incubator temperatures between eclosion and sexual

maturity were more likely to experience complete hatching failure (Figure 3, Table 5A) - with

no eggs hatching in any of the broods where the parents had developed at 28 °C.

390   Fewer larvae hatched when parents had been kept at 25 °C than 22 °C ($F_{1, N=199}$ = 186.72, p =

391   <0.001) (Figure S7), more larvae hatched in block 2, and more larvae hatched when there were

392   more eggs to start with ($F_{1, N=199}$ = 38.935, p = <0.001, Table 5B).

395   *Table 4 - Results of GLMs looking at variables affecting A) hatching success (at least one larva) or*

396   *failure (no larvae) and B) brood size for pairs where at least one larva hatched*

A) Hatching failure					
Independent variable	Estimate	S.E.	χ^2	d.f.	p
Intercept	46.570	8.380			
Evolutionary History (No Care)	-0.238	0.341	0.486	1	0.486
Generation 1 Parental Care Regime (none)	-0.330	0.343	0.931	1	0.335
Temperature	**-1.849**	**0.335**	**323.41**	**1**	**<0.001**
Carcass mass (g)	-0.338	0.233	2.138	1	0.144
Block (2)	0.579	0.340	2.946	1	0.086
Evolutionary History (No Care): Temperature	4.961	314.209	1.569	1	0.210
Gen 1 Parental Care Regime (none): Temperature	-5.158	482.152	1.298	1	0.255
Evolutionary History (No Care): Gen 1 Parental Care Regime (none)	-0.738	0.690	1.149	1	0.284

Block (2):Temperature	-5.420	766.035	1.152	1	0.283
B) Brood size					
Independent variable	**Estimate**	**S.E.**	**F**	**d.f.**	***p***
Intercept	10.392	0.685			
Evolutionary History (No Care)	0.084	0.070	1.428	1	0.234
Generation 1 Parental Care Regime (none)	-0.032	0.070	0.208	1	0.649
Temperature	**-0.366**	**0.030**	**186.72**	**1**	**<0.001**
Carcass mass (g)	-0.082	0.048	2.872	1	0.092
Block (2)	**0.289**	**0.071**	**16.917**	**1**	**<0.001**
Clutch size	**0.029**	**0.005**	**38.935**	**1**	**<0.001**
Evolutionary History (No Care): Temperature	0.035	0.060	0.349	1	0.555
Gen 1 Parental Care Regime (none): Temperature	-0.044	0.059	0.560	1	0.455
Evolutionary History (No Care): Gen 1 Parental Care Regime (none)	-0.133	0.140	0.901	1	0.344
Block (2):Temperature	0.065	0.060	1.163	1	0.282

Figure 3 – The effect of incubator temperature during sexual maturation on hatching failure (1 = at least one larva hatched) or failure (0 = no larvae hatched). Each black datapoint represents one clutch, red points show the mean for each temperature treatment, the thick black lines represent the median for each treatment. N = 383 clutches.

Discussion

We tested experimentally whether a population's past sustained exposure to benign versus harsh social environments predicted its capacity to cope with heat stress during sexual maturation. The key findings of our experiments are summarised in Table 6, for each of the correlates of fitness we measured.

 Table 6 – The key predictors tested, their effects on correlates of fitness measured, the slope estimates for each significant variable, and the hypothesis supported

 by the findings. "None" indicates that the predictor variable did not have a significant effect on the correlate of fitness. "Est" refers to the slope estimate for

 the predictor in the minimum adequate model. Full statistical results are presented in Tables 1-5.

Predictor variables tested	Effect on correlates of fitness							Hypothesis supported
	Survival	Lytic activity	Carcass sphericity	Failure to lay	Clutch size	Hatching failure	Brood size	
Population of origin (No Care or Full Care)	Lower survival in No Care (est: -1.008)	None	None	None	None	None	None	Beneficial mutation
Block	No effect in block 1, lower survival at higher temperature	Stronger lytic activity in block 2 (est for males:	None	Higher failure in block 2 (est: -2.082)	None	None	Larger broods in block 2 (est: 0.289)	Founder effect

	in block 2 (est for block 2: -0.723)	0.164; for females: 0.165)						
Temperature	Lower survival at higher temperature in block 2 (as cell above)	None	None	Higher failure at higher temperature (est: -1.383)	Smaller clutch size at higher temperature (est: -2.717)	Higher hatching failure at higher temperature (est: -1.849)	Smaller brood size at higher temperature (est: -0.366)	N/A

We found that the more benign the social environment previously experienced by a population

the better its survival during sexual maturation – which is consistent with the "beneficial

mutation hypothesis". Individuals derived from the No Care populations had poor survival in

the incubators, even in the most benign thermal environment, compared with those from the

Full Care populations. We have previously shown that individuals in No Care populations carry

a lower mutation load (Pascoal et al. 2022). Perhaps this means that, by chance, they lack the

cryptic genetic variation that enables them to cope with new levels of thermal stress during

sexual maturation (Paaby & Rockman, 2014). Alternatively, perhaps the No Care populations

were intrinsically inferior by this point, and more prone to dying between eclosion and sexual

maturity, regardless of the temperature to which they were exposed. For all other correlates of

fitness we could detect no differences between individuals drawn from the No Care versus Full

Care populations, which argues against this interpretation. Overall, our experimental data allow

us to reject the "mutation load hypothesis" and the "selection filter hypothesis". But perhaps

we have insufficient data to conclude we have found strong support for the "beneficial mutation

hypothesis".

Unexpectedly, we found that experimental block predicted some correlates of fitness (Table

1), which could be due stochastically to a founder effect. The four distinct evolving populations

were established from a single large stock population derived by interbreeding beetles from

four different wild populations. The size of the breeding population was identical for each

block throughout the experiment (n = 100 pairs), as was the maintenance regime, which implies

that any differences between blocks must be attributable to chance differences in genetic

variation when the population was founded. In analyses of earlier generations of the

experimentally evolving populations, we found that selection from social environment

explained divergence between populations in the traits we analysed to a greater extent than

block (e.g. Duarte et al., 2021; Jarrett et al., 2018a, 2018b; Schrader et al., 2015c, 2015a, 2015b,

2017, 2017). By generation 49, however, we find that founder effects are outweighing

differences due to selection from the social environment – at least on the response to heat stress

exposure during sexual maturation.

Founder effects have long been suggested to explain the success of translocation and captive

breeding in conservation programmes (e.g. Brown et al., 2009; Dolman et al., 2021; Miller et

al., 1999). However, in analyses of wild populations it is often hard to disentangle the scale of

founder effects from confounding factors of environmental differences (however subtle) in the

new location (Kolbe et al., 2012; Szűcs et al., 2017) especially when it is difficult to monitor

populations systematically after their release into the wild (Armstrong & Seddon, 2008; Mock

et al., 2004; Seddon et al., 2007; Weeks et al., 2011). Nevertheless, there is some evidence that

founder effects influence the response to selection in wild populations. When *Anolis sagrei*

individuals with long hindlimbs were translocated to seven Bahamian islands with a vegetation

structure known to favour short hindlimbed individuals, they evolved shorter hindlimbs over

four years. However the extent of morphological change was attributable to founder effects

(Kolbe et al., 2012). Perhaps a similar pattern explains the results we have found in this study.

More broadly our study indicates the importance of the initial population size in captive

breeding programmes. It also reinforces the wisdom of using genetic screening to determine

whether the founding population is representative of the source population and has high

heterozygosity (Bragg et al., 2021; Scott et al., 2020), especially in cases where there will no

migration into the population, such as on an isolated island.

For some of the traits we measured (failure to lay, clutch size, failure to hatch) temperature had

such a strong negative effect that it outweighed any effects that could be attributed to the

population's genetic background. This is consistent with previous work across different

species of burying beetle, which found that high temperatures during breeding led to reduced

clutch and brood sizes and a greater likelihood of complete brood failure (Keller et al., 2021;

Moss & Moore, 2021; Ney, 2021; Ong, 2019; Quinby et al., 2020; Sun & Kilner, 2020).

However, in all of these studies the extreme temperature treatment continued throughout

breeding, whereas in this study we elevated temperature only during sexual maturation to better

simulate the natural history of the burying beetle. Our results suggest that there are sensitive

windows in development which affect breeding success even when earlier stages of

reproduction have occurred under benign conditions. In a world with an increasing frequency

and intensity of heat waves (Perkins-Kirkpatrick & Lewis, 2020), this study shows that an

extreme temperature event can also reduce breeding success even if it is only brief, as long as

it coincides with these sensitive windows in development.

There are two likely causes for a reduction in egg laying and egg hatching at higher

temperatures: 1) behavioural – beetles maturing at higher temperatures mated less, and/or 2)

physiological – beetles maturing at higher temperatures had underdeveloped gonads.

Particularly in males, exposure to high temperatures prior to pairing leads to reduced courtship

behaviour in *Aphidius avenae* (Roux et al., 2010) and *Drosophila melanogaster, Drosophila*

simulans and Drosophila mojavensis (Patton & Krebs, 2001). There is also a great deal of

evidence to suggest that thermal stress during sexual maturation can have a physiological effect

on insects. In *Drosophila melanogaster* high temperatures during development cause a range

of spermatid cytological abnormalities (David et al., 2005; Rohmer et al., 2004) and in

Tribolium castaneum cause lower testis volume (Sales et al., 2021). In females, higher

temperatures during development cause smaller oocytes in *Pararge aegeria* (Gibbs et al., 2010)

and cause *Bicyclus anynana* to lay more, smaller eggs (Steigenga & Fischer, 2007). From this

study we are unable to say whether the effects were behavioural, physiological or both, and

whether it was the males or females that drove the patterns of reduced reproductive fitness at

higher temperatures. Further investigations tracking copulatory behaviour, mating thermally

stressed males with control females and vice versa, and dissecting ovaries and testes following

heat stress are needed to answer these questions.

It is widely accepted that a population's standing genetic variation will predict its response to

new environmental challenges. Our study suggests that the social evolutionary history of

burying beetle populations might predict the response to heat stress during sexual maturation

to some extent, although further work is needed before a robust conclusion can be drawn.

However, stochastic genetic variation, due to founder effects or genetic drift, was potentially a

more powerful predictor of population resilience upon exposure to new environments.

Translocation and captive breeding programmes should be designed accordingly to maximise

their chance of success.

References

Araki, H., Cooper, B., & Blouin, M. S. (2007). Genetic Effects of Captive Breeding Cause a

Rapid, Cumulative Fitness Decline in the Wild. *Science, 318*(5847), 100.

https://doi.org/10.1126/science.1145621

Arce, A. N., Johnston, P. R., Smiseth, P. T., & Rozen, D. E. (2012). Mechanisms and fitness

effects of antibacterial defences in a carrion beetle. *Journal of Evolutionary Biology*,

25(5), 930–937. https://doi.org/10.1111/j.1420-9101.2012.02486.x

Armstrong, D. P., & Seddon, P. J. (2008). Directions in reintroduction biology. *Trends in*

Ecology & Evolution, *23*(1), 20–25. https://doi.org/10.1016/j.tree.2007.10.003

Balmford, A. (1996). Extinction filters and current resilience: The significance of past

selection pressures for conservation biology. *Trends in Ecology & Evolution*, *11*(5),

193–196. https://doi.org/10.1016/0169-5347(96)10026-4

Barrett, R. D. H., & Schluter, D. (2008). Adaptation from standing genetic variation. *Trends*

in Ecology & Evolution, *23*(1), 38–44. https://doi.org/10.1016/j.tree.2007.09.008

Bragg, J. G., Yap, J.-Y. S., Wilson, T., Lee, E., & Rossetto, M. (2021). Conserving the

genetic diversity of condemned populations: Optimizing collections and translocation.

Evolutionary Applications, *14*(5), 1225–1238. https://doi.org/10.1111/eva.13192

Brown, S. K., Hull, J. M., Updike, D. R., Fain, S. R., & Ernest, H. B. (2009). Black Bear

Population Genetics in California: Signatures of Population Structure, Competitive

Release, and Historical Translocation. *Journal of Mammalogy*, *90*(5), 1066–1074.

https://doi.org/10.1644/08-MAMM-A-193.1

Capodeanu-Nägler, A., Keppner, E. M., Vogel, H., Ayasse, M., Eggert, A.-K., Sakaluk, S.

K., & Steiger, S. (2016). From facultative to obligatory parental care: Interspecific

variation in offspring dependency on post-hatching care in burying beetles. *Scientific*

Reports, *6*(1), 29323. https://doi.org/10.1038/srep29323

Carroll, S. P., Jørgensen, P. S., Kinnison, M. T., Bergstrom, C. T., Denison, R. F., Gluckman,

P., Smith, T. B., Strauss, S. Y., & Tabashnik, B. E. (2014). Applying evolutionary

biology to address global challenges. *Science*, *346*(6207).

https://doi.org/10.1126/science.1245993

Clutton-Brock, T. H. (1991). *The Evolution of Parental Care* (Vol. 64). Princeton University

Press; JSTOR. https://doi.org/10.2307/j.ctvs32ssj

Cotter, S., & Kilner, R. M. (2010). Sexual division of antibacterial resource defence in

breeding burying beetles, *Nicrophorus vespilloides*. *Journal of Animal Ecology*,

79(1), 35–43. https://doi.org/10.1111/j.1365-2656.2009.01593.x

Crawley, M. J. (2007). Multiple regression. In *The R Book* (pp. 569–591). John Wiley &

Sons, Ltd. https://doi.org/10.1002/9780470515075.ch16

David, J. R., Araripe, L. O., Chakir, M., Legout, H., Lemos, B., Pétavy, G., Rohmer, C., Joly,

D., & Moreteau, B. (2005). Male sterility at extreme temperatures: A significant but

neglected phenomenon for understanding Drosophila climatic adaptations. *Journal of*

Evolutionary Biology, *18*(4), 838–846. https://doi.org/10.1111/j.1420-

9101.2005.00914.x

De Gasperin, O., Duarte, A., Troscianko, J., & Kilner, R. M. (2016). Fitness costs associated

with building and maintaining the burying beetle's carrion nest. *Scientific Reports*, *6*,

35293.

Dolman, P. M., Burnside, R. J., Scotland, K. M., & Collar, N. J. (2021). Captive breeding and

the conservation of the threatened houbara bustards. *Endangered Species Research*,

46, 161–173.

Duarte, A., Rebar, D., Hallett, A. C., Jarrett, B. J. M., & Kilner, R. M. (2021). Evolutionary

change in the construction of the nursery environment when parents are prevented

from caring for their young directly. *Proceedings of the National Academy of*

Sciences, *118*(48), e2102450118. https://doi.org/10.1073/pnas.2102450118

Eggert, A.-K., Reinking, M., & Müller, J. K. (1998). Parental care improves offspring

survival and growth in burying beetles. *Animal Behaviour*, *55*(1), 97–107.

https://doi.org/10.1006/anbe.1997.0588

Elsbeth McPhee, M. (2004). Generations in captivity increases behavioral variance:

Considerations for captive breeding and reintroduction programs. *Biological*

Conservation, *115*(1), 71–77. https://doi.org/10.1016/S0006-3207(03)00095-8

Gibbs, M., Van Dyck, H., & Karlsson, B. (2010). Reproductive plasticity, ovarian dynamics

and maternal effects in response to temperature and flight in Pararge aegeria. *Journal*

of Insect Physiology, *56*(9), 1275–1283.

https://doi.org/10.1016/j.jinsphys.2010.04.009

Graham, D. W., Bergeron, G., Bourassa, M. W., Dickson, J., Gomes, F., Howe, A., Kahn, L.

H., Morley, P. S., Scott, H. M., Simjee, S., Singer, R. S., Smith, T. C., Storrs, C., &

Wittum, T. E. (2019). Complexities in understanding antimicrobial resistance across

domesticated animal, human, and environmental systems. *Annals of the New York*

Academy of Sciences, *1441*(1), 17–30. https://doi.org/10.1111/nyas.14036

Hawkins, N. J., Bass, C., Dixon, A., & Neve, P. (2019). The evolutionary origins of pesticide

resistance. *Biological Reviews*, *94*(1), 135–155. https://doi.org/10.1111/brv.12440

Jarrett, B. J. M., Evans, E., Haynes, H. B., Leaf, M. R., Rebar, D., Duarte, A., Schrader, M.,

& Kilner, R. M. (2018). A sustained change in the supply of parental care causes

adaptive evolution of offspring morphology. *Nature Communications*, *9*(1), 3987.

https://doi.org/10.1038/s41467-018-06513-6

Jarrett, B. J. M., Rebar, D., Haynes, H. B., Leaf, M. R., Halliwell, C., Kemp, R., & Kilner, R.

M. (2018). Adaptive evolution of synchronous egg-hatching in compensation for the

loss of parental care. *Proceedings of the Royal Society B: Biological Sciences*,

285(1885), 20181452. https://doi.org/10.1098/rspb.2018.1452

Jarrett, B. J. M., Schrader, M., Rebar, D., Houslay, T. M., & Kilner, R. M. (2017).

Cooperative interactions within the family enhance the capacity for evolutionary

change in body size. *Nature Ecology & Evolution*, *1*, 0178.

http://dx.doi.org/10.1038/s41559-017-0

Kalinová, B., Podskalská, H., Růžička, J., & Hoskovec, M. (2009). Irresistible bouquet of

death—How are burying beetles (Coleoptera: Silphidae: *Nicrophorus*) attracted by

carcasses. *Naturwissenschaften*, *96*(8), 889–899. https://doi.org/10.1007/s00114-009-

0545-6

Keller, M. L., Howard, D. R., & Hall, C. L. (2021). The thermal ecology of burying beetles:

Temperature influences reproduction and daily activity in Nicrophorus marginatus.

Ecological Entomology, *46*(6), 1266–1272. https://doi.org/10.1111/een.13071

Kolbe, J. J., Leal, M., Schoener, T. W., Spiller, D. A., & Losos, J. B. (2012). Founder Effects

Persist Despite Adaptive Differentiation: A Field Experiment with Lizards. *Science*,

335(6072), 1086–1089. https://doi.org/10.1126/science.1209566

Linksvayer, T. A., & Wade, M. J. (2009). Genes with social effects are expected to harbor

more sequence variation within and between species. *Evolution; International*

Journal of Organic Evolution, *63*(7), 1685–1696. PubMed.

https://doi.org/10.1111/j.1558-5646.2009.00670.x

Lowerre-Barbieri, S., DeCelles, G., Pepin, P., Catalán, I. A., Muhling, B., Erisman, B.,

Cadrin, S. X., Alós, J., Ospina-Alvarez, A., Stachura, M. M., Tringali, M. D.,

Burnsed, S. W., & Paris, C. B. (2017). Reproductive resilience: A paradigm shift in

understanding spawner-recruit systems in exploited marine fish. *Fish and Fisheries*,

18(2), 285–312. https://doi.org/10.1111/faf.12180

Lumley, A. J., Michalczyk, Ł., Kitson, J. J. N., Spurgin, L. G., Morrison, C. A., Godwin, J.

L., Dickinson, M. E., Martin, O. Y., Emerson, B. C., Chapman, T., & Gage, M. J. G.

(2015). Sexual selection protects against extinction. *Nature*, *522*, 470.

Lynch, M. (2016). Mutation and Human Exceptionalism: Our Future Genetic Load. *Genetics*, *202*(3), 869–875. https://doi.org/10.1534/genetics.115.180471

Martin, K., & Wiebe, K. L. (2004). Coping Mechanisms of Alpine and Arctic Breeding Birds: Extreme Weather and Limitations to Reproductive Resilience1. *Integrative and Comparative Biology*, *44*(2), 177–185. https://doi.org/10.1093/icb/44.2.177

Millenium Ecosystem Assessment. (2005). *Ecosystems and Human Well-Being: Synthesis*. World Resources Institute. https://www.millenniumassessment.org/documents/document.356.aspx.pdf

Miller, B., Ralls, K., Reading, R. P., Scott, J. M., & Estes, J. (1999). Biological and technical considerations of carnivore translocation: A review. *Animal Conservation*, *2*(1), 59–68. Cambridge Core. https://doi.org/10.1111/j.1469-1795.1999.tb00049.x

Mock, K. E., Latch, E. K., & Rhodes, O. E. (2004). Assessing losses of genetic diversity due to translocation: Long-term case histories in Merriam's turkey (Meleagris gallopavo merriami). *Conservation Genetics*, *5*(5), 631–645. https://doi.org/10.1007/s10592-004-1849-x

Moss, J. B., & Moore, A. J. (2021). Constrained flexibility of parental cooperation limits adaptive responses to harsh conditions. *Evolution*, *75*(7), 1835–1849. https://doi.org/10.1111/evo.14285

Müller, J. K., & Eggert, A.-K. (1989). Paternity assurance by "helpful" males: Adaptations to sperm competition in burying beetles. *Behavioral Ecology and Sociobiology*, *24*(4), 245–249. https://doi.org/10.1007/BF00295204

Ney, L. (2021). *Impact of temperature stress on reproduction, offspring development, and social immunity in the burying beetle Nicrophorus orbicollis* [University of Nebraska at Omaha]. https://www.proquest.com/docview/2585301523?pq-origsite=gscholar&fromopenview=true

Ong, J. Y. (2019). Coping with Thermal Challenges: Reaction Norms of Life History Traits

in a Burying Beetle with Biparental Care [M.S., Queen's University (Canada)]. In

ProQuest Dissertations and Theses (2186240977). ProQuest Dissertations & Theses

A&I. https://ezp.lib.cam.ac.uk/login?url=https://www.proquest.com/dissertations-

theses/coping-with-thermal-challenges-reaction-norms/docview/2186240977/se-

2?accountid=9851

Otronen, M. (1988). The effect of body size on the outcome of fights in burying beetles

(*Nicrophorus*). *Annales Zoologici Fennici*, *25*(2), 191–201.

Paaby, A. B., & Rockman, M. V. (2014). Cryptic genetic variation: Evolution's hidden

substrate. *Nature Reviews Genetics*, *15*, 247.

Pascoal, S., Mashoodh, R., Shimadzu, H., & Kilner, R. M. (2022). Parental care allows

genetic variation to accumulate by relaxing selection. *BioRxiv*, 2022.05.05.490718.

https://doi.org/10.1101/2022.05.05.490718

Patton, Z. J., & Krebs, R. A. (2001). The Effect of Thermal Stress on the Mating Behavior of

Three Drosophila Species. *Physiological and Biochemical Zoology*, *74*(6), 783–788.

https://doi.org/10.1086/323327

Perkins-Kirkpatrick, S. E., & Lewis, S. C. (2020). Increasing trends in regional heatwaves.

Nature Communications, *11*(1), 3357. https://doi.org/10.1038/s41467-020-16970-7

Pörtner, H. O., Roberts, D. C., Tignor, M., Poloczanska, E. S., Mintenbeck, K., Alegria, A.,

Craig, M., Langsdorf, S., Löschke, S., Möller, V., Okem, A., & Rama, B. (2022).

IPCC, 2022: Climate Change 2022: Impacts, Adaptation, and Vulnerability.

Contribution of Working Group II to the Sixth Assessment Report of the

Intergovernmental Panel on Climate Change. Cambridge University Press.

Quinby, B. M., Belk, M. C., & Creighton, J. C. (2020). Behavioral constraints on local

adaptation and counter-gradient variation: Implications for climate change. *Ecology*

and Evolution, *10*(13), 6688–6701. https://doi.org/10.1002/ece3.6399

R Core Team. (2019). *R: A language and environment for statistical computing*. R

Foundation for Statistical Computing. https://www.r-project.org

Rebar, D., Bailey, N. W., Jarrett, B. J. M., & Kilner, R. M. (2020). An evolutionary switch

from sibling rivalry to sibling cooperation, caused by a sustained loss of parental care.

Proceedings of the National Academy of Sciences, 201911677.

https://doi.org/10.1073/pnas.1911677117

Robert, A. (2009). Captive breeding genetics and reintroduction success. *Biological*

Conservation, *142*(12), 2915–2922. https://doi.org/10.1016/j.biocon.2009.07.016

Rohmer, C., David, J. R., Moreteau, B., & Joly, D. (2004). Heat induced male sterility in

Drosophila melanogaster: Adaptive genetic variations among geographic populations

and role of the Y chromosome. *Journal of Experimental Biology*, *207*(16), 2735–

2743. https://doi.org/10.1242/jeb.01087

Roux, O., Le Lann, C., van Alphen, J. J. M., & van Baaren, J. (2010). How does heat shock

affect the life history traits of adults and progeny of the aphid parasitoid Aphidius

avenae (Hymenoptera: Aphidiidae)? *Bulletin of Entomological Research*, *100*(5),

543–549. Cambridge Core. https://doi.org/10.1017/S0007485309990575

Sales, K., Vasudeva, R., Dickinson, M. E., Godwin, J. L., Lumley, A. J., Michalczyk, Ł.,

Hebberecht, L., Thomas, P., Franco, A., & Gage, M. J. G. (2018). Experimental

heatwaves compromise sperm function and cause transgenerational damage in a

model insect. *Nature Communications*, *9*(1), 4771. https://doi.org/10.1038/s41467-

018-07273-z

Sales, K., Vasudeva, R., & Gage, M. J. G. (2021). Fertility and mortality impacts of thermal

stress from experimental heatwaves on different life stages and their recovery in a

model insect. *Royal Society Open Science*, *8*(3). https://doi.org/10.1098/rsos.201717

Saniotis, A., Henneberg, M., & Mohammadi, K. (2021). Genetic load and biological changes

to extant humans. *Journal of Biosocial Science*, *53*(4), 639–642. Cambridge Core.

https://doi.org/10.1017/S0021932020000413

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years

of Image Analysis. *Nature Methods*, *9*(7), 671–675. PMC.

Schrader, M., Cosby, R., Hesketh, A., Jarrett, B. J. M., & Kilner, R. (2015). A limit on the

extent to which increased egg size can compensate for a poor postnatal environment

revealed experimentally in the burying beetle, *Nicrophorus vespilloides*. *Ecology and*

Evolution, *6*(1), 329–336. https://doi.org/10.1002/ece3.1876

Schrader, M., Jarrett, B. J. M., & Kilner, R. M. (2015a). Parental care masks a density-

dependent shift from cooperation to competition among burying beetle larvae.

Evolution; International Journal of Organic Evolution, *69*(4), 1077–1084. PMC.

https://doi.org/10.1111/evo.12615

Schrader, M., Jarrett, B. J. M., & Kilner, R. M. (2015b). Using Experimental Evolution to

Study Adaptations for Life within the Family. *The American Naturalist*, *185*(5), 610–

619. https://doi.org/10.1086/680500

Schrader, M., Jarrett, B. J. M., Rebar, D., & Kilner, R. M. (2017). Adaptation to a novel

family environment involves both apparent and cryptic phenotypic changes.

Proceedings of the Royal Society B: Biological Sciences, *284*(1862), 20171295.

https://doi.org/10.1098/rspb.2017.1295

Scott, P. A., Allison, L. A., Field, K. J., Averill-Murray, R. C., & Shaffer, B. H. (2020).

Individual heterozygosity predicts translocation success in threatened desert tortoises.

Science, *370*(6520), 1086–1089. https://doi.org/10.1126/science.abb0421

Seddon, P. J., Armstrong, D. P., & Maloney, R. F. (2007). Developing the Science of

Reintroduction Biology. *Conservation Biology*, *21*(2), 303–312.

https://doi.org/10.1111/j.1523-1739.2006.00627.x

Smiseth, P. T., Darwell, C. T., & Moore, A. J. (2003). Partial begging: An empirical model

for the early evolution of offspring signalling. *Proceedings of the Royal Society B:*

Biological Sciences, *270*(1526), 1773–1777. https://doi.org/10.1098/rspb.2003.2444

Snell-Rood, E. C., Burger, M., Hutton, Q., & Moczek, A. P. (2016). Effects of parental care

on the accumulation and release of cryptic genetic variation: Review of mechanisms

and a case study of dung beetles. *Evolutionary Ecology*, *30*(2), 251–265.

https://doi.org/10.1007/s10682-015-9813-4

Steigenga, M. J., & Fischer, K. (2007). Ovarian dynamics, egg size, and egg number in

relation to temperature and mating status in a butterfly. *Entomologia Experimentalis*

et Applicata, *125*(2), 195–203. https://doi.org/10.1111/j.1570-7458.2007.00610.x

Sun, S.-J., & Kilner, R. M. (2020). Temperature stress induces mites to help their carrion

beetle hosts by eliminating rival blowflies. *ELife*, *9*, e55649. PubMed.

https://doi.org/10.7554/eLife.55649

Szűcs, M., Melbourne, B. A., Tuff, T., Weiss-Lehman, C., & Hufbauer, R. A. (2017). Genetic

and demographic founder effects have long-term fitness consequences for colonising

populations. *Ecology Letters*, *20*(4), 436–444. https://doi.org/10.1111/ele.12743

Trumbo, S. T., & Steiger, S. (2020). Finding A Fresh Carcass: Bacterially-Derived Volatiles

And Burying Beetle Search Success. *BioRxiv*, 2020.01.25.919696.

https://doi.org/10.1101/2020.01.25.919696

Weeks, A. R., Sgro, C. M., Young, A. G., Frankham, R., Mitchell, N. J., Miller, K. A., Byrne,

M., Coates, D. J., Eldridge, M. D. B., Sunnucks, P., Breed, M. F., James, E. A., &

Hoffmann, A. A. (2011). Assessing the benefits and risks of translocations in

changing environments: A genetic perspective. *Evolutionary Applications*, *4*(6), 709–

725. https://doi.org/10.1111/j.1752-4571.2011.00192.x

Wickham, H., Averick, M., Bryan, J., Chang, W., D'Agostino McGowan, L., Romain, F.,

Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Lin Pedersen, T., Miller,

E., Milton Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., …

Yutani, H. (2019). Welcome to the tidyverse. *Journal of Open Source Software*,

4(43), 1686. https://doi.org/10.21105/joss.01686

Wingfield, J. C., Kelley, J. P., & Angelier, F. (2011). What are extreme environmental

conditions and how do organisms cope with them? *Current Zoology*, *57*(3), 363–374.

https://doi.org/10.1093/czoolo/57.3.363

You, W., & Henneberg, M. (2018). Cancer incidence increasing globally: The role of relaxed

natural selection. *Evolutionary Applications*, *11*(2), 140–152.

https://doi.org/10.1111/eva.12523

Zizzari, Z. V., & Ellers, J. (2011). Effects of exposure to short-term heat stress on male

reproductive fitness in a soil arthropod. *Journal of Insect Physiology*, *57*(3), 421–426.

https://doi.org/10.1016/j.jinsphys.2011.01.002

Supplementary information

Figure S1 – Breeding protocol experimental design

Figure S2 – Carcass photography protocol (not to scale). Two Canon DSLRs at a 90° angle to each

other were used to photograph each carcass from 30 cm away against a white background. The carcass

was placed on a clear plastic breeding individual box lid (12 x 8 x 1 cm) to avoid the white background

getting contaminated with soil (the lid was wiped between carcasses to maintain a white background

for the top image).

764
765
766 *Script S1 – Carcass roundness script for calculating roundness in ImageJ. The script uses the blue*

767 *channels from both the top and side photos (as this creates the greatest contrast between the carcass*

768 *nest and the background), a threshold filter to separate the carcass from the background and a*

769 *median filter to ignore the smallest black flecks of less than 25 pixels, in case any small soil marks*

770 *remained.*

771
```
772  //  topPath=File.openDialog("Select top-down image"); // get file location
773   //  sidePath=File.openDialog("Select side image"); // get file location
774  Dialog.create("Settings");
775  Dialog.addMessage("Pixel/distance calibration:");
776  Dialog.addNumber("Top-down (pixels per mm)", 53.7523);
777  Dialog.addNumber("Side (pixels per mm)", 30.1806);
778  Dialog.addMessage("\nImage processing:");
779  Dialog.addNumber("Threshold", 70);
780  Dialog.addNumber("Smoothing radius (pixels)", 25);
781  Dialog.addMessage("\nCamera positioning:");
782  Dialog.addCheckbox("Flip image", true);
783  Dialog.show();
784  topPixelsMm = Dialog.getNumber();
785  sidePixelsMm = Dialog.getNumber();; thresholdVal = Dialog.getNumber();;; smoothingVal=
786  Dialog.getNumber();;;; flip = Dialog.getCheckbox();
787  // FILE LOCATIONS
```

```
788    topDir=getDirectory("Directory containing top-down images");
789    topFileList=getFileList(topDir);
790    sideDir=getDirectory("Directory containing side-on images");
791    sideFileList=getFileList(sideDir);
792    // select working directory // list of images in directory
793    // select working directory // list of images in directory
794    if(topFileList.length > sideFileList.length) // error if there are unequal numbers of files
795    exit("Error: unequal number of files in selected folders");
796    if(topFileList.length < sideFileList.length)
797    exit("Error: unequal number of files in selected folders");
798    print("\n_________________________________");
799    print("SETTINGS");
800    print(" Threshold: " + thresholdVal);
801    print(" Smoothing Radius: " + smoothingVal);
802    // START OF BATCH PROCESSING LOOP
803    for(z=0; z<topFileList.length; z++1){ topPath = topDir+topFileList[z];
804    sidePath = sideDir+sideFileList[z];
805    // OPEN & PROCESS TOP IMAGE
806    open(topPath);
807    run("RGB Stack");
808    run("Delete Slice"); // only uses the blue channel (channel with highest contrast)
809    run("Delete Slice");
810    setTool(0); //Rectangle tool
811    waitForUser("Select the Target", "Draw a box to cover the mouse ball\nensure no other dark
812    objects are selected");
813    run("Crop");
814    if(flip==1); // if ticked, flips image so that both are the same way round
815    run("Flip Horizontally");
816    setThreshold(0, thresholdVal);
817    run("Convert to Mask");
818    medianScript = "radius="+smoothingVal+" slice";
819    run("Median...", medianScript); // threshold
820    w = getWidth(); // image dimensions
821    h = getHeight();
822    // SELECT CENTRE
823    doWand(w/2, h/2, 1, "4-connected");
824    List.setMeasurements; // save measurements of perimeter & area
825    topPerim = List.getValue("Perim.");
826    topArea = List.getValue("Area");
827    print("Top Perimeter: " + topPerim + " Top Area: " + topArea);
828    run("Make Inverse"); // invert selection & delete (gets rid of dirt spots etc..)
829    setForegroundColor(255, 255, 255);
830    run("Fill", "slice");
831    run("Make Inverse");
832    run("To Bounding Box");
833    run("Crop");
834    w = getWidth(); // image dimensions
835    h = getHeight();
836    topProfile = newArray(w);
837    for (x=0; x<w; x++){ // pixel column loop
```

```
838    topProfile[x] = 0;
839    for (y=0; y<h; y++){ // counts the number of white pixels in each column
840    } }
841    val = getPixel(x,y);
842    if(val==255)
843    topProfile[x] = topProfile[x] + 1;
844    topProfileCrop = newArray(0);
845    for(a=0; a<topProfile.length; a++){ if(topProfile[a]>0){
846    topProfileCrop = Array.concat(topProfileCrop, topProfile[a]);
847    } }
848    topProfileMm = newArray(topProfileCrop.length);
849    for(a=0; a<topProfileCrop.length; a++){
850    topProfileMm[a] = topProfileCrop[a] / topPixelsMm;
851    setResult("Top Pixels", a, topProfileCrop[a]);
852    setResult("Top Millimetres", a, topProfileMm[a]); }
853    close();
854    // OPEN & PROCESS SIDE IMAGE
855    // make array of millimetre values
856    open(sidePath);
857    run("RGB Stack");
858    run("Delete Slice"); // only uses the blue channel (channel with highest contrast)
859    run("Delete Slice");
860    setTool(0); //Rectangle tool
861    waitForUser("Select the Target", "Draw a box to cover the mouse ball\nensure no other dark
862    objects are selected");
863    run("Crop");
864    setThreshold(0, thresholdVal);
865    run("Convert to Mask");
866    medianScript = "radius="+smoothingVal+" slice";
867    run("Median...", medianScript); // threshold
868    w = getWidth(); // image dimensions
869    h = getHeight();
870    // SELECT CENTRE
871    doWand(w/2, h/2, 1, "4-connected");
872    List.setMeasurements; // save measurements of perimeter & area
873    sidePerim = List.getValue("Perim.");
874    sideArea = List.getValue("Area");
875    print("Side Perimeter: " + sidePerim + " Side Area: " + sideArea);
876    run("Make Inverse"); // invert selection & delete (gets rid of dirt spots etc..)
877    setForegroundColor(255, 255, 255);
878    run("Fill", "slice");
879    run("Make Inverse");
880    run("To Bounding Box");
881    run("Crop");
882    w = getWidth(); // image dimensions
883    h = getHeight();
884    sideProfile = newArray(w);
885    for (x=0; x<w; x++){ // pixel column loop
886    sideProfile[x] = 0;
887    for (y=0; y<h; y++){ // counts the number of white pixels in each column
```

```
888   } }
889   val = getPixel(x,y);
890   if(val==255)
891   sideProfile[x] = sideProfile[x] + 1;
892   sideProfileCrop = newArray(0);
893   for(a=0; a<sideProfile.length; a++){ if(sideProfile[a]>0){
894   sideProfileCrop = Array.concat(sideProfileCrop, sideProfile[a]);
895   } }
896   sideProfileMm = newArray(sideProfileCrop.length);
897   for(a=0; a<sideProfileCrop.length; a++){
898   sideProfileMm[a] = sideProfileCrop[a] / sidePixelsMm;
899   setResult("Side Pixels", a, sideProfileCrop[a]);
900   setResult("Side Millimetres", a, sideProfileMm[a]); }
901   // make array of millimetre values
902   close();
903   // CIRCUMFRENCE CALCULATION
904   arrayRatios = topProfileMm.length / sideProfileMm.length;
905   circumfrenceArray = newArray(sideProfileMm.length);
906   areaArray = newArray(sideProfileMm.length);
907   for(i=0; i<sideProfileMm.length; i++){
908   bLoc = round(arrayRatios*i);
909   b = topProfileMm[bLoc]/2;
910   a = sideProfileMm[i]/2;
911   circumfrenceArray[i] = PI * (a+b) * (1+(3*pow((a-b)/(a+b),2) / (10+pow(4-(3*pow((a-
912   b)/(a+b),2)),0.5)))); 
913   areaArray[i] = PI * a * b;
914   }
915   topLength = topProfileMm.length / topPixelsMm;
916   sideLength = sideProfileMm.length / sidePixelsMm;
917   aveLength = (topLength + sideLength)/2;
918   mmPerSlice = aveLength/sideProfileMm.length; // average length of mouse
919   VolumeSum = 0;
920   AreaSum = 0;
921   for(a=0; a<sideProfileCrop.length; a++){ // loop to count the area and volume of each slice
922   AreaSum = AreaSum + (circumfrenceArray[a]*mmPerSlice);
923   VolumeSum = VolumeSum + (areaArray[a]*mmPerSlice);
924   // setResult("Circumfrence", a, circumfrenceArray[a]);
925   }
926   EndArea = 0;
927   for(a=0; a<sideProfileCrop.length-1; a++)
928   EndArea = EndArea + (pow(pow(areaArray[a]-areaArray[a+1],2),0.5)); // sums end surface
929   values - always positive
930   AreaSum = AreaSum + EndArea;
931   // SPHERICITY CALCULATION - this takes the average value of the area to circumference
932   ratio of the top & side images
933   topCircleArea = PI*pow((topPerim/(2*PI)),2); // this calculates the area a circle would have of
934   the observed perimeter length
935   sideCircleArea = PI*pow((sidePerim/(2*PI)),2);
936   topRoundness = topArea / topCircleArea; // perfect circle area /observed area, so a perfect
937   circle would be 1, going down to zero with worse roundness
```

```
938    sideRoundness = sideArea / sideCircleArea;
939    sphericity = (topRoundness+sideRoundness)/2;
940    // PRINT RESULTS
941    print("\n...........................................................");
942    print("Top-down image: " + topFileList[z]);
943    print("Side-on image: " + sideFileList[z]);
944    print("Sphericity:\t"+ sphericity);
945    print("Volume (mm^3):\t"+VolumeSum);
946    } //end batch processing loop
947    // average roundness
948
949
```

Figure S3 – The process of removing any dark background marks and identity labels, while ensuring that the size and proportion of the carcass ball remained the same, by digitally superimposing white rectangles over any dark objects in Apple Preview. Script S1 removes black flecks of less than 25 pixels but any larger dark shapes, such as clumps of soil, identity labels and unfurled tails, can influence the carcass roundness calculations so these were digitally removed using this process.

Figure S4 – An example of the variation in roundness between carcasses. The left image has a roundness value of 0.2957 and the right image has a value of 0.6031. In Script S1 both the top and side images were used to calculate roundness.

Figure S5 - The effect of experimental block on lytic activity of sexually mature beetles' anal exudate. Lytic activity is defined as the percentage of cell degradation of Micrococcus lysodeikticus *as recorded at 450 nm after 60 minutes. Higher remaining percentages of* M. lysodeikticus *remaining indicate lower exudate lytic activity. Each point represents one beetle (purple = females, black = males), thick*

969   *horizontal lines represent the median for each treatment, whiskers extend to the farthest data point*

970   *which is no more than 1.5 times the interquartile range from the box. N = 75 males and 75 females.*

974   *Figure S6 – The effect of incubator temperature during sexual maturation on clutch size for those pairs*

975   *that laid at least one egg. Each point represents a different clutch, the line represents the fit for the*

976   *linear model, and the shaded areas represent the 95 % confidence intervals. N = 304 clutches.*

Figure S7 – The effect of temperature during sexual maturation and evolutionary population on the number of larvae hatched (brood size). No eggs successfully hatched from beetles kept at 28 °C, so this graph only shows data from 22 °C and 25 °C. Each datapoint represents one clutch, lines represent the reaction norms for evolutionary lines (red = Full Care, blue = No Care) and the shaded areas represent the 95 % confidence intervals. N = 383 broods.