Can recent evolutionary history promote resilience to environmental

change?

Lay summary

Can we use knowledge of a population's evolutionary history to predict how individuals

might cope with environmental change? We investigated whether burying beetles that had

evolved for 45 generations with or without post-hatching parental care differed in their

resilience to extreme temperatures as they developed to sexual maturity. We found limited

evidence that experimental evolution under different regimes of parental care contributed to

thermal resilience, which instead was better explained by chance historical events.

Abstract

Principles of social evolution have long been used retrospectively to interpret social

interactions, but have less commonly been applied predictively to inform conservation and

animal husbandry strategies. We investigate whether differences in developmental

environment, facilitated by divergent social conditions, can predict resilience to

environmental change. Upon exposure to harsh novel environments, populations that

previously experienced more benign social environments are predicted either to suffer fitness

losses (the "mutation load hypothesis" and "selection filter hypothesis") or maintain fitness

(the "beneficial mutation hypothesis"). We tested these contrasting predictions using

populations of burying beetles *Nicrophorus vespilloides* we had evolved experimentally for

45 generations under contrasting social environments by manipulating the supply of post-

hatching parental care. We exposed sexually immature adults from each population to

varying heat stress and measured the effect on survival and reproduction. The greater the

level of parental care previously experienced by a population, the better its survival under

heat stress during sexual maturation. Although this is consistent with the "beneficial mutation

hypothesis", it is also possible that populations that had evolved without post-hatching care

were simply more prone to dying during maturation, regardless of their thermal environment.

Overall, we suggest that stochastic genetic variation, probably due to founder effects, had a

stronger influence on resilience. We discuss the implications for translocation and captive

breeding programmes.

Introduction

In a world changing rapidly through the effects of climate change (Pörtner et al., 2022),

habitat degradation (Millenium Ecosystem Assessment, 2005) and excessive use of

antibiotics (Graham et al., 2019) and pesticides (Hawkins et al., 2019), our ability to predict

how effectively organisms will adapt to new environments is more important than ever. The

principles of social evolution, and how they influence developmental environment, have long

been used retrospectively to understand social interactions (Bourke, 2011). However they

have less commonly been used to inform the development of practical strategies to address

global biological challenges (Carroll et al., 2014). Yet an understanding of a population's

social evolutionary history, by which we mean the type of selective social environment it has

experienced, could predict its resilience to future environmental perturbation, through the

effects that different social environments have on mutation load and selection for particular

alleles (Lumley et al., 2015). This has practical value in its potential to address several major

economic and societal problems, including food security (pest/crop interactions, crop

development), human and animal medicine (antibiotic development, anti-parasite

interventions) and wild animal conservation (choice of populations for translocation schemes,

identification of populations vulnerable to disease and environmental change) (Carroll et al.,

2014).

In theory, historical effects on resilience to the current environment depend on whether the

past developmental environment experienced by the population is benign or harsh - because

this influences the extent of standing genetic variation available. Few studies have tested

experimentally how past exposure to different social environments during development

affects a population's future response to environmental change (but see Lumley et al., 2015).

Benign social interactions, which are usually due to some form of cooperative interaction, are

known to buffer individuals from environmental stress and therefore to reduce the extent to

which mildly deleterious alleles are exposed to natural selection (Linksvayer & Wade, 2009;

Mashoodh et al., 2023; Pascoal et al., 2022). Elevated mutation loads resulting from relaxed

selection have been shown to reduce reproductive success in animals kept in captive breeding

programmes (Araki et al., 2007; McPhee, 2004; Robert, 2009) and have been linked to a

higher prevalence of genetic disorders in human societies where selection is buffered by

widely available healthcare (Lynch, 2016; Saniotis et al., 2021; You & Henneberg, 2018).

The "mutation load hypothesis" therefore predicts that individuals previously exposed to

more benign developmental environments, facilitated by the buffering effect of social

environments which impose fewer selective pressures, should be less resilient to new

environmental stresses. However, whilst most of the mutations that accumulate under relaxed

selection are likely to be (mildly) harmful, some could potentially be beneficial and might

even facilitate greater resilience to environmental change at the population level. The

"beneficial mutation hypothesis" therefore conversely predicts that individuals that develop

in more benign social environments could be more resilient to new environmental stresses, if

by chance they have accumulated cryptic genetic variation that is beneficial when expressed

in these new environments (Barrett & Schluter, 2008; Paaby & Rockman, 2014). At the other

end of the spectrum, a harsh social environment potentially imposes a strong selection filter,

which is likely to reduce the population's mutation load, causing individuals generally to be

fitter (Balmford, 1996). The "selection filter hypothesis" predicts that a history of exposure to

harsher social environments means that populations are more likely to be resilient to

environmental change.

Species in which offspring are facultatively dependent on parental care (Clutton-Brock,

1991) offer a way to test these contrasting predictions. Offspring that receive care are

exposed to a benign social environment during development which, if sustained over many

generations relaxes selection (Snell-Rood et al., 2016). Conversely, offspring that receive no

care over many generations experience a harsher environment during development and

consequently experience stronger selection against deleterious mutations. Our experiments

focus on the burying beetle *Nicrophorus vespilloides*, which exhibits elaborate biparental

care upon which offspring are facultatively dependent (Capodeanu-Nägler et al., 2016). We

know from previous work that offspring exposed to post-hatching care experience a more

benign developmental environment than those that develop without post-hatching care. For

example, Schrader et al. (2017) found that when breeding pairs of beetles were removed

before providing post-hatching care, fewer of them produced at least one surviving larva, and

those larvae that did survive were lighter. We exploited this natural history to test whether,

and how, past social interactions influence future resilience to environmental change.

Burying beetle adults locate the carcasses of small vertebrates using their highly sensitive

antennae (Kalinová et al., 2009; Trumbo & Steiger, 2020) and then fight intrasexually for

ownership (Otronen, 1988). The winning pair then tears the fur or feathers from the carcass,

covers the flesh with antimicrobial oral and anal exudates, rolls it into a ball and buries it

underground (Cotter & Kilner, 2010). The female then lays eggs in the soil surrounding the

carcass nest. When the larvae hatch, 2-3 days later, they crawl to the nest (Müller & Eggert,

1989), where their parents defend them from predators and intra- and interspecific

competitors and feed them trophallactically (Eggert et al., 1998). The duration of post-

hatching care is highly variable and in some cases is non-existent due to the death or

desertion by parents after preparation of the carcass nest (Schrader, Jarrett, et al., 2015a;

Smiseth et al., 2003). In such cases larvae can self-feed (Smiseth et al., 2003) and can survive

(at least in the laboratory) without any post-hatching care.

We evolved two replicate populations of burying beetles in the laboratory under contrasting

regimes of parental care for 45 generations. In the Full Care populations, beetles were

allowed to stay with their offspring throughout development. Individuals in these populations

experienced a more benign environment during development and relaxed selection (Pascoal

et al., 2022). By contrast, in the No Care populations, parents were removed just prior to

larval hatching and so were unable to supply any post-hatching care at all. Therefore

individuals in these populations experienced a harsher social environment and stronger

selection against deleterious mutations (Pascoal et al., 2022).

After 45 generations of experimental evolution, we tested whether there were differences in

the resilience of beetles from these two types of experimental population when they were

exposed to higher temperatures during their juvenile adult stage. We define resilience as the

ability of a population to maintain normal processes, in particular reproductive success, in

response to a harsh novel environment (Lowerre-Barbieri et al., 2017; Martin & Wiebe,

2004; Wingfield et al., 2011). Temperature was chosen as the environmental stressor because

of its ecological relevance in the context of climate change. Insects, being poikilothermic, are

particularly vulnerable to heat stress, and research on burying beetles specifically has shown

that they have reduced breeding success at higher temperatures. We chose to expose the

beetles to the temperature treatments during sexual maturation alone because this is the time

when an adult beetle's gonads are developing (e.g. David et al., 2005; Gibbs et al., 2010;

Sales et al., 2018; Zizzari and Ellers, 2011). Furthermore, Sales et al. (2021) show in flour

beetles *Tribolium castaneum* that this developmental stage is particularly vulnerable to

effects of thermal stress on offspring production. Additionally, the natural history of the

burying beetle will likely render this a particularly sensitive period as sexually immature

individuals cannot stay underground to buffer themselves from heat stress because they must

emerge to feed.

We exposed sexually immature adults to different degrees of heat stress and measured the

effect on the following correlates of fitness: survival to sexual maturity, anal exudate lytic

activity, carcass sphericity, fecundity, and hatching success Duarte et al., 2021). We

predicted that the contrasting regimes of selection imposed by these different social

environments during development for the 44 preceding generations should yield a contrasting

response to heat stress during sexual maturation. The "mutation load hypothesis" and

"selection filter hypothesis" each predict that individuals derived from the No Care

populations should be more resilient to heat stress, whereas the "beneficial mutation

hypothesis" predicts that individuals derived from the No Care populations should be less

resilient to heat stress.

Methods

Burying beetle husbandry in the laboratory: general methods

Burying beetles were bred by placing them in a plastic breeding box (17 x 12 x 6 cm),

partially filled with damp soil (Miracle-Gro Compost), furnished with a 10-15 g mouse

carcass for breeding. Boxes were placed in a cupboard after pairing to simulate underground

conditions. Larvae were counted and weighed 8 days after pairing and each one placed in its

own cell (2 x 2 x 2 cm) within a plastic pupation box (10 x 10 x 2 cm), filled with damp peat.

Sexually immature adults were eclosed around 21 days later and each was placed in its own

individual box (12 x 8 x 2 cm). Adults were fed twice weekly with beef mince until breeding,

which took place 15 days post-eclosion. At all life stages, individuals were kept at 21 °C.

During breeding, pairs were kept in a constantly dark environment, but for all other life

stages they were kept on a 16L: 8D hour light cycle.

Experimental Evolution

The *N. vespilloides* populations used in this study were part of a long-term experimental

evolution project that investigated how populations of burying beetles adapt to the loss of

parental care. There were four experimental populations in all: Full Care (FC) (replicated

twice) and No Care (NC) (replicated twice). These populations were established from a stock

population founded with wild-caught beetles trapped under permit in 2014 from four

woodland sites across Cambridgeshire, UK (Byron's Pool, Gamlingay Woods, Waresley

Woods and Overhall Grove). From this stock population 32 pairs were bred, with the male

and female coming from different sites to minimise the likelihood of inbreeding. The

resulting 671 offspring established the experimental evolution populations. FC populations

were left undisturbed (as described in general methods above) and allowed to stay with their

offspring until dispersal. However, in the NC populations, at 53 h post-pairing, when the

carrion nest was complete but before the larvae had hatched, both parents were removed to

prevent post-hatching care. This procedure was repeated at every generation. The replicate

experimental populations were bred in two separate blocks (FC1/NC1 and FC2/NC2), and the

timing of the lifecycle was staggered between blocks to be 7 days apart. As all the blocks

were derived from the same stock population, FC1 and NC1 were no more related to each

other than FC2 and NC2.

Testing for resilience to heat stress

Experimental generation 1: Partitioning current care environment from evolutionary

history of care

Beetles were taken for analysis at generation 45 of experimental evolution. We partitioned

the effects of current care experienced by individuals (No Care or Full Care), from an

evolutionary history of care experienced by the source population (No Care or Full Care), by

breeding beetles for one generation using the following treatments:

1) Adults from Full Care populations allowed to provide full parental care (Full Care

parents with a Full Care regime) = 73 FC_{FC} pairs

2) Adults from Full Care populations removed at 53 hours after pairing (Full Care

parents with a No Care regime) = 74 FC_{NC} pairs

3) Adults from No Care populations removed at 53 hours after pairing (No Care parents

with a No Care regime) = 74 NC_{NC} pairs

4) Adults from No Care populations allowed to provide full parental care (No Care

parents with a Full Care regime) = 74 NC_{FC} pairs

When larval development was complete, and larvae were starting to disperse away from the

remains of the carrion nest (8 days after pairing), larvae were put in pupation boxes to mature

and reach eclosion, as described above.

Experimental generation 2: Exposure to heat stress

Immediately after eclosion, each experimental individual was placed alone in a small plastic

box (12 x 8 x 6 cm) containing a fine layer of soil. The boxes were placed in incubators

(Panasonic MLR-352H-PE) set to either 22 °C, 25 °C or 28 °C on a 16L: 8D hour light cycle.

The maximum temperature was chosen based on a previous, unpublished pilot experiment in

the laboratory that found a severe reduction in hatching success of beetles reared at 28 °C,

and the 25 °C temperature was chosen to represent an intermediate temperature between one

that was likely to cause severe heat stress and one that was close to our standard laboratory

temperature. Our intention was not to mimic natural thermal conditions closely, but rather

create different levels of heat stress across the treatments within a naturally occurring range.

The boxes were removed from the incubators briefly once after a week so that the beetles

could be fed with beef mince. We then measured the following correlates of fitness for this

generation of beetles:

Survival

To measure the effect of heat stress on survival, after 14 days (at sexual maturity) all

individuals were removed from their incubators and classified as alive or dead.

The surviving individuals were then assigned to one of the following treatments for breeding:

1) 22 °C male x 22 °C female (n = 29 FC_{FC}, 29 FC_{NC}, 28 NC_{NC}, 29 NC_{FC})

2) 25 °C male x 25 °C female (n = 36 FC_{FC}, 35 FC_{NC}, 36 NC_{NC}, 36 NC_{FC})

3) 28 °C male x 28 °C female (n = 32 FC_{FC}, 34 FC_{NC}, 29 NC_{NC}, 30 NC_{FC})

Individuals were only bred within their populations and temperatures, e.g. 22 °C males from

FC_{FC} were only bred with 22 °C females from FC_{FC}, resulting in 12 treatments (Figure S1).

Pairs were placed in breeding boxes containing 350 ml of moist soil and a 10-13 g mouse

carcass, and the box was sealed and kept in a dark environment, as described under 'general

methods' above. All pairs were kept at the standard laboratory temperature of 21 °C,

regardless of their temperature treatment, to isolate the effects of the thermal environment

during sexual maturation on subsequent reproductive success.

Lytic activity

Higher exudate lytic activity is heritable and increases larval survival so it is possible that

there has been differential selection for this during the evolution of the experimental lines

(Arce et al., 2012; Cotter & Kilner, 2010). Additionally, previous research has shown that

when mature adults of other *Nicrophorus* species were subjected to a three-day thermal stress

treatment there were significant contrasting effects on the lytic activity of their oral exudates

(Jacques et al., 2009). Therefore, we would expect to see lower levels of lytic activity in

beetles more negatively affected by higher temperatures.

At 53 h after pairing, we collected anal exudates from parents from a subsample of broods

from each treatment (n = 27 FC_{FC}, 27 FC_{NC}, 22 NC_{NC}, 19 NC_{FC}) by holding each beetle by the

abdomen and pressing a 5 ul glass microcapillary tube (Drummond) to its anus. Each sample

was pipetted immediately into a 1.5 ml Eppendorf tube on dry ice and subsequently stored at

-80 °C until analysis. We quantified the lytic activity of each sample in an automated

microplate reader (BioTek ELx808) by running turbidity assays that measured the

degradation of bacterial cell walls (see Duarte et al., 2021 for full details). Briefly, 1ul of

each exudate sample was diluted in 24 ul of 6.4 pH 0.02 M potassium phosphate buffer and

10ul of this solution was added to a well of a 96-well flat-bottomed microplate (Thermo

Scientific Nunc MicroWell) containing 100 ul of a 1.3 mg ml^{-1} solution of lyophilized

Micrococcus lysodeikticus cells in potassium phosphate buffer. The microplate reader was set

to 25 °C and took an initial reading at 450 nm. Plates then went through a cycle of being

shaken at a medium setting every 10 minutes, followed immediately by a 450 nm reading, for

60 minutes.

Carcass sphericity

Rounder carcasses are less hospitable to parasitism by blowflies (Sun & Kilner, 2020), and

are negatively correlated with parental lifespan, indicating that preparing a rounded carcass

requires significant resources from the parents (De Gasperin et al., 2016). Therefore, we

would expect that if higher temperatures have a negative effect on beetles then they would

make a less spherical carcass.

At 53 h, we also briefly removed the carcass from a subsample of broods (n = 33 FC_{FC}, 31

FC_{NC}, 33 NC_{NC}, 33 NC_{FC}) where the parents had experienced the 22 °C and 28 °C degree

treatments. Each carcass was placed on a white background, alongside a scale for
measurement, and photographed from perpendicular angles using two cameras (Canon
DSLR, with 18-55mm lens) at 30 cm away from the carcass (Figure S2).

Laying failure, clutch size, hatching failure and brood size
At 53 h after pairing, we removed the parents from all 383 broods and counted the number of
eggs laid, through inspection of the underside of the box (following Jarrett et al., 2017;
Schrader et al., 2015c). At 70, 80, 94 and 104 h after pairing we checked the carcass and the
surface of the soil for any first instar larvae, and counted them before discarding them.

Statistical analyses

All statistical tests were conducted in R version 3.6.1 (R Core Team, 2022) and a 0.05
significance threshold was used throughout. Data handling and visualisation were carried out
using base R and the 'tidyverse' suite of R packages (Wickham et al., 2019). Model selection
for the analyses of brood mass and brood size was conducted using the 'dredge' and
'model.avg' functions in the 'MuMIn' R package (Bartoń, 2023). The most parsimonious
model within two AIC_C points of the model with the lowest AIC_C was chosen as the optimal
model. Goodness of fit of optimal models was assessed with the 'DHARMa' package
(Hartig, 2022) or, in the case of beta regression models, the base 'stats' package. Continuous
independent variables in all models were scaled and centred using the 'scale' function in base
R.

Survival

We ran a binomial GLM with survival status as the dependent variable ("0" = died in the

incubator, "1" = survived to sexual maturity), and population evolutionary history (Full Care

or No Care), generation 1 care regime (Full Care or No Care), temperature, sex, block, and

the two-way interactions of these terms, and three-way interactions between population

evolutionary history, block and temperature, and generation 1 care regime and history, block

and temperature as the independent variables in the maximal model. See Table S1 for an

explanation of why each interaction term was included in the models.

Lytic activity

In some of the broods only one of the parents produced enough exudate to allow us to

conduct the lytic activity analysis on their sample. The total useable sample size for exudates

was 150 individuals, and for 110 of these we had a sufficient sample from their partner (55

pairs) and in the other 40 only one partner in the pair could be analysed (we could not collect

enough exudate from the partners of 20 females and 20 males used in the analysis).

We used the percentage change between the 450 nm absorption at the first reading (0

minutes) and at 60 minutes as the dependent variable (as per Duarte et al., 2021). We first ran

a paired t-test to determine whether there was a consistent sex difference within pairs in lytic

activity. As the lytic activity of one individual in a pair may influence its partner's levels, for

males and females we then ran separate beta regression, using the 'betareg' R package

(Cribari-Neto & Zeileis, 2010), with proportion of cells degraded (450 nm reading at 60

minutes/450 nm reading at 0 minutes) in either males' or females' anal exudate as the

dependent variable, and population evolutionary history (Full Care or No Care), generation 1

care regime (Full Care or No Care), temperature, partner's lytic activity, block, the two-way

interactions between temperature and evolutionary history, and sex and temperature, and the

three-way interaction between population evolutionary history, block and temperature, and

generation 1 care regime, block and temperature as independent variables in the maximal

model.

Carcass sphericity

Carcass sphericity was calculated digitally in ImageJ (Schneider et al., 2012), using a macro

script adapted from De Gasperin et al (2016) (Script S1 and Figure S3). The ImageJ script

assumes that if a carcass were perfectly spherical, the two-dimensional images in the carcass

from the top and side would be exact circles. The script compares the actual images to a

perfect circle of the same perimeter and then calculates the difference between the two – such

that a perfectly spherical carcass would have a score of 1 to denote that it fills 100% of the

space of a perfect sphere (Figure S4).

As carcass sphericity was bounded between 0 and 1, we ran a beta regression with carcass

sphericity percentage as the dependent variable and population evolutionary history (Full

Care or No Care), generation 1 care regime (Full Care or No Care), temperature, sex, block,

and the two-way interactions of these terms, and three-way interactions between population

evolutionary history, block and temperature, and generation 1 care regime and history, block

and temperature as the independent variables in the maximal model.

Laying failure, clutch size, hatching failure and brood size

To investigate predictors of fecundity, we first ran a binomial GLM with laying failure ("0" =

for a brood where no eggs were laid and "1" = for a brood where at least one egg was laid) as

the dependent variable and population evolutionary history (Full Care or No Care),

generation 1 care regime (Full Care or No Care), temperature, carcass mass, block, two-way

interactions between population evolutionary history and generation 1 care, population

evolutionary history and temperature, generation 1 care and temperature, population

evolutionary history and block, generation 1 care and block, and block and temperature, and

three-way interactions between population evolutionary history, block and temperature, and

generation 1 care, block and temperature as independent variables.

Then, to examine predictors of clutch size in the broods that did lay eggs, we ran a quasi-

Poisson GLM with clutch size (the number of eggs counted from the underside of the

breeding box) as the dependent variable, and the same independent variables that were used

in the model analysing laying failure vs success.

We then ran two tests to investigate predictors of offspring production. Firstly, we ran a

binomial GLM with brood success ("0" = for a brood with no larvae, "1" = for a brood with

at least one hatching larva) as the dependent variable and the same independent variables that

were used in the binomial model analysing laying failure vs success. Next, for broods where

at least one egg hatched, we ran a quasi-Poisson GLM with brood size (number of larvae

counted on the carcass or soil) as the dependent variable and the same independent variables

that were used in the model analysing laying failure vs success, plus clutch size. Broods were

included in the hatching success model even if no eggs were counted from the underside of

the breeding box, in case by chance a female had laid very few eggs more shallowly in the

soil which then successfully hatched.

Results

Survival

The optimal model retained population evolutionary history, temperature, block (evolutionary

population replicate) and the interaction between temperature and block (Table 1 and S2).

Beetles from populations with an evolutionary history of No Care were more likely to die in

the incubators, regardless of the temperature they experienced in the incubator or the care

they received in generation 1 of the heat stress experiment. Mean survival of Full Care

individuals across blocks was 94%, but 87% for No Care individuals.

To further investigate how block affected survival at higher temperatures we split the dataset

by block and ran generalised linear models with survival probability as the dependent

variable and temperature as the independent variable. In block 1 the impact of temperature

was non-significant ($\chi^2_{1,\ N=618} = 1.820$, $p = 0.177$), but in block 2 there was a significant

negative effect ($\chi^2_{1,\ N=600} = 125.63$, $p = <0.001$). This indicates that beetles exposed to greater

heat stress as they matured sexually were more likely to die, but only in block 2.

Table 1 – Results of binomial model investigating the effects on likelihood of a beetle surviving from
eclosion to sexual maturity in incubators set to 22 °C, 25 °C or 28 °C. Terms retained in the optimal
model based on AIC_C are presented.

Independent variable	Estimate	SE	z	p
Intercept	4.170	0.301	13.857	<0.001
Population evolutionary history (No Care)	-1.008	0.231	-4.362	<0.001

Temperature	-0.335	0.255	-1.315	0.189
Block (2)	-1.041	0.341	-3.055	0.002
Block:temperature	-1.490	0.343	-4.339	<0.001

Figure 1 – The survival of beetles kept in incubators at different levels of heat stress throughout sexual maturation (1 = survived, 0 = died) in relation to A) population evolutionary history and experimental block and B) generation 1 care regime. Points represent data for individual pairs (jittered on the x- and y-axes), lines represent the reaction norms for each treatment and the shaded areas indicate the 95 % confidence intervals. N = 1,281 individuals.

Lytic activity

There was no difference between males and females in lytic activity strength: $t_{54} = 0.446$, $p =$

0.658. However, to take partner lytic activity into account, we still ran separate models for

males and females. Block was retained in the optimal models of both male and female lytic

activity, with individuals having stronger lytic activity in block 2 (Figure 2, Tables 2 and S3).

In the female model, temperature was also retained, with females having lower lytic activity

in their exudates when they developed at higher temperatures. In the male model, generation

1 care regime was also retained, with males that received no post-hatching care having

exudates with lower lytic activity. However, to test the influence of an outlier we re-ran the

analysis with the removal of one datapoint where <55% of *Micrococcus* remained, and the

generation 1 care term was no longer retained, indicating that the effect of generation 1 care

regime was very small (Figure S5, Tables S4 & S5).

Figure 2 – The effect of developmental temperature on male (A & B) and female (C & D) anal exudates in relation to temperature and population evolutionary history and experimental block (A & C) and temperature and generation 1 care regime (B & D). Lytic activity is defined as the percentage of cell degradation of Micrococcus lysodeikticus *as recorded at 450 nm after 60 minutes. Higher remaining percentages of* M. lysodeikticus *remaining indicate lower exudate lytic activity. Each point represents one beetle (jittered on the x- and y-axes), lines represent the reaction norms for each treatment and the shaded areas indicate the 95 % confidence intervals. N = 75 males and 75 females.*

Table 2 – Results of beta regressions investigating the effects on A) male and B) female exudate lytic activity at 53h after pairing. Terms retained in the optimal model based on AIC$_C$ are presented. We have defined lytic activity as the proportion of cell degradation of Micrococcus lysodeikticus *as recorded at 450 nm after 60 minutes.*

Independent variable	Estimate	SE	z	*p*
A) Male exudate				

Intercept	3.972	0.275	14.433	<0.001
Block (2)	-1.368	0.149	-9.160	<0.001
Generation 1 care	0.227	0.125	1.811	0.070
B) Female exudate				
Intercept	3.989	0.299	13.329	<0.001
Block (2)	-1.323	0.160	-8.259	<0.001
Temperature	0.161	0.064	2.522	0.012

Carcass sphericity

442 Generation 1 care regime and temperature were retained in the optimal model. Beetles that

443 received full care prepared rounder carcasses, as did beetles that developed at 22 °C,

444 compared to 28 °C (Figure 3, Tables 3 & S6).

447
Figure 3 – The roundness of carcasses at 53 h after pairing in relation to temperature and A)

population evolutionary history and experimental block and B) generation 1 care regime. A

proportion of 1 would represent a perfectly spherical carcass nest. Points represent data for

individual pairs (jittered on the x- and y-axes), lines represent the reaction norms for each treatment

and the shaded areas indicate the 95 % confidence intervals.

Table 3 – Results of beta regression investigating variation in carcass nest roundness as a proxy for
carcass preparation success. Terms retained in the optimal model based on AIC_C are presented.

Independent variable	Estimate	SE	z	p
Intercept	-0.011	0.044	-0.242	0.809
Generation 1 care	-0.134	0.063	-2.145	0.032
Temperature	-0.252	0.032	-8.009	<0.001

Laying failure and clutch size

Exposure to higher temperatures during sexual maturation was more likely to cause complete

failure to lay, regardless of evolutionary history, and there was a greater chance of complete

laying failure in block 2 (Tables 4A & S7, Figure 4). Neither the population's evolutionary

history of care (Full Care or No Care), nor the care regime experienced in the first generation

of the heat stress experiment, were retained in the optimal model of clutch success vs failure

or clutch size (Tables 4B & S8). In broods where some laying occurred, higher temperatures

led to fewer eggs laid per clutch, particularly in Block 1 (Figure 5).

Table 4 – Results of GLMs looking at variables affecting A) failure to lay: no eggs produced versus at
least one egg laid and B) clutch size for pairs that laid at least one egg. Terms retained in the optimal
model based on AIC_C are presented.

Independent variable	Estimate	SE	z	p
A) Failure to lay				
Intercept	4.488	0.537	8.360	<0.001
Block (2)	-2.082	0.422	-4.933	<0.001
Temperature	-3.287	0.411	-7.994	<0.001

B) Clutch size			
Intercept	2.926	0.034	87.291	<0.001
Block (2)	0.101	0.052	1.929	0.055
Temperature	-0.350	0.033	-10.568	<0.001
Block:temperature	0.131	0.055	2.365	0.019

475
476 *Figure 4 – The laying status (1 = at least 1 egg laid, 0 = no eggs laid) of pairs of beetles kept in*
477 *incubators at different levels of heat stress throughout sexual maturation in relation to A) population*
478 *evolutionary history and experimental block and B) generation 1 care regime. Points represent data*
479 *for individual pairs (jittered on the x- and y-axes), lines represent the reaction norms for each*
480 *treatment and the shaded areas indicate the 95 % confidence intervals.*

483
Figure 5 – The clutch size of pairs of beetles that were kept in incubators at different levels of heat

stress throughout sexual maturation in relation to A) population evolutionary history and

experimental block and B) generation 1 care regime. Only pairs that laid at least one egg are

presented. Points represent data for individual pairs (jittered on the x-axis), lines represent the

reaction norms for each treatment and the shaded areas indicate the 95 % confidence intervals.

Hatching failure and brood size

Neither parental population evolutionary history, nor generation 1 care regime, were retained

in the optimal model of hatching success vs failure (Table S9). Parents kept at higher

incubator temperatures between eclosion and sexual maturity were more likely to experience

complete hatching failure (Figure 6, Tables 5A & S9) – with no eggs hatching in any of the

broods where the parents had developed at 28 °C.

Fewer larvae hatched when their parents had been kept at 25 °C than 22 °C (Figure 7), and

more larvae hatched when there were more eggs to start with (Tables 5B & S10).

Additionally, for pairs where at least one larva hatched, the relationship between temperature

and brood size was different between blocks, with block 1 pairs having more hatching larvae

at 22 °C and block 2 pairs having more larvae at 25 °C (Figure 7B).

Table 5 – Results of GLMs looking at variables affecting A) hatching success (at least one larva) or

failure (no larvae) and B) brood size for pairs where at least one larva hatched. Terms retained in the

optimal model based on AIC_C are presented.

Independent variable	Estimate	SE	z	p
A) Failure to hatch				
Intercept	3.091	0.396	7.817	<0.001
Temperature	-2.741	0.349	-7.855	<0.001

B) Brood size			
Intercept	1.878	0.130	14.405	<0.001
Block (2)	0.289	0.071	4.093	<0.001
Temperature	-0.544	-0.044	-12.416	<0.001
Clutch size	0.029	0.005	6.188	<0.001

509
510

511
512 *Figure 6 – The hatching status (1 = at least 1 hatched larva, 0 = no larvae) of clutches from pairs of*
513 *beetles kept in incubators at different levels of heat stress throughout sexual maturation in relation to*
514 *A) population evolutionary history and experimental block and B) generation 1 care regime. Points*
515 *represent data for individual pairs (jittered on the x- and y-axes), lines represent the reaction norms*
516 *for each treatment and the shaded areas indicate the 95 % confidence intervals.*

519
Figure 7 – The brood size of pairs of beetles that were kept in incubators at different levels of heat
stress throughout sexual maturation in relation to A) experimental evolution population, B)

experimental block, and C) generation 1 care regime. Only pairs with at least one hatching larva are

presented. Points represent data for individual pairs (jittered on the x- and y-axes), lines represent

the reaction norms for each treatment and the shaded areas indicate the 95 % confidence intervals.

Discussion

We tested experimentally whether a population's past evolutionary exposure to contrasting

levels of post-hatching care predicted its capacity to cope with heat stress during sexual

maturation. The key findings of our experiments are summarised in Table 6, for each of the

correlates of fitness we measured.

Table 6 – The key predictors tested, their effects on correlates of fitness measured, and the hypothesis supported by the findings. "None" indicates that the

predictor variable did not have a significant effect on the correlate of fitness. Full statistical results are presented in Tables 1-5.

Predictor variables tested	Effect on correlates of fitness							Hypothesis supported
	Survival	Lytic activity	Carcass sphericity	Failure to lay	Clutch size	Hatching failure	Brood size	
Population evolutionary history (No Care or Full Care)	Lower survival in No Care	None	None	None	None	None	None	Beneficial mutation
Generation 1 care regime (No Care or Full Care)	None	In males, lower lytic activity when they	Lower sphericity when they received no post-hatching	None	None	None	None	N/A

		received no post-hatching care as larvae	care as larvae						
Temperature	Lower survival at higher temperature in block 2	In females, lower lytic activity at higher temperature	Lower sphericity at higher temperature	Higher failure at higher temperature	Smaller clutch size at higher temperature, particularly in block 1	Higher hatching failure at higher temperature	Smaller brood size at higher temperature	N/A	
Block	Lower survival at higher	Stronger lytic activity	None	Higher failure in block 2	Smaller clutch size at higher	None	Larger broods in block 2	Founder effect	

	temperature in block 2	in block 2			temperature, particularly in block 1			

We found that populations that had evolved under Full Care experienced better survival

during sexual maturation across all temperatures. This is consistent with the "beneficial

mutation hypothesis". Nevertheless the effect size was small, which could be due to exposure

to relatively few generations of divergent selection and/or relatively little divergence in

selection pressure since both No Care and Full Care populations developed in the relatively

benign conditions of the lab.

Individuals derived from the No Care evolving populations had poor survival in the

incubators, even in the most benign thermal environment, compared with those from the Full

Care populations. It has previously been shown that individuals in the No Care populations

carry a lower mutation load (Pascoal et al. 2022). Perhaps this means that, by chance, they

lack the cryptic genetic variation that enables them to cope with new levels of thermal stress

during sexual maturation (Paaby & Rockman, 2014). Alternatively, perhaps the No Care

populations were intrinsically inferior by this point, and more prone to dying between

eclosion and sexual maturity, regardless of the temperature to which they were exposed. The

lack of a significant effect of the interaction between population evolutionary history and

temperature on incubator survival is consistent with this alternative interpretation. However,

in all other correlates of fitness we detected no differences between individuals drawn from

the No Care versus Full Care populations, which suggests that the individuals from the No

Care populations were not intrinsically inferior in each of their traits. Overall, our

experimental data allow us to reject the "mutation load hypothesis" and the "selection filter

hypothesis". However, we have insufficient data to conclude we have found strong support

for the "beneficial mutation hypothesis".

Unexpectedly, we found that experimental block predicted some correlates of fitness, which

could be due to a stochastic founder effect. The four distinct evolving populations were

established from a single large stock population derived by interbreeding beetles from four

different wild populations. The size of the breeding population was identical for each block

throughout the experiment (n = 100 pairs), as was the maintenance regime, which implies

that any differences between blocks must be attributable to chance differences in genetic

variation when the population was founded. For example, chance differences in genetic

variation could explain chance differences in the persistence of haplotypes within

populations, since there were no environmental differences that could otherwise explain

variation in reproductive success. In analyses of earlier generations of the experimentally

evolving populations, the experimental social environment (Full Care versus No Care)

explained divergence between populations to a greater extent than did block (e.g. Duarte et

al., 2021; Jarrett et al., 2018a, 2018b; Schrader et al., 2015c, 2015a, 2015b, 2017, 2017). By

generation 46, however, the situation had reversed: block outweighed experimental social

environment in explaining the response to heat stress exposure during sexual maturation.

Founder effects have long been suggested to explain the success of translocation and captive

breeding in conservation programmes (e.g. Brown et al., 2009; Dolman et al., 2021; Miller et

al., 1999). However, in analyses of wild populations it is often hard to disentangle the scale

of founder effects from confounding factors of environmental differences (however subtle) in

the new location (Kolbe et al., 2012; Szűcs et al., 2017) especially when it is difficult to

monitor populations systematically after their release into the wild (Armstrong & Seddon,

2008; Mock et al., 2004; Seddon et al., 2007; Weeks et al., 2011). Nevertheless, there is some

evidence that founder effects influence the response to selection in wild populations. When

Anolis sagrei individuals with long hindlimbs were translocated to seven Bahamian islands

with a vegetation structure known to favour short hindlimbed individuals, they evolved

shorter hindlimbs over four years. However the extent of morphological change was

attributable to founder effects (Kolbe et al., 2012). Perhaps a similar pattern explains the

results we have found in this study. More broadly our study indicates the importance of the

initial population size in captive breeding programmes. It also reinforces the wisdom of using

genetic screening to determine whether the founding population is representative of the

source population and has high heterozygosity (Bragg et al., 2021; Scott et al., 2020),

especially in cases where there will be no migration into the population, such as on an

isolated island. Our findings are of particular interest from a practical perspective considering

that many conservation captive breeding programmes have, by necessity, used smaller

founder populations than ours (Witzenberger & Hochkirch, 2011).

However, for some of the traits we measured (failure to lay, clutch size, failure to hatch) we

found that temperature had such a strong negative effect that it outweighed any effects that

could be attributed to the population's genetic background. This is consistent with previous

work across different species of burying beetle, which found that high temperatures during

breeding led to reduced clutch and brood sizes and a greater likelihood of complete brood

failure (Keller et al., 2021; Moss & Moore, 2021; Ney, 2021; Ong, 2019; Quinby et al., 2020;

Sun & Kilner, 2020). In all of these studies the extreme temperature treatment continued

throughout breeding, whereas in this study we elevated temperature only during sexual

maturation to better simulate the natural history of the burying beetle. Our results suggest that

there are sensitive windows in development which affect breeding success even when

reproduction occurs under benign conditions. In a world with an increasing frequency and

intensity of heat waves (Perkins-Kirkpatrick & Lewis, 2020), this study shows that an

extreme temperature event can reduce breeding success even if it is only brief, as long as it

coincides with these sensitive windows in development.

There are two likely causes for a reduction in egg laying and egg hatching at higher

temperatures: 1) behavioural – beetles maturing at higher temperatures mated less, and/or 2)

physiological – beetles maturing at higher temperatures had underdeveloped gonads.

Particularly in males, exposure to high temperatures prior to pairing leads to reduced

courtship behaviour in *Aphidius avenae* (Roux et al., 2010) and *Drosophila melanogaster,*

Drosophila simulans and Drosophila mojavensis (Patton & Krebs, 2001). There is also a

great deal of evidence to suggest that thermal stress during sexual maturation can have a

physiological effect on insects. In *Drosophila melanogaster* high temperatures during

development cause a range of spermatid cytological abnormalities (David et al., 2005;

Rohmer et al., 2004) and in *Tribolium castaneum* cause lower testis volume (Sales et al.,

2021). In females, higher temperatures during development cause smaller oocytes in *Pararge*

aegeria (Gibbs et al., 2010) and cause *Bicyclus anynana* to lay more, smaller eggs (Steigenga

& Fischer, 2007). From this study we are unable to say whether the effects were behavioural,

physiological or both, and whether it was the males or females that drove the patterns of

reduced reproductive fitness at higher temperatures. Further investigations tracking

copulatory behaviour, mating thermally stressed males with control females and vice versa,

and dissecting ovaries and testes following heat stress are needed to answer these questions.

It is widely accepted that a population's standing genetic variation will predict its response to

new environmental challenges. Our study suggests it is possible that the evolutionary history

of burying beetle populations might predict the response to heat stress during sexual

maturation (because No Care populations had lower survival at all temperatures in the

incubators), although further work is needed before a robust conclusion can be drawn.

However, stochastic genetic variation, due to founder effects or genetic drift, was potentially

a more powerful predictor of population resilience upon exposure to new thermal

environments. Translocation and captive breeding programmes should be designed

accordingly to maximise their chance of success.

References

Araki, H., Cooper, B., & Blouin, M. S. (2007). Genetic Effects of Captive Breeding Cause a

Rapid, Cumulative Fitness Decline in the Wild. *Science*, *318*(5847), 100.

https://doi.org/10.1126/science.1145621

Arce, A. N., Johnston, P. R., Smiseth, P. T., & Rozen, D. E. (2012). Mechanisms and fitness

effects of antibacterial defences in a carrion beetle. *Journal of Evolutionary Biology*,

25(5), 930–937. https://doi.org/10.1111/j.1420-9101.2012.02486.x

Armstrong, D. P., & Seddon, P. J. (2008). Directions in reintroduction biology. *Trends in

Ecology & Evolution*, *23*(1), 20–25. https://doi.org/10.1016/j.tree.2007.10.003

Balmford, A. (1996). Extinction filters and current resilience: The significance of past

selection pressures for conservation biology. *Trends in Ecology & Evolution*, *11*(5),

193–196. https://doi.org/10.1016/0169-5347(96)10026-4

Barrett, R. D. H., & Schluter, D. (2008). Adaptation from standing genetic variation. *Trends

in Ecology & Evolution*, *23*(1), 38–44. https://doi.org/10.1016/j.tree.2007.09.008

Bartoń, K. (2023). *MuMIn: Multi-model inference* (1.47.5) [R]. https://CRAN.R-

project.org/package=MuMIn

Bourke, A. F. G. (2011). *Principles of Social Evolution*. Oxford University Press.

https://doi.org/10.1093/acprof:oso/9780199231157.001.0001

Bragg, J. G., Yap, J.-Y. S., Wilson, T., Lee, E., & Rossetto, M. (2021). Conserving the

genetic diversity of condemned populations: Optimizing collections and translocation.

Evolutionary Applications, *14*(5), 1225–1238. https://doi.org/10.1111/eva.13192

Brown, S. K., Hull, J. M., Updike, D. R., Fain, S. R., & Ernest, H. B. (2009). Black Bear

Population Genetics in California: Signatures of Population Structure, Competitive

Release, and Historical Translocation. *Journal of Mammalogy*, *90*(5), 1066–1074.

https://doi.org/10.1644/08-MAMM-A-193.1

Capodeanu-Nägler, A., Keppner, E. M., Vogel, H., Ayasse, M., Eggert, A.-K., Sakaluk, S.

K., & Steiger, S. (2016). From facultative to obligatory parental care: Interspecific

variation in offspring dependency on post-hatching care in burying beetles. *Scientific*

Reports, *6*(1), 29323. https://doi.org/10.1038/srep29323

Carroll, S. P., Jørgensen, P. S., Kinnison, M. T., Bergstrom, C. T., Denison, R. F., Gluckman,

P., Smith, T. B., Strauss, S. Y., & Tabashnik, B. E. (2014). Applying evolutionary

biology to address global challenges. *Science*, *346*(6207).

https://doi.org/10.1126/science.1245993

Clutton-Brock, T. H. (1991). *The Evolution of Parental Care* (Vol. 64). Princeton University

Press; JSTOR. https://doi.org/10.2307/j.ctvs32ssj

Cotter, S., & Kilner, R. M. (2010). Sexual division of antibacterial resource defence in

breeding burying beetles, *Nicrophorus vespilloides*. *Journal of Animal Ecology*,

79(1), 35–43. https://doi.org/10.1111/j.1365-2656.2009.01593.x

Cribari-Neto, F., & Zeileis, A. (2010). Beta Regression in R. *Journal of Statistical Software*,

34(2), 1–24. https://doi.org/doi:10.18637/jss.v034.i02

David, J. R., Araripe, L. O., Chakir, M., Legout, H., Lemos, B., Pétavy, G., Rohmer, C., Joly,

D., & Moreteau, B. (2005). Male sterility at extreme temperatures: A significant but

neglected phenomenon for understanding Drosophila climatic adaptations. *Journal of*

Evolutionary Biology, *18*(4), 838–846. https://doi.org/10.1111/j.1420-

9101.2005.00914.x

De Gasperin, O., Duarte, A., Troscianko, J., & Kilner, R. M. (2016). Fitness costs associated

with building and maintaining the burying beetle's carrion nest. *Scientific Reports*, *6*,

35293.

Dolman, P. M., Burnside, R. J., Scotland, K. M., & Collar, N. J. (2021). Captive breeding and

the conservation of the threatened houbara bustards. *Endangered Species Research*,

46, 161–173.

Duarte, A., Rebar, D., Hallett, A. C., Jarrett, B. J. M., & Kilner, R. M. (2021). Evolutionary

change in the construction of the nursery environment when parents are prevented

from caring for their young directly. *Proceedings of the National Academy of*

Sciences, *118*(48), e2102450118. https://doi.org/10.1073/pnas.2102450118

Eggert, A.-K., Reinking, M., & Müller, J. K. (1998). Parental care improves offspring

survival and growth in burying beetles. *Animal Behaviour*, *55*(1), 97–107.

https://doi.org/10.1006/anbe.1997.0588

Gibbs, M., Van Dyck, H., & Karlsson, B. (2010). Reproductive plasticity, ovarian dynamics

and maternal effects in response to temperature and flight in Pararge aegeria. *Journal*

of Insect Physiology, *56*(9), 1275–1283.

https://doi.org/10.1016/j.jinsphys.2010.04.009

Graham, D. W., Bergeron, G., Bourassa, M. W., Dickson, J., Gomes, F., Howe, A., Kahn, L.

H., Morley, P. S., Scott, H. M., Simjee, S., Singer, R. S., Smith, T. C., Storrs, C., &

Wittum, T. E. (2019). Complexities in understanding antimicrobial resistance across

domesticated animal, human, and environmental systems. *Annals of the New York*

Academy of Sciences, *1441*(1), 17–30. https://doi.org/10.1111/nyas.14036

Hartig, F. (2022). *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level/Mixed)*

Regression Models (0.4.6) [R]. https://CRAN.R-project.org/package=DHARMa

Hawkins, N. J., Bass, C., Dixon, A., & Neve, P. (2019). The evolutionary origins of pesticide

resistance. *Biological Reviews*, *94*(1), 135–155. https://doi.org/10.1111/brv.12440

Jacques, B. J., Akahane, S., Abe, M., Middleton, W., Hoback, W. W., & Shaffer, J. J. (2009).

Temperature and Food Availability Differentially Affect the Production of

Antimicrobial Compounds in Oral Secretions Produced by Two Species of Burying

Beetle. *Journal of Chemical Ecology*, *35*(8), 871–877.

https://doi.org/10.1007/s10886-009-9680-y

Jarrett, B. J. M., Evans, E., Haynes, H. B., Leaf, M. R., Rebar, D., Duarte, A., Schrader, M.,

& Kilner, R. M. (2018). A sustained change in the supply of parental care causes

adaptive evolution of offspring morphology. *Nature Communications*, *9*(1), 3987.

https://doi.org/10.1038/s41467-018-06513-6

Jarrett, B. J. M., Rebar, D., Haynes, H. B., Leaf, M. R., Halliwell, C., Kemp, R., & Kilner, R.

M. (2018). Adaptive evolution of synchronous egg-hatching in compensation for the

loss of parental care. *Proceedings of the Royal Society B: Biological Sciences*,

285(1885), 20181452. https://doi.org/10.1098/rspb.2018.1452

Jarrett, B. J. M., Schrader, M., Rebar, D., Houslay, T. M., & Kilner, R. M. (2017).

Cooperative interactions within the family enhance the capacity for evolutionary

change in body size. *Nature Ecology & Evolution*, *1*, 0178.

http://dx.doi.org/10.1038/s41559-017-0

Kalinová, B., Podskalská, H., Růžička, J., & Hoskovec, M. (2009). Irresistible bouquet of

death—How are burying beetles (Coleoptera: Silphidae: *Nicrophorus*) attracted by

735 carcasses. *Naturwissenschaften*, *96*(8), 889–899. https://doi.org/10.1007/s00114-009-

736 0545-6

737 Keller, M. L., Howard, D. R., & Hall, C. L. (2021). The thermal ecology of burying beetles:

738 Temperature influences reproduction and daily activity in Nicrophorus marginatus.

739 *Ecological Entomology*, *46*(6), 1266–1272. https://doi.org/10.1111/een.13071

740 Kolbe, J. J., Leal, M., Schoener, T. W., Spiller, D. A., & Losos, J. B. (2012). Founder Effects

741 Persist Despite Adaptive Differentiation: A Field Experiment with Lizards. *Science*,

742 *335*(6072), 1086–1089. https://doi.org/10.1126/science.1209566

743 Linksvayer, T. A., & Wade, M. J. (2009). Genes with social effects are expected to harbor

744 more sequence variation within and between species. *Evolution; International*

745 *Journal of Organic Evolution*, *63*(7), 1685–1696. PubMed.

746 https://doi.org/10.1111/j.1558-5646.2009.00670.x

747 Lowerre-Barbieri, S., DeCelles, G., Pepin, P., Catalán, I. A., Muhling, B., Erisman, B.,

748 Cadrin, S. X., Alós, J., Ospina-Alvarez, A., Stachura, M. M., Tringali, M. D.,

749 Burnsed, S. W., & Paris, C. B. (2017). Reproductive resilience: A paradigm shift in

750 understanding spawner-recruit systems in exploited marine fish. *Fish and Fisheries*,

751 *18*(2), 285–312. https://doi.org/10.1111/faf.12180

752 Lumley, A. J., Michalczyk, Ł., Kitson, J. J. N., Spurgin, L. G., Morrison, C. A., Godwin, J.

753 L., Dickinson, M. E., Martin, O. Y., Emerson, B. C., Chapman, T., & Gage, M. J. G.

754 (2015). Sexual selection protects against extinction. *Nature*, *522*, 470.

755 Lynch, M. (2016). Mutation and Human Exceptionalism: Our Future Genetic Load. *Genetics*,

756 *202*(3), 869–875. https://doi.org/10.1534/genetics.115.180471

757 Martin, K., & Wiebe, K. L. (2004). Coping Mechanisms of Alpine and Arctic Breeding

758 Birds: Extreme Weather and Limitations to Reproductive Resilience1. *Integrative and*

759 *Comparative Biology*, *44*(2), 177–185. https://doi.org/10.1093/icb/44.2.177

Mashoodh, R., Trowsdale, A., Manica, A., & Kilner, R. (2023). Parental care drives the

evolution of molecular genetic variation. *bioRxiv*, 2023.01.09.523216.

https://doi.org/10.1101/2023.01.09.523216

McPhee, M. (2004). Generations in captivity increases behavioral variance: Considerations

for captive breeding and reintroduction programs. *Biological Conservation*, *115*(1),

71–77. https://doi.org/10.1016/S0006-3207(03)00095-8

Millenium Ecosystem Assessment. (2005). *Ecosystems and Human Well-Being: Synthesis*.

World Resources Institute.

https://www.millenniumassessment.org/documents/document.356.aspx.pdf

Miller, B., Ralls, K., Reading, R. P., Scott, J. M., & Estes, J. (1999). Biological and technical

considerations of carnivore translocation: A review. *Animal Conservation*, *2*(1), 59–

68. Cambridge Core. https://doi.org/10.1111/j.1469-1795.1999.tb00049.x

Mock, K. E., Latch, E. K., & Rhodes, O. E. (2004). Assessing losses of genetic diversity due

to translocation: Long-term case histories in Merriam's turkey (Meleagris gallopavo

merriami). *Conservation Genetics*, *5*(5), 631–645. https://doi.org/10.1007/s10592-

004-1849-x

Moss, J. B., & Moore, A. J. (2021). Constrained flexibility of parental cooperation limits

adaptive responses to harsh conditions. *Evolution*, *75*(7), 1835–1849.

https://doi.org/10.1111/evo.14285

Müller, J. K., & Eggert, A.-K. (1989). Paternity assurance by "helpful" males: Adaptations to

sperm competition in burying beetles. *Behavioral Ecology and Sociobiology*, *24*(4),

245–249. https://doi.org/10.1007/BF00295204

Ney, L. (2021). *Impact of temperature stress on reproduction, offspring development, and*

social immunity in the burying beetle Nicrophorus orbicollis [University of Nebraska

at Omaha]. https://www.proquest.com/docview/2585301523?pq-

origsite=gscholar&fromopenview=true

Ong, J. Y. (2019). Coping with Thermal Challenges: Reaction Norms of Life History Traits

in a Burying Beetle with Biparental Care [M.S., Queen's University (Canada)]. In

ProQuest Dissertations and Theses (2186240977). ProQuest Dissertations & Theses

A&I. https://ezp.lib.cam.ac.uk/login?url=https://www.proquest.com/dissertations-

theses/coping-with-thermal-challenges-reaction-norms/docview/2186240977/se-

2?accountid=9851

Otronen, M. (1988). The effect of body size on the outcome of fights in burying beetles

(*Nicrophorus*). *Annales Zoologici Fennici*, *25*(2), 191–201.

Paaby, A. B., & Rockman, M. V. (2014). Cryptic genetic variation: Evolution's hidden

substrate. *Nature Reviews Genetics*, *15*, 247.

Pascoal, S., Mashoodh, R., Shimadzu, H., & Kilner, R. M. (2022). Parental care allows

genetic variation to accumulate by relaxing selection. *bioRxiv*, 2022.05.05.490718.

https://doi.org/10.1101/2022.05.05.490718

Patton, Z. J., & Krebs, R. A. (2001). The Effect of Thermal Stress on the Mating Behavior of

Three Drosophila Species. *Physiological and Biochemical Zoology*, *74*(6), 783–788.

https://doi.org/10.1086/323327

Perkins-Kirkpatrick, S. E., & Lewis, S. C. (2020). Increasing trends in regional heatwaves.

Nature Communications, *11*(1), 3357. https://doi.org/10.1038/s41467-020-16970-7

Pörtner, H. O., Roberts, D. C., Tignor, M., Poloczanska, E. S., Mintenbeck, K., Alegria, A.,

Craig, M., Langsdorf, S., Löschke, S., Möller, V., Okem, A., & Rama, B. (2022).

IPCC, 2022: Climate Change 2022: Impacts, Adaptation, and Vulnerability.

Contribution of Working Group II to the Sixth Assessment Report of the

Intergovernmental Panel on Climate Change. Cambridge University Press.

Quinby, B. M., Belk, M. C., & Creighton, J. C. (2020). Behavioral constraints on local

adaptation and counter-gradient variation: Implications for climate change. *Ecology*

and Evolution, *10*(13), 6688–6701. https://doi.org/10.1002/ece3.6399

R Core Team. (2022). *R: A language and environment for statistical computing* [Computer

software]. R Foundation for Statistical Computing. https://www.r-project.org

Robert, A. (2009). Captive breeding genetics and reintroduction success. *Biological*

Conservation, *142*(12), 2915–2922. https://doi.org/10.1016/j.biocon.2009.07.016

Rohmer, C., David, J. R., Moreteau, B., & Joly, D. (2004). Heat induced male sterility in

Drosophila melanogaster: Adaptive genetic variations among geographic populations

and role of the Y chromosome. *Journal of Experimental Biology*, *207*(16), 2735–

2743. https://doi.org/10.1242/jeb.01087

Roux, O., Le Lann, C., van Alphen, J. J. M., & van Baaren, J. (2010). How does heat shock

affect the life history traits of adults and progeny of the aphid parasitoid Aphidius

avenae (Hymenoptera: Aphidiidae)? *Bulletin of Entomological Research*, *100*(5),

543–549. Cambridge Core. https://doi.org/10.1017/S0007485309990575

Sales, K., Vasudeva, R., Dickinson, M. E., Godwin, J. L., Lumley, A. J., Michalczyk, Ł.,

Hebberecht, L., Thomas, P., Franco, A., & Gage, M. J. G. (2018). Experimental

heatwaves compromise sperm function and cause transgenerational damage in a

model insect. *Nature Communications*, *9*(1), 4771. https://doi.org/10.1038/s41467-

018-07273-z

Sales, K., Vasudeva, R., & Gage, M. J. G. (2021). Fertility and mortality impacts of thermal

stress from experimental heatwaves on different life stages and their recovery in a

model insect. *Royal Society Open Science*, *8*(3). https://doi.org/10.1098/rsos.201717

Saniotis, A., Henneberg, M., & Mohammadi, K. (2021). Genetic load and biological changes

to extant humans. *Journal of Biosocial Science*, *53*(4), 639–642. Cambridge Core.

https://doi.org/10.1017/S0021932020000413

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years

of Image Analysis. *Nature Methods*, *9*(7), 671–675. PMC.

Schrader, M., Cosby, R., Hesketh, A., Jarrett, B. J. M., & Kilner, R. (2015). A limit on the

extent to which increased egg size can compensate for a poor postnatal environment

revealed experimentally in the burying beetle, *Nicrophorus vespilloides*. *Ecology and*

Evolution, *6*(1), 329–336. https://doi.org/10.1002/ece3.1876

Schrader, M., Jarrett, B. J. M., & Kilner, R. M. (2015a). Parental care masks a density-

dependent shift from cooperation to competition among burying beetle larvae.

Evolution; International Journal of Organic Evolution, *69*(4), 1077–1084. PMC.

https://doi.org/10.1111/evo.12615

Schrader, M., Jarrett, B. J. M., & Kilner, R. M. (2015b). Using Experimental Evolution to

Study Adaptations for Life within the Family. *The American Naturalist*, *185*(5), 610–

619. https://doi.org/10.1086/680500

Schrader, M., Jarrett, B. J. M., Rebar, D., & Kilner, R. M. (2017). Adaptation to a novel

family environment involves both apparent and cryptic phenotypic changes.

Proceedings of the Royal Society B: Biological Sciences, *284*(1862), 20171295.

https://doi.org/10.1098/rspb.2017.1295

Scott, P. A., Allison, L. A., Field, K. J., Averill-Murray, R. C., & Shaffer, B. H. (2020).

Individual heterozygosity predicts translocation success in threatened desert tortoises.

Science, *370*(6520), 1086–1089. https://doi.org/10.1126/science.abb0421

Seddon, P. J., Armstrong, D. P., & Maloney, R. F. (2007). Developing the Science of

Reintroduction Biology. *Conservation Biology*, *21*(2), 303–312.

https://doi.org/10.1111/j.1523-1739.2006.00627.x

Smiseth, P. T., Darwell, C. T., & Moore, A. J. (2003). Partial begging: An empirical model

for the early evolution of offspring signalling. *Proceedings of the Royal Society B:*

Biological Sciences, *270*(1526), 1773–1777. https://doi.org/10.1098/rspb.2003.2444

Snell-Rood, E. C., Burger, M., Hutton, Q., & Moczek, A. P. (2016). Effects of parental care

on the accumulation and release of cryptic genetic variation: Review of mechanisms

and a case study of dung beetles. *Evolutionary Ecology*, *30*(2), 251–265.

https://doi.org/10.1007/s10682-015-9813-4

Steigenga, M. J., & Fischer, K. (2007). Ovarian dynamics, egg size, and egg number in

relation to temperature and mating status in a butterfly. *Entomologia Experimentalis*

et Applicata, *125*(2), 195–203. https://doi.org/10.1111/j.1570-7458.2007.00610.x

Sun, S.-J., & Kilner, R. M. (2020). Temperature stress induces mites to help their carrion

beetle hosts by eliminating rival blowflies. *eLife*, *9*, e55649. PubMed.

https://doi.org/10.7554/eLife.55649

Szűcs, M., Melbourne, B. A., Tuff, T., Weiss-Lehman, C., & Hufbauer, R. A. (2017). Genetic

and demographic founder effects have long-term fitness consequences for colonising

populations. *Ecology Letters*, *20*(4), 436–444. https://doi.org/10.1111/ele.12743

Trumbo, S. T., & Steiger, S. (2020). Finding A Fresh Carcass: Bacterially-Derived Volatiles

And Burying Beetle Search Success. *bioRxiv*, 2020.01.25.919696.

https://doi.org/10.1101/2020.01.25.919696

Weeks, A. R., Sgro, C. M., Young, A. G., Frankham, R., Mitchell, N. J., Miller, K. A., Byrne,

M., Coates, D. J., Eldridge, M. D. B., Sunnucks, P., Breed, M. F., James, E. A., &

Hoffmann, A. A. (2011). Assessing the benefits and risks of translocations in

changing environments: A genetic perspective. *Evolutionary Applications*, *4*(6), 709–

725. https://doi.org/10.1111/j.1752-4571.2011.00192.x

Wickham, H., Averick, M., Bryan, J., Chang, W., D'Agostino McGowan, L., Romain, F.,

Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Lin Pedersen, T., Miller,

E., Milton Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., …

Yutani, H. (2019). Welcome to the tidyverse. *Journal of Open Source Software*,

4(43), 1686. https://doi.org/10.21105/joss.01686

Wingfield, J. C., Kelley, J. P., & Angelier, F. (2011). What are extreme environmental

conditions and how do organisms cope with them? *Current Zoology*, *57*(3), 363–374.

https://doi.org/10.1093/czoolo/57.3.363

Witzenberger, K. A., & Hochkirch, A. (2011). Ex situ conservation genetics: A review of

molecular studies on the genetic consequences of captive breeding programmes for

endangered animal species. *Biodiversity and Conservation*, *20*(9), 1843–1861.

https://doi.org/10.1007/s10531-011-0074-4

You, W., & Henneberg, M. (2018). Cancer incidence increasing globally: The role of relaxed

natural selection. *Evolutionary Applications*, *11*(2), 140–152.

https://doi.org/10.1111/eva.12523

Zizzari, Z. V., & Ellers, J. (2011). Effects of exposure to short-term heat stress on male

reproductive fitness in a soil arthropod. *Journal of Insect Physiology*, *57*(3), 421–426.

https://doi.org/10.1016/j.jinsphys.2011.01.002

900