**No evidence that urbanisation encourages wild red foxes (*Vulpes vulpes*) to exploit**

**unfamiliar food-related objects when first discovered**

F. Blake Morton*[1], Marieke Gartner[2], Ellie-Mae Norrie[1], Yacob Haddou[3], Kristy Adaway[1]

[1] Department of Psychology, University of Hull, Hull, UK

[2] Atlanta Zoo, USA

[3] Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow,

Glasgow, UK

**Author Note**

F. Blake Morton 🆔 https://orcid.org/0000-0002-8644-1557

Marieke Gartner 🆔 https://orcid.org/0000-0003-3901-9166

Yacob Haddou 🆔 https://orcid.org/0000-0003-0557-0339

The authors would like to thank everyone involved with the 'British Carnivore Project', particularly Sophie Tait, Josh Chatterton, Alice Turner, Eszter Jardan, Louise Grunnill, and all of the citizen scientists who kindly gave us permission to collect data in their gardens. We are very grateful to the many organisations that helped us acquire permission to access all o fthe other land for the study, particularly staff from the Yorkshire Wildlife Trust, Lincolnshire Wildlife Trust, Scottish Wildlife Trust, The Land Trust, the National Trust, Forestry England (particularly Cath Bashforth), Forestry & Land Scotland, Yorkshire Water, and the city and regional councils (particularly Hull, East Riding of Yorkshire, and Glasgow). Special thanks go to Prof. Hannah Buchanan-Smith (Uni Stirling) and Bruce Gittings (Uni Edinburgh) for their help with various logistics in Scotland, to Dr Alex Weiss (Uni Edinburgh) for statistics advice, and to Dr Carl Soulsbury (Uni Lincoln), Prof. Phyllis Lee (Uni Stirling), and Dr Kevin Parson (Uni Glasgow) for helpful comments and discussions on an earlier draft of the manuscript. Finally, we are grateful to the University of Hull and EU Social Fund Plus for funding.

Correspondence about this article should be directed to Dr Blake Morton, Department of Psychology, University of Hull, HU6 7RX, United Kingdom. E-mail: b.morton@hull.ac.uk.

**Conflict of interest statement:** The authors declare no conflict of interest.

48                                              **Abstract**

Urbanisation is the fastest form of landscape transformation on the planet, but our

understanding of how wildlife responds to such changes is still in its infancy. Red foxes (*Vulpes*

*vulpes*) are renowned for thriving in urban settings where they have many opportunities to exploit

food within or in close proximity to a wide variety of human-made objects (e.g., litter and contents

of outdoor bins). The visibility of such behaviour has likely contributed to urban foxes being

labelled as 'bolder', 'cunning', and a general nuisance to people. Studies in other animals show

that frequent exposure to novel objects can sometimes lead them to overcome their aversion to

unfamiliar objects in general (e.g., stimulus generalisation). However, it is unclear whether this

applies to the willingness of urban fox populations to exploit food-related objects, particularly

when they are first discovered and hence the most novel. To test whether urbanisation encourages

wild foxes to exploit food-related objects, we administered eight types of novel objects containing

food to 87 free-ranging foxes across a large urban-rural gradient (one object per location). One fox

approached and exploited an object in the first instance, while the remaining 86 foxes avoided

them. After two weeks, five foxes exploited the objects, while another 13 foxes touched the objects

but quit before gaining access. Urban variables (e.g., road, greenspace, and human population

density) did not predict whether foxes acknowledged or touched the objects, but their likelihood

of touching them was predicted by the type of novel object encountered. When objects were absent,

foxes willingly consumed the food. Thus, as with rural populations, urban foxes in the current

study displayed an initial aversion to novel objects, suggesting that increased access to food-related

objects through urbanisation does not necessarily encourage all foxes to exploit such resources

when first discovered. Other factors are needed to overcome their initial aversion, which may have

important implications for changing the general perception and management of this species.

**Key words:** innovation, boldness, neophobia, urban wildlife, human-wildlife conflict, behavioural

adaptability, public attitudes, dangerous niche hypothesis

74                                              **Introduction**

75              Urbanisation is currently the fastest form of landscape transformation on the planet (Angel

et al., 2011). Many factors likely contribute to whether or how wildlife can adapt to such

environments (e.g., dispersal, morphology, and dietary generalism) (Thompson et al., 2021). In

terms of opportunistic behaviour, a growing number of studies show that particular behavioural

characteristics (e.g., boldness, neophilia, exploration, persistence, and innovation) can increase

urban animals' willingness to spontaneously or quickly exploit novel resources when first

discovered (Dammhahn et al., 2020; Ducatez et al., 2017; Griffin et al., 2017; Mazza et al., 2021;

Mazza & Guenther, 2021). Such findings suggest that being able and willing to rapidly exploit

novel ecological opportunities, rather than avoid or ignore them, may help at least some species

thrive within urban settings. To date, however, not all studies find differences in opportunistic

behaviour between urban and rural animals for reasons that are still unclear (Griffin et al., 2017;

Vincze & Kovacs, 2022), making it difficult to predict whether (or how) such behaviour is

impacted by urbanisation within and across species. On the one hand, trying to exploit novel

resources whenever they arise may help animals improve their fitness by allowing them to acquire

additional resources before they are taken by competitors (Webster et al., 2009). On the other hand,

the benefits of such behaviour must also be weighed against other factors, such as animals'

opportunities and need for such behaviour (Lee, 1991; Lee & Moura, 2015), as well as the

likelihood of their actions exposing them to risk, such as predators, traps, toxic food, and direct

hostile encounters with people (Schell et al., 2021; Soulsbury & White, 2015).

In the current study, we investigated the relationship between urbanisation and the

willingness of wild red foxes (*Vulpes vulpes*) to exploit novel objects containing food. Red foxes

are one of the most widely distributed carnivorans in the world and are renowned for thriving

within urban landscapes, particularly British cities within the last century (Collenette, 1937; Scott

et al., 2014; Soulsbury et al., 2010; Teagle, 1967; Wilkinson & Smith, 2001). As a generalist

carnivore, urban foxes willingly scavenge anthropogenic food within or in close proximity to a

wide variety of human-made objects, such as discarded litter and the contents of household bins

(Contesse et al., 2004; Doncaster et al., 1990). Since most people live within cities, urban foxes

have greater opportunities to exploit these anthropogenic resources compared to rural populations.

The visibility of such behaviour has likely contributed to urban foxes being labelled as 'bold',

'cunning', and a general nuisance to people (Brand & Baldwin, 2020; Bridge & Harris, 2020).

The dangerous niche hypothesis proposes that the primary benefit of having an aversion to

novelty is protection from danger (Greenberg, 2003), which may be particularly relevant to foxes

because they have a long history of conflict with people (e.g., hunting and pest control) (Soulsbury

& White, 2015). Learning from past experiences can play a crucial role in the willingness of

animals to overcome their aversion to novelty (Greenberg & Mettke-Hofmann, 2001). In terms of

novel objects, studies on a range of animals, including birds, fish, and mammals, have found that

frequent exposure to novel objects can sometimes lead to a reduction in their aversion to unfamiliar

objects in general, which may arise through a combination of different psychological processes

(e.g., habituation, associative learning, object recognition, and stimulus generalisation) (Anderson

et al., 2013; Christensen et al., 2011; Gatto et al., 2022; Meehan & Mench, 2002). To date, it is

unclear whether this applies to urban foxes' willingness to exploit novel food-related objects. For

example, urban foxes maintain natural diets and also exploit food readily found within peoples'

gardens (Contesse et al., 2004; Harris, 1981). Thus, resident urban foxes have access to resources

that are likely more abundant, familiar, and easily accessible to them, which may obviate any need

for exploiting unfamiliar human-made objects.

120             Previous studies have administered novel object tests to urban foxes (Padovani et al.,

2021), but these objects did not contain food which makes it difficult to evaluate opportunistic

behaviour within a foraging context, and data from rural populations are also needed for

comparison. Studies have also found that urban foxes can behave more opportunistically than less

urbanised populations in certain contexts (e.g., unfamiliar bait) (Gil-Fernandez et al., 2020), but

responses to one form of novelty do not necessarily reflect how animals might react to other forms

of novelty, including objects (Miller et al., 2022). Thus, in the current study, we administered novel

food-related objects to urban and rural foxes to determine whether their degree of urbanisation

predicted their willingness to exploit such resources when first discovered.

129                                     **Methods and materials**

130     **Ethical statement**

131             This study was ethically approved by the Animal Welfare Ethics Board of the University

132     of Hull (FHS356), and was carried out in accordance with guidelines outlined by the Association

133     for the Study of Animal Behaviour (ASAB, 2020). No foxes were handled.

134     **Study sites and subjects**

135             We studied 181 locations throughout Scotland and England (Figure 1), including a wide

variety of landscapes in and around different cities (e.g., Glasgow, Edinburgh, Stirling, Leeds,

Hull, Lincoln, Sheffield, and York), including recreational parks, private gardens, tree plantations,

meadows, mixed woodland, coastal and mountainous scrubland, and farmland. Foxes were

unmarked and their participation in the study was entirely voluntary. We gained access to 152 of

these locations by contacting city councils and other organisations that owned land. The remaining

29 locations were private gardens, which we accessed by advertising the study through regional

wildlife groups. Our criteria for including any location in the study were: 1) landowner permission,

2) accessibility to foxes (e.g., no barriers/fences), 3) ability to place our equipment out of public

view to avoid theft or vandalism, and 4) the location could not be $\leq$3.5km from another study area.

This latter criterion was used to reduce the chances of sampling the same fox across more than one

location because $\geq$3.5km is larger than the typical dispersal distance and home range diameter of

British foxes (Soulsbury et al., 2011b; Trewhella et al., 1988). We did not have prior knowledge

of fox presence before contacting landowners, and we included locations in the study regardless

of whether landowners said foxes lived on or near their land.

Figure 1. Distribution of zones in Scotland and England where novel objects were deployed.

**Novel object testing**

To determine whether urban foxes were more likely than rural populations to try to exploit

novel food-related objects, we administered eight types of novel objects (Figure 2) over a 12-

month period between August 2021 and 2022. Only a single object was administered per location.

Each object contained desirable food items (e.g., dried dog food and chicken) (Table S1). Some of

the designs were based on studies in other species (Morton, 2021; Rossler et al., 2020; Thornton

& Samson, 2012; Visalberghi & Limongelli, 1994). Objects varied in terms of design and materials

to test whether the type of novel object impacted foxes' behavioural responses. Foxes could gain

access to the rewards inside objects using persistence and simple behaviours that they would likely

need to exploit human-made objects in the real world (e.g., using their mouth, nose, and/or paws

to bite, push, pull, or lift materials). Because we designed the objects ourselves, we could guarantee

that the specific combination of materials used to build them would be novel to foxes.

Figure 2. Novel objects administered to foxes. Yellow dashed arrows indicate the direction of each

behaviour needed to retrieve the food rewards inside. Task G was never deployed in the field and

hence not depicted in this figure.

The objects were made from basic household materials (e.g., PVC piping, metal screws,

and wooden rods). To access the rewards in Object D, foxes simply had to push through the

aluminium side of the box. Object C had two levels, each containing food. The lid of Object F was

fixed in place and could only be solved by sliding it either to the left or right. Object H had a

hidden axle to allow 360° rotation. Object I was administered with the stick already inside the

pipe; animals merely had to remove the stick using their mouths, which would indirectly rake the

food out. All objects were anchored to the plastic platform and had holes drilled into them to

facilitate odour cues. Ten pegs were used to anchor the platforms to the ground. To further help

foxes detect the objects, we scattered "free food" approximately $\leq$1m away. To test whether the

free food and rewards inside objects had an impact on fox motivation, we varied the free food and

rewards by quality and quantity across all locations (Table S1); each location contained a reward

and 'free food' condition.

A single object was available at each location for $15.5 \pm 1.7$ days.  Researchers were not

present when foxes visited, and we did not touch or replenish the food to avoid unnecessary

disturbance to the objects. Following APHA guidelines, we cleaned objects with antibacterial soap

and 70% alcohol wipes after retrieving them to prevent possible pathogen transmission. We then

washed and dried them prior to redeployment. Forty objects (28.4%) were sprayed with scent

deodoriser to test whether the scent of objects (e.g., human odour) had a significant effect on fox

behaviour.

Since foxes were free-ranging and their participation in the study was entirely voluntary,

some foxes might have avoided our testing locations. Nevertheless, the goal of this study was to

test foxes' willingness to *exploit* novel objects to access the food, which required them to

physically touch the objects (and hence be detected on camera). We therefore based our analysis

on foxes that were at least able and willing to visit the locations.

**Recording fox activity**

At each location, we horizontally placed a 'no glow' (940 nm) infrared motion-sensor

camera (Apeman H45) approximately 4 metres away on a tree trunk. Cameras had a 120 ° sensing

angle and a triggering distance of 20 metres. Video lengths were set to record for 5 minutes, with

a 5 second trigger delay and a 30 second interval in between each video. Camera lenses were

sprayed with defogger and, where possible, minor amounts of understory vegetation were removed

between the camera and object to ensure optimal visibility.

**Behavioural responses to novel objects when first discovered**

We recorded the initial behavioural responses of foxes to the objects using the very first

video of a fox from each location, i.e., when objects were likely to be the most novel to them. All

behaviours were measured before each fox turned away from the object and left the camera's frame

of view for >15 seconds; this further ensured we captured foxes' initial responses and also reduced

the likelihood of us counting multiple foxes as a single individual if, for example, more than one

fox happened to be visiting a location at the same time. We recorded behavioural responses at any

point during the two weeks in which objects were deployed, which captured a larger window of

opportunity for foxes to engage with the objects if, for example, it took them more than one visit

to realise the objects were there.

A myriad of factors can underpin animals' responses to novelty, which are not necessarily

underpinned by any one variable (Lee & Moura, 2015; F. B. Morton et al., 2021; Reader & Laland,

2003). However, our goal was to determine *whether* (not why) subjects were successful at

exploiting novel objects with food when discovered. Thus, during each video, we recorded whether

or not (1, 0) the fox acknowledged the novel object, which we defined in terms of the fox turning

its head to look/smell in the object's direction. We also recorded whether or not (1, 0) the fox

touched the object, for instance, by pushing, pulling, and/or biting it, and whether or not (1, 0) they

ultimately gained access to the rewards inside. Finally, during their initial visits to objects, we

coded whether or not (1, 0) the fox appeared interested but too hesitant to approach the object,

which we defined in terms of the fox stopping short of the object, with legs anchored firmly to the

ground while they stretched their body out to look/smell the object; these behaviours could be

supplemented by the fox occasionally jumping/moving away from the object with their ears

flattened and/or the fox scanning their surroundings while visually inspecting the object.

**Food tests**

Foxes are generalist carnivores and should be highly motivated to consume the food

rewards in our study (Saunders & Harris, 2000). To confirm this, we revisited 30 of our locations

6 months later to leave up to three food conditions, one at a time, on the ground without a novel

object:

• *Condition 1:* 30 chicken-flavoured dried dog food pellets

• *Condition 2:* 15 dried dog food pellets, 15 unsalted peanuts, 1 slice of deli chicken,

240             and 5 sprays of 35mL fish oil mixed with 900mL water

241         • *Condition 3:* 15 dried dog food pellets, 15 unsalted peanuts, 15 mL honey, 15 mL

242             strawberry jam

We then returned every 3 to 7 days for approximately two weeks to either replenish the

same condition or replace it with one of the other three conditions until foxes at each location had

an opportunity to discover at least one of the food conditions. Since our goal was to determine

whether foxes were willing to consume the food items used for novel object testing, we recorded

the following for all fox visits: 1) whether the food was still visible when the fox arrived, 2)

whether the fox acknowledged the presence of the food by directing its head and/or nose in the

exact spot where we left the food, and 3) whether the fox consumed the food, including food

remnants if some of the food was taken beforehand by another species.

**Factors affecting fox behavioural responses to novel objects**

*Methodological variables*

We examined the impact of novel object type because animals often respond differently to

different objects. We examined the effect of seasonality as well as our free food and reward

conditions because this might have impacted foxes' motivation to engage with the objects. We

examined the effect of the deodoriser spray because the scent of the puzzles could have deterred

foxes (e.g., human scent). Since animals' acknowledgement of objects plays a crucial role in

whether animals will have the opportunity to exploit them, we also considered whether fox

behaviour was impacted by the number of days from the time an object was first acknowledged by

a fox until the time we retrieved the equipment. Finally, rewards were sometimes exploited by

rodents and other organisms that were tiny enough to fit through the holes of objects; thus,

whenever possible, we kept records of the presence/absence of rewards at the time of foxes' initial

visits since this might have impacted their ability to detect and engage with the objects. This was

done two ways: 1) by taking a photo of the object whenever researchers visited to switch out the

camera's SD card, and 2) looking at the trail camera footage to see whether food was still present.

Sometimes we could not determine whether food rewards were still present if, for example, the

object was opaque, or we did not return to the location before a fox visited, or the camera footage

was not clear enough for us to see inside the transparent objects.

*Landscape variables*

Most UK residents live within cities and produce many millions of tonnes of waste per

year, which leads to significant problems with litter (DEFRA, 2022). Thus, as discussed, foxes

exposed to relatively higher levels of urbanisation will have greater access to anthropogenic food-

related objects. However, there is no single best way to classify an "urban" versus "rural"

population of animals given that the characteristics of urbanisation are so multi-faceted. Hence, to

allow us to more accurately evaluate the degree of urbanisation likely experienced by foxes across

our study locations, we used a range of variables recommended by Mu et al. (2022), including

human population density, road and greenspace density, land coverage (e.g., cropland), and species

richness. We also included measures of rainfall, temperature, and elevation because, for example,

they factor into cropland suitability.

Landscape data extraction was repeated for a series of circular buffers at 3.5km from the

epicentre of each zone in Figure 1. A Digital Elevation Model raster was sourced from the AWS

Open Data Terrain Tiles through the *elevatr* package at a 200m$^2$ pixel resolution (Hollister et al.,

2021). Average daily mean air temperature over the calendar year (in degrees Celsius) and total

precipitation over the calendar year (in millimetres) were obtained from the *HadUK-Grid* climate

observation dataset for the year 2021, the most recent available data, at a spatial resolution of 1km$^2$

per pixel (Hollis et al., 2019). Human population size data were collected from the *UK gridded*

*population census 2011* at a 1km$^2$ pixel resolution (Reis et al., 2017). Elevation, temperature,

rainfall, and human population size were extracted as the mean raster pixel value within each buffer

size. Road density (in m/m$^2$) within each buffer was computed by sourcing the highway/road class

vector layer from OpenStreetMap (*Planet dump*, 2022). Urban greenspace density (in m$^2$/m$^2$) was

obtained from the *Ordinance Survey Greenspace* vector layer (OPENGREESPACE, 2022).We

extracted percent coverage of five land cover classes by employing the UK Centre for Hydrology

and Ecology Land Cover 2020 product at a 10m$^2$ resolution (C. S. Morton et al., 2021). The raster

is composed of uniquely classified pixels according to categories following the UK Biodiversity

Action Plan, whom we aggregated into five main land cover categories: *urban* (class 20 and 21),

*forest* (class 1 and 2), *grassland* (class 4, 5, 6, 7, 9 and 10), *cropland* (class 3), and *wetland*

environments (class 8, 11, 12, 13, 14, 15, 16, 17, 18 and 19). Percent coverage was computed by

counting how many pixels within each buffer corresponded to each classified land cover and

dividing by the total number of pixels in each buffer. Landscape heterogeneity was also quantified

as the effective number of distinct land covers present in each buffer and computed as the

exponential of the Shannon-Wiener diversity index (Hill's numbers equivalent for q=1) (Chao et

al., 2014; Hill, 1973), resulting in a measure that not only accounts for the absolute richness of

diverse land cover categories but also weighs the relative abundance of each category.

**Statistical analyses**

Cohen's kappa coefficients were calculated between F.B.M. (who developed the

behavioural definitions) and two independent observers (who coded 25% of videos) to ensure that

the definitions were reliable (Table S2). Cohen's kappa coefficients were also calculated between

K.A. (who coded all videos) and F.B.M. (who trained K.A.) using 25% of videos to ensure that K.

A. was reliable prior to coding all of them (Table S3). Finally, Cohen's kappa coefficients were

calculated between 25% of K.A.'s videos at two different time periods (separated by several

months) to test for intra-observer consistency in K.A.'s codings (Table S4).

Landscape variables were calculated using the R programming language version *4.2.0*

within the RStudio IDE version *"Prairie Trillium"* (Team, 2002; Team, 2022). Geospatial

vectorial operations were processed utilizing the *sf* R package (Pebesma, 2018) while raster

extraction employed the *exactextractr* package (Baston, 2021). Data processing was conducted

through the use of the *tidyverse* R packages family (Wickham et al., 2019).

To obtain a global measure of urbanisation from each study location, we entered our

landscape variables into a Principal Components Analysis (PCA) with varimax rotation (Team,

2002). A scree test and parallel analysis were used to determine the number of components to

extract (Horn, 1965; Morton & Altschul, 2019). Item loadings $\geq |.4|$ were defined as salient for the

PCA; items with multiple salient loadings were assigned to the component with the highest

loading.

We used Chi-square tests to determine whether foxes were more likely than not to

acknowledge, touch, and exploit novel objects. We used multiple logistic regressions to test the

effects of our independent variables (i.e., methods and landscape characteristics) on whether foxes

acknowledged, physically touched, and ultimately exploited objects by gaining access to the

rewards inside.

Chi-square tests, Cohen's kappa tests, the PCA, and multiple logistic regressions were

conducted in IBM SPSS (Version 27). All datasets are provided in Datasets S1 and S2.

331                                    **Results**

**The fate of novel objects**

Foxes were recorded at 87 out of 181 locations (Figure 1). One fox successfully exploited

an object in the first instance (Video S1). After two weeks, foxes from another four locations had

gained accessed to the rewards (Video S2). Eleven objects (12.6%) were solved by other species,

while 65 objects (74.7%) had not been solved when researchers returned to retrieve them. The fate

of six objects (6.9%) could not be determined. Objects were more likely to remain unsolved by the

end of the two-week period compared to objects from all other locations, $\chi2$ (1, N = 87) = 21.3, p

< .001.

**Behavioural responses to novel objects**

Cohen's kappa coefficients were $k \geq$ .75 for foxes' acknowledgement and

apprehensiveness towards objects, indicating excellent inter-observer agreement (Tables S2-4).

Out of the 87 locations where foxes were recorded, eight locations were omitted from the

behavioural analysis due to poor visibility or camera malfunctions preventing us from observing

their initial responses to the objects. Of the remaining 79 locations, all foxes were solitary during

their initial visits. Sixty-four of these foxes (81%) acknowledged the novel objects by looking and

sniffing in their direction. Foxes were significantly more likely to acknowledge than ignore the

objects, $\chi^2$ (1, N = 79) = 30.4, p <.0001.

Of the foxes that acknowledged the objects, 27 foxes (42.2%) were too hesitant to approach

in the first instance (Video S3), while 37 foxes (57.8%) avoided the objects without appearing

apprehensive (Video S4). Collectively, foxes were equally likely to hesitate versus not hesitate

when they first discovered the objects, $\chi^2$ (1, N = 64) = 1.56, p = .211.

In terms of their behavioural responses at any point during the two weeks in which objects

were available, we could not determine whether foxes from 9 locations acknowledged them (e.g.,

poor visibility and camera malfunctions). At the remaining 78 locations, objects were

acknowledged at least once by foxes approximately 11.2 ± 3.6 days before we retrieved the

equipment. By the time we retrieved the equipment, foxes from at least 18 of these locations

(23.1%) had physically touched the objects, including the 5 foxes that successfully exploited them

and another 13 foxes that touched the objects but quit before gaining access (Video S5). Foxes that

acknowledged the objects at any point during the two weeks were more likely to avoid the objects

without touching them, $\chi2$ (1, N = 78) = 22.62, p < .001. Foxes that touched the objects were

equally likely to quit versus exploit them, $\chi2$ (1, N = 18) = 3.56, p = .06.

**Food tests**

364            Of the 30 locations where we deployed food without novel objects, foxes were detected at

365    23 locations, and 18 foxes discovered at least one of the food conditions before other animals

366    exploited them. All of these foxes willingly approached and consumed the food (Table 1 and Video

367    S6).

369    Table 1. Food conditions consumed by foxes.

| Location | Condition 1 | Condition 2 | Condition 3 |
|----------|-------------|-------------|-------------|
| 1 | Yes | Yes | Yes |
| 2 | --- | Yes | Yes |
| 3 | --- | Yes | --- |
| 4 | --- | Yes | --- |
| 5 | Yes | Yes | Yes |
| 6 | Yes | Yes | Yes |
| 7 | Yes | Yes | Yes |
| 8 | --- | Yes | Yes |
| 9 | --- | --- | Yes |
| 10 | --- | Yes | Yes |
| 11 | Yes | Yes | Yes |
| 12 | Yes | Yes | Yes |
| 13 | --- | --- | Yes |
| 14 | --- | Yes | Yes |
| 15 | --- | --- | Yes |
| 16 | --- | Yes | --- |
| 17 | --- | Yes | --- |
| 18 | Yes | Yes | Yes |

370            *Note*. "---" = food condition was not administered.

372    **Principal components analysis of landscape characteristics**

373            Across all 181 of our study locations, a PCA of our ecological and urban measures revealed

three components and explained 32.52, 28.71, and 11.98% of the variance, respectively (Table 2,

Figure S1, Table S5). Component 1 was labelled "Wilderness" because it was characterised by

item loadings related to lower levels of cropland and higher levels of natural and remote spaces

(e.g., forests, grasslands, and higher elevations). Component 2 was labelled "Urbanisation"

because it was characterised by higher levels of human, road, and greenspace densities, but lower

levels of cropland. Component 3 was labelled "Biodiversity" because it was characterised by high

levels of landscape heterogeneity and wetlands (i.e., an important habitat for many terrestrial and

aquatic species).

Table 2. Principal components analysis of ecological and urban variables (N = 181 locations).

|  | Varimax-rotated components | | |
| --- | --- | --- | --- |
| Item | PC 1 | PC 2 | PC 3 |
| Elevation | **.892** | -.001 | .16 |
| Rainfall | **.813** | .212 | .125 |
| LCC: Grassland | **.775** | -.117 | .268 |
| LCC: Forest | **.622** | -.14 | -.105 |
| Landscape heterogeneity 1 | .376 | .034 | **.741** |
| Landscape heterogeneity 0 | .236 | -.059 | **.682** |
| Greenspace density | .054 | **.727** | .021 |
| Road density | -.064 | **.965** | -.084 |
| Human population size | -.158 | **.918** | -.103 |
| LCC: Urban | -.183 | **.953** | -.123 |
| LCC: Wetland | -.344 | -.131 | **.758** |
| LCC: Cropland | **-.561** | **-.653** | -.352 |
| Temperature | **-.856** | .185 | -.04 |

384    *Note.* Salient loadings are in boldface. LCC=Land cover category. PC =
385          principal component.
386

**Factors affecting fox behavioural responses to novel objects**

At 56 locations, we were able to determine whether food rewards were still present at the

time of foxes' initial visits, but since rewards were missing at only three of these locations, we

omitted this variable from further analysis given the strong homogeneity of the data. When

considering the first fox video across locations (N = 79 foxes), none of the other methodological

variables or landscape composite scores (derived from each principal component) had a significant

effect on whether foxes acknowledged the novel objects (Table 3). Of the 64 locations where foxes

acknowledged the objects, the methodological variables and landscape composite scores also did

not significantly predict their hesitation to approach the objects when first discovered (Table 4).

Of the locations where foxes acknowledged the objects at some point during the two weeks, type

of novel object was the only predictor of foxes' likelihood of touching them (Table 5), with foxes

being more likely to touch objects B, E, F, H, and I (Figure 2 and 3).

Table 3. Multiple logistic regression predicting whether foxes acknowledged novel objects when

first discovered (N = 79 foxes).

| Variable | Exp (β) | p | 95% CI | |
|---|---|---|---|---|
| | | | Lower | Upper |
| Month | .947 | .605 | .772 | 1.162 |
| Object type | .816 | .095 | .643 | 1.036 |
| Food condition | .935 | .487 | .773 | 1.13 |
| Deodoriser | .792 | .757 | .181 | 3.473 |
| PC 1 (Wilderness) | .985 | .965 | .516 | 1.883 |
| PC 2 (Urbanisation) | 1.039 | .895 | .583 | 1.853 |
| PC 3 (Biodiversity) | 1.186 | .631 | .592 | 2.376 |

402   *Note*. PC = principal component.

409     Table 4. Multiple logistic regression predicting whether foxes behaved apprehensively towards

410     novel objects when first discovered (N = 64 foxes).

| Variable | Exp (β) | p | 95% CI | |
|---|---|---|---|---|
| | | | Lower | Upper |
| Month | .942 | .521 | .784 | 1.131 |
| Object type | .933 | .548 | .743 | 1.171 |
| Food condition | 1.039 | .630 | .888 | 1.217 |
| Deodoriser | 1.71 | .399 | .491 | 5.954 |
| PC 1 (Wilderness) | .684 | .202 | .381 | 1.226 |
| PC 2 (Urbanisation) | 1.146 | .603 | .685 | 1.917 |
| PC 3 (Biodiversity) | .86 | .623 | .471 | 1.57 |

411          *Note*. Significant results in bold. PC = principal component.

413     Table 5. Multiple logistic regression predicting the likelihood of foxes touching a novel object at

414     any point during the two weeks after they were discovered (N = 78 locations).

| Variable | Exp (β) | p | 95% CI | |
|---|---|---|---|---|
| | | | Lower | Upper |
| Month | .919 | .418 | .748 | 1.128 |
| Object type | **.712** | **.027** | **.527** | **.962** |
| Food condition | 1.01 | .917 | .84 | 1.215 |
| Deodoriser | .409 | .184 | .109 | 1.529 |
| Days since fox acknowledged object [a] | .915 | .418 | .738 | 1.135 |
| PC 1 (Wilderness) | 1.28 | .526 | .597 | 2.745 |
| PC 2 (Urbanisation) | .743 | .336 | .406 | 1.36 |
| PC 3 (Biodiversity) | 1.027 | .947 | .467 | 2.261 |

415          *Note*. Significant results in bold. [a] Number of days since first video of a fox
416          acknowledging object (i.e., opportunity to touch/exploit). PC = principal component.
417

419    Figure 3. Proportion of objects touched by foxes (at any point during two weeks) divided by the number of

420    objects (per type) deployed across 64 locations where foxes discovered them. Photos of each object are

421    depicted in Figure 2.

423                                        **Discussion**

We investigated the relationship between urbanisation and the likelihood of wild red foxes

exploiting novel food-related objects when first discovered. Of the foxes that acknowledged the

novel objects, most of them avoided the objects without gaining access. Given that we were able

to determine the fate of all objects, this rules out the possibility that our cameras significantly

missed footage of foxes visiting and exploiting their contents without us knowing it. Collectively,

our findings do not support the hypothesis that urbanisation encourages fox populations to exploit

novel food-related objects when first discovered; rather, as with rural populations, urban foxes

within the current study still displayed an initial aversion to such novelty.

Urban foxes clearly differ from rural populations in significant ways, for instance, in terms

of their diet and skeletal morphology (Parsons et al., 2020; Soulsbury et al., 2010), but our findings

are consistent with a growing number of studies in other species that show urbanisation is not

always a significant predictor of urban-rural differences in opportunistic behaviour, including

responses to novel objects (Griffin et al., 2017; Vincze & Kovacs, 2022). Although we are unable

to comment on other fox populations, our data suggest that foxes within the current study are

unlikely to thrive and maintain viable populations in urban settings by showing a greater

willingness to quickly exploit these novel opportunities in *general* (i.e., all contexts). Indeed, if

that was the case, then more urban foxes would have been willing to touch and exploit the objects

used in the current study.

Foxes always consumed the food rewards when objects were absent despite the presence

of a trail camera, ruling out the possibility that the cameras were a significant deterrent for them

in the novel object condition. Since foxes were willing to consume the food rewards when novel

objects were absent, it also rules out the possibility that food-related motivation explains why foxes

avoided the objects. Finally, as reported in Padovani et al. (2021), we found in our larger multi-

site sample no seasonal differences in foxes' responses to novel objects, which rules out seasonal

differences in food abundance, and hence, variation in their motivation due to hunger.

Further studies are needed to understand how foxes might balance decisions to exploit

versus avoid novel food-related objects, particularly in terms of the benefits, risks, and need for

doing so. Object neophobia, for example, could explain why foxes initially avoided the food when

objects were present, particularly foxes that showed clear signs of apprehension towards the

objects (Greggor et al., 2015; Miller et al., 2022; Travaini et al., 2013). If so, then present

ecological conditions within urban settings included in our study may not favour urban-rural

differences in object neophobia among foxes (Griffin et al., 2017). Factors, such as speed of

habituation, may be crucial for them to overcome their aversion to such novelty (Vincze et al.,

2016). Age, sex, dominance, associative learning, and/or personality are also some of the many

other factors that might influence foxes' decision-making (Fawcett et al., 2017; Griffin et al., 2013;

Padovani et al., 2021; Soulsbury et al., 2011a). As previously discussed, foxes may simply prefer

to exploit easy-access resources wherever possible, similar to what has been proposed for

challenging tasks in some populations of wild raccoons (*Procyon lotor*) (Morton, 2021). Finally,

as with some other species (e.g., corvids), foxes' responses to novel objects may be very nuanced

and depend on the particular context, such as the type of novel object encountered.

*Implications for red fox public perception and management*

Our findings have potentially important implications for changing the public perception

and management of foxes. In terms of public perception, our data support a more nuanced view of

urban fox behaviour by demonstrating that, in some respects, they can be shy and similar to rural

populations. Although further studies are needed, some household surveys show that many

residents label foxes as a 'nuisance' despite never actually experiencing a problem themselves

(Baker et al., 2020; Harris, 1981), which may be explained in part by the patterns of fox behaviour

observed in the current study.

In terms of fox management, many websites recommend chemicals (e.g., ammonia),

ultrasonic noise makers, or more lethal means of controlling nuisance foxes (e.g.,

removal/shooting). Based on our findings, however, we suggest that an alternative and more

humane approach may be to use novel objects as a deterrent. Indeed, foxes from our study were

still unwilling to exploit objects even when the reward quantity was high (e.g., 90 dog food pellets).

It is important to note that fox populations outside the current study may not necessarily respond

to novel objects in the same way, and novel objects will only be effective until foxes overcome

their aversion through habituation. Thus, to establish whether this management practice could be

effective for a given fox population, further research will need to understand, for example, what

characteristics of novel objects are better at deterring those foxes (e.g., size, shape, and proximity

to food), and how long the foxes will continue to avoid novel objects before they need to be

replaced with new ones.

484                                              **Conclusions**

Red foxes thrive within urban settings, but contrary to what has been found in some species,

we found that wild urban foxes were no more likely than rural populations to exploit novel food-

related objects when first discovered. These findings do not support the hypothesis that increased

access to such resources through urbanisation encourages foxes, at least within the current study,

to exploit them when first discovered; instead, other factors are likely needed for urban foxes to

overcome their initial aversion.

491                                              **References**

492   Anderson, M. J., Burpee, T. E., Wall, M. J., & McGraw, J. J. (2013). Exposure to novelty
493          weakens conditioned fear in long-evans rats. *Psicologica, 34*, 59-78.

494
495   Angel, S., Parent, J., Civco, D. L., & Blei, A. M. (2011). *Making room for a planet of cities*.
496          Lincoln Institute of Land Policy.

497
498   ASAB. (2020). Guidelines for the treatment of animals in beahvioural research and teaching.
499          *Animal Behaviour, 159*, i-xi.

500
501   Baker, S. E., Maw, S. A., Johnson, P. J., & Macdonald, D. W. (2020). Not in my backyard:
502          Public perceptions of wildlife and 'pest control' in and around UK homes, and local
503          authority 'pest control' *Animals, 10*, 222.

504
505   Baston, D. (2021). exactexractr: Fast extraction from raster datasets using polygons.

506
507   Brand, A., & Baldwin, M. (2020). Public attitudes to urban foxes in London and the south east.
508          *Mammal Communications, 6*, 34-41.

Bridge, B., & Harris, S. (2020). Do urban red foxes attack people? An exploratory study and review of incidents in Britain. *Human-Wildlife Interactions, 14*, 151-165.

Chao, A., Chiu, C.-H., & Jost, L. (2014). Unifying species diversity, phylogenetic diversity, functional diversity, and related similarity and differentiation measures through hill numbers. *Annual Review of Ecology, Evolution, and Systematics, 45*, 297-324.

Christensen, J. W., Zharkikh, T., & Chovaux, E. (2011). Object recognition and generalisation during habituation in horses. *Applied Animal Behaviour Science, 129*, 83-91.

Collenette, C. L. (1937). *A history of Richmond Park: with an account of its birds and animals*. Sidgwick & Jackson.

Contesse, P., Hegglin, D., Gloor, S., Bontadina, F., & Deplazes, P. (2004). The diet of urban foxes (Vulpes vulpes) and the availability of anthropogenic food in the city of Zurich, Switzerland *Mammalian Biology, 2*, 81-95.

Dammhahn, M., Mazza, V., Schirmer, A., Gottsche, C., & Eccard, J. A. (2020). Of city and village mice: behavioural adjustments of striped field mice to urban environments. *Scientific Reports, 10*, 13056.

DEFRA. (2022). *ENV23 - UK statistics on waste*. Retrieved 23 October, 2022 from https://www.gov.uk/government/statistical-data-sets/env23-uk-waste-data-and-management

Doncaster, C. P., Dickman, C. R., & Macdonald, D. (1990). Feeding ecology of red foxes (Vulpes vulpes) in the city of Oxford, England. *Journal of Mammology, 71*, 188.

Ducatez, S., Audet, J. N., Rodriguez, J. R., Kayello, L., & Lefebvre, L. (2017). Innovativeness and the effects of urbanization on risk-taking behaviors in wild Barbados birds. *Animal Cognition, 20*, 33-42.

Fawcett, J. K., Fawcett, J. M., & Soulsbury, C. D. (2017). Seasonal and sex-specific differences in feeding site attendance by red foxes Vulpes vulpes. *Mammal Study, 42*(2), 117-120.

Gatto, E., Dadda, M., Bruzzone, M., Chiarello, E., de Russi, G., Maschio, M. D., Bisazza, A., & Lucon-Xiccato, T. (2022). Environmental enrichment decreases anxiety-like behavior in zebrafish larvae *Developmental Psychobiology, 64*, e22255.

Gil-Fernandez, M., Harcourt, R., Newsome, T., Towerton, A., & Carthey, A. (2020).
Adaptations of the red fox (Vulpes vulpes) to urban environments in Sydney, Australia.
*Journal of Urban Ecology, 6*, 1-9.

Greenberg, R. (2003). The role of neophobia and neophilia in the development of innovative
behaviour in birds. In S. M. Reader & K. N. Laland (Eds.), *Animal Innovation* (pp. 175-
196). Oxford University Press.

Greenberg, R., & Mettke-Hofmann, C. (2001). Ecological aspects of neophobia and neophilia in
birds. *Current Ornithology, 16*, 119-178.

Greggor, A. L., Thornton, A., & Clayton, N. S. (2015). Neophobia is not only avoidance:
improving neophobia tests by combining cognition and ecology. *Current Opinion in
Behavioral Sciences, 6*, 82-89.

Griffin, A. S., Guez, D., Lermite, F., & Patience, M. (2013). Tracking changing environments:
innovators are fast, but not flexible learners. *PLoS ONE, 8*(12), e84907.

Griffin, A. S., Netto, K., & Peneaux, C. (2017). Neophilia, innovation and learning in an
urbanized world: a critical evaluation of mixed findings. *Current Opinion in Behavioral
Sciences, 16*, 15-22.

Harris, S. (1981). The food of suburban foxes (Vulpes vulpes), with special reference to London.
*Mammal Review, 11*, 151-168.

Hill, M. O. (1973). Diversity and evenness: A unifying notation and its consequences. *Ecology,
54*, 427-432.

Hollis, D., McCarthy, M., Kendon, M., Legg, T., & Simpson, I. (2019). HadUK-Grid - A new
UK dataset of gridded climate observations. *Geoscience Data Journal, 6*, 151-159.

Hollister, J., Shah, T., Robitaille, A. L., Beck, M. W., & Johnson, M. (2021). elevatr: Access
elevation data from various APIs. https://doi.org/doi:10.5281/zenodo.5809645

Horn, J. L. (1965). A rationale and test for the number of factors in factor analysis.
*Psychometrika, 30*, 179-185.

Lee. (1991). Adaptations to environmental change: an evolutionary perspective. In H. O. Box (Ed.), *Primate responses to environmental change* (pp. 39-56). Springer.

Lee, P. C., & Moura, A. (2015). Necessity, unpredictability and opportunity: An exploration of ecological and social drivers of behavioral innovation. In A. B. Kaufman & J. C. Kaufman (Eds.), *Animal Creativity and Innovation* (pp. 317-330). Academic Press.

Mazza, V., Czyperreck, I., Eccard, J. A., & Dammhahn, M. (2021). Cross-context responses to novelty in rural and urban small mammals. *Frontiers in Ecology and Evolution, 9*, 661971.

Mazza, V., & Guenther, A. (2021). City mice and country mice: innovative problem solving in rural and urban noncommensal rodents *Animal Behaviour, 172*, 197-210.

Meehan, C. L., & Mench, J. A. (2002). Environmental enrichment affects the fear and exploratory responses to novelty of young Amazon parrots. *Applied Animal Behaviour Science, 79*, 75-88.

Miller, R., Lambert, M. L., Frohnwieser, A., Brecht, K. F., Bugnyar, T., Crampton, I., Garcia-Pelegrin, E., Gould, K., Greggor, A. L., Izawa, E.-I., Kelly, D. M., Li, Z., Luo, Y., Luong, L. B., Massen, J. J. M., Nieder, A., Reber, S. A., Schiestl, M., Seguchi, A., Sepehri, P., Stevens, J. R., Taylor, A. H., Wang, L., Wolff, L. M., Zhang, Y., & Clayton, N. S. (2022). Socio-ecological correlates of neophobia in corvids. *Current Biology, 32*, P74-85.

Morton, C. S., Marston, R. D., O'Neil, C. G., & Rowland, A. W. (2021). Land Cover Map 2020 (10m classified pixels, GB). https://doi.org/doi:10.5285/35c7d0e5-1121-4381-9940-75f7673c98f7

Morton, F. B. (2021). Do wild raccoons (*Procyon lotor*) use tools? *Animal Cognition, 24*, 433-441.

Morton, F. B., & Altschul, D. (2019). Data reduction analyses of animal behaviour: avoiding Kaiser's criterion and adopting more robust automated methods. *Animal Behaviour, 149*, 89-95.

Morton, F. B., Buchanan-Smith, H. M., Brosnan, S. F., Thierry, B., Paukner, A., Essler, J. L., Marcum, C. S., & Lee, P. C. (2021). Studying animal innovation at the individual level: A ratings-based assessment in capuchin monkeys (Sapajus [Cebus] sp.). *Journal of Comparative Psychology, 135*(2), 258-265.

Mu, H., Li, X., Wen, Y., Huang, J., Du, P., Su, W., Miao, S., & Geng, M. (2022). A global
    record of annual terrestrial human footprint dataset from 2000 to 2018. *Scientific Data,*
    *176*.

OPENGREESPACE, O. (2022). *Contains OS data © Crown Copyright*.

Padovani, R., Shi, Z., & Harris, S. (2021). Are British urban foxes (Vulpes vulpes) "bold"? The
    importance of understanding human-wildlife interactions in urban areas. *Ecology and*
    *Evolution, 11*, 835-851.

Parsons, K. J., Rigg, A., Conith, A. J., Kitchener, A. C., Harris, S., & Zhu, H. (2020). Skull
    morphology diverges between urban and rural populations of red foxes mirroring patterns
    of domestication and macroevolution *Proceedings of the Royal Society B: Biological*
    *Sciences, 287*, 20200763.

Pebesma, E. (2018). Simple features for R: Standardized support for statistical vector data. *The R*
    *Journal, 10*, 439-446.

*Planet dump*. (2022). https://planet.osm.org

Reader, S. M., & Laland, K. N. (2003). *Animal innovation*. Oxford University Press. Table of
    contents only http://www.loc.gov/catdir/toc/fy044/2004296817.html

Reis, S., Liska, T., Steinle, S., Carnell, E., Leaver, D., Roberts, E., Vieno, M., Beck, R., &
    Dragosits, U. (2017). UK gridded population 2011 based on Census 2011 and Land
    Cover Map 2015. NERC Environmental Information Data Centre.

Rossler, T., Mioduszewska, B., O'Hara, M., Huber, L., Prawiradilaga, D. M., & Auersperg, A.
    M. I. (2020). Using an innovation arena to compare wild-caught and laboratory Goffin's
    cockatoos. *Scientific Reports, 10*, 8681.

Saunders, G., & Harris, S. (2000). Evaluation of attractants and bait preferences of captive red
    foxes (Vulpes vulpes). *Wildlife Research, 27*, 237-243.

Schell, C. J., Stanton, L. A., Young, J. K., Angeloni, L. M., Lambert, J. E., Breck, S. W., &
    Murry, M. H. (2021). The evolutionary consequences of human-wildlife conflict in cities.
    *Evolutionary Applications, 14*, 178-197.

Scott, D. M., Berg, M. J., Tolhurst, B. A., Chauvenet, A. L. M., Smith, G. C., Neaves, K.,
Lochhead, J., & Baker, P. J. (2014). Changes in the distribution of red foxes (Vulpes
vulpes) in urban areas in Great Britain: Findings and limitations of a media-driven
nationwide survey. *PLoS ONE, 9*, e99059.

Soulsbury, C. D., Baker, P. J., Iossa, G., & Harris, S. (2010). Red foxes (Vulpes vulpes). In S. D.
Gehrt, S. P. D. Riley, & B. L. Cypher (Eds.), *Urban carnivores: Ecology, conflict, and*
*conservation* (pp. 63-75). John Hopkins University Press.

Soulsbury, C. D., Iossa, G., Baker, P. J., White, P. C. L., & Harris, S. (2011a). Behavioral and
spatial analysis of extraterritorial movements in red foxes (Vulpes vulpes). *Journal of*
*Mammology, 92*, 190-199.

Soulsbury, C. D., Iossa, G., Baker, P. J., White, P. C. L., & Harris, S. (2011b). Behavioral and
spatial analysis of extraterritorial movements in red foxes (Vulpes vulpes). *Journal of*
*Mammology, 92*(1), 190-199.

Soulsbury, C. D., & White, P. (2015). Human-wildlife interactions in urban areas: A review of
conflicts, benefits and opportunities. *Wildlife Research, 42*, 541-553.

Teagle, W. G. (1967). The fox in the London suburbs. *The London Naturalist, 46*, 44-68.

Team, R. (2002). *RStudio: Integrated Development for R.*

Team, R. C. (2022). *R: A language and environment for statistical computing.*

Thompson, M. J., Capilla-Lasheras, P., Dominoni, D. M., Reale, D., & Charmantier, A. (2021).
Phenotypic variation in urban environments: mechanisms and implications *Trends in*
*Ecology & Evolution, 37*, 171-182.

Thornton, A., & Samson, J. (2012, Jun). Innovative problem solving in wild meerkats. *Animal*
*Behaviour, 83*, 1459-1468.

Travaini, A., Vassallo, A. I., Garcia, G. O., Echeverria, A. I., Zapata, S. C., & Nielsen, S. (2013).
Evaluation of neophobia and its potential impact upon predator control techniques: A
study on two sympatric foxes in southern Patagonia. *Behavioural Processes, 92*, 79-87.

Trewhella, W. J., Harris, S., & McAllister, F. E. (1988). Dispersal distance, home-range size and
population density in the red fox (Vulpes vulpes): A quantitative analysis. *Journal of*
*Applied Ecology, 25*, 423-434.

Vincze, E., & Kovacs, B. (2022). Urbanization's effects on problem solving abilities: A meta-
analysis. *Frontiers in Ecology and Evolution, 10*, 834436.

Vincze, E., Papp, S., Preiszner, B., Seress, G., Bokony, V., & Liker, A. (2016). Habituation to
human disturbance is faster in urban than rural house sparrows. *Behavioral Ecology, 27*,
1304-1313.

Visalberghi, E., & Limongelli, L. (1994). Lack of comprehension of cause-effect relations in
tool-using capuchin monkeys (*Cebus apella*). *Journal of Comparative Psychology,*
*108*(1), 15-22.

Webster, M. M., Ward, A. J. W., & Hart, P. J. B. (2009). Individual boldness affects interspecific
interactions in sticklebacks *Behavioral Ecology and Sociobiology, 63*, 511-520.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., Francois, R., Grolemund, G.,
Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M.,
Muller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., Takahashi, K., Vaughan, D.,
Wilke, C., Woo, K., & Yutani, H. (2019). Welcome to the tidyverse. *Journal of Open*
*Source Software, 4*, 1686.

Wilkinson, D., & Smith, G. C. (2001). A preliminary survey for changes in urban fox (Vulpes
vulpes) densities in England and Wales, and implications for rabies control. *Mammal*
*Review, 31*, 107-110.