Bold but not innovative in an urban exploiter, the red fox (*Vulpes vulpes*)

F. Blake Morton*[1], Marieke Gartner[2], Ellie-Mae Norrie[1], Yacob Haddou[3], Carl D. Soulsbury[4],

Kristy A. Adaway[1]

[1] Department of Psychology, University of Hull, Hull, UK

[2] Atlanta Zoo, Atlanta, Georgia, USA

[3] Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow,

Glasgow, UK

[4] School of Life and Environmental Sciences, University of Lincoln, Lincoln, UK

Author Note

F. Blake Morton https://orcid.org/0000-0002-8644-1557

Marieke Gartner https://orcid.org/0000-0003-3901-9166

Yacob Haddou https://orcid.org/0000-0003-0557-0339

Carl D. Soulsbury https://orcid.org/0000-0001-8808-5210

*Author for correspondence: Dr Blake Morton, Department of Psychology, University of Hull,

HU6 7RX, United Kingdom. E-mail: b.morton@hull.ac.uk.

Acknowledgments

The authors would like to thank everyone involved with the 'British Carnivore Project', particularly Sophie Tait, Josh Chatterton, Alice Turner, Eszter Jardan, Dylan Jones, Louise Grunnill, Katherine Sutter, and the citizen scientists who kindly gave us permission to collect data in their gardens. We are very grateful to the many organisations that helped us acquire permission to access other land for the study, particularly staff from the Yorkshire Wildlife Trust, Lincolnshire Wildlife Trust, Scottish Wildlife Trust, The Land Trust, the National Trust, Forestry England (particularly Cath Bashforth), Forestry & Land Scotland, Yorkshire Water, and various city and regional councils (particularly Hull, East Riding of Yorkshire, and Glasgow). Special thanks go to Prof. Hannah Buchanan-Smith (Uni Stirling) and Bruce Gittings (Uni Edinburgh) for their help with various logistics in Scotland, to Dr Alex Weiss (Uni Edinburgh) and Dr Henning Hole (Uni Hull) for statistical discussions, and to Prof. Phyllis Lee (Uni Stirling) and Dr Kevin Parson (Uni Glasgow) for helpful comments on an earlier draft of the manuscript. Finally, FBM is grateful to the University of Hull, UKRI Natural Environment Research Council (NERC) (Grant No. NE/X018342/1), and EU Social Fund Plus for funding.

Conflict of interest statement: The authors declare no conflict of interest.

Highlights:

- The impact of urbanisation on animal adaptability remains unclear

- Bold and innovative behaviour may help some urban species thrive

- We studied wild red foxes' responses to novel food-related objects

- Urban foxes were bolder, but not more innovative, than rural foxes

- Urbanisation may favour bolder, not more innovative, fox behaviour

Abstract

Urbanisation is the fastest form of landscape transformation on the planet, but

researchers' understanding of the relationships between urbanisation and animal adaptability

is still in its infancy. In terms of foraging, bold and innovative behaviours are proposed to help

urban animals access, utilise, and exploit novel anthropogenic food sources. Red foxes

(*Vulpes vulpes*) are one of the best known and widespread urban-dwelling species. However,

despite frequent stories, images, and videos portraying them as "pests" due to their

exploitation of food-related objects (e.g., raiding the contents of outdoor bins), it is unknown

whether they are bolder and more innovative in terms of their likelihood of exploiting these

resources compared to rural populations. In the current study, we gave novel food-related

objects to foxes from 104 locations (one object per location) across a large urban-rural

gradient. To access the food, foxes had to use behaviours necessary for exploiting many food-

related objects in the real world (e.g., biting, pushing, pulling, or lifting human-made materials).

Despite all foxes acknowledging the objects, foxes from 31 locations touched them, while

foxes from 12 locations gained access to the food inside. A principal component analysis of

urban and other landscape variables (e.g., road, greenspace, and human population density)

revealed that urbanisation was significantly and positively related to the likelihood of foxes

touching, but not exploiting, the objects. Thus, while urban foxes may be bolder than rural

populations in terms of their willingness to physically touch novel food-related objects, our

findings are inconsistent with the notion that they are more innovative and pose a general

nuisance to people by exploiting these anthropogenic resources.

Key words: behavioural flexibility, boldness, human-wildlife conflict, neophobia, problem-

solving, urbanisation

Introduction

Urbanisation is the fastest form of landscape transformation on the planet (Angel et al., 2011; Grimm et al., 2008), with 55% of the global human population now living within cities (UN, 2018). Urban environments present wildlife with a range of novel challenges that can include coping with habitat loss and fragmentation (Šálek et al., 2015), increased or novel human disturbances (Rodrigo-Comino et al., 2021), altered competitive interactions (Martin & Bonier, 2018), and new predators or parasites (Guiden et al., 2019; Pedroso-Santos & Costa-Campos, 2020). Species can be characterised based on a gradient of how they adapt to urban environments, including **1)** "*urban avoiders*", which are restricted to non-urban or remnant natural habitats, **2)** "*urban utilisers*", which make occasional use of urban areas, and **3)** "*urban dwellers*", which actively exploit and benefit from urban areas (Fischer et al., 2015). The ability for species to persist and thrive in urban environments is related to a suite of life history, morphological, physiological, behavioural, and cognitive factors (Charmantier et al., 2017; Sol et al., 2014), but researchers' understanding of how animals adapt to urban environments is still in its infancy.

In terms of foraging, species dwelling in urban areas are likely to encounter novel anthropogenic food sources (Murray et al., 2015; Murray et al., 2018) and behavioural traits, particularly boldness and innovation, are proposed to help urban animals access, utilise, and exploit these resources (Dammhahn et al., 2020; Ducatez et al., 2017; Griffin et al., 2017; Mazza et al., 2021; Mazza & Guenther, 2021). Having a greater tendency to innovate can provide urban wildlife with the behavioural flexibility needed to exploit a wide variety of resources (Reader & Laland, 2003). Being more likely to quickly display such behaviour can enable urban wildlife to exploit these opportunities before they are taken by other animals or removed by city cleaners (Webster et al., 2009). To date, however, not all studies find that urban dwellers are bolder and more innovative for reasons that remain unclear (Griffin et al., 2017; Vincze & Kovacs, 2022).

Red foxes (*Vulpes vulpes*) are one of the best-known and widespread urban-dwelling species (Soulsbury et al., 2010). They are an opportunistic generalist omnivore, which enables

them to exploit a diverse range of food items, including mammals, bird, invertebrates, and

plants. In urban areas, foxes will also scavenge a wide variety of anthropogenic food items

from various sources, including bird feeders, compost heaps, bins, and food provisioned by

people (Contesse et al., 2004b; Doncaster et al., 1990; Saunders et al., 1993). Such use of

anthropogenic materials suggests that urban foxes are willing to exploit new feeding

opportunities, but although urban foxes are often *labelled* as being generally bolder than their

rural counterparts, it is unknown whether this is true in all contexts. It is also unknown whether

they are more innovative.

Urban foxes often encounter food-related objects that are temporally, physically, and

spatially "novel" to them, including **1)** continuous changes to the combination of objects found

on streets or in outdoor bins, **2)** objects that look physically different to what animals are

accustomed to seeing (e.g., new or modified containers), and **3)** new or familiar objects found

in unexpected locations (e.g., randomly discarded trash). Such dynamic changes, combined

with frequent encounters, may favour bolder and more innovative behaviour in foxes by

enabling them to use new or modified behaviours (i.e., "innovations") to exploit these

resources, particularly shortly after discovering them (e.g., overnight) (Dammhahn et al., 2020;

Ducatez et al., 2017; Griffin et al., 2017; Mazza et al., 2021; Mazza & Guenther, 2021).

However, despite frequent stories, images, and videos within popular culture portraying urban

foxes as "pests" due to their opportunistic foraging behaviour (Schell et al., 2021; Soulsbury

& White, 2015), it is unclear whether or to what extent such attitudes are due, in part, to their

exploitation of food-related objects, including discarded litter and items found in outdoor bins

(Baker et al., 2020; Harris, 1981).

In the past, studies have given novel objects to urban foxes (Padovani et al., 2021),

but the objects did not contain food and comparisons with rural populations were not made,

making it impossible to evaluate the likelihood of urban foxes behaving bolder and more

innovative within this context. Although urban foxes may be more likely to consume novel bait

(Gil-Fernandez et al., 2020), this does not necessarily reflect how animals react to other forms

of novelty, including human-made objects (Miller et al., 2022). Hence, the current study had

two aims: First, to test whether urban foxes are bolder and more innovative than rural populations in terms of exploiting novel food-related objects, and second, to test whether urban foxes are indeed a general nuisance to people because they exploit these anthropogenic resources.

Methods and materials

Ethical statement

This study was ethically approved by the Animal Welfare Ethics Board of the University of Hull (FHS356), and was carried out in accordance with guidelines outlined by the Association for the Study of Animal Behaviour (ASAB, 2020). No foxes were handled, all trail cameras were placed away from footpaths to minimise public disturbance, and food items used to attract foxes were not harmful if ingested by other animals, including outdoor pets.

Study sites and subjects

We studied 200 locations throughout Scotland and England (Figure 1), including areas in and around different cities (e.g., London, Glasgow, Edinburgh, Stirling, Leeds, Hull, Lincoln, Sheffield, and York). These locations covered a wide variety of landscapes, including recreational parks, private gardens, tree plantations, meadows, mixed woodland, coastal and mountainous scrubland, and farmland. Foxes were unmarked and their participation in the study was entirely voluntary. We gained access to 162 of these locations by contacting city councils and other organisations that owned land. The remaining 38 locations were private gardens, which we accessed by advertising the study through Twitter and regional wildlife groups. Our criteria for including any location in the study included: **1)** landowner permission, **2)** accessibility to foxes (e.g., no barriers/fences), **3)** ability to place our equipment out of public view to avoid theft or vandalism, and **4)** the location could not be $\leq$ 3.5 km from another study area. This latter criterion was used to reduce the chances of sampling the same fox across more than one location because $\geq$ 3.5 km is larger than the typical dispersal distance and home range diameter of British foxes (Soulsbury et al., 2011; Trewhella et al., 1988). We did

175   not have prior knowledge of fox presence before contacting landowners, and we included

176   locations in the study regardless of whether landowners said foxes lived on or near their land.

Figure 1: Distribution of locations where objects were deployed across Scotland and England, with (a) foxes detected (yes or no) and (b) whether

foxes acknowledged, touched, or exploited ("solved") the food-related objects.

Designs and method of administering novel food-related objects

We administered 8 types of food-related objects (Figure 2) across our study locations

between August 2021 and November 2023. Only a single object was administered per

location, and they were available to foxes for 15.5 ± 1.64 days before we removed them.

Although foxes might, of course, respond differently to food-related objects that are left for

longer, two weeks is a very typical timeframe for many food-related objects available to British

urban foxes (e.g., regular street cleaning and bin services every 1-2 weeks).

The objects were made from basic household materials (e.g., PVC piping, metal

screws, and wooden rods). Objects varied in terms of design and materials to ensure that our

data on foxes' behavioural responses were more generalisable and not specific to just one

type of object. Objects were "novel" in terms of their location, which we verified by searching

for similar objects within the surrounding areas. Objects were also novel in terms of their

design, which we assembled ourselves using a unique combination of materials to create

objects that are not widely commercially available, making it highly unlikely that foxes would

have seen those specific combinations before. Each object had a single 'free food' and

'reward' condition (Table S1); the 'free food' was scattered approximately 1m away from each

object. We used different types, combinations, and quantities of food to ensure that our data

on foxes' behavioural responses were more generalisable and not specific to any particular

food. All objects were anchored to the plastic platform and had holes drilled into them to

facilitate odour cues. Tent pegs were used to anchor the platforms to the ground.

Figure 2: Food-related objects administered to foxes. Yellow dashed arrows indicate the direction of each behaviour needed to retrieve the food

rewards inside. Task G was never deployed in the field and hence not depicted in this figure.

Object C had two levels, each containing food. To access the rewards in Object D,

foxes simply had to push through the aluminium side of the box. The lid of Object F was fixed

in place and could only be opened by sliding it either to the left or right. Object H had a hidden

axle to allow 360° rotation. Object I was administered with the stick already inside the pipe;

animals merely had to remove the stick using their mouths, which would indirectly rake the

food out.

Researchers were not present when foxes visited, and we did not touch or replenish

the food to avoid unnecessary disturbance to the objects. Following APHA guidelines, we

cleaned objects with antibacterial soap and 70% alcohol wipes after retrieving them to prevent

possible pathogen transmission. We then washed and dried them prior to redeployment. Forty-

two objects (21%) were sprayed with scent deodoriser to test whether the scent of objects

(e.g., human odour) had a significant effect on fox behaviour.

Since foxes were free-ranging and their participation in the study was entirely

voluntary, some foxes might have avoided our testing locations. Nevertheless, the goal of this

study was to test foxes' likelihood of being bold and innovative enough to *exploit* the objects

within a two-week period, which required them to physically touch the objects (and hence be

detected on camera). We therefore based our analysis on foxes that were at least able and

willing to visit the locations.

238   Recording fox behaviour from trail cameras

At each location, we horizontally placed a 'no glow' (940 nm) infrared motion-sensor

camera (Apeman H45) approximately 4m away on a tree trunk. Cameras had a 120 ° sensing

angle and a triggering distance of 20 m. Video lengths were set to record for 5 min, with a 5 s

trigger delay and a 30 s interval in between each video. Camera lenses were sprayed with

defogger and, where possible, minor amounts of understory vegetation were removed

between the camera and object to ensure optimal visibility.

Measuring urban-rural differences in bold and innovative behaviour

Measuring urban-rural differences in bold and innovative behaviour

A myriad of factors can underpin bold and innovative behaviour, which are not

necessarily due to any single variable (Griffin et al., 2014; Lee & Moura, 2015; F. B. Morton et

al., 2021; Reader & Laland, 2003). Animals, for example, may not use such behaviour to

exploit novel objects if they are too afraid or not hungry. Crucially, however, our goal was to

determine *whether* (not why) subjects would display bold and innovative behaviour to exploit

food-related objects, and so the only way they could do this was by physically engaging with

novel objects themselves.

Foxes could gain access to the food rewards through persistence and by using simple

behaviours used to exploit human-made objects in the real world (e.g., using their mouth,

nose, and/or paws to bite, push, pull, or lift materials). Some of the designs were inspired from

studies of behavioural innovation in other species (Morton, 2021; Rossler et al., 2020;

Thornton & Samson, 2012; Visalberghi & Limongelli, 1994). As with these other studies, we

defined innovation as any behaviour used to operate and successfully gain access to the food

inside each novel object (Figure 2).

To determine whether urban foxes were faster to display bold and innovative

behaviours, we compared differences in urban and rural foxes' likelihood of touching and

exploiting (at any point) the objects. We defined 'touching' objects in terms of foxes pushing,

pulling, licking, and/or biting them, or making physical contact with their nose while smelling

them. We defined 'acknowledging' objects as a fox turning its head to look/smell in the object's

direction. We tested for inter-observer agreement for all behaviours, and there was excellent

agreement (k > .75) between K.A. (who coded all videos), F.B.M. (who developed the

definitions and trained K.A.), and several independent coders (Tables S2-5).

While indeed there might be alternative ways of measuring how quickly foxes display

bold behaviour, such as walking speed or the latency to approach the objects to within a

certain body length, as mentioned before, our research question was related to whether foxes

were bold enough to *exploit* them, which required them to physically touch the objects

regardless of how long it might have taken them to walk up to them. Similarly, while there may

be alternative ways of measuring how quickly foxes innovate, such as the amount of time

spent operating the task until a solution was found, this was not possible due to occasional

camera malfunctions or some of the videos having poor visibility (e.g., fog or raindrops on the

lens); thus, it was more practical, and equally fit for purpose, to analyse how likely urban and

rural fox populations were to exploit the food rewards as a function of how many days since

the objects were discovered.

Food tests

Foxes are generalist carnivores and should be highly motivated to consume the food

rewards in our study (Saunders & Harris, 2000). To confirm this, we revisited 30 of our

locations six months later to leave up to three food conditions, one at a time, on the ground

without an object:

• *Condition 1:* 30 chicken-flavoured dried dog food pellets

• *Condition 2:* 15 dried dog food pellets, 15 unsalted peanuts, 1 slice of deli

chicken, and 5 sprays of 35mL fish oil mixed with 900mL water

• *Condition 3:* 15 dried dog food pellets, 15 unsalted peanuts, 15 mL honey, 15

mL strawberry jam

All of these locations were within the Yorkshire area. We returned every 3 to 7 days

for approximately two weeks to either replenish the same condition or replace it with one of

the other three conditions until foxes at each location had an opportunity to discover at least

one of the food conditions. Since our goal was to determine whether foxes would consume

the food items placed within objects, we recorded the following for all fox visits: **1)** whether the

food was still visible when the fox arrived, **2)** whether the fox acknowledged the presence of

the food by directing its head and/or nose in the exact spot where we left the food, and **3)**

whether the fox consumed the food, including food remnants if some of the food was taken

beforehand by another species.

Factors affecting fox detection and behavioural responses to food-related objects

Methodological variables

We examined the impact of object type and food conditions (Table S1), because these may have impacted foxes' motivation to engage with the objects. We examined the effect of the deodoriser spray because the scent of the puzzles could have deterred foxes (e.g., human scent). Since cameras were not always fully operational (e.g., SD cards full or batteries died), we also examined the impact of the amount of time each camera operated (divided by total days deployment time) after objects were acknowledged by foxes.

Rewards were sometimes exploited by rodents and other organisms that were tiny enough to fit through the holes of objects; thus, whenever possible, we kept records of the presence/absence of rewards at the time of foxes' initial visits since this might have impacted their ability to detect and engage with the objects. This was done two ways: **1)** by taking a photo of the object whenever researchers visited to switch out the camera's SD card, and **2)** looking at the trail camera footage to see whether food was still present. Sometimes we could not determine whether food rewards were still present if, for example, the object was opaque, or we did not return to the location before a fox visited, or the camera footage was not clear enough for us to see inside the transparent objects. At 78 locations, we were able to determine whether food rewards were still present at the time of foxes' initial visits, but since rewards were missing at only 5 (6.5%) of these locations, we omitted this variable from further analysis given the strong homogeneity of the data.

Landscape variables

Most UK residents live within cities and produce many millions of tonnes of waste per year, which leads to significant problems with litter (DEFRA, 2022). Thus, as discussed, foxes exposed to relatively higher levels of urbanisation will have greater access to anthropogenic food-related objects. However, there is no single best way to classify an "urban" versus "rural" population of animals given that the characteristics of urbanisation are so multi-faceted. Hence, to allow us to more accurately evaluate the degree of urbanisation likely experienced

by foxes across our study locations, we used a range of variables recommended by Mu et al. (2022), including human population density, road and greenspace density, land coverage (e.g., cropland), and species richness. We also included measures of rainfall, temperature, and elevation because, for example, they factor into cropland suitability.

Landscape data extraction was repeated for a series of circular buffers at 3.5 km from the epicentre of each zone in Figure 1. A Digital Elevation Model raster was sourced from the AWS Open Data Terrain Tiles through the *elevatr* package at a 200m^2 pixel resolution (Hollister et al., 2021). Average daily mean air temperature over the calendar year (in degrees Celsius) and total precipitation over the calendar year (in millimetres) were obtained from the *HadUK-Grid* climate observation dataset for the year 2021, the most recent available data, at a spatial resolution of 1 km^2 per pixel (Hollis et al., 2019). Human population size data were collected from the *UK gridded population census 2011* at a 1 km^2 pixel resolution (Reis et al., 2017). Elevation, temperature, rainfall, and human population size were extracted as the mean raster pixel value within each buffer size. Road density (in m/m^2) within each buffer was computed by sourcing the highway/road class vector layer from OpenStreetMap (*Planet dump*, 2022). Urban greenspace density (in m^2/m^2) was obtained from the *Ordinance Survey Greenspace* vector layer (OPENGREESPACE, 2022). We extracted percent coverage of five land cover classes by employing the UK Centre for Hydrology and Ecology Land Cover 2020 product at a 10 m^2 resolution (C. S. Morton et al., 2021). The raster is composed of uniquely classified pixels according to categories following the UK Biodiversity Action Plan, which we aggregated into five main land cover categories: *urban* (class 20 and 21), *forest* (class 1 and 2), *grassland* (class 4, 5, 6, 7, 9 and 10), *cropland* (class 3), and *wetland* environments (class 8, 11, 12, 13, 14, 15, 16, 17, 18 and 19). Percent coverage was computed by counting how many pixels within each buffer corresponded to each classified land cover and dividing by the total number of pixels in each buffer. Landscape heterogeneity was also quantified as the effective number of distinct land covers present in each buffer and computed as the exponential of the Shannon-Wiener diversity index (Hill's numbers equivalent for q=1) (Chao et al., 2014; Hill, 1973).

358   Landscape variables were calculated using the R programming language version *4.2.0*

359   within the RStudio IDE version *"Prairie Trillium"* (Team, 2002; Team, 2022). Geospatial

360   vectorial operations were processed utilizing the *sf* R package (Pebesma, 2018) while raster

361   extraction employed the *exactextractr* package (Baston, 2021). Data processing was

362   conducted through the use of the *tidyverse* R packages family (Wickham et al., 2019).

Statistical analyses

To obtain a global measure of urbanisation from each study location, we entered our

landscape variables into a principal component analysis (PCA) with varimax rotation (Team,

2002). A scree test and parallel analysis were used to determine the number of components

to extract (Horn, 1965; Morton & Altschul, 2019). Item loadings $\geq |.4|$ were defined as salient

for the PCA; items with multiple salient loadings were assigned to the component with the

highest loading.

We first tested whether our methodological variables (object type, deodoriser, season,

camera operation time, and food conditions) impacted the likelihood of (a) a fox being detected

or (b) touching the object, using binary logistic regression. We then carried binary generalised

linear mixed effects models (GLMMs) to test the effect of urbanisation on fox behaviour. In our

first model, we tested whether detecting a fox was related to habitat (PCA1, PCA2, PCA3),

with food type included as a random factor. In our second model, we tested whether the fox

touching the object was related to habitat (PCA1, PCA2, PCA3). We also included "*camera*"

(i.e., the proportion of time the camera operated after objects were acknowledged by foxes)

as an additional covariate, and food as a random effect. Finally, for foxes that touched the

objects, we tested whether their ability to access the food inside them was related to habitat

(PCA1, PCA2, PCA3). Again, we included the variable "*camera*" as an additional covariate,

and food as a random effect. All GLMMs were run using the lme4 package (Bates et al., 2015),

with the significance of fixed effects in binomial GLMMS tested using Wald χ^2 tests

implemented in the ANOVA function of the car package (Fox & Weisberg, 2019).

Chi-square tests, Cohen's kappa tests, and the PCA were conducted in IBM SPSS

(Version 27). All other analyses were conducted in R version 4.1.1 (Team, 2021). All data are

provided in Datasets S1 and S2 in the supplementary materials.

388   **Results**

Food tests

Of the 30 locations where we conducted food tests, foxes were detected at 23

locations, and 17 foxes discovered at least one of the food conditions before other animals

exploited them. All of these foxes approached and consumed the food (Table S6 and Video

S2).

Likelihood of foxes touching and exploiting food-related objects

During the period in which objects were deployed, foxes were recorded at 104 (52%)

locations (Figure 1a). Out of the 104 locations where foxes were recorded, it was not possible

to tell whether foxes acknowledged objects at 8 (7.7%) locations due to poor visibility or

camera malfunctions. In all the remaining 96 locations across all habitats, foxes acknowledged

the objects. Foxes went on to touch the objects at 31 locations (32%), and of these, 12 (40%,

1 location could not be determined) exploited the food inside objects (Figure 1b).

Principal component analysis of landscape characteristics

Across our 200 study locations (Figure 1), a PCA of our ecological and urban measures

revealed three components and explained 29.23, 29.05, and 14.47% of the variance,

respectively (Table 2, Figure S1, Table S7). Component 1 was labelled "Wilderness" because

it was characterised by item loadings related to lower levels of cropland and higher levels of

natural and remote spaces (e.g., forests, grasslands, and higher elevations). Component 2

was labelled "Urbanisation" because it was characterised by higher levels of human, road, and

greenspace densities, but lower levels of cropland. Component 3 was labelled "Biodiversity"

because it was characterised by high levels of landscape heterogeneity and wetlands (i.e., an

important habitat for many terrestrial and aquatic species).

Table 2: Principal component analysis of ecological and urban variables (N = 200 locations).

	Varimax-rotated components		
Item	PC 1	PC 2	PC 3
Temperature	**-.811**	.349	-.009
Rainfall	**.827**	.179	.11
Elevation	**.883**	-.008	.167
Human population size	-.238	**.875**	-.091
Greenspace density	.066	**.71**	-.031
Road density	-.143	**.959**	-.068
LCC: Urban	-.227	**.939**	-.103
LCC: Forest	**.629**	-.148	-.094
LCC: Grassland	**.773**	-.131	.267
LCC: Cropland	**-.507**	**-.702**	-.348
LCC: Wetland	-.346	-.133	**.744**
Landscape heterogeneity 0	.203	0	**.716**
Landscape heterogeneity 1	.365	-.031	**.742**

415
416   *Note.* Salient loadings are in boldface. LCC=Land cover category. PC =
         principal component.
417

Effect of methodological variables on the likelihood of fox detection and behaviour

Fox detection on camera was not significantly affected by object type, food condition,

or deodoriser spray (Table 3). Similarly, the likelihood of foxes touching an object was not

related to the object used, deodoriser spray or the proportion of time the camera was

operational (Table 3). There was no significant effect of food condition on the likelihood of

foxes touching objects (Table 3; Figure 3).

431 **Table 3:** Fixed effects from two binary logistic regression models tested using the likelihood

432 ratio χ^2. In each, we tested methodological variables and their likely impact on (a) fox

433 detection and (b) foxes' physical engagement with objects.

Model	Parameter	Likelihood Ratio χ^2	d.f.	P
(a) Fox detection	Object	7.03	7	0.426
	Food	10.91	11	0.451
	Deodoriser	0.12	1	0.731
(b) Fox touches object	Object	3.45	7	0.578
	Food	18.01	11	0.081
	Deodoriser	0.91	1	0.341
	Camera	0.93	1	0.335

434 *Note.* "*Camera*" is the proportion of time the camera operated after objects were
435 acknowledged by foxes.
436

439 **Figure 3:** The proportion of times the foxes touched the objects depending on food condition.

440 Numbers over bars indicated the number of times foxes acknowledged the object for each

441 food condition.

Effect of landscape characteristics on the likelihood of fox detection and touching and exploiting objects

The probability of detecting a fox on camera was significantly lower in more wilderness areas (PCA1: Figure 4a) and greater in more urbanised (PCA2) areas (Table 4; Figure 4b). PCA2 (Urbanisation) was positively associated with foxes touching an object (Table 4; Figure 5), but there was no effect of PCA1 (Wilderness) or PCA3 (Biodiversity) (Table 4). Finally, of those foxes that touched the objects, there was no effect of habitat (PCA1, PCA2, PCA3) on the likelihood of the objects being exploited (Table 4).

Table 4: Fixed effects for three binomial GLMM models tested using Wald χ^2 tests. In each, we tested the impact of landscape characteristics on the likelihood of (a) fox detection and foxes (b) touching and (c) exploiting objects.

Model	Parameter	Wald χ^2	d.f.	P
(a) Fox detection	**PCA 1 (Wilderness)**	**6.07**	**1**	**0.014**
	PCA 2 (Urbanisation)	**46.33**	**1**	**<0.001**
	PCA 3 (Biodiversity)	0.29	1	0.589
(b) Fox touches object	PCA 1 (Wilderness)	1.47	1	0.225
	PCA 2 (Urbanisation)	**9.99**	**1**	**0.002**
	PCA 3 (Biodiversity)	1.48	1	0.224
	Camera	0.04	1	0.844
(c) Fox exploits object	PCA 1 (Wilderness)	2.04	1	0.153
	PCA 2 (Urbanisation)	1.71	1	0.191
	PCA 3 (Biodiversity)	0.63	1	0.426
	Camera	0.15	1	0.697

Note. Significant values are in bold. "*Camera*" is the proportion of time the camera operated after objects were acknowledged by foxes.

Figure 4: The relationship between (a) foxes being detected by camera in relation to the

degree of wilderness (PCA1) and (b) foxes being detected by camera in relation to the

degree of urbanisation (PCA2).

Figure 5: The likelihood of a fox touching a food-related object in relation to the degree of

urbanisation (PCA2).

Discussion

We investigated whether urban foxes are bolder and more innovative than rural populations in terms of exploiting novel food-related objects, and whether such behaviour is consistent with the popular notion that urban foxes are a "pest" because they exploit these anthropogenic resources. Although foxes acknowledged the objects administered in the current study, urbanisation was significantly and positively related to the likelihood of foxes touching, but not exploiting, them. Thus, while urban foxes may be bolder than rural populations in terms of their willingness to physically touch novel food-related objects, our findings are inconsistent with the notion that they are more innovative and pose a general nuisance to people by exploiting them.

Given that we were able to determine the fate of most objects, this rules out the possibility that our cameras significantly missed footage of foxes visiting and exploiting their contents without us knowing it. Foxes always consumed the food rewards when objects were absent despite the presence of a trail camera, ruling out the possibility that the cameras, rather than the food-related objects, were a significant deterrent for them. Since foxes consumed the food rewards when objects were absent, it also rules out the possibility that food-related motivation explains why foxes avoided the objects.

As previously discussed, studies in other species show urban-dwelling animals are more likely than rural populations to physically touch and gain access to novel food-related opportunities (Dammhahn et al., 2020; Ducatez et al., 2017; Griffin et al., 2017; Mazza et al., 2021; Mazza & Guenther, 2021). However, our findings – along with others – illustrate that the relationship between bold and innovative behaviour, particularly with regards to urbanisation, is complex and difficult to generalise across all situations and species (Griffin et al., 2017; Vincze & Kovacs, 2022). Indeed, many other factors likely contribute to whether or how wildlife can adapt to such environments (e.g., dispersal, morphology, and dietary generalist) (Thompson et al., 2021). These studies show, for example, that animals are more innovative in urban environments (field mice, *Apodemus agrarius*; Mazza & Guenther 2021), more innovative in rural environments (spotted hyena: *Crocuta crocuta*; Johnson-Ulrich et al. 2021),

or equally innovative in both (this study; house sparrows: *Passer domesticus*; Papp et al. 2014). Thus, for now, although our study suggests that urbanisation may somehow favour (for whatever reason) bolder behaviour in foxes, such behaviour does not necessarily favour them using innovation to exploit food-related opportunities in all contexts (Griffin et al., 2017). Indeed, if that was the case, then more than just 12 (out of 96) urban foxes should have exploited the objects that were administered in the current study after they were discovered.

There are multiple key factors that may separate bold and innovative behaviour. Evidence from birds, at least, suggests that species that are habitat generalists are better at incorporating novel food into their diet, while dietary generalists are more innovative in terms of how they physically acquire food (Ducatez et al., 2015). Red foxes are both habitat and dietary generalists, so it is unclear whether we would predict greater boldness or greater innovation. Our data suggest that boldness is the key behavioural trait; foxes, regardless of location, always consumed food rewards when objects were absent, but not when objects were present. Object neophobia might explain why some foxes avoided the food-related objects (Greggor et al., 2015; Miller et al., 2022; Travaini et al., 2013). Alternatively, given that food resources in urban environments are also very abundant (Ansell, 2005; Contesse et al., 2004b; Harris, 1981), this could explain why urban foxes were motivated to touch, but not necessarily persist and exploit, the unfamiliar food-related objects used in our study. Finally, individual characteristics such as age, sex, dominance, learning speed, and personality might have contributed to fox decision-making and are therefore worth investigating in the future (Fawcett et al., 2017; Griffin et al., 2013; F. B. Morton et al., 2021; Padovani et al., 2021; Soulsbury et al., 2011).

Despite being labelled as a pest, foxes remain a beloved part of urban fauna across the world (Baker & Harris, 2007; Baker et al., 2020; Brand & Baldwin, 2020; Konig, 2008; Nardi et al., 2020), and so future management needs to balance the co-occurrence of both positive and negative human-wildlife interactions within cities (Soulsbury & White, 2015). Our results contrast the UK popular culture's portrayal of urban foxes as a general 'nuisance' because they exploit food-related objects. Such beliefs may stem from specific, highly publicised cases

or provocative imagery rather than being typical of urban foxes in general. Indeed, most

household surveys (Baker et al., 2004; Harris, 1981), dietary studies (Contesse et al., 2004a),

and direct observations (Plumer et al., 2014) show that the image of foxes foraging from bins

is uncommon, rather than the norm. Even in our study, most foxes were unlikely to exploit

objects when the rewards were relatively large (e.g., 90 dog biscuits). By contrast, our findings

from the "free food" condition as well as other studies (Gil-Fernandez et al., 2020) show that

when anthropogenic resources are more easily accessible (e.g., no physical barriers), urban

foxes may be more likely to exploit such opportunities, which could be due to minimal effort,

risk, or both. We suggest that public perceptions of urban foxes stem from their use of freely-

available resources, rather than their innovative ability to access unfamiliar resources that

require effort.

Conclusions

Red foxes thrive within urban settings, but contrary to what has been observed in some

species, we found that wild urban foxes are, for the most part, no more likely than rural

populations to take advantage of novel food-related objects. Thus, while urban foxes may be

bolder than rural populations in terms of their willingness to physically touch novel food-related

objects, they do not always use innovation to exploit them. The low exploitation rate of food-

related objects found in the current study is also contrary to the notion that urban foxes pose

a general nuisance to people by exploiting these anthropogenic resources, and therefore calls

for a more nuanced view of urban fox behaviour, particularly when it comes to opportunistic

foraging.

References

Angel, S., Parent, J., Civco, D. L., & Blei, A. M. (2011). *Making room for a planet of cities*.
Lincoln Institute of Land Policy.

Ansell, R. J. (2005). *The spatial organisation of a red fox (Vulpes vulpes) population in*
relation to food resources [Doctoral dissertation, University of Bristol].

ASAB. (2020). Guidelines for the treatment of animals in beahvioural research and teaching.
Animal Behaviour, 159, i-xi.
https://doi.org/https://doi.org/10.1016/j.anbehav.2019.11.002

Baker, P., Funk, S., Harris, S., Newman, T., Saunders, G., & White, P. (2004). The impact of human attitudes on the social and spatial organization of urban foxes (Vulpes vulpes) before and after an outbreak of sarcoptic mange. *Proceedings 4th International Urban Wildlife Symposium* 153-163.

Baker, P. J., & Harris, S. (2007). Urban mammals: what does the future hold? An analysis of the factors affecting patterns of use of residential gardens in Great Britain. *Mammal Review, 37*, 297-315.

Baker, S. E., Maw, S. A., Johnson, P. J., & Macdonald, D. W. (2020). Not in my backyard: Public perceptions of wildlife and 'pest control' in and around UK homes, and local authority 'pest control' *Animals, 10*, 222.

Baston, D. (2021). exactextractr: Fast extraction from raster datasets using polygons.

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software, 67*(1), 1-48. https://doi.org/10.18637/jss.v067.i01

Brand, A., & Baldwin, M. (2020). Public attitudes to urban foxes in London and the south east. *Mammal Communications, 6*, 34-41.

Chao, A., Chiu, C.-H., & Jost, L. (2014). Unifying species diversity, phylogenetic diversity, functional diversity, and related similarity and differentiation measures through hill numbers. *Annual Review of Ecology, Evolution, and Systematics, 45*, 297-324.

Charmantier, A., Demeyrier, V., Lambrechts, M., Perret, S., & Grégoire, A. (2017). Urbanization is associated with divergence in pace-of-life in great tits. *Frontiers in Ecology and Evolution, 5*. https://doi.org/https://doi.org/10.3389/fevo.2017.00053

Contesse, P., Hegglin, D., Gloor, S., Bontadina, F., & Deplazes, P. (2004a). The diet of urban foxes (Vulpes vulpes) and the availability of anthropogenic food in the city of Zurich, Switzerland. *Mammalian Biology, 2*, 81-95.

Contesse, P., Hegglin, D., Gloor, S., Bontadina, F., & Deplazes, P. (2004b). The diet of urban foxes (Vulpes vulpes) and the availability of anthropogenic food in the city of Zurich, Switzerland *Mammalian Biology, 2*, 81-95.

Dammhahn, M., Mazza, V., Schirmer, A., Gottsche, C., & Eccard, J. A. (2020). Of city and village mice: behavioural adjustments of striped field mice to urban environments. *Scientific Reports, 10*, 13056.

DEFRA. (2022). *ENV23 - UK statistics on waste*. Retrieved 23 October, 2022 from https://www.gov.uk/government/statistical-data-sets/env23-uk-waste-data-and-management

Doncaster, C. P., Dickman, C. R., & Macdonald, D. (1990). Feeding ecology of red foxes (Vulpes vulpes) in the city of Oxford, England. *Journal of Mammology, 71*, 188.

Ducatez, S., Audet, J. N., Rodriguez, J. R., Kayello, L., & Lefebvre, L. (2017). Innovativeness and the effects of urbanization on risk-taking behaviors in wild Barbados birds. *Animal Cognition, 20*, 33-42.

Ducatez, S., Clavel, J., & Lefebvre, L. (2015). Ecological generalism and behavioural innovation in birds: technical intelligence or the simple incorporation of new foods? *Journal of Animal Ecology, 84*(1), 79-89. https://doi.org/https://doi.org/10.1111/1365-2656.12255

Fawcett, J. K., Fawcett, J. M., & Soulsbury, C. D. (2017). Seasonal and sex-specific differences in feeding site attendance by red foxes Vulpes vulpes. *Mammal Study, 42*(2), 117-120. https://doi.org/https://doi.org/10.3106/041.042.0207

Fischer, J. D., Schneider, S. C., Ahlers, A. A., & Miller, J. R. (2015). Categorizing wildlife responses to urbanization and conservation implications of terminology. *Conservation Biology, 29*, 1246-1248.

Fox, J., & Weisberg, S. (2019). *An {R} companion to applied regression* (Third ed.). Sage. https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Gil-Fernandez, M., Harcourt, R., Newsome, T., Towerton, A., & Carthey, A. (2020). Adaptations of the red fox (Vulpes vulpes) to urban environments in Sydney, Australia. *Journal of Urban Ecology, 6*, 1-9.

Greggor, A. L., Thornton, A., & Clayton, N. S. (2015). Neophobia is not only avoidance: improving neophobia tests by combining cognition and ecology. *Current Opinion in Behavoral Sciences, 6*, 82-89.

Griffin, A. S., Diquelou, M., & Perea, M. (2014). Innovative problem solving in birds: a key role of motor diversity. *Animal Behaviour, 92*, 221-227. https://doi.org/http://dx.doi.org/10.1016/j.anbehav.2014.04.009

Griffin, A. S., Guez, D., Lermite, F., & Patience, M. (2013). Tracking changing environments: innovators are fast, but not flexible learners. *PLoS ONE, 8*(12), e84907.

Griffin, A. S., Netto, K., & Peneaux, C. (2017). Neophilia, innovation and learning in an urbanised world: a critical evaluation of mixed findings. *Current Opinion in Behavoral Sciences, 16*, 15-22.

Grimm, N. B., Faeth, S. H., Golubiewski, N. E., Redman, C. L., Wu, J., Bai, X., & Briggs, J. M. (2008). Global change and the ecology of cities. *Science 319*, 756-760.

Guiden, P. W., Bartel, S. L., Byer, N. W., Shipley, A. A., & Orrock, J. L. (2019). Predator–
prey interactions in the Anthropocene: reconciling multiple aspects of novelty. *Trends*
in Ecology & Evolution, 34, 616-627.

Harris, S. (1981). The food of suburban foxes (Vulpes vulpes), with special reference to
London. *Mammal Review, 11*, 151-168.

Hill, M. O. (1973). Diversity and evenness: A unifying notation and its consequences.
Ecology, 54, 427-432.

Hollis, D., McCarthy, M., Kendon, M., Legg, T., & Simpson, I. (2019). HadUK-Grid - A new
UK dataset of gridded climate observations. *Geoscience Data Journal, 6*, 151-159.

Hollister, J., Shah, T., Robitaille, A. L., Beck, M. W., & Johnson, M. (2021). elevatr: Access
elevation data from various APIs. https://doi.org/doi:10.5281/zenodo.5809645

Horn, J. L. (1965). A rationale and test for the number of factors in factor analysis.
Psychometrika, 30, 179-185.

Johnson-Ulrich, L., Yirga, G., Strong, R. L., & Holekamp, K. E. (2021). The effect of
urbanization on innovation in spotted hyenas. *Animal Cognition, 24*, 1027-1038.

Konig, A. (2008). Fears, attitudes, and opinions of suburban residents with regards to their
urban foxes: a case study in the community of Grunwald - a suburb of Munich.
European Journal of Wildlife Research, 54, 101-109.

Lee, P. C., & Moura, A. (2015). Necessity, unpredictability and opportunity: An exploration of
ecological and social drivers of behavioral innovation. In A. B. Kaufman & J. C.
Kaufman (Eds.), *Animal Creativity and Innovation* (pp. 317-330). Academic Press.

Martin, P. R., & Bonier, F. (2018). Species interactions limit the occurrence of urban-adapted
birds in cities. *Proceedings of the National Academy of Sciences of the United States*
of America, 115, E11495-E11504.

Mazza, V., Czyperreck, I., Eccard, J. A., & Dammhahn, M. (2021). Cross-context responses
to novelty in rural and urban small mammals. *Frontiers in Ecology and Evolution, 9*,
661971.

Mazza, V., & Guenther, A. (2021). City mice and country mice: innovative problem solving in
rural and urban noncommensal rodents. *Animal Behaviour, 172*, 197-210.

Miller, R., Lambert, M. L., Frohnwieser, A., Brecht, K. F., Bugnyar, T., Crampton, I., Garcia-
Pelegrin, E., Gould, K., Greggor, A. L., Izawa, E.-I., Kelly, D. M., Li, Z., Luo, Y.,
Luong, L. B., Massen, J. J. M., Nieder, A., Reber, S. A., Schiestl, M., Seguchi, A.,

Sepehri, P., Stevens, J. R., Taylor, A. H., Wang, L., Wolff, L. M., Zhang, Y., & Clayton, N. S. (2022). Socio-ecological correlates of neophobia in corvids. *Current Biology, 32*, P74-85.

Morton, C. S., Marston, R. D., O'Neil, C. G., & Rowland, A. W. (2021). Land Cover Map 2020 (10m classified pixels, GB). https://doi.org/doi:10.5285/35c7d0e5-1121-4381-9940-75f7673c98f7

Morton, F. B. (2021). Do wild raccoons (*Procyon lotor*) use tools? *Animal Cognition, 24*, 433-441. https://doi.org/https://doi.org/10.1007/s10071-020-01430-y

Morton, F. B., & Altschul, D. (2019). Data reduction analyses of animal behaviour: avoiding Kaiser's criterion and adopting more robust automated methods. *Animal Behaviour, 149*, 89-95.

Morton, F. B., Buchanan-Smith, H. M., Brosnan, S. F., Thierry, B., Paukner, A., Essler, J. L., Marcum, C. S., & Lee, P. C. (2021). Studying animal innovation at the individual level: A ratings-based assessment in capuchin monkeys (Sapajus [Cebus] sp.). *Journal of Comparative Psychology, 135*(2), 258-265. https://doi.org/https://doi.org/10.1037/com0000264

Mu, H., Li, X., Wen, Y., Huang, J., Du, P., Su, W., Miao, S., & Geng, M. (2022). A global record of annual terrestrial human footprint dataset from 2000 to 2018. *Scientific Data, 176*. https://doi.org/https://doi.org/10.1038/s41597-022-01284-8

Murray, M., Cembrowski, A., Latham, A. D. M., Lukasik, V. M., Pruss, S., & St Clair, C. C. (2015). Greater consumption of protein-poor anthropogenic food by urban relative to rural coyotes increases diet breadth and potential for human–wildlife conflict. *Ecography, 38*, 1235-1242.

Murray, M. H., Kidd, A. D., Curry, S. E., Hepinstall-Cymerman, J., Yabsley, M. J., Adams, H. C., Ellison, T., Welch, C. N., & Hernandez, S. M. (2018). From wetland specialist to hand-fed generalist: shifts in diet and condition with provisioning for a recently urbanized wading bird. *Philosophical Transactions of the Royal Society B: Biological Sciences, 373*, 20170100.

Nardi, A., Shaw, B., Brossard, D., & Drake, D. (2020). Public attitudes toward urban foxes and coyotes: the roles of perceived risks and benefits, political idealology, ecological wordview, and attention to local news about urban wildlife. *Human Dimensions of Wildlife, 25*, 405-420.

OPENGREESPACE, O. (2022). *Contains OS data © Crown Copyright*.

Padovani, R., Shi, Z., & Harris, S. (2021). Are British urban foxes (Vulpes vulpes) "bold"? The importance of understanding human-wildlife interactions in urban areas. *Ecology and Evolution, 11*, 835-851.

Papp, S., Vincze, E., Preiszner, B., Liker, A., & Bokony, V. (2014). A comparison of problem-
solving success between urban and rural house sparrows. *Behavioral Ecology and*
Sociobiology, 69.

Pebesma, E. (2018). Simple features for R: Standardized support for statistical vector data.
The R Journal, 10, 439-446.

Pedroso-Santos, F., & Costa-Campos, C. E. (2020). Novel predator-prey interactions of
Rhinella major (Anura: Bufonidae) from an urban area in northern Brazil.
Phyllomedusa: Journal of Herpetology, 19, 287-292.

Planet dump. (2022). https://planet.osm.org

Plumer, L., Davison, J., & DSaarma, U. (2014). Rapid urbanization of red foxes in Estonia:
distribution, behaviour, attacks on domestic animals, and health-risks related to
zoonotic diseases. *PLoS ONE, 9*, e115124.

Reader, S. M., & Laland, K. N. (2003). *Animal innovation*. Oxford University Press. Table of
contents only http://www.loc.gov/catdir/toc/fy044/2004296817.html

Publisher description http://www.loc.gov/catdir/enhancements/fy0620/2004296817-d.html

Reis, S., Liska, T., Steinle, S., Carnell, E., Leaver, D., Roberts, E., Vieno, M., Beck, R., &
Dragosits, U. (2017). UK gridded population 2011 based on Census 2011 and Land
Cover Map 2015. NERC Environmental Information Data Centre.
https://doi.org/https://doi.org/10.5285/0995e94d-6d42-40c1-8ed4-5090d82471e1

Rodrigo-Comino, J., Seeling, S., Seeger, M. K., & Ries, J. B. (2021). Light pollution: A review
of the scientific literature. *The Anthropocene Review*.
https://doi.org/https://doi.org/10.1177/20530196211051209

Rossler, T., Mioduszewska, B., O'Hara, M., Huber, L., Prawiradilaga, D. M., & Auersperg, A.
M. I. (2020). Using an innovation arena to compare wild-caught and laboratory
Goffin's cockatoos. *Scientific Reports, 10*, 8681.
https://doi.org/https://doi.org/10.1038/s41598-020-65223-6

Šálek, M., Drahníková, L., & Tkadlec, E. (2015). Changes in home range sizes and
population densities of carnivore species along the natural to urban habitat gradient.
Mammal Review, 45, 1-14.

Saunders, G., & Harris, S. (2000). Evaluation of attractants and bait preferences of captive
red foxes (Vulpes vulpes). *Wildlife Research, 27*, 237-243.

Saunders, G., White, P. C. L., Harris, S., & Rayner, J. M. V. (1993). Urban foxes (Vulpes
vulpes): food acquisition, time and energy budgeting of a generalized predator.
Symposium of the Zoological Society of London, 65, 215-234.

785
786 Schell, C. J., Stanton, L. A., Young, J. K., Angeloni, L. M., Lambert, J. E., Breck, S. W., &
787   Murry, M. H. (2021). The evolutionary consequences of human-wildlife conflict in
788   cities. *Evolutionary Applications, 14*, 178-197.

789
790 Sol, D., González-Lagos, C., Moreira, D., Maspons, J., & Lapiedra, O. (2014). Urbanisation
791   tolerance and the loss of avian diversity. *Ecology Letters, 17*, 942-950.

792
793 Soulsbury, C. D., Baker, P. J., Iossa, G., & Harris, S. (2010). Red foxes (Vulpes vulpes). In
794   S. D. Gehrt, S. P. D. Riley, & B. L. Cypher (Eds.), *Urban carnivores: Ecology, conflict,*
795   *and conservation* (pp. 63-75). John Hopkins University Press.

796
797 Soulsbury, C. D., Iossa, G., Baker, P. J., White, P. C. L., & Harris, S. (2011). Behavioral and
798   spatial analysis of extraterritorial movements in red foxes (Vulpes vulpes). *Journal of*
799   *Mammology, 92*(1), 190-199. https://doi.org/10.1644/09-MAMM-A-187.1

800
801 Soulsbury, C. D., & White, P. (2015). Human-wildlife interactions in urban areas: A review of
802   conflicts, benefits and opportunities. *Wildlife Research, 42*, 541-553.

803
804 Team, R. (2002). *RStudio: Integrated Development for R.* In

805
806 Team, R. C. (2021). *R: A language and environment for statistical computing.* In R
807   Foundation for Statistical Computing https://www.R-project.org/

808
809 Team, R. C. (2022). *R: A language and environment for statistical computing.* In

810
811 Thompson, M. J., Capilla-Lasheras, P., Dominoni, D. M., Reale, D., & Charmantier, A.
812   (2021). Phenotypic variation in urban environments: mechanisms and implications.
813   *Trends in Ecology & Evolution, 37*, 171-182.

814
815 Thornton, A., & Samson, J. (2012, Jun). Innovative problem solving in wild meerkats. *Animal*
816   *Behaviour, 83*, 1459-1468. <Go to ISI>://WOS:000305082200020

817
818 Travaini, A., Vassallo, A. I., Garcia, G. O., Echeverria, A. I., Zapata, S. C., & Nielsen, S.
819   (2013). Evaluation of neophobia and its potential impact upon predator control
820   techniques: A study on two sympatric foxes in southern Patagonia. *Behavioural*
821   *Processes, 92*, 79-87.

822
823 Trewhella, W. J., Harris, S., & McAllister, F. E. (1988). Dispersal distance, home-range size
824   and population density in the red fox (Vulpes vulpes): A quantitative analysis. *Journal*
825   *of Applied Ecology, 25*, 423-434. https://doi.org/https://doi.org/10.2307/2403834

826
827   UN. (2018). *Urbanisation prospectus 2018*. https://population.un.org/wup/Download/

828
829   Vincze, E., & Kovacs, B. (2022). Urbanisation's effects on problem solving abilities: A meta-
830   analysis. *Frontiers in Ecology and Evolution, 10*, 834436.

831
832   Visalberghi, E., & Limongelli, L. (1994). Lack of comprehension of cause-effect relations in
833   tool-using capuchin monkeys (*Cebus apella*). *Journal of Comparative Psychology,*
834   *108*(1), 15-22. https://doi.org/https://doi.org/10.1037/0735-7036.108.1.15

835
836   Webster, M. M., Ward, A. J. W., & Hart, P. J. B. (2009). Individual boldness affects
837   interspecific interactions in sticklebacks. *Behavioral Ecology and Sociobiology, 63*,
838   511-520.

839
840   Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., Francois, R., Grolemund, G.,
841   Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M.,
842   Muller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., Takahashi, K., Vaughan,
843   D., Wilke, C., Woo, K., & Yutani, H. (2019). Welcome to the tidyverse. *Journal of*
844   *Open Source Software, 4*, 1686.