Optogenetic control of YAP can enhance the rate of wound healing

P. J. Y. Toh[1§], M. Sudol[1,2] and T. E. Saunders[1,3,4*]

[1] Mechanobiology Institute, National University of Singapore, Singapore

[2] Icahn School of Medicine at Mount Sinai, New York City, USA

[3] Institute of Molecular and Cell Biology, A*STAR, Singapore

[4] Warwick Medical School, University of Warwick, United Kingdom

§ Current address: Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and

Medical Sciences, University of Copenhagen, Denmark

* For correspondence: T.E.S. (timothy.saunders@warwick.ac.uk)

Abstract

Tissues need to regenerate to restore function after injury. Yet, this regenerative capacity varies

significantly between organs and between species. For example, in the case of the heart, while

some species retain full regenerative capacity throughout their lifespan, human cardiac cells

display only limited ability to repair injury. Here, we investigate in cell culture the role of the

YAP, a transcriptional co-regulator with a pivotal role in growth, in driving repair after injury.

Utilising an optogenetic version of YAP that enables precise control of pathway activation, we

show that YAP can increase the speed of wound healing in H9c2 cardiomyoblasts. Interestingly,

this is not driven by an increase in proliferation, but by collective cell migration. We

subsequently dissect specific phosphorylation sites in YAP to identify the molecular driver of

accelerated healing. Overall, our results reveal that YAP activation – through controlled

optogenetic activation - can potentially enhance wound healing in a range of conditions.

Keywords: Hippo-YAP, optogenetics, wound healing

Introduction

During myocardial infarction, blood flow to the heart is severely limited or completely cut off by plaque or cholesterol blockages. Without oxygen and nutrients, heart muscles surrounding the region of the blocked vessel begin to necrose. Acute myocardial injury can kill 25% of cardiomyocytes, rendering them non-functional.(1) Pharmacological management after a heart attack only serves to prevent another episode from occurring and there are no known strategies to restore the full capacity of the heart. This in part due to the human heart having a very limited ability to regenerate post injury.(2) This lack of regenerative capability is in contrast to other organs, such as the liver or gut.(3) Further, other organisms show a greater capacity to regenerate injured heart tissue post-embryonic development: young mice (<7 days) can repair heart wounding; and in zebrafish the potential to regenerate heart tissue remains throughout the lifespan.(4,5) Therefore there is a demand to find strategies to induce cardiac regeneration in humans. Though new cardiomyocytes can be synthesised *ex vivo* from embryonic stem cells and organise to form contractile tissues, injected cells have an embryonic lineage and can generate arrythmias.(6,7) Recipients also need to be immunosuppressed for transfusion, which itself is a complex procedure. Alternative strategies to induce human heart tissue repair and regeneration are still heavily sought after.

A strategy to circumvent the above complexity is to stimulate the endogenous proliferation potential of cells in repair. The key idea is to effectively mimic developmental processes, reviving pathways that are switched off postnatally. This approach has seen positive results in transgenic mice models, where activated YAP promotes cardiac regeneration by inducing reparative genetic programs to re-enter the cell cycle and activate proliferative genes in cardiomyocytes.(8–10) The Hippo-YAP pathway is highly implicated in development and

regenerative processes, due to its role in driving cell proliferation, migration and

mechanosensing.(11,12) The pathway is constructed principally from a kinase cascade that

culminates at the transcriptional co-activator YAP and its paralogue TAZ. Phosphorylated

YAP/TAZ triggers its cytoplasmic sequestration(13,14) or degradation,(15) while

unphosphorylated YAP/TAZ can translocate into the nucleus and bind to transcription factors

TEAD1-4.(16–18) This stimulates the expression of a range of genes that typically lead to pro-

survival outcomes in the cell.(19–21) The nucleocytoplasmic distribution of YAP/TAZ is a

key determinant of its activity. Therefore, manipulating the YAP/TAZ localisation can provide

a tool to control the subcellular activation of the Hippo-YAP pathway.

Previous studies have shown that the MAPK/ERK pathway is a key regulator of organ

regeneration(22,23), and crosstalk with the Hippo-YAP pathway in cardiomyocytes reactivates

cell proliferation.(24,25) ERBB2 (a member of the EGFR family) signalling induces a

phosphorylation mark on two non-canonical serine residues on YAP by ERK.(25) The

activating phosphorylation marks on S251 and S333 oppose the conventional deactivating

phosphorylation caused by LATS on S127. The phosphorylation mark at S127 generates a 14-

3-3 binding site, causing the cytoplasmic sequestration of phosphorylated YAP, accelerating

the nuclear export of YAP and inhibiting its co-transcriptional function inside the

nucleus.(26,27) Mutation of serine 127 to alanine has been reported to increase the activity of

YAP by preventing LATS phosphorylation.(28–30) Besides the phosphorylation site at S127,

another protein domain that regulates the localisation of YAP is the C-terminal PDZ-binding

motif,(31) a short motif of only five amino acid residues, -FLTWL at the very carboxyl terminal

end. The PDZ-binding motif allows YAP to bind to proteins containing the PDZ domain such

as ZO-2 and a deletion of this motif impairs the nuclear shuttling of YAP through ZO-2.(31)

However, similar to the S127A mutant, truncation of the PDZ-binding motif does not impair

the induction of TEAD-mediated transcriptional activity, only affecting its subcellular

localisation.(30,32) The YAP paralogue TAZ also contains a C-terminal PDZ-binding motif,

which has been shown to be required for its nuclear localisation.(14) Therefore, the PDZ-

binding motif appears to be a conserved sequence that is integral to the function of YAP/TAZ

as transcriptional co-regulators. However, how these phosphorylation marks and domains alter

the dynamics of YAP signalling – and subsequently its efficacy in regeneration – remain

unknown.

During regeneration, there are two main mechanisms for repair: (1) migration of newly

proliferated cells into the injured area; and (2) cell proliferation within the regenerating zone.

YAP is involved in wound healing, by limiting cytoskeletal and focal adhesion maturation to

promote cell motility.(33) YAP increases migration and invasiveness in a number of cell lines,

often stimulating epithelial-to-mesenchymal transition (EMT).(34) On top of enhancing cell

cycle progression, YAP also promotes cytoskeletal remodelling, responding to the mechanical

change during scar tissue formation post myocardial infarction.(35) YAP has also been shown

to be activated in cardiac fibroblasts after ischemic injury.(36)

While YAP activity has been shown to regenerate the mice heart after injury, caution must be

taken as over-proliferation can cause occlusive vascular diseases.(37) Therefore, having

control over the spatiotemporal localisation of YAP is important. Optogenetics provides such

a method,(38,39) whereby light activation can control protein behaviour, and we have recently

applied this to YAP (optoYAP).(40) Can optogenetic YAP be used to control regeneration after

wounding or injury? optoYAP is potentially a powerful tool to circumvent regenerative

limitations whilst retaining specific spatiotemporal control over its activity. Building upon our

previously reported work,(40) we utilised a range of cell lines, including H9c2 rat

cardiomyoblasts, as *in vitro* models to study how activation of YAP can promote repair post-

injury.[41,42]

Here, we report that activated optoYAP accelerates wound healing in two different cell lines,

driving cell migration that is essential during regeneration. In particular, we show that H9c2

cardiomyoblasts are responsive to optoYAP activation, inducing nuclear accumulation after

light activation. Nuclear optoYAP increases expression of downstream proliferative YAP

target genes *CTGF* and *CYR61*, as well as *TGF-β*, a transforming growth factor involved in

stem cell regulation and differentiation. This suggests that optoYAP can potentially be

applicable in regenerative studies of tissues – including cardiac tissues – by promoting the

renewal and replication of cells post-injury.

The Hippo-YAP pathway crosstalks with ERBB2 signalling during cardiac regeneration,

where ERK induces two activating phosphorylations (S251 and S333) on YAP.(25) To further

understand the potential mechanism for YAP function in regeneration, we generated

phosphomimetic mutants at both serine residues and observed that they had an elevated amount

of nuclear localisation in the dark as compared to wildtype optoYAP. However, the mutants

could still respond to light activation. Building upon the changes in basal distribution of mutant

optoYAP, we mutated the canonical LATS phosphorylation serine residue 127 to alanine,

preventing its inactivating phosphorylation effect on YAP. We observed that optoYAP S127A

was strongly nuclear localised before light activation and the nuclear signal was amplified after

light activation. By removing the C-terminal PDZ-binding motif, another regulatory domain of

YAP, we arrested the intrinsic nuclear localisation of YAP by preventing it from binding to

nuclear importins. However, optogenetic activation was still able to promote nuclear

localisation of this mutant through the NLS present on the optogenetic backbone. Overall, our

results reveal that optogenetic approaches provide a strategy for specific control over YAP

nuclear localisation (and hence activity) in both wildtype YAP and its mutants, thereby making

it a potential tool for driving tissue regeneration.

Results

optoYAP activity in H9c2 cells: Using a previously reported optoYAP for optogenetic control

of YAP nuclear import and export,[40] we transfected H9c2 cells with optoYAP and

characterised the construct in the rat cardiomyoblasts. We subjected transfected cells to

pulsatile light activation with 488 nm laser power at 0.15 mW for 1 second at a 30 second

interval, for a total of 20 minutes. We observed nuclear localisation of the optogenetic construct

after the 20 min activation window (Fig. 1A). After leaving the cells in the dark for a further

20 minutes, optoYAP translocated out of the nucleus and returned to the basal state before light

activation (Fig. 1A). The nuclear/cytoplasmic ratio of optoYAP almost doubled after the 20

min activation (Fig. 1B). Throughout the activation and recovery cycle, we tracked and imaged

the cells over time to obtain the nuclear import and export dynamics (Fig. 1B). The activation

(τ_a) and recovery (τ_r) time constants were calculated to be 4.6 ± 0.6 minutes and 13.5 ± 1.6

minutes respectively (Methods). These values are comparable with those obtained by using the

HEK293T cell line described previously.[40]

Elevated downstream gene expression levels after optoYAP activation: Following the light-

dependent nuclear localisation of optoYAP, we examined the functionality of nuclear localised

optoYAP by measuring the changes in downstream gene expression levels. We quantified the

transcript levels of *CTGF*, *CYR61*, and *TGF-β* in H9c2 cells after 48 hours of pulsatile light

activation (Fig. 1C). *CTGF* and *CYR61* are canonical YAP target genes that provide a readout

for optoYAP activity in the nucleus.(18,29) Additionally, we looked at *TGF-β* transcript levels

as it ©s an inflammation-related growth factor and is frequently involved in the migration and

invasion of tumour cells.(41) YAP has also been shown to be a mediator of *TGF-β*

signalling(42,43) and the pathway has been implicated in cardiomyocyte proliferation and

migration.(44,45)

We observed that the expression levels of all three genes probed were significantly upregulated

in cells containing activated optoYAP as compared to wildtype (WT) H9c2 (Fig. 1C). However,

the transcript levels of *CTGF* and *TGF-β* in transfected cells kept in the dark were also

significantly increased in comparison to wildtype controls, though clearly less than when light

activated. This suggests that there might be small amounts of nuclear optoYAP even without

light activation. However, the upregulation of gene expression across the three genes is

significant when comparing cells kept in the light versus dark. This shows that the optoYAP

construct is functional when nuclear localised and can trigger expression of downstream target

genes.

Figure 1. Characterisation of optoYAP in H9c2 cells. (A) Representative images of H9c2

cells transfected with optoYAP kept in the dark (t = 0 min), then subjected to pulsatile light

activation for 20 min (t = 20 min), followed by recovery in the dark for a further 20 min (t =

40 min). Scale bar, 20 μm. **(B)** Quantification of activation (τ_a) and recovery (τ_r) time constant

of optoYAP. Red line represents average nuclear/cytoplasmic ratio. Vertical dashed line

represents time when 488 nm stimulation ceased. Numbers in brackets represent the 95%

confidence interval of τ. n=12 cells. **(C)** RT-qPCR of YAP downstream genes. Cells were

transfected with optoYAP, and transcript levels were assessed after 48 h of activation.

Expression levels of *CTGF*, *CYR61*, and *TGF-β* were normalised to the housekeeping gene

EIF1B and described as relative fold change. Horizontal bars represent mean and 95%

confidence interval from 6 biological replicates from two independent experiments. NS: not

significant, $*P < 0.05$, $**P < 10^{-2}$, $***P < 10^{-3}$.

optoYAP promotes collective cell migration: We next investigated the role of optoYAP during wound healing, comparing endogenous and optogenetic YAP. Serum-starved cells were seeded in wells 500 μm apart in serum-free media and the""wound" size was quantified every 30 minutes until complete closure. We observed that optoYAP transfected H9c2 cells kept under pulsatile light activation migrated faster over the wound as compared to transfected cells kept in the dark or untransfected H9c2 (Fig. 2A). Wild type H9c2 cells took an average of 66 hours for complete wound closure, in comparison to 60 hours taken for transfected cells in the dark (Fig. 2B). Light activated optoYAP accelerated wound healing, taking an average of 53 hours, a 20% improvement on the wild type cells. Though optoYAP transfected H9c2 cells kept in the dark took a shorter time as compared to wildtype cells, light activation of the optogenetic construct clearly bolstered this behaviour. Extracting the curve slopes in Fig. 2B inset, we see that the activated optoYAP cells have significantly increased rate of closure.

We observed some cell proliferation events during the timelapse, despite the serum starvation before and during the experiment, seen as bright, rounded cell doublets in Fig. 2A. We counted the number of proliferation events that occurred during wound healing and observed that all three conditions had similar number of proliferation events (Fig. 2C). This shows that the differences in the time taken for wound healing observed in Fig. 2B is largely due to cell migration and not proliferation.

Figure 2. Wound healing assay in H9c2 cells. (A) Timelapse images of wildtype H9c2 cells

or cells transfected with optoYAP under different light conditions, taken every 30 min until

gap completely closes. Representative images shown at 10 h intervals. Scale bars, 100 μm. **(B)**

Time taken for H9c2 cells for complete wound healing. Quantification of data in (A) over time.

Bold lines represent mean and dashed lines represent linear regression fitted to the curve. n=10

from two independent experiments. Inset shows the measured slopes under different conditions relative to wildtype (colours as in main plot), error bars are s.d.. **(C)** Number of proliferation events counted during each wound healing experiment in (B). Bars represent mean and error bars represents s.d..

We performed the same wound healing experiment in another cell line, MKN28, to exploit the benefit of a CRISPR $YAP^{-/-}$ background previously reported.(46) This cell line enabled us to precisely delineate the effect of optoYAP without any endogenous YAP present that might confound the above results. Previous studies in the same cell line observed a difference between wildtype and $YAP^{-/-}$ cells in a trans-well migration assay but not during wound healing assays.(46) Therefore, we wanted to determine if our optoYAP construct changes the rate of wound closure in these MKN28 $YAP^{-/-}$ cells.

As expected, the $YAP^{-/-}$ cells took much longer than the wild type or optoYAP transfected cells for wound closure, taking 38 hours as compared to 33 hours for wild type MKN28 cells (Fig. 3A). This shows that endogenous YAP is required for cell motility during wound closure. In MKN28 $YAP^{-/-}$ cells transfected with optoYAP, we noticed that those kept in the dark took a faster time for wound closure (30.5 hours) in comparison to wild type cells. Light activated optoYAP transfected $YAP^{-/-}$ cells had an accelerated rate of wound closure, taking only 29 hours for the wound to close completely. This is supported by analysis of the slope of the curves (Fig. 3A inset). These results support the conclusion that endogenous YAP is involved in cell migration and optoYAP further accelerates wound healing in MKN28 cells. Globally, it was noteworthy that the MKN28 cells took a shorter time for wound closure (<40 hours) as compared to the H9c2 cells (>50 hours).

Figure 3. Wound healing assay in MKN28 variants. (A) Wildtype (WT) MKN28 and

MKN28 *YAP⁻ᐟ⁻* cells transfected with optoYAP were seeded in culture wells 500 μm apart. The

size of the gap was tracked every 30 min until it was completely closed. Bold lines represent

mean and dashed line represents linear regression fitted to curve. n=8 from two independent

experiments. Inset shows slope of lines relative to wildtype (colours as in main plot), with error

bars representing s.d.. **(B)** Number of proliferation events in each experiment in (A). Bars

represent mean and error bars represents s.d..

As with the H9c2 cells, we observed several proliferating MKN28 cells in the cell islands

during wound closure. We counted the number of proliferation events and compared it amongst

the MKN28 variants (Fig. 3B). Unsurprisingly, the *YAP⁻ᐟ⁻* variants had the least amount of

proliferation events (28 ± 7), less than half of the wild type cells (72 ± 6) over the entire time

course. Since YAP plays an important role in proliferation, knocking out YAP naturally

reduces cell proliferation, which might explain the longer time taken for wound closure in

YAP⁻ᐟ⁻ cells as compared to wild type. Both optoYAP transfected *YAP⁻ᐟ⁻* cells under light (45 ±

7) or dark (46 ± 6) conditions had similar number of proliferation events, but still less than wild

type cells. Despite less proliferation observed in optoYAP transfected *YAP⁻ᐟ⁻* cells, the time

taken for wound closure is shorter than wild type cells. This shows that optoYAP speeds up

the rate of wound closure in *YAP*^{-/-} cells, seemingly by inducing cell migration. However, cell

proliferation plays a crucial part in wound closure, and it is important to understand the fine

interplay between proliferation and migration as revealed here by using CRISPR/Cas9-

generated *YAP*^{-/-} MKN28 cells.(46)

Changes in nuclear localisation of optoYAP phosphorylation mutants: Above, we have shown

that YAP plays a role in wound healing, including in cardiomyocytes. Phosphorylation of YAP

on two ERK activating sites appears to be critical in this response,[26] but the dynamics remain

poorly understood. To explore this further, we synthesised three phosphomimetic optoYAP

mutants by substituting serine to aspartic acid on either S251 or S333, or both. We did this in

HEK293T cells due to their ease of transfection over H9c2 or MKN28 cells. We did not observe

distinct nuclear localisation differences amongst the three mutants (Fig. 4A). Comparing the

basal nuclear localisation of the three mutants, we observed that the double mutant had a

significantly higher initial nuclear localisation compared to the S251D mutant, but not the

S333D mutant (Fig. 4B). The mutants all displayed clear response to light activation (Fig. 4B-

D). They also displayed higher initial nuclear localisation over wildtype (Fig. 4E).

Figure 4. Point mutations of optoYAP at ERK phosphorylation serine residues. (A) Three

mutants containing phosphomimetic of ERK target residues at S251, S333 or both S251 and

S333. Representative images of HEK293T cells transfected with different mutant optoYAP

before and after exposure to pulsatile light activation. Scale bars, 10 μm. **(B)** Quantification of

cells from (A), measuring nuclear to cytoplasmic ratio of optoYAP before and after light

activation for each mutant. NS: not significant, ***$P < 10^{-3}$. **(C)** Fraction increase of N/C ratio

after light activation. Box plots represent median and 25th to 75th percentiles. Whiskers show

minimum and maximum points. **(D)** Relative N/C ratio over time during 488 nm activation

amongst optoYAP mutants.©**)** Absolute intensity values of mCherry signal in nuclei over time.

Solid lines represent mean, n=10 cells taken from three independent experiments.

We then looked at canoncial YAP regulatory sites, such as serine 127 and the C-terminal PDZ-

binding motif. We observed that the S127A mutant (referred to as optoYAP S127A) is nuclear

localised even before light activation, contrasting significantly with the wild-type cells (Fig.

5A-B). After light activation, we observed that optoYAP S127A accumulated further within

the nucleus, doubling the nuclear to cytoplasmic ratio of optoYAP S127A as compared to the

dark state (Fig. 5D). This shows that the optoYAP S127A mutant can exert its predominantly

nuclear state but optogenetic activation can intensify this ability and increase optoYAP levels

in the nucleus.

Deletion of the PDZ-binding motif on YAP causes complete YAP exclusion from the

nucleus.(32) We synthesised a truncated optoYAP without the PDZ-binding motif, which we

refer to as optoYAP ΔC. We observed nuclear exclusion of optoYAP ΔC before exposure to

light activation (Fig. 5C). Pulsatile light activation can induce nuclear localisation of optoYAP

ΔC, nearly doubling the amount of nuclear localised optoYAP ΔC. The optoYAP ΔC construct

returns to the cytoplasm after 20 minutes of recovery in the dark, similar to the wildtype and

S127A mutant. Therefore, activation of the optogenetic tool can overcome mutations that drive

YAP out of the nucleus.

Figure 5. optoYAP variants under light activation protocol. Representative images of

mCherry signal in HEK293T cells containing **(A)** wildtype (WT) optoYAP, **(B)** optoYAP with

a point mutation at S127A, or **(C)** truncated optoYAP without the C-terminal PDZ-binding

motif (optoYAP ΔC). The 'Pre activation' panels show cells before exposure to 488 nm light,

and after 20 min of pulsatile activation with 488 nm light, the same cells were imaged as 'Post

activation'. Cells were left in the dark for a 20 min and imaged again, under the 'Recovery'

panel. **(D)** Quantification of nuclear to cytoplasmic ratio of optoYAP in (A-C). Box plots

represent median and 25th to 75th percentiles. Whiskers show minimum and maximum points.

Scale bars, 10 μm. n=22 cells for (A), n=10 cells for (B) and (C), taken from three independent

experiments. *$P < 0.05$, ***$P < 10^{-3}$.

Discussion

Here, we report that optoYAP can accelerate wound healing by promoting cell proliferation and migration in different cell culture systems. We show that activation of YAP can increase the rate of wound repair by 20% in cardiomyocytes. optoYAP, in the absence of endogenous YAP in MKN28 cells, promotes wound healing, accelerating the time taken for closure (Fig. 3A). However, we observed smaller differences in the H9c2 cells between the wildtype and the optoYAP transfected cells (Fig. 2A-B). We further explored the mechanisms that determine the YAP nuclear-cytoplasmic localisation. We demonstrate that the defects in YAP nucleus localisation by perturbation to the S127A residue and C-terminal PDZ-binding motif can be compensated for through use of optoYAP activation. Taken together, our results provide strong evidence that: (i) YAP can play an active role in repair beyond simply proliferation; and (ii) optoYAP can regulate YAP activity, even in the presence of different perturbations to YAP, which may be important for wound repair in a range of cell types.

By counting the number of cell proliferation events during wound closure, we observed a few differences between the MKN28 and H9c2 cell lines. We saw that optoYAP increases the amount of cell proliferation in serum starved MKN28 *YAP*$^{-/-}$ cells, but not to the wild-type levels (Fig. 3B). However, in MKN28 *YAP*$^{-/-}$ cells, the lack of endogenous *YAP* reduces the amount of proliferation in these cells. Despite transfection with optoYAP, the levels of YAP present in MKN28 *YAP*$^{-/-}$ cells might not be the same as endogenous YAP in wild type cells. Therefore, this might explain the differences between the amount of proliferation in the MKN28 variants. On the contrary in H9c2 cells, activated optoYAP does not alter the amount of proliferation in transfected cells (Fig. 2C). This suggests that serum starvation effectively reduces cell proliferation globally. Overall, both cell lines with transfected optoYAP kept in

the dark had comparable numbers of proliferation to their light activated counterparts. This

shows that despite having similar cell proliferation, light activated optoYAP speeds up the time

required for wound closure in both cell lines by enhancing cell migration.

Multiple signalling pathways regulate YAP independently of Hippo, including the epidermal

growth factor receptor (EGFR) signalling pathway.(47) Downstream of EGFR lie the

Ras/Raf/MEK/ERK transduction axis, where ERK is the final kinase activator that regulates

transcription factors. ERK is a key regulator of regeneration and interplay between ERK and

YAP has been shown to be drive YAP activation. By synthesising phosphomimetics of the

ERK phosphorylation sites, we observed some nuclear localisation of optoYAP without light

activation. Nuclear localisation of YAP phosphorylated at both serine residues have been

observed, though a single phosphorylation mark at either S251 or S333 is sufficient to induce

nuclear translocation.(48,49) However, the double mutant is significantly more nuclear

localised than the single mutants (Fig. 4B), which shows the additive effect of phosphorylating

both serine residues. Despite the differences observed amongst the three optoYAP mutants, it

appears that mutating optoYAP at the two ERBB2 activating phosphorylation sites does not

significantly increase nuclear localisation of the optoYAP construct as compared to the

wildtype variant.

Additionally, we looked at canonical YAP regulatory sites, S127 and the PDZ-binding motif,

that changes the cellular distribution of YAP. Despite all three mutants having an increase in

nuclear localisation after light activation, it is evident that the basal level of nuclear localisation

is not equal. The optoYAP S127A mutant (0.68 ± 0.13) has a significantly higher initial nuclear

concentration than wild type (0.44 ± 0.09), since the S127A mutation causes optoYAP to be

predominantly nuclear. Conversely, the optoYAP ΔC mutant (0.36 ± 0.09) has a significantly

lower nuclear localisation compared to wild type (0.44 ± 0.09), which is expected as deletion

of the PDZ-binding motif prevents optoYAP from translocating into the nucleus.(32) Although

the basal levels differ for the three variants, nuclear localisation doubles for all optoYAP

variants and decreases back to their respective basal levels after recovery in the dark (Fig. 5D).

This shows that the optogenetic domains can promote further nuclear translocation despite

mutations that already boosts (S127A) or inhibit (ΔC) nuclear localisation.

The rate of nuclear localisation of the two S127A and ΔC optoYAP mutants have yet to be

studied, which could shed light on the dynamics of nuclear shuttling in the optoYAP mutants.

The rate of nuclear import or export might differ drastically between the optoYAP variants,

depending on the mutation. We hypothesise that the activating S127A mutation increases the

rate of nuclear import but does not affect nuclear export rates, while the negative ΔC mutant

increases the rate of nuclear export but maintains similar import rates as WT. Regardless, it is

evident that mutations in the S127 or PDZ-binding motif changes the initial localisation of

optoYAP, but the activated optogenetic construct can supersede the basal distribution and

induce optoYAP nuclear localisation, nonetheless.

Finally, we note that several optogenetic YAP constructs have been published recently, with

different YAP isoforms and activation protocols. Illes *et al*. first published a tool to control the

subcellular localisation of YAP by tagging it with an optogenetic NLS, shuttling YAP between

the cytoplasm and the nucleus.(50) The authors used the human YAP1 gene without specifying

the details of the splice isoform, and demonstrated that activated YAP increases proliferation

in 2D and 3D cultures, and induces invasion in cancer spheroids.(50) Dowbaj *et al.* presented

optogenetic YAP1, using the human YAP1-2γ isoform,(51) which contains exon 6 and two

WW domains,(52) tethering optogenetic YAP and TAZ to the mitochondria. Their tool allows

the study of protein dynamics by describing the rate of nuclear entry and exit, but does not

demonstrate downstream functionality of nuclear YAP.(51) In our previous work, we reported

a human YAP1-1δ optogenetic tool that translocates between the cytoplasm and nucleus,

demonstrating functional outputs *in vitro* and *in vivo*.(40) Most recently, Meyer *et al.* have

developed a mouse YAP1 tool. They are able to regulate the levels of pluripotency factors Oct4

and Nanog, and hence control the renewal of a naïve stem cell state or preferential

differentiation into the mesendodermal state.(53) The various YAP isoforms signal differently,

where those without exon 6, alpha and beta, can form heterodimers with TAZ.(54) Additionally,

different isoforms have unique and common functions and activate distinct downstream

transcriptional programmes.(55) Taken together, the isoforms of YAP matter when describing

optogenetic constructs, given that each isoform can have unique downstream effects.

The role of YAP in wound healing has been investigated previously, where nuclear localisation

of YAP is observed during healing.(56,57) We demonstrate that optoYAP can accelerate

wound healing as compared to endogenous YAP, providing a tight spatiotemporal control over

its activity. It has been shown that H9c2 cells respond to substrate stiffness and topography,

promoting cell differentiation on a modified ECM synthesized from fibroblast-derived

matrix.(58) With our optogenetic tool, it will be interesting to investigate cell migration in an

environment that mimics physiological conditions by seeding cells on permissive extracellular

matrix, or a stiffened matrix that imitates cholesterol plaques, and tracking wound healing

under such conditions, which will benefit future studies on regeneration.

Materials and Methods

Plasmid construction

Optogenetic plasmid containing YAP (optoYAP) was previously described by Toh et al., 2022.(40) Mutant YAP containing S127A or ΔC modifications were PCR amplified from pBABE (hygro) hYAP1-1δ S127A and pBABE (hygro) hYAP1-1δ ΔC plasmids respectively, both of which were from the library of YAP cDNA constructs of the Sudol laboratory. These constructs are available from 'Addgene' vector resource. Phusion High Fidelity DNA polymerase (ThermoScientific) was used for PCR amplification with primer sequences for each amplification listed in Table 1. PCR products and optoYAP plasmid were cleaved with appropriate FastDigest restriction enzymes (ThermoScientific) for cloning.

Point mutations in hYAP1-1δ were generated with Q5 Site-Directed Mutagenesis Kit (NEB) according to the manufacturer's instructions to synthesise phosphomimic mutants by substituting serine with aspartic acid at two residues, S251 and S333. Positive clones were verified by restriction enzyme digestion and sequencing. Primers used for site-directed mutagenesis (SDM) are listed in Table 1.

Mammalian cell culture and transfection

HEK293T human embryonic kidney cells were grown in DMEM (ThermoScientific) supplemented with 10% (v/v) heat inactivated fetal bovine serum (FBS, Gibco) and 1% (v/v) penicillin/streptomycin (Gibco). MKN28 human gastric adenocarcinoma cells were grown in RPMI (ThermoScientific) supplemented with 10% (v/v) FBS and 1% (v/v) penicillin/streptomycin. H9c2 rat cardiomyocytes (a gift from Dr Manvendra K. Singh's lab)

were grown in DMEM supplemented with 15% (v/v) FBS, 1% (v/v) penicillin/streptomycin

and 1% non-essential amino acids (NEAA, Gibco). All cell cultures were kept in a 37ºC

incubator with 5% CO_2 and passaged every 2-3 days. Cells were routinely checked for

mycoplasma contamination and STR profiled (1st Base).

HEK293T cells were transfected with plasmid DNA using Lipofectamine 2000

(ThermoScientific) using 6 μL of Lipofectamine 2000 reagent in 150 μL opti-MEM

(ThermoScientific) and 3.5 μg of plasmid DNA in 175 μL opti-MEM according to the

manufacturer's instructions. H9c2 cells were transfected with Lipofectamine 2000 as per

HEK293T cells, but with 5 μg of plasmid DNA. MKN28 cells were transfected using K2

transfection system (Biontex).

Fluorescence microscopy

Live imaging was performed on Perkin Elmer spinning disk with a LUCPlanFLN 40x/0.6

numerical aperture (NA) air objective or Nikon Ti-E with Yokogawa W1 spinning disk with a

PLAN APO VC 60x/1.20 NA water immersion objective. Cells were visualised under the

microscope 24 h post transfection in a 37ºC heated and humidified chamber.

Long term imaging of wound healing experiments was performed on widefield Nikon

Biostation IMQ with 10x air objective or Olympus IX81 with a UPLFLN 10x/0.3 NA air

objective. The samples were tracked every 30 min to measure the rate of wound closure until

the gap was completely closed.

Image analysis

The images were analysed with Fiji ImageJ software. The nuclear to cytoplasmic ratio was

measured by drawing a region of interest (ROI) around the cell as well as its nucleus. A macro

plugin written by Richard De Mets was used to mask the nucleus from the cytoplasm to

calculate the mean intensity in the nucleus and cytoplasm based on the ROI drawn. The ROI

was drawn for each cell analysed at every time frame. The wound healing analysis was

performed using a macro written by Ong Hui Ting, which measures the changes in gap area of

the wound over time. Charts were made with Prism 8 software. All figures were assembled

with Adobe Illustrator.

<u>Light activation</u>

Pulsed light activation followed a protocol of 1 s pulse of 0.15 mW 488 nm laser light followed

by a 30 s dark phase. This pulsation is continued for 20 min, which constitutes as 'light

activation'. During recovery, the entire set up is kept in the dark for a further 20 min.

Throughout the entire 40 min of light activation and recovery, imaging was done with a 561

nm laser line to image the mCherry fluorescent protein. For samples kept under dark conditions,

aluminium foil was used to wrap the sample to prevent light contamination.

<u>RT-qPCR</u>

Transfected cells were subjected to 48 h of pulsatile light activation and total RNA was

extracted from cells using RNeasy Plus Mini Kit (QIAGEN) and QIAshredder (QIAGEN)

according to the manufacturer's instructions. RNA was eluted with 40 μL nuclease free water

(NFW) and quantitated using NanoDrop 2000 spectrophotometer. Complementary DNA

(cDNA) was synthesised using SuperScript IV Reverse Transcriptase (ThermoScientific) using

oligo d(T) primers according to the manufacturer's instructions.

Real-time PCR detection were done using SYBR Green PCR Master Mix for 40 cycles in a

Bio-Rad CFX96 thermal cycler. The threshold cycle (Ct) value for each gene was normalised

to the Ct value of a housekeeping gene, *EIF1B*. The relative fold changes were calculated using

ΔΔCt method. The primer sequences for target genes are listed in Table 2.

<u>Wound healing assay</u>

Cells were grown in their respective serum free media for 24 h before 25 000 cells per chamber

were plated into Culture-Insert 2 well in μ-Dish 35 mm, high (ibidi). Cells were kept in serum

free media throughout the experiment. 24 h after seeding, or when cells reach confluency in

the chambers, the culture insert was removed with clean forceps under sterile conditions and

the dish was filled with 2 mL of serum free media for imaging.

<u>Anchorage-independent assay</u>

Anchorage-independent assay was performed using CytoSelect 96-Well Cell Transformation

Assay (Cell Biolabs) according to the manufacturer's instructions. Cells were grown in T25

flasks to 80% confluency, trypsinised and counted with a hemocytometer to obtain the

concentration of live cells. Each well was seeded with 5000 cells and the final cell culture agar

layer solution was 0.4% for soft conditions and 1.2% for stiff conditions, corresponding to a

Young's modulus of <2 kPa(59) and 10 kPa(60) respectively. Fresh medium was changed

every 2 days during incubation and 7 days after seeding, cell colony formation was examined

with EVOS Cell Imaging System (ThermoScientific) under 10X magnification. The total

amount of DNA present in each well was measured with a DNA binding fluorescent dye using

the CyQuant NF Cell Proliferation Assay Kit according to the manufacturer's instructions.

<u>Statistical analysis</u>

Biological replicates refer to independent experimental replicates done on different days with

fresh biological samples and independent transfections and measurements. Technical replicates

refer to repeated measurements of the same sample that represent independent measures of the

random noise associated with protocols or equipment.(61) For comparisons between the same

sample in different light and dark conditions, a paired t-test was used. For comparisons between

two distinct samples, an unpaired t-test was used. For small sample sizes, estimation statistics

was used to show the magnitude of the testing condition.(62)

<u>Curve fitting</u>

Curve fitting for the import and export rates was performed in MATLAB using the curve fitting

function *fit*. Time constant (τ) represents the time scale over which the signal changes before

or after light activation. Nuclear import rate (τ_a) was fitted with $f(x) = a + b\left(1 - e^{-\frac{x}{\tau}}\right)$ and

export rate (τ_r) was fitted with $f(x) = a + b\left(e^{-\frac{x}{\tau}}\right)$, where a is the basal level of signal

intensity; and b is the multiplicative factor representing the change in signal due to light

(de-)activation.

Supplementary Data

Primer name	Sequence	Cloning	Restriction enzyme
PT25 hYAP1-1delta fwd	ATTTATGGTACC ATGGATCCCGGGCAG	Forward primer for S127A and ΔC mutants	KpnI
PT26 hYAP1-1delta S127A rev	CGCGCCGTTTAAACC TATAACCATGTAAG	Reverse primer for S127A	PmeI
PT27 hYAP1-1delta DeltaC rev	CGCGCCGTTTAAACC TAGCTTTCTTTATCTAGC	Reverse primer for ΔC	PmeI
795-2194-YAP-S251D-F1	GATCCACAGGGAGG CGTCATGGGTG	Forward primer for SDM of S251D mutant	-
796-2194-YAP-S251D-R1	CTGGGGAGCCAGGG GTGGTGGCTGT	Reverse primer for SDM of S251D mutant	-
797-2195-YAP-S333D-F1	GATCCCGGGATGTC TCAGGAATTGAG	Forward primer for SDM of S333D mutant	-
798-2195-YAP-S333D-R1	CGACACTGGATTTT GAGTCCCACCATC	Reverse primer for SDM of S333D mutant	-
799-2196-YAP-2mut-F1	CAGCCACCACCCCTGGC TCCCCAGGATCCACAGG GAGGCGTCATGGGTG	Forward primer for SDM of S251D and S333D mutant	-
800-2196-YAP-2mut-R1	CTCAATTCCTGAGACATC CCGGGATCCGACACTGG ATTTTGAGTCCCACCATC	Reverse primer for SDM of S251D and S333D mutant	-

540   **Table 1. List of primers for plasmid cloning.**

Target gene	Forward primer	Reverse primer
EIF1B	TGATGCAACT AAGGGTGACG	CAGTGTCTTT CTGCCGTTCC
CTGF	TGCATCCGTA CTCCCAAAAT	ATGTCTTCAT GCTGGTGCAG
CYR61	TCCCTGTTTT TGGAATGGAG	GAGCACTGGG ACCATGAAGT
TGF-β	TACCTGAACCC GTGTTGCTCTC	GTTGCTGAGGT ATCGCCAGGAA

Table 2. Primers used for RT-qPCR.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analysed during this study are available from the corresponding author

on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was primarily supported by a Singapore Ministry of Education Tier 3 grant

(MOE2016-T3-1-250 002). M.S. was also supported by grants from the National University of

Singapore (R-185-000-2710- 133 & 733) and the Mechanobiology Institute (R-714-018-006-

271). T.E.S. was further supported by an EMBO Global Investigator award and startup funds

from University of Warwick, UK.

Authors' contributions

PJYT, MS and TES conceived the study and design of the work. PJYT performed the

experiments. PJYT and TES did the data analysis. PJYT and TES drafted the work with all

authors approving the final manuscript.

Acknowledgements

We thank Michael Sheetz, in whose lab the work on developing the optogenetic YAP construct

was initiated, and Manvendra Singh for comments on the manuscript.

References

1.   Murry CE, Reinecke H, Pabon LM. Regeneration gaps: observations on stem cells and

cardiac repair. J Am Coll Cardiol. 2006 May;47(9):1777–85.

2.   Sedmera D, Reckova M, DeAlmeida A, Coppen SR, Kubalak SW, Gourdie RG, et al.

Spatiotemporal pattern of commitment to slowed proliferation in the embryonic mouse

heart indicates progressive differentiation of the cardiac conduction system. Anat Rec

A Discov Mol Cell Evol Biol. 2003 Sep;274(1):773–7.

3.   Steinhauser ML, Lee RT. Regeneration of the heart. EMBO Mol Med. 2011 Dec

1;3(12):701–12.

4.   Porrello ER, Mahmoud AI, Simpson E, Hill JA, Richardson JA, Olson EN, et al.

Transient Regenerative Potential of the Neonatal Mouse Heart. Science (1979).

2011;331(6020):1078–80.

5.   Poss KD, Wilson LG, Keating MT. Heart Regeneration in Zebrafish. Science (1979).

2002 Dec 13;298(5601):2188–90.

6.   Chong JJH, Yang X, Don CW, Minami E, Liu YW, Weyers JJ, et al. Human embryonic-
stem-cell-derived cardiomyocytes regenerate non-human primate hearts. Nature. 2014
Jun;510(7504):273–7.

7.   Liu YW, Chen B, Yang X, Fugate JA, Kalucki FA, Futakuchi-Tsuchida A, et al. Human
embryonic stem cell-derived cardiomyocytes restore function in infarcted hearts of non-
human primates. Nat Biotechnol. 2018 Aug;36(7):597–605.

8.   Xin M, Kim Y, Sutherland LB, Murakami M, Qi X, McAnally J, et al. Hippo pathway
effector Yap promotes cardiac regeneration. Proceedings of the National Academy of
Sciences. 2013 Aug 20;110(34):13839–44.

9.   Heallen T, Morikawa Y, Leach J, Tao G, Willerson JT, Johnson RL, et al. Hippo
signaling impedes adult heart regeneration. Development. 2013 Dec;140(23):4683–90.

10.   Leach JP, Heallen T, Zhang M, Rahmani M, Morikawa Y, Hill MC, et al. Hippo pathway
deficiency reverses systolic heart failure after infarction. Nature. 2017
Oct;550(7675):260–4.

11.   Panciera T, Azzolin L, Cordenonsi M, Piccolo S. Mechanobiology of YAP and TAZ in
physiology and disease. Nat Rev Mol Cell Biol. 2017;18(12):758–70.

12.   Fu V, Plouffe SW, Guan KL. The Hippo pathway in organ development, homeostasis,
and regeneration. Curr Opin Cell Biol. 2017;49:99–107.

13.   Basu S, Totty NF, Irwin MS, Sudol M, Downward J. Akt phosphorylates the Yes-
Associated Protein, YAP, to induce interaction with 14-3-3 and attenuation of p73-
mediated apoptosis. Mol Cell. 2003;11(1):11–23.

14.   Kanai F, Marignani PA, Sarbassova D, Yagi R, Hall RA, Donowitz M, et al. TAZ: a
novel transcriptional co-activator regulated by interactions with 14-3-3 and PDZ domain
proteins. EMBO J. 2000 Dec 15;19(24):6778–91.

15. Zhao B, Li L, Tumaneng K, Wang CY, Guan KL. A coordinated phosphorylation by

Lats and CK1 regulates YAP stability through SCFβ-TRCP. Genes Dev. 2010;24(1):72–

85.

16. Vassilev A, Kaneko KJ, Shu H, Zhao Y, DePamphilis ML. TEAD/TEF transcription

factors utilize the activation domain of YAP65, a Src/Yes-associated protein localized

in the cytoplasm. Genes Dev. 2001 May 15;15(10):1229–41.

17. Zhang H, Liu CY, Zha ZY, Zhao B, Yao J, Zhao S, et al. TEAD Transcription Factors

Mediate the Function of TAZ in Cell Growth and Epithelial-Mesenchymal Transition.

Journal of Biological Chemistry [Internet]. 2009 May 15;284(20):13355–62. Available

from: https://doi.org/10.1074/jbc.M900843200

18. Zhao B, Ye X, Yu J, Li L, Li W, Li S, et al. TEAD mediates YAP-dependent gene

induction and growth control. Genes Dev. 2008;22(14):1962–71.

19. Yu FX, Zhao B, Guan KL. Hippo Pathway in Organ Size Control, Tissue Homeostasis,

and Cancer. Cell. 2015;163(4):811–28.

20. Moya IM, Halder G. Hippo–YAP/TAZ signalling in organ regeneration and

regenerative medicine. Nat Rev Mol Cell Biol. 2019;20:211–26.

21. Lai JKH, Toh PJY, Cognart HA, Chouhan G, Saunders TE. DNA-damage induced cell

death in yap1;wwtr1 mutant epidermal basal cells. Elife. 2022 May 1;11.

22. Wen X, Jiao L, Tan H. MAPK/ERK Pathway as a Central Regulator in Vertebrate Organ

Regeneration. Int J Mol Sci. 2022;23(3).

23. De Simone A, Evanitsky MN, Hayden L, Cox BD, Wang J, Tornini VA, et al. Control

of osteoblast regeneration by a train of Erk activity waves. Nature. 2021

Feb;590(7844):129–33.

24.   Bassat E, Mutlak YE, Genzelinakh A, Shadrin IY, Baruch Umansky K, Yifa O, et al.

630   The extracellular matrix protein agrin promotes heart regeneration in mice. Nature.

631   2017;547(7662):179–84.

632   25.   Aharonov A, Shakked A, Umansky KB, Savidor A, Genzelinakh A, Kain D, et al.

633   ERBB2 drives YAP activation and EMT-like processes during cardiac regeneration. Nat

634   Cell Biol. 2020;22(11):1346–56.

635   26.   Oka T, Mazack V, Sudol M. Mst2 and Lats kinases regulate apoptotic function of Yes

636   kinase-associated protein (YAP). Journal of Biological Chemistry.

637   2008;283(41):27534–46.

638   27.   Ege N, Dowbaj AM, Jiang M, Howell M, Hooper S, Foster C, et al. Quantitative

639   Analysis Reveals that Actin and Src-Family Kinases Regulate Nuclear YAP1 and Its

640   Export. Cell Syst. 2018/06/18. 2018 Jun 27;6(6):692-708.e13.

641   28.   Zhao B, Wei X, Li W, Udan RS, Yang Q, Kim J, et al. Inactivation of YAP oncoprotein

642   by the Hippo pathway is involved in cell contact inhibition and tissue growth control.

643   Genes Dev. 2007;21:2747–61.

644   29.   Zhang H, Pasolli HA, Fuchs E. Yes-associated protein (YAP) transcriptional coactivator

645   functions in balancing growth and differentiation in skin. Proceedings of the National

646   Academy of Sciences. 2011;108(6):2270–5.

647   30.   Lamar JM, Stern P, Liu H, Schindler JW, Jiang ZG, Hynes RO. The Hippo pathway

648   target, YAP, promotes metastasis through its TEAD-interaction domain. Proc Natl

649   Acad Sci U S A. 2012 Sep;109(37):E2441-50.

650   31.   Oka T, Remue E, Meerschaert K, Vanloo B, Boucherie C, Gfeller D, et al. Functional

651   complexes between YAP2 and ZO-2 are PDZ domain-dependent, and regulate YAP2

652   nuclear localization and signalling. Biochemical Journal. 2010;432(3):461–78.

32.   Oka T, Sudol M. Nuclear localization and pro-apoptotic signaling of YAP2 require

intact PDZ-binding motif. Genes to Cells. 2009;14(5):607–15.

33.   Mason DE, Collins JM, Dawahare JH, Nguyen TD, Lin Y, Voytik-Harbin SL, et al.

YAP and TAZ limit cytoskeletal and focal adhesion maturation to enable persistent cell

motility. Journal of Cell Biology. 2019 Feb 8;218(4):1369–89.

34.   D'Angelo E, Lindoso RS, Sensi F, Pucciarelli S, Bussolati B, Agostini M, et al. Intrinsic

and Extrinsic Modulators of the Epithelial to Mesenchymal Transition: Driving the Fate

of Tumor Microenvironment. Vol. 10, Frontiers in Oncology. Frontiers Media S.A.;

2020.

35.   Morikawa Y, Zhang M, Heallen T, Leach J, Tao G, Xiao Y, et al. Actin cytoskeletal

remodeling with protrusion formation is essential for heart regeneration in Hippo-

deficient mice. Sci Signal. 2015 May;8(375):ra41.

36.   Mia MM, Cibi DM, Ghani SABA, Singh A, Tee N, Sivakumar V, et al. Loss of Yap/Taz

in cardiac fibroblasts attenuates adverse remodelling and improves cardiac function.

Cardiovasc Res [Internet]. 2022 May 15;118(7):1785–804. Available from:

https://doi.org/10.1093/cvr/cvab205

37.   Zhou J. An emerging role for Hippo-YAP signaling in cardiovascular development. J

Biomed Res. 2014 Jul;28(4):251–4.

38.   Fenno L, Yizhar O, Deisseroth K. The development and application of optogenetics.

Annu Rev Neurosci [Internet]. 2011;34:389–412. Available from:

https://pubmed.ncbi.nlm.nih.gov/21692661

39.   Krueger D, Izquierdo E, Viswanathan R, Hartmann J, Pallares Cartes C, de Renzis S.

Principles and applications of optogenetics in developmental biology. Development

[Internet]. 2019 Oct 22;146(20):dev175067. Available from:

https://doi.org/10.1242/dev.175067

40.   Toh PJY, Lai JKH, Hermann A, Destaing O, Sheetz MP, Sudol M, et al. Optogenetic control of YAP cellular localisation and function. EMBO Rep [Internet]. 2022 Sep 5;23(9):e54401. Available from: https://doi.org/10.15252/embr.202154401

41.   Liu Z, Yi L, Du M, Gong G, Zhu Y. Overexpression of TGF-β enhances the migration and invasive ability of ectopic endometrial cells via ERK/MAPK signaling pathway. Exp Ther Med. 2019;17(6):4457–64.

42.   Szeto SG, Narimatsu M, Lu M, He X, Sidiqi AM, Tolosa MF, et al. YAP/TAZ are mechanoregulators of TGF-b-smad signaling and renal fibrogenesis. Journal of the American Society of Nephrology. 2016;27(10):3117–28.

43.   Qin Z, Xia W, Fisher GJ, Voorhees JJ, Quan T. YAP/TAZ regulates TGF-β/Smad3 signaling by induction of Smad7 via AP-1 in human skin dermal fibroblasts. Cell Communication and Signaling. 2018;16(1):18.

44.   Sorensen DW, van Berlo JH. The Role of TGF-β Signaling in Cardiomyocyte Proliferation. Curr Heart Fail Rep. 2020 Oct;17(5):225–33.

45.   Peng Y, Wang W, Fang Y, Hu H, Chang N, Pang M, et al. Inhibition of TGF-β/Smad3 Signaling Disrupts Cardiomyocyte Cell Cycle Progression and Epithelial–Mesenchymal Transition-Like Response During Ventricle Regeneration. Front Cell Dev Biol. 2021;9.

46.   Qiao Y, Chen J, Lim YB, Finch-Edmondson ML, Seshachalam VP, Qin L, et al. YAP regulates actin dynamics through ARHGAP29 and promotes metastasis. Cell Rep [Internet]. 2017;19(8):1495–502. Available from: http://www.sciencedirect.com/science/article/pii/S2211124717306046

47.   Reddy BVVG, Irvine KD. Regulation of Hippo Signaling by EGFR-MAPK Signaling through Ajuba Family Proteins. Dev Cell. 2013;24(5):459–71.

48. Bui DA, Lee W, White AE, Harper JW, Schackmann RCJ, Overholtzer M, et al.

Cytokinesis involves a nontranscriptional function of the Hippo pathway effector YAP.

Sci Signal. 2016 Mar 1;9(417):ra23–ra23.

49. Yang S, Zhang L, Liu M, Chong R, Ding SJ, Chen Y, et al. CDK1 Phosphorylation of

YAP Promotes Mitotic Defects and Cell Motility and Is Essential for Neoplastic

Transformation. Cancer Res. 2013 Nov 14;73(22):6722–33.

50. Illes B, Fuchs A, Gegenfurtner F, Ploetz E, Zahler S, Vollmar AM, et al. Spatio-selective

activation of nuclear translocation of YAP with light directs invasion of cancer cell

spheroids. iScience [Internet]. 2021;24(3):102185. Available from:

https://www.sciencedirect.com/science/article/pii/S258900422100153X

51. Dowbaj AM, Jenkins RP, Williamson D, Heddleston JM, Ciccarelli A, Fallesen T, et al.

An optogenetic method for interrogating YAP1 and TAZ nuclear–cytoplasmic shuttling.

J Cell Sci [Internet]. 2021 Jul 9;134(13):jcs253484. Available from:

https://doi.org/10.1242/jcs.253484

52. Gaffney CJ, Oka T, Mazack V, Hilman D, Gat U, Muramatsu T, et al. Identification,

basic characterization and evolutionary analysis of differentially spliced mRNA

isoforms of human YAP1 gene. Gene. 2012 Nov;509(2):215–22.

53. Meyer K, Lammers NC, Bugaj LJ, Garcia HG, Weiner OD. Decoding of YAP levels

and dynamics by pluripotency factors. Available from:

https://doi.org/10.1101/2022.10.17.512504

54. Ben C, Wu X, Takahashi-Kanemitsu A, Knight CT, Hayashi T, Hatakeyama M.

Alternative splicing reverses the cell-intrinsic and cell-extrinsic pro-oncogenic

potentials of YAP1. Journal of Biological Chemistry [Internet]. 2020;295(41):13965–

80. Available from:

https://www.sciencedirect.com/science/article/pii/S0021925817497961

55.   Vrbský J, Vinarský V, Perestrelo AR, de La Cruz JO, Martino F, Pompeiano A, et al.

727   Evidence for discrete modes of YAP1 signaling via mRNA splice isoforms in

728   development and diseases. Genomics [Internet]. 2021;113(3):1349–65. Available from:

729   https://www.sciencedirect.com/science/article/pii/S0888754321001014

730   56.   Elbediwy A, Vincent-Mistiaen ZI, Spencer-Dene B, Stone RK, Boeing S, Wculek SK,

731   et al. Integrin signalling regulates YAP and TAZ to control skin homeostasis.

732   Development. 2016 May 15;143(10):1674–87.

733   57.   Lee MJ, Byun MR, Furutani-Seiki M, Hong JH, Jung HS. YAP and TAZ Regulate Skin

734   Wound Healing. Journal of Investigative Dermatology. 2014;134(2):518–25.

735   58.   Suhaeri M, Subbiah R, Van SY, Du P, Kim IG, Lee K, et al. Cardiomyoblast (h9c2)

736   differentiation on tunable extracellular matrix microenvironment. Tissue Eng Part A.

737   2015 Jun;21(11–12):1940–51.

738   59.   Li C, Huang Z, Wang RK. Elastic properties of soft tissue-mimicking phantoms assessed

739   by combined use of laser ultrasonics and low coherence interferometry. Opt Express.

740   2011 May;19(11):10153–63.

741   60.   Nitta T, Haga H, Kawabata K, Abe K, Sambongi T. Comparing microscopic with

742   macroscopic elastic properties of polymer gel. Ultramicroscopy. 2000;82(1):223–6.

743   61.   Blainey P, Krzywinski M, Altman N. Replication. Nat Methods. 2014;11(9):879–80.

744   62.   Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A. Moving beyond P values: data

745   analysis with estimation graphics. Nat Methods. 2019;16(7):565–6.