NUCLEOLAR REORGANIZATION AFTER CELLULAR STRESS IS ORCHESTRATED BY SMN

SHUTTLING BETWEEN NUCLEAR COMPARTMENTS

Shaqraa Musawi[1,2,†], Lise-Marie-Donnio[1,†,*], Zehui Zhao[1], Charlène Magnani[1], Olivier Binda[1,3],

Jianbo Huang[1], Phoebe Rassinoux[1], Arnaud Jacquier[1], Laurent Coudert[1], Patrick Lomonte[1], Cécile

Martinat[4], Laurent Schaeffer[1], Jocelyn Côté[3], Pierre-Olivier Mari[1] and Giuseppina GIGLIA-MARI[1,*]

1. Institut NeuroMyoGène-Pathophysiology and genetics of neuron and muscle

(INMG_PGNM), CNRS UMR 5261, INSERM U1310, Université de Lyon, Université Claude

Bernard Lyon 1, Lyon, France

2. Department of Medical Laboratories Technology, College of Applied Medical Sciences,

Jazan University, Jazan, Saudi Arabia.

3. University of Ottawa, Faculty of Medicine, Department of Cellular and Molecular

Medicine, K1H 8M5, Ottawa, Ontario, CANADA

4. INSERM/UEPS UMR 861, Paris Saclay Université, I-STEM, 91100 Corbeil-Essonnes, France

† These authors contributed equally to this work

* Co-Corresponding authors

<u>**Keywords:**</u>

SMA, SMN, RNAP1, DNA repair, UV lesions, Nucleolus, Coilin, Fibrillarin

ABSTRACT

SMA is an autosomal recessive neuromuscular disease caused by mutations in the multifunctional protein SMN. Within the nucleus, SMN localizes to Cajal bodies (CBs), which have been shown to be associated with nucleoli, nuclear organelles dedicated to the first steps of ribosome biogenesis. The highly organized structure of the nucleolus can be dynamically altered by genotoxic agents. After genotoxic stress, RNAP1, Fibrillarin (FBL) and nucleolar DNA are exported to the periphery of the nucleolus and once DNA repair is fully completed the organization of the nucleolus is restored. We found that SMN is required for the restoration of the nucleolar structure after genotoxic stress. Unexpectedly, during DNA repair, SMN shuttles from the CBs to the nucleolus. This shuttling is important for nucleolar homeostasis and relies on the presence of Coilin and the activity of PRMT1.

INTRODUCTION

The nucleolus is a nuclear membrane-less organelle with a very structured internal organization, which associates with its different functions in ribosomal biogenesis: transcription of ribosomal DNA (rDNA) and early ribosomal RNA (rRNA) maturation [1]. Nucleoli are formed around the rDNAs which are composed of tandem head-to-tail repeats and their structure is thought to be strictly dependent on the transcriptional activity of the RNA polymerase I (RNAP1) [2]. Despite a very structured organization, nucleoli are very dynamic organelles, their shape and number can vary through the cell cycle and many proteins can enter or exit the nucleolus depending on physiological processes or cellular stress responses. This organized structure can be dynamically altered by both genotoxic agents and general cellular stress [3]. For instance, drugs that alter RNAP1 transcription (i.e. cordycepin, actinomycin D, etc.) may cause nucleolar segregation at the periphery of the nucleolus into structures known as nucleolar caps. Furthermore, drugs that block rRNA processing or the topoisomerase II (i.e. doxorubicin) but do not interfere with RNAP1 transcription induce a disruption of the compact nucleolar environment and nucleolar necklaces appear [4].

Amongst different cellular stresses known to modify the nucleolar organization, UV-irradiation has the benefit of being a quick, punctual, and chemically clean method. Moreover, cells are able to repair UV-induced lesions and hence reverse their stress status. UV-induced DNA lesions are repaired by the Nucleotide Excision Repair system (NER)[5]. NER also repairs DNA helix-distorting adducts, including environmental pollutants, the oxidative-damage derived cyclopurines [6] and participates in the repair of the ROS-induced oxidized guanine (8-oxoG) [7].

During UV-irradiation, it has been shown that the nucleolus is not fully disrupted but nucleolar proteins (RNAP1, Fibrillarin [FBL]) and nucleolar DNA are exported to the periphery (for simplicity this phase will be called "displacement") of the nucleolus and when DNA repair is fully completed the proper nucleolar structure is restored (for simplicity this phase will be called "repositioning")[8]. Using a best-candidate approach, we recently found that structural proteins like Nuclear Myosin I (NMI) and **β**-Actin (ActB) seem to play a prominent role in this process [9]. In cells depleted from NMI and ActB, nucleolar structure is not restored and both nucleolar proteins and nucleolar DNA remain at the periphery of the nucleolus, although DNA repair is completed

and transcription is resumed [9]. However, the exact mechanism of NMI and ActB actions on

nucleolar reorganization has not yet been elucidated, probably because many other molecular

actors are still unknown. In order to find a complete molecular mechanism, several structural and

nucleolar proteins have been scrutinized and a certain number have been found to be crucial to

restore a proper nucleolar structure after DNA repair completion. One of these proteins is FBL.

Consequently, we studied whether FBL interacting partners were also involved in this process.

Amongst these different FBL partners, we investigated whether the protein Survival of Motor

Neurons (SMN) was implicated in the restoration of the nucleolar structure after DNA repair

completion.

Spinal Muscular Atrophy (SMA) is an autosomal recessive neuromuscular disease, which

affects neurons that controls the voluntary movement of muscles (motoneurons) [10]. In SMA,

motoneurons are progressively lost leading to progressive muscle wasting and atrophy because

muscles no longer receive signals from the motor neurons of the spinal cord. Children affected

with SMA have symptoms that can vary greatly depending on the age of disease onset and its

severity. Normal activities, such as crawling, walking, maintaining a seated position, controlling

head movements, breathing and swallowing might be affected [10]. With an incidence of 1 in 6,000-

10,000 live births, SMA is the most prevalent hereditary cause of infant mortality [11].

SMA is caused by bi-allelic mutations in the *SMN1* gene (Survival of Motor Neuron: SMN)

and the disease phenotype is modified by the number of copies of a second paralog gene,

SMN2, which is always present in SMA patients [12]. *SMN1* produces a full-length functional version

of the SMN protein whereas in *SMN2*, the absence of exon 7 in most of the transcripts produces

an unstable version of the SMN protein (SMNΔ7). *SMN2* can express about 10-15% of the full-

length protein, which is insufficient to avoid the disease.

SMN is a multifunctional ubiquitous protein involved in many cellular processes, such as

biogenesis and trafficking of ribonucleoproteins, local translation of messenger RNAs, etc [13]. SMN

protein is ubiquitously expressed and is localized to both the cytoplasm and the nucleus. Within

the nucleus, SMN localizes in Gems and Cajal bodies (CBs), which have been shown to associate

with nucleoli [14]. Within CBs, SMN interacts with the protein Coilin [15]. Interestingly, in certain

conditions, SMN is also detected in the nucleoli of mammalian primary neurons and colocalized

with FBL [16]. In addition, a transient colocalization of SMN at the periphery of nucleoli with FBL

after actinomycin D treatment in 10–20% of Hela cells [17,18] suggests that SMN could be present

in the nucleolus under stress conditions. SMN Tudor domain is involved in the binding to RGG

motifs containing proteins such as FBL and Coilin [19].

We investigated the possible role of SMN in nucleolar reorganization during both

displacement and repositioning of RNAP1, during and after DNA repair of UV-induced damage

and generally after stress induction. We show here that in the absence of a functional SMN, both

RNAP1 and FBL remain at the periphery of the nucleolus after DNA repair completion even once

transcription is fully restored. Interestingly, we could reveal that SMN (and its complex) shuttles

to the nucleolus after DNA repair completion. We determine that this shuttling is dependent on

the presence of Coilin and governed by PRMT1-dependent arginine methylation activity.

Additionally, we disclosed that the presence of FBL is important for the proper restoration of

SMN into CBs. Importantly, we could show that SMN cells show a sensitivity to DNA damage and

in particular, to chronic oxidative damage. Our results open up new avenues of research on the

SMN function in nucleolar homeostasis and its impact on SMA pathology.

RESULTS

RNAP1 and FBL repositioning after DNA repair completion are SMN-dependent.

To investigate a possible role of SMN in nucleolar reorganization in response to cellular stress, we investigated whether the previously reported [8] RNAP1 UV-induced displacement and the later repositioning were still happening in the absence of SMN. As SMN-deficient cells we used primary fibroblasts from SMA patients (Figure S1A), SMA iPSC-derived motoneurons (Figure S2) and transformed fibroblasts in which SMN was downregulated by lentiviral transfection of 2 independent inducible shRNAs against SMN 3'UTR (Figure 1A). Using these cell lines, we performed immunofluorescent (IF) assays to detect both RNAP1 and FBL positioning in the absence of damage (No UV), 3 hours post-UV-irradiation (PUVI) (this time point corresponds to the minimum of RNAP1 transcriptional activity as found in [8] and at 40 hours PUVI (this time point corresponds to the RNAP1 full recovery of transcriptional activity and full DNA repair as described in [8]). Wild-type fibroblasts, SV40-transformed MRC5 and primary C5RO were used as positive controls, while Cockayne Syndrome type B (CSB) TC-NER deficient fibroblasts (both transformed and primary; termed CS1AN) were used as negative control, as used in [8]. For iPSC-derived motoneurons, WT and SMA iPSC-derived neuronal progenitors were differentiated into motoneurons [20].

As described in [8], UV-irradiation induced a displacement of both RNAP1 and FBL to the periphery of nucleoli in all cell lines tested (Figures 1B and 1C, Figures S1B and S1C, Figure 2 at 3h PUVI). As expected, in wild-type cells (MRC5, shSCRAMBLE and C5RO) both RNAP1 and FBL recovered their position within the nucleoli at 40 hours PUVI (Figures 1B and 1 C, Figures S1B and S1C). In contrast, in cells depleted of SMN (Figures 1B and 1 C, Sh5-SMN and Sh6-SMN), primary fibroblasts mutated in *SMN1* (Figures S1B and S1C, SMA1) and SMA iPSC-derived motoneurons (Figures S2A and S2B) neither RNAP1 nor FBL recovered the proper position within the nucleoli after DNA repair completion. As previously demonstrated [8], in CSB-deficient cells no return of the RNAP1 and FBL was observed (Figures 1B and 1C, CS1AN-SV) (Figures S1B and S1C, CS1AN).

In CSB-deficient cells, the repositioning of RNAP1 and FBL is impeded because DNA lesions on the transcribed strand of rDNA genes are not properly repaired and RNAP1

transcription is not restored [8]. To investigate whether this was the case in SMN-deficient cells, we

performed an RNA-fish assay detecting the pre-rRNA transcript using a specific probe against

the 47S product (Figure S3A) and could determine (Figures S3B and S1E) and quantify (Figures

1D and S1D) that RNAP1 transcription is restored in SMN-deficient cells at 40 hours PUVI as in

wild-type cells. In parallel, the involvement of SMN in Nucleotide Excision Repair (NER) was

studied by performing UDS (Figure S4A), RRS (Figure S4B) and TCR-UDS (Figures S4C and S4D)

experiments in cells depleted of SMN. Our results clearly show that SMN has no role in NER

(Figure S4).

Taken together, these results indicate that in the absence of SMN, RNAP1 and FBL are

correctly displaced at the periphery of the nucleolus in response to DNA damage but are not

repositioned within the nucleolus once DNA repair reactions are completed and RNAP1

transcription is restored.

A

Western Blot

B

IF RNAP1/FBL

C

D

RNA-FISH 47S

Figure 1: RNAP1 and FBL localization during DNA repair in SMN deficient transformed fibroblasts.

(A) Western Blot of SMN and CSB on whole cell extracts of transformed fibroblasts (MRC5-SV cells). Doxycycline treatment induces the expression of the ShRNA against SMN. (B) Representative confocal microscopy images of immunofluorescence (IF) assay against RNAP1 (green) and FBL (red) in transformed fibroblasts, at different times PUVI. Nuclei and nucleoli are indicated by dashed lines and dotted lines respectively. Scale bar: 5µm (C) Quantification of cell number for RNAP1 localization (inside the nucleolus, outside the nucleolus or mixed localization) at different times PUVI. (D) Quantification of RNA-FISH assay showing the 47S pre-rRNA level after UV-C exposure in transformed fibroblasts. Error bars represent the SEM obtained from at least 27 cells. P-value of student's test compared to No UV condition: ***<0.001

SMN-complex shuttles in the nucleolus and co-localizes with nucleolar proteins after UV irradiation.

We showed that SMN is required for the proper repositioning of RNAP1 and FBL at late time points PUVI. We questioned how SMN could be involved in this mechanism if, usually, it is not present in the nucleoli. In fact, SMN protein is typically located in the cytoplasm and within the nucleus where SMN is found in CBs together with Coilin and in Gems without Coilin. To study the localization of SMN and RNAP1 during the DNA repair process, we initially performed immunofluorescence assays at 3 hours and 40 hours PUVI in wild-type cells. Remarkably, at 40h PUVI, two co-existing populations of cells could be detected: (i) a majority of cells in which RNAP1 is repositioned within the nucleolus and SMN is localized in the CBs/Gems and (ii) a minority of cells in which RNAP1 is still localized at the periphery of the nucleolus and SMN is unusually localized at the periphery of the nucleolus and cannot be detected in CBs/Gems anymore (Figures S5A, S5B, S5C). Because of this result, we decided to extend our analysis by adding a time point intermediate between 3h and 40h PUVI (24 h PUVI) and a time point beyond the 40h PUVI (48h PUVI) and performed the IF assays against SMN and different nucleolar proteins or SMN protein partners (Figure 2). Importantly, we verified the RNAP1 transcriptional activity by

RNA-FISH of 47S at 24h and 48h PUVI and show that RNAP1 transcription is not yet restored at

24h PUVI but it's fully recovered at 48h PUVI (Figure S5D). Surprisingly, at 24h PUVI, our results

revealed the presence of SMN at the periphery of and/or within the nucleolus in the vast majority

of cells (Figure 2A, quantification in 2D). Concomitantly, at 24h PUVI, RNAP1 was found to be

localized at the periphery of the nucleolus (Figure 2A, quantification in 2E). On the other hand,

we observed a complete return to the undamaged condition (RNAP1 within the nucleolus and

SMN in the CBs) at 48h PUVI in the vast majority of cells (Figure 2A, quantification in 2D and 2E).

Despite the presence of SMN and RNAP1 at the nucleolar periphery together at 24 h post-UV,

no co-localization between these two proteins was observed (Figure 2A, panel 24h PUVI).

We showed that the loss of SMN alters FBL localization at 40 h PUVI (Figures 1B and S1B).

Using GST pull-down assays we confirmed that FBL from cell extracts interacts with SMN [17,21,22]

(Figure S5E). Furthermore, using a panel of SMA-linked TUDOR domain mutants, we established

that FBL-SMN interactions require an intact TUDOR domain (Figures S5E and S5F). We, therefore,

examined the localization of SMN and FBL during the DNA repair process (Figure 2B,

quantification in 2F). Interestingly, we observed a substantial co-localization between SMN and

FBL at 24 h PUVI within the nucleolus (Figure 2B, panel 24h PUVI). The co-localization can also

be observed at 40 h PUVI in the population of cells that, at this time point, have not yet restored

the FBL position within the nucleolus (Figure S5B).

In the absence of damage, SMN co-localizes and interacts with Coilin in CBs [15], however

after DNA damage induction, it has been shown that CBs are disrupted [23]. Because of this

evidence, we examined the localization of Coilin and its interactions with SMN during the DNA

repair process. Interestingly, Coilin is localized to the periphery of the nucleolus already at 3h

PUVI (Figure 2C, panel 3h PUVI) and remains in this location at 24h PUVI (Figure 2C, panel 24h

PUVI) and at 40h PUVI (Figure 2C, panel 40h PUVI) in the subset of cells that did not yet reposition

RNAP1. Remarkably, when Coilin is at the periphery of the nucleolus, no colocalization with SMN

is observed (Figure 2C).

To investigate whether the shuttling of SMN within the nucleolus is specific to UV damage

or also happening when cells are submitted to other types of damage, we treated MRC5-SV cells

with KBrO$_3$, known to produce a majority of 8-oxo-Guanine lesions [24] and examined the

localization of SMN. When treated with KBrO$_3$, SMN shuttling into the nucleolus was also

observed (Figure S6A). Additionally, to investigate whether the SMN shuttling is a general event

in stress response, we treated the MRC5-SV cells with Cordycepin, an RNAP1 transcription-

blocking drug (FigureS6B) and checked the localization of SMN (Figure S6C). After cordycepin

treatment (3 hours), SMN shuttling was observed into the nucleolus but with a kinetic that was

faster than the one measured after damage induction (Figure S6C).

Finally, in order to visualize SMN shuttling in living cells, we produced a Cherry-SMN cell

line by transfecting a Cherry-SMN expressing plasmid into the inducible sh6-SMN cell line and

selecting a suitable clone that presented the correct SMN localization and level of expression

(Figure S7A). After depletion of the endogenous SMN, by induction of the shRNA against SMN

with doxycycline, we UV-irradiated and performed time-lapse imaging over the following 48

hours, with intervals of 1 hour per image (Figure S7B). During the time-lapse, we could detect

the fading of the focal Cherry-SMN signal at around 3 to 6 hours PUVI and the concomitant

appearance of Cherry SMN around the nucleoli. This signal lasted at least 24 hours for some cells

and disappeared at around 32 hours PUVI with the reappearance of the focal pattern (CBs or

Gems) around 40 hours PUVI (Figure S7B).

Figure 2: Localization of SMN and its partners during DNA repair

Representative microscopy images of immunofluorescence (IF) assay in MRC5 cells showing, after 16J/m² UV-C irradiation, the localization of SMN (red) and (A) RNAP1, (B) FBL or (C) Coilin (green) at different times PUVI. Nuclei and nucleoli are indicated by dashed lines and dotted lines respectively. Scale bar represents 5 µm.

Quantification of cells number for localization of (D) SMN and (G) Coilin (in Cajal Bodies [CBs] or Gems, at the periphery of the nucleolus or mixed localization), of (E) RNAP1 and (F) FBL (inside the nucleolus, outside the nucleolus or mixed localization) at different times PUVI.

SMN complex shuttles at the nucleolus after UV irradiation.

Interestingly and differently from Coilin (Figure 2C), at 3h PUVI, SMN is still visible in a focal pattern within the nucleus, reminiscent of Gems. Because of this Coilin-independent localization and to investigate whether SMN shuttles within the nucleolus as an individual protein or as a complex, we investigated whether Gemin2, 3, 4 and 5 (subunits of the SMN complex) change location after UV-irradiation. Remarkably, we could show that Gemin 3, Gemin 4 and Gemin 5 interact with SMN all along this process of displacement and repositioning (Figures 3B, 3C, 3D and corresponding quantifications in 3F, 3G and 3H), suggesting that the whole SMN complex is likely involved in the shuttling, or at least some components of the complex. It should be noted that Gemin5 is also present in foci that are SMN negative, these foci were visible all along the nucleolar reorganization process (Figure 3D). Interestingly, Gemin 2 presented a different localization than SMN in both undamaged and damaged conditions. Gemin 2 was found to be present in both CBs/Gems and in the nucleolus in undamaged cells (Figure 3A and corresponding quantification 3E). After damage induction, the proportion of CBs/Gems containing Gemin2 decreased and concomitantly the nucleolar localization increased. To exclude that this nucleolar localization could be the result of a non-specific signal within the nucleolus, we quantified the Gemin2 signal within the nucleolus and observed that the signal increased at 24 to 48 hours PUVI (Figure S8). It is important to note that at 48 hours PUVI, most Gemin 2 was still found within the nucleolus (Figure 3E), while SMN was mainly localized in CBs/Gems (Figure 2D).

254   These results might indicate that some SMN partners follow exactly the same localization of SMN

255   during damage-induced nucleolar reorganization, while others might be delayed in some steps

256   of this process.

Figure 3: Localization of Gemin2, Gemin3, Gemin4 and Gemin5 during DNA repair

Representative microscopy images of immunofluorescence (IF) assay in MRC5 cells showing, after 16J/m² UV-C irradiation, the localization of SMN (red) and (A) GEMIN3, (B) GEMIN3, (C) GEMIN4 or (D) GEMIN2 (green) at different times PUVI. Nuclei and nucleoli are indicated by dashed lines and dotted lines respectively. Scale bar represents 5 μm.

Quantification of cells number for localization of (E) GEMIN3, (F) GEMIN4, (G) GEMIN5 or (H) GEMIN2 (in Cajal Bodies [CBs] or Gems, at the periphery of the nucleolus or mixed localization), at different times PUVI.

(I) Quantification of fluorescent signal in the nucleolus from the IF against GEMIN2. Error bars represent the standard error of the mean (SEM) obtained from at least 80 cells. P-value of Student's test compared to No UV condition. **<0.005, ***<0.001.

SMN interacts with FBL, but not with coilin, in the nucleolus after UV irradiation.

We showed that SMN co-localizes with FBL within the nucleolus at 24h PUVI (Figure 2B panel 24h PUVI). To assess the *in vitro* interaction between SMN and FBL after UV-C exposure, we performed a GST pull-down assay using cell extracts untreated or UV-treated at different time points (Figure S8A). We observed that UV treatment seemed to enhance FBL-SMN interactions, while GST alone failed to associate with FBL even if more GST alone than GST-SMN was used in pull-down assays. To confirm this result *in vivo*, we performed a Proximity Ligation Assay (PLA) on wild-type cells at different times PUVI (Figures S8B and S8C). The majority of the cells at 24h PUVI presented a strong PLA signal specifically in the nucleolus between SMN and FBL (Figure S8C) which could be quantified as a specific interaction signal (Figure S8B).

Surprisingly, no colocalization was found between SMN and Coilin during the shuttling process, neither at 3 hours nor later on (Figure 2 C). SMN and Coilin did colocalize in CBs before damage induction and 48 hours post damage induction. To confirm this result, we performed PLA on wild-type cells at different times PUVI (Figures S8D and S8E) and we could not detect any added interaction/proximity between SMN and Coilin during damage-induced nucleolar reorganization.

Coilin is required for SMN import into the nucleolus and for the nucleolar rearrangement following UV-C exposure.

We observed that Coilin localized to the nucleolus during UV damage (at 3h PUVI) prior to SMN (Figures 2C and 2G). These results led us to hypothesize that Coilin might be the factor that recruits SMN to the nucleolus. To test this idea, we depleted cells of Coilin by using a pool of four specific siRNAs and two individual siRNAs (Figure S9A) and performed IF of SMN and RNAP1 on wild-type cells before damage and at different time PUVI (Figure 4A) in the presence or absence of Coilin. Interestingly, what could be noticed was that the number of cells presenting SMN foci was drastically diminished when Coilin was downregulated (Figure S9B), this result is consistent with the fact that CBs are dismantled in the absence of Coilin. This would mean that, in this condition, visible SMN foci would more likely correspond to Gems. To investigate whether Gems would shuttle to the nucleolus, we excluded from the quantification cells that did not present SMN foci. Our results show that, without Coilin, the shuttling of SMN within the nucleolus is impaired and SMN remains mostly in Gems (Figure 4A and corresponding quantification Figures 4C and S9C). Consequently, in cells deficient for Coilin, the nucleolar reorganization after DNA damage and repair is not restored and RNAP1 remains at the periphery of the nucleolus at 48h PUVI (Figures 4A and 4C and S9C), indicating that the shuttling of SMN and the presence of Coilin are both important to ensure the re-establishment of the nucleolar reorganization.

To investigate whether the absence of Coilin would impact the interaction between SMN and FBL, we performed PLA and IF of SMN and FBL on wild-type cells before damage and at different times PUVI in the presence or absence of Coilin (Figures 4D, 4E and 4F). Our results show that without Coilin, SMN remains outside of the nucleolus and no co-localization with FBL was observed (Figure 4D). As a consequence, the interaction between SMN and FBL observed at 24h PUVI (Figures S8B and S8C) is lost in Coilin-depleted cells (Figures 4E and 4F).

A

No UV   3h PUVI   24h PUVI   48h PUVI

IF RNAP1/SMN

siMock

siCoilin

B SMN localization

siMock   siCoilin

Population (%)

No 3 24 48   No 3 24 48
PUVI (h)   PUVI (h)

■ Gems ■ Mixed ■ Nucleolus

C RNAP1 localization

siMock   siCoilin

Population (%)

No 3 24 48   No 3 24 48
PUVI (h)   PUVI (h)

■ Inside ■ Mixed ■ Outside

D

No UV   3h PUVI   24h PUVI   48h PUVI

IF FBL/SMN

siMock

siCoilin

E PLA SMN-FBL

No UV   3h PUVI   24h PUVI

siMock

siCoilin

F Quantification PLA SMN-FBL

Relative fluorescence (a.u.)

□ siMock ■ siCoilin

***   **   ***

No UV   3h   24h
Time post UV

Figure 4: SMN shuttling is Coilin-dependent

*(A and D) Representative microscopy images of immunofluorescence (IF) assay showing the localization of SMN (red) and (A) RNAP1 or (D) FBL (green) at different times PUVI in cells transfected with siMock or siCoilin pool. Nuclei and nucleoli are indicated by dashed lines and dotted lines respectively. Scale bar: 5µm. (B and C) Quantification of cells number for localization of (B) SMN (in Cajal Bodies [CBs] or Gems, at the periphery of the nucleolus or mixed localization) and (C) RNAP1 (inside the nucleolus, outside the nucleolus or mixed localization) at different times PUVI in cells transfected with siMock or siCoilin pool. (E) Representative microscopy images of PLA showing the interaction between SMN and FBL at different times PUVI in cells transfected with siMock or siCoilin. Scale bar: 5µm. (F) Quantification of fluorescent signal in the nucleus against the couple SMN-FBL from PLA experiment in cells transfected with siMock or siCoilin pool after UV-C irradiation. P-value of Student's test compared to siMock condition. **<0.005, ***<0.001.*

FBL is required for SMN export from the nucleolus and for the nucleolar rearrangement following UV-C exposure.

As Coilin, FBL is an essential partner of SMN and a nucleolar protein that can methylate rDNAs and Histones within the nucleolus [21,25]. We showed that FBL and SMN interact in cell extracts by GST (Figures S5E and S8A) and *in vivo* by PLA (Figures S8B and S8C), notably stronger after SMN shuttling into the nucleolus (24h PUVI). As the SMN shuttling is dependent on Coilin (Figure 4), we wondered whether FBL depletion would also play a role in this shuttling process. To test this hypothesis, we depleted cells of FBL by using a specific siRNA pool (Figure S10A) or 2 independent siRNAs (Figure S10B) and performed IF of SMN and RNAP1 on wild-type cells before damage and at different times PUVI (Figure 5A, corresponding quantifications in Figures 5B and 5C) in presence or absence of FBL. Our results show that without FBL, the shuttling of SMN within the nucleolus is altered, namely SMN localizes at the periphery of the nucleolus already at 3h PUVI and stays at the periphery of the nucleolus at all time points (Figures 5A, 5B and S10B). Importantly, in cells deficient for FBL, the nucleolar reorganization after DNA damage

and repair is not restored and RNAP1 remains at the periphery of the nucleolus at 48h PUVI

(Figures 5A, 5C and S10C), indicating that the shuttling of SMN within the nucleolus and the

presence of FBL are both important to ensure the re-establishment of the nucleolar

reorganization.

To verify if and how the absence of FBL affects the localization and interaction between

SMN and Coilin, we performed IF and PLA of SMN and Coilin on wild-type cells before damage

and at different times PUVI (Figures 5D, 5E and 5F) in the presence or absence of FBL. Our results

show that, without FBL, the interaction between SMN and Coilin is detectable at all times PUVI

(Figures 5E and 5F) and both SMN and Coilin are localized at the periphery of the nucleolus

already at 3h PUVI, this localization does not change at 24h or 40h PUVI (Figure 5D). These

findings show that FBL is also a critical player in SMN shuttling and in nucleolar reorganization

following UV irradiation and DNA repair.

Figure 5: The release of SMN from the nucleolus is FBL-dependent

(A and D) Representative microscopy images of immunofluorescence (IF) assay showing the

localization of SMN (red) and (A) RNAP1 or (D) COILIN (green) at different times PUVI in cells

transfected with siMock or siFBL pool. Nuclei and nucleoli are indicated by dashed lines and

dotted lines respectively. Scale bar: 5µm. (B and C) Quantification of cells number for localization

of (B) SMN (in Cajal Bodies [CBs] or Gems, at the periphery of the nucleolus or mixed localization)

and (C) RNAP1 (inside the nucleolus, outside the nucleolus or mixed localization) at different

times PUVI in cells transfected with siMock or siFBL pool. (E) Representative microscopy images

of PLA showing the interaction between SMN and COILIN at different times PUVI in cells

transfected with siMock or siFBL pool. Scale bar: 5µm. (F) Quantification of fluorescent signal in

the nucleus against the couple SMN-COILIN from PLA experiment in cells transfected with

siMock or siFBL pool after UV-C irradiation. P-value of Student's test compared to siMock

*condition. ***<0.001.*

SMN mutants are deficient in nucleolar reorganization after DNA damage induction and repair

SMN protein has different domains that are important for its functionality, interactions

with partners and stability (Figure S5F). In order to study how different SMN domains impact the

nucleolar reorganization function of SMN, we have focused on two SMN mutants: (i) a mutant in

the Tudor domain (E134K) and (ii) a mutant in the YG domain (Y272C). The Tudor domain is

important for the interaction of SMN with FBL and Coilin [21], while the YG domain is essential for

the oligomerization of SMN [26] which is more stable than the monomeric form.

We investigated whether these mutated SMN might have a defect in nucleolar

reorganization after DNA damage induction or in the shuttling of SMN to and/or from the

nucleolus. We produced three GFP-SMN cell lines by transfecting into the inducible sh6-SMN

cell line (i) a wild-type GFP-tagged SMN (hereafter GFP-SMN); (ii) a mutant GFP-tagged SMN,

bearing the mutation in the Tudor domain E134K (hereafter GFP-SMN[E134K]); (iii) a mutant GFP-

tagged SMN, bearing the mutation in the YG domain Y272C (hereafter GFP-SMN[Y272C]). After the

selection of suitable clones, we verified the amount of GFP-SMN and the mutant SMN proteins

produced and we could determine that the amount of mutant SMN was lower than the GFP-SMN

protein (Figure S11A). After depletion of the endogenous SMN, by induction of the sh6-SMN

RNA, we UV-irradiated SMN wild type and mutant expressing clones and we performed IF against

RNAP1 coupled to the direct visualization of the GFP-SMN, GFP-SMN^{E134K} or GFP-SMN^{Y272C}

signal. Wild-type and mutant SMN proteins could be visualized in CBs/Gems in undamaged cells

(Figures S11B and S11C). Nevertheless, the Tudor mutant GFP-SMN^{E134K} expressing cells failed

to restore CBs/Gems at 48h PUVI and the nucleolar reorganization was not restored, RNAP1

remaining at the periphery of the nucleolus at 48h PUVI (Figures S11B, S11C and S11D).

Interestingly, these results are reminiscent of the results observed in the absence of FBL (Figures

5A, 5B and 5C). The YG mutant GFP-SMN^{Y272C} failed to properly shuttle to the nucleolus and

remained mostly in CBs/Gems all along the nucleolar reorganization process (Figures S11B and

S11C) and the nucleolar reorganization was not restored at 48h PUVI (Figures S11B and S11D).

These results are reminiscent of the results observed in the absence of COILIN (Figures 4A, 4B

and 4C). To resume, both SMN mutants studied, show that a functional SMN is needed to

properly ensure nucleolar homeostasis after DNA damage induction and that in different ways

SMN shuttling is perturbed by these mutations.

PRMT1 activity mediates the nucleolar shuttling of SMN

One of the activities of SMN is to bind, via the Tudor domain, Arginine methylated

proteins [27,28]. Arginine methylation is a widespread post-translational modification that can occur

in histones and non-histone proteins [29–32]. The enzymes catalyzing the transfer of a methyl group

to Arginine residues are part of a family called the PRMTs (Protein Arginine Methyl Transferases).

PRMTs can mono-methylate Arginine residues (MMA) or di-methylate Arginine residues either

symmetrically (SDMA) or asymmetrically (ADMA). Because these proteins affect SMN functions

but also the interaction of SMN with Coilin [33], we wondered whether one of the PRMTs could

affect, disturb or enhance SMN shuttling after DNA damage. Using GST pull-down assay, we

observed that PRMT1 interacts with SMN (Figure S12A). We thus decided to deplete cells from

PRMT1 by siRNA silencing (Figure S12B) and performed IF against SMN and FBL. We found that

PRMT1 inhibited the shuttling of SMN to the nucleolus (Figures 6A and 6B). In fact, in PRMT1-depleted cells, SMN remains mostly in CBs/Gems. This SMN localization after DNA damage is reminiscent of the one observed in Coilin-depleted cells (Figure 4A).

To verify whether the perturbation of SMN shuttling observed in Figure 6A is due to the physical depletion of PRMT1 or the inhibition of the ADMA methylase activity, we treated the cells with the PRMT-class I specific inhibitor MS023 [34] and the more specific and potent PRMT1 inhibitor, Furamidine[35] (Figures 6D, 6E, S12C and S12D) prior to DNA damage and IF assays. We could verify that the inhibition of the methylase activity of PRMTs from class I and specifically of PRMT1 perturbs SMN shuttling (Figures 6D, 6E and S12C) and that SMN remains mostly in CBs/Gems as when PRMT1 is depleted (Figure 6A). Because the SMN complex requires the protein arginine methyltransferase 5 (PRMT5) to assemble Sm core structures of spliceosomal U-rich small nuclear ribonucleoprotein particles[36], we investigated whether depletion of PRMT5 would affect SMN shuttling to the nucleolus. We down-regulated PRMT5 by siRNA (Figure S13D), induced DNA damage and perform IF against SMN and RNAP1 at different times PUVI (Figure S13A). Our results show that depletion of PRMT5 does not influence SMN shuttling (Figures S13A and S13B), nor the nucleolar homeostasis after DNA damage and repair (Figures S13A and S13C).

To investigate how the depletion of PRMT1 affects the interactions between SMN and its partners (Coilin and FBL), we performed PLA assays and measured a stronger interaction of SMN and FBL before damage induction and at 3h PUVI which corresponds with the beginning of SMN shuttling at the nucleolus (Figures 7A and 7B). We also quantified a stronger interaction of SMN with Coilin when PRMT1 is depleted (Figures 7C and 7D) before damage induction and up to 48h PUVI.

Remarkably, we showed by IF that PRMT1 is also shuttling during nucleolar reorganization in wild-type cells and could be detected at 6h and up to 48h PUVI within the nucleolus or at the periphery of the nucleolus (Figures S14A and S14B). Importantly, the shuttling of PRMT1 at the nucleolus is dependent on the SMN protein as in SMN-depleted cells (Sh6-SMN), PRMT1 is not detected inside the nucleoli but remains mainly nuclear or localized at the periphery of the nucleolus (Figures S14A and S14B).

443   Interestingly, we could quantify an increased level of PRMT1 after damage induction and

we could show that this increase is partially dependent on the presence of a functional SMN

(Figures S14C, S15A and S15B).

Figure 6: PRMT1 activity mediates the nucleolar shuttling of SMN

(A) Representative microscopy images of immunofluorescence (IF) assay showing the localization

of SMN (red) and FBL (green) at different times PUVI in cells transfected with siMock or siPRMT1

pool. Nuclei and nucleoli are indicated by dashed lines and dotted lines respectively. Scale bar:

5µm. (B and C) Quantification of cells number for localization of (B) SMN (in Cajal Bodies [CBs]

or Gems, at the periphery of the nucleolus or mixed localization) and (C) FBL (inside the

nucleolus, outside the nucleolus or mixed localization) at different times PUVI in cells transfected

with siMock or siPRMT1 pool. (D) Representative microscopy images of IF assay showing the

localization of SMN (red) and FBL (green) in MRC5 cells treated with DMSO, MS023 or

Furamidine followed by 16J/m² UV-C irradiation. Nuclei and nucleoli are indicated by dashed

lines and dotted lines respectively. Scale bar: 5µm. (E) Quantification of cells number for

localization of SMN (in Cajal Bodies [CBs] or Gems, at the periphery of the nucleolus or mixed

localization) at different times PUVI in cells treated with DMSO, MS023 or Furamidine.

Figure 7: PRMT1 remodels the interaction of SMN with FBL and COILIN

(A and C) Representative microscopy images of PLA showing the interaction between SMN and

FBL (A) or between SMN and COILIN (C) at different times PUVI in cells transfected with siMock

or siPRMT1 pool. Scale bar: 5µm. (B and D) Quantification of fluorescent signal in the nucleus

against the couple SMN-FBL (B) or SMN-COILIN (D) from PLA experiment in cells transfected

with siMock or siPRMT1 pool after UV-C irradiation. P-value of Student's test compared to

*siMock condition. ***<0.001.*

SMN cells are sensitive to DNA damage

We demonstrated that a functional SMN is important for proper nucleolar homeostasis

after DNA damage repair and wondered whether this newly discovered function could have a

biological consequence on SMN cells and *in fine* on SMA patients. In order to investigate whether

DNA damage could impact the survival of SMN cells we performed clonogenic assays on sh6-

SMN cells with and without induction of the sh expression and compared the results with a DNA

repair-deficient cell lines (CS1AN-SV) mutated in the CSB protein, involved in transcription

coupled-NER (Figure 8A). Our results show that SMN-depleted cells are moderately sensitive to

UV damage as their sensitivity to UV irradiation is intermediate between an SMN wild-type cell

line and a DNA repair-deficient cell line.

Because we demonstrated that SMN shuttling is also happening during oxidative damage

repair (Figure S6A), we wondered whether SMN cells would be sensitive to chronic oxidative

damage. We usually culture rodents, human patients or mutated cell lines in physioxia conditions

(3% of oxygen) to avoid the supraphysiological oxygen levels disturbing normal cellular

metabolism and increasing the cellular concentration of reactive oxygen species [37]. To induce

chronic oxidative damage in SMN-depleted cells, we culture them in a standard incubator at 20%

oxygen and compare their ability to form colonies. As with many established cell lines, Human

SV40-immortalized fibroblasts MRC5 (MRC5-SV) have a better clonogenicity in physioxia

conditions (Figure 8B). However, MRC5-SV expressing shRNA SMN has a decreased

clonogenicity compared to MRC5-SV that expresses a mock shRNA (Figure 8B), which

demonstrates that the absence of SMN *per se* hinders the ability of cells to form clones.

Importantly, when SMN-depleted cells are grown in supraphysiological oxygen levels their

clonogenicity is more drastically reduced, compared with the physioxia culture conditions (Figure

8C).

Figure 8: SMN-deficient cells are sensitive to DNA damage

(A) Survival curve as determined by the colony-forming ability to UV-C of cells expressing (+Dox) or not (-

Dox) Sh6-SMN. (B) Clonogenic assay of cells cultured with different quantities of oxygen (3% or 20%) in

the presence (MRC5 or ShSCRB) or absence of SMN (Sh5-SMN and Sh6-SMN).

DISCUSSION

One of the most fascinating and nearly unexplored areas in the DNA repair field is how cells restore their cellular activities after the completion of all the reactions that allow cells to eliminate DNA lesions. Most DNA lesions block transcription and replication [5] and although we have extensive knowledge of how cells recognize and repair DNA lesions, very little is known about how cells restart these cellular processes. In post-mitotic cells, restoration of the damage-induced block of transcription is essential for cell survival.

We have previously shown that RNAP1 transcription is blocked after UV lesions and that TC-NER pathway is responsible for the repair of UV-lesions on ribosomal DNA [8]. Importantly, UV-damages impact the organization of the nucleolus and during DNA repair both nucleolar DNA and RNAP1 are displaced at the periphery of the nucleolus [8]. Interestingly, although RNAP1 transcription restarts when UV-lesions on the transcribed strand are repaired, the return of the RNAP1 within the nucleolus is dependent on the proper repair of DNA lesions on the untranscribed nucleolar DNA [8]. In this particular case (GG-NER deficient cells), RNAP1 transcription restarts in a non-canonical compartment, the periphery of the nucleolus, and this anomaly might influence the proper ribosome biogenesis. Therefore, restoring the proper nucleolar structure and organization might be important for cellular viability or the efficiency of cellular processes. The recovery of a normal nucleolar structure after DNA damage repair is not a passive process and requires the presence of some key proteins [9], although their exact mechanistic role has not been established yet.

In a quest to find the exact molecular mechanism for the re-establishment of the nucleolar organization after DNA repair completion, we set up a best-candidate approach that guided us to inspect the consequences of different nucleolar proteins' interactors depletion. One of these candidates was the protein SMN, which is a particularly interesting protein to scrutinize because of the known interaction with the nucleolar protein FBL via its Tudor domain [17].

SMN is part of a multiprotein complex that includes Gemins2-8 and Unrip [38] and acts like a molecular chaperone needed to assemble small ribonucleoproteins (snRNPs) which are part of the spliceosome [39] and small nucleolar ribonucleoprotein particles (snoRNPs) involved in

posttranscriptional processing of ribosomal RNA [21]. Additionally, SMN has been also found implicated in transcription [40], histone mRNA processing [41], mRNAs trafficking [42] and translation [43].

We have shown here that SMN plays a role in nucleolar homeostasis after DNA repair completion. In fact, silencing or mutations (Tudor domain and oligomerization mutants) of the SMN protein hinder the proper return of RNAP1 within the nucleolus (Figure 9), despite the restart of RNAP1 transcription (Figure 1, Figures S2, S3 and S11). Contrary to what has been observed in GG-NER deficient cells, the RNAP1 mislocalization after DNA repair completion is not induced by the presence of unrepaired DNA lesions because we could show that SMN deficient cells are proficient in the NER pathway, repairing UV-lesions on both transcribed and untranscribed regions of the genome (Figure S4).

The multifunctional SMN is localized in both the cytoplasm and in nuclear bodies (Cajal Bodies and Gems) and in addition, SMN can also be detected in the nucleolus of neurons [16] although the function of SMN in nucleolus remains enigmatic. Interestingly, during the process of nucleolar reorganization, we observed a shuttling of SMN from CBs/Gems to the periphery of the nucleolus or within the nucleolus during RNAP1 displacement and a return to CBs/Gems after the RNAP1 repositioning (Figures 2, 9). SMN shuttles to the nucleolus together with its partners Gemin 2, Gemin 3, Gemin 4 and Gemin 5 (Figure 3). While Gemin 3 and 4 followed the same SMN localization, the classical SMN partner Gemin 2 [44] was observed to have both CBs/Gems and nucleolar localization. While SMN/Gemin2 foci disappeared during RNAP1 displacement, the nucleolar signal of Gemin 2 increased reaching the maximum after RNAP1 repositioning and Gemin2 foci did not reappear. This result might suggest that SMN complex is not exactly the same in CBs/Gems when performing snRNPs assembly and in the nucleolus during DNA repair-induced nucleolar reorganization. These results also suggest that the restoration of fully functional CBs/Gems containing Gemin2 might be delayed in time, probably further delaying the restoration of the other functions of the SMN complex. Notably, Gemin 5 was observed in both SMN positive and negative foci which could be because a large fraction of Gemin5 protein is found outside of the SMN complex [45]. Interestingly, SMN negative foci of Gemin5 were present at any time point throughout the nucleolar reorganization process, which might indicate that SMN

negative foci of Gemin5 do not participate in the shuttling of SMN complex and in nucleolar homeostasis.

SMN shuttling happens in replicative and post-mitotic cells (motoneurons) and in different stress situations: UV-induced DNA damage, RNAP1 transcriptional block and drug-induced oxidative damage. Notably, we should highlight that the kinetics of SMN shuttling might vary depending on different types of stresses but also on the culture conditions, the amount and quality of serum, supraphysiological oxygen levels and the type of culture media. The kinetic of SMN shuttling should be defined for each type of stress/damage and culture condition.

SMN shuttling is dependent on both Coilin and FBL (Figures 4, 5 and 9). In Coilin-depleted undamaged cells, we observed a high number of cells in which no CBs or other foci were visible, this might be because, without Coilin, CBs are disrupted and just Gems are visible. In the absence of Coilin, we observe an increased number of Gems and the number of cells in which SMN is at the nucleolus is reduced. Interestingly, in Coilin-depleted cells RNAP1 repositioning is hindered, indicating that in the absence of Coilin nucleolar reorganization is compromised. In FBL-depleted cells, SMN remains at the periphery of the nucleolus at all time points and RNAP1 does not recover the proper localization within the nucleolus but remains at the periphery of the nucleolus.

Because FBL interacts with SMN via its Tudor domain [17] and SMN interacts mainly with methylated arginine residues, we explored the possibility that one of the PRMTs would be responsible for SMN shuttling within the nucleolus. Indeed, the activity of PRMT1, responsible for ADMA, is essential to recruit SMN to the nucleolus (Figures 6 and 9). In contrast, depletion of PRMT5, which is responsible for the assembly of Sm proteins to the SMN complex did not influence the shuttling of SMN, probably indicating that the SMN function in nucleolar homeostasis is independent of the snRNPs biogenesis process[46] (Figure S13).

Interestingly, depletion of Coilin, FBL and PRMT1 also modifies the interactions between SMN, Coilin and FBL, by increasing their proximities (Figures 5, 6 and 7) which might indicate that specific methylations are important to maintain an equilibrium of interactions between these three partners.

Remarkably PRMT1 shuttles within the nucleolus after DNA damage induction as SMN

(Figure 7). This shuttling within the nucleolus is SMN-dependent. Although we do not know if the

substrate of PRMT1 is a protein that directly interacts with SMN, we showed that methylation is

essential for SMN shuttling. It is plausible to assume that because FBL is methylated by PRMT1

[47], FBL might be the substrate methylated during the process of RNAP1 repositioning within the

nucleolus after the completion of DNA repair.

Figure 9: SMN shuttling during DNA damage/repair dependent nucleolar reorganization.

In the future years, it will be important to unveil the precise mechanism of action of SMN

within the nucleolus, by investigating what are the partners of SMN in this organelle after DNA

damage and how they function together with SMN, how CBs/Gems disassemble and reassemble,

how they restore their activity and what are the consequences on splicing, ribosome biogenesis

and finally translation fidelity. More generally, it will be important to continue to study how cells

maintain and/or restore nucleolar homeostasis after cellular stresses, such as DNA damage.

We believe that our study will also open new avenues of research that could improve the

well-being of SMA patients. As for other multifunctional proteins, it is difficult to identify exactly

which function of SMN is responsible for the phenotype observed in SMA patients. Likely, the

SMA phenotype could be due to a combination of different defects that would synergistically

impact cellular activities. Our results present here a newly discovered SMN function in

maintaining nucleolar homeostasis after DNA damage and repair. Importantly, we showed that

SMN-depleted cells show a moderate but remarkable sensitivity to UV-induced damage and

most importantly to chronic oxidative damage (Figure 8) which is what motoneurons would have

to deal with all along patients' life. This result is also important because they point to the fact that

not just the DNA repair process is important for cell viability but also all the processes that restore

cellular functions after DNA repair completion.

Our findings may directly impact the life and well-being of SMA patients [48]. In fact, in cells

and motoneurons of SMA patients, exogenous and endogenous DNA damage might

progressively and lastingly disrupt the nucleolar structure and disturb ribosome biogenesis

leading to perturbed protein translation [49]. This defect may contribute to the neurodegenerative

phenotype of SMA motoneurons. If this hypothesis is true, SMA patients could be advised to

prevent deleterious DNA damage to avoid a reorganization of the nucleolus that would affect

proper protein production within neurons. Although exogenous DNA damage is partially

avoidable, endogenous DNA damage is inevitable and one way to reduce its overload is to follow

a diet rich in antioxidants. This nutritional approach combined with a healthy lifestyle, avoiding

exogenous damage, such as cigarette smoke, pollutants, and potentially harmful molecules, may

retard the degeneration of motoneurons and thus SMA progression.

MATERIALS AND METHODS

Cells culture

SV40-immortalized human fibroblasts, wild-type (MRC5-SV) and CSB-deficient (CS1AN-SV), were cultured in DMEM supplemented with 10% fetal bovine serum (FBS) and 1% penicillin and streptomycin (P/S) and incubated at 37°C with 20% O2 and 5% CO2.

The MRC5-SV + Sh cells were obtained by transduction of lentiviral particles produced (as described https://www.addgene.org/protocols/plko/#Ehttps://www.addgene.org/protocols/plko/ - E) from piSMART hEF1α/turboGFP (Dharmacon) doxycycline-inducible lentiviral system containing a Short Hairpin (Sh) Scramble (VSC6572), Sh5-SMN (V3IHSHEG_4923340; mature antisense: TAAACTACAACACCCTTCT) or Sh6-SMN (V3IHSHEG_5297527; mature antisense: TTCAAATTTTCTCAACTGC). ShSMN both target telomeric *SMN1* and centromeric *SMN2* copies of the gene. The cells were cultured in DMEM supplemented with 10% FBS and 1% P/S; maintained in 100 ng/ml puromycin. Sh expression is induced with 100 ng/ml doxycycline. The cells were incubated at 37°C with 20% O_2 and 5% CO_2.

Treatment

DNA damage was inflicted by UV-C light (254 nm, 6-Watt lamp). Cells were globally irradiated with a UV-C dose of 16 J/m² or locally irradiated with a UV-C dose of 100 J/m² through a filter with holes of 5µm. After irradiation, cells were incubated at 37°C with 5% CO2 for different periods of time.

The PRMT inhibitor, MS023 (ab223596) diluted in DMSO, was added at 1µM in the medium 20h before irradiation. The PRMT1 inhibitor, Furamidine (ab287098) diluted in DMSO, was added at 1µM in the medium 15h before irradiation.

Transfection of small interfering RNAs (siRNAs)

Cells were seeded in six-well plates with coverslip and allowed to attach. Cells were transfected two times in an interval of 24h with siRNA using Lipofectamine® RNAiMAX reagent (Invitrogen; 13778150) according to the manufacturer' protocol. Experiments were performed

between 24h and 72h after the second transfection. Protein knock-down was confirmed for each

experiment by western blot. The small interfering RNA (siRNAs) used in this study are: siMock,

Horizon, D-001206-14 (10nM); siCOILIN, Horizon, M-019894-01 (5nM); siFBL, Horizon, L-011269-

00 (10nM); siPRMT1, Horizon, L-010102-00 (10nM). The final concentration used for each siRNA

is indicated in parentheses. All siRNA are a pool of four different siRNA.

Protein extraction

Cells were collected using trypsin or by scraping and centrifuged 10 min at 1400 rpm. To

verify siRNA efficiency, the coverslip need for the experiment was displaced before fixation and

cells that remained in the dish were collected. The extraction of total proteins was performed

using the PIERCE RIPA buffer (Thermo, #89900) complemented with EDTA-free cOmplete PIC

(ROCHE).

Western blot

Proteins were separated on a SDS-PAGE gel composed of bisacrylamide (37:5:1), and

then transferred onto a polyvinylidene difluoride membrane (PVDF, 0.45 μm; Millipore). The

membrane was blocked in PBS-T (PBS and 0.1 % Tween 20) with 5 % milk and incubated for 2h

at room temperature (RT) or overnight at 4°C with the primary antibodies diluted in milk PBS-T

(see table of antibody). Subsequently, the membrane was washed with PBS-T (3× 5-10 min) and

incubated with the following secondary antibody diluted 1/5000 in milk PBS-T: Goat anti-rabbit

IgG HRP conjugate (170-6515; BioRad) or Goat anti-mouse IgG HRP conjugate (170-6516;

BioRad). After the same washing procedure, protein bands were visualized via

chemiluminescence (ECL Enhanced Chemo Luminescence; Pierce ECL Western Blotting

Substrate) using the ChemiDoc MP system (BioRad).

Cytostripping

To improve the nuclear signal of immunofluorescence signal, the cytoplasm of the cells

was removed before fixation. After two washes with cold PBS, cells were incubated on ice 5 min

with cold cytoskeleton buffer (10mM PIPES pH6,8; 100mM NaCl; 300mM Sucrose; 3mM MgCl2;

1mM EGTA; 0,5% Triton-X100) followed by 5min with cold cytostripping buffer (10mM Tris HCL

pH7,4; 10mM NaCl; 3mM MgCl2; 1% Tween 40; 0,5% sodium deoxycholate). After three gentle

washes with cold PBS, cells were fixed.

Immunofluorescence

Cells were grown on coverslips, washed with PBS at RT, and fixed with 2%

paraformaldehyde (PFA) for 15min at 37° C. Cells were permeabilized with PBS 0.1 % Triton X-

100 (3X short + 2X 10 min washes). Blocking of non-specific signals was performed with PBS+

(PBS, 0.5 % BSA, 0.15 % glycine) for at least 30 min. Then, coverslips were incubated with primary

antibody diluted in PBS+ for 2h at room temperature (RT) or overnight at 4°C in a moist chamber.

After several washes with PBS + 0.1% Triton X-100 (3X short + 2X 10 min) and a quick washed

with PBS+, cells were incubated for 1h at RT in a moist chamber with the following secondary

antibody coupled to fluorochrome and diluted 1/400 in PBS+ : Goat anti-mouse Alexa Fluor®

488 [A11001, Invitrogen] or 594 [A11005] and Goat anti-rabbit Alexa Fluor® 488 [A11008] or 594

[A11012]. After the same washing procedure but with PBS, coverslips were finally mounted using

Vectashield with DAPI (Vector Laboratories).

Proximity ligation assay

PLA experiments were done using Duolink™ II secondary antibodies and detection kits

(Sigma-Aldrich, #DUO92002, #DUO92004, and #DUO92008) according to the manufacturer's

instructions. In brief, cells were fixed and permeabilized with the same procedure as

immunofluorescence followed by incubation in PLA blocking buffer for 1h at 37 °C. After

blocking, cells were incubated overnight at 4°C with primary antibodies diluted in PLA Antibody

Diluent. After washes with PLA buffer A (1 short + 3x5min), cells were incubated with PLUS and

MINUS PLA probes for 1h at 37 °C. After the same washing procedures with PLA buffer A, if

probes were in close proximity (<40 nm), they were ligated together to make a closed circle

thanks to the incubation of 30min at 37°C with the Duolink™ ligation solution. Then, after the

same washing procedures, the DNA is amplified and detected by fluorescence 594 thanks to the

incubation of 100min at 37°C with the Duolink™ amplification solution. After washing with PLA buffer

B (1 short + 2x10min), coverslips were mounted using Vectashield with DAPI (Vector Laboratories).

RNA Fluorescence In Situ Hybridization (RNA-FISH)

Cells were grown on coverslips and globally irradiated for different times. Then, cells were

washed with PBS at RT, and fixed with 4% PFA for 15min at 37° C. After two washes with PBS,

cells were permeabilized with PBS+ 0.4 % Triton X-100 for 7min at 4° C. Cells were washed

rapidly with PBS before incubation for at least 30min with pre-hybridization buffer (15%

formamide in 2X SSPE pH8.0 [0.3M NaCl, 15.7mM $NaH_2PO_4.H_2O$ and 2.5mM EDTA]). 35ng of

probe was diluted in 70µl of hybridization mix (2X SSPE, 15% formamide, 10% dextran sulfate

and 0.5mg/ml tRNA). Hybridization of the probe was conducted overnight at 37°C in a humidified

environment. Subsequently, cells were washed twice for 20 min with prehybridization buffer, then

once for 20 min with 1X SSPE. After extensive washing with PBS, the coverslips were mounted

with Vectashield containing DAPI (Vector Laboratories). The probe sequence (5' to 3') is Cy5-

AGACGAGAACGCCTGACACGCACGGCAC. At least 30 cells were imaged for each condition

of each cell lines.

Primary Antibodies

The following primary antibodies were used:

Antibody against	Manufacturer	Catalog Nr	Source	IF/PLA	WB
Alpha-Tubulin	Sigma	T6074	Mouse		1/50 000
Coilin	Proteintech	10967-1-AP	Rabbit	1/500	1/5000
CSB	Santa Cruz Biotechnology	sc398022	Mouse		1/100
FBL	Abcam	ab5821	Rabbit	1/500	1/500
GEMIN2	ABclonal	A3082	Rabbit	1/500	
GEMIN3	Santa Cruz Biotechnology	sc374373	Mouse	1/1000	
GEMIN4	Santa Cruz Biotechnology	sc365424	Mouse	1/1000	

GEMIN5	Proteintech	24897-1-AP	Rabbit	1/500	
PRMT1	Abcam	ab190892	Rabbit	1/5000	1/1000
RNAP1	Santa-cruz	sc48385	Mouse	1/500	
SMN	Proteintech	11708-1-AP	Rabbit	1/100	
SMN	BD Biosciences	610646	Mouse	1/500	1/5000

Images acquisition and analysis and statistics

Confocal images of the cells were obtained on a Zeiss LSM 880 NLO confocal laser
scanning inverted microscope using either a Plan-Apochromat 63x/1.4 or 40x/1.3 oil immersion
objective. Other images were obtained using an upright Zeiss Z1 Imager with a Plan-Apochromat
40x/0.95 objective. Time-lapse imaging of live cells for 48 hours was achieved on a Zeiss
AxioObserver 7 microscope with an Alpha Plan-Apochromat 63x/1.46 oil objective, equipped
with a dedicated incubator system set at 37°C and 5% CO_2.

Images of the cells for each experiment were obtained with the same microscopy system
and constant acquisition parameters. Images were analyzed with Image J software. For all images
of this study, nuclei and nucleoli were delimited with dashed and dotted lines respectively, using
DAPI staining.

Protein localization data represented as composite bar graphs was performed by manually
sorting cells based on protein localization. The number of cells analyzed per time point or
condition was between 100 and 300 cells, except for primary cells, motoneuons and GFP-SMN
expressing cells (50 to 100 cells).

All experiments have been performed at least three times (biological replicates). Error
bars represent the Standard Error of the Mean (SEM) of the biological replicates. Excel was used
for statistical analysis and plotting of all the numerical data. Statistics were performed using a
Student's test to compare two different conditions (siMock vs. siRNA X or No UV irradiation vs.
after irradiation) with the following parameters: two-tailed distribution and two-sample unequal
variance (heteroscedastic).

Acknowledgments

751   Dedicated to the mothers and fathers of children affected with SMA. We would like to

752   thank Laura Trinkle-Mulcahy for the generous gift of the mCherry-N1. For access to the confocal

753   microscope, we acknowledge the contribution of the imaging facility CIQLE (https://ciqle.univ-

754   lyon1.fr - Centre d'imagerie Quantitative Lyon-Est - SFR Santé Lyon-Est (UAR3453 CNRS, US7

755   INSERM, UCBL), a member of LyMIC (Lyon Multiscale Imaging Center).

Conflicts of interest

758   The authors disclose no potential conflict of interest.

Funding

This work was supported by AFM-Téléthon (MyoNeurALP 1 and 2) and by Institut National du

Cancer (InCa, PLBIO19-126).

SM was funded by Jazan university, Jazan, Saudi Arabia.

PL and OB are funded by an AFM-Téléthon grant (MyoNeurALP) and by the Joint Collaborative

Research Program between the Centre for Neuromuscular Disease (University of Ottawa) and

Institut NeuroMyoGène (INMG - Claude Bernard Université Lyon 1).

JC is funded by a Canadian Institutes of Health Research (CIHR) grant (MOP 123381) and

CureSMA Canada.

REFERENCES

1. Dubois, M.-L. & Boisvert, F.-M. The Nucleolus: Structure and Function. in *The Functional Nucleus* (eds. Bazett-Jones, D. P. & Dellaire, G.) 29–49 (Springer International Publishing, 2016). doi:10.1007/978-3-319-38882-3_2.

2. Hernandez-Verdun, D. Assembly and disassembly of the nucleolus during the cell cycle. *Nucl. Austin Tex* **2**, 189–194 (2011).

3. Boulon, S., Westman, B. J., Hutten, S., Boisvert, F.-M. & Lamond, A. I. The nucleolus under stress. *Mol. Cell* **40**, 216–227 (2010).

4. Louvet, E., Junéra, H. R., Le Panse, S. & Hernandez-Verdun, D. Dynamics and compartmentation of the nucleolar processing machinery. *Exp. Cell Res.* **304**, 457–470 (2005).

5. Giglia-Mari, G., Zotter, A. & Vermeulen, W. DNA damage response. *Cold Spring Harb. Perspect. Biol.* **3**, a000745 (2011).

6. Brooks, P. J. The case for 8,5'-cyclopurine-2'-deoxynucleosides as endogenous DNA lesions that cause neurodegeneration in xeroderma pigmentosum. *Neuroscience* **145**, 1407–1417 (2007).

7. Kumar, N. *et al.* Global and transcription-coupled repair of 8-oxoG is initiated by nucleotide excision repair proteins. *Nat. Commun.* **13**, 974 (2022).

8. Daniel, L. *et al.* Mechanistic insights in transcription-coupled nucleotide excision repair of ribosomal DNA. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E6770–E6779 (2018).

9. Cerutti, E. *et al.* β-Actin and Nuclear Myosin I are responsible for nucleolar reorganization during DNA Repair. 646471 Preprint at https://doi.org/10.1101/646471 (2019).

10. Talbot, K. & Tizzano, E. F. The clinical landscape for SMA in a new therapeutic era. *Gene Ther.* **24**, 529–533 (2017).

11. Monani, U. R. Spinal muscular atrophy: a deficiency in a ubiquitous protein; a motor neuron-specific disease. *Neuron* **48**, 885–896 (2005).

12. Calucho, M. *et al.* Correlation between SMA type and SMN2 copy number revisited: An analysis of 625 unrelated Spanish patients and a compilation of 2834 reported cases. *Neuromuscul. Disord. NMD* **28**, 208–215 (2018).

13. Singh, R. N., Howell, M. D., Ottesen, E. W. & Singh, N. N. Diverse role of survival motor neuron protein. *Biochim. Biophys. Acta Gene Regul. Mech.* **1860**, 299–315 (2017).

14. Trinkle-Mulcahy, L. & Sleeman, J. E. The Cajal body and the nucleolus: 'In a relationship' or 'It's complicated'? *RNA Biol.* **14**, 739–751 (2017).

15.   Hebert, M. D., Szymczyk, P. W., Shpargel, K. B. & Matera, A. G. Coilin forms the bridge
between Cajal bodies and SMN, the spinal muscular atrophy protein. *Genes Dev.* **15**, 2720–
2729 (2001).

16.   Wehner, K. A. *et al.* Survival motor neuron protein in the nucleolus of mammalian
neurons. *Brain Res.* **945**, 160–173 (2002).

17.   Jones, K. W. *et al.* Direct interaction of the spinal muscular atrophy disease protein SMN
with the small nucleolar RNA-associated protein fibrillarin. *J. Biol. Chem.* **276**, 38645–38651
(2001).

18.   Renvoisé, B. *et al.* Distinct domains of the spinal muscular atrophy protein SMN are
required for targeting to Cajal bodies in mammalian cells. *J. Cell Sci.* **119**, 680–692 (2006).

19.   Thandapani, P., O'Connor, T. R., Bailey, T. L. & Richard, S. Defining the RGG/RG motif.
Mol. Cell **50**, 613–623 (2013).

20.   Jacquier, A. *et al.* Severe congenital myasthenic syndromes caused by agrin mutations
affecting secretion by motoneurons. *Acta Neuropathol. (Berl.)* **144**, 707–731 (2022).

21.   Pellizzoni, L., Baccon, J., Charroux, B. & Dreyfuss, G. The survival of motor neurons
(SMN) protein interacts with the snoRNP proteins fibrillarin and GAR1. *Curr. Biol. CB* **11**, 1079–
1088 (2001).

22.   Whitehead, S. E. *et al.* Determinants of the Interaction of the Spinal Muscular Atrophy
Disease Protein SMN with the Dimethylarginine-modified Box H/ACA Small Nucleolar
Ribonucleoprotein GAR1 *. *J. Biol. Chem.* **277**, 48087–48093 (2002).

23.   Cioce, M., Boulon, S., Matera, A. G. & Lamond, A. I. UV-induced fragmentation of Cajal
bodies. *J. Cell Biol.* **175**, 401–413 (2006).

24.   Kawanishi, S. & Murata, M. Mechanism of DNA damage induced by bromate differs
from general types of oxidative stress. *Toxicology* **221**, 172–178 (2006).

25.   Tessarz, P. *et al.* Glutamine methylation in histone H2A is an RNA-polymerase-I-
dedicated modification. *Nature* **505**, 564–568 (2014).

26.   Gupta, K. *et al.* Assembly of higher-order SMN oligomers is essential for metazoan
viability and requires an exposed structural motif present in the YG zipper dimer. *Nucleic Acids*
Res. **49**, 7644–7664 (2021).

27.   Friesen, W. J., Massenet, S., Paushkin, S., Wyce, A. & Dreyfuss, G. SMN, the product of
the spinal muscular atrophy gene, binds preferentially to dimethylarginine-containing protein
targets. *Mol. Cell* **7**, 1111–1117 (2001).

28.   Sprangers, R., Groves, M. R., Sinning, I. & Sattler, M. High-resolution X-ray and NMR

structures of the SMN Tudor domain: conformational variation in the binding site for

symmetrically dimethylated arginine residues. *J. Mol. Biol.* **327**, 507–520 (2003).

29.   Fulton, M. D., Brown, T. & Zheng, Y. G. Mechanisms and Inhibitors of Histone Arginine

Methylation. *Chem. Rec. N. Y. N* **18**, 1792–1807 (2018).

30.   Bedford, M. T. & Richard, S. Arginine methylation an emerging regulator of protein

function. *Mol. Cell* **18**, 263–272 (2005).

31.   Di Lorenzo, A. & Bedford, M. T. Histone arginine methylation. *FEBS Lett.* **585**, 2024–

2031 (2011).

32.   Lee, Y.-H. & Stallcup, M. R. Minireview: protein arginine methylation of nonhistone

proteins in transcriptional regulation. *Mol. Endocrinol. Baltim. Md* **23**, 425–433 (2009).

33.   Hebert, M. D., Shpargel, K. B., Ospina, J. K., Tucker, K. E. & Matera, A. G. Coilin

methylation regulates nuclear body formation. *Dev. Cell* **3**, 329–337 (2002).

34.   Eram, M. S. *et al.* A Potent, Selective, and Cell-Active Inhibitor of Human Type I Protein

Arginine Methyltransferases. *ACS Chem. Biol.* **11**, 772–781 (2016).

35.   Yan, L. *et al.* Diamidine compounds for selective inhibition of protein arginine

methyltransferase 1. *J. Med. Chem.* **57**, 2611–2622 (2014).

36.   Prusty, A. B. *et al.* Impaired spliceosomal UsnRNP assembly leads to Sm mRNA down-

regulation and Sm protein degradation. *J. Cell Biol.* **216**, 2391–2407 (2017).

37.   Stuart, J. A. *et al.* How Supraphysiological Oxygen Levels in Standard Cell Culture Affect

Oxygen-Consuming Reactions. *Oxid. Med. Cell. Longev.* **2018**, 1–13 (2018).

38.   Otter, S. *et al.* A Comprehensive Interaction Map of the Human Survival of Motor

Neuron (SMN) Complex. *J. Biol. Chem.* **282**, 5825–5833 (2007).

39.   Pellizzoni, L., Yong, J. & Dreyfuss, G. Essential Role for the SMN Complex in the

Specificity of snRNP Assembly. *Science* **298**, 1775–1779 (2002).

40.   Zhao, D. Y. *et al.* SMN and symmetric arginine dimethylation of RNA polymerase II C-

terminal domain control termination. *Nature* **529**, 48–53 (2016).

41.   Tisdale, S. *et al.* SMN is essential for the biogenesis of U7 small nuclear

ribonucleoprotein and 3'-end formation of histone mRNAs. *Cell Rep.* **5**, 1187–1195 (2013).

42.   Fallini, C., Donlin-Asp, P. G., Rouanet, J. P., Bassell, G. J. & Rossoll, W. Deficiency of the

Survival of Motor Neuron Protein Impairs mRNA Localization and Local Translation in the

Growth Cone of Motor Neurons. *J. Neurosci. Off. J. Soc. Neurosci.* **36**, 3811–3820 (2016).

43.   Sanchez, G. *et al.* A novel function for the survival motoneuron protein as a translational

regulator. *Hum. Mol. Genet.* **22**, 668–684 (2013).

44.   Cauchi, R. J. SMN and Gemins: 'We are family' … or are we?: Insights into the
partnership between Gemins and the spinal muscular atrophy disease protein SMN. *BioEssays*
32, 1077–1089 (2010).

45.   Martinez-Salas, E., Embarc-Buh, A. & Francisco-Velilla, R. Emerging Roles of Gemin5:
From snRNPs Assembly to Translation Control. *Int. J. Mol. Sci.* **21**, 3868 (2020).

46.   Meister, G. *et al.* Methylation of Sm proteins by a complex containing PRMT5 and the
putative U snRNP assembly factor pICln. *Curr. Biol. CB* **11**, 1990–1994 (2001).

47.   Lin, C.-H., Huang, H.-M., Hsieh, M., Pollard, K. M. & Li, C. Arginine methylation of
recombinant murine fibrillarin by protein arginine methyltransferase. *J. Protein Chem.* **21**, 447–
453 (2002).

48.   Lomonte, P., Baklouti, F. & Binda, O. The Biochemistry of Survival Motor Neuron Protein
Is Paving the Way to Novel Therapies for Spinal Muscle Atrophy. *Biochemistry* **59**, 1391–1397
(2020).

49.   Karyka, E. *et al.* SMN-deficient cells exhibit increased ribosomal DNA damage. *Life Sci.*
Alliance **5**, e202101145 (2022).