# The transcriptional repressor Opi1 modulates the DNA Damage Response by downregulation of inositol pyrophosphates in *Saccharomyces cerevisiae*

Giovanna Marques Panessa[*]; Marina Rodrigues Pires[*]; Rodrigo Rodrigues Felix[*;]
Rafaella Jekabson[*]; Eduardo Tassoni Tsuchida[†]; Nadja Cristhina de Souza-Pinto[‡];
Fernanda Marques da Cunha[*] and José Renato Rosa Cussiol[*]

[*] Departamento de Bioquímica, Escola Paulista de Medicina, Universidade Federal de São Paulo, São Paulo, Brasil

[†] Departamento de Genética e Biologia Evolutiva, Instituto de Biociências, Universidade de São Paulo, São Paulo, Brasil

[‡] Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, São Paulo, Brasil

**Running title:** Inositol metabolism crosstalk with DDR

**Keywords:** DNA Damage Response, Replication stress, Opi1, inositol pyrophosphates

**Corresponding author:**

José Renato R. Cussiol

Rua Três de Maio, 100

Instituto de Farmacologia e Biologia Molecular

Escola Paulista de Medicina – Universidade Federal de São Paulo

São Paulo, SP 04044-020

t. +55 11 5576-4444 Voip 1085

e-mail: cussiol@unifesp.br

2                                              **Abstract**

Inositol is a six-carbon sugar that functions as a precursor for signaling molecules such

as phosphoinositides and inositol polyphosphates, which are involved in the regulation of

important biological processes such as energy metabolism, environmental stress

response, phosphate signaling, among others. Given its role in a myriad of signaling

pathways, regulation of inositol synthesis is essential for cellular homeostasis. In budding

yeast, transcription of genes involved in inositol metabolism is regulated by the

transcriptional repressor Opi1, which repress transcription of genes containing cis-acting

inositol-sensitive upstream activation sequences (UAS$_{INO}$).  Upon genotoxic stress, cells

activate the DNA Damage Response (DDR) to coordinate DNA repair and cell cycle

progression. It has been proposed that inositol containing molecules might act as

modulators of the DDR, but evidences are still scarce. Herein, we report that *opi1Δ* cells

fail to downregulate the inositol phosphate pathway leading to sensitivity to genotoxins

and replication defects. Moreover, cells lacking Opi1 show decreased gamma-H2A levels

which might indicate that Opi1 contributes to the activation of the DDR kinases Mec1/Tel1

(ATR/ATM in mammals). Importantly, we show that deletion of the inositol pyrophosphate

kinase Kcs1 (IP6K1/2/3 in mammals), which leads to inhibition of inositol pyrophosphate

synthesis, rescues the MMS sensitivity and replication defects of *opi1Δ* cells. Further,

overexpression of Kcs1 recapitulates the MMS sensitivity of cells lacking Opi1. Therefore,

we propose that cells must downregulate inositol pyrophosphate synthesis during

replication stress in order to trigger an effective DNA Damage Response.

**Introduction**

The DNA molecule is constantly exposed to endogenous and exogenous

genotoxins, leaving it subjected to a multitude of lesions that, if left unchecked, can lead

to genomic instability, a hallmark of cancer and other diseases (Jackson and Bartek

2009). Under such circumstances, a comprehensive DNA Damage Response (DDR) is

activated to tackle the consequences of DNA lesion accumulation, such as through DNA

repair, telomere maintenance, chromatin remodeling, inhibition of DNA synthesis and cell

cycle progression, transcription reprogramming, among others (Hanawalt 2015; Lanz *et al.*

2019; Cussiol *et al.* 2020). Although DDR has been extensively studied at the molecular

level, recent studies have proposed that the DDR is connected with metabolism of

biomolecules such as carbohydrates and lipids (Simpson-Lavy *et al.* 2015; Yi *et al.* 2017;

Ferrari *et al.* 2017). Interestingly, it was proposed many years ago that might exist a

crosstalk between the DDR and inositol metabolism in eukaryotes (Zewail *et al.* 2003). In

line with that, many proteins from the inositol pathway are phosphorylated in response to

DNA damage (Zhou *et al.* 2016; Lanz *et al.* 2021). Inositol metabolites such as inositol

polyphosphates (IPs) and inositol pyrophosphates (PP-IPs) have been implicated in cell

cycle regulation and DNA damage repair (Banfic *et al.* 2013, 2016; Jadav *et al.* 2013), but the

molecular basis for these effects is unknown. Moreover, it was recently proposed that

phosphatidylinositol phosphate lipids (PIPs) are enriched in the nucleus after DNA

damage and are important mediators of ATR signaling in mammalian cells (Wang *et al.*

2017). However, little is known about how DNA damage can regulate inositol metabolism

and vice-versa.

Inositol is synthesized from glucose and is an essential precursor of a broad range

of molecules that play important roles in cell signaling (Wilson *et al.* 2013; Henry *et al.* 2014;

Shah *et al.* 2017). It has been proposed that the concentration of intracellular inositol is

critical for higher eukaryotes, as perturbation of inositol levels is correlated with human

diseases such as bipolar disorder, Alzheimer's disease, diabetes, and cancer (Winegrad

1987; Fruen and Lester 1991; Shi *et al.* 2006). In budding yeast, inositol biosynthesis is fine-

tuned by the transcriptional repressor Opi1. In absence of inositol, Opi1 is localized to the

perinuclear ER (endoplasmic reticulum) membrane establishing interactions with the

integral membrane protein Scs2 and the phospholipid precursor phosphatidic acid (PA)

(Craven and Petes 2001; Brickner and Walter 2004; Gaspar *et al.* 2017; Hofbauer *et al.* 2018). As

a result, the heterodimeric transcriptional activator Ino2-Ino4 binds to a cis-acting inositol-

sensitive upstream activation sequences (UAS$_{INO}$) and upregulates expression of several

genes related to phospholipid metabolism including the inositol-3-phosphate synthase

(*INO1*), which promotes inositol *de novo* synthesis. Once intracellular inositol

concentration increases, PA is redirected to the synthesis of phosphatidylinositol (PI)

leading to its exhaustion. Subsequently, Opi1 migrates to the nucleus where it binds to

the heterodimeric transcriptional activator Ino2-Ino4 and promotes transcriptional

repression through the interaction with the proteins Sin3 and Cyc8, creating a scaffold for

the recruitment of histone deacetylase (HDAC) complexes  (Wagner *et al.* 2001; Jäschke *et*

*al.* 2011; Kliewe *et al.* 2017b) (Figure 1A). Deletion of Opi1 leads to constitutive activation

of a large number of genes, most of which are regulated by Ino2-Ino4 (Santiago and

Mamoun 2003; Jesch *et al.* 2005; Hoppen *et al.* 2005). Many of these genes are involved in

phospholipid biosynthesis, although UAS$_{INO}$ motifs are found in several genes related to

other metabolic processes, which might suggest that Ino2-Ino4 controls the expression

of distinct biological processes (Wimalarathna *et al.* 2011). Importantly, deletion of Opi1

leads to constitutive expression of *INO1* with overproduction of inositol even when cells

are supplemented with inositol (Graves and Henry 2000). Interestingly,

phosphoproteomic analysis in budding yeast showed that the transcriptional repressor

Opi1 is phosphorylated in a Mec1/Tel1-dependent manner after MMS exposure (Balint *et*

*al.* 2015; BastosdeOliveira *et al.* 2015; Lanz *et al.* 2021), which might suggest that Opi1 is

involved in the DDR.

Herein, we propose that the transcriptional repressor Opi1 mediates, at least in part, the

inositol pathway-DDR crosstalk. We find that cells lacking Opi1 under genotoxic stress

show enhanced sensitivity, increased expression of *INO1*, cell cycle defects with delay in

S-phase progression and decreased histone H2A phosphorylation. Interestingly, deletion

of the inositol pyrophosphate kinase Kcs1 in an *opi1Δ* strain rescues MMS sensitivity and

cell cycle defects. Moreover, overexpression of Kcs1 in wild-type cells phenocopies the

genotoxin sensitivity of an *opi1Δ* strain. Together, these results indicate that modulation

of PP-IPs synthesis during replication stress is part of the DDR and a key factor for cell

survival in budding yeast.

## Material and Methods

**Yeast strains and plasmids.** Strains generated in this study are isogenic from BY4741
and S288C (where indicated). All yeast strains and plasmids used in this study are
described in Table S1 and Table S2, respectively. All yeast strains were generated using
yeast genetic tools for gene disruption and epitope tagging (Longtine *et al.* 1998). All yeast
transformations were performed using the lithium acetate method (Gietz *et al.* 1992; Gietz
and Woods 2006). Knockout strains were confirmed by genotyping using primers and
epitope tagged strains were confirmed by western blotting. Restriction digestion cloning
was used to generate the plasmids for this study. Briefly, a PCR product containing 250bp
of the *OPI1* promoter + *OPI1* ORF with 3xHA sequence fused to the C-terminus + *ADH1*
terminator sequence was cloned into the pRS416 vector using XhoI and SacII (New
England Biolabs). For Ino1 and Kcs1 overexpression, the ORF coding for *INO1* and *KCS1*
were PCR amplified from yeast genomic DNA using primers and cloned into pYES2-NTC
using EcoRI and XhoI (New England Biolabs) under the control of the GAL1 promoter.

**Growth conditions.** Cells were grown in YPD (1% yeast extract, 2% peptone, 2%
dextrose) and plasmid-bearing cells were grown in synthetic complete media lacking
uracil (SC-URA) (0.17% YNB w/o aminoacids - Difco), 0.5% ammonium sulfate (MP
Biomedicals) 0.07% CSM-URA dropout mix (Sunrise Science), 2% dextrose (Sigma). For
experiments performed in the absence of inositol, SC-INO medium was prepared using
an YNB without inositol (MP Biomedical). For expression of *INO1* under the control of the

GAL1 promoter, cells were grown overnight in SC-URA medium with 2% dextrose and

were plated in SC-URA plates in the presence of 2% dextrose (Sigma) or 2% galactose

(Difco).

**Serial dilution assays.** Overnight cultures were inoculated in the appropriate medium to

an optical density ($OD_{600nm}$) 0.1 and cells were grown until log phase and normalized to

an $OD_{600nm}$ of 1.0. Fourfold serial dilutions were spotted on yeast plates and grown for 1–

3 days at 30°C in the presence or absence of drugs. Plates were manually inspected and

digitalized in an Uvitec Alliance 4.7.

**Real Time Quantitative PCR (RT-qPCR) Analysis.** Cells were grown overnight in YPD

and diluted next day to an $OD_{600nm}$ of 0.1 in synthetic complete medium either

supplemented or lacking inositol (SC+INO or SC-INO). Cells were grown for at least two

cell cycle divisions and immediately harvested at 4°C, washed with sterile ultra-pure $H_2O$,

and kept at -80°C. Alternatively, log phase cells were treated with MMS (Sigma-Aldrich)

0.1% for 1 hour. Total RNA from $5x10^7$ cells was extracted using RNeasy mini kit

(Qiagen). One $\mu$g total RNA was used for cDNA synthesis using Quantinova Reverse

Transcription kit (Qiagen). Real time PCR was performed in an ABI Prism 7500 (Applied

Biosystems) using Quantinova Probe PCR kit (Qiagen) in the presence of Taqman®

probes for *INO1* (Sc04136910_s1) and *ACT1* (Sc04120488_s1) (Thermo Fisher

Scientific). The *ACT1* gene served as an internal standard for normalization as it was

previously shown that there is no change in mRNA levels either after inositol depletion

(Ye *et al.* 2013; Gaspar *et al.* 2017) or MMS treatment (Gasch *et al.* 2001). All reactions were

made with three biological replicates. Relative values of mRNA are described as fold

change relative to a control condition indicated in the experiment.

**Western blot.** Whole-cell lysates were denatured with 3x SDS sample buffer (188mM

Tris-HCL pH 6.8; 3% SDS; 30% glycerol; 0.01% bromophenol blue, 1x Bond-Breaker

TCEP solution – Thermo Fischer Scientific) and resolved on SDS-PAGE gels. Proteins

were transferred onto PVDF membranes (Amersham Hybond P – GE Healthcare).

Phosphorylated histone H2A (gamma-H2A) and epitope tagged proteins were probed

using specific antibodies: anti-gamma H2A (Ab17353-Abcam, 1:2500 dilution in TBST

and 5% non-fat dry milk), anti-HA (12CA5; Roche, 1:10000 in TBST and 2% non-fat dry

milk). Anti-PGK1 was used as a housekeeping control (22C5D8, Abcam, 1:15000 dilution

in TBST and 2% non-fat dry milk). After primary antibody, PVDF membranes were

incubated in the presence of ECL HRP-linked secondary antibody (NA931-GE, 1:10000

dilution in TBST and 2% non-fat dry milk). Blots were developed using the Amershan ECL

Prime detection reagent and imaged in a Uvitec Alliance 4.7

**Cell cycle synchronization.** For arrest of cells in G1, yeast cells were grown in YPD

medium at 30°C until log phase followed by $\alpha$-factor (Zymo Research) treatment (30 ng/ml

for *bar1Δ* background strains) for 2 h. To release from G1 arrest, cells were centrifuged

and resuspended in fresh medium in the presence of MMS 0.033% (where indicated).

For recovery of cells from MMS treatment, cells were treated with MMS 0.033% for 2 h,

harvested and resuspended in fresh medium (YPD).

**Flow cytometry.** Logarithmic growing yeast cells were collected, harvested, fixed in 1 ml of 70% ethanol (Sigma-Aldrich), and incubated overnight at 4°C. Cells were then centrifuged, and residual ethanol was dried in a speed-vac. After that, samples were solubilized in sodium citrate buffer (LabSynth) (50 mM, pH 7.2) and sonicated (three cycles of 3 sec, amplitude 30%) to release cell clumps. Samples were then incubated with 100 µg of RNAseA (Invitrogen) for 1 h at 37°C followed by incubation with 500 µg of Proteinase K (Invitrogen) for 1 h at 42 °C. Then, 1 µL of SYTOX Green (Thermo Fisher Scientific) was added to the samples and incubated for 2 h at 4°C protected from the light. Data were acquired using a BD Accuri C6 Flow Cytometer.

**Respiratory capacity assay.** Oxygen consumption rates were measured using high-resolution O2K Fluorespirometer (Oroboros, Innsbruck, Austria). For each assay, $10^6$ cells were added to the Oroboros chamber containing a final volume of 2.0 ml of YPEG medium at 30°C under agitation. Briefly, exponentially growing cells in YPD were harvested, resuspended in respiratory medium (YPEG; 2% glycerol and 1% ethanol), and incubated for at least 3 hours at 30°C. Finally, cells were transferred to O2K chambers for measurement of $O_2$ consumption rates. After basal respiration assessment, 1µM of the mitochondrial uncoupler FCCP (Sigma-Aldrich) was added to measure maximal respiration. At the end, 2µM of antimycin A (Sigma-Aldrich) was added to inhibit mitochondrial respiration. Non-mitochondrial oxygen consumption rates were subtracted from all measurements. Optimal concentrations of FCCP and antimycin A were previously determined by titration.

**Microscopy analysis.** Images were acquired in a Zeiss LSM 780 microscope. Laser lines

at 561nm were used for the detection of Opi1-GFP tagged protein in yeast cells. For yeast

Opi1-GFP analysis, cells were grown in synthetic complete media (SC +INO) until the log

phase (OD = 0.3) and shifted to SC -INO for 1 hour. Finally, MMS (0.1%) was added to

the cells and incubated for 30 min at 30°C. At each condition, cells were washed in sterile

water and resuspended in fresh synthetic complete media lacking tryptophan (SC-TRP).

Live yeast cultures were mounted on an agarose slide pad (1.2% agarose in SC-TRP

media) and >100 cells were scored for each replicate. Cells were manually counted for

Opi1-GFP localization (nuclear, transition from the nucleus to the ER and ER localization).

Statistics were calculated based on the mean and SEM derived from two independent

replicates.

219 **Data availability**
220
221

222 Strains and plasmids are available upon request. Tables S1 and S2 contain the list of

223 yeast strains and plasmids used in this study, respectively. Figure S1 contains

224 supplemental data in support of Figure 3A and B. Figure S2 contains supplemental data

225 in support of Figure 4. Figure S3 and S4 contains supplemental data in support of Figure

226 5.

                                                              **Results**

**Opi1 is important for cell survival during genotoxic stress**

To determine whether cells lacking Opi1 are more sensitive to DNA damage, we

performed sensitivity assays in the presence of genotoxins. We found that *opi1Δ* cells

showed sensitivity to genotoxins such as MMS and bleomycin (Figure 1B-C), thus

indicating that Opi1 is important for the response to DNA damage. To confirm that this

sensitivity to MMS was specific to *OPI1* deletion, we constructed a pRS416 plasmid

expressing Opi1-HA from its endogenous promoter (Figure 1D) and show that Opi1-HA

expression rescues the MMS sensitivity in *opi1Δ* (Figure 1E). Furthermore, deletion of

genes that counteract Opi1 repressing functions, such as *INO1* and *SCS2*, does not lead

to MMS sensitivity (Figure 1B). These results suggest that downregulation of inositol

levels through Opi1 are important to trigger an effective response to genotoxic stress.

**Cells lacking Opi1 show replication defects and decreased gamma-H2A levels**

Since *opi1Δ* mutants are sensitive to genotoxic stress induced by MMS (Figure

1B), we sought to investigate if these cells have defects in the DDR. First, we treated

exponentially growing cells with MMS 0.033% for 2 hours to induce an intra S-phase

arrest followed by release in fresh medium. Remarkably, *opi1Δ* cells arrest in early S-

phase rather than mid S-phase and showed impaired S-phase progression during

recovery from MMS treatment when compared to a wild-type strain (Figure 2A). Because

cells lacking Opi1 show replication defects in the presence of MMS, we monitored

gamma-H2A levels by immunoblot using a phospho-specific antibody. Upon replication

stress, the sensor kinases Mec1 and Tel1 catalyze the phosphorylation of long tracks of

histone H2A at serine residue 129 (gamma-H2A) surrounding the DNA lesion creating a

platform for the recruitment of multiple DNA damage response proteins (Rogakou *et al.*

1998; Downs *et al.* 2000), thus serving as a readout for DDR activation (Fernandez-Capetillo

*et al.* 2004). Interestingly, gamma-H2A levels were significantly lower in *opi1Δ* relatively to

wild-type after 2 hours of MMS treatment, but no significant difference was observed on

kinetics of gamma-H2A disappearance during recovery (Figure 2B). Therefore, this result

suggests that DNA damage signaling is downregulated in *opi1Δ*. To further address if

DDR is downregulated in *opi1Δ* cells, we monitored the progressive phosphorylation of

histone H2A upon MMS treatment. We arrested cells in G1 with $\alpha$-factor and released in

fresh YPD medium in the presence of MMS. Remarkably, cells lacking Opi1 have a

delayed histone H2A phosphorylation and decreased intensity of gamma-H2A signal in

latter time points (Figure 2C). Additionally, cells lacking Opi1 show slower S-phase

progression in the presence of MMS (Figure 2D). Together, these results indicate that

Opi1 is important for cells to cope with DNA damage during replication.

**Opi1 migrates to the nucleus upon genotoxic stress induced by MMS**

Knowing that transcriptional repression function of Opi1 is dependent on its cellular

localization and because its absence causes MMS sensitivity (Figure 1B), we investigated

Opi1 localization after genotoxic stress. Using a strain expressing Opi1-GFP, we

recapitulated Opi1 cellular localization during inositol metabolism (Figure 3A panels i and

ii). MMS treatment of cells supplemented with 100$\mu$M of inositol (+INO) did not cause any

alteration in Opi1-GFP intensity or localization (data not shown). Surprisingly, MMS

treatment of cells kept without inositol supplementation (-INO) induced a rapid

translocation of Opi1-GFP to the nucleus, suggesting that the nuclear localization of Opi1

is important to cope with genotoxic stress (Figure 3A-B and Supplementary Figure 1).

Importantly, yeast cells treated with MMS in SC-INO are viable (Figure 3C), which implies

that Opi1 does not fully repress *INO1* in these conditions, otherwise cells would die.

Subsequently, to understand if Opi1 migration to the nucleus in the presence of MMS

causes gene repression, we looked at *INO1* expression during MMS treatment by RT-

qPCR analysis. While MMS treatment does not change *INO1* expression in wild-type cells

supplemented with inositol (+INO), there is a substantial decrease in *INO1* expression in

the absence of inositol (-INO). Strikingly, *INO1* expression significantly increases in an

*opi1Δ* strain upon MMS treatment both in +INO and -INO conditions (Figure 3D). These

results show that under genotoxic stress Opi1 migrates to the nucleus to repress *INO1*

expression in order to downregulate inositol synthesis. Because Opi1 represses genes

containing UAS$_{INO}$, it is possible that in its absence, several other genes are also

upregulated upon MMS treatment.

**The constitutive activation of Ino2-Ino4 responsive genes is the cause of MMS**

**sensitivity in cells lacking Opi1**

Since Opi1 repress the transcriptional activator Ino2-Ino4 in the presence of

inositol, we investigated if *INO2* and *INO4* deletions could rescue the MMS sensitivity of

cells lacking Opi1. Remarkably, *opi1Δino2Δ* and *opi1Δino4Δ* strains are rescued for the

MMS sensitivity, which is an indication that constitutive expression of Ino2-Ino4

responsive genes is causing the MMS sensitivity (Figure 4A and 4B). Of note, it has been

shown that the lack of Opi1 compromises mitochondrial function due to respiratory

defects related to a decrease in mitochondrial membrane cardiolipin content (Luévano-

Martínez *et al.* 2013). Interestingly, deletion of *INO2* completely restores the respiratory

capacity of cells lacking Opi1 (Figure 4C). To elucidate if the rescue of MMS sensitivity is

related to a recovery of mitochondrial function we sought to investigate if *INO2* deletion

would also rescue the MMS sensitivity of *opi1Δ* in respiratory conditions (rich medium

containing glycerol 3% as a carbon source). However, while *opi1Δino2Δ* fully rescues

MMS sensitivity in fermentative conditions (Figure 4A), there is only a partial rescue in

YPG (Figure 4D). It is known that ROS production during MMS treatment is exacerbated

in the presence of high glucose (2%) with inhibition of glycolytic enzymes leading to

growth arrest (Kitanovic *et al.* 2009; Yazgan and Krebs 2012). ROS leakage from

mitochondria can be ameliorated by reducing glucose concentration in the medium.

Therefore, we repeated the MMS sensitivity assay in the presence of 0.5% glucose with

the expectation that if the lack of Opi1 affects the oxidative stress response we would see

a rescue of MMS sensitivity. However, there was no change in MMS sensitivity

(Supplementary Figure 2A). Furthermore, treatment with $H_2O_2$ does not sensitize *opi1Δ*

cells (Supplementary Figure 2B). Therefore, we conclude that constitutive activation of

genes regulated by Ino2-Ino4 affects both the response to genotoxic stress and

mitochondrial function, but probably by independent mechanisms.

**Opi1 regulates inositol pyrophosphates synthesis during the DDR**

It is well established that the expression of *INO1* is positively regulated by Ino2-

Ino4 and massively increases in the absence of Opi1 (Figure 3D) (Graves and Henry 2000;

Santiago and Mamoun 2003; Jesch *et al.* 2005). Consequently, we deleted *INO1* in an *opi1Δ*

background with the expectation that it would phenocopy the MMS resistance of an

*opi1Δino2Δ*. Surprisingly, an *opi1Δino1Δ* double mutant showed growth defects and

increased MMS sensitivity (Figure 5A). This negative genetic interaction between Opi1

and Ino1 suggests that upregulation of one or more genes repressed by Opi1 in the

absence of Ino1 negatively impacts the cell. To test if constitutive expression of *INO1* is

causing MMS sensitivity, we overexpressed *INO1* by cloning it under the control of the

strong inducible *GAL1* promoter with the expectation that it would phenocopy the MMS

sensitivity of an *opi1Δ* strain. Expression of *INO1* from the *GAL1* promoter rescues the

inositol auxotrophy of cells lacking *INO1* (Supplementary Figure 3A), but cells

overexpressing *INO1* do not show sensitivity to MMS, which is an indication that

increased levels of inositol do not affect the DDR (Supplementary Figure 3B).

Besides Ino1, Ino2-Ino4 upregulate the expression of several other genes

counteracting Opi1 suggesting that constitutive expression of another Ino2-Ino4 target in

*opi1Δ* cells is causing MMS sensitivity. Therefore, we conducted a screening of a small

set of genes which expression is dependent on Ino2-Ino4 and that are upregulated in

*opi1Δ* cells (Santiago and Mamoun 2003). Among them, we selected genes involved in

the phosphatidylinositol and phosphatidylcholine synthesis (Itr1 - inositol transporter,

Opi3 - phospholipid methyltransferase and Cki1 - choline kinase), polyphosphate

metabolism (Pho84 - high-affinity inorganic phosphate transporter) and inositol

pyrophosphate synthesis (Kcs1 - inositol pyrophosphate kinase). Similar to what was

observed for Ino1, deletion of the inositol transporter Itr1, phospholipid methyltransferase

Opi3, choline kinase Cki1 and inorganic phosphate transporter Pho84 also showed a

negative genetic interaction with Opi1 (Supplementary Figure 4). On the other hand,

deletion of Kcs1 in an *opi1Δ* strain significantly rescued the MMS sensitivity and the S-

phase progression after replication stress induced by MMS (Figure 5B and C). In budding

yeast, Kcs1 catalyzes the synthesis of inositol pyrophosphates. Inositol polyphosphates

$IP_5$ and $IP_6$ generated by Ipk2 and Ipk1 respectively are substrates of Kcs1 that catalyzes

the pyrophosphorylation of the C5 $\alpha$-phosphate to 5-PP-$IP_4$ and 5-PP-$IP_5$ respectively

(Saiardi *et al.* 1999). These results, suggest that overproduction of PP-IPs by Kcs1 is most

probably the cause of genotoxin sensitivity (Figure 5B and C). Interestingly, cells lacking

the Siw14 phosphatase that counteracts Kcs1-dependent PP-IPs synthesis have a mild

sensitivity to MMS (Figure 5D). It has been shown that *siw14Δ* mutants have increased

levels of 5-PP-$IP_5$ (Steidle *et al.* 2016). Consistent with that, wild-type cells overexpressing

Kcs1 phenocopy an *opi1Δ* strain with cell growth delay and sensitivity to MMS (Figure

5E). Taken together, our results suggest that increased PP-IPs synthesis due to the lack

of the transcriptional repressor Opi1 leads to impaired capacity to cope with replication

stress induced by MMS.

398                                          **Discussion**

In this manuscript, we demonstrated that the transcriptional repressor Opi1

modulates the expression of Ino2-Ino4 during genotoxic stress. Importantly, we propose

that Opi1 controls PP-IPs levels through downregulation of genes from inositol phosphate

pathway. Failure to downregulate PP-IPs in an *opi1Δ* strain causes downregulation of the

DDR leading to sensitivity to genotoxic agents due to increased replication stress,

reinforcing the notion that inositol metabolism affects the DDR.

**PP-IPs overproduction affects the response to replication stress**

The fact that Opi1 regulates the expression of several genes involved in different

metabolic pathways makes the task of identifying the cause of genotoxin sensitivity very

challenging. However, we found that deletion of the transcription factor Ino2-Ino4 and the

PP-IPs kinase Kcs1 restores viability of cells lacking Opi1. In budding yeast, PP-IPs are

synthetized from the inositol polyphosphates $IP_5$ and $IP_6$ in a reaction catalyzed by the

PP-IPs kinases Kcs1 and Vip1 (Saiardi *et al.* 1999; Mulugu *et al.* 2007; Lee *et al.* 2008). Kcs1

catalyzes the conversion of $IP_5$ and $IP_6$ to different isomers of PP-IPs such as 5-PP-$IP_4$

and 5-PP-$IP_5$ (Saiardi *et al.* 1999). Kcs1 and its mammalian orthologs IP6Ks have been

implicated in several biological processes such as cell cycle progression, DNA repair,

telomeric length, nutrient signaling, environmental stress response, polyphosphate

metabolism among others (Saiardi *et al.* 2005; Auesukaree *et al.* 2005; Szijgyarto *et al.* 2011;

Banfic *et al.* 2013, 2016; Worley *et al.* 2013; Wilson *et al.* 2013). It has been previously shown

that *KCS1* possesses UAS$_{INO}$ sequences in its promoter region and that its expression is

upregulated when inositol concentration increases (Wimalarathna *et al.* 2011).

Corroborating the notion that PP-IPs synthesis is upregulated in cells lacking Opi1, Kcs1

protein levels were shown to increase in *opi1Δ* cells (Ye *et al.* 2013). Additionally, the

inositol polyphosphate multikinase Ipk2 that catalyzes the synthesis of $IP_4$ and $IP_5$ also

contains $UAS_{INO}$ sequences in its promoter region, and its expression is dependent on

Ino2-Ino4 (Wimalarathna *et al.* 2011). Therefore, an increase in PP-IPs levels might result

whether from increased Kcs1 or Ipk2 expression. It is interesting to note that while Ino1

overexpression does not sensitize cells to MMS, that of Kcs1 does (Supplementary Figure

3B and Figure 5E respectively), suggesting that Ino1 overexpression does not lead to an

increase in PP-IPs synthesis, which might indicate that there is a rate-limiting step

reaction downstream of inositol-3-phophate synthesis restraining PP-IPs concentration in

the cell.

Finally, is interesting to note that Opi1 shows negative genetic interactions with

genes involved in inositol synthesis and transport (Ino1 and Itr1); phosphatidylcholine

synthesis (Opi3 and Cki1) and phosphate metabolism (Pho84) (Figure 5A and Supp.

Figure 4). For instance, deletion of the inorganic phosphate transporter Pho84 leads to a

synthetic lethal/sick phenotype (Supp. Figure 4). Because inositol and phosphate

metabolism are intimately related (Auesukaree *et al.* 2005; Mulugu *et al.* 2007; Lee *et al.* 2008;

Kliewe *et al.* 2017a), this observation might give new insights for further exploration of

these pathways.

**Genotoxin sensitivity in *opi1Δ* is directly associated with DNA damage**

Deletion of the transcriptional repressor Opi1 can induce pleiotropic effects in the

cell. Therefore, we conducted a careful analysis to rule out the possibility that the

genotoxin sensitivity of *opi1Δ* cells was due to a deregulation of other metabolic pathways

not directly associated with the DDR. It was previously shown that cells lacking Opi1

possess short telomeres (Askree *et al.* 2004). In contrast, cells lacking Kcs1 have longer

telomeres (York *et al.* 2005). This finding could indicate that deletion of Kcs1 rescues MMS

sensitivity due to restoration of telomere length. However, this is probably not the case as

it was shown that telomere length does not affect cellular fitness nor yeast sensitivity to

DNA damage (Harari *et al.* 2017). Consistent with this, PP-IPs were also shown to

modulate cell cycle independently of telomere length (Banfic *et al.* 2016). MMS can also

damage mitochondrial DNA leading to direct inhibition of the respiratory chain with

increased ROS leakage, oxidative inactivation of glycolytic enzymes and cell cycle arrest

(Kitanovic *et al.* 2009). Moreover, deletion of Opi1 decreases mitochondrial cardiolipin

content and it was proposed that overproduction of inositol might affect cardiolipin

production probably through inhibition by phosphorylation of the Phosphatidylglycerol

phosphate synthase, Pgs1 (Luévano-Martínez *et al.* 2013). We found, however, that

genotoxin sensitivity of cells lacking Opi1 is probably not due to a defect in mitochondria

homeostasis because: i) diauxic shift induced by incubation with lower glucose levels

does not ameliorate MMS sensitivity (Supp. Figure 2A); ii) *opi1Δ* does not show sensitivity

to $H_2O_2$ (Supp. Figure 2B); iii) deletion of *INO2* rescues the respiratory defects of cells

lacking Opi1, but does not rescue the MMS sensitivity in the presence of a respiratory

carbon source (Figure 4D). Despite that, we cannot entirely rule out the possibility that

the genotoxin sensitivity of *opi1Δ* is dependent on mitochondria because it is known that

PP-IPs can also modulate mitochondrial respiration and ROS signaling (Onnebo and

Saiardi 2009; Szijgyarto *et al.* 2011; Worley *et al.* 2013). Furthermore, MMS treatment can

also induce ER stress leading to activation of the UPR (*Unfolded Protein* Response) and

it was shown that Ino2-Ino4 also regulate expression of genes involved in the UPR

(Jelinsky and Samson 1999; Jesch *et al.* 2005; Thibault *et al.* 2012). Nevertheless, it was shown

that the absence of Opi1 causes an expansion of the ER membrane, alleviating the ER

stress induced by DTT or tunicamycin in an UPR independent manner (Schuck *et al.* 2009).

**A working model for modulation of PP-IPs synthesis during replication stress**

**induced by MMS**

How does inositol pyrophosphates deregulation affect DNA replication leading to

genotoxin sensitivity? It is established that PP-IPs modulate protein function either

through non-enzymatic pyrophosphorylation of proteins, or allosteric regulation through

its interaction with protein domains (Wilson *et al.* 2013). However, up until now, only a few

targets have been characterized. Therefore, we can speculate that proteins related to the

DDR might being modified by PP-IPs. Of note, it has been shown that Kcs1 is important

for S-phase progression, as PP-IPs levels increase after pheromone induced arrest by

an unknown mechanism (Banfic *et al.* 2013). In fact, it was previously demonstrated that

Kcs1 is required for DNA hyperrecombination phenotype in yeast cells with defects in

protein kinase C1 (Pkc1) (Luo *et al.* 2002). Moreover, it was shown in mammalian cells

that the orthologous of Kcs1 (IP6K1/2/3) are important for homologous recombination

repair (Jadav *et al.* 2013). Curiously, deletion of Kcs1 also confers sensitivity to MMS,

suggesting that depletion of PP-IPs, which synthesis is dependent on Kcs1, is also

important for the DDR (Figure 5B and (Zhou *et al.* 2016)). Since MMS-induced lesions can

be bypassed during replication by homologous recombination (HR) mechanisms (Cussiol

*et al.* 2017), it is tempting to speculate that PP-IPs targets proteins related to HR in order

to modulate the repair or bypass of MMS-induced replication blocks. Moreover, it has

been shown that inositol pyrophosphates can compete with PIP$_3$ (PtdIns(3,4,5)P3) for its

binding to pleckstrin-homology (PH) domains in proteins (Luo *et al.* 2003). In line with that,

a recent report has revealed that PIP$_2$ rapidly accumulates in the nucleus after DNA

damage in mammals and is important to mediate ATR recruitment to DNA lesions (Wang

*et al.* 2017). Therefore, accumulation of inositol pyrophosphate has the potential to

outcompete PIPs leading to downregulation of DNA damage signaling. This hypothesis

can explain the lower levels of gamma-H2A observed in *opi1Δ* mutants (Figure 2B and

C). Consequently, a decrease in Mec1/Tel1 signaling in cells lacking Opi1 might affect

the ability of the cells to cope with replication stress.

We propose a model in which concerted downregulation of inositol

pyrophosphates is important to cope with increased replication stress through a robust

DNA Damage Response (Figure 6). In the presence of genotoxic stress, Opi1

translocates to the nucleus to repress the inositol phosphate pathway, probably through

inhibition of *KCS1* and/or *IPK2* gene expression. In cells lacking Opi1, Ino2-Ino4

constitutively promotes transcription of genes from the inositol phosphate pathway,

leading to the accumulation of Kcs1-dependent inositol pyrophosphates. PP-IPs

accumulation might then modulate DDR targets either by allosteric interaction or

pyrophosphorylation. Because replication stress is a hallmark of cancer cells (Hills and

Diffley 2014; Gaillard *et al.* 2015) and the PP-IP pathway is highly conserved from yeast to

humans, the finding that PP-IP accumulation sensitizes cells to genotoxic stress might be

relevant for future exploration in higher eukaryotes, especially in cancer cell lines.

535
536                                    **Acknowledgements**

538         We thank Jacilene Barbosa and Natalia Bromberg for technical support. Dr.

539   Aparecida Sadae Tanaka and Dr. Maria Luiza Vilela Oliva for kindly allowing access to

540   their laboratory and equipment. Dr. Francisco Bastos de Oliveira and Dr. Gustavo

541   Monteiro Silva for comments and suggestions, Dr. Marcus Bustamante Smolka and Dr.

542   Luis Eduardo Soares Netto for donating plasmid and yeast strains.

543
544
545                                          **Funding**

This work is supported by a J.R.C grant from FAPESP (2018/05417-0); G.M.P.

grant from FAPESP (2021/04887-6); R.J grant from FAPESP (2019/25497-1); F.M.C

grant from FAPESP (2019/09732-0) and N.C.S-P grant from FAPESP (2017/04372-0)

Author contributions: J.R.C. wrote the paper, designed the study, performed

experiments and data analysis. G.M.P., M.R.P., R.J, E.T.T. performed experiments, data

analysis and helped revised the manuscript. F.M.C. designed and performed experiments

for measuring oxygen consumption and helped revised the manuscript. N.C.S-P provided

reagents, infrastructure and helped revised the manuscript.

556
557
558                                  **Conflicts of interest**

560   None declared

**References**

**References**

Askree S. H., T. Yehuda, S. Smolikov, R. Gurevich, J. Hawk, *et al.*, 2004 A genome-wide screen
for Saccharomyces cerevisiae deletion mutants that affect telomere length. Proc Natl Acad
Sci U S A 101: 8658–63. https://doi.org/10.1073/pnas.0401263101
Auesukaree C., H. Tochio, M. Shirakawa, Y. Kaneko, and S. Harashima, 2005 Plc1p, Arg82p,
and Kcs1p, Enzymes Involved in Inositol Pyrophosphate Synthesis, Are Essential for
Phosphate Regulation and Polyphosphate Accumulation in Saccharomyces cerevisiae. J
Biol Chem 280: 25127–25133. https://doi.org/10.1074/jbc.M414579200
Balint A., T. Kim, D. Gallo, J. R. Cussiol, F. M. Bastos De Oliveira, *et al.*, 2015 Assembly of
Slx4 signaling complexes behind DNA replication forks. EMBO Journal 34: 2182–2197.
https://doi.org/10.15252/embj.201591190
Banfic H., A. Bedalov, J. York, and D. Visnjic, 2013 Inositol Pyrophosphates Modulate S Phase
Progression after Pheromone-induced Arrest in Saccharomyces cerevisiae. J Biol Chem
288: 1717–1725. https://doi.org/10.1074/jbc.M112.412288
Banfic H., V. Crljen, L.-S. Vesna, V. Dembitz, H. Lalic, *et al.*, 2016 Inositol pyrophosphates
modulate cell cycle independently of alteration in telomere length. Adv Biological Regul
60: 22–28. https://doi.org/10.1016/j.jbior.2015.09.003
BastosdeOliveira F. M., D. Kim, J. R. Cussiol, J. Das, M. Jeong, *et al.*, 2015 Phosphoproteomics
Reveals Distinct Modes of Mec1/ATR Signaling during DNA Replication. Mol Cell 57.
https://doi.org/10.1016/j.molcel.2015.01.043
Brickner J. H., and P. Walter, 2004 Gene recruitment of the activated INO1 locus to the nuclear
membrane. PLoS Biol 2: e342. https://doi.org/10.1371/journal.pbio.0020342
Craven R. J., and T. D. Petes, 2001 The Saccharomyces cerevisiae suppressor of choline
sensitivity (SCS2) gene is a multicopy Suppressor of mec1 telomeric silencing defects.
Genetics 158: 145–54. https://doi.org/10.1093/genetics/158.1.145
Cussiol J. R., D. Dibitetto, A. Pellicioli, and M. B. Smolka, 2017 Slx4 scaffolding in
homologous recombination and checkpoint control: lessons from yeast. Chromosoma 126:
45–58. https://doi.org/10.1007/s00412-016-0600-y
Cussiol J. R. R., B. L. Soares, and F. M. B. de Oliveira, 2020 From yeast to humans:
Understanding the biology of dna damage response (DDR) kinases. Genet Mol Biol 43.
https://doi.org/10.1590/1678-4685-gmb-2019-0071
Downs J. A., N. F. Lowndes, and S. P. Jackson, 2000 A role for Saccharomyces cerevisiae
histone H2A in DNA repair. Nature 408: 1001–4. https://doi.org/10.1038/35050000
Fernandez-Capetillo O., A. Lee, M. Nussenzweig, and A. Nussenzweig, 2004 H2AX: the histone
guardian of the genome. DNA Repair (Amst) 3: 959–967.
https://doi.org/10.1016/j.dnarep.2004.03.024
Ferrari E., C. Bruhn, M. Peretti, C. Cassani, W. V. Carotenuto, *et al.*, 2017 PP2A Controls
Genome Integrity by Integrating Nutrient-Sensing and Metabolic Pathways with the DNA
Damage Response. Mol Cell 67: 266-281.e4. https://doi.org/10.1016/j.molcel.2017.05.027
Fruen B. R., and B. R. Lester, 1991 Inositol and inositol 1,4,5-trisphosphate content of Down
syndrome fibroblasts exhibiting enhanced inositol uptake. FEBS Lett 295: 43–47.
https://doi.org/10.1016/0014-5793(91)81380-Q

Gaillard H., T. García-Muse, and A. Aguilera, 2015 Replication stress and cancer. Nat Rev
605           Cancer 15: 276–289. https://doi.org/10.1038/nrc3916

606    Gasch A. P., M. Huang, S. Metzner, D. Botstein, S. J. Elledge, *et al.*, 2001 Genomic expression
607           responses to DNA-damaging agents and the regulatory role of the yeast ATR homolog
608           Mec1p. Mol Biol Cell 12: 2987–3003. https://doi.org/10.1091/mbc.12.10.2987

609    Gaspar M. L., Y.-F. Chang, S. A. Jesch, M. Aregullin, and S. A. Henry, 2017 Interaction
610           between repressor Opi1p and ER membrane protein Scs2p facilitates transit of phosphatidic
611           acid from the ER to mitochondria and is essential for INO1 gene expression in the presence
612           of choline. J Biol Chem 292: 18713–18728. https://doi.org/10.1074/jbc.M117.809970

613    Gietz D., S. A. Jean, R. A. Woods, and R. H. Schiestl, 1992 Improved method for high efficiency
614           transformation of intact yeast cells. Nucleic Acids Res 20: 1425.

615    Gietz R. D., and R. A. Woods, 2006 Yeast transformation by the LiAc/SS Carrier DNA/PEG
616           method. Methods Mol Biol 313: 107–120. https://doi.org/10.1385/1-59259-958-3:107

617    Graves J. A., and S. A. Henry, 2000 Regulation of the yeast INO1 gene: The products of the
618           INO2, INO4 and OPI1 regulatory genes are not required for repression in response to
619           inositol. Genetics 154: 1485–1495.

620    Hanawalt P. C., 2015 Historical perspective on the DNA damage response. DNA Repair (Amst)
621           36: 2–7. https://doi.org/10.1016/j.dnarep.2015.10.001

622    Harari Y., S. Zadok-Laviel, and M. Kupiec, 2017 Long telomeres do not affect cellular fitness in
623           yeast. mBio 8: 1–9. https://doi.org/10.1128/mBio.01314-17

624    Henry S. A., M. L. Gaspar, and S. A. Jesch, 2014 The response to inositol: Regulation of
625           glycerolipid metabolism and stress response signaling in yeast. Chem Phys Lipids 180: 23–
626           43. https://doi.org/10.1016/j.chemphyslip.2013.12.013

627    Hills S. A., and J. F. X. Diffley, 2014 DNA replication and oncogene-induced replicative stress.
628           Curr Biol 24: R435-44. https://doi.org/10.1016/j.cub.2014.04.012

629    Hofbauer H. F., M. Gecht, S. C. Fischer, A. Seybert, A. S. Frangakis, *et al.*, 2018 The molecular
630           recognition of phosphatidic acid by an amphipathic helix in Opi1. J Cell Biol 217: 3109–
631           3126. https://doi.org/10.1083/jcb.201802027

632    Hoppen J., A. Repenning, A. Albrecht, S. Geburtig, and H.-J. Schüller, 2005 Comparative
633           analysis of promoter regions containing binding sites of the heterodimeric transcription
634           factor Ino2/Ino4 involved in yeast phospholipid biosynthesis. Yeast 22: 601–613.
635           https://doi.org/10.1002/yea.1209

636    Jackson S. P., and J. Bartek, 2009 The DNA-damage response in human biology and disease.
637           Nature 461: 1071–1078.

638    Jadav R. S., M. V. L. Chanduri, S. Sengupta, and R. Bhandari, 2013 Inositol pyrophosphate
639           synthesis by inositol hexakisphosphate kinase 1 is required for homologous recombination
640           repair. J Biol Chem 288: 3312–21. https://doi.org/10.1074/jbc.M112.396556

641    Jäschke Y., J. Schwarz, D. Clausnitzer, C. Müller, and H.-J. Schüller, 2011 Pleiotropic
642           corepressors Sin3 and Ssn6 interact with repressor Opi1 and negatively regulate
643           transcription of genes required for phospholipid biosynthesis in the yeast Saccharomyces
644           cerevisiae. Mol Genet Genomics 285: 91–100. https://doi.org/10.1007/s00438-010-0589-5

645    Jelinsky S. A., and L. D. Samson, 1999 Global response of Saccharomyces cerevisiae to an
646           alkylating agent. Proc Natl Acad Sci U S A 96: 1486–91.
647           https://doi.org/10.1073/pnas.96.4.1486

Jesch S. A., X. Zhao, M. T. Wells, and S. A. Henry, 2005 Genome-wide analysis reveals inositol,
not choline, as the major effector of Ino2p-Ino4p and unfolded protein response target gene
expression in yeast. J Biol Chem 280: 9106–18. https://doi.org/10.1074/jbc.M411770200
Kitanovic A., T. Walther, M. O. Loret, J. Holzwarth, I. Kitanovic, *et al.*, 2009 Metabolic
response to MMS-mediated DNA damage in Saccharomyces cerevisiae is dependent on the
glucose concentration in the medium. FEMS Yeast Res 9: 535–51.
https://doi.org/10.1111/j.1567-1364.2009.00505.x
Kliewe F., J. Kumme, M. Grigat, S. Hintze, and H.-J. J. Schüller, 2017a Opi1 mediates
repression of phospholipid biosynthesis by phosphate limitation in the yeast Saccharomyces
cerevisiae. Yeast 34: 67–81. https://doi.org/10.1002/yea.3215
Kliewe F., M. Engelhardt, R. Aref, and H.-J. Schüller, 2017b Promoter recruitment of
corepressors Sin3 and Cyc8 by activator proteins of the yeast Saccharomyces cerevisiae.
Curr Genet 63: 739–750. https://doi.org/10.1007/s00294-017-0677-8
Lanz M. C., D. Dibitetto, and M. B. Smolka, 2019 DNA damage kinase signaling: checkpoint
and repair at 30 years. EMBO J 38: e101801. https://doi.org/10.15252/embj.2019101801
Lanz M. C., K. Yugandhar, S. Gupta, E. J. Sanford, V. M. Faça, *et al.*, 2021 In-depth and 3-
dimensional exploration of the budding yeast phosphoproteome. EMBO Rep 22.
https://doi.org/10.15252/embr.202051121
Lee Y. S., K. Huang, F. A. Quiocho, and E. K. O'Shea, 2008 Molecular basis of cyclin-CDK-
CKI regulation by reversible binding of an inositol pyrophosphate. Nat Chem Biol 4: 25–
32. https://doi.org/10.1038/nchembio.2007.52
Longtine M. S., A. McKenzie, D. J. Demarini, N. G. Shah, A. Wach, *et al.*, 1998 Additional
modules for versatile and economical PCR-based gene deletion and modification in
Saccharomyces cerevisiae. Yeast 14: 953–961. https://doi.org/10.1002/(SICI)1097-
0061(199807)14:10<953::AID-YEA293>3.0.CO;2-U
Luévano-Martínez L. A., P. Appolinario, S. Miyamoto, S. Uribe-Carvajal, and A. J.
Kowaltowski, 2013 Deletion of the transcriptional regulator opi1p decreases cardiolipin
content and disrupts mitochondrial metabolism in Saccharomyces cerevisiae. Fungal Genet
Biol 60: 150–8. https://doi.org/10.1016/j.fgb.2013.03.005
Luo H. R., A. Saiardi, H. Yu, E. Nagata, K. Ye, *et al.*, 2002 Inositol Pyrophosphates Are
Required for DNA Hyperrecombination in Protein Kinase C1 Mutant Yeast. Biochemistry
41: 2509–2515. https://doi.org/10.1021/bi0118153
Luo H. R., Y. E. Huang, J. C. Chen, A. Saiardi, M. Iijima, *et al.*, 2003 Inositol Pyrophosphates
Mediate Chemotaxis in Dictyostelium via Pleckstrin Homology Domain-PtdIns(3,4,5)P3
Interactions. Cell 114: 559–572. https://doi.org/10.1016/S0092-8674(03)00640-8
Mulugu S., W. Bai, P. C. Fridy, R. J. Bastidas, J. C. Otto, *et al.*, 2007 A Conserved Family of
Enzymes That Phosphorylate Inositol Hexakisphosphate. Science (1979) 316: 106–109.
https://doi.org/10.1126/science.1139099
Onnebo S., and A. Saiardi, 2009 Inositol pyrophosphates modulate hydrogen peroxide signalling.
Biochem J 423: 109–118. https://doi.org/10.1042/BJ20090241
Rogakou E. P., D. R. Pilch, A. H. Orr, V. S. Ivanova, and W. M. Bonner, 1998 DNA double-
stranded breaks induce histone H2AX phosphorylation on serine 139. J Biol Chem 273:
5858–68. https://doi.org/10.1074/jbc.273.10.5858
Saiardi A., H. Erdjument-Bromage, A. M. Snowman, P. Tempst, and S. H. Snyder, 1999
Synthesis of diphosphoinositol pentakisphosphate by a newly identified family of higher

inositol polyphosphate kinases. Curr Biol 9: 1323–6. https://doi.org/10.1016/s0960-
9822(00)80055-x
Saiardi A., A. C. Resnick, A. M. Snowman, B. Wendland, and S. H. Snyder, 2005 Inositol
pyrophosphates regulate cell death and telomere length through phosphoinositide 3-kinase-
related protein kinases. Proc Natl Acad Sci U S A. https://doi.org/10.1073/pnas.0409322102
Santiago T. C., and C. ben Mamoun, 2003 Genome expression analysis in yeast reveals novel
transcriptional regulation by inositol and choline and new regulatory functions for Opi1p,
Ino2p, and Ino4p. J Biol Chem 278: 38723–30. https://doi.org/10.1074/jbc.M303008200
Schuck S., W. A. Prinz, K. S. Thorn, C. Voss, and P. Walter, 2009 Membrane expansion
alleviates endoplasmic reticulum stress independently of the unfolded protein response. J
Cell Biol 187: 525–36. https://doi.org/10.1083/jcb.200907074
Shah A., S. Ganguli, J. Sen, and R. Bhandari, 2017 Inositol Pyrophosphates: Energetic,
Omnipresent and Versatile Signalling Molecules. J Indian Inst Sci 97: 23–40.
Shi Y., A. N. Azab, M. N. Thompson, and M. L. Greenberg, 2006 Inositol phosphates and
phosphoinositides in health and disease. Subcell Biochem 39: 265–92.
https://doi.org/10.1007/0-387-27600-9_11
Simpson-Lavy K. J., A. Bronstein, M. Kupiec, and M. Johnston, 2015 Cross-Talk between
Carbon Metabolism and the DNA Damage Response in S. cerevisiae. Cell Rep 12: 1865–
75. https://doi.org/10.1016/j.celrep.2015.08.025
Steidle E. A., L. S. Chong, M. Wu, E. Crooke, D. Fiedler, *et al.*, 2016 A Novel Inositol
Pyrophosphate Phosphatase in Saccharomyces cerevisiae: Siw14 PROTEIN
SELECTIVELY CLEAVES THE β-PHOSPHATE FROM 5-DIPHOSPHOINOSITOL
PENTAKISPHOSPHATE (5PP-IP5). J Biol Chem 291: 6772–83.
https://doi.org/10.1074/jbc.M116.714907
Szijgyarto Z., A. Garedew, C. Azevedo, and A. Saiardi, 2011 Influence of inositol
pyrophosphates on cellular energy dynamics. Science (1979) 334: 802–805.
https://doi.org/10.1126/science.1211908
Thibault G., G. Shui, W. Kim, G. C. McAlister, N. Ismail, *et al.*, 2012 The Membrane Stress
Response Buffers Lethal Effects of Lipid Disequilibrium by Reprogramming the Protein
Homeostasis Network. Mol Cell 48: 16–27. https://doi.org/10.1016/j.molcel.2012.08.016
Wagner C., M. Dietz, J. Wittmann, A. Albrecht, and H. J. Schüller, 2001 The negative regulator
Opi1 of phospholipid biosynthesis in yeast contacts the pleiotropic repressor Sin3 and the
transcriptional activator Ino2. Mol Microbiol 41: 155–66. https://doi.org/10.1046/j.1365-
2958.2001.02495.x
Wang Y.-H., A. Hariharan, G. Bastianello, Y. Toyama, G. v. Shivashankar, *et al.*, 2017 DNA
damage causes rapid accumulation of phosphoinositides for ATR signaling. Nat Commun 8:
2118. https://doi.org/10.1038/s41467-017-01805-9
Wilson M. S. C., T. M. Livermore, and A. Saiardi, 2013 Inositol pyrophosphates: between
signalling and metabolism. Biochem J 452: 369–79. https://doi.org/10.1042/BJ20130118
Wimalarathna R., C.-H. Tsai, and C.-H. Shen, 2011 Transcriptional control of genes involved in
yeast phospholipid biosynthesis. J Microbiol 49: 265–73. https://doi.org/10.1007/s12275-
011-1130-1
Winegrad A. I., 1987 Banting Lecture 1986: Does a Common Mechanism Induce the Diverse
Complications of Diabetes? Diabetes 36: 396–406. https://doi.org/10.2337/diab.36.3.396

Worley J., X. Luo, and A. P. Capaldi, 2013 Inositol pyrophosphates regulate cell growth and the
environmental stress response by activating the HDAC Rpd3L. Cell Rep 3: 1476–82.
https://doi.org/10.1016/j.celrep.2013.03.043
Yazgan O., and J. E. Krebs, 2012 Mitochondrial and nuclear genomic integrity after oxidative
damage in Saccharomyces cerevisiae. Front Biosci (Landmark Ed) 17: 1079–93.
https://doi.org/10.2741/3974
Ye C., W. M. M. S. Bandara, and M. L. Greenberg, 2013 Regulation of inositol metabolism is
fine-tuned by inositol pyrophosphates in Saccharomyces cerevisiae. J Biol Chem 288:
24898–908. https://doi.org/10.1074/jbc.M113.493353
Yi C., J. Tong, P. Lu, Y. Wang, J. Zhang, *et al.*, 2017 Formation of a Snf1-Mec1-Atg1 Module
on Mitochondria Governs Energy Deprivation-Induced Autophagy by Regulating
Mitochondrial Respiration. Dev Cell 41: 59-71.e4.
https://doi.org/10.1016/j.devcel.2017.03.007
York S. J., B. N. Armbruster, P. Greenwell, T. D. Petes, and J. D. York, 2005 Inositol
diphosphate signaling regulates telomere length. J Biol Chem 280: 4264–9.
https://doi.org/10.1074/jbc.M412070200
Zewail A., M. W. Xie, Y. Xing, L. Lin, P. F. Zhang, *et al.*, 2003 Novel functions of the
phosphatidylinositol metabolic pathway discovered by a chemical genomics screen with
wortmannin. Proc Natl Acad Sci U S A 100: 3345–50.
https://doi.org/10.1073/pnas.0530118100
Zhou C., A. E. H. Elia, M. L. Naylor, N. Dephoure, B. A. Ballif, *et al.*, 2016 Profiling DNA
damage-induced phosphorylation in budding yeast reveals diverse signaling networks. Proc
Natl Acad Sci U S A 113: E3667–E3675. https://doi.org/10.1073/pnas.1602827113

**Figure Legends**

**Figure 1. Cells lacking Opi1 show sensitivity to genotoxic stress.** (A) Working model

for how Opi1 repress transcription of target genes; (B-C) Effect of deletion of genes

involved in inositol metabolism in sensitivity to genotoxins; (D) Immunoblot showing

expression of Opi1-HA from a pRS416 plasmid; (E) A pRS416 plasmid expressing Opi1

from its endogenous promoter complement an *opi1Δ* strain; For (B, C and E), fourfold

serial dilutions were spotted on the media indicated in the figure and plates were grown

for 2–3 days at 30°C.

**Figure 2. Cells lacking Opi1 show defects upon replication stress induced by MMS.**

(A) S-phase progression analysis of wild-type and *opi1Δ* strains. Cells were treated with

0.033% MMS for 2 hours and then release in fresh YPD medium; (B and C) Immunoblot

analysis of gamma-H2A in the indicated strains. For (B), cells were treated with 0.033%

MMS for 2 hours and then release in fresh YPD medium until 6 hours. For (C), cells were

arrested in G1 with $\alpha$-factor and then released in medium containing MMS. Immunoblots

were probed with anti-gamma-H2A and anti-Pgk1 (loading control) antibodies as

described in *Material and methods*. (D) S-phase progression analysis of wild-type and

*opi1Δ* strains. Cells were arrested in G1 with $\alpha$-factor and then released in medium

containing MMS.

**Figure 3. Opi1 is required to repress inositol synthesis during genotoxic stress.** (A-

B) Effect of inositol availability and MMS treatment on Opi1-GFP localization. Cellular

localization of Opi1-GFP was scored manually and the means and standard deviations

from two independent experiments are shown; (C) Cells treated with MMS in medium

lacking inositol are still viable; (D) Effect of inositol availability and MMS treatment on

*INO1* expression by RT-qPCR. Relative values of mRNA are described as fold change

relative to a control condition (*wt* + INO). For C, fourfold serial dilutions were spotted on

the media indicated in the figure and plates were grown for 2–3 days at 30°C. For D,

experiments were made with three biological replicates and statistical significance of the

differences observed was calculated using one one-way ANOVA with Tukey's multiple

comparisons test. Of note, statistical significances between *wt + INO (MMS) vs. opi1Δ +*

*INO (MMS) (p = 0.003); wt - INO (MMS) vs. opi1Δ - INO (MMS) (p = 0.0006); opi1Δ - INO*

*vs. opi1Δ - INO (MMS) (p = 0.0129).*

**Figure 4. Deletion of the transcriptional activators Ino2-Ino4 rescues genotoxin**

**sensitivity and mitochondrial defects of cells lacking Opi1.** (A-B) Deletion of *INO*2

and *INO*4 completely rescue the MMS sensitivity of cells lacking Opi1; (C) Deletion of

*INO*2 completely restores mitochondrial respiratory capacity of cells lacking Opi1; (D) –

Deletion of *INO*2 does not completely rescue the MMS sensitivity of cells lacking Opi1 in

respiratory medium (YPG). For A, B and D, fourfold serial dilutions were spotted on the

media indicated in the figure and plates were grown for 2–3 days at 30°C.

**Figure 5. Deregulation of inositol pyrophosphate synthesis in cells lacking Opi1 is**

**causing genotoxin sensitivity.** (A) Deletion of Ino1 does not rescues the MMS

sensitivity of cells lacking Opi1; (B-C) Deletion of the inositol pyrophosphate kinase Kcs1

rescues the MMS sensitivity and replication defects in cells lacking Opi1. (D) Cells lacking

Siw14 are sensitive to MMS; (E) Overexpression of Kcs1 causes MMS sensitivity. For (A-

B, D and E), fourfold serial dilutions were spotted on the medium indicated in the figure

and plates were grown for 2–3 days at 30°C.

823
824    **Figure 6. Proposed model illustrating how Opi1 modulate PP-IPs synthesis during**
825    **genotoxic stress.**

**Figure 1**

A

B

C

D

E

# Figure 2

# Figure 3

A

Exp. growing yeast culture (+ Inositol) → Release from media supplemented with inositol → MMS treatment

(SC + INOSITOL)   1 hour   (SC - INOSITOL)

+ INOSITOL · 5 µm

- INOSITOL · 5 µm

- INOSITOL + MMS · 5 µm

B

Frequency of cells (%)

- Nuclear
- Transition
- Perinuclear

t0 +INO · t0 -INO · 15' -INO (MMS) · 30' -INO (MMS)

C

Control · MMS 0.01%

SC + Inositol · *wild-type* · *opi1Δ*

SC - Inositol · *wild-type* · *opi1Δ*

D

*wt* + INO
*wt* + INO (MMS)
*wt* - INO
*wt* - INO (MMS)
*opi1Δ* + INO
*opi1Δ* + INO (MMS)
*opi1Δ* - INO
*opi1Δ* - INO (MMS)

*INO1* mRNA relative expression

# Figure 4

A

| | Control (YPD) | MMS 0.015% | MMS 0.0175% | Bleomycin 5µg/ml |
|---|---|---|---|---|
| *wild-type* | | | | |
| *opi1Δ* | | | | |
| *ino2Δ* | | | | |
| *opi1Δino2Δ* | | | | |

B

| | Control (YPD) | MMS 0.015% | MMS 0.0175% |
|---|---|---|---|
| *wild-type* | | | |
| *ino4Δ* | | | |
| *opi1Δ* | | | |
| *opi1Δino4Δ* | | | |

C

D

| | | YPG | |
|---|---|---|---|
| | Control | MMS 0.005% | MMS 0.01% |
| *wild-type* | | | |
| *opi1Δ* | | | |
| *ino2Δ* | | | |
| *opi1Δino2Δ* | | | |

**Figure 5**

A

B

C

D

E

**Figure 6**