Subcytoplasmic location of translation controls protein output

Ellen L. Horste[1,2], Mervin M. Fansler[2,3], Ting Cai[2], Xiuzhen Chen[2], Sibylle Mitschka[2], Gang
Zhen[2], Flora C. Y. Lee[4,5], Jernej Ule[4,5], and Christine Mayr[1,2,3†]

[1]Gerstner Sloan Kettering Graduate School of Biomedical Sciences, New York, NY 10065, USA
[2]Cancer Biology and Genetics Program, Sloan Kettering Institute, New York, NY 10065, USA
[3]Tri-Institutional Training Program in Computational Biology and Medicine, Weill-Cornell
Graduate College, New York, NY 10021, USA
[4]UK Dementia Research Institute, King's College London, London, SE5 9NU, UK
[5]The Francis Crick Institute, 1 Midland Road, London NW1 1AT, UK

[†]Correspondence: mayrc@mskcc.org

Summary

The cytoplasm is highly compartmentalized, but the extent and consequences of
subcytoplasmic mRNA localization in non-polarized cells are largely unknown. We determined
mRNA enrichment in TIS granules (TGs) and the rough endoplasmic reticulum (ER) through
particle sorting and isolated cytosolic mRNAs by digitonin extraction. When focusing on non-
membrane protein-encoding mRNAs, we observed that 52% have a biased transcript
distribution across these compartments. Compartment enrichment is determined by a
combinatorial code based on mRNA length, exon length, and 3′UTR-bound RNA-binding
proteins. Compartment-biased mRNAs differ in the functional classes of their encoded proteins:
TG-enriched mRNAs encode low-abundance proteins with strong enrichment of transcription
factors, whereas ER-enriched mRNAs encode large and highly expressed proteins.
Compartment localization is an important determinant of mRNA and protein abundance, which
is supported by reporter experiments showing that redirecting cytosolic mRNAs to the ER
increases their protein expression. In summary, the cytoplasm is functionally compartmentalized
by local translation environments.

Keywords

Cytoplasmic organization; spatial regulation of protein synthesis; cytoplasmic compartments;
condensates; TGs; mRNA localization; gene expression regulation; functional
compartmentalization; translation environment; translation regulation; compartment-dependent
mRNA abundance regulation; rough endoplasmic reticulum membrane; gene architecture;
mRNA length; exon length; 3′UTR; RNA-binding proteins; CLIP analysis; TIS11B; TIAL1;
LARP4B; TIS11B knockout

Introduction

In polarized cells such as neurons, intestinal epithelial cells, or cells of the early fly embryo, the
majority of mRNAs have a distinct spatial localization pattern. [1-5] mRNA localization enables the
local control of protein production and activity. [6-8] In non-polarized cells, mRNA localization has
primarily been studied for membrane proteins, which mostly localize to the endoplasmic
reticulum (ER) or to mitochondria. [9-12] Whereas the rough ER is established as major site of
local protein synthesis for membrane and secretory proteins, [9,10,13] the cytoplasm is
compartmentalized by additional membrane-bound and membraneless organelles. [14-17] Some of
these compartments may enable the generation of unique biochemical translation

environments, which have been suggested to be crucial for protein interaction partner selection
during protein synthesis, [16,18-20] but it is currently largely unknown if the location of protein
synthesis also matters for protein output.

TIS granules (TGs) represent one such unique translation compartment, which promotes the co-
translational formation of protein complexes. [16,19] TGs are formed by the RNA-binding protein
(RBP) TIS11B together with its bound mRNAs. [16,21] *TIS11B* mRNA is ubiquitously expressed, [22]
suggesting that TGs are widespread. TGs are present under steady-state cultivation conditions
and form a network-like structure that is intertwined with the rough ER. [16,21] To investigate the
broader biological significance of TGs, we set out to determine the mRNAs enriched in TGs, the
neighboring rough ER, and the surrounding cytosol.

As TIS11B protein is present in cells in two states (Fig. 1A), (i) as soluble protein in the cytosol,
and (ii) as phase-separated TG network, [16,21] we decided to use fluorescent particle sorting [23] to
identify mRNAs enriched in TGs. We also applied fluorescent particle sorting to isolate ER-
enriched mRNAs and extracted cytosolic mRNAs using digitonin. We focused our analysis on
mRNAs that encode non-membrane proteins and found more than 3600 mRNAs to be
consistently enriched in one of the three compartments. mRNAs enriched in each compartment
share similar mRNA architectures which differ substantially between compartments.
Compartment-enriched mRNAs also differed significantly in production and degradation rates as
well as in the functional classes and expression levels of their encoded proteins. TIS11B
knockout (KO) and reporter experiments support a model by which a combinatorial code
consisting of mRNA architecture features together with 3′UTR-bound RBPs, including TIS11B,
TIA1/L1, and LARP4B, largely determines the compartment-biased mRNA localization pattern.
Intriguingly, we observed that redirecting cytosolic mRNAs to the ER controls protein
expression, which indicates that protein abundance regulation is spatially regulated in the
cytoplasm.

Results

Approach to determine subcytoplasmic mRNA localization

We set out to identify mRNAs that are localized in non-polarized human HEK293T cells under
steady-state cultivation conditions. We focused on three major unenclosed cytoplasmic
compartments—TGs, a condensate network formed by the RBP TIS11B, the cytosolic surface
of the ER, and the soluble part of the cytoplasm known as the cytosol (Fig. 1B). For simplicity,
we consider here the sum of the three compartments as the universe of cytoplasmic mRNAs.

To identify TG-enriched (TG+) and ER-enriched (ER+) mRNAs, we performed fluorescent
particle sorting followed by RNA-seq. After co-transfecting cells with mCherry-TIS11B and GFP-
SEC61B to label TGs and rough ER, respectively, we used mechanical lysis and differential
centrifugation to isolate the cytoplasmic membrane fraction, followed by flow cytometry-based
sorting of fluorescent particles (Fig. S1A, S1B). DAPI staining allowed us to identify and discard
TG and ER particles that were still associated with nuclei. We used confocal microscopy to
assess the purity of the particles. While ER particles generally did not contain mCherry-TIS11B,
most of the TG particles contained GFP-SEC61B, consistent with the intimate association of TG
and ER inside cells (Fig. 1C). Using western blot analysis, we observed that both particles
contain similar amounts of ER proteins such as Calnexin and GFP-SEC61B, but TG particles
contain 13-fold more mCherry-TIS11B than ER particles (Fig. S1C-E). As TG are defined by the
presence of TIS11B, [16] we reasoned that the strong overrepresentation of TIS11B in TG
particles would allow us to identify relative enrichments of mRNAs between the compartments.

For subsequent RNA-seq experiments, we sorted TG particles from the TIS11B+SEC61B+
population and we sorted ER particles from the TIS11B−SEC61B+ population (Fig. S1B).

To isolate cytosolic mRNAs, we used digitonin extraction. [24] The extracted cytosol was not
contaminated by nuclei or the ER, but it contained cytosolic proteins, including GAPDH which
was used as a positive control (Fig. S1C). It also contained TIS11B, which was expected since
soluble TIS11B is known to be present in the cytosol. We performed RNA-seq to determine the
mRNA composition in the three fractions and focused our analysis on protein-coding mRNAs.
The biological replicate samples of each compartment correlated well (Fig. S1F).

mRNAs that encode membrane or secretory proteins largely localize to the ER membrane

We set out to investigate if the relative mRNA transcript distribution differs across the three
compartments. For each gene, we determined a compartment-specific localization score. This
score is calculated using the RPKM value obtained in each of the three compartments
respectively and dividing it by the sum of the RPKM values in all three compartments. Thus,
each gene is assigned three localization scores that correspond to the fraction of its transcripts
localizing to each of the three compartments: TGs, the ER, and the cytosol.

It is accepted that most mRNAs that encode membrane or secretory proteins are translated on
the ER. [10,11,13] To account for the difference in distribution, we plotted the localization scores of
mRNAs that encode membrane/secretory proteins separately from the rest of the mRNAs that
encode non-membrane proteins (Fig. S1G). In line with previous analyses, we find preferential
partitioning of mRNAs encoding membrane/secretory proteins in the ER samples (Fig. S1G).
[10,11,13] To examine if our compartment isolation method is valid, we compared it with datasets
derived from three alternative isolation methods. [9,11,13] In our analysis, we consider 69% ($N =$
1,476) of membrane/secretory proteins to be enriched on the ER (Fig. S1H, Table S1). When
comparing our results with mRNAs identified by biochemical fractionation, MERFISH or APEX-
seq, we detected between 80-90% overlap and our data showed a quantitative relationship with
data obtained by APEX-seq (Fig. S1I, S1J). [9,11,13] These results strongly support the validity of
our purification strategy for mRNAs that encode membrane/secretory proteins.

Half of mRNAs that encode non-membrane proteins have a biased transcript distribution
in the cytoplasm

It was unclear whether non-membrane protein encoding mRNAs are biased in their localization.
Not surprisingly, we found that these mRNAs have more evenly distributed localization scores
across the three compartments (Fig. S1G). To faithfully compare absolute differences in mRNA
distribution across the three compartments, the relative size of each compartment needs to be
considered. However, this parameter is currently unknown. Therefore, instead, we calculated
the relative enrichment of mRNAs within each compartment. We considered an mRNA
compartment-enriched, if its mean localization score across biological replicates was at least
1.25-fold higher than the median localization score of the compartment samples (Fig. 1D-F).
Based on this criterion, we identified 1246 TG+ mRNAs, 919 ER+ mRNAs, and 1481 mRNAs
enriched in the cytosol (CY+), which were non-overlapping (Fig. 1D-F, Table S1). The remaining
3369 mRNAs were not enriched in a single compartment and were considered to have an
unbiased localization pattern (Fig. 1D-F, Fig. S1K). Fig. 1D-F illustrate that the distribution of
TG+, ER+, or CY+ localization scores are significantly different from the localization scores of
mRNAs with unbiased localization patterns. Since localization scores across the three
compartments sum to 1, an mRNA enriched in one compartment is relatively de-enriched in the
other two (Fig. S1L). Based on this strategy, 52% of mRNAs that encode non-membrane

proteins are significantly enriched in one of the three subcytoplasmic compartments in steady-
state conditions.

As a recent study also analyzed the relative distribution of mRNA transcripts across subcellular
compartments, we compared our compartment enrichment data with their results. [25] Although
their dataset was generated by density gradient centrifugation in a different cell line, the two
compartment enrichment datasets strongly agree in a qualitative and quantitative manner (Fig.
S1M), suggesting that our isolation method as well as our strategy to define compartment-
enriched mRNAs are valid. As non-membrane protein encoding mRNAs with biased transcript
distributions in the cytoplasm have not been systematically characterized, we focused all
subsequent analyses on mRNAs that encode non-membrane proteins.

Validation of compartment-enriched mRNAs by single-molecule RNA-FISH

We further validated the mRNAs designated as compartment-enriched by performing single-
molecule (sm) RNA-FISH on endogenous mRNAs (Table S2). [26] To distinguish between TG+
and ER+ mRNAs, we performed smRNA-FISH together with co-transfection of BFP-TIS11B and
GFP-SEC61B to simultaneously visualize mRNA puncta, TGs, and the rough ER (Fig. 1G, 1H,
Fig. S2A-G). Here, we considered an mRNA to have an unbiased localization pattern if its
transcript distribution correlated with compartment size. As proxy for relative compartment size,
we used the areas of the maximum projection of the fluorescent signals for each compartment
and compared them to the whole cell area. Based on the compartment size distribution across
186 cells, for unbiased mRNAs, we expect that 11% of transcripts localize to TGs and 29% of
transcripts localize to the ER (Fig. 1I, 1J).

For 3/3 TG+ mRNAs, we observed a significant enrichment of mRNA puncta in TGs, but not on
the ER (Fig. 1G, 1I, 1J, Fig. S2A, S2B, S2H, S2I). For the five ER+ mRNAs tested, the mRNA
puncta of 4/5 mRNAs were significantly enriched on the ER and for all five, we observed a 2-4-
fold higher fraction of mRNA puncta that co-localized with the ER compared to TGs (Fig. 1H-K,
Fig. S2C-I).

Cytosolic mRNAs were isolated through digitonin extraction. This means that CY+ mRNAs
localize to the soluble part of the cytoplasm and are not attached to cytoplasmic structures,
including membranes or the cytoskeleton. As steady-state smRNA-FISH can only inform on co-
localization and not attachment, we validated CY+ mRNAs by performing smRNA-FISH before
and after digitonin extraction and calculated the fraction of retained mRNAs. We observed
significantly greater retention of both TG+ and ER+ mRNAs compared to CY+ mRNAs, which
were depleted by about 90% following digitonin treatment (Fig. 1L, 1M, Fig. S3A-C). This
confirms that CY+ mRNAs predominantly localize to the soluble part of the cytoplasm. Taken
together, as we successfully validated 10/11 mRNAs that were designated to be TG+ or ER+ or
CY+ (Fig. 1N), we conclude that about half (52%) of mRNAs encoding non-membrane proteins
are enriched in distinct subcytoplasmic compartments.

mRNA and protein levels strongly correlate with the location of translation

Next, we characterized the features of compartment-enriched mRNAs and found substantial
differences in their steady-state mRNA and protein levels (Fig. 2A, 2B, Fig. S4A, S4B). We
observed that TG+ mRNAs have the lowest steady-state expression levels and encode proteins
with the lowest expression levels (Fig. 2A, 2B). To examine if the low mRNA levels are caused
by high mRNA degradation rates, we estimated mRNA half-lives by analyzing Precision Run-On
sequencing (Pro-seq) and RNA-seq data (Fig. 2C, 2D, Fig. S4C-E). [27,28] Pro-seq values can be

treated as transcription rates and RNA-seq data can be viewed as a measure of RNA
concentration to estimate RNA decay rates required for a steady-state equilibrium. [28] For TG+
mRNAs, we observed that the low steady-state mRNA levels were not primarily caused by a low
mRNA stability. Instead, these mRNAs had the lowest transcription rates, suggesting that these
mRNAs are either produced at a low rate or have high cotranscriptional degradation rates (Fig.
2C, 2D, Fig. S4D, S4E). [29] CY+ mRNAs had the highest degree of mRNA turnover with both
high production and degradation rates (Fig. 2C, 2D). ER+ mRNAs encode proteins with the
highest expression levels, particularly when normalizing to their intermediate steady-state
mRNA levels (Fig. 2A, 2B).

In addition, we observed that the compartment-enriched mRNAs differed substantially in their
gene architectures (Fig. 2E-H, Fig. S4F-K). ER+ mRNAs encode the largest proteins with a
median size of 840 amino acids, nearly three-times larger than proteins encoded by CY+
mRNAs (Fig. 2E). The difference in protein size was reflected in the large differences in exon
number and mRNA length between ER+ and CY+ mRNAs (Fig. 2F, Fig. S4J, S4K). The median
length of ER+ mRNAs is 4600 nucleotides (nt), whereas the median length of CY+ mRNAs is
2000 nt. Not surprising, CY+ mRNAs also have the shortest 3′UTRs (Fig. 2G). TG+ mRNAs are
uniquely characterized by large coding sequence (CDS) exons with a median size of 200 nt,
compared to a median exon size of 133 nt for the remaining mRNAs (Fig. 2H). Further analysis
revealed that the majority of TG+ mRNAs have a gene architecture that is similar to the gene
architecture of *ZFP36L1* (encoding TIS11B), which is characterized by a short first exon and a
long last exon that contains ~95% of its coding sequence (Fig. 2I).

Moreover, compartment-enriched mRNAs encode substantially different functional gene
classes. [30] Consistent with the low steady-state protein expression levels, TG+ mRNAs were
strongly enriched in proteins containing zinc fingers and in transcription factors, which are
known to have low expression (Fig. 2J). [31] In contrast, ER+ mRNAs encode large and highly
abundant proteins, and they were enriched in helicases, cytoskeleton-binding proteins, and
chromatin regulators (Fig. 2K). CY+ mRNAs often encode smaller proteins involved in the
regulation of translation or splicing (Fig. 2L).

TGs support active translation

TIS granules may constitute a specialized translation environment for nuclear proteins that
require low expression levels, such as transcription factors (Fig. 2A, 2B, 2J). [31] In order to
provide evidence for active translation in TIS granules, we applied the SunTag system to
simultaneously visualize mRNAs and their nascent proteins in TGs and in the cytosol (Fig. S3D,
S3E). [32] We confirmed that TGs represent a translation environment for mRNAs. [16,19] We
observed that the number of mRNA foci in TGs was five-fold lower compared to the cytosol.
However, the proportion of mRNA translated was similar in TGs and the cytosol (Fig. S3F,
S3G). Taken together, our data show that TGs are sites of active translation and that the low
expression level of TG-translated proteins is predominantly a result of their low nuclear gene
expression (Fig. 2A, 2C).

Differential 3′UTR binding of several RBPs correlates with compartment enrichment of mRNAs

Our next goal was to identify the RBPs responsible for mRNA enrichment in the three
compartments (Fig. 1D-F). As TIS11B is the scaffold protein of TGs, [16] we performed iCLIP of
TIS11B in HEK293T cells (Fig. S5A, S5B). We confirmed that the top binding motif of TIS11B in
3′UTRs of mRNAs is the canonical AU-rich element (UAUUUA) (Fig. S5C). To perform a

comprehensive analysis on localization regulators, we included additional CLIP datasets. [33,34]
Altogether, we analyzed CLIP data from 170 RBPs and found that 24 of them showed an
enrichment of binding sites in 3′UTRs of mRNAs belonging to transcripts that preferentially
localize to one of the three compartments (Table S4). We applied logistic regression and
identified seven RBPs whose binding contributed most significantly to mRNA enrichment in the
three compartments. They include TIS11B, HuR, PUM2, HNRNPC, TIA1/L1, LARP4B and
METAP2 (Fig. 3A). As a previous CLIP analysis showed that peaks for TIA1 and TIAL1 cannot
be distinguished, [35] we used the sum of peaks from TIA1 and TIAL1 to obtain the values for
TIA1/L1. According to this regression analysis, the presence of TIS11B, HuR, PUM2, and
HNRNPC on mRNAs correlates with TG enrichment, the presence of TIA1/L1 correlates with
ER enrichment, and the presence of LARP4B or METAP2 correlates with cytosol enrichment
(Fig. 3A).

mRNA architecture features together with RBPs generate a combinatorial code for
subcytoplasmic transcript distribution

As our analysis revealed that 2154 mRNAs (30.7%) that encode non-membrane proteins were
not bound by any of the seven RBPs (Fig. S5D), we considered additional regulatory factors for
mRNA localization. Among these mRNAs, we observed that mRNA length correlated strongly
with the ER and CY localization scores, but in opposite directions, suggesting that long mRNAs
are associated with ER localization (Fig. 3B). Similarly, average CDS exon length correlated
strongly and in a positive manner with the TG localization score, but negatively with the CY
localization score (Fig. 3B).

When including mRNA and exon length in the logistic regression, we observed that mRNA
architecture features contribute strongly to mRNA enrichment in each of the three
compartments (Fig. 3C, Table S4). Our data indicate that the propensity of an mRNA for
compartment localization depends on both architectural features and the presence of 3′UTR-
bound RBPs. To better understand the rules for mRNA localization to the compartments, we
plotted the propensity for TG enrichment and integrated RBPs that were positively or negatively
associated with TG enrichment together with exon length (Fig. 3D). Regardless of exon length,
binding of LARP4B/METAP2 decreased the TG localization score (Fig. 3D, 3E). For mRNAs
with average exon length, TIS11B binding strongly increased their localization propensity to
TGs. A similar TG localization propensity was observed for mRNAs with long exons that were
not bound by TIS11B. The strongest TG enrichment was found for mRNAs with long exons that
were bound by TIS11B, indicating that TIS11B and exon length have additive effects (Fig. 3D,
3E).

The two features that best promote mRNA localization to the ER are mRNA length and 3′UTR-
bound TIA1/L1 (Fig. 3F, 3G). Short mRNAs increase their ER localization propensity by the
presence of TIA1/L1. Long mRNAs that are not bound by TIA1/L1 have a similar propensity for
ER localization, which can be further increased by the presence of TIA1/L1 (Fig. 3F, 3G). In
contrast, shorter mRNAs not bound by any RBP or that are bound by LARP4B/METAP2 have a
high propensity to localize to the cytosolic fraction (Fig. 3G, 3H). Taken together, our data
suggest the existence of a combinatorial code, that integrates mRNA and exon length with the
presence of RBPs, to determine subcytoplasmic mRNA localization (Fig. 3E, 3G).

mRNA transcript distribution changes upon TIS11B deletion

Next, we set out to experimentally test the proposed mRNA localization code by investigating
the influence of TIS11B and TIA1/L1 on subcytoplasmic mRNA transcript distribution. We

generated HEK293T cells with an inducible knockout (KO) of TIS11B, isolated ER particles and
extracted the cytosol (Fig. S5E, S5F, Table S5). To examine where mRNAs designated as TG+
localize in the absence of TGs, we identified the top 20% of mRNA localization changes to the
ER and the cytosol and intersected them with mRNAs designated as TG+ (Fig. 4A). As only two
compartments were isolated, increased mRNA localization to the ER means decreased
cytosolic localization and vice versa (Fig. 4A).

We did not find specific RBPs associated with the localization-changing mRNAs, because TG+
mRNAs are mostly bound by TIS11B and only a small minority of them (13% and 15%) are
LARP4B or TIA1/L1 targets (Table S5). However, the TG+ mRNAs that increased their cytosolic
localization upon TIS11B KO were the shortest, encoded the smallest proteins, and had the
shortest exon length (Fig. 4B-E). In contrast, TG+ mRNAs that increased their ER localization
upon deletion of TIS11B were significantly longer, encoded the largest proteins, and had longer
exons (Fig. 4B-E).

These results support a model in which mRNA architecture features set up a 'default' steady-
state mRNA transcript distribution pattern in the cytoplasm which can be overcome or reinforced
through the binding of RBPs. Short mRNAs with average exon length localize to TGs when
bound by TIS11B but in the absence of TIS11B they revert to the transcript distribution
established by mRNA architecture and the remaining bound RBPs, in this case the cytosol (Fig.
3E). Similarly, longer TG+ mRNAs that encode the largest proteins localize to the ER upon loss
of TIS11B (Fig. 3G). Currently, the 'readers' of the mRNA architecture features are unknown.

3′UTR-bound TIAL1 promotes localization of non-membrane protein-encoding mRNAs to the ER

We observed that ER+ mRNAs were enriched in 3′UTR-bound TIA1/L1, which had not been
previously reported (Fig. 3A, 3C, 3F). To test the influence of TIA1/L1 on mRNA localization, we
set out to investigate mRNA localization changes to the three compartments in TIA1/L1 double
KO cells. [36] However, as previously reported, these cells showed a high rate of cell death, which
prevented us from obtaining high-quality particles.

To validate TIA1/L1-dependent mRNA localization to the ER, we used the MS2 tethering system
to mimic 3′UTR-binding of TIA1/L1 (Fig. 4F). We generated a *GFP-THAP1* reporter mRNA that
contained MS2-binding sites as 3′UTR. [37-39] Coexpression of mCherry-tagged MS2 coat protein
(MCP) fused to TIAL1 tethers TIAL1 to the 3′UTR of the reporter mRNA (Fig. 4F). As a control,
mCherry-tagged MCP was tethered to the *GFP-THAP1* reporter mRNA.

Coexpression of the reporter mRNA and MCP resulted in evenly distributed cytosolic expression
of both MCP protein and reporter mRNA (Fig. 4F-H). In contrast, coexpression of the reporter
mRNA and MCP-TIAL1 resulted in perinuclear, reticulated expression of MCP-TIAL1 with the
mRNA reporter predominantly localizing to the rough ER (Fig. 4F-H). Colocalization was
assessed by RNA-FISH of the GFP-tagged reporter mRNA and simultaneous visualization of
the rough ER through fluorescently tagged SEC61B. Using line diagrams of the fluorescence
intensities, we quantified the overlap between the reporter mRNAs and the ER (Fig. 4I). In the
presence of MCP-TIAL1, we observed higher correlation coefficients between the reporter
mRNA and the ER (Fig. 4J). This result indicated that 3′UTR-bound TIAL1 was sufficient to
induce localization of non-membrane protein encoding mRNAs to the rough ER surface.

3′UTR-bound TIAL1 increases protein expression

For endogenous mRNAs, we observed that ER+ mRNAs encode the highest expressed
proteins (Fig. 2B). Moreover, mRNAs predominantly bound by TIA1/L1 encode proteins with
higher expression levels compared with other mRNAs (Fig. 5A). Using our mRNA reporter (Fig.
4F), we investigated the contribution of TIAL1 to steady-state protein expression. We used
FACS to measure GFP protein expression of the mRNA reporter with or without tethering of
TIAL1 to the 3′UTR (Fig. S6A-C). We observed a 3.5-fold increase in protein expression upon
3′UTR-tethering of TIAL1 compared to tethering of MCP alone (Fig. 5B, 5C). Increased GFP
protein expression was not due to an increase in mRNA abundance (Fig. 5D). We confirmed the
TIA1/L1-dependent increase in protein expression using a second GFP reporter (Fig. S6D-F).
As TIAL1 promotes translation of mRNAs on the ER membrane, it was unclear if increased
protein expression was caused by TIAL1 or by a potentially unique translation environment
provided by the rough ER membrane. For example, it was reported that mRNAs that encode
non-membrane proteins contain 1.4-fold more ribosomes when translated on the ER membrane
than when translated in the cytosol. [40]

TIAL1 cooperates with the rough ER environment to promote protein expression

To disentangle the effects of TIAL1 and the ER membrane on protein expression, we tethered
the reporter mRNA directly to the ER surface by fusing MCP to SEC61B, a subunit of the
translocon complex in the rough ER (Fig. 5E). MCP-SEC61B perfectly colocalized with the ER
and recruited reporter mRNAs to the ER (Fig. 5F, S6G-I). However, reporter protein expression
only increased by 1.25-fold compared to the tethering of MCP alone and did not increase mRNA
abundance of the reporter (Fig. 5G-I). We used a second ER localization reporter by fusing
MCP to TRAPα, which represents a different subunit of the translocon complex and obtained a
similar result. We observed an increase in protein expression by 1.5-fold when the reporter
mRNA was tethered to TRAPα (Fig. S6J-M). These results suggested that the ER membrane
environment has a significant but small stimulatory effect on translation.

Next, we investigated if the TIAL1-dependent increase in protein expression is intrinsic to TIAL1
or if it depends on its localization to the ER membrane. We added a CAAX motif to TIAL1 to
localize the TIAL1-bound mRNA reporter to the plasma membrane instead of the ER membrane
(Fig. 5J). The CAAX signal is a prenylation motif that efficiently localized MCP and MCP-TIAL1
to the plasma membrane (Fig. 5K). [32] Translation of the TIAL1-bound mRNA reporter at the
plasma membrane increased protein expression by 1.8-fold (Fig. 5L, 5M). As translation of the
TIAL1-bound reporter at the ER membrane resulted in two-fold higher protein expression than
its translation at the plasma membrane (Fig. 5M), our result suggested that TIAL1 cooperated
with the environment on the rough ER membrane to promote protein expression.

As the RBPs bound to the reporter mRNA were identical in these experiments, our results
demonstrate that the subcytoplasmic location of translation controls steady-state protein
expression levels by two-fold when comparing plasma and ER membranes. This relationship
was also observed for endogenous mRNAs, where TIA/L1 bound mRNAs were associated with
high protein output in every compartment, but with the highest protein yields being observed in
the ER compartment (Fig. 5N).

The repressive effect of cytosolic TIS11B on protein expression is overcome by its
localization to rough ER membrane

Next, we examined if the environment on the rough ER membrane also promotes protein
expression of mRNAs bound by other RBPs, including TIS11B (Fig. 6A, 6B). In cells expressing
mCherry-TIS11B fusion constructs, about 30% form TGs at steady state (Fig. S7A, S7B). [16]

However, we noticed that addition of MCP to TIS11B fusion constructs resulted in limited TG
formation and predominant expression of TIS11B in the cytosol (Fig. S7A, S7B). In the cytosolic
state, binding of MCP-TIS11B to the reporter mRNA repressed reporter protein expression by
two-fold, compared to tethering of MCP alone (Fig. 6C, 6D). This decrease in protein expression
was partially caused by a TIS11B-dependent decrease in mRNA level (Fig. 6E), consistent with
previous reports that suggested that cytosolic TIS11B represses the expression of certain
cytokine and cell cycle mRNAs. [41-43] In contrast, fusing TIS11B to MCP-SEC61B localizes
TIS11B and the bound reporter mRNA to the rough ER (Fig. 6A, 6B), which overcomes the
repressive effect of cytosolic TIS11B and increased protein expression two-fold (Fig. 6A-E). The
two-fold increase in protein expression was recapitulated with a second reporter and indicates
that the repressive effect on protein expression mediated by cytosolic TIS11B is overcome by
translation of the TIS11B-bound mRNA on the ER (Fig. 6D, Fig. S7C-E).

Model

Taken together, we observed that mRNAs that are uniquely enriched in one of three
cytoplasmic compartments differ substantially in their architectural features, in the RBPs bound
to them, and in the expression levels and functional classes of their encoded proteins (Fig. 7).
TG+ mRNAs are characterized by the longest CDS exons and TIS11B binding to the 3′UTR.
These mRNAs encode the lowest abundance proteins with a strong enrichment of transcription
factors. In contrast, although TGs are intertwined with the rough ER, ER+ mRNAs are the
longest, are predominantly bound by TIA1/L1, and encode highly abundant large proteins. CY+
mRNAs are the shortest and encode small and highly abundant proteins. They are bound by
LARP4B/METAP2 and have high production and degradation rates (Fig. 7). Moreover, by using
mRNA reporters, we showed that relocation of protein synthesis from the cytosol to the ER
increases protein expression, indicating that the location of translation influences protein output
(Fig. 7).

Discussion

We determined the distribution of endogenous mRNA transcripts across three cytoplasmic
compartments, including TGs, the rough ER, and the cytosol under steady-state conditions. Our
RNA-seq results, which were validated by smRNA-FISH, suggest that approximately half of all
mRNAs are uniquely enriched in one of these three cytoplasmic compartments.

Functionally related genes are translated in unique compartments

One of our most striking findings was that within each investigated compartment a different
group of functionally related mRNAs is translated (Fig. 2). Moreover, the compartment-enriched
mRNAs have vastly different gene architectures and are characterized by substantially different
production and degradation rates as well as the expression levels of their encoded proteins (Fig.
2). These features are consistent with the compartment-enriched gene groups, indicating that
the cytoplasm is strongly partitioned into different functional and regulatory compartments that
are not enclosed by membranes.

Surprisingly, we observed that transcription factors are substantially enriched among TG+
mRNAs (Fig. 2J). This unexpected result can be explained by the previous observation that
transcription factors are often present in low abundance [31] and we found that TG+ mRNAs
encode the proteins with the lowest expression levels (Fig. 2B). Moreover, many transcription

factors have an unusual gene architecture with longer than average coding exons. Together
with TIS11B binding, this was the strongest feature for mRNA enrichment in TGs (Fig. 3C-E).
Interestingly, both characteristics correlate with features associated with low mRNA abundance
levels, but whereas TIS11B-binding correlates negatively with pre-mRNA production rates
(Spearman's correlation coefficient R = -0.26), exon length negatively correlates with mRNA
half-life (Spearman's correlation coefficient R = -0.34). [44] The unique gene architecture together
with predominant binding of TIS11B provides an explanation for why TGs enrich for low-
abundance mRNAs.

In contrast, ER+ mRNAs encode the largest proteins with the highest expression levels. These
include helicases, cytoskeleton-bound proteins, and chromatin regulators (Fig. 2). It is possible
that anchoring of ribosomes on the ER membrane may facilitate the protein synthesis of very
large proteins. Moreover, it is notable that despite the intertwinement of TGs and the rough ER,
the compartment-enriched mRNAs encode proteins that differ substantially in their expression
levels, which are the lowest for TG+ mRNAs and the highest for ER+ mRNAs.

It was previously shown that localization to the ER membrane of certain non-membrane protein
encoding mRNAs increases their translation, [9,40] and we confirmed this result. In addition, we
describe a new role for TIAL1 in the regulation of translation, as TIAL1 binding substantially
increased mRNA translation (Fig. 5C). So far, TIA1 and TIAL1 have mostly been described as
regulators of pre-mRNA splicing and as translational repressors in the context of cellular stress,
where they assemble into stress granules. [45,46] However, in the absence of stress, TIA1/L1 has
been reported to promote polysome association which supports our findings. [36,47] For both
reporter mRNAs and endogenous mRNAs, we observed that the presence of TIAL1 increased
protein expression in all compartments, but only in the context of the ER did we observe a
cooperative effect on translation (Fig. 5M, 5N). The factor that cooperates with TIAL1 on the ER
to upregulate translation is currently unknown. Importantly, our reporter results demonstrate that
a change in the location of protein synthesis within the cytoplasm strongly influences protein
output, indicating that a change in mRNA localization can alter protein abundance.

A combinatorial code of mRNA architecture features and 3′UTR-bound RBPs controls
subcytoplasmic mRNA transcript distribution

RBPs play an established role in mRNA localization. [1,7] Additionally, we observed a strong
association of mRNA architecture features with mRNA transcript localization to the three
cytoplasmic compartments (Fig. 3C). It is most likely that mRNA length, CDS length, and CDS
exon length are not direct regulators of mRNA localization but that specific factors, including
currently unknown RBPs, read-out this information. In addition, we speculate that mRNA
architecture influences mRNP size, conformation, and packaging [48,49] and that these biophysical
features may act as additional determinants of subcytoplasmic mRNA localization. This idea is
supported by previous insights into *oskar* mRNA localization, where the deposition of the exon
junction complex, which is involved in mRNP packaging, [48,49] was found to be required for
proper mRNA localization in the cytoplasm. [50]

We present a model for the regulation of subcytoplasmic transcript distribution that is based on
a combinatorial code generated by mRNA architecture features together with the bound RBPs,
where individual components act in an additive manner (Fig. 3E, 3G). This model was tested
experimentally by analyzing the localization propensity of TG+ mRNAs upon deletion of TIS11B.
The obtained results confirmed the contribution of mRNA architectural features to mRNA
localization and suggest that the binding of RBPs can overcome the default localization pattern
established by the mRNA architecture features (Fig. 4A-E).

Is it biologically relevant if only 20% of transcripts localize to TGs?

Based on the estimated size of TGs (Fig. 1I), we expect that 11% of mRNA transcripts localize to TGs by chance. Using smRNA-FISH on three individual TG+ mRNAs we observed a two-fold enrichment in TGs, meaning that on average 22% of these transcripts localize to TGs. This raises the important question: whether it matters biologically that a minority population of transcripts for a given mRNA localizes to a certain compartment.

This question was addressed in a follow-up project, where we investigated the biological consequences of *MYC* mRNA, which is a TG+ mRNA, when it was translated in TGs or the cytosol. [19] We observed that several MYC protein complexes were only formed when *MYC* mRNA was translated in TGs and not when it was translated in the cytosol. The TG-dependent protein complexes formed co-translationally and had functional consequences for MYC target gene expression in the nucleus. TG-translated MYC induced different target genes than cytosol-translated MYC. [19] Our results indicate biological relevance, even when only a fraction of transcripts are translated in TGs.

In summary, our study revealed a surprisingly high degree of cytoplasmic compartmentalization. This is the basis for the translation of functionally related proteins in defined environments that strongly affect mRNA and protein expression. Our results highlight the contribution of spatial regulation whose consequences go beyond the effects mediated by the mRNA-bound proteins. In the future, our findings may provide the basis for biotechnology applications that make use of engineered 3′UTR sequences to boost protein expression in experimental settings or to increase protein production of mRNA vaccines.

Limitations of our study

The exact compartment sizes of TGs, the rough ER, and the cytosol are currently unknown and can only be estimated. However, using two different methods for the identification of compartment-enriched mRNAs yielded highly similar enrichment values.

To obtain sufficient material for TG and ER particle sorting, we used transfected, fluorescently labeled proteins instead of endogenous proteins. In cells that highly express TIS11B, in the future, TG particle sorting may be possible using endogenous, fluorescently tagged TIS11B.

The use of spike-ins to isolated compartments obtained from defined cell numbers may have enabled us to perform absolute enrichments versus the relative enrichment analyses that we report here. Moreover, all analyses were performed at the gene level. Alternative 3′UTR isoforms are known to differentially localize and therefore, we would expect to obtain a higher resolution for compartment enrichment of transcripts if instead of genes alternative 3′UTR isoforms are analyzed. [38,51] However, with our purification strategy we did not obtain sufficient mRNA quantities to perform the analysis at the level of alternative 3′UTR isoforms.

Acknowledgements

This work was funded by the NIH Director's Pioneer Award (DP1-GM123454), the R35GM144046 NIH grant, a grant from the Pershing Square Sohn Cancer Research Alliance, a grant from the G. Harold and Leila Y. Mathers Foundation, and the MSK Core Grant (P30 CA008748) to CM. J.U. received funding from the ERC under the European Union Horizon 2020 Research and Innovation Program (835300-RNPdynamics) and The Francis Crick Institute

receives its core funding from Cancer Research UK (CC0102), the UK Medical Research
Council (CC0102), and the Wellcome Trust (CC0102). An NIH F31 (CA254335-01) fellowship
was awarded to E.L.H.

We thank Yevgeniy Romin and Eric Chan from the Molecular Cytology Core Facility for help
with microscopy and image analysis. We thank Kathleen Daniels from the Flow Cytometry Core
Facility (now Sana Biotechnologies) for help with particle sorting and Liana Boraas at Yale
University School of Medicine for help with endogenous RNA-FISH. We thank Weirui Ma
(Zhejiang University) for experimental guidance and mentorship of E.L.M. and Alban Ordureau
for analysis of the TMT mass spectrometry data. We thank Yang Luo (MSKCC) and Ben
Kleaveland (WCMC) for helpful discussions and critical reading of the manuscript.

Author contributions

E.L.H. performed all experiments, except the mass spectrometry analysis whose samples were
prepared by X.C. and the TIS11B KO cells were generated by S.M. F.C.Y.L. and J.U. performed
and analyzed the TIS11B iCLIP experiment. M.M.F performed the logistic regression and
provided the gene architecture features. T.C. analyzed the RNA-seq data. G.Z. analyzed the
CLIP data with input from C.M. E.L.H. and C.M. conceived the project, designed the
experiments, and wrote the manuscript with input from all authors.

Declaration of Interests

The authors declare no competing interests.

Methods

STAR methods table

Cell lines

Generation of a doxycycline inducible *TIS11B* knockout cell line (*TIS11B* KO)

Doxycycline inducible Cas9 (iCas9) HEK293T cells were generated by infecting cells with
lentivirus containing a Cas9-P2A-GFP expression cassette under a doxycycline inducible
promoter as described previously. [52] During consecutive rounds of fluorescence-activated cell
sorting, we selected a cell pool exhibiting robust induction of Cas9/GFP expression after
doxycycline treatment (100 ng/ml for 24 hours), and low levels of leaky transgene expression in
the absence of the drug.

Next, we transduced iCas9 cell lines with a lentiviral construct harboring a pair of guide RNAs
either targeting *TIS11B* or gRNAs that target an intergenic region. To generate these constructs,
we adapted the plentiGuide-puro vector. [53] to incorporate a second guide RNA expression
cassette as described previously. [54] For this purpose, the plasmid was digested with BsmBI
(FastDigest Esp3I, Thermo Fisher Scientific) and a synthetic 391 bp double-stranded DNA
fragment encoding 5′-(1st gRNA/scaffold/H1 promoter/2nd gRNA)-3′ was inserted using the
NEBuilder HiFi assembly system (NEB). Synthetic DNA fragments were ordered from Genewiz
and sequences are listed in Table S6. The assembled vector DNA was used to transform
chemically competent Stbl3 bacteria cells (Invitrogen), and correct vector clones were identified
by Sanger sequencing.

Lentivirus was generated in HEK293T cells using standard methods and 200 μl of viral
supernatant was used to transduce iCas9 cells in a 6-well dish together with 8 μg/ml polybrene.
Transduced cells were subjected to puromycin selection (1 μg/ml) for five days and resistant
cells were aliquoted and frozen for all further experiments. Finally, for induction of gene
knockouts, *TIS11B* KO and corresponding control cells (with gRNAs targeting an intergenic
region) were treated with doxycycline (100 ng/ml) for five days, after which TIS11B protein
expression was evaluated by western blotting, ER particle sorting and digitonin extraction was
performed.

Detailed methods

Constructs

Fluorescently-tagged TIS11B and SEC61B constructs. The eGFP/mCherry/BFP fusion
constructs for TIS11B and SEC61B expression were described previously. [16] They were
generated in the pcDNA3.1-puro expression vector. The TIS11B and SEC61B coding regions
were PCR amplified from HeLa cDNA and inserted downstream of eGFP/mCherry/BFP using
BsrGI/EcoRI or BsrGI/HindIII restriction sites, respectively.

Constructs to generate the mRNA localization reporter. To investigate the influence of RBPs on
mRNA localization of a GFP mRNA reporter, RBPs were fused to MCP and tethered to a GFP
mRNA reporter containing MS2 binding sites as 3′UTR. [37,38] To investigate mRNA localization-
dependent protein expression of the GFP mRNA reporter, a CAAX sequence was fused to MCP
or to MCP-RBP fusions.

GFP mRNA reporter. To generate the GFP mRNA reporter, the GFP-BIRC3-MS2-SU [39] vector
was used the BIRC3 coding region was replaced with the THAP1 coding region. It was PCR
amplified from the GFP-THAP1 vector using THAP1-MS2 F and THAP1-MS2 R primers and

inserted between the BsrGI and AgeI sites. The SU fragment was removed with HindIII and
XhoI and blunt end ligated, resulting in GFP-THAP1-MS2.

MCP-mCherry RBP fusion constructs. To generate MCP-mCherry, the MCP coding sequence
was PCR amplified from UBC NLS-HA-2XMCP-tagRFPt vector (Addgene 64541) using MCP F
and MCP R primers and inserted in-frame, upstream of mCherry (mCherry lacking a start
codon) between BmtI and BamHI sites in pcDNA3.1-puro-mCherry vector. [16] To generate MCP-
mCherry-TIS11B and MCP-mCherry-TIAL1, their coding sequences were inserted in-frame,
downstream of mCherry between the BsrGI and XbaI sites. The TIS11B coding sequence was
amplified from pcDNA3.1-puro-GFP-TIS11B using TIS11B MCP F and TIS11B MCP R primers
and the TIAL1 coding sequence was PCR amplified from pFRT_TO_FlagHA_TIAL1 (Addgene
106090) using TIAL1 MCP F and TIAL1 MCP R primers.

MCP-mCherry fusion constructs with subcellular localization signals. To generate pcDNA3.1-
puro-MCP-mCherry-SEC61B, the MCP-mCherry coding sequence was cut from MCP-mCherry
vector using BmtI and BsrGI and pasted in-frame, upstream of SEC61B in pcDNA3.1-mCherry-
SEC61B (replacing mCherry). To generate the TIS11B-MCP-mCherry-SEC61B vector, TIS11B
coding sequence was PCR amplified from pcDNA3.1-puro-GFP-TIS11B using TIS-SEC F and
TIS-SEC R primers and pasted in-frame, upstream of MCP into the BmtI site in the MCP-
mCherry-SEC61B vector. To generate TRAPα-MCP-mCherry, the TRAPα coding sequence
(encoded by the *SSR1* gene) was PCR amplified from HeLa cDNA using TRAPα MCP F and
TRAPa MCP R and inserted in-frame, upstream of MCP in the pcDNA3.1-puro-MCP-mCherry
vector.

For plasma membrane localization, the CAAX prenylation signal was added to the C-terminus of
MCP-mCherry or MCP-mCherry-TIAL1. The CAAX coding sequence was purchased as a gene
fragment from Azenta as described [32] and PCR amplified using TIAL1 CAAX F and CAAX R
primers. It was inserted in-frame using the BsrGI and ApaI sites, located downstream of
mCherry to generate pcDNA3.1-puro-MCP-mCherry-CAAX. It was inserted in-frame using
EcoNI and ApaI sites to generate MCP-mCherry-TIAL1-CAAX.

SunTag constructs were described previously. [32]

Isolation of subcytoplasmic compartments

Transfection. HEK293T cells were seeded in six 10 cm dishes (particle sorting) or one well from
a 6-well plate (cytosol extraction) at 80% confluency in antibiotic free media. After 24 hours,
cells were transfected by calcium phosphate with either 3 µg mCherry-TIS11B or 1 µg GFP-
SEC61B per dish (particle sorting), or 500ng mCherry-TIS11B (cytosol extraction).

Particle purification. 20 hours after transfection, cells were rinsed once with ice-cold PBS,
scraped in 10 ml ice-cold PBS, and pelleted at 300 x g. Pellets from two plates were
resuspended in 1 ml ice-cold hypotonic isolation buffer (225 mM mannitol, 75 mM sucrose, 20
mM Tris-HCl pH 7.4, 0.1 mM EDTA). Cells were lysed with 50 strokes in a 1 ml dounce-
homogenizer with pestle on ice in order to shear the nuclei from the ER. Nuclei were pelleted
with a two-minute spin at 600 x g. The supernatant contains the cytoplasmic membrane fraction,
which was pelleted with a 15-minute spin at 7000 x g and resuspended in ice-cold PBS for
fluorescent particle sorting.

Fluorescent particle sorting. Particles were sorted on a BD FACSAria III cell sorter equipped
with a 70 µm nozzle. The forward-scatter threshold was decreased from 5,000 to 800 in order to
visualize subcellular particles. Particles were first detected by fluorescence using the 594 nm
and 488 nm excitation lasers, for mCherry-TIS11B and GFP-SEC61B respectively, and 405 nm
excitation laser for DAPI. A sorting gate was drawn on particles that were either mCherry-
positive or GFP-positive, but DAPI-negative, to exclude any remaining nuclei. Sorting was

performed in purity mode with an average speed of 150 particles/second. Particles were sorted
directly into 1 ml of TRIzol solution in Eppendorf tubes, holding 180,000 particles per tube. RNA
extraction was performed for each tube separately and total RNA for each sample was
combined for library preparation. Two biological replicates for each particle prep were
sequenced. For each replicate, about 1.5 million TIS11B granule particles and 2.0 million ER
particles were collected.

Cytosol extraction. The cytosol was extracted as previously described. [24] HEK293T cells
transfected were plated in a six-well plate at 80% confluency. After 24 hours, cells were rinsed
once in the dish with ice-cold PBS. After aspirating PBS, 300 µl ice-cold digitonin solution (40
µg/ml digitonin, 150 mM NaCl, 20 mM HEPES pH 7.4, 0.2 mM EDTA, 2 mM DTT, 2 mM MgCl2)
was added and incubated on a shaker at 4°C for ten minutes. After incubation, the digitonin-
derived cytosolic extract was pipetted from the plate and spun at 20,000 x g for one minute to
pellet any floating cells. 200 µl of cytosolic extract was added to 1 ml TRIzol solution for RNA
extraction.

RNA-seq library preparation

RiboGreen RNA Reagent (ThermoFisher) was used for RNA quantification and quality control
was performed by Agilent BioAnalyzer. 50-500 ng of total RNA underwent polyA selection and
TruSeq library preparation according to instructions provided by Illumina (TruSeq Stranded
mRNA LT Kit, catalog # RS-122-2102), with eight cycles of PCR. Samples were barcoded and
run on a HiSeq 4000 in a PE50 run, using the HiSeq 3000/4000 SBS Kit (Illumina). An average
of 27 million paired reads was generated per sample.

Western Blotting

For whole cell lysate preparation, cells were trypsinized and washed twice with PBS and lysed
in 2x Laemmli Sample buffer (Alfa Aesar, J61337). For cytosolic lysate, cytosol was extracted
with digitonin as described above and one volume of 2x Laemmli Sample buffer was added.
Laemmli lysates were boiled for 10 min at 95°C. Samples were subjected to SDS-PAGE on
NuPAGE 4%–12% Bis-Tris gradient protein gel (Invitrogen). Imaging was captured on the
Odyssey DLx imaging system (Li-Cor). Quantification was performed using ImageJ. The
antibodies used are listed in the Key Resources Table.

TIS11B iCLIP

Transfection. HEK293T cells were seeded in 10 cm dishes at 80% confluency in antibiotic free
media. After 24 hours, cells were transfected by calcium phosphate with either 3 µg GFP-
TIS11B or 1.5 µg GFP-only per dish.

Sample preparation. 20 hours after transfection, cells were rinsed once with ice-cold PBS and 6
ml of fresh PBS was added to each plate before crosslinking. Cells were irradiated once with
150 mJ/cm^2 in a Spectroline UV Crosslinker at 254 nm. Irradiated cells were scraped into
Eppendorf tubes, spun at 500 x g for one minute, and snap-frozen. Crosslinked cell pellets were
lysed in iCLIP lysis buffer (50 mM Tris-HCl pH 7.4, 100 mM NaCl, 1% Igepal CA-630 (Sigma
I8896), 0.1% SDS, 0.5% sodium deoxycholate), sonicated with the Bioruptor Pico for 10 cycles
30 seconds ON/30 seconds OFF, and supplemented with 0.5 U of RNase I per 1 mg/ml lysate
for RNA fragmentation. Lysates were pre-cleared by centrifugation at 20,000 x g at 4°C. A mix
of Protein A/G Dynabeads (50 µl of each per sample, Life Technologies) were coupled to 10 µg
of rabbit anti-GFP antibody (Abcam ab290). TIS11B protein-RNA complexes were
immunoprecipitated from 1 ml of crosslinked lysate and washed with high salt and PNK buffer
(NEB). RNA was repaired by 3′ dephosphorylation and ligated to L3-IR adaptor on beads. [55]
Excess adaptor was removed by incubation with 5′ deadenylase and the exonuclease RecJf
(NEB). TIS11B protein-RNA complexes were eluted from the beads by heating at 70°C for one

minute. The complexes were then visualized via the infrared-labeled adaptor, purified with SDS-
PAGE, and transferred to nitrocellulose membrane. cDNA was synthesized with Superscript IV
Reverse Transcriptase (Life Technologies) and circularized by CircLigase II. Circularized cDNA
was purified with AmPURE bead-based purification (A63880, Beckman Coulter), amplified by
PCR and sequenced by Novaseq.

RNA-FISH

Single molecule RNA-FISH for endogenous mRNAs. Probe design. Primary probes were
designed using the ProbeDealer package in MATLAB. [56] Each primary probe contains 30
transcript-targeting nucleotides preceded by 20 common nucleotides that are complementary to
the secondary probe. At least 30 probes were designed for each transcript, purchased in a pool
from IDT. The secondary probes are 5′ conjugated to AlexaFluor 633 and were purchased from
IDT.

Transfection. Prior to cell seeding, 35 mm glass cover slips were sterilized with ethanol then
incubated in 1 µg/ml fibronectin in PBS at room temperature for one hour. Cover slips were
rinsed in PBS and HeLa cells were seeded at 100,000 per coverslip. 24 hours after seeding,
cells were co-transfected with 250 ng BFP-TIS11B and 100ng of GFP-SEC61B using
Lipofectamine 3000 (Invitrogen).

Sample preparation. 20 hours after transfection, cells were rinsed once with PBS then fixed in
4% paraformaldehyde for 10 minutes at room temperature. All steps were performed at room
temperature if not otherwise noted. Cells were rinsed twice with PBS and permeabilized with
0.5% Triton-X solution for 10 minutes. Cells were rinsed twice with PBS and incubated for five
minutes in pre-hybridization buffer (2xSSC, 50% formamide). Cells were incubated in primary
probe hybridization solution (40 µM primary probe, 2xSSC, 50% formamide, 10% dextran
sulfate (Sigma), 200 µg/ml yeast tRNA (Sigma), 1:100 Murine RNase Inhibitor (NEB)), for at
least 15 hours at 37°C. To remove excess or unbound primary probes, cells were then rinsed
twice in 2xSSC + 0.1% Tween for 15 minutes at 60°C then once more for 15 minutes at room
temperature. Cells were incubated in secondary probe solution (4 nM secondary probe, 2xSSC,
50% ethylene carbonate, 1:100 Murine RNase Inhibitor) for 30 minutes in the dark. Secondary
probes were rinsed twice in 50% ethylene carbonate, 2xSSC solution for five minutes then
mounted with Prolong Diamond mounting solution (Invitrogen).

Cytosol extraction. To visualize and validate CY+ versus TG+ or ER+ endogenous mRNAs,
HeLa cells were seeded as described above, then incubated in 2 ml digitonin solution described
above (40 µg/ml digitonin, 150 mM NaCl, 20 mM HEPES pH 7.4, 0.2 mM EDTA, 2 mM DTT, 2
mM MgCl2) for 10 min at 4°C. Digitonin solution was removed, coverslips were rinsed with 2 ml
PBS, and RNA-FISH was performed as described above. Mounting media with DAPI was used
to visualize nuclei (Invitrogen P36931).

Validation of TG+ and ER+ mRNAs using smRNA-FISH. We performed smRNA-FISH on
endogenous mRNAs (Table S2) while simultaneously visualizing TGs and the ER. We
considered an mRNA to have an unbiased localization pattern if its transcript distribution
correlated with the cytoplasmic compartment sizes. As a proxy for the relative compartment
sizes, we used the area occupied by TGs or the ER compared to the whole cell area, obtained
from the maximum projection of the fluorescent signals in 186 cells. We used FIJI to delineate
the whole cell border with the fluorescent signal from RNA-FISH. For TGs, the fluorescent signal
from BFP-TIS11B and for the ER the fluorescent signal from GFP-SEC61B both obtained from
the maximum intensity Z-projections was used to delineate each compartment. Where there
was overlap between the TG mask and the ER mask, the ER was subtracted, and the region
was defined as TG. In this way the compartments are mutually exclusive. The mask area of
each compartment was quantified and read out as a proportion of the total cell area. Across all

cells, the median size of TGs was estimated to be 11% of the cell size, whereas the median ER
size was estimated to be 29% of the cell size (Fig 1I, 1J). Therefore, for mRNAs with an
unbiased transcript distribution, we expect that typically 11% of transcripts colocalize with TGs
and 29% colocalize with the ER.

To determine mRNA transcripts enriched in TG or ER, smRNA-FISH foci were counted using
the maxima function and the total number of foci per cell are quantified. Next, all foci are
overlaid with the TG mask and the ER mask to identify mRNAs that colocalize with each
compartment. To determine if an mRNA is compartment enriched, we tested if its observed
compartment distribution differs from the expected distribution based on compartment size
using a Mann Whitney test. The code for the image analysis is available (see below).

Of note, this analysis does not distinguish between nuclear and cytoplasmic mRNA localization.
For 7/8 mRNAs this does not influence the outcome because the mRNA signal in the nucleus is
negligible or non-existent. However, smRNA-FISH probes for endogenous *TES* produce high
nuclear background signal. In this case, the prominence value, used to define local maxima to
call foci, is increased such that nuclear noise does not substantially influence foci quantification
(Fig. S2G).

<u>Validation of CY+ mRNAs by smRNA-FISH after digitonin extraction.</u> To distinguish CY+
mRNAs from TG+ or ER+ mRNAs, we performed smRNA-FISH on endogenous mRNAs in
untreated and digitonin treated cells, as previously reported. [57] The total number of mRNA foci
per cell is calculated using the maxima function in FIJI. Next, thresholding is applied to DAPI
fluorescence to generate a nuclear mask. Total mRNA foci are overlaid with the DAPI mask and
nuclear foci are subtracted from the total, yielding cytoplasmic foci. Cytoplasmic foci are
quantified for at least 10 cells per condition per experiment. For each experiment, the mean
fraction of transcripts retained is calculated as the average cytoplasmic foci per digitonin-treated
cell divided by the average cytoplasmic foci per untreated cell. At least three separate
experiments per mRNA were performed.

<u>RNA-FISH after transfection of constructs.</u> RNA-FISH experiments probing for GFP-fusion
constructs were performed as described previously. [16] Stellaris FISH probes for eGFP with
Quasar 670 Dye were used.

<u>Line profile analysis.</u> To quantify colocalization of ER (GFP-SEC61B) and mRNA (AF633)
fluorescence signals, line profiles were generated with FIJI (ImageJ). For each cell, 2-4 straight
lines were drawn to cross the ER in different directions, indicated by the white arrows shown in
the figures. Fluorescence signal along the straight line of the ER and the mRNA reporter was
calculated for each channel using the plot profile tool in FIJI. The values of the Pearson's
correlation coefficient r were calculated using Excel. Perfect correlation of protein-mRNA is
indicated by $r = 1$, perfect exclusion is indicated by $r = -1$, and random distribution is indicated
by $r = 0$.

Confocal microscopy

Confocal imaging was performed using ZEISS LSM 880 with Airyscan super-resolution mode or
Nikon CSU-W1 with SoRa super-resolution mode. A Plan-Apochromat 63x/1.4 (Zeiss) or
60x/1.49 (Nikon) Oil objective was used. For live cell imaging, cells were incubated with a
LiveCell imaging chamber (Zeiss, Nikon) at 37°C and 5% CO2 and imaged in cell culture media.
Excitations were performed sequentially using 405, 488, 594 or 633 nm laser wavelength and
imaging conditions were experimentally optimized to minimize bleed-through. Z-stack images
were captured with the interval size of 0.2 μm. Images were prepared with FIJI (ImageJ)
software.

TMT mass spectrometry

To obtain protein expression levels, TMT mass spectrometry analysis was performed on
HEK293T cells cultivated in steady-state conditions. Cells were trypsinized and washed three
times with ice-cold PBS. Pelleted cells were snap-frozen in liquid nitrogen. Cell pellets were
lysed with 200 µl buffer containing 8 M urea and 200 mM EPPS (pH at 8.5) with protease
inhibitor (Roche) and phosphatase inhibitor cocktails 2 and 3 (Sigma). Benzonase (Millipore)
was added to a concentration of 50 µg/ml and incubated at room temperature for 15 min
followed by water bath sonication. Samples were centrifuged at 14,000 g at 4°C for 10 min, and
supernatant extracted. The Pierce bicinchoninic acid (BCA) protein concentration assay was
used to determine protein concentration. Protein disulfide bonds were reduced with 5 mM tris
(2-carboxyethyl) phosphine at room temperature for 15 min, and alkylated with 10 mM
iodoacetamide at room temperature for 30 min in the dark. The reaction was quenched with 10
mM dithiothreitol at room temperature for 15 min. Aliquots of 100 µg were taken for each sample
and diluted to 100 µl with lysis buffer. Samples were subject to chloroform/methanol
precipitation as previously described. [58] Pellets were reconstituted in 200 mM EPPS buffer and
digested with Lys-C (1:50 enzyme-to-protein ratio) and trypsin (1:50 enzyme-to-protein ratio),
and digested at 37°C overnight.

Peptides were TMT-labeled as described. [58] Briefly, peptides were TMT-tagged by the addition
of anhydrous ACN and TMTPro reagents (16plex) for each respective sample and incubated for
1 hour at room temperature. A ratio check was performed by taking a 1 µl aliquot from each
sample and desalted by StageTip method [59]. TMT tags were then quenched with hydroxylamine
to a final concentration of 0.3% for 15 min at room temperature. Samples were pooled 1:1
based on the ratio check and vacuum-centrifuged to dryness. Dried peptides were reconstituted
in 1 ml of 3% ACN/1% TFA, desalted using a 100 mg tC18 SepPak (Waters), and vacuum-
centrifuged overnight.

Peptides were centrifuged to dryness and reconstituted in 1 ml of 1% ACN/25mM ABC.
Peptides were fractionated into 48 fractions. Briefly, an Ultimate 3000 HPLC (Dionex) coupled to
an Ultimate 3000 Fraction Collector using a Waters XBridge BEH130 C18 column (3.5 um 4.6 x
250 mm) was operated at 1 ml/min. Buffer A, B, and C consisted of 100% water, 100% ACN,
and 25mM ABC, respectively. The fractionation gradient operated as follows: 1% B to 5% B in
1 min, 5% B to 35% B in 61 min, 35% B to 60% B in 5 min, 60% B to 70% B in 3 min, 70% B to
1% B in 10 min, with 10% C the entire gradient to maintain pH. The 48 fractions were then
concatenated to 12 fractions, (i.e. fractions 1, 13, 25, 37 were pooled, followed by fractions 2,
14, 26, 38, etc.) so that every 12th fraction was used to pool. Pooled fractions were vacuum-
centrifuged and then reconstituted in 1% ACN/0.1% FA for LC-MS/MS.

Fractions were analyzed by LC-MS/MS using a NanoAcquity (Waters) with a 50 cm (inner
diameter 75 µm) EASY-Spray Column (PepMap RSLC, C18, 2 µm, 100 Å) heated to 60°C
coupled to an Orbitrap Eclipse Tribrid Mass Spectrometer (Thermo Fisher Scientific). Peptides
were separated by direct injection at a flow rate of 300 nl/min using a gradient of 5 to 30%
acetonitrile (0.1% FA) in water (0.1% FA) over 3 hours and then to 50% ACN in 30 min and
analyzed by SPS-MS3. MS1 scans were acquired over a range of m/z 375-1500, 120K
resolution, AGC target (standard), and maximum IT of 50 ms. MS2 scans were acquired on
MS1 scans of charge 2-7 using isolation of 0.5 m/z, collision-induced dissociation with activation
of 32%, turbo scan, and max IT of 120 ms. MS3 scans were acquired using specific precursor
selection (SPS) of 10 isolation notches, m/z range 110-1000, 50K resolution, AGC target
(custom, 200%), HCD activation of 65%, max IT of 150 ms, and dynamic exclusion of 60 s.

Visualization of translation in TGs

The SunTag system was used to visualize mRNA translation in the cytosol and the TGER
domain. Stable expression of td-PP7-3xmCherry (Addgene 74926) and scFv-GCN4-sfGFP
(Addgene 60907) was achieved by generating virus in HEK293T cells and transducing HeLa
cells. Cells were seeded on 3.5 cm glass bottom dishes (Cellvis, D35-20-1-N). 20 hours later,
cells were transfected with either the SunTag vector expressing KIF18B (Addgene 74928) or
SunTag-FOS-UTR. At 15 hours post transfection, cells were treated with 100 ng/ml doxycycline
for one hour to induce SunTag expression. Confocal imaging was performed as described
above. Colocalization of foci was quantified using FIJI.

mRNA localization-dependent GFP protein expression

Transfection. HeLa cells were seeded in 12-well plates at 80% confluency and transfected with
250 ng GFP-THAP1-MS2 and 250 ng of the MCP-mCherry fusion constructs indicated in the
figure (Lipofectamine 3000, Invitrogen). When indicated, GFP-THAP1 or GFP-BIRC3-MS2-SU
was used instead of GFP-THAP1-MS2. At 13-15 hours post transfection, cells were analyzed by
FACS. For RNA-FISH experiments, cells were seeded at 80% confluency in 4-well slide
chambers (Millipore Sigma) and cotransfected with 75 ng GFP-THAP1-MS2, 100 ng BFP-
SEC61B, and 75 ng of the indicated MCP-mCherry fusion constructs.

FACS analysis to measure GFP protein expression. Cells were trypsinized, washed once in
complete media, then resuspended in FACS buffer (PBS plus 1% FCS). At least 5,000 cells
were measured on a BD LSR-Fortessa Cell Analyzer and FACS data were analyzed using
FlowJo software. GFP protein expression corresponds to GFP mean fluorescence intensity
(MFI). To determine the effect of MCP-tethered RBPs on protein output of the GFP reporter
mRNA, only cells that were successfully cotransfected with both the MCP-mCherry fusion and
the GFP reporter constructs were analyzed. To do so, the double-positive cells
(mCherry+/GFP+) were gated, and all single positive and unstained cells were excluded from
the analysis. The reported GFP MFI was calculated from the double-positive cells.
Untransfected cells were used to draw the gates for mCherry+ or GFP+ cells.

qPCR analysis to measure *GFP* mRNA abundance. Cells were trypsinized, washed once in
complete media, then resuspended in FACS buffer (PBS plus 1% FCS). To determine the effect
of MCP-tethered RBPs on GFP reporter mRNA stability, cells were sorted based on expression
of both the MCP-mCherry fusion and the GFP reporter constructs. The BD FACSAria III cell
sorter was used to collect 50,000 cells from each co-transfected population. Cells were sorted
directly into 1 ml of TRIzol solution in Eppendorf tubes for total RNA was extraction. cDNA
synthesis was performed on 200 ng of RNA per sample using the SuperScript IV VILO ezDNase
Master Mix (Invitrogen). ezDNase enzyme was included to eliminate plasmid DNA
contamination. To measure the relative expression levels of reporter mRNA by qRT-PCR,
FastStart Universal SYBR Green Master Mix (ROX) from Roche was used together with GFP-
qPCR F/R primers. GAPDH was used as a housekeeping gene.

Data analysis

RNA-seq of subcytoplasmic fractions from HEK293T cells

RNA-seq. Alignment was generated in Dragen v3.10 (Illumina) against the hg38-alt-masked-v2
reference acquired from GENCODE v43 with default parameters. Gene expression analysis
was performed using HOMER v4.11 software. [60] The mean RPKM values of all biological
replicates were calculated and used for downstream analyses. Only protein-coding genes were
analyzed. A gene was considered expressed if the RPKM value is 3 or greater.

Classification of membrane/secretory proteins versus non-membrane proteins. Information on
the presence of transmembrane domains or a signal sequence was obtained from uniprot. All
expressed genes were separated into mRNAs that encode membrane/secretory proteins or
non-membrane proteins. If a protein contains a signal sequence but not a transmembrane
domain, it is considered as secretory protein. All proteins with transmembrane domains are
considered membrane proteins and all remaining proteins are classified as non-membrane
proteins. Among the 9155 mRNAs expressed in HEK293T cells, 2140 were classified as
membrane/secretory proteins, whereas 7015 were classified as non-membrane proteins (Table
S1).

Compartment-specific localization scores. The sum of RPKM values obtained from TG particles,
ER particles, and the cytosol was considered as total cytoplasmic mRNA expression. For each
gene, the mean compartment-specific RPKM value was divided by the total cytoplasmic mRNA
expression. As a result, each gene is assigned three localization scores that correspond to the
fraction of its transcripts that localize to each of the three compartments: TGs, the ER, and the
cytosol.

Compartment-specific enrichment of mRNAs that encode membrane/secretory proteins. We
considered an mRNA to be ER-enriched if the ratio of localization scores (ER/TG) was greater
than 1.25 and classified it as TG-enriched if it was smaller than 0.8. The median localization
score of membrane/secretory mRNAs in the cytosol was 0.09. If the cytosolic localization score
of an mRNA was greater than 0.36, it was considered enriched in the cytosol. If the ER and TG-
specific localization scores were similar and the cytosolic partition coefficient was smaller than
0.18, the mRNA was assigned to the ER, whereas it was considered not localized if the
cytosolic localization score was smaller than 0.18 (Fig. S1H).

Compartment-specific enrichment of mRNAs that encode non-membrane proteins. To faithfully
compare differences in mRNA distribution across the three compartments, it is necessary to
know the relative size distribution of the three compartments. However, this parameter is
currently unknown. Therefore, instead of comparing the localization scores across samples, we
determined the most enriched mRNAs within each compartment. We considered an mRNA
compartment-enriched, if its average localization score (from biological replicates) was at least
1.25-fold higher than the median localization score of its corresponding compartment samples.
For TG particles, the median localization score was 0.32, for ER particles, it was 0.30, and for
the cytosol, the median localization score was 0.34. If the enrichment was observed in two
compartments, the mRNA was assigned to the compartment with the higher value. With this
strategy, we identified 1246 TG+ mRNAs, 919 non-overlapping ER+ mRNAs, and 1481 CY+
mRNAs. The remaining 3369 mRNAs (48%) do not have a compartment-biased mRNA
localization pattern and were called (unbiased).

Justification of the cut-off used to determine compartment-enriched mRNAs. A minimum cut-off
of 1.25-fold higher than the median localization score corresponds to approximately one
standard deviation. The compartment-enriched mRNAs differed substantially in their functional
and architectural features (Fig. 2). We generated subgroups among the compartment-enriched
mRNAs that represent the top, middle, and bottom-enriched subgroups (Fig. S4). Even when
focusing on the bottom-enriched groups (which are close to the cut-off used), the differences in
functional and architectural features across the compartment-enriched groups were still highly
significant (Fig. S4). The cut-off is further justified as we were able to validate 10/11 mRNAs
considered to be compartment enriched with an independent method. Moreover, we
demonstrate that TG-translated MYC has biological effects, despite *MYC* mRNA being found in
the bottom enriched TG+ group. [19]

mRNA transcript distribution in HEK293T TIS11B KO cells

We focused on the analysis of mRNAs that encode non-membrane proteins (Table S5). The
mean RPKM values of the biological replicates of digitonin-extracted samples and the ER
particles were calculated for TIS11B KO cells and their corresponding control HEK293T cells. A
gene was considered expressed if the average RPKM value in the ER and in the cytosol
samples was greater than 3 RPKM (N = 6229). The compartment-specific localization scores
were calculated and the difference in localization scores between TIS11B KO and control
samples were calculated for ER and cytosol. The top 20% of genes with a localization change
towards ER or the cytosol were intersected with genes considered as TG+ (N = 1246) and
further analyzed with respect to their bound RBPs and architectural features.

mRNA and protein features of the localized mRNAs

RPKM values of mRNAs were obtained from RNA-seq data of unfractionated HEK293T cells
and were determined for the compartment-biased mRNAs. Pro-seq and RNA-seq from HEK293
cells were obtained from GEO (GSE140365: PRO-seq; GSE142895: RNA-seq). [27] Raw reads
were processed by trimmomatic (version: 0.39) to trim low-quality ends (average quality per
base < 15, 4 bp window) and adapters. [61] Trimmed reads were mapped to the human genome
(hg19) using hisat2 (version: 2.1.0). [62] Reads mapped to each gene were counted by
featureCounts (version: 1.6.4). [63] To estimate mRNA stability rates, log2-normalized counts of
Pro-seq data were divided by the log2-normalized RNA-seq data, as described previously [28].
3′UTR length of each mRNA was obtained from Ref-seq. The longest 3′UTR isoform of each
gene is reported. mRNA length, CDS length, average CDS exon length, and total exon number
of genes were determined using transcripts from the Matched Annotation from
the NCBI and EMBL-EBI (MANE) [64] human version 1.2. For each gene, the transcript with
longest mRNA length was selected. Protein length was calculated by dividing CDS length by
three.

Proteomics protein expression analysis

Protein expression was obtained from TMT-based quantitative mass spectrometry analysis of
HEK293T cells. Precursor protein abundance was calculated for each protein and scaled to the
TMT abundance for each channel. Relative abundance was then calculated by averaging the
condition-specific biological replicates. In brief, mass spectra were processed using Protein
Discoverer 2.5 (ThermoFisher) using the Minora algorithm (set to default parameters) for
precursor quantification and using a TMTpro workflow for TMT-based quantification. Database
searching included all canonical entries from the human Reference Proteome UniProt database
(SwissProt – 2022-03), as well as an in-house curated list of contaminants. The identification of
proteins was performed using the SEQUEST-HT engine against the database using the
following parameters: a tolerance level of 10 ppm for MS^1 and 0.6 Da for MS^2 post-recalibration
and the false discovery rate of the Percolator decoy database search was set to 1%. Trypsin
was used as the digestion enzyme, two missed cleavages were allowed, and the minimal
peptide length was set to 7 amino acids. Carbamidomethylation of cysteine residues (+57.021
Da) was set as static modifications, while oxidation of methionine residues (+15.995 Da) was
set as a variable modification. The final protein-level FDR was set to 1%. Precursor abundance
quantification was determined based on intensity, and the minimum replicate feature parameter
was set at 50%. Proteins were quantified based on unique and razor peptides and proteins with
less than two different peptides were excluded. For TMT-based quantification, similar search
parameters were used, with the addition of TMTpro tags on lysine residues and peptide N
termini (+304.207 Da) set as static modifications. For TMTpro-based reporter ion quantitation,
the summed signal-to-noise (S:N) ratio for each TMT channel was extracted, and the closest
matching centroid to the expected mass of the TMT reporter ion was found (integration
tolerance of 0.003 Da). PSMs with poor quality, MS^3 spectra with TMT reporter ion channels
missing, or isolation specificity less than 0.7, or with less than 70% of SPS masses matching to

the identified peptides, or with an average TMT reporter summed signal-to-noise ratio that was
less than 10 or had no MS^3 spectra were excluded from quantification. We exported the results
of protein identification and quantification to Excel, including the TMT-based reporter ion
quantitation. Additionally, we extracted the MS^1 precursor abundance for each protein (Minora
algorithm), which indicates its relative abundance in the tryptic sample. Each MS^1-based
abundance measured should be a representation of the sum of all the respective TMT-labeled
peptides combined. Therefore, for a rudimentary metric of protein abundance across samples,
we divided the total MS^1-abundance for individual proteins by their respective TMT summed
signal-to-noise ratio to each TMT channel.

CLIP data analysis

iCLIP analysis of TIS11B in HEK293T cells. Raw fastq files were demultiplexed using the iCount
python package (https://icount.readthedocs.io). 5′ and 3′ adapters were trimmed by Cutadapt. [65]
Trimmed reads were mapped to human genome using STAR and reads mapping to tRNA/rRNA
were discarded. [66] Crosslink sites were called from bam files using the "xlsites" function of
iCount. CLIP-seq analysis was carried out on the iMaps platform
(https://imaps.genialis.com/iclip), where peak calling was performed by analysing cDNA counts
at crosslink sites using Paraclu. [67] Motif analysis was carried out using HOMER software.
Enrichment was calculated within the genomic coordinates of a total of 57,714 TIS11B CLIP
peaks found in 3′UTRs. Total peaks: 190,920; peaks in 3′UTRs: 57,714.

POSTAR3 CLIP data. CLIP data on 168 RBPs were downloaded from Postar3 [34] and peak
counts that overlapped with annotated 3′UTRs from Ref-seq in all mRNAs that encode non-
membrane proteins were recorded. For each RBP, the median number of 3′UTRs CLIP peaks
was calculated and all 3′UTRs with peaks counts greater than the median were considered as
targets. Based on the fraction of mRNAs that are considered compartment-specific (TG: 17.8%;
ER 13.1%; CY: 21.1%; unbiased: 48.0%), we determined the expected number of target genes
for each compartment. If the observed number of targets divided by the expected number of
targets in a compartment was greater than 1.5, the RBP was added to our short-list (Table S4).
As TIS11B and TIA1/L1 are known to bind to AU-rich sequences, we added the processed
PAR-CLIP data of the LARP4B RBP as it was reported to bind to AU-rich elements. [33]

Logistic regression. The R package `nnet` (v7.3-17) was used to fit logistic regression models to
predict the subcytoplasmic mRNA localization of non-membrane proteins. An initial model used
CLIP peak counts from the RBPs on the short list (N = 24). A second model used the top seven
RBPs from the first model fit and added mRNA length and average CDS exon length.
Covariates with missing values were imputed as zeros. All covariates were first `sqrt`
transformed and then standardized. The 'unbiased' category was used as the base level. The R
package `broom` (v0.8.0) was used to compute t-test statistics for the model coefficients. The
code is available on github (github.com/Mayrlab/tiger-seq).

Confirmation of the logistic regression. To validate the contribution of each individual RBP, we
used more stringent criteria to determine their targets. Among all mRNAs that encode non-
membrane proteins with at least one CLIP peak in the 3′UTR, we considered the top third of
mRNAs as targets of each RBP (TIS11B: 1781 targets; TIA1/L1: 1313 targets; LARP4B: 1621
targets; METAP2: 256 targets; HuR: 1124 targets; PUM2: 427 targets; HNRNPC: 232 targets).
mRNAs only bound by LARP4B or METAP2 are LARP4B/METAP2 targets and not bound by
another RBP (from the seven RBPs investigated), N = 717. mRNAs predominantly bound by
TIS11B are TIS11B targets exclusively bound by TIS11B or co-bound by TIA1/L1, with
TIS11B/TIA1/L1 ≥ 2 (N = 834). mRNAs predominantly bound by TIA1/L1 are TIA1/L1 targets
exclusively bound by TIA1/L1 or co-bound by TIS11B but TIS11B/TIA1/L1 < 2 (N = 634).

Intersection of membrane/secretory mRNAs with previous datasets. <u>APEX-seq.</u> The
mRNAs that are coexpressed in our RNA-seq dataset (N = 9155 mRNAs) and the ER
membrane-localized mRNAs from the APEX-seq dataset (N = 1045) were determined. [11] The
overlapping 845 mRNAs were intersected with the mRNAs that encode membrane/secretory
proteins found to be ER+ in our analysis (N = 1476). We detected 673 mRNAs which
correspond to 80% of all APEX-seq mRNAs that are considered to be ER membrane (ERM)-
enriched. The universe used to test for enrichment were all mRNAs that encode non-membrane
proteins (N = 2140). <u>Biochemical fractionation.</u> A similar analysis was performed for the
fractionation dataset from Reid (2012). [9] Among the 385 coexpressed mRNAs that are enriched
on the ER according to Reid, we detected 308 in our ER+ fraction when focusing on
membrane/secretory protein encoding mRNAs. This group represents 80% of all ER-enriched
mRNAs detected by Reid. <u>MERFISH.</u> In the MERFISH dataset, which was generated in U2OS
cells, 1037 mRNAs are considered ER-enriched. Among them, N = 571 are co-expressed in our
dataset and considered mRNAs encoding membrane/secretory proteins. Among the 571 co-
expressed mRNAs we consider 511 as ER+, which corresponds to 89%. Among the ER-de-
enriched mRNAs (Log2FC nonER vs ER = -0.34), only 69 mRNAs encode membrane/secretory
proteins. Among the 69 mRNAs, we consider 8 as ER+, which corresponds to 11.6%. [13]

Intersection of mRNAs that encode non-membrane proteins with a previous dataset. The
relative distribution of mRNA transcripts across subcellular compartments, including the
membrane fraction, phase-separated granules, and the cytosol was determined using density
gradient centrifugation in U2OS cells. [25] The number of co-expressed mRNAs that encode non-
membrane proteins was N = 6557, which corresponds to 93% of our dataset. This dataset
determines the proportion of transcripts that localize to the different fractions. For co-expressed
TG+ mRNAs (N = 1153), ER+ mRNAs (N = 839) and CY+ mRNAs (N = 1400), we plotted the
proportion of mRNAs that localize to phase-separated granules, to the membrane fraction, and
to the cytosol in the LoRNA dataset in U2OS cells.

Gene ontology analysis

Gene ontology (GO) analysis was performed using DAVID. [30]

Further statistical analysis

Statistical parameters are reported in the figures and figure legends, including the definitions
and exact values of N and experimental measures (mean ± std or boxplots depicting median,
25th and 75th percentile (boxes) and 5% and 95% confidence intervals (error bars). Pair-wise
transcriptomic feature comparisons were performed using a two-sided Mann-Whitney test. For
more than two samples, a Kruskal-Wallis test was performed. For transcriptomic analyses,
statistical significance is indicated by asterisks *, $0.05 > P > 1 \times 10^{-9}$; **, $1 \times 10^{-10} > P > 1 \times 10^{-20}$;
, $1 \times 10^{-21} > P > 1 \times 10^{-80}$; *, $1 \times 10^{-81} > P > 0$. Exact P values are listed in Table S3.
Enrichment was determined using a X^2 test. The P value was calculated using a two-sided
Fisher's exact test. When indicated, a two-sided t-test with assumption of equal variance was
applied. Statistical significance for experimental data is indicated by asterisks *, $P < 0.05$, **, $P <$
0.01, ***, $P < 0.001$, ****, $P < 0.0001$.

Data and code availability

The data of the proteomics experiment were deposited in the MassIVE repository (dataset
identifier MSV000092176). The RNA-seq samples obtained from the subcytoplasmic
fractionation and the TIS11B iCLIP data obtained from HEK293T cells are available at GEO
(Accession number: GSE215770). The code for logistic regression is available on github
(github.com/Mayrlab/tiger-seq). Raw western blot data, raw imaging data and scripts for
analysis are deposited at Mendeley (https://data.mendeley.com/datasets/nmt7ppsp8r/1).

Figure Legends

Figure 1. Strategy to identify compartment-enriched mRNAs.

1A. Confocal live cell imaging of HeLa cells after transfection of mCherry (mC)-TIS11B and
GFP-SEC61B to visualize TGs and the rough ER. Scale bar, 5 µm.

1B. Schematic of a cell with three cytoplasmic compartments.

1C. As in (A) but showing fluorescent TG (left) and ER (right) particles. Scale bar, 5 µm.

1D. The transcript localization scores for the TG samples are shown for all mRNAs that encode
non-membrane proteins (N = 7015), for the mRNAs defined as TG+ (N = 1246), and for mRNAs
considered to have an unbiased transcript distribution (N = 3369) to illustrate that the relative
transcript distribution between TG+ and unbiased mRNAs is substantially different. Mann
Whitney test, P = 0.

1E. As in (D), but the transcript localization scores for the ER samples are shown for all mRNAs
that encode non-membrane proteins (N = 7015), for the mRNAs defined as ER+ (N = 919), and
for mRNAs considered to have an unbiased transcript distribution (N = 3369). Mann Whitney
test, $P = 1 \times 10^{-123}$.

1F. As in (D), but the transcript localization scores for the cytosolic fractions are shown for all
mRNAs that encode non-membrane proteins (N = 7015), for the mRNAs defined as CY+ (N =
1481), and for mRNAs considered to have an unbiased transcript distribution (N = 3369). Mann
Whitney test, P = 0.

1G. SmRNA-FISH of endogenous TG+ mRNA *BAG3* (green) in HeLa cells. TIS granules (BFP-
TIS11B, blue) and the ER (GFP-SEC-61B, magenta) were simultaneously visualized. Bottom
panel shows a 5x zoom-in of the area indicated by dashed white box. White circles indicate
colocalization of mRNA puncta with TGs, whereas dashed white circles indicate colocalization
with the ER. Representative images are shown. Scale bar, 5 µm.

1H. As in (G), but smRNA-FISH of the endogenous ER+ mRNA *ALDH18A1* is shown.

1I. Quantification of TG-localizing smRNA-FISH foci of three TG+ and five ER+ endogenous
mRNAs. The white box plot indicates the expected fraction of mRNA transcripts based on the
TG compartment size, obtained from of 186 cells. mRNAs defined as TG+ are enriched in TGs
(Mann Whitney test, ***, (TG), $P = 5 \times 10^{-11}$), whereas mRNAs defined as ER+ are not enriched.
Additional images are shown in Fig. S2A-F and values for the individual mRNAs are shown in
Fig. S2H.

1J. As in (I) but quantification of ER-localizing smRNA-FISH foci of three TG+ and five ER+
endogenous mRNAs. The white box plot indicates the expected fraction of mRNA transcripts
based on the ER compartment size, obtained from of 186 cells. mRNAs defined as ER+ are
enriched on the ER (Mann Whitney test, ***, (TG), $P = 1 \times 10^{-6}$), whereas mRNAs defined as
TG+ are not enriched. Values for the individual mRNAs are shown in Fig. S2I.

1K. The ratio of smRNA-FISH foci colocalizing with the ER compared to the foci colocalizing
with TGs is shown for the TG+ and ER+ endogenous mRNAs from (I) and (J). T-test for
independent samples, *, P = 0.044.

1L. smRNA-FISH foci of endogenous mRNAs in HeLa cells before (−) and after (+) digitonin
extraction. Cell boundaries are indicated by the dotted lines. Representative images are shown.
Scale bar, 5 µm.

1M. Quantification of (L). Shown is the fraction of digitonin-resistant smRNA-FISH foci of
endogenous mRNAs as mean ± std of three independent experiments. Number of cells

analyzed, *MLST8*, *N* = 70; *SF3A2*, *N* = 67; *MAP2K2*, *N* = 48; *ALDH18A1*, *N* = 63; *TES, N* = 81;
IDH, *N* = 127; *BAG3*, *N* = 187; *DUSP1*, *N* = 162; *DNAJB1, N* = 138. Additional images are
shown in Fig. S3A-C. T-test for independent samples, *, *P* < 0.041.

1N. Summary of the smRNA-FISH validation for mRNAs defined as compartment enriched.

1101

Figure 2. Characteristics of compartment-enriched mRNAs.

2A. Steady-state mRNA abundance levels obtained from whole cell lysates. TG+, *N* = 1246;
ER+, *N* = 919, CY+, *N* = 1481; unbiased, *N* = 3369. Mann Whitney test: *, 0.05 >*P*> 1 x 10^{-9}; **,
1 x 10^{-10} >*P*> 1 x 10^{-20}; ***, 1 x 10^{-21} >*P*> 1 x 10^{-80}; ****, 1 x 10^{-81} >*P*> 0. Exact *P* values are listed
in Table S3. RPKM, reads per kilobase of transcript per million reads mapped.

2B. As in (A), but steady-state protein levels obtained from whole cell lysates are shown. TG+,
N = 469; ER+, *N* = 638; CY+, *N* = 833; unbiased, *N* = 2001.

2C. As in (B), but Pro-seq levels are shown, which indicate transcription rates. TG+, *N* = 1222;
ER+, *N* = 896; CY+, *N* = 1425; unbiased, *N* = 3268.

2D. As in (C), but estimated mRNA half-lives are shown.

2E. As in (A), but protein size distributions are shown. AA, amino acid

2F. As in (A), but mRNA length distributions are shown.

2G. As in (A), but 3′UTR length distributions are shown.

2H. As in (A), but average CDS exon length distributions are shown.

2I. Gene model of *ZFP36L1* (TIS11B) showing its unusual CDS exon length distribution. Tall
boxes indicate CDS exons and the narrow boxes indicate the 5′ and 3′UTRs.

2J. Gene ontology analysis for TG+ mRNAs. Shown are the top six functional gene classes and
their Benjamini-Hochberg adjusted *P* values for categories that are significantly and uniquely
enriched in TG+ mRNAs. The Benjamini-Hochberg adjusted *P* values for the same categories
for ER+ and CY+ mRNAs are shown for comparison.

2K. As in (J), but for ER+ mRNAs.

2L. As in (J), but for CY+ mRNAs.

1124

Figure 3. mRNA architecture features together with RBPs determine the subcytoplasmic
transcript distribution.

3A. 3′UTR-bound RBPs that are positively or negatively associated with compartment-enriched
mRNAs. Shown are the -log10 transformed *P* values obtained from logistic regression (see
Table S4).

3B. Pearson's correlation coefficients of mRNA and exon length with compartment localization
scores (LS).

3C. As in (A) but integrating 3′UTR-bound RBPs from (A) and the mRNA architecture features
'mRNA length' and 'CDS exon length'.

3D. The propensity to localize to TGs is shown for mRNAs stratified by exon length and bound
RBPs. No RBP (*N* = 1498), bound only by LARP4B or METAP2 (*N* = 717) or only by TIS11B (*N*
= 834). Mann Whitney test was performed. *P* value categories as in Fig. 2A.

3E. Model showing the additive effects of exon length and RBPs on the mRNA localization
propensity to TGs or the cytosol. RBPs can have positive (check) or negative (x) effects. Shown
as in Fig. 2I.

3F. As in (D) but shown is the propensity of mRNAs to localize to the ER, stratified by mRNA
length and the bound RBPs. Bound only by TIA1/L1 ($N = 634$).

3G. As in (E) but showing the additive effects of mRNA length and RBPs on the mRNA
localization propensity to the ER or the cytosol. RBPs can have positive or negative effects.

3H. As in (F) but shown is the propensity of mRNAs to localize to the cytosol. mRNAs bound
only by LARP4B or METAP2 ($N = 717$) or only by TIS11B ($N = 834$).

1146

Figure 4. Experimental validation of the regulators of subcytoplasmic mRNA transcript
distribution.

4A. TG+ mRNAs are shown and are color-coded based on their change in compartment
localization in TIS11B KO cells compared with control cells. No change ($N = 508$); up in the ER
($N = 197$); up in CY ($N = 217$).

4B. Length distribution of mRNAs from (A). Mann Whitney test was performed. *P* value
categories as in Fig. 2A. Exact *P* values are listed in Table S3.

4C. As in (B) but shown is protein size distribution.

4D. As in (B) but shown is CDS exon length distribution.

4E. As in Fig. 3E but shown are the mRNA features of TG+ mRNAs that change their
localization upon TIS11B KO.

4F. Schematic of the mRNA reporter used to validate the effect of a single 3′UTR-bound RBP
on mRNA localization. The *GFP-THAP1* reporter mRNA contains MS2 hairpins as 3′UTR, which
allow binding of the cotransfected MS2 coat protein (mCherry-tagged MCP). Fusion of TIAL1 to
MCP tethers it to the 3′UTR of the reporter mRNA. mC, mCherry.

4G. Confocal live cell imaging of HeLa cells expressing the indicated constructs. Scale bar, 5
µm.

4H. RNA-FISH (teal) of the GFP reporter mRNA from (F) in HeLa cells coexpressing the
indicated MCP-fusion construct together with BFP-SEC61B to visualize the rough ER
(magenta). Representative confocal images are shown. Scale bar, 5 µm.

4I. Line profiles of the fluorescence intensities obtained from the arrows shown in (H) together
with the obtained Pearson's correlation coefficients.

4J. Quantification of the Pearson's correlation coefficients between the *GFP-THAP1* reporter
mRNA and the rough ER in the experiment shown in (H). Two line profiles were generated for
each cell. For MCP, $N = 26$ cells and for MCP-TIAL1 $N = 21$ were analyzed. The horizontal line
denotes the median and the error bars denote the 25th and 75th percentiles. Mann-Whitney test,
****, $P < 0.0001$.

1174

Figure 5. 3′UTR-bound TIAL1 cooperates with the rough ER membrane environment to
increase protein expression.

5A. Protein abundance of mRNAs stratified by RBP binding. No RBP (*N* = 126), bound only by
TIS11B (*N* = 267), bound only by TIA1/L1 (*N* = 232). Mann Whitney test was performed. *P* value
categories as in Fig. 2A.

5B. GFP protein expression measured by FACS in HeLa cells using the *GFP-THAP1* reporter
mRNA with and without TIAL1 tethering. Representative histograms are shown. GFP-negative
cell populations are shown as dotted lines.

5C. Quantification of the experiment shown in (B). Shown is the mean ± std of five independent
experiments. T-test for independent samples, ****, *P* = 0.0003.

5D. Quantification of mRNA level of the experiment from (B). Shown is the mean ± std of three
independent experiments. T-test for independent samples, NS, not significant.

5E. Schematic of a *GFP-THAP1* mRNA reporter that investigates the influence of subcellular
mRNA localization on protein expression. Fusion of MCP to SEC61B localizes the GFP reporter
mRNA (shown as in Fig. 4F) to the rough ER membrane, whereas MCP alone localizes it to the
cytosol.

5F. Confocal live cell imaging of HeLa cells expressing the indicated constructs. Scale bar, 5
µm.

5G. As in (B) but the *GFP-THAP1* reporter mRNA was used with and without SEC61B tethering.

5H. Quantification of the experiment from (G). Shown is the mean ± std of four independent
experiments. T-test for independent samples, **, *P* =0.0026.

5I. Quantification of mRNA level in the experiment from (G). Shown is the mean ± std of three
independent experiments. T-test for independent samples, NS, not significant.

5J. As in Fig. 4F, but addition of a prenylation signal (CAAX) localizes the TIAL1-bound *GFP-*
THAP1 reporter mRNA to the plasma membrane. In the absence of CAAX, the TIAL1-bound
reporter mRNA localizes to the rough ER.

5K. Confocal live cell imaging of HeLa cells expressing the indicated constructs. Scale bar, 5
µm.

5L. As in (B) but the *GFP-THAP1* reporter mRNA was tethered with the indicated constructs.

5M. Quantification of the experiment from (L). Shown is the mean ± std of four independent
experiments. T-test for independent samples, ****, *P* < 0.0006, **, *P* = 0.002.

5N. Endogenous mRNAs bound by TIA1/L1 encode higher expressed proteins than mRNAs not
bound by any RBP. The largest TIA1/L1-associated increase was observed for ER+ mRNAs.
Mann Whitney test was performed. *P* value categories as in Fig. 2A.

1209

Figure 6. Localization of cytosolic mRNAs to the rough ER membrane increases their
protein expression.

6A. Schematic of a *GFP-THAP1* reporter mRNA bound by TIS11B that allows investigation of
localization-dependent GFP protein expression. Fusion of MCP and TIS11B localizes the mRNA
reporter to the cytosol, whereas the TIS11B-MCP-SEC61B fusion localizes the mRNA to the
rough ER membrane.

6B. Confocal live cell imaging of HeLa cells expressing the constructs from (A). Scale bar, 5 µm.

6C. As in Fig. 5B, but the *GFP-THAP1* reporter mRNA was tethered with the indicated MCP-
fusion constructs.

6D. Quantification of the experiment from (C). Shown is the mean ± std of four independent
experiments. T-test for independent samples, ****, $P < 0.0001$, **, $P = 0.003$, NS, not significant.

6E. Quantification of mRNA level in the experiment from (C). Shown is the mean ± std of three
independent experiments. T-test for independent samples, *, $P = 0.037$; NS, not significant.

1223

1224 **Figure 7. Model.**

1225 Model showing features of endogenous mRNAs with biased subcytoplasmic transcript
1226 distribution. Summarized are also the changes in protein abundance upon relocalization of the
1227 reporter from the cytosol to the ER (see text for details). The horizontal arrow indicates no
1228 change.

1229

1230

Supplementary Figures

Figure S1. Strategy to determine subcytoplasmic mRNA localization.

S1A. Cell fractionation strategy to obtain the cytoplasmic membrane fraction. Transfected
HEK293T cells were lysed in hypotonic buffer, followed by douncing and differential
centrifugation at 600 g to pellet nuclei. The supernatant was subsequently spun at 7000 g. The
pellet contains the cytoplasmic membrane fraction that was used for subsequent fluorescent
particle sorting.

S1B. FACS plot showing the gating strategy to obtain mCherry-TIS11B+/GFP-SEC61B+ TG
particles and mCherry-TIS11B−/GFP-SEC61B+ ER particles. The particles were costained with
DAPI to segregate TG and ER particles from nuclear contamination (DAPI high). The DAPI low
particles were sorted.

S1C. Immunoblot showing markers used to evaluate the quality of the three compartment
fractions. WCL is the unfractionated whole cell lysate. CY is the digitonin-extracted cytosol. Pre-
sort is the membrane-enriched cytoplasmic lysate from which TG and ER particles are sorted,
which is the final step in (A). 50K TG indicates 50,000 sorted TG particles and 50K ER indicates
50,000 sorted ER particles. H3 antibody was used as a marker for nuclear components,
GAPDH was used as marker for cytosolic proteins, and Calnexin and GFP-SEC61B were used
as ER markers. *, signal from mCherry-TIS11B.

S1D. Quantification of endogenous TIS11B and mCherry-TIS11B from (C) together with
additional biological replicates. Shown is the ratio of protein abundance in 50K TG over 50K ER
as mean ± std of three independent experiments.

S1E. As in (D), but quantification of endogenous Calnexin and GFP-SEC61B. Shown is the ratio
of protein abundance in 50K ER over 50K CY as mean ± std of three independent experiments.

S1F. Correlation of log2-transformed RPKM values obtained by RNA-seq for biological
replicates of subcytoplasmic compartments. The Pearson correlation coefficients are shown.

S1G. Baseline distribution of localization scores across the three investigated cytoplasmic
compartments is shown separately for mRNAs that encode membrane/secretory proteins and
mRNAs that encode non-membrane proteins.

S1H. Distribution of localization scores in each fractionation sample for compartment-enriched
mRNAs that encode membrane/secretory proteins.

S1I. Overlap of ER+ mRNAs that encode membrane/secretory proteins (N = 1476) defined by
us (particle sorting) with previous datasets that used alternative isolation methods. In the APEX-
seq dataset 78% of ER-localized mRNAs overlap with our ER+ mRNAs. In the fractionation
dataset, 80.5% of ER-localized mRNAs overlap with our ER+ mRNAs. Among the MERFISH
ER-localized mRNAs 89% overlap with our ER+ mRNAs and only 11.6% of our ER+ mRNAs
overlap with the ER-de-enriched mRNAs obtained by MERFISH.

S1J. ER+ mRNAs that encode membrane/secretory proteins (N = 1476) defined by us are
significantly enriched on the ER membrane (ERM) according to APEX-seq, whereas CY+
mRNAs that encode membrane/secretory proteins show a significantly lower ERM enrichment.
Similarly, CY+ mRNAs that encode membrane/secretory proteins have significantly higher
APEX2 enrichment scores in the cytosol than mRNAs considered ER+ by us. Mann Whitney
test: $P < 2\times10^{-7}$.

S1K. Ternary plot showing compartment-enriched mRNAs. Each dot represents an mRNA that
is color-coded as in Fig. 1D-F. mRNAs in the center (light grey) are considered to have an
unbiased transcript distribution.

S1L. Distribution of localization scores in each fractionation sample for compartment-enriched
mRNAs that encode non-membrane proteins.

S1M. Validation of mRNAs that encode non-membrane proteins and are defined as
compartment-enriched by us in comparison with the LoRNA dataset. Our TG+ mRNAs are
highest enriched in the mRNAs that LoRNA identifies in the phase-separated granule fraction.
Our ER+ mRNAs are highest enriched in the mRNAs that LoRNA identifies in the membrane
fraction. Our CY+ mRNAs are highest enriched in the mRNAs that LoRNA identifies in the
cytosol. Mann Whitney test was performed. Exact P values are listed in Table S3.

1284

Figure S2. Validation of endogenous TG+ and ER+ mRNAs by smRNA-FISH.

S2A. SmRNA-FISH of endogenous TG+ mRNA *DUSP1* (green) in HeLa cells. TGs (BFP-
TIS11B, blue) and the ER (GFP-SEC-61B, magenta) were simultaneously visualized. The
maximum projection of the fluorescent signals is shown. Bottom panel shows a 5x zoom-in of
the area indicated by the white dashed box. White circles indicate colocalization of mRNA
puncta with TGs, whereas dashed circles indicate colocalization with the ER. Representative
images are shown. Scale bar, 5 μm.

S2B. As in (A), but smRNA-FISH of endogenous TG+ mRNA *DNAJB1* is shown.

S2C. As in (A), but smRNA-FISH of endogenous ER+ mRNA *PLA2GA4* is shown.

S2D. As in (A), but smRNA-FISH of endogenous ER+ mRNA *IDH1* is shown.

S2E. As in (A), but smRNA-FISH of endogenous ER+ mRNA *TES* is shown.

S2F. As in (A), but smRNA-FISH of endogenous ER+ mRNA *ACTN4* is shown.

S2G. As in (E) but shown is a 2x magnification of the area indicated by the white dashed box.
The image illustrates the high background signal in the nucleus observed when probing for
endogenous *TES*. For quantification of this mRNA, the noise tolerance value used in the
puncta-calling function was increased to limit nuclear background noise. The white circles
indicate nuclear foci that are included in the quantification at this tolerance level. Scale bar, 5
μm.

S2H. Colocalization of TGs and smRNA-FISH foci of three TG+ (beige) and five ER+ (blue)
endogenous mRNAs. The beige box plot indicates the expected fraction of mRNA transcripts

based on the TG compartment size distribution, obtained from 186 cells. Number of cells
analyzed for RNA-FISH: *BAG3* (*N* = 17), *DUSP1* (*N* = 30), *DNAJB1* (*N* = 25), *PLA2GA4* (*N* =
17), *IDH1* (*N* = 18), *ALDH18A1* (*N* = 22), *TES* (*N* = 34), *ACTN4* (*N* = 23). Mann Whitney test
shows that the mRNA transcript distribution of 3/3 TG+ mRNAs (beige) is significantly higher
than what would be expected for unbiased mRNAs. *P* value categories as in Figure 2A, exact *P*
values listed in Table S3.

S2I. As in (H), but colocalization of the ER and smRNA-FISH foci of three TG+ (beige) and five
ER+ (blue) endogenous mRNAs. The blue box plot indicates the expected fraction of mRNA
transcripts based on the ER compartment size distribution, obtained from of 186 cells. Mann
Whitney test shows that the mRNA transcript distribution of 4/5 ER+ mRNAs (blue) is
significantly higher than what would be expected for unbiased mRNAs.

1316

Figure S3. Validation of CY+ mRNAs by smRNA-FISH after digitonin extraction.

S3A. Shown are smRNA-FISH foci of endogenous *SF3A2* mRNA and *MAP2K2* mRNA in HeLa
cells before (-) and after (+) digitonin extraction. *SF3A2* and *MAP2K2* are considered CY+
mRNAs. Cell boundaries are indicated by the dotted lines. Representative images are shown.
Scale bar, 5 µm.

S3B. As in (A), but smRNA-FISH foci of endogenous *TES* mRNA and *IDH1* mRNA are shown,
which are considered ER+ mRNAs.

S3C. As in (A), but smRNA-FISH foci of endogenous *DUSP1* mRNA and *DNAJB1* mRNA are
shown, which are considered TG+ mRNAs.

S3D. Schematic of the reporter mRNA used with the SunTag system to measure nascent
protein synthesis. CDS, coding sequence. The KIF18B construct was used previously. [32]

S3E. Confocal imaging of HeLa cells stably expressing SunTag reporter proteins svFc-GFP and
mCherry-tagged PP7 protein (mC-PP7) co-transfected with two constructs (i) BFP-TIS11B to
visualize TGs and (ii) SunTag-labeled mRNA encoding KIF18B with PP7-binding sites in the
3′UTR. The *KIF18B* mRNA is visualized by mC-PP7 binding (teal) whereas the KIF18B protein
is visualized by svFc-GFP binding (magenta). Foci with co-localization of mRNA and protein
represent nascent protein synthesis and are indicative of active translation. A representative
example is shown. The white box indicated area is shown at 6x magnification in the lower panel.
White arrows indicate actively translating mRNA, yellow arrow indicates non-translating mRNA.
Scale bar, 5 µm (top panel), 1 µm (bottom panel).

S3F. Quantification of the experiment from (E). Shown are the number of mRNA foci in TGs or
the cytosol (CY) using the Suntag reporter from (D). Each dot represents a cell. *N* = 24 cells
were analyzed.

S3G. As in (F), but shown are the mRNAs that are actively translated in each compartment,
which were identified by counting the teal and magenta-double positive foci.

1342

Figure S4. Characteristics of compartment-enriched mRNAs are shown for subgroups.

S4A. To justify the cut-off used to determine compartment-enriched mRNAs (Fig. 1D-F), we
show the data from Figure 2 in more detail. Steady-state mRNA abundance levels obtained
from whole cell lysates is shown for unbiased mRNAs (*N* = 3369), three subgroups of TG+
mRNAs (top, middle, bottom, *N* = 415 each); three subgroups of ER+ mRNAs (top, middle,
bottom, *N* = 306 each); three subgroups of CY+ mRNAs (top, middle, bottom, *N* = 493 each).
Even when focusing on the bottom-enriched groups (which are close to the cut-off used), the

differences across the compartment-enriched groups are still highly significant. Mann Whitney
tests were performed. *P* value categories as in Fig. 2A. Exact *P* values are shown in Table S3.
RPKM, reads per kilobase of transcript per million reads mapped.

S4B. As in (A), but steady-state protein levels obtained from whole cell lysates are shown.

S4C. As in Fig. 2C, but steady-state mRNA abundance levels of compartment-enriched mRNAs
obtained by RNA-seq from whole cell lysates of HEK293 cells are shown. This sample was
used together with the Pro-seq sample to estimate mRNA half-lives.

S4D. As in (A), but Pro-seq levels are shown.

S4E. As in (A), but estimated mRNA half-lives are shown.

S4F. As in (A), but protein size distributions are shown.

S4G. As in (A), but mRNA length distributions are shown.

S4H. As in (A), but 3′UTR length distributions are shown.

S4I. As in (A), but average CDS exon length distributions are shown.

S4J. As in Fig. 2A, but the number of exons per mRNA is shown.

S4K. As in (A), but the number of exons per mRNA is shown.

1365

Figure S5. Analyses of TIS11B CLIP data and TIS11B KO samples.

S5A. Gel showing samples used for iCLIP of GFP-tagged TIS11B. The region outlined in red
was used for iCLIP sample preparation.

S5B. TIS11B iCLIP tag distribution obtained from HEK293T cells.

S5C. The top five motifs that were enriched within TIS11B peaks in 3′UTRs compared to all
nucleotides in 3′UTRs. Shown are *P* values obtained by HOMER.

S5D. The fraction of mRNAs bound by at least one RBP (from Fig. 3A) for the different groups
of compartment-enriched mRNAs is shown.

S5E. Immunoblot of TIS11B in control cells and TIS11B KO HEK293T cells. H3 was used as
loading control.

S5F. Correlation of log2-transformed RPKM values obtained by RNA-seq for biological
replicates of sorted ER particles or digitonin-extracted cytosol samples for control and TIS11B
KO cells. The Pearson correlation coefficients are shown.

1379

Figure S6. mRNA localization-dependent protein expression of the GFP reporter.

S6A. Gating strategy to assess GFP protein expression of the reporter mRNA by FACS. Left
panel shows the ungated population of HeLa cells coexpressing MCP-mCherry and the *GFP-*
THAP1-MS2 reporter, separated by size (forward scatter) and granularity (side scatter). The
black circle indicates the live cells that were used for subsequent analysis. Middle panel, the
GFP- and mCherry-double positive population was gated to obtain the GFP mean fluorescence
values (MFI, right panel) which corresponds to the reported GFP protein expression values.

S6B. As in (A), but HeLa cells coexpressing MCP-mCherry-TIAL1 and the *GFP-THAP1-MS2*
reporter.

S6C. As in Fig. 4F and 5C, but the MS2 sites in the GFP reporter were omitted. Top: coexpression of MCP-mCherry-TIAL1 does not result in the binding of MCP to the reporter mRNA without MS2 sites. This experiment serves as control for the effect of TIA1L1 overexpression on reporter mRNA expression. Bottom: Quantification of the experiment with the GFP mRNA reporter lacking the MS2 binding sites. Shown is the mean ± std of three independent experiments. T-test for independent samples, NS, not significant.

S6D. Schematic of a second mRNA reporter used to validate the effect of a single 3′UTR-bound RBP on protein expression. The *GFP-BIRC3* reporter mRNA contains the BIRC3 coding region and MS2 hairpins as 3′UTR, which allow binding of the co-transfected MS2 coat protein (mCherry-tagged MCP). Fusion of TIAL1 to MCP tethers TIAL1 to the 3′UTR of the reporter mRNA. mC, mCherry.

S6E. GFP protein expression of the reporter mRNA from (D) in HeLa cells, coexpressing the indicated MCP-fusion constructs, measured by FACS. Representative histograms are shown. GFP-negative cell populations are shown as dotted lines.

S6F. Quantification of the experiment shown in (E). Shown is the mean ± std of five independent experiments. T-test for independent samples, **, $P = 0.005$.

S6G. RNA-FISH of the GFP reporter mRNA (teal) from Fig. 5E in HeLa cells coexpressing MCP-mCherry-SEC61B (magenta) to visualize colocalization between the mRNA and the rough ER membrane. Representative confocal images are shown. Scale bar, 5 µm.

S6H. Line profiles of the fluorescence intensities of the arrows from (G).

S6I. Quantification of the experiment from (G). Two line profiles were generated for each cell. The Pearson's correlation coefficients between the reporter mRNA and the ER were determined. For MCP, $N = 26$ cells were analyzed, for MCP-SEC61B, $N = 26$ cells were analyzed. The horizontal line denotes the median and the error bars denote the 25^{th} and 75^{th} percentiles. Mann-Whitney test, ****, $P < 0.0001$.

S6J. Schematic of a second GFP-tagged mRNA reporter that investigates the influence of subcellular mRNA localization on protein expression. Fusion of MCP to TRAPα localizes the GFP reporter mRNA to the ER membrane, whereas MCP alone localizes it to the cytosol.

S6K. Confocal live cell imaging of HeLa cells expressing mC-tagged TRAPα-MCP. Scale bar, 5 µm.

S6L. GFP protein expression of the reporter mRNAs from (J) coexpressing the indicated MCP-fusion constructs in HeLa cells measured by FACS. Representative histograms are shown. The histograms on the left indicate GFP-negative cell populations.

S6M. Quantification of the experiment from (L). Shown is the mean ± std of four independent experiments. T-test for independent samples, ****, $P < 0.0001$.

Figure S7. Redirecting mRNA localization from the cytosol to the rough ER overcomes the repressive effect of a bound RBP.

S7A. Confocal live cell imaging of HeLa cells expressing the indicated constructs. Shown are representative images with TGs or cytosolic TIS11B. mC, mCherry. Scale bar, 5 µm.

S7B. Quantification from (A). The fraction of HeLa cells with TGs is shown after transfection of the indicated TIS11B fusion constructs. $N = 165$ cells were analyzed for mCherry-TIS11B and $N = 198$ cells were analyzed for MCP-mCherry-TIS11B. MCP-mCherry-TIS11B largely prevents TG formation.

S7C. Schematic of a second TIS11B-bound mRNA reporter that allows investigation of mRNA
localization-dependent GFP expression. The coding region of the reporter is provided by BIRC3,
followed by MS2 binding sites. Tethering of MCP or TIS11B localizes the mRNA reporter to the
cytosol, whereas the MCP-TIS11B-SEC61B fusion localizes the mRNA reporter to the rough
ER.

S7D. GFP protein expression of the reporter mRNA from (C) in HeLa cells, coexpressing the
indicated MCP-fusion constructs, measured by FACS. Representative histograms are shown.
The histograms on the left indicate GFP-negative cell populations.

S7E. Quantification of the experiment from (D). Shown is the mean ± std of four independent
experiments. T-test for independent samples, ****, $P < 0.0001$, MCP vs TIS11B-SEC61B: $P =$
0.058; NS).

1444
1445

1446 **References**

1447
1. Huttelmaier, S., Zenklusen, D., Lederer, M., Dictenberg, J., Lorenz, M., Meng, X.,
Bassell, G.J., Condeelis, J., and Singer, R.H. (2005). Spatial regulation of beta-actin
translation by Src-dependent phosphorylation of ZBP1. Nature *438*, 512-515.
10.1038/nature04115.
2. Lecuyer, E., Yoshida, H., Parthasarathy, N., Alm, C., Babak, T., Cerovina, T., Hughes,
T.R., Tomancak, P., and Krause, H.M. (2007). Global analysis of mRNA localization
reveals a prominent role in organizing cellular architecture and function. Cell *131*, 174-
187. S0092-8674(07)01022-7 [pii]10.1016/j.cell.2007.08.003.

3. Moor, A.E., Golan, M., Massasa, E.E., Lemze, D., Weizman, T., Shenhav, R., Baydatch,
S., Mizrahi, O., Winkler, R., Golani, O., et al. (2017). Global mRNA polarization regulates
translation efficiency in the intestinal epithelium. Science *357*, 1299-1303.
10.1126/science.aan2399.
4. Tushev, G., Glock, C., Heumuller, M., Biever, A., Jovanovic, M., and Schuman, E.M.
(2018). Alternative 3' UTRs Modify the Localization, Regulatory Potential, Stability, and
Plasticity of mRNAs in Neuronal Compartments. Neuron *98*, 495-511 e496.
10.1016/j.neuron.2018.03.030.
5. Glock, C., Biever, A., Tushev, G., Nassim-Assir, B., Kao, A., Bartnik, I., Tom Dieck, S.,
and Schuman, E.M. (2021). The translatome of neuronal cell bodies, dendrites, and
axons. Proc Natl Acad Sci U S A *118*. 10.1073/pnas.2113929118.
6. Moretti, F., Rolando, C., Winker, M., Ivanek, R., Rodriguez, J., Von Kriegsheim, A.,
Taylor, V., Bustin, M., and Pertz, O. (2015). Growth Cone Localization of the mRNA
Encoding the Chromatin Regulator HMGN5 Modulates Neurite Outgrowth. Mol Cell Biol
35, 2035-2050. 10.1128/MCB.00133-15.
7. Buxbaum, A.R., Haimovich, G., and Singer, R.H. (2015). In the right place at the right
time: visualizing and understanding mRNA localization. Nature reviews. Molecular cell
biology *16*, 95-109. 10.1038/nrm3918.
8. Biever, A., Donlin-Asp, P.G., and Schuman, E.M. (2019). Local translation in neuronal
processes. Current opinion in neurobiology *57*, 141-148. 10.1016/j.conb.2019.02.008.
9. Reid, D.W., and Nicchitta, C.V. (2012). Primary role for endoplasmic reticulum-bound
ribosomes in cellular translation identified by ribosome profiling. J Biol Chem *287*, 5518-
5527. 10.1074/jbc.M111.312280.

10.   Jan, C.H., Williams, C.C., and Weissman, J.S. (2014). Principles of ER cotranslational
translocation revealed by proximity-specific ribosome profiling. Science *346*, 1257521.
10.1126/science.1257521.
11.   Fazal, F.M., Han, S., Parker, K.R., Kaewsapsak, P., Xu, J., Boettiger, A.N., Chang, H.Y.,
and Ting, A.Y. (2019). Atlas of Subcellular RNA Localization Revealed by APEX-Seq.
Cell *178*, 473-490.e426. 10.1016/j.cell.2019.05.027.
12.   Qin, W., Myers, S.A., Carey, D.K., Carr, S.A., and Ting, A.Y. (2021). Spatiotemporally-
resolved mapping of RNA binding proteins via functional proximity labeling reveals a
mitochondrial mRNA anchor promoting stress recovery. Nature communications *12*,
4980. 10.1038/s41467-021-25259-2.
13.   Xia, C., Fan, J., Emanuel, G., Hao, J., and Zhuang, X. (2019). Spatial transcriptome
profiling by MERFISH reveals subcellular RNA compartmentalization and cell cycle-
dependent gene expression. Proc Natl Acad Sci U S A *116*, 19490-19499.
10.1073/pnas.1912459116.
14.   Banani, S.F., Lee, H.O., Hyman, A.A., and Rosen, M.K. (2017). Biomolecular
condensates: organizers of cellular biochemistry. Nature reviews. Molecular cell biology
18, 285-298. 10.1038/nrm.2017.7.
15.   Shin, Y., and Brangwynne, C.P. (2017). Liquid phase condensation in cell physiology
and disease. Science *357*. 10.1126/science.aaf4382.
16.   Ma, W., and Mayr, C. (2018). A Membraneless Organelle Associated with the
Endoplasmic Reticulum Enables 3'UTR-Mediated Protein-Protein Interactions. Cell *175*,
1492-1506 e1419. 10.1016/j.cell.2018.10.007.
17.   Peeples, W., and Rosen, M.K. (2021). Mechanistic dissection of increased enzymatic
rate in a phase-separated compartment. Nature chemical biology *17*, 693-702.
10.1038/s41589-021-00801-x.
18.   Gasparski, A.N., Mason, D.E., Moissoglu, K., and Mili, S. (2022). Regulation and
outcomes of localized RNA translation. Wiley interdisciplinary reviews. RNA *13*, e1721.
10.1002/wrna.1721.
19.   Luo, Y., Pratihar, S., Horste, E.H., Mitschka, S., Mey, A.S.J.S., Al-Hashimi, H.M., and
Mayr, C. (2023). mRNA interactions with disordered regions control protein activity.
bioRxiv, 2023.2002.2018.529068. 10.1101/2023.02.18.529068.
20.   Gasparski, A.N., Moissoglu, K., Pallikkuth, S., Meydan, S., Guydosh, N.R., and Mili, S.
(2023). mRNA Location and Translation Rate Determine Protein Targeting to Dual
Destinations. bioRxiv, 2023.2004.2024.538105. 10.1101/2023.04.24.538105.
21.   Ma, W., Zhen, G., Xie, W., and Mayr, C. (2021). In vivo reconstitution finds multivalent
RNA-RNA interactions as drivers of mesh-like condensates. eLife *10*.
10.7554/eLife.64252.
22.   Han, X., Zhou, Z., Fei, L., Sun, H., Wang, R., Chen, Y., Chen, H., Wang, J., Tang, H.,
Ge, W., et al. (2020). Construction of a human cell landscape at single-cell level. Nature
581, 303-309. 10.1038/s41586-020-2157-4.
23.   Hubstenberger, A., Courel, M., Benard, M., Souquere, S., Ernoult-Lange, M., Chouaib,
R., Yi, Z., Morlot, J.B., Munier, A., Fradet, M., et al. (2017). P-Body Purification Reveals
the Condensation of Repressed mRNA Regulons. Mol Cell *68*, 144-157 e145.
10.1016/j.molcel.2017.09.003.
24.   Liu, X., and Fagotto, F. (2011). A method to separate nuclear, cytosolic, and membrane-
associated signaling molecules in cultured cells. Science signaling *4*, pl2.
10.1126/scisignal.2002373.
25.   Villanueva, E., Smith, T., Pizzinga, M., Elzek, M., Queiroz, R.M.L., Harvey, R.F.,
Breckels, L.M., Crook, O.M., Monti, M., Dezi, V., et al. (2022). A system-wide
quantitative map of RNA and protein subcellular localisation dynamics. bioRxiv,
2022.2001.2024.477541. 10.1101/2022.01.24.477541.

26. Boraas, L., Hu, M., Thornton, L., Vejnar, C.E., Zhen, G., Giraldez, A.J., Mayr, C., Wang,
S., and Nicoli, S. (2021). Non-coding function for mRNAs in Focal Adhesion Architecture
and Mechanotransduction. bioRxiv, 2021.2010.2004.463097.
10.1101/2021.10.04.463097.
27. Patel, R.K., West, J.D., Jiang, Y., Fogarty, E.A., and Grimson, A. (2020). Robust
partitioning of microRNA targets from downstream regulatory changes. Nucleic Acids
Res *48*, 9724-9746. 10.1093/nar/gkaa687.
28. Blumberg, A., Zhao, Y., Huang, Y.-F., Dukler, N., Rice, E.J., Chivu, A.G., Krumholz, K.,
Danko, C.G., and Siepel, A. (2021). Characterizing RNA stability genome-wide through
combined analysis of PRO-seq and RNA-seq data. BMC biology *19*, 30.
10.1186/s12915-021-00949-x.
29. Smalec, B.M., Ietswaart, R., Choquet, K., McShane, E., West, E.R., and Churchman,
L.S. (2022). Genome-wide quantification of RNA flow across subcellular compartments
reveals determinants of the mammalian transcript life cycle. bioRxiv,
2022.2008.2021.504696. 10.1101/2022.08.21.504696.
30. Huang da, W., Sherman, B.T., and Lempicki, R.A. (2009). Systematic and integrative
analysis of large gene lists using DAVID bioinformatics resources. Nature protocols *4*,
44-57. 10.1038/nprot.2008.211.
31. Vaquerizas, J.M., Kummerfeld, S.K., Teichmann, S.A., and Luscombe, N.M. (2009). A
census of human transcription factors: function, expression and evolution. Nature
reviews. Genetics *10*, 252-263. 10.1038/nrg2538.
32. Yan, X., Hoek, T.A., Vale, R.D., and Tanenbaum, M.E. (2016). Dynamics of Translation
of Single mRNA Molecules In Vivo. Cell *165*, 976-989. 10.1016/j.cell.2016.04.034.
33. Küspert, M., Murakawa, Y., Schäffler, K., Vanselow, J.T., Wolf, E., Juranek, S.,
Schlosser, A., Landthaler, M., and Fischer, U. (2015). LARP4B is an AU-rich sequence
associated factor that promotes mRNA accumulation and translation. Rna *21*, 1294-
1305. 10.1261/rna.051441.115.
34. Zhao, W., Zhang, S., Zhu, Y., Xi, X., Bao, P., Ma, Z., Kapral, Thomas H., Chen, S.,
Zagrovic, B., Yang, Yucheng T., and Lu, Zhi J. (2021). POSTAR3: an updated platform
for exploring post-transcriptional regulation coordinated by RNA-binding proteins.
Nucleic Acids Research *50*, D287-D294. 10.1093/nar/gkab702.
35. Wang, Z., Kayikci, M., Briese, M., Zarnack, K., Luscombe, N.M., Rot, G., Zupan, B.,
Curk, T., and Ule, J. (2010). iCLIP predicts the dual splicing effects of TIA-RNA
interactions. PLoS Biol *8*, e1000530. 10.1371/journal.pbio.1000530.
36. Meyer, C., Garzia, A., Mazzola, M., Gerstberger, S., Molina, H., and Tuschl, T. (2018).
The TIA1 RNA-Binding Protein Family Regulates EIF2AK2-Mediated Stress Response
and Cell Cycle Progression. Mol Cell *69*, 622-635 e626. 10.1016/j.molcel.2018.01.011.
37. Bertrand, E., Chartrand, P., Schaefer, M., Shenoy, S.M., Singer, R.H., and Long, R.M.
(1998). Localization of ASH1 mRNA particles in living yeast. Mol Cell *2*, 437-445.
38. Berkovits, B.D., and Mayr, C. (2015). Alternative 3' UTRs act as scaffolds to regulate
membrane protein localization. Nature *522*, 363-367. 10.1038/nature14321.
39. Lee, S.H., and Mayr, C. (2019). Gain of Additional BIRC3 Protein Functions through 3'-
UTR-Mediated Protein Complex Formation. Mol Cell *74*, 701-712 e709.
10.1016/j.molcel.2019.03.006.
40. Voigt, F., Zhang, H., Cui, X.A., Triebold, D., Liu, A.X., Eglinger, J., Lee, E.S., Chao, J.A.,
and Palazzo, A.F. (2017). Single-Molecule Quantification of Translation-Dependent
Association of mRNAs with the Endoplasmic Reticulum. Cell reports *21*, 3740-3753.
10.1016/j.celrep.2017.12.008.
41. Stoecklin, G., Colombi, M., Raineri, I., Leuenberger, S., Mallaun, M., Schmidlin, M.,
Gross, B., Lu, M., Kitamura, T., and Moroni, C. (2002). Functional cloning of BRF1, a
regulator of ARE-dependent mRNA turnover. EMBO J *21*, 4709-4718.

42. Lykke-Andersen, J., and Wagner, E. (2005). Recruitment and activation of mRNA decay
enzymes by two ARE-mediated decay activation domains in the proteins TTP and BRF-
1. Genes Dev *19*, 351-361. 10.1101/gad.1282305.

43. Galloway, A., Saveliev, A., Lukasiak, S., Hodson, D.J., Bolland, D., Balmanno, K.,
Ahlfors, H., Monzon-Casanova, E., Mannurita, S.C., Bell, L.S., et al. (2016). RNA-
binding proteins ZFP36L1 and ZFP36L2 promote cell quiescence. Science *352*, 453-
459. 10.1126/science.aad5978.

44. He, P.C., Wei, J., Dou, X., Harada, B.T., Zhang, Z., Ge, R., Liu, C., Zhang, L.S., Yu, X.,
Wang, S., et al. (2023). Exon architecture controls mRNA m(6)A suppression and gene
expression. Science *379*, 677-682. 10.1126/science.abj9090.

45. Kedersha, N.L., Gupta, M., Li, W., Miller, I., and Anderson, P. (1999). RNA-binding
proteins TIA-1 and TIAR link the phosphorylation of eIF-2 alpha to the assembly of
mammalian stress granules. The Journal of cell biology *147*, 1431-1442.
10.1083/jcb.147.7.1431.

46. Gilks, N., Kedersha, N., Ayodele, M., Shen, L., Stoecklin, G., Dember, L.M., and
Anderson, P. (2004). Stress granule assembly is mediated by prion-like aggregation of
TIA-1. Molecular biology of the cell *15*, 5383-5398. 10.1091/mbc.e04-08-0715.

47. Mazan-Mamczarz, K., Lal, A., Martindale, J.L., Kawai, T., and Gorospe, M. (2006).
Translational repression by RNA-binding protein TIAR. Mol Cell Biol *26*, 2716-2727.
10.1128/MCB.26.7.2716-2727.2006.

48. Pacheco-Fiallos, B., Vorländer, M.K., Riabov-Bassat, D., Fin, L., O'Reilly, F.J., Ayala,
F.I., Schellhaas, U., Rappsilber, J., and Plaschka, C. (2023). mRNA recognition and
packaging by the human transcription-export complex. Nature *616*, 828-835.
10.1038/s41586-023-05904-0.

49. Bonneau, F., Basquin, J., Steigenberger, B., Schäfer, T., Schäfer, I.B., and Conti, E.
(2023). Nuclear mRNPs are compact particles packaged with a network of proteins
promoting RNA-RNA interactions. Genes Dev. 10.1101/gad.350630.123.

50. Hachet, O., and Ephrussi, A. (2004). Splicing of oskar RNA in the nucleus is coupled to
its cytoplasmic localization. Nature *428*, 959-963. 10.1038/nature02521.

51. Cheng, L.C., Zheng, D., Zhang, Q., Guvenek, A., Cheng, H., and Tian, B. (2021).
Alternative 3' UTRs play a widespread role in translation-independent mRNA association
with the endoplasmic reticulum. Cell reports *36*, 109407. 10.1016/j.celrep.2021.109407.

52. Cao, J., Wu, L., Zhang, S.M., Lu, M., Cheung, W.K., Cai, W., Gale, M., Xu, Q., and Yan,
Q. (2016). An easy and efficient inducible CRISPR/Cas9 platform with improved
specificity for multiple gene targeting. Nucleic Acids Res *44*, e149. 10.1093/nar/gkw660.

53. Sanjana, N.E., Shalem, O., and Zhang, F. (2014). Improved vectors and genome-wide
libraries for CRISPR screening. Nat Methods *11*, 783-784. 10.1038/nmeth.3047.

54. Thomas, J.D., Polaski, J.T., Feng, Q., De Neef, E.J., Hoppe, E.R., McSharry, M.V.,
Pangallo, J., Gabel, A.M., Belleville, A.E., Watson, J., et al. (2020). RNA isoform screens
uncover the essentiality and tumor-suppressor activity of ultraconserved poison exons.
Nat Genet *52*, 84-94. 10.1038/s41588-019-0555-z.

55. Zarnegar, B.J., Flynn, R.A., Shen, Y., Do, B.T., Chang, H.Y., and Khavari, P.A. (2016).
irCLIP platform for efficient characterization of protein-RNA interactions. Nat Methods
13, 489-492. 10.1038/nmeth.3840.

56. Hu, M., Yang, B., Cheng, Y., Radda, J.S.D., Chen, Y., Liu, M., and Wang, S. (2020).
ProbeDealer is a convenient tool for designing probes for highly multiplexed
fluorescence in situ hybridization. Sci Rep *10*, 22031. 10.1038/s41598-020-76439-x.

57. Cui, X.A., Zhang, H., and Palazzo, A.F. (2012). p180 promotes the ribosome-
independent localization of a subset of mRNA to the endoplasmic reticulum. PLoS Biol
10, e1001336. 10.1371/journal.pbio.1001336.

58. Navarrete-Perea, J., Yu, Q., Gygi, S.P., and Paulo, J.A. (2018). Streamlined Tandem
Mass Tag (SL-TMT) Protocol: An Efficient Strategy for Quantitative (Phospho)proteome
Profiling Using Tandem Mass Tag-Synchronous Precursor Selection-MS3. Journal of
proteome research *17*, 2226-2236. 10.1021/acs.jproteome.8b00217.
59. Rappsilber, J., Mann, M., and Ishihama, Y. (2007). Protocol for micro-purification,
enrichment, pre-fractionation and storage of peptides for proteomics using StageTips.
Nature protocols *2*, 1896-1906. 10.1038/nprot.2007.261.
60. Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre,
C., Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and B cell
identities. Mol Cell *38*, 576-589. 10.1016/j.molcel.2010.05.004.
61. Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for
Illumina sequence data. Bioinformatics *30*, 2114-2120. 10.1093/bioinformatics/btu170.
62. Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based
genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol
37, 907-915. 10.1038/s41587-019-0201-4.
63. Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. Bioinformatics *30,* 923-930.
10.1093/bioinformatics/btt656.
64. Morales, J., Pujar, S., Loveland, J.E., Astashyn, A., Bennett, R., Berry, A., Cox, E.,
Davidson, C., Ermolaeva, O., Farrell, C.M., et al. (2022). A joint NCBI and EMBL-EBI
transcript set for clinical genomics and research. Nature *604*, 310-315. 10.1038/s41586-
022-04558-8.
65. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput
sequencing reads. 2011 *17*, 3. 10.14806/ej.17.1.200.
66. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*, 15-21. 10.1093/bioinformatics/bts635.
67. Frith, M.C., Valen, E., Krogh, A., Hayashizaki, Y., Carninci, P., and Sandelin, A. (2008).
A code for transcription initiation in mammalian genomes. Genome Res *18,* 1-12.
10.1101/gr.6831208.

1662

A Difference in localizaiton score (TIS11B KO vs control); Up in ER (down in CY) N = 197; Up in CY (down in ER) N = 217; Log2 (RPKM) control

B mRNA length (kb); No change, Up in ER, Up in CY; NS * *

C Protein length (AA x 1000); No change, Up in ER, Up in CY; * * *

D Exon length (kb); No change, Up in ER, Up in CY; * NS *

E No change; Up in ER; Up in CY

F GFP-THAP1 MCP mC; GFP-THAP1 MCP mCTIAL1

G MCP; MCP-TIAL1

H GFP-THAP1-MS2 / BFP-SEC61B / Merge; MCP; MCP-TIAL1

I MCP R= 0.31; MCP-TIAL1 R= 0.65; Fluorescence intensity; BFP-SEC61B; GFP-THAP1-MS2

J Correlation coefficient; **** ; MCP-TIAL1; MCP

A

B
MCP   MCP-TIS11B   TIS11B-MCP-SEC61B

C

D

E

**Localization change
of mRNA reporter**

$$CY \xrightarrow{\text{+ TIAL1}} ER \quad \begin{array}{l} \text{mRNA} \leftrightarrow \\ \text{Protein} \Uparrow \end{array}$$

**Features of endogenous mRNAs
with biased subcytoplasmic transcript distribution**

	TG+	ER+	CY+
mRNA length	Intermediate	Long	Short
Protein length	Intermediate	Long	Short
CDS exon length	Long	Short	Short
Exon number	Low	High	Low
Localizing RBP	TIS11B	TIA1/L1	LARP4B, METAP2
mRNA metabolism	Low production	-	High production, low stability
Encoded proteins	Low abundance, transcription factors	High abundance, large size	High abundance, small size