Autophagy is required to prevent early pro-inflammatory responses and neutrophil recruitment during *Mycobacterium tuberculosis* infection without affecting pathogen replication in macrophages

Rachel L. Kinsella*, Jacqueline M. Kimmey[#], Asya Smirnov, Reilly Woodson, Margaret R. Gaggioli, Sthefany M. Chavez, Darren Kreamalmeyer, Christina L. Stallings*

Department of Molecular Microbiology, Center for Women's Infectious Disease Research, Washington University School of Medicine, St. Louis, MO 63110, USA

[#]Current address: Department of Microbiology and Environmental Toxicology, UC Santa Cruz, Santa Cruz, CA, USA

*Correspondence: Rachel L. Kinsella: rlkinsella@wustl.edu, 314-286-0277

Christina L. Stallings: stallings@wustl.edu, 314-286-0276

Keywords: autophagy, *Mycobacterium tuberculosis*, macrophage, dendritic cells, neutrophils, inflammation, T_H17

Running title: Autophagy prevents neutrophil accumulation during infection

ABSTRACT

The immune response to *Mycobacterium tuberculosis* infection determines tuberculosis disease outcomes, yet we have an incomplete understanding of what immune factors contribute to a protective immune response. Neutrophilic inflammation has been associated with poor disease prognosis in humans and in animal models during *M. tuberculosis* infection and, therefore, must be tightly regulated. ATG5 is an essential autophagy protein that is required in innate immune cells to control neutrophil-dominated inflammation and promote survival during *M. tuberculosis* infection, however, the mechanistic basis for how ATG5 regulates neutrophil recruitment is unknown. To interrogate what innate immune cells require ATG5 to control neutrophil recruitment during *M. tuberculosis* infection, we used different mouse strains that conditionally delete *Atg5* in specific cell types. We found that ATG5 is required in CD11c$^+$ cells (lung macrophages and dendritic cells) to control the production of proinflammatory cytokines and chemokines during *M. tuberculosis* infection, which would otherwise promote neutrophil recruitment. This role for ATG5 is autophagy-dependent, but independent of mitophagy, LC3-associated phagocytosis, and inflammasome activation, which are the most well-characterized ways that autophagy proteins regulate inflammation. In addition to the increase in proinflammatory cytokine production during *M. tuberculosis* infection, loss of ATG5 in innate immune cells also results in an early induction of T$_H$17 responses. Despite prior published *in vitro* cell culture experiments supporting a role for autophagy in controlling *M. tuberculosis* replication in macrophages, loss of autophagy does not affect *M. tuberculosis* burden in macrophages *in vivo* and, therefore, the effects of autophagy on inflammatory responses occur without changes in pathogen numbers. These findings reveal new roles for autophagy proteins in lung resident macrophages and dendritic cells that are required to suppress inflammatory responses that are associated with poor control of *M. tuberculosis* infection.

INTRODUCTION

According to the World Health Organization, 10 million people fell ill with *Mycobacterium tuberculosis* infection and 1.5 million people died of tuberculosis (TB) in 2020, marking the first increase in TB-associated deaths in over a decade (1). Whether a person controls the initial *M. tuberculosis* infection or develops active TB disease is directly impacted by the type of immune response elicited in the infected individual (2). Therefore, better understanding of what constitutes a protective versus non-protective immune response to *M. tuberculosis* infection is critical for developing better therapies and prevention measures to fight this deadly disease. Genetic mouse models have provided invaluable insight into the immunological processes that are required for control of *M. tuberculosis* infection. Infection of mice through the aerosol route leads to phagocytosis of *M. tuberculosis* by alveolar macrophages, initiating an inflammatory response and recruitment of innate immune cells to the lung (2). *M. tuberculosis* replicates within these innate immune cells until antigen specific T cells traffic to the lung where they activate the innate immune cells to restrain *M. tuberculosis* replication and suppress inflammation. *M. tuberculosis* establishes a chronic infection in wild-type (WT) mice, which survive for over a year with this infection.

Atg5fl/fl-LysM-Cre mice, which delete the *Atg5* gene specifically in macrophages, inflammatory monocytes, some dendritic cells (DCs), and neutrophils, are severely susceptible to *M. tuberculosis* infection (3–5), highlighting ATG5 as a critical component of a protective immune response to *M. tuberculosis*. *M. tuberculosis* infected *Atg5fl/fl-LysM-Cre* mice fail to control bacterial replication and succumb to infection by 40 days post-infection (dpi) (3–5). The uncontrolled *M. tuberculosis* replication is associated with an early (by 14 dpi) and sustained influx of neutrophils in the *M. tuberculosis* infected *Atg5fl/fl-LysM-Cre* mice. Depletion of neutrophils during *M. tuberculosis* infection in *Atg5fl/fl-LysM-Cre* mice rescues the susceptibility and extends their survival (3), demonstrating that the neutrophil-dominated inflammation contributed to their susceptibility. In general, higher abundance of neutrophils during *M.*

tuberculosis infection have been associated with worse disease outcomes in mice (6–13) and humans

(12,14–17). Therefore, understanding the regulatory mechanisms that govern neutrophil recruitment and

accumulation during *M. tuberculosis* infection could be key for manipulating inflammatory responses to

better control TB.

ATG5 is required for the intracellular pathway of autophagy, a process by which cytoplasmic

contents are targeted to the lysosome for degradation (18,19). Initiation of autophagy involves phagophore

formation from the endoplasmic reticulum, which is mediated by the ULK1 complex (ULK1/ULK2,

ATG13, FIP200, ATG101) and the PI3 kinase complex (ATG14L, BECLIN1, VPS15, and

VPS34)(20,21). Elongation of the autophagosomal double membrane depends on two ubiquitin-like

conjugation systems. In the first system, ATG12 is activated by ATG7, transferred to ATG10, and

covalently attached to ATG5. The second ubiquitin-like component is LC3 (microtubule-associated

protein 1 light chain 3), which is conjugated to phosphatidylethanolamine, generating the membrane

bound form called LC3-II through the actions of ATG7 and ATG3. ATG5-ATG12 facilitates LC3

lipidation through its interactions with ATG3, while ATG16L1 specifies the localization of LC3

conjugation to the autophagosome membrane (18,19,21,22). The autophagosome membrane is then

completed and targeted for fusion with the lysosome where the autophagosome cargo are degraded. In

addition, ATG5 also functions outside of autophagy, including during *M. tuberculosis* infection (3),

although these activities remain poorly understood. Recent work supports an autophagy-dependent role

for ATG5 in LysM[+] innate immune cells in suppressing neutrophil recruitment to the lungs during *M.*

tuberculosis infection (23). Using *in vitro* cell culture experiments, multiple groups have reported that

macrophages require autophagy to control *M. tuberculosis* replication by targeting the pathogen to the

lysosome (xenophagy) as well as to prevent necrosis following days of infection in culture (5,23–29).

However, to date there is no evidence that xenophagy functions in this capacity *in vivo*. Therefore, the

mechanistic basis for how loss of ATG5 results in early and exaggerated recruitment of neutrophils during

M. tuberculosis infection *in vivo* remains unknown.

In this manuscript, we dissect the role for ATG5 in regulating neutrophil recruitment and

accumulation during *M. tuberculosis* infection *in vivo*. We find that ATG5 functions with other autophagy

proteins specifically in CD11c$^+$ lung macrophages and DCs to limit the production of cytokines and

chemokines that otherwise promote neutrophil influx to the lung early in *M. tuberculosis* infection. We

demonstrate that loss of autophagy in macrophages and DCs does not affect *M. tuberculosis* burden in

these cell types *in vivo* and instead changes the inflammatory response to the infection at time points

before cell survival is compromised. In addition, ATG5 is required in lung macrophages and DCs to limit

IL-17A production from CD4$^+$ T cells. Together, our studies reveal new roles for ATG5 and other

autophagy proteins in regulating inflammatory responses during infection, which with further dissection

could provide insight into pathways that may be targeted to effectively promote protective immune

responses during TB.

RESULTS

ATG5 is required in CD11c$^+$ lung macrophages and DCs to control neutrophil recruitment and

accumulation early during *M. tuberculosis* infection *in vivo*.

M. tuberculosis infection of *Atg5$^{fl/fl}$-LysM-Cre* mice results in the recruitment of a higher number of

neutrophils in the lungs at 14 dpi as compared to *Atg5$^{fl/fl}$* controls, despite equivalent bacterial burdens at

this time point (3). There are also no differences in the abundance of non-neutrophil cell types in *Atg5$^{fl/fl}$-*

LysM-Cre and *Atg5$^{fl/fl}$* mice at 14 dpi (3). This indicates that specifically neutrophils are accumulating in

M. tuberculosis-infected *Atg5$^{fl/fl}$-LysM-Cre* mice due to a defect in inflammatory responses to infection

and not due to higher burden. To determine which LysM$^+$ cells required ATG5 to control the early influx

of neutrophils into the lungs during *M. tuberculosis* infection, we compared bacterial burdens and neutrophil inflammation in *Atg5$^{fl/fl}$-LysM-Cre*, *Atg5$^{fl/fl}$-Mrp8-Cre* (deletion in neutrophils)*, *Atg5$^{fl/fl}$-Cd11c-Cre* (deletion in lung macrophages and DCs), and *Atg5$^{fl/fl}$* controls at 14 dpi. At 14 dpi, the *Atg5$^{fl/fl}$-LysM-Cre* and *Atg5$^{fl/fl}$-Cd11c-Cre* mice, but not *Atg5$^{fl/fl}$-Mrp8-Cre* mice, had higher levels of neutrophil inflammation in the lungs as compared to *Atg5$^{fl/fl}$* controls **(Fig 1A and 1B)**. The degree of increased neutrophil frequency was similar in *Atg5$^{fl/fl}$-LysM-Cre* and *Atg5$^{fl/fl}$-Cd11c-Cre* mice, indicating that loss of *Atg5* in CD11c$^+$ cells, but not neutrophils, leads to the early influx of neutrophils into the lungs during *M. tuberculosis* infection. At 14 dpi, none of the mouse strains harbored increased *M. tuberculosis* burden in their lungs **(Fig 1C)**, indicating that the increase in neutrophil abundance in *Atg5$^{fl/fl}$-Cd11c-Cre* mice is not due to elevated bacterial burden and reflects a dysregulated inflammatory response to infection.

To determine if the higher levels of neutrophils in the lungs of *M. tuberculosis*-infected *Atg5$^{fl/fl}$-CD11c-Cre* mice was due to elevated neutrophil abundance in circulation prior to or during infection, we monitored neutrophil frequency in the blood in uninfected and 14 dpi *Atg5$^{fl/fl}$* and *Atg5$^{fl/fl}$-CD11c-Cre* mice. There was no significant difference in the frequency of neutrophils in the blood of uninfected or 14 dpi *Atg5$^{fl/fl}$* and *Atg5$^{fl/fl}$-CD11c-Cre* mice **(S1A Fig)**, suggesting that the accumulation of neutrophils in the lungs of *M. tuberculosis*-infected *Atg5$^{fl/fl}$-CD11c-Cre* mice was due to specific recruitment of neutrophils to the site of infection or an inability to clear neutrophils from the lung. To begin to investigate this latter possibility, we monitored whether dead neutrophils were accumulating in the lungs of *Atg5$^{fl/fl}$-CD11c-Cre* mice during *M. tuberculosis* infection by analyzing neutrophil viability at 14 dpi by flow cytometry **(S1B Fig)**. We did not observe a significant difference in the frequency of viable neutrophils at 14 dpi in *Atg5$^{fl/fl}$* and *Atg5$^{fl/fl}$-CD11c-Cre* mice, indicating that increased neutrophil inflammation in the lungs of *Atg5$^{fl/fl}$-CD11c-Cre* mice was not due to differences in neutrophil viability.

The higher levels of neutrophils in the lungs of *Atg5^{fl/fl}-LysM-Cre* and *Atg5^{fl/fl}-Cd11c-Cre* mice were sustained through 21 dpi **(Fig 1D)**. However, only *Atg5^{fl/fl}-LysM-Cre* mice, and not *Atg5^{fl/fl}-Cd11c-Cre* mice, had higher bacterial burdens in the lungs at 21 dpi **(Fig 1E)**, similar to as previously reported (3). Loss of *Atg5* in neutrophils results in increased susceptibility to *M. tuberculosis* infection in some, but not all, *Atg5^{fl/fl}-Mrp8-Cre* mice (3). The susceptible *Atg5^{fl/fl}-Mrp8-Cre* mice accumulate higher neutrophil numbers and bacterial burdens in their lungs at 21 dpi **(Fig 1D and 1E)** (3). Therefore, loss of *Atg5* in neutrophils is likely contributing to the higher burdens in the lungs of *Atg5^{fl/fl}-LysM-Cre* mice at 21 dpi. These data indicate that ATG5 has a role in CD11c⁺ lung macrophages and DCs to regulate early recruitment of neutrophils, but not the control of *M. tuberculosis* replication during *M. tuberculosis* infection at 14 and 21 dpi.

To determine how the loss of Atg5 in CD11c⁺ cells and the resulting early influx of neutrophils into the lungs affected host susceptibility, we monitored survival in *M. tuberculosis* infected *Atg5^{fl/fl}-Cd11c-Cre* mice as compared to *Atg5^{fl/fl}* controls. *Atg5^{fl/fl}-Cd11c-Cre* mice succumbed to *M. tuberculosis* infection between 100 and 150 dpi, which was significantly earlier than *Atg5^{fl/fl}* controls (median survival time of 259 dpi) **(Fig 1F)**, but not as early as *Atg5^{fl/fl}-LysM-Cre* mice (succumb 30-40 dpi (3)). These data demonstrate that ATG5 is required in CD11c⁺ lung macrophages and DCs to control early neutrophil recruitment and promote survival following *M. tuberculosis* infection.

The role for ATG5 in lung macrophages and DCs in regulating neutrophil recruitment is dependent on other autophagy proteins.

Deletion of multiple different autophagy genes in all LysM⁺ innate immune cells can result in increased neutrophil recruitment to the lung during *M. tuberculosis* infection (23). However, we previously showed that at least one role for ATG5 in LysM⁺ innate immune cells in controlling *M. tuberculosis* infection is

autophagy-independent (3). Therefore, it is not known if the role for ATG5 specifically in CD11c$^+$ lung

macrophages and DCs is the same as described when broadly deleting *Atg5* in all LysM$^+$ cells. To

determine whether the regulation of neutrophil recruitment by ATG5 in CD11c$^+$ lung macrophages and

DCs was dependent on other autophagy proteins or represented the autophagy-independent role for ATG5

during *M. tuberculosis* infection, we monitored neutrophil abundance in the lungs of mice lacking

expression of another essential autophagy protein, BECLIN 1, in CD11c$^+$ cells (*Becn1$^{fl/fl}$-Cd11c-Cre*) at

14 dpi by flow cytometry. Similar to *Atg5$^{fl/fl}$-Cd11c-Cre* mice, *Becn1$^{fl/fl}$-Cd11c-Cre* mice also exhibited

elevated neutrophil frequency in the lung at 14 dpi relative to *Becn1$^{fl/fl}$* control mice **(Fig 2A)**, despite no

difference in bacterial burden **(Fig 2B)**. In addition, analysis of *M. tuberculosis* infected *Atg16l1$^{fl/fl}$-LysM-*

Cre and *Becn1$^{fl/fl}$-LysM-Cre* mice also revealed higher levels of neutrophils in the lungs at 14 dpi relative

to controls, without higher bacterial burdens **(Fig 2C and 2D)**.

In addition to their role in canonical autophagy, the proteins ATG5, BECLIN 1, and ATG16L1 are

also required for the process of LC3 associated phagocytosis (LAP), where LC3 is recruited to the

phagosome, resulting in LC3$^+$ single membrane vesicles that traffic to the lysosome for degradation. LAP

can dampen inflammatory responses through efferocytosis, pathogen removal, stimulating inhibitory

immune-receptor signaling, and reducing auto-antigen levels (30–33). In contrast to canonical autophagy,

LAP uses RUBICON and UVRAG instead of ATG14 in the PI3K complex and does not depend on ULK1

(33,34). To distinguish between whether ATG5, BECLIN 1 and ATG16L1 were functioning through

autophagy or LAP to regulate neutrophil recruitment during *M. tuberculosis* infection, we infected mice

lacking RUBICON expression, a protein specifically required for LAP. *Rubicon$^{-/-}$* mice had no difference

in neutrophil accumulation or bacterial burdens as compared to WT controls following *M. tuberculosis*

infection **(Fig 2C and 2D)**, indicating that LAP is not required to control neutrophil inflammation during

M. tuberculosis infection. Importantly, BECLIN 1 and ATG5 function at different steps of autophagy.

Therefore, the requirement of both BECLIN 1 and ATG5 suggests that both the initiation and elongation

steps of autophagy are required in CD11c$^+$ cells to control neutrophil recruitment early during *M.*

tuberculosis infection. Targeting of pathogens to the lysosome via autophagy is termed xenophagy and

multiple studies have reported roles for xenophagy in controlling *M. tuberculosis* replication in

macrophages and DCs in cell culture *in vitro* (5,23–29). However, in addition to there being no differences

in bacterial burden in the lungs at 14 dpi **(Fig 1C and 2D)** (3), there was no significant difference in the

proportion of macrophages, eosinophils, neutrophils, inflammatory monocytes, or DCs that were infected

with *M. tuberculosis* **(Fig 2E)** and no difference in the *M. tuberculosis* burden in autophagy-deficient cells

in the lungs of *Atg5$^{fl/fl}$-LysM-Cre* and *Atg5$^{fl/fl}$* mice at 14 dpi **(Fig 2F).** Loss of xenophagy in bone marrow

derived macrophages *in vitro* has also been associated with increased necrosis during *M. tuberculosis*

infection (23). However, we did not detect a difference in the viability of macrophages, inflammatory

monocytes, eosinophils, neutrophils or DCs in the lungs of *Atg5$^{fl/fl}$* and *Atg5$^{fl/fl}$-Cd11c-Cre* mice at 14 dpi

(Fig 2G, S1B Fig), although we cannot rule out effects on the balance of different cell death pathways**.**

These data support that the role for autophagy in CD11c$^+$ lung macrophages and DCs during *M.*

tuberculosis infection *in vivo* is independent of xenophagy regulating *M. tuberculosis* replication. In

addition, these data show that the elevated neutrophil abundance in *M. tuberculosis*-infected *Atg5$^{fl/fl}$-*

LysM-Cre mice is not due to differences in the cell viability or the cell types infected with *M. tuberculosis*

but instead is driven by an imbalanced inflammatory response.

Autophagy regulates proinflammatory responses in macrophages during *M. tuberculosis* infection.

We have previously shown that lungs of *Atg5$^{fl/fl}$-LysM-Cre* mice at 14 dpi contain higher levels of G-CSF

and IL-17A than control mice (3), cytokines that promote neutrophil development and recruitment. At this

time point, the primary CD11c^{+} cell types that are infected by *M. tuberculosis* are the lung macrophages,

encompassing alveolar and interstitial macrophages (35,36). Therefore, we hypothesized that autophagy

could be suppressing the production of these cytokines from *M. tuberculosis* infected macrophages. We

tested this hypothesis by culturing bone marrow derived macrophages (BMDMs) from *Atg5$^{fl/fl}$*, *Atg5$^{fl/fl}$*-

LysM-Cre, Atg16l1$^{fl/fl}$, Atg16l1$^{fl/fl}$-LysM-Cre, Becn1$^{fl/fl}$ and *Becn1$^{fl/fl}$-LysM-Cre* mice and infecting with *M.*

tuberculosis in vitro before monitoring cytokine and chemokine production using a cytokine bead array

(Bio-Rad) on the supernatants from infected cultures **(Fig 3A-3F and S2 Fig)**. Of the 23 cytokines tested,

we detected significantly higher levels of IL-1β, G-CSF, KC, TNF-α and RANTES from the *Atg5$^{fl/fl}$*-

LysM-Cre macrophage cultures compared to controls at 24 hpi **(Fig 3A-3E)**, despite no difference in

bacterial burdens or BMDM viability at this time point **(Fig 3F and S2A Fig)**, indicating that the

heightened proinflammatory response of *Atg5$^{-/-}$* BMDMs was not in response to increased antigen or

macrophage cell death. The levels of these cytokines were only different following *M. tuberculosis*

infection and not in mock infected cultures, indicating that the increased pro-inflammatory responses were

infection-induced. The higher levels of G-CSF and KC, both pro-inflammatory signals associated with

neutrophil inflammation (37–39), produced from *Atg5*-deficient macrophages in response to *M.*

tuberculosis infection were dose dependent and confirmed by ELISA **(Fig 3H and 3I)**. Similar to *Atg5$^{fl/fl}$*-

LysM-Cre BMDMs, *Atg16l1$^{fl/fl}$-LysM-Cre* BMDMs also produced higher levels of IL-1β, G-CSF and

TNF-α following *M. tuberculosis* infection *in vitro* **(Fig 3A, 3B and 3D)**. *Becn1$^{fl/fl}$-LysM-Cre* BMDMs

also produced more IL-1β and G-CSF following *M. tuberculosis* infection *in vitro* compared to control

cells **(Fig 3A, 3B and S2 Fig)** despite no difference in *M. tuberculosis* burden at this time point **(Fig 3F)**.

In addition, *Becn1$^{fl/fl}$-LysM-Cre* BMDMs produced higher levels of IL-6, MIP-1α, MIP-1β, and MCP-1

following *M. tuberculosis* infection **(Fig 3G and S2 Fig)**. IL-6 in particular is associated with neutrophil

recruitment (40–43) and also trended higher in *M. tuberculosis*-infected *Atg5$^{fl/fl}$-LysM-Cre* BMDMs **(Fig**

3G), so we further analyzed the levels of IL-6 produced by *M. tuberculosis*-infected *Atg5^{fl/fl}-LysM-Cre*

BMDMs using an ELISA (**Fig 3J**). These analyses revealed a dose-dependent increase of IL-6 secretion

in *M. tuberculosis*-infected *Atg5^{fl/fl}-LysM-Cre* BMDMs. We were able to detect more cytokines and

chemokines being differentially produced by *M. tuberculosis* infected BMDMs than what was previously

detected in the lungs of *Atg5^{fl/fl}-LysM-Cre* mice at 14dpi (3). This is likely due to the dilution of signals

produced by macrophages in the context of the total lung homogenate, decreasing our sensitivity to detect

macrophage specific responses. In addition, it is possible that the inflammatory responses of BMDMs

differ from lung macrophages and DCs during *M. tuberculosis* infection. Nonetheless, loss of expression

of the autophagy proteins ATG5, ATG16L1, or BECLIN 1 in macrophages both *in vivo* and *in vitro* results

in higher levels of cytokines and chemokines that are associated with neutrophil recruitment and

accumulation following *M. tuberculosis* infection relative to controls, indicating that canonical autophagy

is required in macrophages to control proinflammatory responses during *M. tuberculosis* infection without

affecting *M. tuberculosis* burden in macrophages.

Autophagy suppresses neutrophil recruitment early during *M. tuberculosis* infection independent

of mitophagy and inflammasome activation.

Autophagy has been shown to suppress proinflammatory responses by negatively regulating

inflammasome activation indirectly through regulation of NFκB signaling and directly by degrading pro-

IL-1β and clearance of inflammasome components (44–46), which can otherwise promote pro-

inflammatory responses, IL-1β secretion and neutrophil recruitment (44,45,47–49). Indeed, *Atg5^{fl/fl}-LysM-*

Cre, *Atg16l1^{fl/fl}-LysM-Cre* and *Becn1^{fl/fl}-LysM-Cre* BMDMs produce significantly more IL-1β in response

to *M. tuberculosis* infection *in vitro* at 24 hpi compared to control BMDMs (**Fig 3A**), supporting that loss

of autophagy has resulted in increased inflammasome activation. The primary inflammasome activated

during *M. tuberculosis* infection of macrophages is the NLRP3 inflammasome, which consists of the NOD-, LRR- and pyrin-domain containing protein 3 (NLRP3) sensor, ASC adaptor, and CASPASE 1(50– 53). TLR engagement and NFκB activation during *M. tuberculosis* infection constitute the priming step of inflammasome activation, resulting in increased expression of pro-IL-1β and NLRP3 (54,55). Phagocytosis of *M. tuberculosis* and subsequent Esx-1-dependent plasma membrane damage leading to potassium efflux is the second signal promoting NLRP3 inflammasome formation, which mediates CASPASE 1 activation followed by IL-1β processing and secretion (26, 29).

To determine whether the increased neutrophil inflammation following *M. tuberculosis* infection in autophagy-deficient mice results from increased inflammasome activation, we crossed *Caspase1/11$^{-/-}$* mice to *Atg5$^{fl/fl}$-LysM-Cre* and *Becn1$^{fl/fl}$-LysM-Cre* mice and monitored neutrophil abundance in the lungs at 14 dpi. *Caspase1/11$^{-/-}$/Atg5$^{fl/fl}$-LysM-Cre and Caspase1/11$^{-/-}$/Becn1$^{fl/fl}$-LysM-Cre* mice had similar neutrophil abundances and bacterial burdens in the lungs at 14 dpi as *Atg5$^{fl/fl}$-LysM-Cre mice* and *Becn1$^{fl/fl}$-LysM-Cre* mice, respectively **(Fig 4A and 4B)**, indicating that increased neutrophil recruitment in the absence of autophagy is occurring independent of CASPASE1/11. *Caspase1/11* deletion also did not extend the survival of *Atg5$^{fl/fl}$-LysM-Cre* mice, indicating that increased inflammasome activation does not contribute to the early susceptibility of these mice **(Fig 4C)**.

Autophagy has also been shown to suppress inflammatory responses via the process of mitophagy, where autophagy targets old and damaged mitochondria to the lysosome for degradation (56,57). The build-up of damaged or dysfunctional mitochondria in the absence of autophagy results in loss of mitochondrial membrane potential and the release of reactive oxygen species (ROS), mitochondrial DNA, and ATP to the cytosol where it can lead to oxidative damage, inflammasome activation, and pro-inflammatory cytokine production (57–61). Mitophagy requires the canonical autophagy proteins,

including ATG5, ATG16L1, and BECLIN 1, as well as PARKIN and PTEN-induced kinase 1 (PINK1)

(62). PINK1 accumulates on damaged mitochondria and activates the mitochondrial E3 ubiquitin ligase,

PARKIN, to ubiquitinylate damaged mitochondria (58,61). Optineurin and NDP52 are the main

mitophagy receptors that interact with the ubiquitinylated mitochondria and LC3, leading to

autophagosome engulfment of the mitochondria (58,63). To investigate whether loss of mitophagy could

contribute to higher neutrophil accumulation in the lungs following *M. tuberculosis* infection, we

measured neutrophil frequency in the lungs at 14 dpi by flow cytometry in *Parkin*$^{-/-}$ and *Pink1*$^{-/-}$ mice

relative to WT controls. There was no difference in neutrophil abundance or bacterial burdens in *M.*

tuberculosis-infected *Parkin*$^{-/-}$ or *Pink1*$^{-/-}$ mice relative to WT mice at 14 dpi **(Fig 4D and 4E)**, indicating

that mitophagy is not required to control neutrophil recruitment early during *M. tuberculosis* infection. To

determine if mitophagy is required in macrophages to control proinflammatory cytokine and chemokine

production during *M. tuberculosis* infection we generated BMDMs from *Parkin*$^{-/-}$, *Pink1*$^{-/-}$, and WT mice

and infected the macrophages with *M. tuberculosis* for 24 hours. We measured cytokine and chemokine

levels from mock and *M. tuberculosis* infected cultures using the cytokine bead array (Bio-Rad). Unlike

in autophagy-deficient BMDMs, there were no differences in IL-6, IL-1β, G-CSF, KC, TNF-α or

RANTES production by *M. tuberculosis* infected *Parkin*$^{-/-}$ and *Pink1*$^{-/-}$ macrophages at 24 hpi compared

to WT macrophages (**S3 Fig**), nor any differences in bacterial burden (**S3 Fig**). Therefore, loss of

mitophagy in macrophages does not result in higher levels of inflammation during *M. tuberculosis*

infection.

ATG5 is required to suppress early T$_H$17 responses in the lungs during *M. tuberculosis* infection.

The higher levels of IL-17A observed in the lungs of *Atg5*$^{fl/fl}$-*LysM-Cre* mice at 14 dpi with *M. tuberculosis*

relative to controls was not reproduced by BMDMs infected with *M. tuberculosis* for 24 hours (**S2 Fig**).

Although there are many reasons to explain this, one possibility is that the macrophages were not the

source of IL-17A *in vivo*. We investigated what cell type was expressing higher levels of IL-17A in the

Atg5^{fl/fl}-LysM-Cre mice during *M. tuberculosis* infection by crossing the *Atg5^{fl/fl}* and *Atg5^{fl/fl}-LysM-Cre*

mice with an IL-17A reporter mouse that expresses GFP under the IL-17A promoter (Jax # 018472). We

infected *Il17a-GFP/Atg5^{fl/fl}* and *Il17a-GFP/Atg5^{fl/fl}-LysM-Cre* mice with *M. tuberculosis* and monitored

GFP expression as a proxy of IL-17A expression in immune cells at 14 dpi. The only cell type we

reproducibly detected >0.5% of the cells expressing GFP were CD4$^+$ T cells. Similar to previous studies

with *Atg5^{fl/fl}* and *Atg5^{fl/fl}-LysM-Cre* mice, there was no difference in total CD4$^+$ T cell numbers in the

lungs of *Il17a-GFP/Atg5^{fl/fl}* and *Il17a-GFP/Atg5^{fl/fl}-LysM-Cre* mice at 14 dpi **(Fig 5A)** (3). However, a

greater frequency and number of the CD4$^+$ T cells in the lungs of *Il17a-GFP/Atg5^{fl/fl}-LysM-Cre* mice at

14 dpi were IL-17-GFP$^+$ compared to *Il17a-GFP/Atg5^{fl/fl}* mice **(Fig 5B and 5C)**. These data indicate that

CD4$^+$ T cells contribute to the higher levels of IL-17A in the lungs of *M. tuberculosis* infected *Atg5^{fl/fl}-*

LysM-Cre mice and ATG5 is required in innate immune cells to negatively regulate T$_H$17 responses during

M. tuberculosis infection.

IL-17A drives neutrophil influx by promoting the production of neutrophil chemokines MIP-1α

and KC and through activation of endothelial cells (64–66). Therefore, the increased T$_H$17 responses could

be responsible for the early influx and accumulation of neutrophils in the lungs of *M. tuberculosis*-infected

Atg5^{fl/fl}-LysM-Cre mice. To determine if the increased expression of IL-17A by T cells was responsible

for the enhanced influx of neutrophils at 14 dpi in *Atg5^{fl/fl}-LysM-Cre* mice, we depleted CD4$^+$ T cells by

administering antibodies specific for CD4 from day -2 to 14 dpi **(Fig 5D and 5E)**. At 14 dpi, we harvested

the lungs for enumeration of *M. tuberculosis* burden and neutrophil abundance and found that there was

no effect of CD4$^+$ T cell depletion on either readout **(Fig 5F and 5G)**. In addition, blocking IL-17A

signaling by administering an anti-IL-17A antibody from day -1 to 14 dpi **(Fig 5H)** did not affect *M.*

tuberculosis burden **(Fig 5I)** or neutrophil abundance **(Fig 5J)** in *Atg5$^{fl/fl}$-LysM-Cre* mice and *Atg5$^{fl/fl}$* mice

at 14 dpi. Therefore, although ATG5 is required in innate immune cells to suppress IL-17A expression in

T cells, this role does not contribute to differences in neutrophil accumulation early during *M. tuberculosis*

infection.

DISCUSSION

Atg5$^{fl/fl}$-LysM-Cre mice are extremely susceptible to *M. tuberculosis* infection, where neutrophils

accumulate in the lungs of infected *Atg5$^{fl/fl}$-LysM-Cre* mice by 14 dpi and are sustained at high levels until

the mice succumb to the infection between 30-40 dpi (3). It was previously unknown how ATG5 imparted

control of neutrophil recruitment to the lungs during *M. tuberculosis* infection. We have discovered that

ATG5 is required in CD11c$^+$ lung macrophages and DCs to regulate proinflammatory cytokine production

and neutrophil accumulation in the lungs during *M. tuberculosis* infection. This role for ATG5 is shared

with ATG16L1 and BECLIN 1, but not RUBICON, indicating it is autophagy-dependent and does not

involve LAP. *Atg5$^{fl/fl}$-CD11c-Cre* mice succumb to *M. tuberculosis* infection at a similar time as

Atg16l1$^{fl/fl}$-LysM-Cre and *Atg7$^{fl/fl}$-LysM-Cre* mice (23), suggesting that loss of the role for autophagy that

we have identified in CD11c$^+$ lung macrophages and DCs is responsible for the susceptibility of

Atg16l1$^{fl/fl}$-LysM-Cre and *Atg7$^{fl/fl}$-LysM-Cre* mice to *M. tuberculosis*. These data demonstrate that

autophagy is required in CD11c$^+$ cells to control inflammatory response during *M. tuberculosis* infection

and promote survival through the chronic phase of infection. We were able to reproduce the heightened

proinflammatory responses in *M. tuberculosis* infected autophagy-deficient BMDMs *in vitro*, suggesting

that autophagy specifically suppresses inflammatory responses from macrophages during *M. tuberculosis*

infection, although this does not rule out a similar role in DCs *in vivo*. Alveolar macrophages are among

the first cells to encounter *M. tuberculosis* in the airways and orchestrate the initial response to infection, recruiting other innate immune cells to the lung (35,36). We postulate that similar to the BMDMs, autophagy-deficient alveolar macrophages overproduce pro-inflammatory signals during *M. tuberculosis* infection, leading to increased neutrophil recruitment.

The increased levels of cytokines and chemokines produced by autophagy-deficient macrophages was dependent on *M. tuberculosis* infection, demonstrating that pathogen detection was required. However, the heightened proinflammatory responses in autophagy-deficient macrophages occurred in the absence of differences in bacterial burden, indicating that the enhanced inflammatory response is not due to increased antigen. Although multiple studies have shown effects of loss of autophagy on *M. tuberculosis* burden in macrophages and DCs *in vitro* in cell culture (5,23–29), there was no difference in *M. tuberculosis* burden in these cell types *in vivo* when we detected higher levels of proinflammatory cytokine production and neutrophil influx. Therefore, the role for autophagy in regulating proinflammatory cytokine production from CD11c[+] lung macrophages and DCs during *M. tuberculosis* infection is independent of xenophagy controlling *M. tuberculosis* replication. We also ruled out the involvement of CASPASE1/11-dependent inflammasome activity and mitophagy in autophagy-dependent regulation of neutrophil accumulation during *M. tuberculosis* infection, leaving open the question of how autophagy regulates macrophage proinflammatory responses during infection.

Our understanding of the number of cellular pathways regulated by autophagy continues to grow and there are many possible mechanisms by which autophagy could regulate inflammatory responses in macrophages (18). One possibility is that targeting of autophagy components to intracellular *M. tuberculosis* prevents sensing of *M. tuberculosis* by cytosolic pattern recognition receptors and subsequent proinflammatory cytokine and chemokine production, without impacting *M. tuberculosis* replication. Autophagy can also regulate levels of IL-1β (44,62,67), IL-6 (32,68,69) and TNF-α (68,69) downstream

of pathogen sensing, in part by negatively regulating NFκB activation (70–72). Autophagy can both

directly promote autophagic cell death and apoptosis as well as negatively regulate RIPK3-dependent

necrotic cell death, all which can impact inflammatory responses (23,73–77). Although we did not detect

overall differences in the viability of autophagy deficient cells at the time points we observed effects on

cytokine and chemokine production *in vivo* or *in vitro*, the mechanism of cell death in *Atg5$^{fl/fl}$* and *Atg5$^{-/-}$*

cells during *M. tuberculosis* infection may vary and could impact the inflammatory responses (73,74,78).

Autophagy may also be required in macrophages for the efficient removal of dead or infected neutrophils

from the lung and resolution of inflammation. The best studied process for removal of dead cells by

phagocytes is efferocytosis (79). LAP has specifically been shown to contribute to efferocytosis (80), and

although we rule out a role for LAP in regulating neutrophil accumulation during *M. tuberculosis*

infection, in remains possible that processing of dead cells may be attenuated in the absence of autophagy.

Another possible mechanism for how autophagy regulates inflammatory responses from macrophages

during *M. tuberculosis* infection involves the process of ER-phagy. ER-phagy is induced under conditions

of ER stress, accumulation of unfolded proteins, and during infection (81,82). ER-phagy restrains ER

stress responses by targeting excess or damaged endoplasmic reticulum to autophagosomes for

degradation (81), but in the absence of autophagy, ER stress activates NFκB-dependent transcription of

inflammatory cytokines, such as IL-1β, IL-6, IL-18 and TNF-α (82).

We also discovered that ATG5 was required in innate immune cells to suppress early T_H17

responses during *M. tuberculosis* infection. At this point, we do not know if the increased abundance of

IL-17$^+$ CD4$^+$ T cells in *M. tuberculosis* infected *Atg5$^{fl/fl}$-LysM-Cre* mice is due to an autophagy-dependent

or independent role for ATG5, but there are multiple other reports of autophagy regulating T_H17

responses. The effect of loss of *Atg5* in innate immune cells on IL-17A expression from T cells may be

explained by the requirement for autophagy in DCs to negatively regulate surface expression of disintegrin

and metalloproteinase domain-containing protein 10 (ADAM10). ADAM10 cleaves its substrate ICOSL and lower levels of ICOSL leads to decreased ICOSL-ICOS interactions between DCs and T cells, resulting in less $CD25^{hi}$ $CD4^+$ T regulatory cells and more $IL-17^+$ $CD4^+$ T cells (83). In addition, autophagy negatively regulates T_H17 differentiation by reducing IL-23 and IL-1β levels, which promote T_H17 differentiation and IL-17A secretion (4,84–86). Nonetheless, depleting $CD4^+$ T cells or blocking IL-17A did not rescue the increased neutrophil accumulation in the lungs of *M. tuberculosis* infected *Atg5[fl/fl]-LysM-Cre* mice at 14 dpi, suggesting that the hyper-inflammatory responses from infected autophagy-deficient macrophages and DCs is sufficient to recruit excessive neutrophils early during infection. However, it is possible that the heightened T_H17 responses in *M. tuberculosis* infected *Atg5[fl/fl]-LysM-Cre* mice have a longer-term impact on the increased susceptibility of these mice and could contribute to the susceptibility of *Atg5[fl/fl]-Cd11c-Cre* mice during the chronic phase of infection.

Our data support a role for autophagy in restricting neutrophil accumulation in the lung but does not rule out additional contributions from autophagy independent roles for ATG5, BECLIN 1 and ATG16L1. In particular, loss of *Atg5, Becn1* or *Atg16l1* in $LysM^+$ cells led to different degrees of elevated neutrophil frequency in the lung at 14 dpi relative to control mice, where loss of *Atg5* results in the greatest increase in neutrophil abundance. This could in part be due to differences in efficiency of gene deletion in the different lines (23). In addition, we have previously identified an autophagy independent role for ATG5 in neutrophils, where loss of ATG5 in neutrophils can result in early lethality during *M. tuberculosis* infection (3). Therefore, loss of *Atg5* in $LysM^+$ cells, which includes neutrophils, may result in higher neutrophil frequency in the lung at 14 dpi than loss of *Becn1* or *Atg16l1* because of this autophagy-independent role for ATG5 in neutrophils. We hypothesize that the combination of the newly discovered role for autophagy in $CD11c^+$ lung macrophages and DCs to regulate inflammatory responses and an autophagy-independent role for ATG5 in neutrophils collectively allow for control of *M. tuberculosis*

infection, where loss of both functions results in the extreme susceptibility of *Atg5^{fl/fl}-LysM-Cre* mice to

M. tuberculosis infection. Higher abundances of neutrophils have been associated with poor disease

prognosis and treatment outcomes in TB patients (14–16). Therefore, our new findings and future

dissection of the ATG5-dependent mechanisms of regulating neutrophil recruitment to the lungs during

M. tuberculosis infection will provide critical insight into how to promote protective immune responses

to TB.

MATERIALS AND METHODS

Mice

All flox mice (*Atg5^{fl/fl}*, *Atg16l1^{fl/fl}*, *Becn1^{fl/fl}*) used in this study have been described previously (3,87–89)

and colonies are maintained in an enhanced barrier facility. LysM-Cre (Jax #004781), Cd11c-Cre (Jax

#007567), Mrp8-Cre (Jax #021614) from the Jackson Laboratory were crossed to specific flox mice.

Il17a-IRES-GFP-KI (Jax # 018472) reporter mice were bred to *Atg5^{fl/fl}-LysM-Cre* and *Atg5^{fl/fl}* mice to

generate the *Il17a-GFP/Atg5^{fl/fl}-LysM-Cre* and *Il17a-GFP/Atg5^{fl/fl}* lines. *Rubicon^{-/-}* (Jax # 032581) and

WT litter mates were provided by D. Green and J. Martinez (90). Caspase 1/11^{-/-} (Jax #016621) were bred

to *Atg5^{fl/fl}-LysM-Cre* and *Becn1^{fl/fl}-LysM-Cre* mice. *Parkin^{-/-}* (Jax # 006582)(91), *Pink1^{-/-}* (Jax # 017946)

(56), and WT control mice were a gift from Dr. Jonathan Brestoff at Washington University School of

Medicine. Male and female littermates (aged 6-12 weeks) were used and were subject to randomization.

A minimum of 3 mice were used per experiment and each experiment was performed twice. Statistical

consideration was not used to determine mouse sample sizes. The mice were housed and bred at

Washington University in St. Louis in specific pathogen-free conditions in accordance with federal and

university guidelines, and protocols were approved by the Animal Studies Committee of Washington

University.

Infection of mice with *M. tuberculosis* and measurement of bacterial burden in the lungs

M. tuberculosis Erdman expressing GFP (*Mtb*-GFP (10,92)) was used in all experiments except experiments with the *Il-17a-GFP/Atg5^{fl/fl}-LysM-Cre* reporter mice when WT Erdman was used. *M. tuberculosis* was cultured at 37°C in 7H9 (broth) or 7H11 (agar) (Difco) medium supplemented with 10% oleic acid/albumin/dextrose/catalase (OADC), 0.5% glycerol, and 0.05% Tween 80 (broth). Cultures of GFP expressing *M. tuberculosis* were grown in the presence of kanamycin (20μg/mL) to ensure plasmid retention. *M. tuberculosis* cultures in logarithmic growth phase (OD600 = 0.5–0.8) were washed with PBS + 0.05% Tween-80, sonicated to disperse clumps, and diluted in sterile water before delivering 100 CFUs of aerosolized *M. tuberculosis* per lung using an Inhalation Exposure System (Glas-Col). Within 2 hours of each infection, lungs were harvested from at least two control mice, homogenized, and plated on 7H11 agar to determine the input CFU dose. At 14 dpi, *M. tuberculosis* titers were determined by homogenizing the superior, middle, and inferior lobes of the right lung and plating serial dilutions on 7H11 agar. Colonies were counted after 3 weeks of incubation at 37°C in 5% CO_2.

Flow cytometry from blood and infected lungs

Blood was collected by cardiac puncture into K2EDTA anticoagulant tubes (BD, 365974). Red blood cell lysis was performed by adding 2ml of ACK lysis buffer (Gibco, A10492-01) per 100μL of EDTA-treated blood for 5 minutes at room temperature. Cells were pelleted at 500xg for 5 minutes and then resuspended in 50uL of PBS+ 2% HI-FBS + 2mM EDTA in the presence of Fc receptor blocking antibody (BioLegend, 101302). Cells were labelled with antibodies as described below. Lungs were perfused with sterile PBS and digested for 1 hour with 625μg/mL collagenase D (Roche 11088875103) and 75U/mL DNase I (Sigma D4527). Cells were quenched with PBS + 2% heat-inactivated (HI)-FBS, + 2mM EDTA and passed

through a 70μM filter. Cells were stained with Zombie-violet or Zombie-NIR in PBS at 1:2000 dilution

for 5 minutes at room temperature prior to resuspending in PBS + 2% HI-FBS + 2mM EDTA in the

presence of Fc receptor blocking antibody (BioLegend, 101302) for blocking. Cells were labelled with

antibodies at a 1:200 dilution using the following mouse markers: CD11b_BV605 or PerCP-Cy5.5 (clone

M1/70), CD45_AF700 (BioLegend, 103259), Ly6G_PE-Cy7 or AF647 (clone 1A8), MHCII_Spark blue

550 (BioLegend, 107662), CD62L_Pe/Cy5 (BioLegend, 104410), CD44_BV510 (BioLegend, 103044),

CD11c_PerCP (BioLegend, 117325), Ly6C_BV605 (BioLegend, 128036), CD4_BV570 (clone RM4-5),

TCRb_BV421 (clone H57-597), CD19_Pacific blue (BioLegend, 115526), Zombie-violet or -NIR

(BioLegend, 423113 or 423105), SiglecF_BV480 (BD Biosciences, 746668), MHC-II_Sparkblue550

(BioLegend, 107662), CD64_PerCP-eFluor 710 (eBiosciences, 46061482) and MerTK_PE/Cy7

(eBiosciences, 25575182). Cells were incubated for 20 minutes at 4^{o}C with antibodies, washed and then

fixed in 4% paraformaldehyde (Electron Microscope Sciences) for 20 minutes at room temperature. Flow

cytometry was performed on an LSR-Fortessa (BD Bioscience) or an Aurora (Cytek Biosciences, with 4

laser 16V-14B-10YG-8R configuration) and analyzed using FlowJo software (Tree Star). Absolute cell

counts were determined using Precision count beads (BioLegend) or volumetric-based counting on the

Aurora. Gating strategies to identify *Mtb*-GFP^{+} myeloid cell populations, viable lung neutrophils, CD4^{+}

T cells and IL-17-GFP^{+} CD4^{+} T cells are in supplementary figure S4A-S4C.

<u>Culturing and infection of bone marrow derived macrophages (BMDMs)</u>

BMDMs were generated by flushing femurs and tibias of mice and culturing the cells in DMEM, 20% HI-

FBS, 10% supernatant from 3T3 cells overexpressing M-CSF + 1% MEM non-essential amino acids

(Cellgro 25-025-CI), 2mM L-glutamine, 100U/mL penicillin and 100μg/mL streptomycin (Sigma P4333)

at 37^{o}C in 5% CO_2 in non-TC treated plates. After 6 days non-adherent cells were removed and 1x10^{6}

adherent macrophages were seeded per well in 6 well non-TC treated plates in DMEM, 10% HI-FBS, 1%

MEM non-essential amino acids and 2 mM L-glutamine. BMDMs were rested overnight at 37^{o}C in 5%

CO$_2$. *M. tuberculosis* was grown to an OD of 0.6-0.8, washed with PBS twice, sonicated to disperse

clumps, centrifuged at 55xg for 10 minutes to remove clumps and resuspended in antibiotic-free BMDM

media. Macrophages were infected at an MOI of 10 by centrifuging the cells at 200xg for 10 minutes.

BMDMs were washed with PBS twice to remove free *M. tuberculosis* and fresh BMDM media was added

to each well. Cells were incubated at 37^{o}C and 5% CO$_2$ for 24 hours. To determine CFU counts, the cells

were lysed with 0.05% triton X-100, serially diluted, and plated onto 7H11 agar and incubated for 21 days

when bacterial colonies were counted. At 24 hours post-infection (hpi) supernatants were stored at -80^{o}C

for cytokine analysis. To assess BMDM viability, cells were harvested at 24 hpi by gently scrapping,

washed with PBS twice, stained with Zombie-violet in PBS at 1:2000 dilution for 5 minutes at room

temperature, washed once with PBS, and fixed in 4% PFA for 20 minutes at room temperature before

performing flow cytometry. Flow cytometry was performed on an LSR-Fortessa (BD Bioscience) and

analyzed using FlowJo software (Tree Star). The gating strategy to identify viable BMDMs at 24 hpi is

depicted in Supplementary Figure 1D.

Cytokine analysis

BMDM supernatants were filtered through a 0.22 µm filter twice to remove *M. tuberculosis* and analyzed

using the BioPlex-Pro Mouse Cytokine 23-Plex Immunoassay (Bio-Rad) as per the manufacturer's

instructions. ELISAs were performed according to the manufacturer's instructions (R&D systems):

KC/CXCL1 (DY453), IL-6 (DY406) and G-CSF (DY414).

IL-17A blocking and T cell depletion

To neutralize IL-17A, 100µg of InVivo monoclonal anti-IL-17A (Bio X Cell, BE0173) neutralizing

antibody was administered to *Atg5fl/fl* and *Atg5fl/fl-LysM-Cre* mice by intraperitoneal (i.p.) injection every

other day starting at 1 day prior to infection, with the final dose delivered at 13 dpi, similar to published

protocols (93). Control mice received 100µg of IgG from mouse serum (Sigma, I5381) by i.p. injection

every other day starting 1 day prior to infection and finishing on 13 dpi. To deplete $CD4^+$ T cells from

mice, 250µg of anti-mouse CD4 (Leinco Technologies, C1333) was administered by i.p. injection at 2

days prior to infection, 5 dpi and 12 dpi. Control mice received 250µg of IgG from rat serum (Sigma,

18015) i.p. on 2 days prior to infection, 5 dpi and 12 dpi.

<u>Data and statistics</u>

All experiments were performed at least twice. When shown, multiple samples represent biological (not

technical) replicates of mice randomly sorted into each experimental group. No blinding was performed

during animal experiments. Animals were only excluded when pathology unrelated to *M. tuberculosis*

infection was present (i.e. bad teeth leading to weight loss). Determination of statistical differences was

performed with Prism (GraphPad Software, Inc.) using log-rank Mantel-Cox test (survival), unpaired two-

tailed t-test (to compare two groups with similar variances), or one-way ANOVA with Šídák Multiple

Comparison test (to compare more than two groups). When used, center values and error bars represent

the mean +/- S.E.M. In all figures, all significant differences are indicated by asterisks: * *$P<0.05$, ** *

$P<0.01$, *** $P<0.001$, **** $P<0.0001$. Non-significant comparisons of particular interest are noted as

n.s.

ACKNOWLEDGEMENTS

The authors thank Dr. Jonathan Brestoff at Washington University School of Medicine for generously

providing the *Parkin^-/-* and *Pink1^-/-* mice. This work was supported by NIH grants R01 AI132697 and U19

AI142784, a Burroughs Wellcome Fund Investigators in the Pathogenesis of Infectious Disease Award,

and the Philip and Sima Needleman Center for Autophagy Therapeutics and Research to C.L.S., a Potts

Memorial Foundation postdoctoral fellowship to R.L.K., and a National Science Foundation Graduate

Research Fellowship DGE-1143954 and the NIGMS Cell and Molecular Biology Training Grant

GM007067 to J.M.K.

AUTHOR CONTRIBUTIONS

534   The experiments were designed by R.L.K., J.M.K., and C.L.S. The experiments were executed by R.L.K.,

535   J.M.K., and A.S. with assistance from R.W., M.R.G. and S.M.C. D.K. bred and maintained the mouse

536   colonies. The manuscript was written by R.L.K. and C.L.S.

FIGURE LEGENDS

Figure 1. ATG5 is required in CD11c$^+$ cells to regulate the early influx of neutrophils during *M.

***tuberculosis* infection *in vivo*.** (**A**) Representative flow cytometry plots of neutrophils

(CD45$^+$Ly6G$^+$CD11b$^+$) at 14 days post-infection (dpi) from *Atg5^fl/fl*, *Atg5^fl/fl-LysM-Cre*, *Atg5^fl/fl-CD11c-*

Cre and *Atg5^fl/fl-Mrp8-Cre* mice. (**B**) Proportion of CD45$^+$ cells that are neutrophils in the lung at 14 dpi

in *Mtb*-GFP infected *Atg5^fl/fl* (n=15), *Atg5^fl/fl-LysM-Cre* (n=8), *Atg5^fl/fl-CD11c-Cre* (n=10), and *Atg5^fl/fl-*

Mrp8-Cre (n=6) mice. Neutrophil frequency is reported as a ratio relative to the average neutrophil

frequency in *Atg5^fl/fl* control mice at 14 dpi within a given experiment. (**C**) Lung burden from the right

lung at 14 dpi in *Mtb*-GFP infected *Atg5^fl/fl* (n=15), *Atg5^fl/fl-LysM-Cre* (n=8), *Atg5^fl/fl-CD11c-Cre* (n=10),

and *Atg5^fl/fl-Mrp8-Cre* (n=6) mice. (**D**) Proportion of CD45$^+$ cells that are neutrophils in the lung at 21 dpi

in *Mtb*-GFP infected *Atg5$^{fl/fl}$* (n=19)*, Atg5$^{fl/fl}$-LysM-Cre* (n=6)*, Atg5$^{fl/fl}$-CD11c-Cre* (n=6) and *Atg5$^{fl/fl}$*-

Mrp8-Cre (n=7) mice. Neutrophil frequency is reported as a ratio relative to the average neutrophil

frequency in *Atg5$^{fl/fl}$* control mice at 21 dpi within a given experiment. Susceptible and healthy *Atg5$^{fl/fl}$*-

Mrp8-Cre mice are defined as done previously where susceptible *Atg5$^{fl/fl}$-Mrp8-Cre* mice have lost more

than 5% of their pre-infection body weight by 20 dpi and healthy *Atg5$^{fl/fl}$-Mrp8-Cre* mice have lost less

than 5% of their pre-infection body weight at 20 dpi (3). **(E)** Lung burden from the right lung at 21 dpi in

Mtb-GFP infected *Atg5$^{fl/fl}$* (n=19)*, Atg5$^{fl/fl}$-LysM-Cre* (n=6)*, Atg5$^{fl/fl}$-CD11c-Cre* (n=6), and *Atg5$^{fl/fl}$*-*Mrp8*-

Cre (n=7) mice. **(F)** Kaplan Meier curve of survival proportions during *Mtb*-GFP infection of *Atg5$^{fl/fl}$*

(n=9) and *Atg5$^{fl/fl}$-CD11c-Cre* (n=9) mice. Statistical differences were determined by a log-rank Mantel-

Cox test (F) or one-way ANOVA and Šídák multiple comparison test (B-E). * P < 0.05, ** P < 0.01, ***

P < 0.001, **** P < 0.0001. Differences that are not statistically significant are designated as ns. Pooled

data from at least two separate experiments is graphed where each data point is from one biological

replicate.

Figure 2. The role for ATG5 in lung macrophages and DCs in regulating neutrophil recruitment is

dependent on other autophagy proteins but does not involve control of pathogen infection or

burden. (A) Proportion of CD45^{+} cells that are neutrophils (CD45^{+}Ly6G^{+}CD11b^{+}) in the lung at 14 dpi

in *Mtb*-GFP infected *Becn1$^{fl/fl}$* (n=8) or *Becn1$^{fl/fl}$-CD11c-Cre* (n=11) mice. Neutrophil frequency is

reported as a ratio relative to the average neutrophil frequency in *Becn1$^{fl/fl}$* control mice at 14 dpi. **(B)**

Lung burden from the right lobes of the lung at 14 dpi in *Mtb*-GFP infected *Becn1$^{fl/fl}$* (n=8) or *Becn1$^{fl/fl}$*-

CD11c-Cre (n=11) mice. **(C)** Neutrophil frequency of CD45^{+} cells reported as a ratio to the average

neutrophil frequency in floxed control mice at 14 dpi in *Mtb*-GFP infected *Atg5$^{fl/fl}$* (n=6)*, Atg5$^{fl/fl}$-LysM-*

Cre (n=5)*, Becn1$^{fl/fl}$* (n=6)*, Becn1$^{fl/fl}$-LysM-Cre* (n=6)*, Atg16l1$^{fl/fl}$* (n=7)*, Atg16l1$^{fl/fl}$-LysM-Cre* (n=7) mice.

Neutrophil frequencies at 14 dpi in *Mtb*-GFP infected *Rubicon*[-/-] (n=6) mice were compared to wildtype

C57BL/6J (n=6) mice. **(D)** Lung burden from right lobes of the lung at 14 dpi in *Mtb*-GFP infected *Atg5*[fl/fl]

(n=6), *Atg5*[fl/fl]-*LysM-Cre* (n=5), *Becn1*[fl/fl] (n=6), *Becn1*[fl/fl]-*LysM-Cre* (n=6), *Atg16l1*[fl/fl] (n=7), *Atg16l1*[fl/fl]-

LysM-Cre (n=7)*, wildtype C57BL/6J (n=6) and *Rubicon*[-/-] (n=6) mice. **(E,F)** The proportion of *M.*

tuberculosis infected (*Mtb*-GFP[+]) and the median fluorescence intensity of *Mtb*-GFP in infected alveolar

macrophage (Alv. Mac.), eosinophils (Eos.), neutrophils (Neut.), inflammatory monocytes (Inf. Mono.),

CD11b[+] DC, CD103[+] DC, and non-alveolar macrophages in the lung in *Atg5*[fl/fl] (n=7) and *Atg5*[fl/fl]-*LysM-*

Cre (n=7) mice at 14 dpi. **(G)** The proportion of viable (Zombie[-]) alveolar macrophages, eosinophils,

neutrophils, inflammatory monocytes, CD11b[+] DC, CD103[+] DC and non-alveolar macrophages in the

lung in *Atg5*[fl/fl] (n=8) and *Atg5*[fl/fl]-*CD11c-Cre* (n=7) mice at 14 dpi. Statistical differences were determined

by student t-test to compare the LysM-Cre or CD11c-Cre mice to their respective floxed control and

Rubicon[-/-] mice to wildtype C57BL/6J mice (A-G).* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P <$

0.0001. Differences that are not statistically significant are designated as ns. Pooled data from at least two

separate experiments is graphed where each data point is from one biological replicate.

Figure 3. Autophagy regulates proinflammatory responses in macrophages during *M. tuberculosis*

infection. (A) Cytokine bead array data to quantify cytokines in culture supernatants from *Atg5*[fl/fl], *Atg5*[fl/fl]-

LysM-Cre[-], Atg16l1[fl/fl]*, Atg16l1*[fl/fl]-*LysM-Cre, Becn1*[fl/fl] or *Becn1*[fl/fl]-*LysM-Cre* BMDMs mock-treated or

infected with *Mtb*-GFP at an MOI of 10 for 24 hours. BMDMs generated from at least 3 mice were tested

in duplicate to quantify cytokine production. **(A)** IL-1β, **(B)** G-CSF, **(C)** KC, **(D)** TNF-α, and **(E)** RANTES

levels at 24 hpi are shown. **(F)** BMDM CFU counts at 24 hpi. **(G)** IL-6 levels at 24 hpi are shown. **(H)**

KC, **(I)** G-CSF, and **(J)** IL-6 levels at 24 hpi in *Atg5*[fl/fl] (n=8) and *Atg5*[fl/fl]-*LysM-Cre* (n=8) BMDMs

infected with *M. tuberculosis* at an MOI of 10 or 20 for 24 hours determined by ELISA. Statistical

differences were determined by student t-test to compare the autophagy gene-deficient cells to their

respective floxed control cells (A-J). * P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001. Differences

that are not statistically significant are designated as ns. Cytokine levels below detection limits are

designated as dbl. Each data point is one biological replicate, and the samples were generated from at least

two separate experiments.

Figure 4. Autophagy suppresses neutrophil recruitment independent of mitophagy and

inflammasome activation during *M. tuberculosis* infection. (A) Proportion of CD45$^+$ cells that are

neutrophils (CD45$^+$Ly6G$^+$CD11b$^+$) in the lung at 14 dpi in *Mtb*-GFP infected *Becn1$^{fl/fl}$* (n=5), *Becn1$^{fl/fl}$-*

LysM-Cre (n=7), *Caspase1/11$^{-/-}$/Becn1$^{fl/fl}$* (n=9), *Caspase1/11$^{-/-}$/Becn1$^{fl/fl}$-LysM-Cre* (n=13), *Atg5$^{fl/fl}$* (n=5),

Atg5$^{fl/fl}$-LysM-Cre (n=5), *Caspase1/11$^{-/-}$/Atg5$^{fl/fl}$* (n=4), or *Caspase1/11$^{-/-}$/Atg5$^{fl/fl}$-LysM-Cre* (n=5) mice

reported as a ratio relative to the average neutrophil frequency in corresponding floxed control mice. **(B)**

Lung burden at 14 dpi from right lobes of the lung in *Mtb*-GFP infected *Becn1$^{fl/fl}$* (n=5), *Becn1$^{fl/fl}$-LysM-*

Cre (n=7), *Caspase1/11$^{-/-}$/Becn1$^{fl/fl}$* (n=9), *Caspase1/11$^{-/-}$/Becn1$^{fl/fl}$-LysM-Cre* (n=13), *Atg5$^{fl/fl}$* (n=5),

Atg5$^{fl/fl}$-LysM-Cre (n=5), *Caspase1/11$^{-/-}$/Atg5$^{fl/fl}$* (n=4), or *Caspase1/11$^{-/-}$/Atg5$^{fl/fl}$-LysM-Cre* (n=5) mice.

The legend in 4A should be used for 4B too. **(C)** Kaplan Meier curve of survival proportions during *Mtb*-

GFP infection of *Atg5$^{fl/fl}$*, *Atg5$^{fl/fl}$-LysM-Cre*, *Caspase1/11$^{-/-}$/Atg5$^{fl/fl}$*, and *Caspase1/11$^{-/-}$/Atg5$^{fl/fl}$-LysM-Cre*

mice. **(D)** Proportion of CD45$^+$ cells that are neutrophils (CD45$^+$Ly6G$^+$CD11b$^+$) in the lung at 14 dpi in

Mtb-GFP infected wildtype (n=4), *Parkin$^{-/-}$* (n=3) or *Pink1$^{-/-}$* (n=3) mice reported as a ratio relative to the

average neutrophil frequency in wildtype mice. **(E)** Lung burden from the right lobe of the lung at 14 dpi

in *Mtb*-GFP infected wildtype (n=4), *Parkin$^{-/-}$* (n=3) or *Pink1$^{-/-}$* (n=3) mice. Statistical differences were

determined by log-rank Mantel-Cox test (C) and one-way ANOVA and Šídák multiple comparison test

(A-B, D-E) * P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001. Differences that are not statistically

significant are designated as ns. Pooled data from at least two separate experiments is graphed where each

data point is from one biological replicate.

Figure 5. ATG5 is required to suppress early T$_H$17 responses during *M. tuberculosis* infection (A)

The number of CD4$^+$ T cells (CD45$^+$ TCRβ$^+$ CD4$^+$) in *Atg5$^{fl/fl}$* and *Atg5$^{fl/fl}$-LysM-Cre* mice are reported as

the total cells per lung in the left lobe at 14 dpi. **(B)** The frequency of IL-17-GFP$^+$ CD4$^+$ T cells in the

lung (CD45$^+$ TCRβ$^+$ CD4$^+$ IL-17-GFP$^+$) of *Atg5$^{fl/fl}$* and *Atg5$^{fl/fl}$-LysM-Cre* mice are reported as the

percentage of CD4$^+$ T cells that are IL-17-GFP positive at 14 dpi. **(C)** The number IL-17-GFP$^+$ CD4$^+$ T

cells in *Atg5$^{fl/fl}$* and *Atg5$^{fl/fl}$-LysM-Cre* mice are reported as the total cells per lung in the left lobe at 14 dpi.

(D) Schematic depicting the timing of CD4-depletion antibody injections. **(E)** The number of CD4$^+$ T

cells (CD45$^+$ TCRβ$^+$ CD4$^+$) in *Mtb*-GFP infected *Atg5$^{fl/fl}$* and *Atg5$^{fl/fl}$-LysM-Cre* mice are reported as the

total cells per lung in the left lobe at 14 dpi following antibody treatment. **(F)** Lung burden from the right

lobes of the lung at 14 dpi in *Mtb*-GFP infected *Atg5$^{fl/fl}$* or *Atg5$^{fl/fl}$-LysM-Cre* mice that received isotype or

CD4-depletion antibodies. **(G)** Proportion of CD45$^+$ cells that are neutrophils (CD45$^+$Ly6G$^+$CD11b$^+$) in

the lung at 14 dpi in *Mtb*-GFP infected *Atg5$^{fl/fl}$* or *Atg5$^{fl/fl}$-LysM-Cre* mice that received CD4-depletion or

isotype treatment. Neutrophil frequency is reported as a ratio relative to the average neutrophil frequency

in *Atg5$^{fl/fl}$* control mice at 14 dpi. **(H)** Schematic depicting delivery of the IL-17 neutralizing antibody

treatments. **(I)** Lung burden from the right lobes of the lung at 14 dpi in *Mtb*-GFP infected *Atg5$^{fl/fl}$* or

Atg5$^{fl/fl}$-LysM-Cre mice that received isotype or IL-17 neutralizing antibodies. **(J)** Proportion of CD45$^+$

cells that are neutrophils (CD45$^+$Ly6G$^+$CD11b$^+$) in the lung at 14 dpi in *Mtb*-GFP infected *Atg5$^{fl/fl}$* or

Atg5$^{fl/fl}$-LysM-Cre mice that received isotype or IL-17 neutralizing antibodies. Neutrophil frequency is

reported as a ratio relative to the average neutrophil frequency in *Atg5$^{fl/fl}$* control mice at 14 dpi. Statistical

differences were determined by student t-test (A) or one-way ANOVA and Šídák multiple comparison

test (A-C, E-G, I and J). * P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001. Differences that are not

statistically significant are designated as ns. Pooled data from at least two separate experiments is graphed

where each data point is from one biological replicate.

Supplemental Figure 1. ATG5 is not required in CD11c$^+$ cells to regulate neutrophil viability in the

lung or accumulation of neutrophils in the blood during *M. tuberculosis* infection. (A) The proportion

of lung neutrophils (CD45$^+$ Ly6G$^+$ CD11b$^+$) that are viable (Zombie$^-$) in *M. tuberculosis* infected *Atg5$^{fl/fl}$*

(n=8) and *Atg5$^{fl/fl}$-CD11c-Cre* (n=7) mice at 14 dpi. **(B)** The proportion of CD45$^+$ cells in the blood that

are neutrophils in *Atg5$^{fl/fl}$* (n=4 in naïve and n=3 in 14 dpi) and *Atg5$^{fl/fl}$-CD11c-Cre* (n=4) mice at 14 dpi

and uninfected (naïve) mice. Statistical differences were determined by student t-test comparing the

genotypes within a particular cell type or treatment group. * P < 0.05, ** P < 0.01, *** P < 0.001, **** P

< 0.0001. Differences that are not statistically significant are designated as ns. Pooled data from at least

two separate experiments are graphed where each data point is from one biological replicate.

Supplemental Figure 2. Levels of viability, cytokines, and chemokines in autophagy-deficient

macrophages during *M. tuberculosis* infection. (A) The proportion of *Atg5$^{fl/fl}$* (n=5) and *Atg5$^{fl/fl}$-LysM-*

Cre (n=5) BMDMs that are viable (Zombie$^-$) at 24 hpi in mock and *Mtb*-GFP treated groups. Cytokine

bead array data from mock treated and *Mtb*-GFP infected BMDMs. *Atg5$^{fl/fl}$, Atg5$^{fl/fl}$-LysM-Cre, Atg16l1$^{fl/fl}$,*

Atg16l1$^{fl/fl}$-LysM-Cre, Becn1$^{fl/fl}$ and *Becn1$^{fl/fl}$-LysM-Cre* BMDMs were cultured for 24 hpi and cytokine

levels were measured in the spent media from mock treated or infected macrophages. BMDMs from at

least 3 mice were tested in duplicate to quantify the cytokines in the bead array. All cytokine and

chemokine data that are not significantly different between *Atg5$^{fl/fl}$-LysM-Cre* and *Atg5$^{fl/fl}$* mice are

reported here. **(B)** IL-1α, **(C)** IL-4, **(D)** IL-10, **(E)** IL-12(p40), **(F)** IL-12(p70), **(G)** IFN-γ, **(H)** MIP1β,

G(**I**) M-CSF, M(**J**) IP1α, (**K**) MCP-1, (**L**) Eotaxin and (**M**) IL-17 levels at 24 hpi. Statistical differences

were determined by one-way ANOVA and Šídák multiple comparison test (A) and student t-test

comparing the autophagy deficient macrophage with its floxed control within a treatment condition (B-

M). * P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001. Cytokine levels below detection limits are

designated as dbl. Differences that are not statistically significant are designated as ns. Each data point is

one biological replicate, and the samples were generated from at least two separate experiments.

Supplemental Figure 3. Mitophagy is not required in macrophages to regulate proinflammatory

responses during *M. tuberculosis* infection. WT*, Parkin*[-/-] and *Pink1*[-/-]*,* BMDMs were cultured for 24

hpi and cytokine levels were measured by cytokine bead array in the media from mock treated or *Mtb*-

GFP infected macrophages. (**A**) IL-1β, (**B**) G-CSF, (**C**) IL-6, (**D**) KC, (**E**) RANTES and (**F**) TNF-α levels

at 24 hpi (**G**) BMDM CFU counts from 24 hpi. BMDMs from at least 3 mice were tested in duplicate to

quantify cytokines in the bead array. Each point is one biological replicate. Statistical differences were

determined by one-way ANOVA and Šídák multiple comparison test (A-G). * P < 0.05, ** P < 0.01, ***

P < 0.001, **** P < 0.0001. Cytokine levels below detection limits are designated as dbl. Statistical

differences that are not significant are designated as ns. Each data point is one biological replicate, and

the samples were generated from at least two separate experiments.

Supplemental Figure 4. Gating strategies for flow cytometry identification of myeloid cells and IL-

17 expressing T cells in the lung or viable BMDMs. Representative flow cytometry plots depicting the

gating strategy used to identify *Mtb*-GFP[+] myeloid cells (**A**), viable lung neutrophils (**B**) and IL-17-GFP[+]

CD4[+] T cells (**C**) in the lung at 14 dpi. (**D**) Representative flow cytometry plots showing the gating strategy

to identify viable BMDMs at 24 hpi. FSC-A, forward scatter area; SSC-A, side scatter area; FSC-H,

forward scatter height; FSC-W, forward scatter width, Alv. Mac., alveolar macrophages; Eos.,

eosinophils; DC, dendritic cell, and Inf. Mono., inflammatory monocytes.

REFERENCES

1.   World Health Organization. Regional Office for South-East Asia. WHO global Tuberculosis report 2021
[Internet]. New Delhi: World Health Organization. Regional Office for South-East Asia; 2022 [cited 2022 Sep
9]. Available from: https://apps.who.int/iris/handle/10665/361162

2.   Kinsella R, Zhu D, Harrison G, Mayer Bridwell A, Prusa J, Chavez S, et al. Perspectives and Advances in the
Understanding of Tuberculosis. Annu Rev Pathol. 2021 Jan 24;16:377–408.

3.   Kimmey JM, Huynh JP, Weiss LA, Park S, Kambal A, Debnath J, et al. Unique role for ATG5 in neutrophil-
mediated immunopathology during M. tuberculosis infection. Nature. 2015 Dec;528(7583):565–9.

4.   Castillo EF, Dekonenko A, Arko-Mensah J, Mandell MA, Dupont N, Jiang S, et al. Autophagy protects against
active tuberculosis by suppressing bacterial burden and inflammation. Proc Natl Acad Sci. 2012 Nov
13;109(46):E3168–76.

5.   Watson RO, Manzanillo PS, Cox JS. Extracellular M. tuberculosis DNA Targets Bacteria for Autophagy by
Activating the Host DNA-Sensing Pathway. Cell. 2012 Aug;150(4):803–15.

6.   Mishra BB, Lovewell RR, Olive AJ, Zhang G, Wang W, Eugenin E, et al. Nitric oxide prevents a pathogen-
permissive granulocytic inflammation during tuberculosis. Nat Microbiol. 2017 Jul;2(7):17072.

7.   Mishra BB, Rathinam VAK, Martens GW, Martinot AJ, Kornfeld H, Fitzgerald KA, et al. Nitric oxide controls
the immunopathology of tuberculosis by inhibiting NLRP3 inflammasome–dependent processing of IL-1β.
Nat Immunol. 2013 Jan;14(1):52–60.

8.   Nandi B, Behar SM. Regulation of neutrophils by interferon-γ limits lung inflammation during tuberculosis
infection. J Exp Med. 2011 Oct 24;208(11):2251–62.

9.   Niazi MKK, Dhulekar N, Schmidt D, Major S, Cooper R, Abeijon C, et al. Lung necrosis and neutrophils reflect
common pathways of susceptibility to *Mycobacterium tuberculosis* in genetically diverse, immune
competent mice. Dis Model Mech. 2015 Jan 1;dmm.020867.

10.   Nair S, Huynh JP, Lampropoulou V, Loginicheva E, Esaulova E, Gounder AP, et al. Irg1 expression in myeloid
cells prevents immunopathology during M. tuberculosis infection. J Exp Med. 2018 Apr 2;215(4):1035–45.

11.   Dorhoi A, Yeremeev V, Nouailles G, Weiner J, Jörg S, Heinemann E, et al. Type I IFN signaling triggers
immunopathology in tuberculosis-susceptible mice by modulating lung phagocyte dynamics. Eur J Immunol.
2014 Aug;44(8):2380–93.

12. Moreira-Teixeira L, Stimpson PJ, Stavropoulos E, Hadebe S, Chakravarty P, Ioannou M, et al. Type I IFN
exacerbates disease in tuberculosis-susceptible mice by inducing neutrophil-mediated lung inflammation
and NETosis. Nat Commun. 2020 Dec;11(1):5566.

13. Moreira-Teixeira L, Tabone O, Graham CM, Singhania A, Stavropoulos E, Redford PS, et al. Mouse
transcriptome reveals potential signatures of protection and pathogenesis in human tuberculosis. Nat
Immunol. 2020 Apr;21(4):464–76.

14. Carvalho ACC, Amorim G, Melo MGM, Silveira AKA, Vargas PHL, Moreira ASR, et al. Pre-Treatment
Neutrophil Count as a Predictor of Antituberculosis Therapy Outcomes: A Multicenter Prospective Cohort
Study. Front Immunol. 2021 Jul 2;12:661934.

15. Han Y, Kim SJ, Lee SH, Sim YS, Ryu YJ, Chang JH, et al. High blood neutrophil-lymphocyte ratio associated
with poor outcomes in miliary tuberculosis. J Thorac Dis. 2018 Jan;10(1):339–46.

16. Lowe DM, Bandara AK, Packe GE, Barker RD, Wilkinson RJ, Griffiths CJ, et al. Neutrophilia independently
predicts death in tuberculosis: Table 1–. Eur Respir J. 2013 Dec;42(6):1752–7.

17. Berry MPR, Graham CM, McNab FW, Xu Z, Bloch SAA, Oni T, et al. An interferon-inducible neutrophil-driven
blood transcriptional signature in human tuberculosis. Nature. 2010 Aug;466(7309):973–7.

18. Kinsella RL, Nehls EM, Stallings CL. Roles for Autophagy Proteins in Immunity and Host Defense. Vet Pathol.
2018 May;55(3):366–73.

19. Kimmey JM, Stallings CL. Bacterial Pathogens versus Autophagy: Implications for Therapeutic Interventions.
Trends Mol Med. 2016 Dec;22(12):1060–76.

20. Lamb CA, Yoshimori T, Tooze SA. The autophagosome: origins unknown, biogenesis complex. Nat Rev Mol
Cell Biol. 2013 Dec;14(12):759–74.

21. Mizushima N, Yoshimori T, Ohsumi Y. The Role of Atg Proteins in Autophagosome Formation. Annu Rev Cell
Dev Biol. 2011 Nov 10;27(1):107–32.

22. Mizushima N, Noda T, Yoshimori T, Tanaka Y, Ishii T, George MD, et al. A protein conjugation system
essential for autophagy. Nature. 1998 Sep;395(6700):395–8.

23. Golovkine GR, Roberts AW, Morrison HM, Rivera-Lugo R, McCall RM, Nilsson H, et al. Autophagy restricts
Mycobacterium tuberculosis during acute infection in mice. Nat Microbiol [Internet]. 2023 Apr 10 [cited
2023 Apr 11]; Available from: https://www.nature.com/articles/s41564-023-01354-6

24. Manzanillo PS, Ayres JS, Watson RO, Collins AC, Souza G, Rae CS, et al. The ubiquitin ligase parkin mediates
resistance to intracellular pathogens. Nature. 2013 Sep;501(7468):512–6.

25. Köster S, Upadhyay S, Chandra P, Papavinasasundaram K, Yang G, Hassan A, et al. *Mycobacterium
tuberculosis* is protected from NADPH oxidase and LC3-associated phagocytosis by the LCP protein CpsA.
Proc Natl Acad Sci [Internet]. 2017 Oct 10 [cited 2023 Apr 4];114(41). Available from:
https://pnas.org/doi/full/10.1073/pnas.1707792114

26. Aylan B, Bernard EM, Pellegrino E, Botella L, Fearns A, Athanasiadi N, et al. ATG7 and ATG14 restrict
cytosolic and phagosomal Mycobacterium tuberculosis replication in human macrophages. Nat Microbiol
[Internet]. 2023 Mar 23 [cited 2023 Apr 4]; Available from: https://www.nature.com/articles/s41564-023-
01335-9

27. Dutta RK, Kathania M, Raje M, Majumdar S. IL-6 inhibits IFN-γ induced autophagy in Mycobacterium
tuberculosis H37Rv infected macrophages. Int J Biochem Cell Biol. 2012 Jun;44(6):942–54.

28. Deretic V. Autophagy in Tuberculosis. Cold Spring Harb Perspect Med. 2014 Nov 1;4(11):a018481–a018481.

29. Sakowski ET, Koster S, Portal Celhay C, Park HS, Shrestha E, Hetzenecker SE, et al. Ubiquilin 1 Promotes IFN-
γ-Induced Xenophagy of Mycobacterium tuberculosis. Lewinsohn DM, editor. PLOS Pathog. 2015 Jul
30;11(7):e1005076.

30. Heckmann BL, Green DR. LC3-associated phagocytosis at a glance. J Cell Sci. 2019 Mar 1;132(5):jcs231472.

31. Wan J, Weiss E, Ben Mkaddem S, Mabire M, Choinier PM, Picq O, et al. LC3-associated phagocytosis protects
against inflammation and liver fibrosis via immunoreceptor inhibitory signaling. Sci Transl Med. 2020 Apr
15;12(539):eaaw8523.

32. Inomata M, Xu S, Chandra P, Meydani SN, Takemura G, Philips JA, et al. Macrophage LC3-associated
phagocytosis is an immune defense against *Streptococcus pneumoniae* that diminishes with host aging. Proc
Natl Acad Sci. 2020 Dec 29;117(52):33561–9.

33. Martinez J, Almendinger J, Oberst A, Ness R, Dillon CP, Fitzgerald P, et al. Microtubule-associated protein 1
light chain 3 alpha (LC3)-associated phagocytosis is required for the efficient clearance of dead cells. Proc
Natl Acad Sci. 2011 Oct 18;108(42):17396–401.

34. Sanjuan MA, Dillon CP, Tait SWG, Moshiach S, Dorsey F, Connell S, et al. Toll-like receptor signalling in
macrophages links the autophagy pathway to phagocytosis. Nature. 2007 Dec;450(7173):1253–7.

35. Cohen SB, Gern BH, Delahaye JL, Adams KN, Plumlee CR, Winkler JK, et al. Alveolar Macrophages Provide an
Early Mycobacterium tuberculosis Niche and Initiate Dissemination. Cell Host Microbe. 2018 Sep;24(3):439-
446.e4.

36. Rothchild AC, Olson GS, Nemeth J, Amon LM, Mai D, Gold ES, et al. Alveolar macrophages generate a
noncanonical NRF2-driven transcriptional response *to Mycobacterium tuberculosis* in vivo. Sci Immunol.
2019 Jul 5;4(37):eaaw6693.

37. Vieira S, Lemos H, Grespan R, Napimoga M, Dal-Secco D, Freitas A, et al. A crucial role for TNF-a in mediating
neutrophil influx induced by endogenously generated or exogenous chemokines, KC/CXCL1 and LIX/CXCL5.
Br J Pharmacol. 2009;11.

38. Bendall LJ, Bradstock KF. G-CSF: From granulopoietic stimulant to bone marrow stem cell mobilizing agent.
Cytokine Growth Factor Rev. 2014 Aug;25(4):355–67.

39. Semerad CL, Liu F, Gregory AD, Stumpf K, Link DC. G-CSF Is an Essential Regulator of Neutrophil Trafficking
from the Bone Marrow to the Blood. Immunity. 2002 Oct;17(4):413–23.

40. Pollara G, Turner CT, Rosenheim J, Chandran A, Bell LCK, Khan A, et al. Exaggerated IL-17A activity in human
in vivo recall responses discriminates active tuberculosis from latent infection and cured disease. Sci Transl
Med. 2021 May 5;13(592):eabg7673.

41. Sasaki Y, Guo YM, Goto T, Ubukawa K, Asanuma K, Kobayashi I, et al. IL-6 Generated from Human
Hematopoietic Stem and Progenitor Cells through TLR4 Signaling Promotes Emergency Granulopoiesis by
Regulating Transcription Factor Expression. J Immunol. 2021 Aug 2;ji2100168.

42. Hu S, He W, Du X, Yang J, Wen Q, Zhong XP, et al. IL-17 Production of Neutrophils Enhances Antibacteria
Ability but Promotes Arthritis Development During Mycobacterium tuberculosis Infection. EBioMedicine.
2017 Sep;23:88–99.

43. Fielding CA, McLoughlin RM, McLeod L, Colmont CS, Najdovska M, Grail D, et al. IL-6 Regulates Neutrophil
Trafficking during Acute Inflammation via STAT3. J Immunol. 2008 Aug 1;181(3):2189–95.

44. Saitoh T, Akira S. Regulation of inflammasomes by autophagy. J Allergy Clin Immunol. 2016 Jul;138(1):28–36.

45. Harris J, Hartman M, Roche C, Zeng SG, O'Shea A, Sharp FA, et al. Autophagy Controls IL-1β Secretion by
Targeting Pro-IL-1β for Degradation. J Biol Chem. 2011 Mar;286(11):9587–97.

46. Pang Y, Wu L, Tang C, Wang H, Wei Y. Autophagy-Inflammation Interplay During Infection: Balancing
Pathogen Clearance and Host Inflammation. Front Pharmacol. 2022 Feb 22;13:832750.

47. Ko JH, Yoon SO, Lee HJ, Oh JY. Rapamycin regulates macrophage activation by inhibiting NLRP3
inflammasome-p38 MAPK-NFκB pathways in autophagy- and p62-dependent manners. Oncotarget. 2017
Jun 20;8(25):40817–31.

48. Shi CS, Shenderov K, Huang NN, Kabat J, Abu-Asab M, Fitzgerald KA, et al. Activation of autophagy by
inflammatory signals limits IL-1β production by targeting ubiquitinated inflammasomes for destruction. Nat
Immunol. 2012 Mar;13(3):255–63.

49. Pyrillou K, Burzynski LC, Clarke MCH. Alternative Pathways of IL-1 Activation, and Its Role in Health and
Disease. Front Immunol. 2020 Dec 18;11:613170.

50. McElvania TeKippe E, Allen IC, Hulseberg PD, Sullivan JT, McCann JR, Sandor M, et al. Granuloma Formation
and Host Defense in Chronic Mycobacterium tuberculosis Infection Requires PYCARD/ASC but Not NLRP3 or
Caspase-1. Tyagi AK, editor. PLoS ONE. 2010 Aug 20;5(8):e12320.

51. Beckwith KS, Beckwith MS, Ullmann S, Sætra RS, Kim H, Marstad A, et al. Plasma membrane damage causes
NLRP3 activation and pyroptosis during Mycobacterium tuberculosis infection. Nat Commun. 2020
Dec;11(1):2270.

52. Abdalla H, Srinivasan L, Shah S, Mayer-Barber KD, Sher A, Sutterwala FS, et al. Mycobacterium tuberculosis
Infection of Dendritic Cells Leads to Partially Caspase-1/11-Independent IL-1β and IL-18 Secretion but Not to
Pyroptosis. Kwaik YA, editor. PLoS ONE. 2012 Jul 24;7(7):e40722.

53. Mayer-Barber KD, Barber DL, Shenderov K, White SD, Wilson MS, Cheever A, et al. Cutting Edge: Caspase-1
Independent IL-1β Production Is Critical for Host Resistance to *Mycobacterium tuberculosis* and Does Not
Require TLR Signaling In Vivo. J Immunol. 2010 Apr 1;184(7):3326–30.

54. Lin KM, Hu W, Troutman TD, Jennings M, Brewer T, Li X, et al. IRAK-1 bypasses priming and directly links
TLRs to rapid NLRP3 inflammasome activation. Proc Natl Acad Sci. 2014 Jan 14;111(2):775–80.

55. Yu Qiao,Peng Wang,Jianni Qi,Lei Zhang,Chengjiang Gao. TLR-induced NF-jB activation regulates NLRP3
expression in murine macrophages. 1016 Feb 10;2012(586):1022–6.

56. Goldberg MS, Fleming SM, Palacino JJ, Cepeda C, Lam HA, Bhatnagar A, et al. Parkin-deficient Mice Exhibit
Nigrostriatal Deficits but Not Loss of Dopaminergic Neurons. J Biol Chem. 2003 Oct;278(44):43628–35.

57. Lazarou M, Sliter DA, Kane LA, Sarraf SA, Wang C, Burman JL, et al. The ubiquitin kinase PINK1 recruits
autophagy receptors to induce mitophagy. Nature. 2015 Aug;524(7565):309–14.

58. Killackey SA, Philpott DJ, Girardin SE. Mitophagy pathways in health and disease. J Cell Biol. 2020 Nov
2;219(11):e202004029.

59. Zhang W, Ren H, Xu C, Zhu C, Wu H, Liu D, et al. Hypoxic mitophagy regulates mitochondrial quality and
platelet activation and determines severity of I/R heart injury. eLife. 2016 Dec 20;5:e21407.

60. Wang S han, Martin SM, Harris PS, Knudson CM. Caspase Inhibition Blocks Cell Death and Enhances
Mitophagy but Fails to Promote T-Cell Lymphoma. Chandra D, editor. PLoS ONE. 2011 May 17;6(5):e19786.

61. Nguyen TN, Padman BS, Lazarou M. Deciphering the Molecular Signals of PINK1/Parkin Mitophagy. Trends
Cell Biol. 2016 Oct;26(10):733–44.

62. Harris J, Deen N, Zamani S, Hasnat MA. Mitophagy and the release of inflammatory cytokines.
Mitochondrion. 2018 Jul;41:2–8.

63. Song Y, Zhou Y, Zhou X. The role of mitophagy in innate immune responses triggered by mitochondrial
stress. Cell Commun Signal. 2020 Dec;18(1):186.

64. Griffin GK, Newton G, Tarrio ML, Bu D xiu, Maganto-Garcia E, Azcutia V, et al. IL-17 and TNF-α Sustain
Neutrophil Recruitment during Inflammation through Synergistic Effects on Endothelial Activation. J
Immunol. 2012 Jun 15;188(12):6287–99.

65. Roussel L, Houle F, Chan C, Yao Y, Bérubé J, Olivenstein R, et al. IL-17 Promotes p38 MAPK-Dependent
Endothelial Activation Enhancing Neutrophil Recruitment to Sites of Inflammation. J Immunol. 2010 Apr
15;184(8):4531–7.

66. Halwani R, Al-Muhsen S, Hamid Q. T Helper 17 Cells in Airway Diseases. Chest. 2013 Feb;143(2):494–501.

67. Ding WX, Jaeschke H. Autophagy in macrophages regulates the inflammasome and protects against liver
injury. J Hepatol. 2016 Jan;64(1):16–8.

68. Bussi C, Peralta Ramos JM, Arroyo DS, Gaviglio EA, Gallea JI, Wang JM, et al. Autophagy down regulates pro-
inflammatory mediators in BV2 microglial cells and rescues both LPS and alpha-synuclein induced neuronal
cell death. Sci Rep. 2017 Mar 3;7(1):43153.

69. Peng X, Wang Y, Li H, Fan J, Shen J, Yu X, et al. ATG5-mediated autophagy suppresses NF-κB signaling to limit
epithelial inflammatory response to kidney injury. Cell Death Dis. 2019 Mar 15;10(4):253.

70. Chang CP, Su YC, Hu CW, Lei HY. TLR2-dependent selective autophagy regulates NF-κB lysosomal
degradation in hepatoma-derived M2 macrophage differentiation. Cell Death Differ. 2013 Mar;20(3):515–
23.

71. Liu K, Zhang L, Zhao Q, Zhao Z, Zhi F, Qin Y, et al. SKP2 attenuates NF-κB signaling by mediating IKKβ
degradation through autophagy. Su B, editor. J Mol Cell Biol. 2018 Jun 1;10(3):205–15.

72. Yang Q, Liu TT, Lin H, Zhang M, Wei J, Luo WW, et al. TRIM32-TAX1BP1-dependent selective autophagic
degradation of TRIF negatively regulates TLR3/4-mediated innate immune responses. Philpott DJ, editor.
PLOS Pathog. 2017 Sep 12;13(9):e1006600.

73. Lim J, Park H, Heisler J, Maculins T, Roose-Girma M, Xu M, et al. Autophagy regulates inflammatory
programmed cell death via turnover of RHIM-domain proteins. eLife. 2019 Jul 9;8:e44452.

74. Bock FJ, Riley JS. When cell death goes wrong: inflammatory outcomes of failed apoptosis and mitotic cell
death. Cell Death Differ. 2023 Feb;30(2):293–303.

75. Tang D, Kang R, Coyne CB, Zeh HJ, Lotze MT. PAMPs and DAMPs: signal 0s that spur autophagy and
immunity. Immunol Rev. 2012 Sep;249(1):158–75.

76. Zhao X, Khan N, Gan H, Tzelepis F, Nishimura T, Park SY, et al. Bcl-xL mediates RIPK3-dependent necrosis in
M. tuberculosis-infected macrophages. Mucosal Immunol. 2017 Nov;10(6):1553–68.

77. Yousefi S, Perozzo R, Schmid I, Ziemiecki A, Schaffner T, Scapozza L, et al. Calpain-mediated cleavage of Atg5
switches autophagy to apoptosis. Nat Cell Biol. 2006 Oct;8(10):1124–32.

78. Mázló A, Jenei V, Burai S, Molnár T, Bácsi A, Koncz G. Types of necroinflammation, the effect of cell death
modalities on sterile inflammation. Cell Death Dis. 2022 May 2;13(5):423.

79. Doran AC, Yurdagul A, Tabas I. Efferocytosis in health and disease. Nat Rev Immunol. 2020 Apr;20(4):254–
67.

80. Asare PF, Roscioli E, Hurtado PR, Tran HB, Mah CY, Hodge S. LC3-Associated Phagocytosis (LAP): A
Potentially Influential Mediator of Efferocytosis-Related Tumor Progression and Aggressiveness. Front
Oncol. 2020 Aug 5;10:1298.

81. Moretti J, Roy S, Bozec D, Martinez J, Chapman JR, Ueberheide B, et al. STING Senses Microbial Viability to
Orchestrate Stress-Mediated Autophagy of the Endoplasmic Reticulum. Cell. 2017 Nov;171(4):809-823.e13.

82. Chipurupalli S, Samavedam U, Robinson N. Crosstalk Between ER Stress, Autophagy and Inflammation. Front
Med. 2021 Nov 5;8:758311.

83. Niven J, Madelon N, Page N, Caruso A, Harlé G, Lemeille S, et al. Macroautophagy in Dendritic Cells Controls
the Homeostasis and Stability of Regulatory T Cells. Cell Rep. 2019 Jul;28(1):21-29.e6.

84. Peral de Castro C, Jones SA, Ní Cheallaigh C, Hearnden CA, Williams L, Winter J, et al. Autophagy Regulates
IL-23 Secretion and Innate T Cell Responses through Effects on IL-1 Secretion. J Immunol. 2012 Oct
15;189(8):4144–53.

85. Cho ML, Kang JW, Moon YM, Nam HJ, Jhun JY, Heo SB, et al. STAT3 and NF-κB Signal Pathway Is Required for
IL-23-Mediated IL-17 Production in Spontaneous Arthritis Animal Model IL-1 Receptor Antagonist-Deficient
Mice. J Immunol. 2006 May 1;176(9):5652–61.

86. Sutton C, Brereton C, Keogh B, Mills KHG, Lavelle EC. A crucial role for interleukin (IL)-1 in the induction of IL-
17–producing T cells that mediate autoimmune encephalomyelitis. J Exp Med. 2006 Jul 10;203(7):1685–91.

87. Gawriluk TR, Hale AN, Flaws JA, Dillon CP, Green DR, Rucker EB. Autophagy is a cell survival program for
female germ cells in the murine ovary. REPRODUCTION. 2011 Jun;141(6):759–65.

88. Hara T, Nakamura K, Matsui M, Yamamoto A, Nakahara Y, Suzuki-Migishima R, et al. Suppression of basal
autophagy in neural cells causes neurodegenerative disease in mice. Nature. 2006 Jun;441(7095):885–9.

89. Hwang S, Maloney NS, Bruinsma MW, Goel G, Duan E, Zhang L, et al. Nondegradative Role of Atg5-Atg12/
Atg16L1 Autophagy Protein Complex in Antiviral Activity of Interferon Gamma. Cell Host Microbe. 2012
Apr;11(4):397–409.

90. Martinez J, Malireddi RKS, Lu Q, Cunha LD, Pelletier S, Gingras S, et al. Molecular characterization of LC3-
associated phagocytosis reveals distinct roles for Rubicon, NOX2 and autophagy proteins. Nat Cell Biol. 2015
Jul;17(7):893–906.

91. Kitada T, Pisani A, Porter DR, Yamaguchi H, Tscherter A, Martella G, et al. Impaired dopamine release and
synaptic plasticity in the striatum of *PINK1* -deficient mice. Proc Natl Acad Sci. 2007 Jul 3;104(27):11441–6.

92. Huynh JP, Lin CC, Kimmey JM, Jarjour NN, Schwarzkopf EA, Bradstreet TR, et al. Bhlhe40 is an essential
repressor of IL-10 during Mycobacterium tuberculosis infection. J Exp Med. 2018 Jul 2;215(7):1823–38.

93. Domingo-Gonzalez R, Das S, Griffiths KL, Ahmed M, Bambouskova M, Gopal R, et al. Interleukin-17 limits
hypoxia-inducible factor 1α and development of hypoxic granulomas during tuberculosis. JCI Insight. 2017
Oct 5;2(19):e92973.

A) Neutrophils 14 dpi

$Atg5^{fl/fl}$ — 27.5
$Atg5^{fl/fl}$-$LysM$-Cre — 47.0
$Atg5^{fl/fl}$-$CD11c$-Cre — 56.7
$Atg5^{fl/fl}$-$Mrp8$-Cre — 22.0

Ly6G / CD11b

B) Ratio of neutrophil frequency to $Atg5^{fl/fl}$

ns
****
****

□ $Atg5^{fl/fl}$
■ $Atg5^{fl/fl}$-$LysM$-Cre
■ $Atg5^{fl/fl}$-$CD11c$-Cre
■ $Atg5^{fl/fl}$-$Mrp8$-Cre

C) Lung CFU ($\log_{10}$)

ns

□ $Atg5^{fl/fl}$
■ $Atg5^{fl/fl}$-$LysM$-Cre
■ $Atg5^{fl/fl}$-$CD11c$-Cre
■ $Atg5^{fl/fl}$-$Mrp8$-Cre

D) Ratio of neutrophil frequency to $Atg5^{fl/fl}$

ns
**
***
****

□ $Atg5^{fl/fl}$
■ $Atg5^{fl/fl}$-$LysM$-Cre
■ $Atg5^{fl/fl}$-$CD11c$-Cre
■ $Atg5^{fl/fl}$-$Mrp8$-Cre susceptible
■ $Atg5^{fl/fl}$-$Mrp8$-Cre healthy

E) Lung CFU ($\log_{10}$)

ns
ns
***
ns
****
**

□ $Atg5^{fl/fl}$
■ $Atg5^{fl/fl}$-$LysM$-Cre
■ $Atg5^{fl/fl}$-$CD11c$-Cre
■ $Atg5^{fl/fl}$-$Mrp8$-Cre susceptible
■ $Atg5^{fl/fl}$-$Mrp8$-Cre healthy

F) Survival during *M. tuberculosis* infection

Probability of Survival / Days post-infection

□ $Atg5^{fl/fl}$ median survival 259 dpi (n=9)
■ $Atg5^{fl/fl}$-$CD11c$-Cre median survival 107 dpi (n=9)

****

A)

B)

C)

D)

E)

F)

G)

Legend:
- $Atg5^{fl/fl}$
- $Atg5^{fl/fl}$-LysM-Cre
- $Atg16l1^{fl/fl}$
- $Atg16l1^{fl/fl}$-LysM-Cre
- $Becn1^{fl/fl}$
- $Becn1^{fl/fl}$-LysM-Cre

A)

Ratio of neutrophil frequency to flox control

- *Becn1^{fl/fl}*
- *Becn1^{fl/fl}-LysM-Cre*
- *Caspase 1/11^{-/-} / Becn1^{fl/fl}*
- *Caspase 1/11^{-/-} / Becn1^{fl/fl}-LysM-Cre*
- *Atg5^{fl/fl}*
- *Atg5^{fl/fl}-LysM-Cre*
- *Caspase 1/11^{-/-} / Atg5^{fl/fl}*
- *Caspase 1/11^{-/-} / Atg5^{fl/fl}-LysM-Cre*

B)

Lung CFU ($\log_{10}$)

C)

Probability of Survival

Days post infection (dpi)

- *Atg5^{fl/fl}* median survival 185 dpi (n=8)
- *Atg5^{fl/fl}-LysM-Cre* median survival 28 dpi (n=8)
- *Caspase1/11^{-/-}/Atg5^{fl/fl}* median survival 253 dpi (n=6)
- *Caspase1/11^{-/-}/Atg5^{fl/fl}-LysM-Cre* median survival 35 dpi (n=9)

D)

Ratio of neutrophil frequency to WT

- WT
- *Parkin^{-/-}*
- *Pink1^{-/-}*

E)

Lung CFU ($\log_{10}$)

- WT
- *Parkin^{-/-}*
- *Pink1^{-/-}*

A) # CD4+ T cells/lung (10^5)
ns
Il-17a-GFP/Atg5^fl/fl (n=5)
Il-17a-GFP/Atg5^fl/fl-LysM-Cre (n=5)

B) % IL-17-GFP+ CD4+ T cells
*
Il-17a-GFP/Atg5^fl/fl (n=5)
Il-17a-GFP/Atg5^fl/fl-LysM-Cre (n=5)

C) # IL-17-GFP+ CD4+ T cells/lung (10^4)
*
Il-17a-GFP/Atg5^fl/fl (n=5)
Il-17a-GFP/Atg5^fl/fl-LysM-Cre (n=5)

D) Anti-CD4 or isotype
Infected   Sacrificed
-2 0   5   13 14
Days post-infection

E) CD4+ cells/lung (10^5)
*
ns   **
ns
Atg5^fl/fl (n=3) Isotype
Atg5^fl/fl-LysM-Cre (n=3) Isotype
Atg5^fl/fl (n=3) CD4 depletion
Atg5^fl/fl-LysM-Cre (n=4) CD4 depletion

F) Lung CFU (log10)
ns
Atg5^fl/fl (n=3) Isotype
Atg5^fl/fl-LysM-Cre (n=3) Isotype
Atg5^fl/fl (n=3) CD4 depletion
Atg5^fl/fl-LysM-Cre (n=4) CD4 depletion

G) Ratio of neutrophil frequency to *Atg5^fl/fl*
ns
ns
*   ***
Atg5^fl/fl (n=3) Isotype
Atg5^fl/fl-LysM-Cre (n=3) Isotype
Atg5^fl/fl (n=3) CD4 depletion
Atg5^fl/fl-LysM-Cre (n=4) CD4 depletion

H) Anti-IL-17 or isotype
Infected   Sacrificed
-1 0 1 3 5 7 9 11 13 14
Days post-infection

I) Lung CFU (log10)
ns
Atg5^fl/fl (n=5) Isotype
Atg5^fl/fl-LysM-Cre (n=7) Isotype
Atg5^fl/fl (n=5) IL-17 blocking
Atg5^fl/fl-LysM-Cre (n=8) IL-17 blocking

J) Ratio of neutrophil frequency to *Atg5^fl/fl*
ns
ns
**   ****
Atg5^fl/fl (n=5) Isotype
Atg5^fl/fl-LysM-Cre (n=7) Isotype
Atg5^fl/fl (n=5) IL-17 blocking
Atg5^fl/fl-LysM-Cre (n=8) IL-17 blocking