Dynamics of Mitochondrial NAD+ Import Reveal Preference for Oxidized Ligand and Substrate Led Transport

Shivansh Goyal [1] and Xiaolu A. Cambronne [1,2,3]

[1] Department of Molecular Biosciences, [2] LiveStrong Cancer Institute, University of Texas at Austin, Austin, TX 78712

[3] Correspondence: lulu@austin.utexas.edu

Abstract

SLC25A51 is a member of the mitochondrial carrier family (MCF) but lacks key residues that have been attributed to the mechanism of other nucleotide MCF transporters. Thus, how SLC25A51 transports NAD$^+$ across the inner mitochondrial membrane remains unclear. To elucidate its mechanism, we used Molecular Dynamic simulations to study reconstituted SLC25A51 homology models in lipid bilayers. We observed spontaneous binding of cardiolipin phospholipids to three distinct sites on the exterior of SLC25A51's central pore and found that mutation of these sites impaired transporter activity. We also observed that stable formation of the required matrix gate was controlled by a single salt bridge. Using simulation data and in-cell activity assays we identified binding sites in SLC25A51 for NAD$^+$ and showed that its binding was guided by an electrostatic interaction between NAD$^+$ and a negatively charged patch in the pore. In turn, interaction of NAD$^+$ with interior residue E132 guided the ligand to dynamically engage and weaken the salt bridge gate, representing a ligand-induced initiation of transport.

Significance

NAD$^+$ is an intermediary metabolite whose multiple functions are entwined with respiration, catabolism, and stress responses in cells. Previous sensor measurements had indicated that its continuous biosynthesis was required to sustain mitochondrial matrix levels in respiring cells, and SLC25A51 was identified as the required importer of NAD$^+$ across the inner mitochondrial membrane. However, SLC25A51 has minimal amino acid homology to other nucleotide carriers. By combining modeling approaches and experimental assays, this work provides mechanistic insight into how human SLC25A51 recognizes its ligand, how the transporter can be regulated by its lipid environment, and an observation of ligand-induced gate opening. This represents the first description of the ligand binding site for any NAD$^+$ transporter.

Introduction

SLC25A51 is an essential gene that is ubiquitously expressed across all mammalian cells. Its protein controls concentrations of NAD$^+$ in the mitochondrial matrix by directly importing the oxidized dinucleotide, and thus mammalian mitochondria depend on SLC25A51 for sustaining NAD$^+$ levels (1–3). Loss of SLC25A51 resulted in loss of cellular respiration and oxidative flux of the tricarboxylic acid cycle. SLC25A51 further impacted the concentrations of NAD-derived metabolites and cofactors, including NADH and NADPH (1–3).

There is little understood, nevertheless, about the underlying mechanisms of SLC25A51 or how the transporter interacts with oxidized NAD$^+$, compared to NADH or related molecules such as nicotinamide or the mononucleotide. SLC25A51 is a member of the mitochondrial carrier family (MCF) that localizes to the inner mitochondrial membrane (1–4). By homology, its structure harbors a pseudo-tri-symmetrical pore that is characteristic of the family (4–7). The trice repeated domain comprises two transmembrane helices joined by a short matrix helix parallel to the matrix membrane. Based on cytoplasmic and matrix-open structures of the related ADP/ATP nucleotide carriers, it is surmised that members of this structurally related family function through an alternating-access mechanism (5–11).

SLC25A51 has low conservation (<20%) in its amino acid sequence compared to its functional homologues in *S. cerevisiae* and *A. thaliana* that selectively import mitochondrial NAD$^+$ (*SI Appendix*, Fig. S1) (4, 12, 13). In phylogenetic analyses, SLC25A51 did not cluster with other mammalian nucleotide carriers in the family (7, 14). Thus, it remains unanswered how SLC25A51 engages and transports its ligand.

In this work we used Molecular Dynamics (MD) simulations to study how SLC25A51 homology models engage the NAD$^+$ ligand under the constraints of thermodynamics and in the context of a lipid environment. We tested our observations in intact cells with mutational analysis and measurements of in-situ free mitochondrial NAD$^+$ concentrations using a LigA-based NAD$^+$ biosensor (15). This study provides the first insight into the binding site of an NAD$^+$

transporter, identifies a mechanism of ligand-induced transport, and reveals cardiolipin-binding as a regulator of SLC25A51 activity.

Results

Stable modeling of SLC25A51 in a lipid bilayer.

To study how SLC25A51 recognizes NAD^+ for import, we first generated apo models of human SLC25A51 in its outward-facing, cytoplasmic (c-state) conformation using Swiss-Model and AlphaFold2 modeling [16, 17]. Both model structures adopted the characteristic MCF fold and were superimposable upon the solved bovine ANT crystal structure (PDB ID: 1OKC) [5]. Using MolProbity, we determined that the quality of the equilibrated AlphaFold2 and Swiss-Model structures were in the 100th and 97th percentile, respectively [18]. Model quality was further estimated using pLDDT for AlphaFold2 (*SI Appendix*, Fig. S2*A*) and QMEANDisCo analysis for both AlphaFold2 and Swiss-Model with global scores of 0.55 ± 0.05 and 0.56 ± 0.05, respectively (*SI Appendix*, Fig. S2*B and C*) [16, 19]. A direct comparison, nevertheless, revealed a difference of 3.5 Å root-mean-square deviation (RMSD) in atomic positions between the structures. Closer inspection revealed several residues in the central pore that were hydrophobically buried in one model but exposed to the hydrophilic pore in the other (*SI Appendix*, Fig. S2*D*). Therefore, we decided to test both models in this study.

Each of the apo models was individually embedded in a 86 x 86 Å^2 swatch of lipid bilayer using Charmm-GUI that recapitulated the composition of phospholipid components in mammalian inner mitochondrial membrane (POPC:POPE:CL=2:3:2) [20–24]. The system was hydrated and modeled with an ionic concentration of 150 mM KCl. Each model was analyzed in triplicate, and each replicate was ~1 µs in duration (*SI Appendix,* Table S1). We observed that the mean RMSD over time among the replicates for both stabilized c-state models were 4.1 ± 0.3 Å and 2.9 ± 0.2 Å for the Swiss-Model and AlphaFold2 models, respectively (*SI Appendix,* Fig. S3*A*, *B* and Table S1). In parallel, as a control for the approach, we generated a homology

model for c-state human SLC25A4 (ANT1, ADP/ATP carrier) based on the solved bovine structure (PDB ID: 1OKC) (5). The model for SLC25A4 (RMSD 2.8 Å) displayed a similar stability compared to the SLC25A51 c-state models over the course of the simulations (*SI Appendix,* Fig. S3C). We further calculated root-mean-square fluctuation (RMSF) values for all c-state models, and as expected, observed less flexibility in transmembrane regions compared to loop regions (*SI Appendix,* Fig. S3 *D-F*). Together, the data indicated that the experimental model systems were stable.

Cardiolipin was required for SLC25A51 activity.

With both the AlphaFold2 and Swiss-Model models, we observed spontaneous recruitment of cardiolipin molecules to three sites on SLC25A51 (Fig. 1*A* and Movie S1). Notably, all lipids were initially positioned at random and cardiolipin molecules were not docked to SLC25A51. We observed cardiolipin molecules arriving from different initial positions, displacing other types of phospholipids initially bound, and consistently limiting their interactions to the same three distinct sites on SLC25A51. Together, the data suggested site-specific recruitment of cardiolipin to SLC25A51.

Cardiolipin binding was mediated by polar interactions between its phosphates and exterior-facing residues on the transporter at Binding Site 1 (T65, R66, R270 and G271), Binding Site 2 (N156, T157, R82 and G83), and Binding Site 3 (S249, R174, G175) (Fig. 1*B*). The observed sites correlated in position with cardiolipin binding motifs [F/Y/W δ G] and [F/Y/W K/R G] identified from ADP/ATP transporter studies (5, 8, 9, 25–28). All binding sites were on the exterior of the central pore facing the matrix leaflet and positioned at the junctions between the three repeated protein domains that comprised SLC25A51's pore (Fig. 1*A* and *B*). Each cardiolipin molecule spanned an even-numbered helix from one domain and the matrix helix of the neighboring domain. At sites 1 and 3, cardiolipin bridged adjacent domains through extensive engagement of each of its phosphates (Fig. 1*B*). At Site 2, a single phosphate of the

cardiolipin molecule bridged the domains in 3 out of 4 replicates (Fig. 1*B*). This suggested at least two binding modalities for cardiolipin and that cardiolipin may engage SLC25A51 asymmetrically.

To determine whether the observed cardiolipin binding was required for the activity of SLC25A51, we expressed a genetically-encoded ratiometric fluorescent sensor for free NAD^+ in the mitochondria of HeLa cells (1, 15, 29). Fig. 1*C* depicts the inverted fluorescence intensity of the sensor that is a readout of relative NAD^+ levels. Acute ectopic expression of [Flag]SLC25A51 resulted in increased steady-state levels of free mitochondrial NAD^+ compared to empty vector, demonstrating that this assay could measure the activity of SLC25A51 of importing NAD^+ in intact and respiring cells (Fig. 1*C*). We introduced point mutations to impair SLC25A51's interaction with cardiolipin by blunting the electrostatic interactions at each binding sites (R82Q, R174Q, and R270Q). These sites were exterior to the transporter's pore, distinct from the active site, and not obviously involved in any interhelical interactions. Mutation of all three sites, resulted in loss of SLC25A51 activity (Fig. 1*C*), indicating that cardiolipin is likely required for SLC25A51 activity.

The matrix gate is comprised of a single salt bridge interaction

To study the control of SLC25A51 opening to the matrix, we examined all possible inter-helical interactions at the position where gated salt-bridge networks have been identified in the related MCF carriers (4, 6, 30). In SLC25A51, there were only two potential interhelical interactions: a putative hydrogen bond between Q52 and Q142, and a putative ionic salt-bridge between E139 and K236 (Fig. 2*A*). Time evolution of each interaction over 1000 ns in both AlphaFold2 and Swiss-Model models indicated that the hydrogen bond (Q52-Q142) was not formed (combined mean percent occupancy, m.p.o. 2% ± 0.7%) while the salt bridge interaction (E139-K236) was stable (combined m.p.o., 93% ± 3%) (Fig. 2*B* and *C*).

We used the mitochondrial NAD$^+$ sensor assay to test whether these interactions were required for SLC25A51 activity. Loss of the gate is predicted to favor the inward-facing, matrix (m-state) conformation, which would not be amenable to NAD$^+$ import. Expression of SLC25A51 variants with an ablated salt bridge (K236Q, K236A, E139Q, or E139A point mutations) were not able to increase steady-state levels of free mitochondrial NAD$^+$ levels compared to empty vector control (Fig. 2*D*). This indicated that formation of the E139-K236 salt bridge was required for SLC25A51 activity. In line with the simulation results, disruption of the putative hydrogen bond with Q52A or Q142A did not have any major impact on SLC25A51 activity (Fig. 2*D*). Furthermore, conserved residues in the consensus matrix gate motif (PX[D/E]XX[K/R]XXQ)—including L55 and N233 whose equivalents were identified as critical in the yeast and plant mitochondrial carriers (31)—were not required for SLC25A51 activity in intact cells (Fig. 2*D*). Together, the modeling and the biochemical assays indicated that a single interhelical salt-bridge between E139 and K236 formed the matrix gate.

Glutamine braces are not required for SLC25A51 activity

In many MCF carriers, glutamine braces play important roles in regulating the matrix gate (6). We observed an analogously positioned glutamine residue Q240 in SLC25A51 to putatively brace the identified E139-K236 salt-bridge (Fig. 2*A* and *B*). Modeling analyses was used to test the stability of either a Q240-E139 or Q240-K236 interaction. We did not observe consistent interactions in either model (Q240-E139 combined m.p.o. of 5% ± 3%; Q240-K236 combined m.p.o of 45% ± 16%) (Fig. 2*E*). In agreement, mutation of Q240A, or analogous glutamine residues Q59 and Q146 in adjacent domains, minimally impaired SLC25A51 activity (Fig. 2*D*). Together, our data demonstrated that the identified salt-bridge gate in SLC25A51 can function without stabilization from glutamine braces, distinctly unique from many other MCF carriers (6).

Arginine cap residues contribute to regulating SLC25A51 activity.

Cap residues at the matrix opening of the transporter have been proposed to help

stabilize the c-state of MCF carriers by neutralizing c-terminal negative dipoles of converging

odd-numbered helices or by forming stabilizing ionic interactions with matrix helices (32, 33). In

SLC25A51's modeling, we observed three arginine residues, R57, R238, and R140, positioned

to putatively serve as cap residues (*SI Appendix,* Fig. S4*A*). While mutation of all three residues

(R57Q, R140Q, and R238Q) destabilized the protein as observed by Western blot (*SI Appendix,*

Fig. S4*B*), individual mutations were better tolerated and each partially impaired SLC25A51

activity (*SI Appendix,* Fig. S4*C*). In agreement, we observed stable interactions over time

between R140 and the backbone of residue Q240 (m.p.o 95% ± 3% AlphaFold2 model, m.p.o.

80% ± 16% Swiss-Model), as well as between R238 and the backbone of residue Q59 (m.p.o.

70% ± 10%, Swiss-Model) (*SI Appendix,* Fig. S4*D-F*).

NAD⁺ import can occur independently of a stable cytoplasmic gate

SLC25A51 is predicted to function via an alternating-access mechanism similarly to

other MCF family members (4, 7). We thus generated an m-state model (MolProbity 94[th]

percentile)—using Swiss-Homology Modeling and the solved structure of *T. thermophila*

ADP/ATP carrier (PDB ID: 6GCI) (25)—to help identify the cytoplasmic gate. Performed in

triplicate (~1 µs / replicate, *SI Appendix,* Table S1), we observed consistent and stable

formation of a putative ionic gate between K198 and E291 (m.p.o. 95% ± 5%) in an analogous

plane to the identified cytoplasmic gate in the ADP/ATP carrier (*SI Appendix,* Fig. S5*A* and *B*).

This suggested that SLC25A51 has single salt bridge gates on each side of its pore. Mutation of

the cytoplasmic gate is expected to favor the c-state conformation. Interestingly, introduction of

double mutation K198Q, E291Q did not impair SLC25A51 activity (*SI Appendix,* Fig. S5*C*)

unlike previously observed for ADP/ATP carrier (11, 25). This indicated that a c-state

conformation was sufficient for importing NAD$^+$ and that a stable m-state was not required for continuous uptake of NAD$^+$.

Putative tyrosine brace interactions were also observed but with lowered percent occupancy (E103-Y290 m.p.o. 68% ± 24%; R194-E291 m.p.o. 63% ± 8.5%) (*SI Appendix,* Fig. S5*A* and *B*). In agreement with the idea that a stable m-state was not essential, many of the disruptive mutations had minimal effect on SLC25A51 activity (*SI Appendix,* Fig. S5*C*). The exception was that R194A and R194Y blocked SLC25A51 activity, however mutation of its binding partner E291 did not result in any impairment, so any contribution from R194 was likely independent of its brace function. Together, the data indicated that cytoplasmic gate and brace features were not critical to SLC25A51's import function.

NAD$^+$ is orientated by charge to bind at three conserved contact points

To study how SLC25A51 engages its ligand, we unbiasedly modeled the binding of NAD$^+$ onto both c-state models. We used the Autodock 4.2 algorithm (34, 35) to generate a series of poses ordered by calculated binding energy scores. We chose the poses with the lowest binding energy scores from each model to continue with analyses using MD simulations.

With both models (*SI Appendix,* Table S1 Simulations 10-18), we observed a prompt (within 100 ns) and consistent positioning of the nicotinamide ring (Fig. 3*A*, *B* and Movies S2, S3). The central ligand-binding pore of SLC25A51 is characterized by positive charges arising from inward facing residues K91, R182 and R278 on the even-numbered helices, and a distinct negatively-charged region from acidic residue E132 on helix 3 (Fig. 3*A*). We observed in both models that the positively-charged nicotinamide ring of oxidized NAD$^+$ consistently oriented away from the basic residues in the binding site and toward E132 (Fig. 3*A*, *B* and Movies S2, S3). Ablation of E132's charge impaired SLC25A51 activity, but a E132D variant retained function (Fig. 3*C*). E132 is conserved across SLC25A51 orthologs (*SI Appendix,* Fig. S1*A*) (4).

Following positioning of the nicotinamide ring, we observed the nicotinamide ribose (NR) moiety to form the following consistent contacts (in 8 out of 9 simulations with both Swiss-Model and AlphaFold2 models) (*SI Appendix,* Table S1). Borrowing from the nomenclature used to describe the ADP/ATP binding pocket (7, 36, 37), we observed that residue T94 in contact site 1 and N183 in contact site 2 stabilized the NR moiety through specific hydrogen bonds with its carboxamide group and ribose (2' and 3' OH) respectively (Fig. 3*D*). R278 (contact site 3), R182, and K91 engaged the negative charges from the phosphates of NAD$^+$ (Fig. 3*D* and Movies S2, S3).

We mutated and tested the requirement of every conserved residue in the three contact sites using the sensor assay (Fig. 3*E* and *SI Appendix,* Fig. S1) (4, 6). For contact site 1, we found that SLC25A51 variants with a T94V or K91Q mutation impaired SLC25A51 import with minimal effects on protein stability. Mutating adjacent contact site residues (Q90 and M98) that were not observed to participate in binding of the NR moiety minimally impaired SLC25A51 despite being conserved residues (Fig. 3*E*). Similarly, we confirmed that R182 and N183 were required for site 2 but not neighboring residue S186, and we confirmed that R278 was required for site 3 (Fig. 3*E*).

NAD$^+$ directly engaged with the gate in SLC25A51

While modeling the binding of NAD$^+$, we unexpectedly observed two simulations wherein the ligand traversed deeper into the pore (Movie S4 and *SI Appendix,* Table S1 simulations 16 and 17). To determine whether an extended simulation could provide additional insights, we continued with a pose that originated from the unbiased docking with the Swiss-Homology model but had scored among lowest binding energy scores (*SI Appendix,* Table S1 simulations 19-22). This pose positioned the nicotinamide ring deeper in the pocket.

In all four replicate simulations 19-22, we observed channeling of the nicotinamide ring from E132 to E139 (Fig. 4*A*, *B* and Movie S4). In both model structures residue E139, of the

salt-bridge gate, was positioned below E132 on the same helix 3. We also observed that as the

nicotinamide ring interacted with E139 (85%, 48%, 69% and 93% occupancy), the phosphates

on NAD$^+$ could interact with K236, the other half of the salt bridge. Resultingly, formation of the

E139-K236 salt bridge was negatively correlated with the NAD$^+$ phosphate-K236 interaction ($r^2 =$

-0.192, -0.045, -0.681 and -0.336 per replicate, $p < 0.001$), as well as with the nicotinamide ring

-E139 interaction ($r^2 =$ -0.298, -0.053, -0.091 and -0.342 calculated per replicate, $p < 0.001$) (Fig.

4C, D and Movie S5).

Discussion

The molecular characterization of mitochondrial carriers has been a persistent challenge

with crystallographic structures currently available only for the bovine and fungal ADP/ATP

carriers (5, 8, 9, 25, 38). SLC25A51's lack of close homology to the previously identified yeast

and plant NAD$^+$ transporters delayed its de-orphanization as a mammalian mitochondrial NAD$^+$

transporter (*SI Appendix,* Fig. S1). Using a combination of modeling, simulations, and

biochemical assays, we have gained insight into SLC25A51's binding of NAD$^+$, its dependency

on cardiolipin for function, and a ligand induced mechanism that disrupted the matrix gate.

A unique aspect in NAD$^+$'s binding to SLC25A51 was the initial orientation of its oxidized

nicotinamide ring via electrostatic interactions. We observed that the delocalized positive charge

on the ring was drawn to the negative surface on helix 3 formed by E132 and simultaneously

repelled from basic residues K91, R182, and R278 in the same plane (Fig. 3A, Movie S2 and

S3). This positioned NAD$^+$ such that the nicotinamide ribose moiety engaged contact sites 1 and

2 consistently, and the negative charges on NAD$^+$'s phosphate groups were satisfied and

engaged with the positively charged sidechains in the pore (e.g., R278 in contact site 3) (Fig.

3D). In addition to the negative charge on E132, we confirmed that the residues comprising the

identified contact sites were required for SLC25A51 activity (Fig. 3C and E). The data was

consistent with previous docking and mutational analyses (2, 4). We observed that all three

contact sites in SLC25A51's pore must be engaged to orient the ligand such that it can travel deeper into the pore. This may explain why nicotinamide, or its mononucleotide were unable to efficiently compete with NAD$^+$ for transport (1, 2).

Positioned below NAD$^+$'s initial binding site in SLC25A51 was a single interhelical salt-bridge that embodied the matrix gate (Fig. 2). This salt-bridge interaction was the only consistent interhelical bond formed within SLC25A51's PX[D/E]XX[K/R] motifs, and it was required for activity (Fig. 2*D*) (6). Following the initial binding of NAD$^+$, the nicotinamide ring was channeled to the residue E139, which is located directly below E132 and comprises half of the matrix gate (Movie S4). The negatively charged phosphate groups on NAD$^+$ in turn engaged the other half of the gate, residue K236, and together this disrupted an otherwise stable gate interaction (Fig. 4 and Movie S5). However, it will still need to be determined whether this interaction can proceed to an occluded state or to the transport of the molecule. An analogous mechanism was proposed for the ADP/ATP carrier involving interference of the gate via the N6 atom of the adenine moiety on the nucleotide or the guanidino NH1 or NH2 of a conserved binding site arginine residue (36). We found additional features in SLC25A51 such as an arginine-cap that functioned to further stabilize the matrix gate (*SI Appendix*, Fig. S4), but we also identified that the c-state was sufficient for continuous import of NAD$^+$ when the cytoplasmic gate was weakened indicating that its relatively weak gate could be readily opened as suggested previously (*SI Appendix*, Fig. S5) (7).

NAD$^+$'s interaction with the matrix gate depended on the positive charge of its nicotinamide ring being positioned close to the acidic residue of the gate. This implies that without this positive charge, even with ideal positioning of the ring, the transport process may be halted. This model is consistent with the electrostatic funneling observed with ADP/ATP carriers (39–44). Moreover, previous studies have found that AMP, dUTP, NADH, GDP and GTP could bind to the ADP/ATP carrier, suggesting that the dynamics of transport in cells involves an additional differentiation step to discriminate between bound molecules (45, 46). Our substrate-

induced model may explain why NAD$^+$ can be transported at micromolar concentrations but that NADH can compete for SLC25A51 uptake at millimolar concentrations (2). Apart from an oxidized nicotinamide ring, NADH is structurally identical and likely can engage all binding sites in SLC25A51. Nevertheless, cytosolic ratios—which due to non-discriminating porin channels in the outer membrane—represent what is experienced by SLC25A51 in cells. Free NADH is approximately 100 to 1000-fold less abundant than free NAD$^+$ (47–50). In this scenario the relatively subtle E132 may be enough to create a molecular discrimination based on electrostatic charge. Additionally, the overall charge of an NADH or an NAD$^+$ molecule would be differentially masked in the positive pore of SLC25A51 (6). With an extra positive charge from its nicotinamide ring, NAD$^+$ bound to SLC25A51 retains a net positive charge in the binding site is likely to be favored by the negative potential in the matrix, compared to the neutral charge of bound NADH in the same site (6).

To transport a ligand such as NAD$^+$ with an asymmetric structure, one may either expect the pore to be matched in asymmetrical shape or to have flexibility. We found evidence suggesting that SLC25A51 required cardiolipin for its activity (Fig. 1). Cardiolipin has been found to support MCF activities in a variety of ways (28, 51–53). The data supporting cardiolipin recruitment at specific sites was consistent and collected unbiasedly. We observed two modalities of cardiolipin binding, and site 2 cardiolipin tended to bind non-canonically using only one of its phosphates. A potential consequence is that this would asymmetrically impact the pore in a dynamic manner that is potentially regulatable.

Overall, this study provided unique insights into the binding and transport mechanism of an NAD$^+$ carrier. It is a launching point for future characterization and structural studies to answer remaining questions such as what controls the directionality of NAD$^+$ import, whether SLC25A51 requires an antiport molecule akin to other carriers, and to elucidate the full dynamics of the SLC25A51 transporter. A structural understanding will accelerate the development of modulators targeting SLC25A51 in various diseases.

Material and Methods

Detailed material and methods are provided in the supplementary file.

Competing Interests

The authors declare no competing interests

Acknowledgements

We thank Akhilesh Paspureddi and Dr. Ron Elber for valuable discussions and help with simulation analyses throughout this study. Supported by the Texas Advanced Computing Center, NIH DP2 GM126897, the Pew Charitable Trust, and CPRIT RP210079.

References

1. T. S. Luongo, *et al.*, SLC25A51 is a mammalian mitochondrial NAD+ transporter. *Nature* **588**, 174–179 (2020).

2. N. Kory, *et al.*, MCART1/SLC25A51 is required for mitochondrial NAD transport. *Sci. Adv.* **6** (2020).

3. E. Girardi, *et al.*, Epistasis-driven identification of SLC25A51 as a regulator of human mitochondrial NAD import. *Nat. Commun. 2020 111* **11**, 1–9 (2020).

4. M. Ziegler, *et al.*, Welcome to the Family: Identification of the NAD+ Transporter of Animal Mitochondria as Member of the Solute Carrier Family SLC25. *Biomol. 2021, Vol. 11, Page 880* **11**, 880 (2021).

5. E. Pebay-Peyroula, *et al.*, Structure of mitochondrial ADP/ATP carrier in complex with carboxyatractyloside. *Nat. 2003 4266962* **426**, 39–44 (2003).

6. J. J. Ruprecht, E. R. S. Kunji, Structural Mechanism of Transport of Mitochondrial Carriers. *Annu. Rev. Biochem.* **90**, 535–558 (2021).

7. A. J. Robinson, C. Overy, E. R. S. Kunji, The mechanism of transport by mitochondrial carriers based on analysis of symmetry. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 17766–17771 (2008).

8. J. J. Ruprecht, *et al.*, Structures of yeast mitochondrial ADP/ATP carriers support a domain-based alternating-access transport mechanism. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E426–E434 (2014).

9. H. Nury, *et al.*, Structural basis for lipid-mediated interactions between mitochondrial ADP/ATP carrier monomers. *FEBS Lett.* **579**, 6031–6036 (2005).

10. R. Springett, M. S. King, P. G. Crichton, E. R. S. Kunji, Modelling the free energy profile of the mitochondrial ADP/ATP carrier. *Biochim. Biophys. Acta - Bioenerg.* **1858**, 906–914 (2017).

11. M. S. King, M. Kerr, P. G. Crichton, R. Springett, E. R. S. Kunji, Formation of a

cytoplasmic salt bridge network in the matrix state is a fundamental step in the transport mechanism of the mitochondrial ADP/ATP carrier. **1857**, 14–22 (2016).

12. S. Todisco, G. Agrimi, A. Castegna, F. Palmieri, Identification of the Mitochondrial NAD+ Transporter in Saccharomyces cerevisiae*. *J. Biol. Chem.* **281**, 1524–1531 (2006).

13. F. Palmieri, *et al.*, Molecular identification and functional characterization of Arabidopsis thaliana mitochondrial and chloroplastic NAD+ carrier proteins. *J. Biol. Chem.* **284**, 31249–31259 (2009).

14. F. Palmieri, The mitochondrial transporter family SLC25: Identification, properties and physiopathology. *Mol. Aspects Med.* **34**, 465–484 (2013).

15. X. A. Cambronne, *et al.*, Biosensor reveals multiple sources for mitochondrial NAD+. *Science (80-.).* **352**, 1474–1477 (2016).

16. J. Jumper, *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nat. 2021 5967873* **596**, 583–589 (2021).

17. A. Waterhouse, *et al.*, SWISS-MODEL: Homology modelling of protein structures and complexes. *Nucleic Acids Res.* **46**, W296–W303 (2018).

18. C. J. Williams, *et al.*, MolProbity: More and better reference data for improved all-atom structure validation. *Protein Sci.* **27**, 293–315 (2018).

19. G. Studer, *et al.*, QMEANDisCo-distance constraints applied on model quality estimation. *Bioinformatics* **36**, 1765–1771 (2020).

20. E. M. Mejia, G. M. Hatch, Mitochondrial phospholipids: role in mitochondrial function. *J. Bioenerg. Biomembr. 2015 482* **48**, 99–112 (2015).

21. J. Comte, B. Maïsterrena, D. C. Gautheron, Lipid composition and protein profiles of outer and inner membranes from pig heart mitochondria. Comparison with microsomes. *BBA - Biomembr.* **419**, 271–284 (1976).

22. S. Jo, T. Kim, V. G. Iyer, W. Im, CHARMM-GUI: A web-based graphical user interface for CHARMM. *J. Comput. Chem.* **29**, 1859–1865 (2008).

23. J. Lee, *et al.*, CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. *J. Chem. Theory Comput.* **12**, 405–413 (2016).

24. E. L. Wu, *et al.*, CHARMM-GUI membrane builder toward realistic biological membrane simulations. *J. Comput. Chem.* **35**, 1997–2004 (2014).

25. J. J. Ruprecht, *et al.*, The Molecular Mechanism of Transport by the Mitochondrial ADP/ATP Carrier. *Cell* **176**, 435-447.e15 (2019).

26. A. L. Duncan, J. J. Ruprecht, E. R. S. Kunji, A. J. Robinson, Cardiolipin dynamics and binding to conserved residues in the mitochondrial ADP/ATP carrier. *Biochim. Biophys. Acta - Biomembr.* **1860**, 1035–1045 (2018).

27. P. G. Crichton, *et al.*, Trends in thermostability provide information on the nature of substrate, inhibitor, and lipid interactions with mitochondrial carriers. *J. Biol. Chem.* **290**, 8206–8217 (2015).

28. N. Senoo, *et al.*, Cardiolipin, conformation, and respiratory complex-dependent oligomerization of the major mitochondrial ADP/ATP carrier in yeast. *Sci. Adv.* **6**, 780–808 (2020).

29. J. M. Eller, *et al.*, Flow Cytometry Analysis of Free Intracellular NAD+ Using a Targeted Biosensor. *Curr. Protoc. Cytom.* **88**, e54 (2019).

30. D. R. Nelson, C. M. Felix, J. M. Swanson, Highly conserved charge-pair networks in the mitochondrial carrier family. *J. Mol. Biol.* **277**, 285–308 (1998).

31. D. V. Miniero, M. Monné, M. A. Di Noia, L. Palmieri, F. Palmieri, Evidence for Non-Essential Salt Bridges in the M-Gates of Mitochondrial Carrier Proteins. *Int. J. Mol. Sci. 2022, Vol. 23, Page 5060* **23**, 5060 (2022).

32. Q. Yi, *et al.*, Molecular dynamics simulations on apo ADP/ATP carrier shed new lights on the featured motif of the mitochondrial carriers. *Mitochondrion* **47**, 94–102 (2019).

33. C. L. Pierri, F. Palmieri, A. De Grassi, Single-nucleotide evolution quantifies the

importance of each site along the structure of mitochondrial carriers. *Cell. Mol. Life Sci.* **71**, 349–364 (2014).

34.   S. Forli, *et al.*, Computational protein–ligand docking and virtual drug screening with the AutoDock suite. *Nat. Protoc. 2016 115* **11**, 905–919 (2016).

35.   G. M. Morris, *et al.*, Software news and updates AutoDock4 and AutoDockTools4: Automated docking with selective receptor flexibility. *J. Comput. Chem.* **30**, 2785–2791 (2009).

36.   E. R. S. Kunji, A. J. Robinson, The conserved substrate binding site of mitochondrial carriers. *Biochim. Biophys. Acta - Bioenerg.* **1757**, 1237–1248 (2006).

37.   A. J. Robinson, E. R. S. Kunji, Mitochondrial carriers in the cytoplasmic state have a common substrate binding site. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 2617–2622 (2006).

38.   E. R. S. Kunji, M. Harding, Projection structure of the atractyloside-inhibited mitochondrial ADP/ATP carrier of Saccharomyces cerevisiae. *J. Biol. Chem.* **278**, 36985–36988 (2003).

39.   Y. Wang, E. Tajkhorshid, Electrostatic funneling of substrate in mitochondrial inner membrane carriers. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 9598–9603 (2008).

40.   D. Heidkämper, V. Müller, D. R. Nelson, M. Klingenberg, Probing the role of positive residues in the ADP/ATP carrier from yeast. The effect of six arginine mutations on transport and the four ATP versus ADP exchange modes. *Biochemistry* **35**, 16144–16152 (1996).

41.   M. Monné, F. Palmieri, E. R. S. Kunji, The substrate specificity of mitochondrial carriers: Mutagenesis revisited. *http://dx.doi.org/10.3109/09687688.2012.737936* **30**, 149–159 (2013).

42.   E. M. Krammer, *et al.*, High-Chloride Concentrations Abolish the Binding of Adenine Nucleotides in the Mitochondrial ADP/ATP Carrier Family. *Biophys. J.* **97**, L25–L27 (2009).

43.   F. Dehez, E. Pebay-Peyroula, C. Chipot, Binding of ADP in the mitochondrial ADP/ATP

carrier is driven by an electrostatic funnel. *J. Am. Chem. Soc.* **130**, 12725–12733 (2008).

44. V. Mavridou, *et al.*, Substrate binding in the mitochondrial ADP/ATP carrier is a step-wise process guiding the structural changes in the transport cycle. *Nat. Commun. 2022 131* **13**, 1–12 (2022).

45. H. Majd, *et al.*, Screening of candidate substrates and coupling ions of transporters by thermostability shift assays. *Elife* **7** (2018).

46. J. Mifsud, *et al.*, The substrate specificity of the human ADP/ATP carrier AAC1. *http://dx.doi.org/10.3109/09687688.2012.745175* **30**, 160–168 (2013).

47. D. H. Williamson, P. Lund, H. A. Krebs, "The Redox State of Free Nicotinamide-Adenine Dinucleotide in the Cytoplasm and Mitochondria of Rat Liver" (1967).

48. Q. Zhang, D. W. Piston, R. H. Goodman, Regulation of corepressor function by nuclear NADH. *Science (80-.).* **295**, 1895–1897 (2002).

49. Y. P. Hung, J. G. Albeck, M. Tantama, G. Yellen, Imaging cytosolic NADH-NAD + redox state with a genetically encoded fluorescent biosensor. *Cell Metab.* **14**, 545–554 (2011).

50. Y. Zhao, *et al.*, Genetically encoded fluorescent sensors for intracellular NADH detection. *Cell Metab.* **14**, 555–566 (2011).

51. M. Schlame, Protein crowding in the inner mitochondrial membrane. *Biochim. Biophys. Acta - Bioenerg.* **1862**, 148305 (2021).

52. Y. Lee, C. Willers, E. R. S. Kunji, P. G. Crichton, Uncoupling protein 1 binds one nucleotide per monomer and is stabilized by tightly bound cardiolipin. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 6973–6978 (2015).

53. S. Ghosh, *et al.*, An essential role for cardiolipin in the stability and function of the mitochondrial calcium uniporter. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 16383–16390 (2020).

Figure Legends

Fig. 1. Cardiolipin binding impacts SLC25A51 function.

(A) Observed positioning of bound cardiolipin (wheat). Differential hues in SLC25A51 delineate the three repeated domains that comprise its structure.

(B) Representative poses depicting identified interactions at sites 1-3. Yellow dashes indicate interactions < 3.5 Å. At site 2, the binding of a single cardiolipin phosphate was often observed.

(C) (left) Relative concentrations of free mitochondrial NAD^+ measured using a ratiometric biosensor and ectopic expression of either Control empty vector, Wildtype [Flag]SLC25A51, or mutant [Flag]SLC25A51 harboring R82Q, R174Q, R270Q point mutations. Dashed horizontal line indicates baseline for unchanged ratiometric measurements at 1; mean ± SD, n = 4, ANOVA < 0.01 with post-hoc Dunnett's Test *p < 0.001. (right)* Anti-Flag Western Blot of HeLa cells expressing indicated SLC25A51 variants or empty vector; β-actin, loading control.

Fig. 2. A single salt bridge stabilizes the NAD^+-accessible, outward facing conformation.

(A) Graphical representation of the matrix gate with dash lines indicating possible polar non-covalent interactions. Interactions were either indicated (red) or unsupported (grey) by data.

(B) Relative position of the matrix gate in a cartoon representation of SLC25A51; tested glutamine brace residues (white) and interactions (dashed lines).

(C) Time evolution of distance between indicated side chains for putative gate interactions. Interaction cutoff of 3.5 Å, dashed line.

(D) Free mitochondrial NAD^+ levels measured using a ratiometric biosensor in HeLa cells expressing Control empty vector (n=25), Wildtype [Flag]SLC25A51 (n=25), and indicated mutants. Dashed horizontal line indicates baseline at 1. Data is mean ± SD, n = 4 - 6,

ANOVA $p < 0.001$ with post-hoc Dunnett's test *$p < 0.05$, **$p < 0.01$. *(bottom)* Anti-Flag

Western Blot from HeLa whole cell lysates express either empty vector or indicated

FlagSLC25A51 variants; β-actin, loading control.

(E) Time evolution of distance between indicated side chains of putative brace interactions.

Interaction cutoff at 3.5 Å, dashed line.

Fig. 3. E132 and conserved contact sites comprise an NAD$^+$ binding site

(A) Electrostatic environment of the substrate binding site (APBS as Pymol plugin).

Representative positioning of NAD$^+$ from simulation 17 at 1230 ns (*SI Appendix*, Table S1).

(B) Minimal distance over time between the nicotinamide ring and E132 in simulations 17

(Swiss-Model) and 12 (AlphaFold2) (*SI Appendix*, Table S1). Interaction cutoff at 4 Å,

dashed horizontal line.

(C) Ratiometric sensor measurements of free mitochondrial NAD$^+$ levels for FlagSLC25A51

mutants E132D (n=4) and E132A (n=4) expressed in HeLa cells; Control empty vector (n=8)

and Wildtype FlagSLC25A51 (n=8). Dashed horizontal line indicates baseline at 1. Data is

mean ± SD, ANOVA $p < 0.001$ with post-hoc Dunnett's test *$p < 0.001$. *(bottom)* Anti-Flag

Western Blot; β-actin, loading control.

(D) Bound NAD$^+$ (orange) in the ligand binding pocket is shown in both side and enlarged views.

Non-covalent interactions < 3.5 Å, dashed yellow lines.

(E) Free mitochondrial NAD$^+$ levels measured using a ratiometric biosensor in HeLa cells that

expressed Control empty vector, Wildtype FlagSLC25A51, or mutant variants as indicated.

Dashed horizontal line indicates baseline at 1; data is mean ± SD, n = 4-7, unless otherwise

indicated, ANOVA $p < 0.001$, post-hoc Dunnett's test *$p < 0.001$. *(bottom)* Anti-Flag Western

Blot; β-actin, loading control.

Fig. 4. Substrate induced disruption of the matrix gate.

(A) Minimum distance between the nicotinamide ring and E139 over time, simulation 17 (*SI Appendix*, Table S1).

(B) 2-dimentional projection (Ligplot+) of NAD$^+$ and its interaction with E132 and the E139-K236 salt bridge.

(C) Time evolutions of the E139-K236 salt bridge (teal) compared to the nicotinamide ring and E139 interaction (red); simulation 19 (*SI Appendix*, Table S1)

(D) Time evolutions of the E139-K236 salt bridge (teal) compared to the phosphates in NAD$^+$ and K236 (grey); simulation 19 (*SI Appendix*, Table S1)

Figure 1

Figure 2

Figure 3

Figure 4