Pairing Metagenomics and Metaproteomics to Characterize Ecological Niches and Metabolic Essentiality of gut microbiomes

Short running title: Proteins' ecological and metabolic roles

Tong Wang[1,†], Leyuan Li[2,3,†], Daniel Figeys[3,*], Yang-Yu Liu[1,4*]

[1]*Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA 02115, USA.*
[2]*China Key Laboratory of Medical Proteomics, Beijing Proteome Research Center, National Center for Protein Sciences (Beijing), Beijing Institute of Lifeomics, Beijing 102206, China.*
[3]*School of Pharmaceutical Sciences, Ottawa Institute of Systems Biology, Faculty of Medicine, University of Ottawa, Ottawa, ON K1H8M5, Canada.*
[4]*Center for Artificial Intelligence and Modeling, the Carl R. Woese Institute of Genomic Biology, University of Illinois at Urbana-Champaign, Champaign, IL 61820, USA.*

† These authors contributed equally to this work.
* Correspondence should be addressed to Y.-Y.L. (yyl@channing.harvard.edu) or D.F. (dfigeys@uottawa.ca). The physical mailing address of Y.-Y.L. is: Yang-Yu Liu, Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, 181 Longwood Ave, Boston, MA 02115, USA. The physical mailing address of D.F. is: Daniel Figeys, School of Pharmaceutical Sciences, Ottawa Institute of Systems Biology, Faculty of Medicine, University of Ottawa, 451 Smyth Rd, Ottawa, ON K1H8M5, Canada

Abstract

The genome of a microorganism encodes its potential functions that can be implemented through expressed proteins. It remains elusive how a protein's selective expression depends on its metabolic essentiality to microbial growth or its ability to claim resources as ecological niches. To reveal a protein's metabolic or ecological role, we developed a computational pipeline, which pairs metagenomics and metaproteomics data to quantify each protein's gene-level and protein-level functional redundancy simultaneously. We first illustrated the idea behind the pipeline using simulated data of a consumer-resource model. We then validated it using real data from human and mouse gut microbiome samples. In particular, we analyzed ABC-type transporters and ribosomal proteins, confirming that the metabolic and ecological roles predicted by our pipeline agree well with prior knowledge. Finally, we performed *in vitro*

cultures of a human gut microbiome sample and investigated how oversupplying various sugars involved in ecological niches influences the community structure and protein abundance. The presented results demonstrate the performance of our pipeline in identifying proteins' metabolic and ecological roles, as well as its potential to help us design nutrient interventions to modulate the human microbiome.

Keywords

Metagenomics; Metaproteomics; Functional Redundancy; Ecological Niche; Metabolic Essentiality; Gut Microbiome

Introduction

Metagenomic sequencing has enabled the measurement of the genomic content and functional potential of microbial communities at an unprecedented rate, aiding the understanding of their role in host health[1–3] and biogeochemical cycling[4–6]. While various computational approaches based on these genomes quantify interactions within microbial communities[7–12] and analyze the functional redundancy and functional stability of microbial communities[10–12], they focus on potential rather than actual function, as microorganisms only express a subset of genes as proteins[13]. Recent advancements in high-throughput metaproteomics allow us to quantify protein abundances in human gut microbiomes[14], offering insights into gene expression in response to environmental changes when paired with metagenomic data.

From the metabolic perspective, some genes and their encoded proteins are indispensable for cell metabolism under any conditions, and microbial growth halts without these essential functions—aminoacyl-tRNA synthetase[15,16], ribosomal proteins[17–19], and enzymes involved in glycolysis[20,21]. From the ecological perspective, gene expression is influenced by ecological selection, with specific proteins indicating which resources a microbe can utilize and defining its ecological niche. For instance, *E. coli* prefers glucose over lactose due to the repressed expression of lactose-utilizing enzymes, even though it can use both sugars[22,23]. Such specialization of consuming one resource caused by the selective gene expression may reduce the niche overlap with other species and allow microbial coexistence,

as seen with two *E. coli* strains where one expresses acetyl-coenzyme synthetase (Acs)[24,25] to

consume acetate produced by the other[26–29].

Understanding the selective expression of microbial genes is an outstanding question

in microbiology. Does the behavior of selective expression of microbial genes differ between

metabolic function (e.g., essential for microbial growth metabolism) and ecological function

(e.g., claiming resources as a niche)? To answer this, we developed a computational method

to analyze paired metagenomic and metaproteomic[30–33,14] data, constructing the gene content

network (GCN) or protein content network (PCN) --- a bipartite graph that connects microbial

taxa to their genes or expressed proteins, respectively (Figs. 1a&b). For each gene and its

encoding protein, we compared its gene-level (or protein-level) functional redundancy (FR),

revealing each protein family's metabolic or ecological role. Our method, validated with several

gut microbiome data, accurately predicts that ABC-type transporters are related to ecological

niches[34–36] , and ribosomal proteins are essential[17–19]. Finally, we performed *in vitro* culture

experiments using human gut microbiome samples to investigate how oversupplying sugars

involved in ecological niches influence community structure and protein expression.

Materials and Methods

***In-vitro* human gut microbiota culture and metaproteomics.** Three healthy individual

microbiota samples were collected and biobanked. The frozen microbiome samples were

cultured in our optimized culture medium[37] with or without the presence of different sugars in

technical triplicates, and were taken at different times for optical density and metaproteomic

analyses. For single-strain samples, proteins were extracted with 4% SDS 8M urea buffer in

100 mM Tris-HCl buffer, followed by precipitation and acetone washing. Proteins were digested

with trypsin desalted[38] for LC-MS/MS analysis using an Orbitrap Exploris 480 mass

spectrometer. For the cultured microbiomes, an automated process extracted and purified

proteins, which were then digested, desalted, and quantified using TMT11plex[39], ensuring

mixed representation in labeling to avoid bias. Samples underwent a 2-hour LC gradient and

were analyzed by mass spectrometry. More details can be found in Supplemental Methods.

Datasets. Metagenomics data from four individual microbiomes were obtained from the

previous MetaPro-IQ study[14,33] (accessible from the NCBI sequence read archive (SRA) under

the accession of SRP068619) and the same samples were reanalyzed by an ultra-deep

metaproteomics approach[14] via the PRIDE partner repository[40] with the dataset identifier

PXD027297). Proteomics dataset of the cultured singles strain samples has been deposited to

ProteomeXchange Consortium with the identifier PXD037923. Metaproteomic dataset of the

RapidAIM-cultured microbiome samples has been deposited to ProteomeXchange Consortium

(identifier PXD037925). The metaproteomic dataset of the mouse gut microbiome comprising

twenty gut microbes is derived from a previous study[41] that was deposited to

ProteomeXchange Consortium with the dataset identifier PXD009535 and to MassIVE with the

dataset identifier MSV000082287.

Database search and data processing. Proteomics database searches used FASTA

databases of the individual strains downloaded from NCBI and MaxQuant[42] 1.6.17.0 for

analysis, without the label-free quantification. Metaproteomic database searches of cultured

microbiome samples were performed using MetaLab V2.2[43], MaxQuant option was used to

search the TMT dataset against the IGC database of the human gut microbiome. The resulting

data table was normalized using R package MSstatsTMT , and missing values were imputed

using R package DreamAI[45]. The "fraction" of each taxon-specific protein is computed by

dividing the protein intensity by the sum of the intensities of all proteins assigned to the same

taxon. The log2 fold change of each protein is obtained by taking log2 of the ratio between its

fraction in the treatment group (with added sugars) and its fraction in the control group (without

added sugars).

Statistics. To calculate correlation throughout the study, we used Pearson's correlation

coefficient. All statistical tests were performed using standard numerical and scientific

computing libraries in the Python programming language (version 3.7.1) and Jupyter Notebook

(version 6.1).

Results

Specialist function, niche function, and essential function

Here, we define three types of functions for protein families that we would like to categorize: (1) "*Specialist function*": specialized by only a few taxa and not widely shared within a community. (2) "*Niche function*": arising from ecological competition, widespread among genomes of numerous taxa but selectively expressed under specific ecological conditions. (3) "*Essential function*": metabolically indispensable for and widely shared by many taxa within a microbial community. We emphasize that our definition is not exhaustive; some proteins may display attributes of multiple categories or not align precisely with any single category.

Using a simple hypothetical example of two competing species (Fig. 1a, b), we demonstrated the three function types: (1) The blue protein is a specialist function since it is solely encoded in the pink species' genome; (2)The red protein belongs to a niche function due to its selective expression by the yellow species even though the protein is encoded in the genomes of both species; (3) The green protein is an essential function because both species need it for biomass synthesis. In coexistence, the pink species specializes in the blue resource, avoiding competition with the yellow species for the red resource.

GCN, PCN, and network degree

We can identify the functional types of proteins in this hypothetical case by comparing the structure of the GCN and PCN (Fig. 1a, b). For example, consider the protein responsible for converting red resource to green metabolite (red broken circle in Fig. 1a, b), its degree in the GCN $k_{GCN} = 2$, while its degree in the PCN $k_{PCN} = 1$. This degree reduction is due to distinct ecological niches being occupied by two species when they are cocultured. By contrast, the protein responsible for assimilating critical green metabolites (green broken circle in Fig. 1a, b) into biomass does not show a degree reduction ($k_{GCN} = k_{PCN} = 2$) because it is essential for microbial growth. Similarly, since the blue protein is only specialized by the pink species, its $k_{GCN} = k_{PCN} = 1$. Thus, three function types occupy different regions in the k_{GCN} vs. k_{PCN} plot (Fig. 1c).

Quantifying gene- and protein-level functional redundancy of each gene and its encoded protein

However, the network degree does not consider the significant impact of the microbial taxonomic profile, which details the makeup of a microbial community. This profile is represented by $\boldsymbol{p} = (p_1, \ldots, p_N)$, where p_i is the relative abundance of taxon-i and $\sum_{i=1}^{N} p_i = 1$. For a given gene and its encoded protein, we can define its gene-level FR ($\mathrm{FR_g}$) and protein-level FR ($\mathrm{FR_p}$) within this sample as

$$\mathrm{FR_g} = \sum_{i=1}^{N} \sum_{j \neq i}^{N} \left(1 - d_{ij}^{\mathrm{GCN}}\right) p_i p_j, \tag{1}$$

and

$$\mathrm{FR_p} = \sum_{i=1}^{N} \sum_{j \neq i}^{N} \left(1 - d_{ij}^{\mathrm{PCN}}\right) p_i p_j. \tag{2}$$

d_{ij}^{GCN} (or d_{ij}^{PCN}) is the distance between taxon-i and taxon-j based on their genomic capacity to express this gene (or the presence of the protein). For simplicity, we assume d_{ij}^{GCN} is binary, i.e., $d_{ij}^{\mathrm{GCN}} = 0$ if and only if both taxa share the potential to express the gene, and $d_{ij}^{\mathrm{GCN}} = 1$ otherwise. $d_{ij}^{\mathrm{PCN}} = 0$ if and only if both taxa have expressed the protein. Here, we define $\mathrm{FR_g}$ and $\mathrm{FR_p}$ for each protein, different from our previous studies where FR was calculated by including all genes or proteins in a microbial community[12,14].

Comparing $\mathrm{FR_g}$ and $\mathrm{FR_p}$ provides deeper insight into proteins' function types. For the red protein in our hypothetical example, $d_{12}^{\mathrm{GCN}} = 0$ and $d_{12}^{\mathrm{PCN}} = 1$ because both species share the potential to express the gene while only the yellow species have expressed it (Fig. 1a, b). As a result, $\mathrm{FR_g} = 2(1-0)p_1 p_2 = 2p_1 p_2$ and $\mathrm{FR_p} = 2(1-1)p_1 p_2 = 0$ (Fig. 1d, e). Following the same analysis, $\mathrm{FR_g} = \mathrm{FR_p} = 2p_1 p_2$ for the green protein, and $\mathrm{FR_g} = \mathrm{FR_p} = 0$ for the blue protein (Fig. 1d, e). Different from composition-independent $\mathrm{k_{GCN}}$ and $\mathrm{k_{PCN}}$, $\mathrm{FR_g}$ and $\mathrm{FR_p}$ take the microbial composition into account and thus are more ecologically meaningful. Notably, a more uneven abundance distribution would lead to smaller $\mathrm{FR_g}$ and $\mathrm{FR_p}$ ($p_1 = p_2 = 0.5$ in Fig. 1d; $p_1 = 0.9$ and $p_2 = 0.1$ in Fig. 1e). The influence of relative abundances on FR can be mitigated by using the normalized FR: nFR = FR / TD, where TD $= 1 - \sum_i p_i^2$ (Supplementary Fig. 1; see Supplementary Information for the definition).

Overview of our computational pipeline

Following the idea of comparing $\mathrm{FR_g}$ with $\mathrm{FR_p}$, we developed a computational pipeline to assign the function types (specialist, niche, or essential) to protein families based on the paired metagenome and metaproteome (Fig. 1f). This pipeline starts with DNA sequences from

metagenomes and peptides sequences from metaproteomes. Using the "protein-peptide
bridge" approach that maps peptides to their taxonomic origins and protein families (i.e.,
orthologous protein clusters), it generates the GCN, PCN, and taxonomic profile, from which
we compute FR_g and FR_p. Details about this approach can be found in the Supplementary
Information. Finally, based on the scatterplot of FR_g vs. FR_p, each protein family is categorized
into one of the three function types. Note that the computational pipeline assigns the function
type without leveraging the known biological functions. Instead, we validate these assignments
against the knowledge about biological functions.
196
Illustration of our computational pipeline using synthetic data
To illustrate the pipeline's workflow, we utilized synthetic data generated by a consumer-
resource model (CRM). Each niche (or specialist) function is modeled as the consumption of
a unique and externally supplied resource (Fig. 2a1), whose loss would make a species unable
to consume the corresponding resource (Fig. 2a2, a3). The loss of an essential function is
modeled as reducing a species' growth rate by 5% (Fig. 2a4).
For each species, each niche (specialist, or essential) function was assigned to the
species' genome with probability p_n (p_s, or p_e), respectively (Fig. 2b, left). We set $p_n = p_e = 0.7$
to ensure that we cannot distinguish niche functions from essential functions only based on
their k_{GCN}. We set $p_s = 0.2 < p_n = p_e$ so that specialist functions were assigned to fewer
species than niche and essential functions. Species' actual expressed functions were
determined by randomly sampling a subset of its potential functions (Fig. 2b, middle). This
behavior of sub-sampling was observed when we cultured single microbial strains in different
environments (Supplementary Fig. 21). We simulated community dynamics until reaching a
steady state, for which we constructed the PCN of the surviving species (Fig. 2b, right; see
Supplementary information for technical details). More technical details of CRM are in
Supplementary information.
In our model with 10,000 species and 20 functions for each of the three function types,
each species randomly sampled a subset of potential functions (Fig. 2c, left) to express (Fig.
2c, middle). We demonstrated a simulation example with 35 species surviving in the final
steady state after the community assembly initialized with 10,000 species (Fig. 2c, right).
We applied the taxonomic profile, GCN, and PCN for the surviving 35 species to our
computational pipeline, finding that the three modeled protein function types were correctly
classified as three clusters (60 out of 60 were correct) by the Gaussian mixture model in both

the comparison of network degree (Fig. 2d) and FR (Fig. 2e). We emphasize that the observed
three functional clusters arise from community assembly. When we randomly picked 35
species (same as the number of surviving species) from the initial pool with equal abundances
without assembly, niche functions cannot be distinguished from essential functions (Fig. 2f, g).
Even when assigning the same randomly picked 35 species with the same abundances of
surviving species, we still cannot differentiate these two functional types (Supplementary Fig.
2). Our findings held when varying (1) the number of species and functions or (2) model
parameters p_n, p_s, and p_e, affirming the method's robustness in distinguishing function types
(Supplementary Figs. 3, 4, and 5).

Three protein functional clusters observed in human gut microbiomes

Next, we validated our computational pipeline on real data of human mucosal-luminal interface
samples previously collected from the ascending colon of four children[14,33]. Here we focused
on the genus level and annotated the identified proteins from metagenomic and metaproteomic
data via the clusters of orthologous genes (COGs) database[46,47]. We chose the genus level
due to widely shared peptide sequences across species (Supplementary Fig. 6). We searched
metagenomic reads and metaproteomic peptides against the integrated gene catalog (IGC)
database of the human gut microbiome[48] to generate the GCN and PCN[14], and took the
intersected COGs between the two networks. Taxonomic assignment was performed using the
"protein-peptide bridge" method as described previously[14]. Our analysis centers on subject
HM454, identifying 1,542 intersected COGs in both the GCN and PCN and obtaining a
taxonomic profile of 85 genera using MetaPhlAn2[49]. The connectance (i.e., the number of
edges divided by the maximal number of possible edges) of the GCN (or PCN) is 0.220 (or
0.049), respectively (Fig. 3a, b). The GCN displayed a higher nestedness (nestedness metric
NODF[50]=0.667) than the PCN (NODF=0.453). More details about data processing and NODF
are in Supplementary Information.
The network degree analysis revealed a general decline from GCN to PCN, with 804
out of 1,542 COGs having $\mathrm{k_{PCN}} < 0.2\mathrm{k_{GCN}}$ (Fig. 3c). This decline greatly influences FR but
does not fully explain why many COGs have $\mathrm{FR_p} \sim 0$ (744 out of 1,542 have $\mathrm{FR_p} < 0.01$ in Fig.
3d) and $\mathrm{k_{PCN}}$ nonlinearly correlates with $\mathrm{FR_p}$ (Fig. 3e). For example, for L-arabinose isomerase
(COG2160), its $\mathrm{k_{PCN}}$ (7) is fairly close to $\mathrm{k_{GCN}}$ (8), but its $\mathrm{FR_p}$ (0.04) is much lower than $\mathrm{FR_g}$
(0.23) since the genus *Blautia* (relative abundance = 22%) did not express L-arabinose
isomerase, even if it has this capacity encoded in its genome.

Using the Gaussian mixture model fitted on simulated data, we categorized all protein
families into three clusters (Figs. 3c&d). Although clusters on real data are not as distinct as
on simulated data, the relative positioning of the three clusters (shaded areas in Fig. 3c, d)
agrees well with our hypothesis (Fig. 1). The weaker clustering might result from a greater
variation in k_{GCN} (or FR_g) for real data (Fig. 3c, d) than that for simulated data (Fig. 2d, e).

Some COGs have $FR_p > FR_g$ (Fig. 3c, d), contradicting the sub-sampling argument for
the gene expression. FR_p should not exceed FR_g if the PCN was a proper subgraph of the
GCN. This contradiction may stem from limitations in metagenomic sequencing and
metaproteomic identification depths, as both metagenomics and metaproteomics require
sufficient depth to detect genes or proteins, respectively. We tested how the lower detection
capability of metaproteomics or metagenomics influences the FR by varying the protein or
gene abundance percentile threshold (PAPT or GAPT), which denotes the percentage of most
abundant proteins or genes being kept. As PAPT or GAPT decreases, FR_p or FR_g drops
respectively (Supplementary Fig. 15). When GAPT decreases, we observed more proteins
with FR_p greater than their FR_g.

Validating three functional clusters observed in human gut microbiomes

Our computational pipeline accurately assigns functional clusters for protein families, agreeing
with their known biological functions. For example, COG0539 (ribosomal protein S1) was
assigned as the essential function, which is essential for translational initiation[17–19,51,52]. Another
example is the assignment of COG1116 (ABC-type nitrate/sulfonate/bicarbonate transport
system)[34] as a niche function, whose expression has been shown to be selectively enriched
for a few microbial species[53].

The pipeline's classifications were systematically validated against well-established
biological roles of specific protein families: (1) ABC-type transporters are niche proteins due to
their connection with ecological metabolic niches[34–36]; (2) Ribosomal proteins are essential
proteins because they are indispensable for the microbial growth[54,55]; (3) PTS
(phosphotransferase system) proteins are specialist proteins because an evolutionary study
has shown that various species within the same genus even possess a different set of PTS
proteins[56]. To evaluate our pipeline's performance, we quantified its accuracy in assigning
these protein families (ABC-type transporters, ribosomal proteins, or PTS proteins) against the
assumed "ground-truth" functions (niche, essential, or specialist functions, respectively).

For HM454, our computational pipeline based on the FR_g/FR_p plot correctly categorizes 81 of 122 COGs belonging to ABC-type transporters, ribosomal proteins, or PTS proteins. In comparison, when classifying functions based on the k_{GCN}/k_{GCN} plot, 74 COGs were correctly assigned, slightly worse than that based on the FR_g/FR_p plot. Specifically, 26 of 53 COGs belonging to ABC-type transporters are classified as niche functions. The fraction of ribosomal proteins classified to the cluster of essential functions is 83.0%(=44/53). For the PTS proteins, among the identified 16 COGs, 11 are classified as specialist functions.

Alternative clustering and classification methods

Alternatively, we explored the unsupervised K-mean clustering with K=3, which captured the three representative functional clusters with their positions agreeing with our expectations (Supplementary Fig. 13). We also designed a supervised classifier based on quadratic discriminant analysis (QDA). QDA, trained on ABC-type transporters, PTS proteins, and ribosomal proteins as the niche, specialist, and essential functions, generated clusters closely resembling those from the Gaussian mixture model (Supplementary Fig. 14). For HM454, the K-mean clustering categorizes 48 of these 122 COGs that are ABC-type transporters, ribosomal proteins, or PTS proteins into clusters respectively representing niche, essential, or specialist functions (i.e., the accuracy is 39.3%). For the QDA classifier, the accuracy is 59.0%(=72/122). Thus, we select the Gaussian mixture model as the classification method because of its superior accuracy (66.4%=81/122).

Comparing FR_g with FR_p identifies ecological niches and metabolic essentiality

We focused on analyzing ABC-type transporters[34–36] and ribosomal proteins[17–19]. ABC-type transporters are energy-requiring transporter proteins that allow microbes to exploit specific niches like glucose uptake[34–36]. For HM454, we indeed found that k_{GCN} for all ABC-type transporters are much larger than their k_{PCN} (Fig. 4a). Similarly, we also found that their FR_g are much larger than their FR_p, classifying many transporter proteins as niche functions (Fig. 4b). Some transporter proteins were classified as specialist functions (blue dots in Fig. 4b) due to the specialization on the gene level, which is carried to the protein level. Some transporter proteins were classified as essential functions (green dots in Fig. 4b). One example is the ABC-type Fe^{3+}/spermidine/putrescine transporter (COG3842), as iron is essential for bacteria to function as a co-factor in iron-containing proteins[57,58].

Ribosomal proteins, critical for protein synthesis and microbial growth[54,55], showed little

variance between k_{GCN} and k_{PCN} (the mean and the standard deviation of the relative difference

$\frac{k_{GCN}-k_{PCN}}{k_{GCN}}$ is 0.09 $\pm$ 0.41; Fig. 4e), with most correctly classified as essential (44 out 53 COGs

in Fig. 4f). Notably, two ribosomal proteins (L28 and L34), which have been reported as non-

essential to microbes such as *E. coli*[17,52,59], were accurately classified as non-essential proteins

(red dots in Fig. 4e). Certain specialized ribosomal proteins in microbial genomes continue to

be specialized on the protein level and thus were classified as specialist functions.

Alternatively, we looked at the distribution of network degrees (Figs. 4c&g) and FR

(Figs. 4d&h). For ABC-type transporters, the distribution of k_{PCN} is close to 0 (median of 2),

while the median of k_{GCN} is 25. For ribosomal proteins, the distribution of k_{PCN} (median is 12) is

similar to k_{GCN} (median is 14). For ABC-type transporters, the distribution of FR_p is close to 0

(with a median ~ 0.01), while the median of FR_g is around 0.30. For ribosomal proteins, the

distribution of FR_p (median ~ 0.20) is similar to the distribution of FR_g (median ~ 0.21).

The similar patterns are also true for the other three individuals (Supplementary Figs.

8-10). Variations in the FR_g /FR_p plot across individuals (Supplementary Fig. 8-10) are likely

due to differences in gut environments, diets, and microbial composition. Note that HM503 is

an outlier due to its lower diversity (36 genera versus the average of 56.7) (Supplementary Fig.

11). Similarly, the Shannon diversity index for HM503 is only 1.41, lower than other individuals

(mean and standard deviation are 2.05 and 0.18). The lower diversity in HM503 leads to fewer

taxa owning the same function on average, resulting in lower FR_g and FR_p values.

Extending our analysis to additional protein families, we discovered that proteins linked

to glycolysis, RNA polymerase, and the TCA cycle displayed patterns similar to ribosomal

proteins (Figs. 4i&j). By contrast, proteins associated with sugar utilization, glycolyl hydrolase,

and aromatic amino acid biosynthesis exhibited patterns more akin to ABC-type transporters.

We also analyzed the classification results for the unknown COG functions (with the category

'S: Function Unknown'). Out of the 104 unknown COGs in our analysis, the majority are

classified as specialist (43/104) or niche functions (40/104), with a smaller portion identified as

essential functions (21/104).

We also confirmed our results using the KEGG Orthology (KO) annotation[60–63], which

has a lower annotation rate (78%) than COG (92%). For HM454, our computational pipeline

categorizes 75 of these 126 KOs that are ABC-type transporters, ribosomal proteins, or PTS

proteins into clusters respectively representing niche, essential, or specialist functions, similar

to the classification accuracy based on the COG. The contrasting difference between ABC-

type transporters and ribosomal proteins is well preserved (see Supplementary Fig. 7).

Additionally, the distribution of FR_p shows a dramatic difference across KO groups

(Supplementary Fig. 12). Some ecologically strongly selected KO groups such as ABC

transporters have small FR_p (Supplementary Fig. 12). As a comparison, proteins from

aminoacyl-tRNA biosynthesis[15,16], glycolysis[20,21], and ribosomes[17–19] have large FR_p and huge

variations within each group (Supplementary Fig. 12).

Validating our method on the mouse gut microbiome

In testing our method's feasibility in other microbial communities, we leveraged a

metaproteomic dataset from mice gavaged with a synthetic microbiome comprising twenty

sequenced bacteria[41]. Since this study lacks paired metagenomes, we used its 16S rRNA gene

sequencing data with individual genomes to infer its metagenome. Here we focused on the

strain level because peptides of different strains in this simple synthetic gut microbiome can

be distinguished. We relied on the comparison between FR_g and FR_p to generate the

distribution of functional clusters across many protein families for this dataset (Fig. 5n). The

results mirrored those of human gut microbiomes, especially the contrasting patterns between

ABC-type transporters and ribosomal proteins (Fig. 5).

368
Response of community and protein abundance to the introduction of sugars

After identifying niche functions through our computational pipeline, we explored using

nutrients associated with niche functions to manipulate the community structure. In ecology, a

niche is often defined as an abiotic and biotic factor that supports the survival of species[9,64–66].

Therefore, niche functions are associated with corresponding limiting resources involved in

those functions. For example, COG1879 (ABC-type sugar transporter) is categorized as a

niche function due to microbial competition for sugars. Here, we leveraged the *in vitro*

community and studied how expression levels of ATP-type transporters respond to supplied

sugars so that a microbial taxon can achieve a better living strategy.

Using the RapidAIM V2.0 approach[67], which replicates the functional profiles of

individual gut microbiomes *in vitro*[37], we cultured three individual human gut microbiota

samples and used a semi-automated metaproteomics workflow to observe how taxon-specific

proteins respond to the presence of glucose, fructose, and kestose (Fig. 6a). Samples were

cultured in technical triplicates, and protein abundances were quantified at 0, 1, 5, 12, and 24

hours using 11-plex tandem mass tag (TMT11plex)[39] for a total of 189 samples. We analyzed

the Bray–Curtis dissimilarity of metaproteomes over time, finding that more complex sugars

induce more pronounced alterations in protein profiles (Supplementary Fig. 16).

To reflect the effect of introduced sugars on protein abundances, we used log2 of fold

change in normalized protein abundances/intensities (see Supplemental Methods for details)

between the treatment and control group (Fig. 6). We hypothesized that excessive sugars

remove the growth limitation on carbon resources, prompting microbes to upregulate other

transporters for uptaking more other scarce resources (e.g., nitrogen or amino acids) for better

growth (Fig. 6a). We analyzed log2 fold changes of ABC-type transporters 5 hours later after

sugar introduction (Figs. 6b-d), with most COGs close to zero. Among seven significantly

influenced COGs, COG1126 (ABC-type polar amino acid transport system) is the only one that

is revealed to be a niche function. Focusing on COG1126, we found that it is specialized by

the genus *Holdemanella*. *Holdemanella* benefits from upregulating COG1126, as the

proportion of *Holdemanella* proteins significantly increases from 13.5%(± 0.06%) for the control

to 15.8 %(± 0.08%) with the added glucose (p-value = 0.04, Mann-Whitney U test applied).

We observed that adding fructose, glucose and fructose, or kestose alters ABC-type

transporters' expression similarly to glucose alone (Fig. 6). The correlation in log2 fold changes

of ABC-type transporters between different added sugars is significant (Supplementary Fig.

18). Notably, complex sugars trigger more significant fold changes in microbial protein

expression. This pattern persists for metaproteomic measurements 12 and 24 hours later,

while the fold changes 1 hour later are less significant (Supplemental Fig. 19; p-value < 0.01

for four sugar-adding scenarios, Mann-Whitney U test applied). Additionally, the

overwhelmingly positive log2 fold changes of ribosomal proteins (Supplementary Fig. 20; p-

value $< 10^{-4}$ for four sugar-adding scenarios, one-sample Wilcoxon test applied) probably

implies faster microbial growth when simple sugars are supplied[68,69].

Discussion

We developed a computational pipeline to classify protein families as specialist, niche, and

essential functions by comparing FR_g with FR_p . This approach supplements traditional

methods that test metabolic essentiality by gene knockout[17–19] and identify limiting resources

by measuring biomass changes upon resource supplies[70–73]. We first illustrated this method on

synthetic data and then validated it using real datasets of human and mouse gut microbiomes.

We acknowledge a limitation in our validation process—the reliance on limited available

literature. Hence, our classification should be seen as a preliminary framework that is open to
refinement as the investigation of protein families' functions improves.

Our findings bridge the gap between the ecological niche theory, which posits that each
resource (or niche) can only be occupied by one species for steady-state conditions[66,74,75], and
the FR revealed by shared functions among microbial genomes[11,12]. We solved this dilemma
by showing niche proteins usually have very small FR_p and large FR_g. Additionally, our
ecological framework combines genomic capacity and protein functions together by introducing
species with sub-sampled functions. The model framework accounts for selective expression
due to different environmental conditions[76–78] or evolved strains with distinct metabolic
niches[79,26,27], reconciling phenotype-focused ecological models with genetic data.

The observed case of FR_p > FR_g for some COGs could stem from using MetaProIQ, a
general microbiome catalog, for metaproteome analysis. Although MetaProIQ facilitates the
identification of proteins from various gut microbes, the general search against it may cause
anomalies. In contrast, directly searching metaproteomic data against gene calls from the
paired metagenome may deliver more accurate identifications. However, this strategy suffers
when the metagenomic sequencing is incomplete, leading to undetected proteins due to
missing genes. In our human gut microbiome datasets, limited sequencing depth might cause
such issues. For the synthetic mouse gut communities with complete genomes for each
microbial strain, we matched metaproteome to microbial genomes and did not encounter any
case of FR_p > FR_g, highlighting the effectiveness of this approach in contexts with complete
genomic data.

In this work, we only validated our pipeline on gut-related biomes due to the limited
accessibility of paired metagenome-metaproteome in other environments. In nutrient-poor
environments, where ribosomal proteins are less expressed[80,81], essential proteins like
ribosomes may be harder to detect, causing potential detection biases. This aspect warrants
further investigation to ensure the robustness and applicability of our method in diverse
ecological settings. Other technical limitations can also impact clustering accuracy. Smaller
ribosomal proteins L28 and L34 could be detected less frequently in metaproteomics, and post-
translational modifications may also result in missed cleavage and identification of peptides.
Advances such as metaproteomics-assembled proteomes (MAPs)[82] may improve taxon-
specific functional annotations and the accuracy of our clustering outcomes.

448
449

Data and code availability. All code for simulations used in this manuscript can be found at https://github.com/wt1005203/ecological_niches.

Acknowledgements. We thank Dr. Janice Mayne for the help with providing biobanking samples, Dr. Zhibin Ning for the help with running mass spectrometry and Dr. Joeselle Serrana for the help with processing the metagenomic data. D.F. acknowledges grants from Natural Sciences and Engineering Research Council of Canada (NSERC), and the Government of Canada through Genome Canada and the Ontario Genomics Institute (OGI-114 & OGI-149). D.F. acknowledges a Distinguished Research Chair from the University of Ottawa. Y.-Y.L. is supported by grants R01AI141529, R01HD093761, RF1AG067744, UH3OD023268, U19AI095219 and U01HL089856 from the National Institutes of Health, USA; a pilot grant from the Biology of Trauma Initiative of Broad Institute, USA; and the Office of the Assistant Secretary of Defense for Health Affairs, through the Traumatic Brain Injury and Psychological Health Research Program (Focused Program Award) under award no. (W81XWH-22-S-TBIPH2), endorsed by the Department of Defense, USA.

Author contributions. Y.-Y.L. and D.F. supervised the study. T.W. and Y.-Y.L. conceived the project. All authors designed the research. L.L. prepared and curated the empirical data as well as performed all wet-lab experiments. T.W. analyzed all data and developed the ecological model. T.W. wrote the initial manuscript. All authors edited and approved the manuscript.

Competing Interests. The authors declare no competing interests. D.F. co-founded MedBiome Inc., a clinical microbiomics company.

IRB determination/approval statement. The sample collection of human gut microbiomes was approved by the Research Ethics Board of the Children's Hospital of Eastern Ontario (CHEO), Ottawa, ON, Canada. The written informed consent forms were obtained from their parents.

References

References

1.   Tyson, G. W., Chapman, J., Hugenholtz, P., Allen, E. E., Ram, R. J., Richardson, P.
M. *et al.* Community structure and metabolism through reconstruction of microbial genomes
from the environment. *Nature* **428**, 37–43 (2004).
2.   Flint, H. J., Scott, K. P., Louis, P. & Duncan, S. H. The role of the gut microbiota in
nutrition and health. *Nature Reviews Gastroenterology & Hepatology* **9**, 577–589 (2012).
3.   Lloyd-Price, J., Arze, C., Ananthakrishnan, A. N., Schirmer, M., Avila-Pacheco, J.,
Poon, T. W. *et al.* Multi-omics of the gut microbial ecosystem in inflammatory bowel
diseases. *Nature* **569**, 655–662 (2019).
4.   Paerl, H. W. & Pinckney, J. L. A Mini-review of Microbial Consortia: Their Roles in
Aquatic Production and Biogeochemical Cycling. *Microb Ecol* **31**, 225–247 (1996).
5.   Falkowski, P. G., Fenchel, T. & Delong, E. F. The Microbial Engines That Drive
Earth's Biogeochemical Cycles. *Science* **320**, 1034–1039 (2008).
6.   Louca, S., Parfrey, L. W. & Doebeli, M. Decoupling function and taxonomy in the
global ocean microbiome. *Science* **353**, 1272–1277 (2016).
7.   Levy, R. & Borenstein, E. Metabolic modeling of species interaction in the human
microbiome elucidates community-level assembly rules. *PNAS* **110**, 12804–12809 (2013).
8.   Pacheco, A. R., Moel, M. & Segrè, D. Costless metabolic secretions as drivers of
interspecies interactions in microbial ecosystems. *Nature Communications* **10**, 103 (2019).
9.   Fahimipour, A. K. & Gross, T. Mapping the bacterial metabolic niche space. *Nature*
Communications **11**, 4887 (2020).
10.   Louca, S., Jacques, S. M., Pires, A. P., Leal, J. S., Srivastava, D. S., Parfrey, L. W. *et*
al. High taxonomic variability despite stable functional structure across microbial
communities. *Nat Ecol Evol* **1**, 15 (2016).
11.   Louca, S., Polz, M. F., Mazel, F., Albright, M. B., Huber, J. A., O'Connor, M. I. *et al.*
Function and functional redundancy in microbial systems. *Nature Ecology & Evolution* **2**, 936
(2018).
12.   Tian, L., Wang, X. W., Wu, A. K., Fan, Y., Friedman, J., Dahlin, A. *et al.* Deciphering
functional redundancy in the human microbiome. *Nature Communications* **11**, 6217 (2020).
13.   Franzosa, E. A., Morgan, X. C., Segata, N., Waldron, L., Reyes, J., Earl, A. M. *et al.*
Relating the metatranscriptome and metagenome of the human gut. *PNAS* **111**, E2329-2338
(2014).
14.   Li, L., Wang, T., Ning, Z., Zhang, X., Butcher, J., Serrana, J. M. *et al.* Revealing
proteome-level functional redundancy in the human gut microbiome using ultra-deep
metaproteomics. *Nat Commun* **14**, 3428 (2023).
15.   Ibba, M. & Soll, D. Aminoacyl-tRNA synthesis. *Annu Rev Biochem* **69**, 617–650
(2000).
16.   Parker, D. J., Lalanne, J. B., Kimura, S., Johnson, G. E., Waldor, M. K., & Li, G. W.
Growth-Optimized Aminoacyl-tRNA Synthetase Levels Prevent Maximal tRNA Charging.
Cell Syst **11**, 121-130.e6 (2020).
17.   Shoji, S., Dambacher, C. M., Shajani, Z., Williamson, J. R. & Schultz, P. G.
Systematic chromosomal deletion of bacterial ribosomal protein genes. *J Mol Biol* **413**, 751–
761 (2011).
18.   Dabbs, E. R. Mutants lacking individual ribosomal proteins as a tool to investigate
ribosomal properties. *Biochimie* **73**, 639–645 (1991).

19.   Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M. *et al.*
Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio
collection. *Mol Syst Biol* **2**, 2006.0008 (2006).
20.   Romano, A. H. & Conway, T. Evolution of carbohydrate metabolic pathways.
Research in Microbiology **147**, 448–455 (1996).
21.   Yan, Y. *Engineering Microbial Metabolism For Chemical Synthesis: Reviews And*
Perspectives. (World Scientific, 2017).
22.   Jacob, F. & Monod, J. Genetic regulatory mechanisms in the synthesis of proteins. *J*
Mol Biol **3**, 318–356 (1961).
23.   Okano, H., Hermsen, R., Kochanowski, K. & Hwa, T. Regulation underlying
hierarchical and simultaneous utilization of carbon substrates by flux sensors in Escherichia
coli. *Nat Microbiol* **5**, 206–215 (2020).
24.   Kumari, S., Beatty, C. M., Browning, D. F., Busby, S. J., Simel, E. J., Hovel-Miner,
G., & Wolfe, A. J. Regulation of Acetyl Coenzyme A Synthetase inEscherichia coli. *Journal*
of Bacteriology **182**, 4173–4179 (2000).
25.   Starai, V. J. & Escalante-Semerena, J. C. Acetyl-coenzyme A synthetase (AMP
forming). *Cell Mol Life Sci* **61**, 2020–2030 (2004).
26.   Rosenzweig, R. F., Sharp, R. R., Treves, D. S. & Adams, J. Microbial evolution in a
simple unstructured environment: genetic differentiation in Escherichia coli. *Genetics* **137**,
903–917 (1994).
27.   Treves, D. S., Manning, S. & Adams, J. Repeated evolution of an acetate-crossfeeding
polymorphism in long-term populations of Escherichia coli. *Mol Biol Evol* **15**, 789–797
(1998).
28.   Lin, H., Castro, N. M., Bennett, G. N. & San, K.-Y. Acetyl-CoA synthetase
overexpression in Escherichia coli demonstrates more efficient acetate assimilation and lower
acetate accumulation: a potential tool in metabolic engineering. *Appl Microbiol Biotechnol*
71, 870–874 (2006).
29.   Kinnersley, M. A., Holben, W. E. & Rosenzweig, F. E Unibus Plurum: genomic
analysis of an experimentally evolved polymorphism in Escherichia coli. *PLoS Genet* **5**,
e1000713 (2009).
30.   Penzlin, A., Lindner, M. S., Doellinger, J., Dabrowski, P. W., Nitsche, A., & Renard,
B. Y. Pipasic: similarity and expression correction for strain-level identification and
quantification in metaproteomics. *Bioinformatics* **30**, i149-156 (2014).
31.   Ram, R. J., VerBerkmoes, N. C., Thelen, M. P., Tyson, G. W., Baker, B. J., Blake, R.
C. *et al.* Community proteomics of a natural microbial biofilm. *Science* **308**, 1915–1920
(2005).
32.   VerBerkmoes, N. C., Denef, V. J., Hettich, R. L. & Banfield, J. F. Systems biology:
Functional analysis of natural microbial consortia using community proteomics. *Nat Rev*
Microbiol **7**, 196–205 (2009).
33.   Zhang, X., Ning, Z., Mayne, J., Moore, J. I., Li, J., Butcher, J. *et al.* MetaPro-IQ: a
universal metaproteomic approach to studying human and mouse gut microbiota. *Microbiome*
4, 31 (2016).
34.   Steinsiek, S. & Bettenbrock, K. Glucose transport in Escherichia coli mutant strains
with defects in sugar transport systems. *J Bacteriol* **194**, 5897–5908 (2012).
35.   Fath, M. J. & Kolter, R. ABC transporters: bacterial exporters. *Microbiol Rev* **57**, 995–
1017 (1993).

36.   Nikaido, H. Maltose transport system of Escherichia coli: an ABC-type transporter.
FEBS Lett **346**, 55–58 (1994).
37.   Li, L., Abou-Samra, E., Ning, Z., Zhang, X., Mayne, J., Wang, J. *et al.* An in vitro
model maintaining taxon-specific functional activities of the gut microbiome. *Nat Commun*
10, 4146 (2019).
38.   Zhang, X., Li, L., Mayne, J., Ning, Z., Stintzi, A., & Figeys, D. Assessing the impact
of protein extraction methods for human gut metaproteomics. *Journal of Proteomics* **180**,
120–127 (2018).
39.   Creskey, M., Li, L., Ning, Z., Fekete, E. E., Mayne, J., Walker, K. *et al.* An economic
and robust TMT labeling approach for high throughput proteomic and metaproteomic
analysis. 2022.07.30.502163 Preprint at https://doi.org/10.1101/2022.07.30.502163 (2022).
40.   Perez-Riverol, Y., Bai, J., Bandla, C., García-Seisdedos, D., Hewapathirana, S.,
Kamatchinathan, S. *et al.* The PRIDE database resources in 2022: a hub for mass
spectrometry-based proteomics evidences. *Nucleic Acids Research* **50**, D543–D552 (2022).
41.   Patnode, M. L., Beller, Z. W., Han, N. D., Cheng, J., Peters, S. L., Terrapon, N. *et al.*
Interspecies Competition Impacts Targeted Manipulation of Human Gut Bacteria by Fiber-
Derived Glycans. *Cell* **179**, 59-73.e13 (2019).
42.   Tyanova, S., Temu, T. & Cox, J. The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. *Nat Protoc* **11**, 2301–2319 (2016).
43.   Cheng, K., Ning, Z., Zhang, X., Li, L., Liao, B., Mayne, J., & Figeys, D. MetaLab 2.0
Enables Accurate Post-Translational Modifications Profiling in Metaproteomics. *J. Am. Soc.*
Mass Spectrom. **31**, 1473–1482 (2020).
44.   Huang, T., Choi, M., Tzouros, M., Golling, S., Pandya, N. J., Banfai, B. *et al.*
MSstatsTMT: Statistical Detection of Differentially Abundant Proteins in Experiments with
Isobaric Labeling and Multiple Mixtures. *Mol Cell Proteomics* **19**, 1706–1723 (2020).
45.   Ma, W., Kim, S., Chowdhury, S., Li, Z., Yang, M., Yoo, S. *et al.* DreamAI: algorithm
for the imputation of proteomics data. 2020.07.21.214205 Preprint at
https://doi.org/10.1101/2020.07.21.214205 (2021).
46.   Tatusov, R. L., Galperin, M. Y., Natale, D. A. & Koonin, E. V. The COG database: a
tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res* **28**, 33–
36 (2000).
47.   Tatusov, R. L., Fedorova, N. D., Jackson, J. D., Jacobs, A. R., Kiryutin, B., Koonin, E.
V. *et al.* The COG database: an updated version includes eukaryotes. *BMC Bioinformatics* **4**,
41 (2003).
48.   Li, J., Jia, H., Cai, X., Zhong, H., Feng, Q., Sunagawa, S. *et al.* An integrated catalog
of reference genes in the human gut microbiome. *Nat Biotechnol* **32**, 834–841 (2014).
49.   Segata, N., Waldron, L., Ballarini, A., Narasimhan, V., Jousson, O., & Huttenhower,
C. Metagenomic microbial community profiling using unique clade-specific marker genes.
Nat Methods **9**, 811–814 (2012).
50.   Almeida-Neto, M., Guimarães, P., Guimarães Jr, P. R., Loyola, R. D. & Ulrich, W. A
consistent metric for nestedness analysis in ecological systems: reconciling concept and
measurement. *Oikos* **117**, 1227–1239 (2008).
51.   Sørensen, M. A., Fricke, J. & Pedersen, S. Ribosomal protein S1 is required for
translation of most, if not all, natural mRNAs in Escherichia coli in vivo11Edited by D.
Draper. *Journal of Molecular Biology* **280**, 561–569 (1998).

52. Galperin, M. Y., Wolf, Y. I., Garushyants, S. K., Vera Alvarez, R. & Koonin, E. V.
Nonessential Ribosomal Proteins in Bacteria and Archaea Identified Using Clusters of
Orthologous Genes. *Journal of Bacteriology* **203**, e00058-21 (2021).
53. Bakhti, S. Z. & Latifi-Navid, S. Oral microbiota and Helicobacter pylori in gastric
carcinogenesis: what do we know and where next? *BMC Microbiology* **21**, NA-NA (2021).
54. Cech, T. R. The Ribosome Is a Ribozyme. *Science* **289**, 878–879 (2000).
55. Xue, S. & Barna, M. Specialized ribosomes: a new frontier in gene regulation and
organismal biology. *Nat Rev Mol Cell Biol* **13**, 355–369 (2012).
56. Barabote, R. D. & Saier, M. H. Comparative Genomic Analyses of the Bacterial
Phosphotransferase System. *Microbiology and Molecular Biology Reviews* **69**, 608–634
(2005).
57. Yilmaz, B. & Li, H. Gut Microbiota and Iron: The Crucial Actors in Health and
Disease. *Pharmaceuticals* **11**, 98 (2018).
58. Seyoum, Y., Baye, K. & Humblot, C. Iron homeostasis in host and gut bacteria – a
complex interrelationship. *Gut Microbes* **13**, 1874855 (2021).
59. Bubunenko, M., Baker, T. & Court, D. L. Essentiality of Ribosomal and Transcription
Antitermination Proteins Analyzed by Systematic Gene Replacement in Escherichia coli.
Journal of Bacteriology **189**, 2844–2853 (2007).
60. Kanehisa, M. A database for post-genome analysis. *Trends Genet* **13**, 375–376 (1997).
61. Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic*
Acids Res **28**, 27–30 (2000).
62. Mao, X., Cai, T., Olyarchuk, J. G. & Wei, L. Automated genome annotation and
pathway identification using the KEGG Orthology (KO) as a controlled vocabulary.
Bioinformatics **21**, 3787–3793 (2005).
63. Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M. & Tanabe, M. KEGG as a
reference resource for gene and protein annotation. *Nucleic Acids Res* **44**, D457-462 (2016).
64. E, H. G. The multivariate niche. *Cold Spring Harbor Symposia on Quantitative*
Biology **22**, 415–421 (1957).
65. Leibold, M. A. The Niche Concept Revisited: Mechanistic Models and Community
Context. *Ecology* **76**, 1371–1382 (1995).
66. Holt, R. D. Bringing the Hutchinsonian niche into the 21st century: Ecological and
evolutionary perspectives. *PNAS* **106**, 19659–19665 (2009).
67. Li, L., Ning, Z., Zhang, X., Mayne, J., Cheng, K., Stintzi, A., & Figeys, D. RapidAIM:
a culture- and metaproteomics-based Rapid Assay of Individual Microbiome responses to
drugs. *Microbiome* **8**, 33 (2020).
68. You, C., Okano, H., Hui, S., Zhang, Z., Kim, M., Gunderson, C. W. *et al.*
Coordination of bacterial proteome with metabolism by cyclic AMP signalling. *Nature* **500**,
301–306 (2013).
69. Basan, M., Hui, S., Okano, H., Zhang, Z., Shen, Y., Williamson, J. R., & Hwa, T.
Overflow metabolism in Escherichia coli results from efficient proteome allocation. *Nature*
528, 99–104 (2015).
70. Tilman, D. Resource Competition between Plankton Algae: An Experimental and
Theoretical Approach. *Ecology* **58**, 338–348 (1977).
71. Wandersman, C. & Delepelaire, P. Bacterial iron sources: from siderophores to
hemophores. *Annu Rev Microbiol* **58**, 611–647 (2004).
72. Hibbing, M. E., Fuqua, C., Parsek, M. R. & Peterson, S. B. Bacterial competition:
surviving and thriving in the microbial jungle. *Nature Reviews Microbiology* **8**, 15–25 (2010).

73.   Smith, R. L. & Smith, T. M. *Elements of Ecology*. (Benjamin Cummings, San
Francisco, 2003).
74.   Hutchinson, G. E. The Paradox of the Plankton. *The American Naturalist* **95**, 137–145
(1961).
75.   Marsland, R., Cui, W. & Mehta, P. The Minimum Environmental Perturbation
Principle: A New Perspective on Niche Theory. *The American Naturalist* **196**, 291–305
(2020).
76.   Cappelletti, V., Hauser, T., Piazza, I., Pepelnjak, M., Malinovska, L., Fuhrer, T. *et al.*
Dynamic 3D proteomes reveal protein functional alterations at high resolution in situ. *Cell*
184, 545-559.e22 (2021).
77.   Schreiber, F., Littmann, S., Lavik, G., Escrig, S., Meibom, A., Kuypers, M. M., &
Ackermann, M. Phenotypic heterogeneity driven by nutrient limitation promotes growth in
fluctuating environments. *Nat Microbiol* **1**, 1–7 (2016).
78.   Gutierrez-Ríos, R. M., Freyre-Gonzalez, J. A., Resendis, O., Collado-Vides, J., Saier,
M., & Gosset, G. Identification of regulatory network topological units coordinating the
genome-wide transcriptional response to glucose in Escherichia coli. *BMC Microbiology* **7**, 53
(2007).
79.   Goyal, A., Bittleston, L. S., Leventhal, G. E., Lu, L. & Cordero, O. X. Interactions
between strains govern the eco-evolutionary dynamics of microbial communities. *Elife* **11**,
e74987 (2022).
80.   Basan, M., Zhu, M., Dai, X., Warren, M., Sévin, D., Wang, Y. P., & Hwa, T. Inflating
bacterial cells by increased protein synthesis. *Molecular Systems Biology* **11**, 836 (2015).
81.   Serbanescu, D., Ojkic, N. & Banerjee, S. Nutrient-Dependent Trade-Offs between
Ribosomes and Division Protein Synthesis Control Bacterial Cell Size and Growth. *Cell*
Reports **32**, 108183 (2020).
82.   Armengaud, J. Metaproteomics to understand how microbiota function: The crystal
ball predicts a promising future. *Environmental Microbiology* **25**, 115–125 (2023).
83.   Zhang, X., Walker, K., Mayne, J., Li, L., Ning, Z., Stintzi, A., & Figeys, D. Evaluating
live microbiota biobanking using an ex vivo microbiome assay and metaproteomics. *Gut*
Microbes **14**, 2035658 (2022).
84.   Kanehisa, M., Sato, Y. & Morishima, K. BlastKOALA and GhostKOALA: KEGG
Tools for Functional Characterization of Genome and Metagenome Sequences. *J Mol Biol*
428, 726–731 (2016).
85.   MacArthur, R. Species packing and competitive equilibrium for many species.
Theoretical Population Biology **1**, 1–11 (1970).
86.   Goyal, A. & Maslov, S. Diversity, Stability, and Reproducibility in Stochastically
Assembled Microbial Ecosystems. *Phys. Rev. Lett.* **120**, 158102 (2018).
87.   Goldford, J. E., Lu, N., Bajić, D., Estrela, S., Tikhonov, M., Sanchez-Gorostiaga, A. *et*
al. Emergent simplicity in microbial community assembly. *Science* **361**, 469–474 (2018).
88.   Marsland III, R., Cui, W., Goldford, J., Sanchez, A., Korolev, K., & Mehta, P.
Available energy fluxes drive a transition in the diversity, stability, and functional structure of
microbial communities. *PLoS Comput Biol* **15**, e1006793 (2019).
89.   Li, Z., Liu, B., Li, S. H. J., King, C. G., Gitai, Z., & Wingreen, N. S. Modeling
microbial metabolic trade-offs in a chemostat. *PLoS Comput Biol* **16**, e1008156 (2020).
90.   Dubinkina, V., Fridman, Y., Pandey, P. P. & Maslov, S. Multistability and regime
shifts in microbial communities explained by competition for essential nutrients. *Elife* **8**,
e49720 (2019).

91.   Abrams, P. A. Character displacement and niche shift analyzed using consumer-
resource models of competition. *Theor Popul Biol* **29**, 107–160 (1986).
92.   Ispolatov, I. & Doebeli, M. A note on the complexity of evolutionary dynamics in a
classic consumer-resource model. *Theor Ecol* **13**, 79–84 (2020).
93.   Arthur, R. M. Species Packing, and What Competition Minimizes. *PNAS* **64**, 1369–
1371 (1969).
94.   Hardin, G. The competitive exclusion principle. *Science* **131**, 1292–1297 (1960).
95.   Atmar, W. & Patterson, B. D. The measure of order and disorder in the distribution of
species in fragmented habitat. *Oecologia* **96**, 373–382 (1993).
713
714

Figure 1: **Protein functions involved in determining ecological niches are postulated to**
have larger discrepancies between the gene-level functional redundancy FR_g and
protein-level functional redundancy FR_p. Here we use a hypothetical example with three
representative proteins (3 broken circles with complementary shapes to their substrates) to
demonstrate this point. **a,** Schematic of the genomic capacity of two microbial taxa (pink oval
vs yellow indented oval). Two resources (red pentagon and blue triangle) are externally
supplied to the community. The green metabolite can be transformed from the red or blue
resource and further utilized in biomass synthesis. The pink taxon has the capacity of
converting either supplied resource into the green metabolite (red and blue arrows), while the
yellow taxon can only convert the red resource (red arrow). **b,** Schematic of expressed proteins
for two microbial taxa after their competition in the same community. After the competition, the
reduced resource conflict (represented by the pink taxon choosing the blue resource as the
sole one to consume) can promote their coexistence. Gene content network (GCN) and protein
content network (PCN) can be used to capture genomic capacity and expressed protein
functions for all taxa. Alternatively, this network can be represented as incidence matrices on
the bottom (grey areas imply the existence of edges connecting taxa to proteins). **c-d,** The
comparison between k_{GCN} and k_{PCN} or between FR_g and FR_p helps to classify proteins into
three protein functional types: specialist function, essential function, and niche function. In the
calculation of FR_g and FR_p, we assume equal abundances of the two species, i.e., $p_1 = p_2 =$
0.5. **e,** The comparison between FR_g and FR_p when $p_1 = 0.9$ and $p_2 = 0.1$. **f,** The pipeline of
assigning the functions (specialist, niche, or essential) to protein families based on the paired
metagenomes and metaproteomes. Each individual's gut microbiome sample was subjected
to DNA and protein extraction. Then a protein-peptide bridge approach can be used for
generating the GCN based on the metagenome and PCN based on the metaproteome. When
matched metagenomes are available, taxonomic and functional annotations of the
metagenomes can be used for PCN generation. Based on the generated GCN, PCN, and
taxonomic profile, FR_g and FR_p can be computed and used for the function assignment.

Figure 2: **Three protein functional clusters (specialist function, essential function, and**
niche function) considered in the community assembly model form three distinct
clusters when the network degree and functional redundancy are compared between
the GCN and PCN in model-generated synthetic data. a1-a4, Three types of functions
modeled have different ecological and metabolic roles. The niche function (red proteins) and
specialist function (blue proteins) are modeled as abilities to consume externally supplied
resources. The role of essential functions (green proteins) is considered as a reduction in the
overall growth rate for each missing essential function. **b,** A schematic diagram of the
community assembly. Species (ovals and indented ovals) with expressed gene functions
selected via the sub-sampling of their genomic capacity. Then all species are co-cultured
together to simulate their ecological competition. **c,** A simulation example of the community
assembly, and the construction of GCN and PCN for the survived species. **d-e,** The
comparison of network degree and functional redundancy respectively based on the GCN and
PCN of survived species in the simulation example in panel-c. A Gaussian mixture model with
3 clusters is used to identify 3 protein functional clusters. Ellipses around clusters cover areas
one standard deviation away from their means. **f-g,** The comparison of network degree and
functional redundancy respectively based on the GCN and PCN of 35 species randomly
selected from the 10,000 species in the initial pool. All points/functions are colored red (niche
functions), green (essential functions), and blue (specialist functions) according to their types
of functions in the model. k_{GCN} (or k_{PCN}) is the network degree of each function in the GCN (or
PCN). FR_g (or FR_p) is the functional redundancy of each function on the gene level (or protein
level), respectively.

Figure 3: **Real data of the human gut microbiome showing three clusters on the plot that compares FR$_g$ with FR$_p$.** Metagenome and metaproteome of subject HM454 mucosal-luminal interface samples[33] were used to construct GCN and PCN, respectively. **a,** The GCN shows if a genus owns (or doesn't own) a COG as its genomic capacity, which is colored in black (or white). The GCN matrix is ordered to have decreasing network degrees for both genera and COGs. **b,** The PCN shows if a genus expresses (or doesn't express) a COG as its protein function, which is colored in black (or white). The PCN matrix follows the same order as the GCN. **c,** Differences in network degree for most COGs are large. k$_{GCN}$ is the network degree of each COG in the GCN (i.e. the number of genera owning each COG in the GCN). k$_{PCN}$ is the network degree of each COG in the PCN (i.e. the number of genera owning each COG in the PCN). **d,** FR$_g$ is larger than FR$_p$ for most COGs. Three clusters with three distinct colors (blue, red, and green) are predicted by the Gaussian mixture model with 3 clusters fitted on synthetic data. The transparent large circles represent centroids of three clusters. **e,** The relationship between FR$_p$ and network degree of PCN for COGs is not monotonic.

Figure 4: **Comparison of network degree and functional redundancy between the gene and protein level for many protein families including ABC-type transporters and ribosomal proteins from the human gut microbiome. a,** Network degrees in GCN are larger than network degrees in PCN for most ABC-type transporter COGs. k_{GCN} (or k_{PCN}) is the network degree of each COG in the GCN (or PCN). **b,** FR_g is larger than FR_p for most ABC-type transporter COGs. **c-d,** The distribution of network degrees and functional redundancies

(violin plots and boxplots) for ABC-type transporter COGs show a significantly huge reduction
from k_{GCN} to k_{PCN} or from FR_g to FR_p. **e,** Network degrees in GCN are comparable with that in
PCN for most ribosomal protein COGs. **f,** FR_g is comparable with FR_p for most ribosomal
protein COGs. Points in scatter plots are colored by the same colors used in Fig. 3d. **g-h,** The
distribution of network degrees and functional redundancies (violin plots and boxplots) for
ribosomal protein COGs show no significant reduction from k_{GCN} to k_{PCN} or from FR_g to FR_p. **i,**
The fraction of assigned specialist, niche, or essential functions based on comparing network
degrees k_{GCN} and k_{PCN} for many protein families. **j,** The fraction of assigned specialist, niche,
or essential functions based on comparing functional redundancies FR_g and FR_p for many
protein families. In all boxplots, the middle white dot is the median, the lower and upper hinges
correspond to the first and third quartiles, and the black line ranges from the 1.5 × IQR (where
IQR is the interquartile range) below the lower hinge to 1.5 × IQR above the upper hinge. All
violin plots are smoothed by a kernel density estimator and 0 is set as the lower bound. All
statistical analyses were performed using the two-sided Mann-Whitney-Wilcoxon U Test with
Bonferroni correction between genomic capacity (GCN) and protein functions (PCN). P values
obtained from the test is divided into 5 groups: (1) $p > 0.05$ (ns), (2) $0.01 < p \leq 0.05$ (*), (3)
$10^{-3} < p \leq 0.01$ (**), (4) $10^{-4} < p \leq 10^{-3}$ (***), and (5) $p \leq 10^{-4}$ (****). Network degree
comparison of ABC transporters: $p = 7.11 \times 10^{-16}$. Network degree comparison of ribosomal
proteins: proteins: $p = 0.10$. Redundancy comparison of ABC transporters: $p = 2.19 \times 10^{-11}$.
Redundancy comparison of ribosomal proteins: $p = 1.00$.

Figure 5: **Comparison of network degree and functional redundancy between the gene**
and protein level for many protein families including ABC-type transporters and
ribosomal proteins from the synthetic mouse gut microbial community. a, Comparison
between network degrees of all COGs in the GCN (k_{GCN}) and network degrees of all COGs in
the PCN (k_{PCN}). **b,** Comparison between functional redundancies of all COGs in the GCN (FR_g)
and functional redundancies of all COGs in the PCN (FR_p). **c-d,** The distribution of network
degrees and functional redundancies (violin plots and boxplots) for all COGs. **e-h,** Comparison
between k_{GCN} and k_{PCN}, comparison between FR_g and FR_p, the distribution of network degrees,
and the distribution of functional redundancies for ABC-type transporter COGs. **i-l,** Comparison
between k_{GCN} and k_{PCN}, comparison between FR_g and FR_p, the distribution of network degrees,
and the distribution of functional redundancies for ribosomal protein transporter COGs. **m,** The
fraction of assigned specialist, niche, or essential functions based on comparing network
degrees k_{GCN} and k_{PCN} for many protein families. **n,** The fraction of assigned specialist, niche,
or essential functions based on comparing functional redundancies FR_g and FR_p for many
protein families. In all boxplots, the middle white dot is the median, the lower and upper hinges
correspond to the first and third quartiles, and the black line ranges from the 1.5 × IQR (where
IQR is the interquartile range) below the lower hinge to 1.5 × IQR above the upper hinge. All
violin plots are smoothed by a kernel density estimator and 0 is set as the lower bound. All
statistical analyses were performed using the two-sided Mann-Whitney-Wilcoxon U Test with
Bonferroni correction between genomic capacity (GCN) and protein functions (PCN). P values
obtained from the test is divided into 5 groups: (1) $p > 0.05$ (ns), (2) $0.01 < p \leq 0.05$ (*), (3)
$10^{-3} < p \leq 0.01$ (**), (4) $10^{-4} < p \leq 10^{-3}$ (***), and (5) $p \leq 10^{-4}$ (****).

Figure 6: **Microbes modify their expression for ABC-type transporters to adapt to added**
sugars. All heatmaps share the same color bar on the right. **a,** Schematic of *in-vitro* cultures
of a collected human gut microbiome. In the treatment group, one sugar is added to the
community. Metaproteomic measurements 5 hours later were used to compare the intensity of
each taxon-specific protein using the log2 fold change of each protein's fraction (i.e. normalized
intensity over each genus) from the treatment group divided by that from the control group.
Log2 fold changes of ABC-type transporters were computed 5 hours after (**b**) glucose, (**c**)
fructose, (**d**) kestose, or (**e**) glucose and fructose is added. The transported metabolites for
each COG are added to the brackets.