Higher-order epistasis shapes natural variation in germ stem cell niche activity

Sarah R Fausett[1,#a,*], Asma Sandjak[1,2], Bénédicte Billard[1,2], & Christian Braendle[1]*

[1]Université Côte d'Azur, CNRS, Inserm, IBV, Nice, France
[#a]Department of Biology and Marine Biology, University of North Carolina Wilmington, Wilmington, NC, USA

[2]These authors contributed equally to this work

*Corresponding authors
Email: fausetts@uncw.edu
Email: braendle@unice.fr

Abstract
Natural quantitative variation in developmental processes, such as cell proliferation, must be driven by allelic variation. Yet, the genotype-phenotype relationships underlying such trait variation are understudied due to their inherent complexity. Taking advantage of the simple *Caenorhabditis elegans* germline stem cell system, we used a quantitative genetic approach to characterize natural differences in germ stem cell niche activity of two distinct wild isolates, measured as differences in germline progenitor zone (PZ) size. Using quantitative trait locus (QTL) analysis, we detected multiple candidate loci, including two large-effect loci on chromosomes II and V. Resolving the chromosome V QTL, we discovered that the isolate with a smaller PZ exhibits a unique 148 base pair deletion in the promoter region of the Notch ligand, *lag-2*, a central signal promoting germ stem cell fate and proliferation. As predicted, introducing this deletion into the isolate with a large PZ resulted in significantly smaller PZ size. Unexpectedly, re-introducing the deleted ancestral sequence in the isolate with a smaller PZ caused PZ size to reduce even further. We show that these seemingly contradictory phenotypic effects are due to, partly antagonistic, epistatic interactions among the *lag-2* promoter, the chromosome II QTL, and additional loci elsewhere in the genome. While we identified an obvious causal polymorphism explaining natural variation in germ stem cell niche activity, the effects of this variant became unpredictable due to higher-order epistasis. Studying the genetic architecture of quantitative developmental systems without taking into account its natural variation may be misleading, emphasizing the need for a better integration of developmental and quantitative genetics.

Keywords
germ stem cell, QTL, natural variation, *C. elegans*, epistasis, evolution, Delta/Notch, proliferation, *lag-2*

Introduction

Fine-tuning of cellular proliferation is a fundamental aspect of organismal development and tissue homeostasis, often coordinated by stem cell niches. Even small perturbations in stem cell niche activity can deregulate tissue growth and maintenance to cause pathologies (Morrison and Spradling, 2008). Dissecting the molecular genetic mechanisms of stem cell niches has, therefore, become a major focus of biological research. While developmental genetic studies of stem cell niche function in animals have unraveled underlying key molecular regulatory mechanisms, whether and how stem cell activity is modulated by genetic variation segregating in natural populations remains largely unaddressed. If it exists, how does such allelic variation contribute to variation in stem cell niche activity? Do known genes involved in stem cell niche signaling harbor this variation? And to what extent can natural variation in stem cell niche activity be explained by effects of single, large-effect genetic variants versus polygenic contributions of small-effect variants? Most quantitative traits are considered to be complex, involving a polygenic architecture, with genetic variants not only acting additively but also in an interactive manner. Such epistasis, also termed gene-gene (G x G) interactions, corresponds to non-additive interactions between allelic variants at different genomic loci (Phillips, 2008). Strong polygenicity and epistasis have been observed for most quantitative phenotypes across distant taxa (Boyle et al., 2017; Campbell et al., 2018; Jakobson and Jarosz, 2020; Mackay, 2014; Sackton and Hartl, 2016; Weinreich et al., 2013). Although experimentally difficult to characterize, molecular and quantitative genetic analyses also suggest the presence of widespread epistatic interactions when measuring developmental phenotypes (e.g. Barkoulas et al., 2013; Chandler et al., 2017; Duveau and Félix, 2012; Gibson and Hogness, 1996; Huang et al., 2012; Koneru et al., 2021; Paaby et al., 2015; Steiner et al., 2007; Vanhaeren et al., 2014; Vonesch et al., 2016). So far, however, there is no information on how specific interactions between natural alleles cause quantitative variation in animal stem cell systems.

Germline stem cell (GSC) systems are fundamental to metazoan development and reproduction by maintaining immortal germ cell populations in an undifferentiated state and by integrating many genetic and environmental cues that adjust the production of germ cell progenitors (Hubbard and Schedl, 2019; Laws and Drummond-Barbosa, 2017). Genetic research and comparative evo-devo studies have uncovered a diversity of GSC systems across distant taxa (Agata et al., 2006; Extavour and Akam, 2003; Lehmann, 2012), but whether the activity of these systems also shows more subtle, quantitative variation in natural populations of the same species is currently not known. Genomic and developmental genetic analysis of closely related species (e.g. within the genus *Drosophila* or the nematode genus *Caenorhabditis*) indicate that principal features of the GSC niche are largely conserved (Karp and Greenwald, 2003; Kocsisova et al., 2022; Poullet et al., 2015; Rudel and Kimble, 2001; Toomey and Frydman, 2014). Nevertheless, population-genetic studies in *Drosophila* show that central GSC genes can harbor surprisingly high levels of allelic variation, even within species, suggesting that these genes evolve rapidly, often due to positive selection (Bubnell et al., 2022; Choi and Aquadro, 2015; Flores et al., 2015). Therefore, despite their involvement in a fundamental developmental process, regulatory genes of GSC niches do not seem to be evolutionarily constrained. What remains unclear is how observed natural allelic variation translates into phenotypic variation in GSC niche activity.

To study the genetic basis of natural quantitative variation in GSC niche activity, we use the GSC system in *C. elegans*, which has served as a simple *in vivo* system to study stem cell niche function, involving a set of well-defined molecular signaling pathways (Crittenden and Kimble, 2008; Hubbard and Schedl, 2019; Kimble and Crittenden, 2007). The *C. elegans* hermaphrodite germline consists of two symmetrical arms with distal germ cell progenitor zones (PZ), also termed mitotic or proliferative zones, that include the stem cells, as well as progenitor cells in mitosis and meiotic S phase (Figure 1A). Germ cells differentiate into gamete progenitors through meiotic stages as they progress toward the proximal end of the arm (Figure 1A)

(Crittenden and Kimble, 2008; Hubbard and Schedl, 2019). The key regulatory signals are expressed by the Distal Tip Cell (DTC), a somatic gonadal cell that caps and makes contact with cells in the PZ (Figure 1A). These signals are the Delta/Serrate/LAG-2 (DSL-family) ligands, LAG-2 and APX-1, which activate Notch (GLP-1) receptors in distal germ cells, promoting the stem cell fate and inhibiting entry into meiosis (Austin and Kimble, 1989; Berry et al., 1997; Gao and Kimble, 1995; Henderson et al., 1994; Nadarajan et al., 2009; Tax et al., 1994; Yochem and Greenwald, 1989). The DTC with its contacts thus constitutes a stem cell niche (Hubbard and Schedl, 2019). GLP-1/Notch signaling is necessary and sufficient for the maintenance of the germ stem cell pool (Hubbard and Schedl, 2019; Kimble and Crittenden, 2007). Germ cells progressively enter meiosis as they move proximally and lose contact with the DTC. This is controlled by a network of RNA regulatory proteins, PUF (Pumilio and FBF) RNA-binding proteins, which promote self-renewal downstream GLP-1/Notch, and RNA regulatory pathways (GLD-1, GLD-2, SCF^{PROM-1}), which promote entry into meiosis (Chen et al., 2020; Crittenden and Kimble, 2008; Hansen and Schedl, 2013; Haupt et al., 2020; Hubbard and Schedl, 2019; Kimble and Crittenden, 2007). Additional signals from gonadal sheath cells may further modulate *C. elegans* germ cell proliferation and differentiation (Gopal et al., 2021; Gordon, 2020; Killian and Hubbard, 2005; Li et al., 2022; McCarter et al., 1997; Starich et al., 2014; Tolkin and Hubbard, 2021). Germline PZ cell number is thus determined by the interplay of distal proliferative activity and the spatio-temporal transition into the proximal meiotic state. Moreover, proliferative activity of the *C. elegans* germline is highly sensitive to variation in physiology and the external environment (e.g. differing in response nutrient quality and quantity, temperature, or social environment) (Angelo and Van Gilst, 2009; Aprison and Ruvinsky, 2016; Hubbard and Schedl, 2019; Poullet et al., 2015; Seidel and Kimble, 2015; Sowa et al., 2015). Environmental variation modulates GSC proliferation via metabolic and sensory signaling pathways (TGF-β, TOR, and insulin), which act both dependently and independently of niche-mediated Delta/Notch signaling (Dalfó et al., 2012; Hubbard et al., 2013; Korta et al., 2012; Michaelson et al., 2010; Pekar et al., 2017; Thondamal et al., 2014). The *C. elegans* germ stem cell system is thus highly plastic, capable of fine-tuning its activity in response to subtle environmental changes. Whether natural genetic variation can similarly modulate activity of this stem cell system is not known. However, several reports suggest that the size of germline progenitor pool varies not only among different *Caenorhabditis* species but also among distinct *C. elegans* wild isolates (Poullet et al., 2016, 2015)*.*

In this study, we aimed to quantify and genetically characterize natural variation in *C. elegans* germ stem cell niche activity. We show that germline progenitor zone (PZ) size – here, a proxy for germ stem cell niche activity – differs between genetically and geographically distinct wild isolates under identical standard lab conditions. We used a QTL mapping approach to identify genomic regions associated with natural variation in PZ size, concentrating on two wild isolates with pronounced differences in PZ size. Genetic mapping revealed four candidate QTL including two large-effect loci on chromosomes II and V that act additively to explain ~32% of the variation in PZ size among the lines of the mapping set. We discovered that the QTL region on chromosome V harbors an INDEL variant in the promoter region of the DSL ligand *lag-2*, which causally contributes to variation in PZ size. However, we also discovered that the phenotypic effects of this variant are strongly modulated through epistatic interactions with the QTL on chromosome II and additional, unknown genomic loci. Taken together, our results indicate that complex epistatic interactions are important contributors to natural variation in germ stem cell niche activity and provide first insights into the quantitative genetic architecture of an animal stem cell system.

RESULTS
Germline progenitor zone size varies among natural *Caenorhabditis* populations.
Under standard laboratory conditions, the germline progenitor zone (PZ) of the reference wildtype *C. elegans* strain (N2) typically contains ~250 cells during early adulthood (Hubbard and Schedl,

2019). Imaging through the entire PZ and counting the total number of germ cells is time-consuming and inefficient for screening large numbers of strains. To quantify natural variation in PZ size among many wild isolates of *Caenorhabditis*, we first developed a method to approximate PZ size by quantifying the trans-sectional area of the PZ in a single fluorescent image of a whole-mount DAPI-stained gonad (Figure 1B). PZ measurements derived by this method correlated well with individual counts of PZ nuclei from image stacks through the entire PZ (adj. R^2 = 0.757, p<2.2e-16) (Figure 1C, Table S1). We examined several wild isolates of *C. elegans* and *C. briggsae* from around the globe and discovered significant inter- and intraspecific variation in PZ size when measured in standard lab conditions (Figure 1D, Table S2).

Figure 1. Germline progenitor zone size varies among natural *Caenorhabditis* populations.
A) The germline progenitor zone (PZ), also termed proliferative zone or mitotic region, located at the distal end of each gonad arm, contains mitotically dividing stem and progenitor cells. The germ stem cell (GSC) pool is maintained through Delta/Notch signaling by the somatic Distal Tip Cell (DTC), which enwraps distal PZ cells. The Delta/Serrate/LAG-2 (DSL-family) ligands, LAG-2 and APX-1, activate Notch (GLP-1) in germ cells to promote proliferation while also preventing meiotic entry. Germ cells differentiate into gamete progenitors through meiotic stages as they progress toward the proximal end of the arm. B) PZ size was used as a proxy for PZ cell number. Whole-mount worms were stained with DAPI and imaged under epifluorescence. The PZ was identified as previously described (Crittenden and Kimble, 2008) and isolated (B'). The size (trans-sectional area) of the PZ was defined and calculated by setting a fluorescence threshold in ImageJ (B"). C) PZ size (μm^2/1000) measured in this way correlates well with cell number as counted by hand from z-stacks (adj R^2 = 0.757, p<2.2e-16). D) PZ size (μm^2/1000) was measured in select wild isolates of *C. elegans* and *C. briggsae*. Lowercase letters indicate significant (p<0.05) Tukey-adjusted pairwise contrasts: isolates that share the same letters do not exhibit a significant difference in PZ size. E) World map showing the geographical origin of examined wild isolates. For statistical models and results see Tables S1-S2.

Germ cell proliferative activity differs between wild isolates JU1200 and JU751

Among the wild isolates we found to vary in PZ size, JU1200 and JU751 showed significant and highly reproducible differences. JU1200 shows high genetic similarity with the laboratory strain N2 (Cook et al., 2017). As young adult hermaphrodites, JU1200 (Scotland) exhibited consistently more progenitor cells than JU751 (France) (Figure 1D). We assayed PZ cell number in these two isolates from late larval to reproductive adult stages. While PZ cell number did not differ at the mid-L4 stage, JU751 showed significantly reduced PZ cell number relative to JU1200 across two stages of early adulthood (Figure 2A, Table S3).

Given that PZ size may not only be affected by proliferative activity but also by the transition rate towards the meiotic fate, we assayed mitotic activity using a 15-minute EdU (5-ethynyl-2-deoxyuridine) pulse to directly label and quantify proliferation rate in the PZ prior to the adult stage. At the mid-L4 stage, when PZ cell number in JU751 and JU1200 were not significantly different (Figure 2A), JU751 exhibited significantly reduced counts of EdU+ cells, and thus lower germ cell proliferative activity than JU1200 (Figure 2B, Table S4). Differences in germ cell proliferation thus contribute to the observed difference in PZ size between JU751 and JU1200.

Figure 2. Germ cell proliferative activity differs between wild isolates JU1200 and JU751.
A) Number of PZ nuclei at mid L4 (mL4), young adult (YA) with 1-10 eggs *in utero*, 24 hours post-mid L4 (mL4+24), and 24 hours post-young adult (YA+24). Nuclei were counted in z-stacks of extruded DAPI-stained gonads. Bold lowercase letters indicate significant (p<0.05) pairwise contrasts such that groups sharing the same letters are not significantly different. B) Number of EdU+ nuclei in the PZ after a 15-minute pulse of EdU at mid L4 (*** p < 0.001). For statistical models and results, see Tables S3-S4. C) Representative images of the PZ in whole-mount DAPI-stained worms at mid-L4+24h. D) Representative images of EdU staining at the mid-L4 stage. Dotted lines mark the proximal boundary of the PZ. Scale bars are 20μm.

QTL mapping identifies multiple QTL explaining variation in PZ size

To characterize the genetic architecture underlying the observed differences in PZ size between JU751 and JU1200, we performed Quantitative Trait Locus (QTL) mapping based on phenotyping of F2 recombinant inbred lines (RILs) derived from these two isolates (Figures 3A and 3B). The RILs were constructed by selfing F1 hermaphrodites from reciprocal parental crosses and inbreeding the F2 lines for twelve generations to produce a panel of lines homozygous at each locus for one of the two parental genotypes (Figure 3B). We measured PZ size of young adult hermaphrodites (1-10 eggs in the uterus) in 70 RILs and mapped the phenotypic differences based on a genetic map derived from the recombination frequencies of 140 SNP markers throughout the genome (Figure 3A, Figure S1, Table S5). PZ size varied continuously across RILs and showed transgressive segregation (i.e. extremes exceeding the parental phenotypes) (Figure 3A). We therefore employed a multi-QTL modeling approach using the R package, R/qtl (Broman and Saunak, 2009). As a first step, we performed a single QTL scan, an algorithm that calculates the likelihood that the genotype at any *single* locus can explain the phenotype data. This scan revealed a large-effect locus in the center of chromosome II with a 1.5-LOD interval of ~7.25 Mb (Figure 3C, green line). Next, we used a "two-QTL" algorithm to determine the likelihood that any *pair* of loci is associated with the phenotype. This revealed an additional locus on the left arm of chromosome V (~2.6 Mb) acting additively with the chromosome II (abbreviated QII) QTL (Figure 3D-F). Single-QTL mapping with the QII QTL as a covariate also revealed the chromosome V QTL (abbreviated CV) (Figure 3C, magenta line). A final two-QTL scan that specifically allowed loci to interact with opposing effects revealed two more potential QTL on chromosomes I (~1 Mb) and X (~2.7 Mb) (Figure 3G-I). We then performed multi-QTL model selection, using these four loci as candidates. We performed the model selection manually and used a model-building and pruning tool provided in R/qtl. This algorithm begins naïvely or with a specified model, adding loci through successive genome scans and assigning each model a penalized LOD score. It then

prunes the model to achieve the simplest form with the highest penalized LOD score. A representative set of all the models tested, along with their penalized LOD scores, are shown in Figure 3J. We used both approaches and determined that our RIL data best support a model in which C2 and C5 act additively to determine PZ size. This model explained ~32% of the variation in the RIL data (Table S6).

Figure 3. QTL mapping identifies multiple QTL explaining variation in PZ size
A) PZ size (μm^2/1000) measured in 70 F2 recombinant inbred lines (RILs) derived from a cross between wild isolates JU1200 (red) and JU751 (blue). B) Scheme illustrating the generation and analysis of the F2 RIL panel derived from the two parental isolates with different PZ size. Example: The dotted black line indicates the location of a theoretical QTL. C) Results of single-QTL scans that interrogate each location on the genetic map for an association with the phenotype. The logarithm of odds (LOD score) is shown on the y-axis and the horizontal lines indicates the significance threshold for each scan as determined by permutation analysis (5000 permutations). The genetic map for each linkage group is shown on the X-axis. While the green line shows the results of a naïve scan, the magenta line shows the results of a scan in which SNP marker 14 on chr II (the peak in the magenta curve) is included as a covariate. In this scan, an additional QTL on chr V is detected. D) Results from a two-QTL scan that interrogates each pair of loci for a significant effect on the phenotype. The top half show the maximum LOD scores (LOD_a) comparing a strictly additive model for each pair of loci on chr II and V to the null model. The bottom half shows the LOD scores comparing the additive model with the single-QTL model for each pair of loci (LOD_{av1}). LOD_{av1} scores above threshold (white, as determined by permutation analysis) indicate an improvement in fit over the single-QTL model. E) Interaction plot for a pair of markers that fall within the significant peak in the bottom half of panel D (chr V, marker 1 and chr II, marker 14). F) Approximate locations and sizes (based on 1.5 LOD intervals) of the QTL on chr II and V. G) Results from a two-QTL scan that interrogates each pair of loci for a significant effect on the phenotype, allowing for interactions (opposing effects on phenotype). The top half shows the maximum LOD scores (LOD_i) comparing a full two-QTL model for each pair of loci on chr I and X to the additive model. The bottom half shows the LOD scores comparing the full model (allowing for interactions) with the single-QTL model for each pair of loci (LOD_{fv1}). LOD_i scores above threshold (white, as determined by permutation analysis) indicate evidence for an interaction between two loci, whereas LOD_{fv1} scores above the threshold indicate an improvement in fit of the full QTL model over the single-QTL model. H) Interaction plot for a pair of markers that fall within the significant peak in the top half of panel G (chr I, marker 14 and chr X, marker 2). I) Approximate locations and sizes (based on 1.5 LOD intervals) of the candidate QTL on chr I and X. J) Graphic representation of multi-QTL model selection. Each node represents a candidate QTL (QII=chrII@35cM, QV=chr V@0cM, QI=chrI@23cM, QXa=chrX@16cM, QXb=chrX@2.5cM) and each spoke represents an interaction. Values below each model are the penalized LOD scores. Scores above zero indicate that the model performs better than the global null of zero QTL. For additional information on the mapping and statistical methods used, see Experimental Procedures and Figure S1. For multi-QTL model statistics see Table S6. Analyses were carried out on raw data (PZ Area in pixels) and y-axes in (A), (E), and (H) were scaled to μm^2 for presentation (0.1008 μm^2/pixel).

Validation of the chromosome II QTL through generation of near-isogenic lines (NILs)
To validate the effect of the chromosome II QTL (QII) on PZ size, we generated near-isogenic lines (NILs). Two RILs containing JU751 sequence in QII were backcrossed for 10 generations to JU1200 to produce two NILs containing JU1200 sequence in various segments of the QTL (Figure 4A). Two RILs containing JU1200 sequence in QII were backcrossed for 10 generations to JU751 to produce five NILs containing JU1200 sequence in various segments of the QTL (Figure 4B). NILs were generated by selecting for the presence of parent-specific PCR products in the QTL region. We assayed PZ size several times in each of the NILs. The NILs established in the JU1200 background showed subtle effects on PZ size, which were not always consistent across experimental assays (blocks) (Data S1). We, therefore, analyzed the aggregated results from across nine experiments using generalized linear models to account for experimental block effect and found that we could detect a small but significant reduction in PZ size (Figure 4C, Table S7). The NILs in the JU751 background showed larger and more consistent differences in PZ size, confirming the effect of QII on PZ size (Figure 4D, Table S8). Moreover, recombination events that had occurred during the backcrossing procedure allowed us to narrow down the QTL region from 7.25 Mb to ~1.37 Mb (Figure 4B). Searching for JU751-JU1200 polymorphisms in this reduced genomic region uncovered a total of ~676 variants (122 potential high-impact variants, i.e. variants predicted by a BCSQ algorithm to likely alter gene function, for example, through frameshift or nonsense mutations) in 236 distinct genes and 10 INDELs (Data S1) (Cook et al., 2017). However, we did not identify a strong candidate variant for further analysis.

Figure 4. Validation of the chromosome II QTL through generation of near-isogenic lines (NILs)
A) Generation of NILs in the JU1200 background. Diagram showing the original QII QTL interval of ~7.25Mb on chr II, with the dashed vertical line indicating the approximate location of the LOD peak. Two RILs (RIL54, RIL127) containing JU751 sequence in QII were backcrossed for 10 generations to JU1200 to produce two NILs (NIC1697, NIC1701). Blue: JU751 genotype, red: JU1200 genotype. B) Generation of NILs in the JU751 background. Diagram showing the original QII interval of ~7.25Mb on chr II, with the dashed vertical line indicating the approximate location of the LOD peak of the original QTL. Two RILs (RIL128, 71) containing JU1200 sequence in QII were backcrossed for 10 generations to JU751 to produce five NILs (NIC1671, NIC1672, NIC1673, NIC1675, NIC1676. Blue: JU751 genotype, red: JU1200 genotype. C) PZ size measurements in NILs established in the JU1200 background shown in (A) and the parental isolates. D) PZ size measurements in NILs established in the JU751 background shown in (B) and the parental isolates. The reduced QII interval was determined by asking whether each NIL showed a significant increase in PZ size as compared to JU751. All NILs except NIC1676 had larger PZs than JU751. Therefore, the region over which the JU1200 segment in these NILs overlap was considered the reduced QTL interval (~1.37Mb). Analysis was carried out on raw data (PZ area in pixels), and the y-axes were scaled to $\mu m^2/1000$ for presentation (0.0504 μm^2/pixel). For statistical models and results see Tables S7 and S8 (* $0.01 < p \leq 0.05$, ** $0.05 < p \leq 0.01$, *** $p \leq 0.001$).

A 148 base pair deletion in the upstream regulatory region of the DSL ligand *lag-2* is a candidate causal variant in the chromosome V QTL

The 2.6Mb region of QV contained only 21 presumptive candidate variants (SNPs, INDELS) contributing to PZ size differences between JU751 and JU1200 (Data S1). Inspecting these variants in detail revealed one major candidate: a deletion in JU751 upstream of the genomic region encoding the delta-like ligand, *lag-2*, a primary regulator of germ cell proliferation and differentiation in the Notch pathway (Cook et al., 2017; Hubbard and Schedl, 2019). We named this deletion *lag-2(cgb1007);* abbreviated from here forward as *lag-2p(-)* whereas we refer to genotypes without this deletion as *lag-2p(+)*. This deletion spans 148 bp and is located in the *lag-2* promoter region, 2441 bp upstream of the transcription initiation site (Figures 5A and S2). The 3.3kb upstream promotor region of *lag-2* contains six HLH-2 binding sites (E-boxes), which are necessary for robust *lag-2* transcription in the DTC (Chesney et al., 2009; Karp and Greenwald, 2003). The *lag-2(cgb1007)* deletion contains one of these conserved HLH-2 binding sites as well as most of a 30bp poly(G) repeat (Figure 5A, S2, S3). This deletion could explain reduced germ cell proliferation observed in JU751 due to reduced *lag-2* transcription via the loss of a specific binding site for the transcription factor HLH-2 (Chesney et al., 2009).

Examining whole-genome sequence data of a global panel of over 1500 *C. elegans* wild isolates (Cook et al., 2017; Lee et al., 2021), we found the *lag-2(cgb1007)* deletion to be unique to the isolate JU751. A similar, but larger, deletion was found in two isolates from Asia, GXW1 and JU4073 (Figure S3C and S3D). Since no other isolates were found to carry the *lag-*

2(cgb1007) deletion, we wondered if this deletion could have arisen after isolation, during laboratory culture. We therefore examined additional isolates (for which no whole-genome sequence data exist) that were collected from the same habitat (compost) and location (Le Perreux-sur-Marne, France) as JU751 in 2004 and 2005 (Barrière and Félix, 2007). All other isolates (N=5) that were collected together with JU751 from this locality on the same day shared a highly similar haplotype and also carried the *lag-2(cgb1007)* deletion. In contrast, all examined isolates with different haplotypes, collected at other dates throughout 2004 and 2005 (N=13) did not carry the deletion (Vigne et al., 2021a)(Figure S3). We conclude that the *lag-2(cgb1007)* deletion was acquired before isolation in the laboratory and this allele is likely rare in natural populations.

Figure 5. A 148 base pair deletion in the upstream regulatory region of the DSL ligand *lag-2* has background and stage-specific effects on *lag-2* expression and PZ size.

A) Diagram of the QV QTL (black bar) and a 5 kb region within it containing the *lag-2* gene and its 3 kb upstream sequence. The well-characterized *lag-2* promoter contains several known transcription factor binding sites including six HLH-2 E-boxes (dark blue)(Chesney et al., 2009; Karp and Greenwald, 2003). JU751 has a 148 bp deletion (red) overlapping the third HLH-2 binding site and most of a 30bp polyG repeat (orange). The locations of all promoter elements and UCSC phastCons statistics (green) were taken from the wormbase.org (version WS283) JBrowse tool. B) The size of the PZ ($\mu m^2/1000$) in the parents and allelic replacement lines (ARLs) at the young adult stage. C) Representative images used for smFISH quantification of *lag-2* transcripts in the DTC at mid L3 (*lag-2* mRNA green, DAPI [DNA] white, scale bar 10μm). C) The gonad arm is outlined by a dashed line. The arrow marks the DTC. C') Higher magnification of C. Arrow indicates the DTC. Arrowheads indicate individual puncta, which were quantified to determine *lag-2* expression levels. D-F) Quantification of *lag-2* mRNA puncta via smFISH. Crossbars and error bars represent estimated marginal means +/- standard error from a single generalized linear model. Data were split into three graphs for clarity. Groups with different lowercase letters are significantly different (p<0.05) Tukey-adjusted pairwise contrasts. Analysis was carried out on raw data (PZ area in pixels), and the y-axes were scaled to $\mu m^2/1000$ for presentation (0.0504 μm^2/pixel). For statistical models and results see Tables S9-S10.

Restoring the genomic region of the *lag-2(cgb1007)* deletion does not increase but reduces PZ size in JU751

To validate that the *lag-2(cgb1007)* deletion genomic region contributes to differences in germline PZ size between JU751 and JU1200, we created reciprocal allelic replacement lines (ARLs) using CRISPR-*Cas9* genome editing. First, we introduced the deletion into JU1200, resulting in the strain JU1200*lag-2p(-)*. Second, we restored the deleted *lag-2* promoter region in JU751, resulting in the strain JU751*lag-2p(+)*. As predicted, removing the 148 bp fragment in the JU1200 background strongly reduced PZ size (Figure 5B). In contrast, and contrary to expectations, restoring the corresponding fragment in the JU751 background did not increase but further reduced PZ size (Figure 5B and Table S9). Therefore, relatively smaller PZ size in JU751 cannot be explained by the *lag-2(cgb1007)* deletion alone. In other words, the effects of the *lag-2(cgb1007)* deletion are strongly dependent on the genetic background. This is in line with our QTL analysis, in which the QV QTL was only detected when accounting for variation at the QII QTL (Figure 3C). Furthermore, while *lag-2(cgb1007)* has clear effects on PZ size and, therefore contributes to the overall effect QV, we cannot rule out other causal variants within QV.

Differences in *lag-2* expression recapitulate genotypic differences in PZ size

Given the result above, we next wanted to directly determine if and how observed genotypic differences in PZ size can be explained by differences in *lag-2* expression. Using single molecule fluorescence *in situ* hybridization (smFISH) (Raj et al., 2010, 2008), we quantified *lag-2* transcripts in the DTCs of the parental isolates and reciprocal ARLs, JU1200*lag-2p(-)* and JU751*lag-2p(+)*. We first measured *lag-2* expression in parental isolates and found that expression was considerably higher during larval development than in the adult stage, with strongest expression during the early L3 stage (Figure S4 and Table S12). Quantifying *lag-2* expression in parental isolates and reciprocal ARLs across three larval stages, we found that at the early L3 stage (but not at mid-L2 or at mid-L4), JU1200 showed significantly higher *lag-2* expression relative to JU751 (Figure 5D and Table S10), consistent with its larger PZ size. Remarkably, relatively higher expression of *lag-2* in JU1200 at the L3 stage was completely abolished in JU1200*lag-2p(-)*, which showed very similar expression levels as JU751 across all three developmental stages (Figure 5E, Table S10). These results are consistent with past results showing that the *lag-2* promotor E-boxes are required for robust *lag-2*::GFP transgene expression during the L3 stage (Chesney et al., 2009). Inserting the 148 bp *lag-2* promoter sequence into the JU751 background did not result in higher *lag-2* expression at L3 relative to JU751. Instead, *lag-2* expression in JU751*lag-2p(+)* was lower at L3 with similar trends in expression changes among the stages (Figure 5F, Table S10). This result confirmed that the unexpected reduction in PZ size of JU751*lag-2p(+)* is indeed causally linked to reduced *lag-2* expression. Overall, these smFISH measurements show that *lag-2* expression (at the L3 stage) tightly recapitulates observed genotypic differences in PZ size at the early adult stage and confirm that the effects of the *lag-2(cgb1007)* deletion are strongly dependent on the genetic background. Analysis of the parental isolates also shows that natural differences in *C. elegans* germ stem cell niche activity can be directly linked to differences in the expression of a core signaling component, the Notch ligand *lag-2.*

Higher-order epistatic interactions shape natural variation in PZ size

Given the apparent epistatic interactions between and genetic background, we built a more exhaustive model to account for additional two- and three-way interactions among 1) central portions of the chromosome II QTL isolated in our NILs ("C2"—with genotypes *C2-JU1200* or *C2-JU751*), 2) the *lag-2* promoter variant ("C5"—with genotypes *lag-2p(+)* or *lag-2p(-))*, and 3) the genetic background ("BGND"—JU1200 or JU751). To do this, we generated a large data set containing all eight possible genotype combinations (labeled i to viii) (Figure 6A). Analyzing this data set revealed both subtle and dramatic two- and three-way interactions (Figure 6B to 6E). To explore these data in detail, we built a generalized linear model, using a "top-down" model

selection strategy to optimize the Bayesian information criterion (BIC) and the adjusted deviance accounted for in the model (D^2, (Márcia Barbosa et al., 2013)) (Table S11). The optimized model indicated that PZ size is best explained by all three main effects and all of their interactions (i.e., fully-crossed). In addition, the model includes a fixed main block effect and interactions to account for variation across different experimental blocks. The model accounts for 33.4% of the adjusted deviance and fits the data well (χ^2 =2796.5 on 2763 df; p = 0.324) (Table S11).

To interpret the model, it is helpful to compare the estimated marginal means (Figure 6B, Table S11B). Examining the genetic interactions of the two loci and genetic background reveals widespread and strong epistasis explaining variation in PZ size. As mentioned earlier (Figure 5B), the *lag-2(cgb1007)* deletion in otherwise isogenic parental backgrounds, has opposing phenotypic effects: the deletion strongly reduces PZ size in JU1200 but increases PZ size in JU751 (Figure 6B). Examining the epistatic interactions shows that the effects of the *lag-2(cgb1007)* variant are highly contingent on both C2 and genetic background, so that the *lag-2* deletion may have negative, positive or no phenotypic consequences across the set of genotypes (Figures 6B-E). Similarly, the phenotypic effects of C2 are strongly dependent on the genetic context: *C2-JU751* reduces PZ size in every genotype (Figure 6B—i vs. ii, v vs. vii, and vi vs. viii) *except JU1200*$_{C5\text{-}lag\text{-}2p(-)}$ (Figure 6B—iii vs. iv). The presence of the *lag-2* deletion dampens the effect of *C2-JU751* in the JU1200 background (Figure 6E' and 6E"), which could imply that variants within C2 act, at least in part, through this region of the *lag-2* promoter. As one might expect, the combined effect of all other unidentified variants in the genetic background has the greatest influence on PZ size overall. Yet, this effect does vary remarkably depending on the presence of *lag-2p* deletion and genotypes at C2. In general, the JU751 background strongly reduces PZ size (Figure 6C', 6C"), but there is no significant difference between *JU1200*$_{C2\text{-}JU1200}$ and *JU751*$_{C2\text{-}JU1200}$ when the *lag-2* deletion is present (Figure 6B—iv vs. v, and 6C'). We interpret this to mean that, in presence of the *lag-2* deletion (*lag-2p(-)*), *C2-JU1200* variants exert a strong positive effect to maintain PZ size despite the average negative effect of variants in the JU751 background (Figure 6D", red line). Conversely, *C2-JU751* variants cannot maintain this positive effect resulting in background sensitivity (Figure 6D", blue line). Finally, without the *lag-2p* deletion (*lag-2p(+)*), the *C2-JU751* variants do not modify the strong negative effect of the JU751 background (Figure 6D' vs 6D" red lines).

Our model shows that higher-order epistatic interactions contribute to variation in *C. elegans* PZ size. Importantly, some of the observed phenotypic effects are masked when only two-way interactions are considered (for example, Figure 6E vs. E' and E"). These results highlight the need for caution when interpreting the phenotypic effect of a variant or allele, as the effect may be highly contingent on other variants in the genetic background. We also stress that even our interpretations of the effects of C2 and the genetic background must be considered with care as they are aggregate effects, which may be influenced by underlying epistatic interactions and, possibly, additive effects of closely linked loci.

Figure 6. Higher-order epistatic interactions shape natural variation in PZ size
A) Scheme depicting the generation of the eight genotypes used in the interactions analysis. Modifications were made to either create or replace the *lag-2(cgb1007)* deletion in JU1200 and JU751 background NILs. The successfully modified lines were then backcrossed to the parental lines to isolate the CRISPR modifications in the parental backgrounds. Genotypes are labeled i-viii for clarity. B) Estimated marginal means +/- SE of PZ size from a generalized linear model describing a data set containing all eight genotypes. JU1200 and JU751 genotypes are indicated by red and blue, respectively, and are indicated for each of the loci and background below the chart. Lowercase letters indicate significant (p<0.05) Tukey-adjusted pairwise contrasts such that groups that share a letter are not significantly different. C-E) The same model data as in (A) presented to show interactions among the loci and background. Graphs with black axes show only two-way interactions. The third dimension is represented as a split into two graphs with red and blue axes indicating the genotypes for the locus given in the bottom left. Two-way interaction p-values are given in the graphs. Analysis was carried out on raw data (PZ area in pixels), but the y-axes were scaled to μm^2 for presentation (0.0504 μm^2/pixel) (see Table S11 for model details).

DISCUSSION

Our study provides insights into the natural variation and quantitative genetic architecture of a germ stem cell system. Examining only a handful of *C. elegans* wild isolates, we were able to detect significant differences in germline progenitor zone (PZ) size, indicative of variation in germ stem cell niche activity. Our main findings are: (1) Natural variation in PZ between two isolates maps to multiple, partly interacting QTL. (2) Resolving the QTL on chromosome V identifies a deletion in the promoter region of the DSL ligand *lag-2*, a known key signal promoting the germ stem cell fate via the Notch pathway. (3) Natural quantitative variation in *C. elegans* germline proliferative activity arises through modulation of transcriptional activity of core signaling elements of the germ stem cell niche. (4) Using introgression and allelic replacement lines, we show that main effects and interactions of the two QTL on chromosomes II and V (*lag-2* deletion) are partly antagonistic and highly contingent on the genetic background. (5) Hence, higher-order epistatic interactions shape natural variation in germ stem cell niche activity.

Epistatic interactions underlie additive genetic variance

Performing QTL mapping on a panel of F2 RILs, derived from two parental isolates with divergent PZ size, we detected four, partly interacting QTL. In the best model identified by our multi-QTL model selection approach, the QII and QV QTL act additively to explain 32% of the phenotypic variance in the examined RIL population. However, analysis of introgression and allelic replacement lines uncovered extensive epistatic interactions among these loci and the genetic background. This finding is not surprising given that gene-gene interactions contribute to the component of additive genetic variance (V_A) measured at the population level, that is, high V_A variance need not reflect additive gene action (Cheverud and Routman, 1995; Falconer, 1996; Forsberg et al., 2017; Hill et al., 2008; Huang and Mackay, 2016; Lachowiec et al., 2015; Mäki-Tanila and Hill, 2014; Monnahan and Kelly, 2015; Sackton and Hartl, 2016). In particular, measures of V_A due to individual loci will strongly depend on allele frequencies of all loci that are epistatic to each other. Hence, measures of V_A (and other genetic variance components, including the epistatic variance) provide little to no information about underlying epistatic interactions, and hence, genetic trait architecture, even when QTLs have been detected. Characterizing the molecular nature of QTLs and their interaction in controlled genetic backgrounds is therefore essential.

Germ stem cell niche activity is a complex trait

Our observations suggest that *C. elegans* germ stem cell niche activity is a typical complex trait involving a polygenic architecture. Additional observations support this view indirectly: (1) PZ size varies continuously, and often subtly, in natural populations (this study); (2) mutations in large number of genes modulate PZ size and these genes exhibit diverse functions that not only act in germ stem cell proliferation, cell cycle progression and differentiation, but also in diverse metabolic or sensory processes (Hubbard and Schedl, 2019), (3) PZ size shows stochastic variability (e.g. inter-individual variation within the same environment) and strong sensitivity to diverse environmental factors (this study; (Hubbard and Schedl, 2019). Together with our findings, these observations suggest that *C. elegans* PZ size is a higher-order phenotype that likely integrates effects of allelic variation at many loci. In the case of our example, focusing on two target isolates with pronounced differences in PZ size, we also identified large-effect loci or variants (C2 and C5) whose effects were partly antagonistic and, overall, strongly dependent on the genetic context. Together, and in line with previous studies, this suggests that the effects of large-effects variants, even if resolved at the molecular level (such as the *lag-2(cgb1007)* deletion), can be misleading if epistatic interactions in the native background are ignored (Bernstein et al., 2019; Campbell et al., 2018).

Natural variation in *lag-2* expression
Resolving the QV QTL region identified a unique 148 bp deletion in the *lag-2* promoter region of JU751, *lag-2(cgb1007).* This deletion removes one of six E-box sites required for binding of HLH-2, a positive regulator of *lag-2* mediated germ cell proliferation and germline expansion (Chesney et al., 2009; Karp and Greenwald, 2003). Mutating these E-box sites was shown to result in reduced *lag-2::GFP* transgene expression (and subsequently reduced germ cell proliferation), specifically during the L3 stage (Chesney et al., 2009). We therefore expected the *lag-2(cgb1007)* to cause reduced *lag-2* expression due to reduced HLH-2 binding activity. Consistent with this scenario, introducing this deletion into JU1200 (with a large PZ size) significantly reduced both PZ size and *lag-2* expression in the DTC (Figure 5B and 5E). In contrast, restoring the corresponding ancestral sequence (JU1200) of this deletion in JU751 (with a small PZ size) further reduced, rather than increasing, PZ size and *lag-2* expression (Figure 5B and 5F). The *lag-2* deletion allele on its own therefore exerts opposing effects depending on the genetic background (sign epistasis). However, examining the interaction with the chromosome II QTL shows that *C2-JU1200* can dampen this reversal (Figure 6C" vs. C"). In addition, the *lag-2* deletion modifies the effect of C2 by dampening the negative effects of *II-JU751* in both backgrounds (Figure 6E' vs. E"). This indicates that C2 interacts with the *lag-2* promoter region, specifically in the region of the *lag-2(cgb1007)* deletion. We do not have any hypotheses regarding the molecular nature of this interaction as none of the genes (HLH-2, LIN-39, UNC-130, DAF-3/DAF-5, CES-1) with confirmed binding sites or response elements in the *lag-2* promoter region are located within the C2 QTL region. Resolving the QTL will, therefore, be required to dissect this genetic interaction. Overall, the *lag-2* deletion containing an HLH-2 binding site previously shown to positively effect *lag-2* expression in N2, can have positive, negative, or neutral effects depending on the genetic context (Figure 6).

Although it remains unclear if and how the *lag-2(cgb1007)* deletion contributes to reduced PZ size in JU751, our smFISH measurements of *lag-2* transcripts in the DTC provide unambiguous support that differences in *lag-2* transcriptional activity contribute to observed differences in germ stem cell niche activity observed between JU751 and JU1200. *lag-2* smFISH measurements closely mirrored PZ size measurements not only in parental isolates but also in reciprocal ARLs, in which the *lag-2* promoter fragment was manipulated (Figure 5B). Identification and characterization of this natural variant shows that natural variation in PZ size can occur through direct modification of core signals of the *C. elegans* stem cell niche. While it is clear that this contributes to the overall effect of the QV QTL, we cannot rule out contributions from other unidentified variants in the QTL region.

Limitations of QTL mapping and interpretation of epistatic interactions
Here we focused on the analysis of two large-effect QTLs and their interactions, detected using a rather small panel of RILs (N=70). Therefore, detecting small-effect and additional epistatic interactions were unfeasible given the relatively low statistical power. Nevertheless, a specific search for interacting loci with opposing effects yielded two candidate QTL on chromosomes I and X (Figure 3H). Further analysis of this antagonistic interaction will likely be limited by its relatively small effect size and the already complex interactions between the C2, C5, and the genetic background. Another technical limitation of our study is evident in that our efforts to isolate causal variants through introgression lines (NILs) were only partially successful. While we were able to narrow down the presumptive QII QTL interval from 7.25 Mb to 1.37 Mb, its effect on PZ size became more difficult to detect as the QTL interval was being reduced, especially in the JU1200 background (Figure 4A). Moreover, there was a relatively high number of polymorphisms between JU1200 and JU751 in this region (Data S1). Therefore, the C2 QTL likely harbors multiple loci affecting PZ size variation, potentially also including closely linked, antagonistically loci, which seem to be common in *C. elegans* (Bernstein et al., 2019).

Beyond the technical limitations mentioned above, a number of other issues complicate the interpretation of our findings. First, and foremost, artificial mapping populations, such as the F2 RIL panel used here, may generate novel epistatic interactions through disruption of linked genomic regions. This is particularly relevant for the predominantly selfing *C. elegans*, with low effective recombination leading to strong linkage disequilibrium (Cutter et al., 2009; Lee et al., 2021; Noble et al., 2021b; Rockman et al., 2010). Most prominently, crosses between wild isolates of selfing *Caenorhabditis* have uncovered consistent, strong outbreeding depression due to widespread genetic incompatibilities reducing survival and reproduction (Ben-David et al., 2021; Dolgin et al., 2007; Gimond et al., 2013; Noble et al., 2021b; Seidel et al., 2008). Such epistatic interactions of novel allelic combinations generated in artificial *C. elegans* populations are thus particularly likely to affect traits that involve a polygenic architecture (Bernstein et al., 2019; Campbell et al., 2018; Chandler et al., 2017; Gibson and Dworkin, 2004; Noble et al., 2021a). In other words, epistatic interactions observed in artificial mapping panels need not necessarily reflect epistatic interactions that occur in the wild; nevertheless, their study provides insight into the genetic architecture underlying trait variation.

Fitness relevance of PZ size and its natural variation
While activity of any germ stem cell system is obviously relevant for organismal reproductive fitness, it is not clear if and how observed natural variation in germ cell proliferation (and PZ size) might translate into variation in reproductive fitness, in particular, because *C. elegans* primarily reproduces through self-fertilizing hermaphrodites that sequentially produce sperm, then oocytes for the remainder of life. In laboratory conditions, this causes *C. elegans* fecundity to be limited by the amount of self-sperm (~250) initially produced. PZ size is generally measured, as in our study, during the early adult stage, so that likely most, if not all, of the observed progenitor cells will never develop into mature oocytes and become fertilized under selfing. Although adult PZ size is not a direct proxy for future reproductive potential of selfing hermaphrodites, PZ size also reflects proliferative activity of earlier larval stages, as illustrated by our quantifications of proliferative activity (Figures 2B and 5D). Hence, variation in adult PZ size may reflect differential reproductive investment during larval development, which may trade-off with reduced energy allocation to somatic development.

With regard to reproductive fitness differences of the target isolates in our study, JU751 and JU1200, we have previously shown that selfing JU751 hermaphrodites have significantly reduced brood size relative to JU1200 (Vigne et al., 2021b). This reduction is not caused by differential sperm production but partly due to a major-effect variant causing early matricidal hatching in JU751 (Vigne et al., 2021b). Still, even after correcting for this genetic variant in JU751, brood size remains significantly smaller in JU751 relative to JU1200 (Vigne et al., 2021b). Possibly, reduced germ cell proliferation (and smaller adult PZ) in JU751 could thus reflect a lower reproductive investment, in line with a much shortened reproductive (and life) span. Of course, this scenario is highly speculative and observed differences need not be necessarily adaptive. Similarly, we ignore whether any of the detected QTL, including the *lag-2(cgb1007)* deletion, are under selection. Even if selectively advantageous it remains unclear if QTL were selected because of their effects on germline proliferative activity or unknown pleiotropic effects.

Perspective
There is now ample evidence showing that complex, often highly polygenic, epistatic interactions commonly contribute to phenotypic variation, including variation in developmental systems. Dissecting the molecular nature of such epistatic interactions remains intrinsically difficult, even when limiting efforts to two or three loci. Together with past research, our analysis of natural genetic variation in the activity of the *C. elegans* germ stem cell niche illustrates this problem, reinforcing the view that generalizations on trait architecture based on the isolated study of individual mutations in single genetic backgrounds are potentially misleading. Combining

statistical and molecular genetic approaches is, therefore, essential to uncover and describe, often idiosyncratic, epistatic interactions shaping trait architecture in natural populations.

ACKNOWLEDGEMENTS

We would like to thank Nausicaa Poullet, Anne Vielle and Clotilde Gimond for contributions to experimental work during the early stages of this project. We thank Clotilde Gimond, Laure Mignerot and Joao Picao Osorio for experimental advice, discussion and helpful comments on previous versions of the manuscript. CB acknowledges funding by the Fondation ARC sur la recherche sur le cancer (PJA 20161205047). SRF was supported by a post-doctoral fellowship from the City of Nice, France (Ville de Nice: Aides Individuelles Jeunes Chercheurs). We acknowledge additional financial support by the Centre National de la Recherche Scientifique (CNRS), the Institut national de la santé et de la recherche médicale (Inserm), and the Université Côte d'Azur (UCA).

AUTHOR CONTRIBUTIONS

Conceptualization, S.R.F and C.B; Methodology, S.R.F, A.S. C.B.; Investigation, S.R.F., B.B., and A.S.; Formal Analysis and Visualization, S.R.F.; Writing—Original Draft, S.R.F.; Writing—Review and Editing, S.R.F., C.B.; Supervision and Project Administration, C.B.; Funding Acquisition, C.B., S.R.F.

DECLARATION OF INTERESTS

The authors declare no competing interests.

MATERIALS and METHODS

Experimental model

All *C. elegans* strains used in this study are listed in the Key Resources Table. Standard *C. elegans* methods (Brenner, 1974; Stiernagle, 2006; Wood, 1988) were used to maintain all strains (2.5% agar NGM plates seeded with *E. coli* strain OP50).

Method details
Quantification germline progenitor zone nuclei

Images of dissected gonads fixed in 100% ice cold methanol and stained with DAPI (VECTASHIELD® Antifade Mounting Medium with DAPI) were taken using an Olympus BX61 microscope (RRID:SCR_020343) with a CoolSnap HQ2 camera. Z-stacks with 1μm increments were performed in the DAPI channel at a 40x magnification. The nuclei were counted by hand using ImageJ (RRID:SCR_003070). First, the PZ was defined as the region adjacent to the distal tip cell ending at the transition zone boundary where two or more crescent shape nuclei per row of germ cells can be observed (Crittenden and Kimble, 2008). The stack of images was cropped at the progenitor zone to transition zone boundary and counted.

Quantification of germ cell mitosis

Mitotic germ cell proliferation was quantified in L4 larvae using the Click-IT EdU kit from Thermofisher (Cat: C10338) according to the manufacturer's instructions unless otherwise noted. Briefly, worms were synchronized via hypochlorite treatment (1:1:2 Bleach:1M NaOH:ddH2O for 6 min then spin and wash 3x in M9) and eggs were allowed to hatch overnight in M9 at 20°C. Eggs were plated at a density of ~400 eggs/plate and allowed to develop until L4. Mid-L4 larvae (and later young adults with 1-10 eggs in the uterus) were isolated, washed, and allowed to soak in 100uL of 20mM EdU in M9 buffer for exactly 15 minutes (sufficient time to stain 50-75% of the PZ in young adults). Worms were quickly washed in M9 to remove EdU, and the gonads were released by cutting on adherent glass slides (Fisherbrand™ Superfrost™ Plus Microscope Slides). A small amount of levamisole was used to anesthetize the worms. Dissected gonads were carefully washed on the slides, fixed in 4% PFA, and washed again. Washes were performed

by application of small volumes of solution directly to the tissue on the slides followed by careful aspiration using a mouth pipette. After a final wash in dH2O, tissue was dried on a slide warmer (35°C for 5 min) to adhere tissue to slides. Slides were soaked in 100% MeOH at -20C overnight. The following day, tissues were rehydrated in PBST and EdU staining was performed using the Click-IT EdU kit according to the manufacturer's instructions. The Click-IT staining protocol was repeated once and then slides were mounted with VECTASHIELD® Antifade Mounting Medium with DAPI. Imaging was performed through a 40x objective on an Olympus BX61 microscope (RRID:SCR_020343) with a CoolSnap HQ2 camera. Z-stacks were taken though the distal gonad (one per worm). EdU-positive and DAPI-stained PZ nuclei were counted by hand.

Estimation of PZ size

Quantifying PZ germ cell number can be a rate-limiting step in the scoring of PZ size among many samples. To overcome this obstacle, we developed a simple method of estimating PZ size and used this as a proxy for PZ cell number (Figure 1B and 1C). In brief, lines were synchronized by hypochlorite treatment (1:1:2 Bleach:1M NaOH:ddH2O for 6 min then spun and washed 3x in M9). Eggs were allowed to hatch in M9 buffer at 20C overnight. The following day, the eggs were plated at a density of ~400 worms/plate on standard NGM media seeded with OP50. Larvae were allowed to develop into young adults at 20C. Young adults were collected by washing the plates when the majority of animals showed 1-10 eggs in the uterus. Samples were washed in M9 and fixed in ice-cold methanol for at least 5 min. To prepare samples for imaging, worms were rehydrated in M9 and applied to glass slides with Vectashield mounting medium containing DAPI (Vector Laboratories, Burlingame, CA). The distal germlines of young adult hermaphrodites (containing 1-10 embryos in the uterus) were imaged through a 40x objective on an Olympus BX61 microscope (RRID:SCR_020343) with a CoolSnap HQ2 camera. One gonad arm was imaged per worm, that being the arm which happened to be closer to the objective lens. A single image was taken for each worm, with the plane of section capturing most of the nuclei on one side of the gonad such that crescent-shaped nuclei were visible, and the shape of the gonad in that plane was representative of any plane (Figure 1B). PZ size was then quantified for each image. The progenitor zone was defined as the region adjacent to the distal tip cell ending at the transition zone boundary where two or more crescent-shaped nuclei per row of germ cells can be observed (Crittenden and Kimble, 2008). The PZ was cropped away from all other tissues using ImageJ, and the threshold tool was used to maximize the number of PZ nuclei highlighted while minimizing highlighted background. The number of pixels highlighted was recorded as "PZ Area". This method of estimation correlates well with hand counts of PZ nuclei (Figure 1C).

RIL phenotyping and QTL mapping

We quantified PZ size in a set of 70 SNP-genotyped RILs derived from the two parents, JU1200 and JU751 (Billard et al., 2020; Vigne et al., 2021b). Lines were divided into six scoring blocks and scored over the course of eight weeks. PZ size was normalized across blocks using a correction factor. The correction factor was derived from the least square means of each block (lsmeans package v. 2.30-0) to each observation (correction factor = $LSM_{BlockX}/LSM_{Block1}$). Normalized PZ size was mapped using the software package R/qtl (v. 1.50)(Broman and Saunak, 2009). First, a genetic linkage map was derived from the SNP genotypes of all 70 RILs. Then, single QTL and two-QTL standard interval mapping based on hidden Markov models was used to identify candidate QTL. LOD thresholds were set to be the maximum genome wide LOD scores derived from 5000 random permutations of the data. These candidate QTL served as the starting point for multi-QTL modeling in which the R/qtl software generates "penalized LOD scores" for each tested model. We selected the model with the highest penalized LOD score as per (Broman and Saunak, 2009) (Figure 3J).

NIL generation and chr II QTL genotyping

To validate the genomic interval of the QTL on chromosome II, we constructed nearly isogenic lines (NILs). Two RILs with JU1200 sequence in the QTL region (RIL128 and RIL71) and two RILs with JU751 sequence in the QTL region (RIL127 and RIL54) were backcrossed for 10-12 generations to isolate the region in the opposite parent's background. Lines were selected at each generation based on PCR genotyping of several INDELS in the chromosome II QTL (See Table S13 for primers and Resources Table for lines created).

Generation of allelic replacement lines (ARLs) and CRISPR/Cas-9 gene editing

In order to validate the candidate INDEL upstream of *lag-2* in the chromosome V QTL, we created reciprocal allelic replacement lines (ARLs) using CRISPR/Cas-9 editing using a protocol similar to that described by (Paix et al., 2015). To introduce the deletion into the JU1200 background, we selected a NIL containing JU751 sequence through most of the chromosome II QTL (NIC1701) and injected sgRNAs (Synthego Corp.) to induce a double cut around the insertion site in the *lag-2* promoter (Figure S2, Table S13). We co-injected a single-stranded donor oligonucleotide to act as the repair template (Table S13). *dpy-10* was used in a co-CRISPR strategy according to (Paix et al., 2015). Lines containing successful replacements in *lag-2p* (C1-2 and C1-21) were identified by PCR genotyping of the deletion and sanger sequencing. These lines were then crossed to JU1200 to segregate away the chromosome II containing JU751 sequence leaving us with lines containing both the JU751 version of the chromosome II QTL and the *lag-2p* deletion (NIC1713, NIC1714, NIC1715) as well as a line containing only the *lag-2p* deletion in the JU1200 background (NIC1720). To replace the *lag-2p* deletion in the JU751 background, we selected a NIL containing JU1200 sequence in the chromosome II QTL (NIC1672) and injected guide RNAs (Synthego Corp.) (Table S13) to induce a double cut around the insertion site. We co-injected a double-stranded PCR product to act as the repair template (Table S13). *dpy-10* was used in a co-CRISPR strategy according to (Paix et al., 2015). Lines containing successful replacements (C2-12 and C2-22) were identified by PCR genotyping of the deletion and Sanger sequencing. These lines were then crossed to JU751 to segregate away the chromosome II containing JU1200 sequence, leaving us with lines containing both the JU1200 version of the chromosome II QTL and the *lag-2p* insertion (NIC1716, NIC1717, NIC1719) as well as lines containing only the *lag-2p* insertion in the JU751 background (NIC1723, NIC1724, NIC1725). For a list of all lines created, see Resources Table.

Measurement of lag-2 transcripts via smFISH

Worms were bleach synchronized as above and grown at 20ºC until the appropriate developmental stage. Worms were then washed off plates and fixed in 1mL fresh fixative (4% formaldehyde in PBS) for 40 min. Worms were washed twice in PBS, resuspended in 1mL 70% ethanol, and stored at 4°C for several days. A *lag-2* RNA probe set (Barkoulas et al. 2013) coupled to AF594 (Custom Stellaris Fish Probes, Biosearch Tech, Teddington UK) was resuspended in RNase-free TE buffer (pH8) to make a 100µM stock then diluted 1/30 (working solution) in RNase-free water. Fixed worms were washed 3 min in 1mL wash solution (10% deionized formamide and 2x Saline Sodium Citrate buffer (Ambion) in RNase-free water). Worms were then resuspended in 100µL hybridization solution (10% formamide, 10% dextran sulfate, 2x SSC) + 1 µL of the *lag-2* probe working solution and incubated in the dark at 30°C overnight. The following day, samples were rinsed, washed 30 min at 30°C in wash solution, and stained 30 min at 30°C in 1mL wash solution + DAPI (7.5 µg/mL final, Sigma). Finally, samples were washed twice in PBS and mounted on glass slides in Prolong diamond antifade mounting medium (Eugene, OR). Worms were imaged under epifluorescence on an inverted Zeiss Z.1 microscope (RRID:SCR_020919) with a 100x oil immersion objective. The same illumination and exposure settings (Zen 3.2 Blue Edition software) were used for all samples. Z-stacks were taken through the DTC with a step size of 0.4µm and fluorescent puncta were quantified by hand using FIJI software (NIH, Bethesda, MD, http://rsb.info.nih.gov/ij/)

Quantification and statistical analysis

Statistical analyses were performed using R Statistical Software (R Core Team, 2021) (R version 4.2.0, RStudio 2022.02.3 build 492, RRID:SCR_001905) using generalized linear mixed models. Data were fitted to either gaussian or negative binomial (nbinom2) models using the glmmTMB package v. 1.1.3 (Brooks et al., 2017). Residual diagnostics were performed using the DHARMa package v. 0.4.5 (RRID:SCR_022136) according to the developer's guidelines (Hartig, 2022). When model selection was warranted, the Bayesian information criterion (BIC) was used to select the optimal model. The emmeans package v. 1.7.4-1 (RRID:SCR_018734) was used to calculate model estimated marginal means, standard errors, and Tukey-corrected p-values for pairwise contrasts (Length, 2022). For model fitting of the interaction data set corresponding to Figure 6, D^2 was calculated using the modEvA package v. 3.0 (Márcia Barbosa et al., 2013). Specific details of each data analysis are described in the supplemental table legends, which display the statistical results (Tables S1-S12). Other R packages used for data manipulation and plotting included tidyverse v. 1.3.1 (Wickham et al., 2019) (RRID:SCR_019186), MASS v. 7.3-57 (Venables et al., 2002) (RRID:SCR_019125), ggbeeswarm v. 0.6.0 (Clarke and Sherrill-Mix, 2017), rnaturalearth and rnaturalearthdata v. 0.1.0 (South, 2022), rgeos v. 0.5-9 (Bivand and Rundel, 2021), and sf v. 1.0-7 (Pebesma, 2018).

RESOURCES TABLE

REAGENT or	SOURCE	IDENTIFIER
Antibodies		
Bacterial and virus strains		
Escherichia coli strain OP50	*Caenorhabditis* Genetics Center (https://cgc.umn.edu)	RRID:WB-STRAIN: WBStrain00041969
Biological samples		
Chemicals, peptides, and recombinant proteins		
Critical commercial assays		
Click-iT™ EdU Cell Proliferation Kit for Imaging, Alexa Fluor™ 555 dye	Invitrogen™	Cat. # C10338
Deposited data		
Experimental models: Cell lines		
Experimental models: Organisms/strains		
AF16, *C. briggsae* wild isolate	*Caenorhabditis* Genetics Center (https://cgc.umn.edu)	RRID:WB-STRAIN:WBStrain00040935
CB4856, *C. elegans* wild isolate	*C. elegans* Natural Diversity Resource (https://elegansvariation.org)	RRID:WB-STRAIN:WBStrain00004602
CX11262, *C. elegans* wild isolate	*C. elegans* Natural Diversity Resource (https://elegansvariation.org)	RRID:WB-STRAIN:WBStrain00005299
HK104, *C. briggsae* wild isolate	*Caenorhabditis* Genetics Center (https://cgc.umn.edu)	RRID:WB-STRAIN:WBStrain00041077
JU576, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00022973
JU577, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00022974
JU584, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00022981
JU636, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	/RRID:WB-STRAIN:WBStrain00023012
JU637, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00023013
JU644, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00023020
JU656, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00023028
JU660, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00023032
JU751, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00023065
JU752, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00023066
JU753, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00023067

JU754, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00023068
JU755, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:
JU756, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:
JU775, *C. elegans* wild isolate	*C. elegans* Natural Diversity Resource (https://elegansvariation.org)	CeNDR (https://elegansvariation.org)/RRID:WB-STRAIN:WBStrain00023072
JU815, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00023097
JU816, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00023098
JU817, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00023099
JU818, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00023100
JU821, *C. elegans* wild isolate	M.-A. Félix (Barrière and Félix, 2007)	RRID:WB-STRAIN:WBStrain00023103
JU1200, *C. elegans* wild isolate	*C. elegans* Natural Diversity Resource (https://elegansvariation.org)	RRID:WB-STRAIN:WBStrain00023141
JU1491, *C. elegans* wild isolate	*C. elegans* Natural Diversity Resource (https://elegansvariation.org)	RRID:WB-STRAIN:WBStrain00023219
N2, *C. elegans* lab reference strain	*Caenorhabditis* Genetics Center (https://cgc.umn.edu)	RRID:WB-STRAIN:WBStrain00000001
NIC601, *C. elegans* F2 RIL(1) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC606, *C. elegans* F2 RIL(6) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC610, *C. elegans* F2 RIL(10) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC617, *C. elegans* F2 RIL(17) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC619, *C. elegans* F2 RIL(19) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC621, *C. elegans* F2 RIL(21) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC623, *C. elegans* F2 RIL(23) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC624, *C. elegans* F2 RIL(24) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC625, *C. elegans* F2 RIL(25) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC626, *C. elegans* F2 RIL(26) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC628, *C. elegans* F2 RIL(28) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A

NIC630, *C. elegans* F2 RIL(30) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC631, *C. elegans* F2 RIL(31) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC633, *C. elegans* F2 RIL(33) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC636, *C. elegans* F2 RIL(36) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC640, *C. elegans* F2 RIL(40) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC643, *C. elegans* F2 RIL(43) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC646, *C. elegans* F2 RIL(46) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC647, *C. elegans* F2 RIL(47) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC649, *C. elegans* F2 RIL(49) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC650, *C. elegans* F2 RIL(50) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC653, *C. elegans* F2 RIL(53) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC654, *C. elegans* F2 RIL(54) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC655, *C. elegans* F2 RIL(55) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC660, *C. elegans* F2 RIL(60) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC661, *C. elegans* F2 RIL(61) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC662, *C. elegans* F2 RIL(62) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC664, *C. elegans* F2 RIL(64) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC665, *C. elegans* F2 RIL(65) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC666, *C. elegans* F2 RIL(66) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC667, *C. elegans* F2 RIL(67) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC668, *C. elegans* F2 RIL(68) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC671, *C. elegans* F2 RIL(71) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC675, *C. elegans* F2 RIL(75) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC678, *C. elegans* F2 RIL(78) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC680, *C. elegans* F2 RIL(80) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A

NIC681, *C. elegans* F2 RIL(81) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC685, *C. elegans* F2 RIL(85) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC686, *C. elegans* F2 RIL(86) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC687, *C. elegans* F2 RIL(87) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC691, *C. elegans* F2 RIL(91) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC692, *C. elegans* F2 RIL(92) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC694, *C. elegans* F2 RIL(94) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC696, *C. elegans* F2 RIL(96) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC697, *C. elegans* F2 RIL(97) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC698, *C. elegans* F2 RIL(98) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC699, *C. elegans* F2 RIL(99) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC700, *C. elegans* F2 RIL(100) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC702, *C. elegans* F2 RIL(102) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC703, *C. elegans* F2 RIL(103) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC704, *C. elegans* F2 RIL(104) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC705, *C. elegans* F2 RIL(105) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC707, *C. elegans* F2 RIL(107) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC709, *C. elegans* F2 RIL(109) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC710, *C. elegans* F2 RIL(110) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC711, *C. elegans* F2 RIL(111) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC713, *C. elegans* F2 RIL(113) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC716, *C. elegans* F2 RIL(116) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC717, *C. elegans* F2 RIL(117) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC721, *C. elegans* F2 RIL(121) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC727, *C. elegans* F2 RIL(127) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A

NIC728, *C. elegans* F2 RIL(128) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC729, *C. elegans* F2 RIL(129) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC730, *C. elegans* F2 RIL(130) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC731, *C. elegans* F2 RIL(131) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC734, *C. elegans* F2 RIL(134) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC735, *C. elegans* F2 RIL(135) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC736, *C. elegans* F2 RIL(136) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC739, *C. elegans* F2 RIL(139) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC740, *C. elegans* F2 RIL(140) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC741, *C. elegans* F2 RIL(141) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC742, *C. elegans* F2 RIL(142) JU751 x JU1200	C. Braendle (Vigne et al., 2021b)	N/A
NIC1671, *C. elegans* NIL, JU751 background	This study	N/A
NIC1672, *C. elegans* NIL, JU751 background	This study	N/A
NIC1673, *C. elegans* NIL, JU751 background	This study	N/A
NIC1675, *C. elegans* NIL, JU751 background	This study	N/A
NIC1676, *C. elegans* NIL, JU751 background	This study	N/A
NIC1697, *C. elegans* NIL, JU1200 background	This study	N/A
NIC1701, *C. elegans* NIL, JU1200 background	This study	N/A
NIC1702, *C. elegans* NIL, JU1200 background	This study	N/A
NIC1713, *C. elegans* ARL, JU1200 background	This study	N/A
NIC1714, *C. elegans* ARL, JU1200 background	This study	N/A
NIC1715, *C. elegans* ARL, JU1200 background	This study	N/A
NIC1716, *C. elegans* ARL, JU751 background	This study	N/A
NIC1717, *C. elegans* ARL, JU751 background	This study	N/A
NIC1719, *C. elegans* ARL, JU751 background	This study	N/A

NIC1720, *C. elegans* ARL, JU1200 background	This study	N/A
NIC1723, *C. elegans* ARL, JU751 background	This study	N/A
NIC1724, *C. elegans* ARL, JU751 background	This study	N/A
NIC1725, *C. elegans* ARL, JU751 background	This study	N/A
NIC2091, *C. elegans* NIL, JU1200 background	This study	N/A
MY16, *C. elegans* wild isolate	*C. elegans* Natural Diversity Resource (https://elegansvariation.org)	RRID:WB-STRAIN:WBStrain00027662
Oligonucleotides		
Oligonucleotides used for generation of ARLs, See Table S13.	This study	N/A
Genotyping primers, See Table S13	This study	N/A
lag-2 RNA probe for smFISH, See Table S13	Custom Stellaris Fish Probes, Biosearch Tech, Teddington UK	N/A
Recombinant DNA		
Software and algorithms		
ImageJ	(Schneider et al., 2012)	https://imagej.net/downloads
FIJI	(Schindelin et al., 2012)	https://imagej.net/downloads
R Statistical Software v. 4.2.0	(R Core Team, 2021)	RRID:SCR_001905
RStudio 2022.02.03 build 492	(RStudio Team, 2020)	RRID:SCR_000432
glmmTMB package for R (v. 1.1.3)	(Brooks et al., 2017)	https://cran.r-project.org/web/packages/glmmTMB/index.html
DHARMa package for R (v. 0.4.5)	(Hartig, 2022)	RRID:SCR_022136
Emmeans package for R (v. 1.7.4-1)	(Length, 2022)	RRID:SCR_018734
modEvA package for R (v. 3.0)	(Márcia Barbosa et al., 2013)	https://cran.r-project.org/web/packages/modEvA/index.html
R/qtl package for R (v 1.50)	(Broman and Saunak, 2009)	RRID:SCR_009085

References

Agata, K., Nakajima, E., Funayama, N., Shibata, N., Saito, Y., Umesono, Y., 2006. Two different evolutionary origins of stem cell systems and their molecular basis. Semin. Cell Dev. Biol., Olfaction 17, 503–509. https://doi.org/10.1016/j.semcdb.2006.05.004

Angelo, G., Van Gilst, M.R., 2009. Starvation Protects Germline Stem Cells and Extends Reproductive Longevity in C. elegans. Science 326, 954–958. https://doi.org/10.1126/science.1178343

Aprison, E.Z., Ruvinsky, I., 2016. Sexually Antagonistic Male Signals Manipulate Germline and Soma of C. elegans Hermaphrodites. Curr. Biol. 26, 2827–2833. https://doi.org/10.1016/j.cub.2016.08.024

Austin, J., Kimble, J., 1989. Transcript analysis of glp-1 and lin-12, homologous genes required for cell interactions during development of C. elegans. Cell 58, 565–571. https://doi.org/10.1016/0092-8674(89)90437-6

Barkoulas, M., van Zon, J.S., Milloz, J., van Oudenaarden, A., Félix, M.-A., 2013. Robustness and Epistasis in the C. elegans Vulval Signaling Network Revealed by Pathway Dosage Modulation. Dev. Cell 24, 64–75. https://doi.org/10.1016/j.devcel.2012.12.001

Barrière, A., Félix, M.-A., 2007. Temporal Dynamics and Linkage Disequilibrium in Natural Caenorhabditis elegans Populations. Genetics 176, 999–1011. https://doi.org/10.1534/genetics.106.067223

Ben-David, E., Pliota, P., Widen, S.A., Koreshova, A., Lemus-Vergara, T., Verpukhovskiy, P., Mandali, S., Braendle, C., Burga, A., Kruglyak, L., 2021. Ubiquitous Selfish Toxin-Antidote Elements in Caenorhabditis Species. Curr. Biol. 31, 990-1001.e5. https://doi.org/10.1016/j.cub.2020.12.013

Bernstein, M.R., Zdraljevic, S., Andersen, E.C., Rockman, M.V., 2019. Tightly linked antagonistic-effect loci underlie polygenic phenotypic variation in *C. elegans*. Evol. Lett. 3, 462–473. https://doi.org/10.1002/evl3.139

Berry, L.W., Westlund, B., Schedl, T., 1997. Germ-line tumor formation caused by activation of glp-1, a Caenorhabditis elegans member of the Notch family of receptors. Development 124, 925–936. https://doi.org/10.1242/dev.124.4.925

Billard, B., Vigne, P., Braendle, C., 2020. A Natural Mutational Event Uncovers a Life History Trade-Off via Hormonal Pleiotropy. Curr. Biol. 30, 4142-4154.e9. https://doi.org/10.1016/j.cub.2020.08.004

Bivand, R., Rundel, C., 2021. rgeos: Interface to Geometry Engine - Open Source ('GEOS'). https://r-forge.r-project.org/projects/rgeos/.

Boyle, E.A., Li, Y.I., Pritchard, J.K., 2017. An Expanded View of Complex Traits: From Polygenic to Omnigenic. Cell 169, 1177–1186. https://doi.org/10.1016/j.cell.2017.05.038

Brenner, S., 1974. THE GENETICS OF CAENORHABDITIS ELEGANS. Genetics 77, 71–94. https://doi.org/10.1093/genetics/77.1.71

Broman, K.W., Saunak, S., 2009. A Guide to QTL Mapping with R/qtl. Springer Science+Business Media, LLC.

Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A., Skaug, H.J., Machler, M., Bolker, B.M., 2017. glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. R J. 9, 378–400. https://doi.org/10.3929/ethz-b-000240890

Bubnell, J.E., Ulbing, C.K.S., Fernandez Begne, P., Aquadro, C.F., 2022. Functional Divergence of the bag-of-marbles Gene in the Drosophila melanogaster Species Group. Mol. Biol. Evol. 39, msac137. https://doi.org/10.1093/molbev/msac137

Campbell, R.F., McGrath, P.T., Paaby, A.B., 2018. Analysis of Epistasis in Natural Traits Using Model Organisms. Trends Genet. 34, 883–898. https://doi.org/10.1016/J.TIG.2018.08.002

Chandler, C.H., Chari, S., Kowalski, A., Choi, L., Tack, D., DeNieu, M., Pitchers, W., Sonnenschein, A., Marvin, L., Hummel, K., Marier, C., Victory, A., Porter, C., Mammel, A., Holms, J., Sivaratnam, G., Dworkin, I., 2017. How well do you know your mutation? Complex effects of genetic background on expressivity, complementation, and ordering of allelic effects. PLOS Genet. 13, e1007075. https://doi.org/10.1371/journal.pgen.1007075

Chen, J., Mohammad, A., Pazdernik, N., Huang, H., Bowman, B., Tycksen, E., Schedl, T., 2020. GLP-1 Notch—LAG-1 CSL control of the germline stem cell fate is mediated by transcriptional targets lst-1 and sygl-1. PLOS Genet. 16, e1008650. https://doi.org/10.1371/JOURNAL.PGEN.1008650

Chesney, M.A., Lam, N., Morgan, D.E., Phillips, B.T., Kimble, J., 2009. C. elegans HLH-2/E/Daughterless controls key regulatory cells during gonadogenesis. Dev. Biol. 331, 14–25. https://doi.org/10.1016/j.ydbio.2009.04.015

Cheverud, J.M., Routman, E.J., 1995. Epistasis and its contribution to genetic variance components. Genetics 139, 1455–1461. https://doi.org/10.1093/genetics/139.3.1455

Choi, J.Y., Aquadro, C.F., 2015. Molecular Evolution of Drosophila Germline Stem Cell and Neural Stem Cell Regulating Genes. Genome Biol. Evol. 7, 3097–3114. https://doi.org/10.1093/gbe/evv207

Clarke, E., Sherrill-Mix, S., 2017. ggbeeswarm: Categorical Scatter (Violin Point) Plots. R package version 0.6.0.

Cook, D.E., Zdraljevic, S., Roberts, J.P., Andersen, E.C., 2017. CeNDR, the Caenorhabditis elegans natural diversity resource. Nucleic Acids Res. 45, D650–D657. https://doi.org/10.1093/nar/gkw893

Crittenden, S.L., Kimble, J., 2008. Analysis of the C. elegans Germline Stem Cell Region, in: Hou, S.X., Singh, S.R. (Eds.), Germline Stem Cells, Methods in Molecular Biology™. Humana Press, Totowa, NJ, pp. 27–44. https://doi.org/10.1007/978-1-60327-214-8_2

Cutter, A.D., Dey, A., Murray, R.L., 2009. Evolution of the Caenorhabditis elegans Genome. Mol. Biol. Evol. 26, 1199–1234. https://doi.org/10.1093/molbev/msp048

Dalfó, D., Michaelson, D., Hubbard, E.J.A., 2012. Sensory Regulation of the C. elegans Germline through TGF-β-Dependent Signaling in the Niche. Curr. Biol. 22, 712–719. https://doi.org/10.1016/j.cub.2012.02.064

Dolgin, E.S., Charlesworth, B., Baird, S.E., Cutter, A.D., 2007. Inbreeding and Outbreeding Depression in Caenorhabditis Nematodes. Evolution 61, 1339–1352. https://doi.org/10.1111/j.1558-5646.2007.00118.x

Duveau, F., Félix, M.-A., 2012. Role of Pleiotropy in the Evolution of a Cryptic Developmental Variation in Caenorhabditis elegans. PLOS Biol. 10, e1001230. https://doi.org/10.1371/journal.pbio.1001230

Extavour, C.G., Akam, M., 2003. Mechanisms of germ cell specification across the metazoans: epigenesis and preformation. Development 130, 5869–5884. https://doi.org/10.1242/dev.00804

Falconer, D.S., 1996. Introduction to Quantitative Genetics. Pearson Education.

Flores, H.A., DuMont, V.L.B., Fatoo, A., Hubbard, D., Hijji, M., Barbash, D.A., Aquadro, C.F., 2015. Adaptive Evolution of Genes Involved in the Regulation of Germline Stem Cells in Drosophila melanogaster and D. simulans. G3 GenesGenomesGenetics 5, 583–592. https://doi.org/10.1534/g3.114.015875

Forsberg, S.K.G., Bloom, J.S., Sadhu, M.J., Kruglyak, L., Carlborg, Ö., 2017. Accounting for genetic interactions improves modeling of individual quantitative trait phenotypes in yeast. Nat. Genet. 49, 497–503. https://doi.org/10.1038/ng.3800

Gao, D., Kimble, J., 1995. APX-1 can substitute for its homolog LAG-2 to direct cell interactions throughout Caenorhabditis elegans development. Proc. Natl. Acad. Sci. 92, 9839–9842. https://doi.org/10.1073/pnas.92.21.9839

Gibson, G., Dworkin, I., 2004. Uncovering cryptic genetic variation. Nat. Rev. Genet. 5, 681–690. https://doi.org/10.1038/nrg1426

Gibson, G., Hogness, D.S., 1996. Effect of Polymorphism in the Drosophila Regulatory Gene Ultrabithorax on Homeotic Stability. Science 271, 200–203. https://doi.org/10.1126/science.271.5246.200

Gimond, C., Jovelin, R., Han, S., Ferrari, C., Cutter, A.D., Braendle, C., 2013. Outbreeding Depression with Low Genetic Variation in Selfing Caenorhabditis Nematodes. Evolution 67, 3087–3101. https://doi.org/10.1111/evo.12203

Gopal, S., Amran, A., Elton, A., Ng, L., Pocock, R., 2021. A somatic proteoglycan controls Notch-directed germ cell fate. Nat. Commun. 12, 6708. https://doi.org/10.1038/s41467-021-27039-4

Gordon, K., 2020. Recent Advances in the Genetic, Anatomical, and Environmental Regulation of the C. elegans Germ Line Progenitor Zone. J. Dev. Biol. 2020 Vol 8 Page 14 8, 14. https://doi.org/10.3390/JDB8030014

Hansen, D., Schedl, T., 2013. Stem Cell Proliferation Versus Meiotic Fate Decision in Caenorhabditis elegans, in: Schedl, T. (Ed.), Germ Cell Development in C. Elegans, Advances in Experimental Medicine and Biology. Springer, New York, NY, pp. 71–99. https://doi.org/10.1007/978-1-4614-4015-4_4

Hartig, F., 2022. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level/Mixed) Regression Models.

Haupt, K.A., Law, K.T., Enright, A.L., Kanzler, C.R., Shin, H., Wickens, M., Kimble, J., 2020. A PUF Hub Drives Self-Renewal in Caenorhabditis elegans Germline Stem Cells. Genetics 214, 147–161. https://doi.org/10.1534/GENETICS.119.302772

Henderson, S.T., Gao, D., Lambie, E.J., Kimble, J., 1994. lag-2 may encode a signaling ligand for the GLP-1 and LIN-12 receptors of C. elegans. Dev. Camb. Engl. 120, 2913–2924.

Hill, W.G., Goddard, M.E., Visscher, P.M., 2008. Data and Theory Point to Mainly Additive Genetic Variance for Complex Traits. PLOS Genet. 4, e1000008. https://doi.org/10.1371/journal.pgen.1000008

Huang, W., Mackay, T.F.C., 2016. The Genetic Architecture of Quantitative Traits Cannot Be Inferred from Variance Component Analysis. PLOS Genet. 12, e1006421. https://doi.org/10.1371/journal.pgen.1006421

Huang, X., Effgen, S., Meyer, R.C., Theres, K., Koornneef, M., 2012. Epistatic Natural Allelic Variation Reveals a Function of AGAMOUS-LIKE6 in Axillary Bud Formation in Arabidopsis. Plant Cell 24, 2364–2379. https://doi.org/10.1105/tpc.112.099168

Hubbard, E.J.A., Korta, D.Z., Dalfó, D., 2013. Physiological Control of Germline Development, in: Schedl, T. (Ed.), Germ Cell Development in C. Elegans, Advances in Experimental Medicine and Biology. Springer, New York, NY, pp. 101–131. https://doi.org/10.1007/978-1-4614-4015-4_5

Hubbard, E.J.A., Schedl, T., 2019. Biology of the *Caenorhabditis elegans* Germline Stem Cell System. Genetics 213, 1145–1188. https://doi.org/10.1534/genetics.119.300238

Jakobson, C.M., Jarosz, D.F., 2020. What Has a Century of Quantitative Genetics Taught Us about Nature's Genetic Tool Kit? Annu. Rev. Genet. 54, 439–464. https://doi.org/10.1146/ANNUREV-GENET-021920-102037

Karp, X., Greenwald, I., 2003. Post-transcriptional regulation of the E/Daughterless ortholog HLH-2, negative feedback, and birth order bias during the AC/VU decision in C. elegans. Genes Dev. 17, 3100–3111. https://doi.org/10.1101/gad.1160803

Killian, D.J., Hubbard, E.J.A., 2005. Caenorhabditis elegans germline patterning requires coordinated development of the somatic gonadal sheath and the germ line. Dev. Biol. 279, 322–335. https://doi.org/10.1016/j.ydbio.2004.12.021

Kimble, J., Crittenden, S.L., 2007. Controls of germline stem cells, entry into meiosis, and the sperm/oocyte decision in Caenorhabditis elegans. Annu. Rev. Cell Dev. Biol. 23, 405–433. https://doi.org/10.1146/annurev.cellbio.23.090506.123326

Kocsisova, Z., Bagatelas, E.D., Santiago-Borges, J., Lei, H.C., Egan, B.M., Mosley, M.C., Schneider, D.L., Schedl, T., Kornfeld, K., 2022. Notch signaling in germ line stem cells controls reproductive aging in C. elegans. https://doi.org/10.1101/2022.03.04.482923

Koneru, S.L., Hintze, M., Katsanos, D., Barkoulas, M., 2021. Cryptic genetic variation in a heat shock protein modifies the outcome of a mutation affecting epidermal stem cell development in C. elegans. Nat. Commun. 12, 3263. https://doi.org/10.1038/s41467-021-23567-1

Korta, D.Z., Tuck, S., Hubbard, E.J.A., 2012. S6K links cell fate, cell cycle and nutrient response in C. elegans germline stem/progenitor cells. Development 139, 859–870. https://doi.org/10.1242/dev.074047

Lachowiec, J., Shen, X., Queitsch, C., Carlborg, Ö., 2015. A Genome-Wide Association Analysis Reveals Epistatic Cancellation of Additive Genetic Variance for Root Length in Arabidopsis thaliana. PLOS Genet. 11, e1005541. https://doi.org/10.1371/journal.pgen.1005541

Laws, K.M., Drummond-Barbosa, D., 2017. Control of Germline Stem Cell Lineages by Diet and Physiology, in: Arur, S. (Ed.), Signaling-Mediated Control of Cell Division : From Oogenesis to Oocyte-to-Embryo Development. Springer International Publishing, Cham, pp. 67–99. https://doi.org/10.1007/978-3-319-44820-6_3

Lee, D., Zdraljevic, S., Stevens, L., Wang, Y., Tanny, R.E., Crombie, T.A., Cook, D.E., Webster, A.K., Chirakar, R., Baugh, L.R., Sterken, M.G., Braendle, C., Félix, M.-A., Rockman, M.V., Andersen, E.C., 2021. Balancing selection maintains hyper-divergent haplotypes in Caenorhabditis elegans. Nat. Ecol. Evol. 5, 794–807. https://doi.org/10.1038/s41559-021-01435-x

Lehmann, R., 2012. Germline Stem Cells: Origin and Destiny. Cell Stem Cell 10, 729–739. https://doi.org/10.1016/j.stem.2012.05.016

Length, R., 2022. emmeans: Estimated Marginal Means, aka Least-Squares Means.

Li, X., Singh, N., Miller, C., Washington, I., Sosseh, B., Gordon, K.L., 2022. The C. elegans gonadal sheath Sh1 cells extend asymmetrically over a differentiating germ cell population in the proliferative zone. eLife 11, e75497. https://doi.org/10.7554/eLife.75497

Mackay, T.F.C., 2014. Epistasis and quantitative traits: using model organisms to study gene–gene interactions. Nat. Rev. Genet. 15, 22–33. https://doi.org/10.1038/nrg3627

Mäki-Tanila, A., Hill, W.G., 2014. Influence of Gene Interaction on Complex Trait Variation with Multilocus Models. Genetics 198, 355–367. https://doi.org/10.1534/genetics.114.165282

Márcia Barbosa, A., Real, R., Muñoz, A.-R., Brown, J.A., 2013. New measures for assessing model equilibrium and prediction mismatch in species distribution models. Divers. Distrib. 19, 1333–1338. https://doi.org/10.1111/ddi.12100

McCarter, J., Bartlett, B., Dang, T., Schedl, T., 1997. Soma–Germ Cell Interactions in Caenorhabditis elegans:Multiple Events of Hermaphrodite Germline Development Require the Somatic Sheath and Spermathecal Lineages. Dev. Biol. 181, 121–143. https://doi.org/10.1006/dbio.1996.8429

Michaelson, D., Korta, D.Z., Capua, Y., Hubbard, E.J.A., 2010. Insulin signaling promotes germline proliferation in C. elegans. Development 137, 671–680. https://doi.org/10.1242/dev.042523

Monnahan, P.J., Kelly, J.K., 2015. Epistasis Is a Major Determinant of the Additive Genetic Variance in Mimulus guttatus. PLOS Genet. 11, e1005201. https://doi.org/10.1371/journal.pgen.1005201

Morrison, S.J., Spradling, A.C., 2008. Stem Cells and Niches: Mechanisms That Promote Stem Cell Maintenance throughout Life. Cell 132, 598–611. https://doi.org/10.1016/j.cell.2008.01.038

Nadarajan, S., Govindan, J.A., McGovern, M., Hubbard, E.J.A., Greenstein, D., 2009. MSP and GLP-1/Notch signaling coordinately regulate actomyosin-dependent cytoplasmic streaming and oocyte growth in C. elegans. Dev. Camb. Engl. 136, 2223–2234. https://doi.org/10.1242/dev.034603

Noble, L.M., Rockman, M.V., Teotónio, H., 2021a. Gene-level quantitative trait mapping in Caenorhabditis elegans. G3 GenesGenomesGenetics 11, jkaa061. https://doi.org/10.1093/g3journal/jkaa061

Noble, L.M., Yuen, J., Stevens, L., Moya, N., Persaud, R., Moscatelli, M., Jackson, J.L., Zhang, G., Chitrakar, R., Baugh, L.R., Braendle, C., Andersen, E.C., Seidel, H.S., Rockman, M.V., 2021b. Selfing is the safest sex for Caenorhabditis tropicalis. eLife 10, e62587. https://doi.org/10.7554/eLife.62587

Paaby, A.B., White, A.G., Riccardi, D.D., Gunsalus, K.C., Piano, F., Rockman, M.V., 2015. Wild worm embryogenesis harbors ubiquitous polygenic modifier variation. eLife 4, e09178. https://doi.org/10.7554/eLife.09178

Paix, A., Folkmann, A., Rasoloson, D., Seydoux, G., 2015. High Efficiency, Homology-Directed Genome Editing in Caenorhabditis elegans Using CRISPR-Cas9 Ribonucleoprotein Complexes. Genetics 201, 47–54. https://doi.org/10.1534/genetics.115.179382

Pebesma, E., 2018. Simple Features for R: Standardized Support for Spatial Vector Data. R J. 10, 439–446. https://doi.org/10.32614/RJ-2018-009

Pekar, O., Ow, M.C., Hui, K.Y., Noyes, M.B., Hall, S.E., Hubbard, E.J.A., 2017. Linking the environment, DAF-7/TGFβ signaling and LAG-2/DSL ligand expression in the germline stem cell niche. Dev. Camb. Engl. 144, 2896–2906. https://doi.org/10.1242/dev.147660

Phillips, P.C., 2008. Epistasis - The essential role of gene interactions in the structure and evolution of genetic systems. Nat. Rev. Genet. https://doi.org/10.1038/nrg2452

Poullet, N., Vielle, A., Gimond, C., Carvalho, S., Teotónio, H., Braendle, C., 2016. Complex heterochrony underlies the evolution of Caenorhabditis elegans hermaphrodite sex allocation. Evolution 70, 2357–2369. https://doi.org/10.1111/evo.13032

Poullet, N., Vielle, A., Gimond, C., Ferrari, C., Braendle, C., 2015. Evolutionarily divergent thermal sensitivity of germline development and fertility in hermaphroditic Caenorhabditis nematodes. Evol. Dev. 17, 380–397. https://doi.org/10.1111/ede.12170

R Core Team, 2021. R: A language and environment for staistical computing.

Raj, A., Rifkin, S.A., Andersen, E., van Oudenaarden, A., 2010. Variability in gene expression underlies incomplete penetrance. Nature 463, 913–918. https://doi.org/10.1038/nature08781

Raj, A., van den Bogaard, P., Rifkin, S.A., van Oudenaarden, A., Tyagi, S., 2008. Imaging individual mRNA molecules using multiple singly labeled probes. Nat. Methods 5, 877–879. https://doi.org/10.1038/nmeth.1253

Rockman, M.V., Skrovanek, S.S., Kruglyak, L., 2010. Selection at Linked Sites Shapes Heritable Phenotypic Variation in C. elegans. Science 330, 372–376. https://doi.org/10.1126/science.1194208

RStudio Team, 2020. RStudio: Integrated Development for R.

Rudel, D., Kimble, J., 2001. Conservation of glp-1 Regulation and Function in Nematodes. Genetics 157, 639–654. https://doi.org/10.1093/genetics/157.2.639

Sackton, T.B., Hartl, D.L., 2016. Genotypic Context and Epistasis in Individuals and Populations. Cell 166, 279–287. https://doi.org/10.1016/j.cell.2016.06.047

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: an open-source platform for biological-image analysis. Nat. Methods 9, 676–682. https://doi.org/10.1038/nmeth.2019

Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of image analysis. Nat. Methods 9, 671–675. https://doi.org/10.1038/nmeth.2089

Seidel, H.S., Kimble, J., 2015. Cell-cycle quiescence maintains Caenorhabditis elegans germline stem cells independent of GLP-1/Notch. eLife 4, e10832. https://doi.org/10.7554/eLife.10832

Seidel, H.S., Rockman, M.V., Kruglyak, L., 2008. Widespread Genetic Incompatibility in C. Elegans Maintained by Balancing Selection. Science 319, 589–594. https://doi.org/10.1126/science.1151107

South, A., 2022. rnaturalearth: World Map Data from Natural Earth. https://docs.ropensci.org/rnaturalearth.

Sowa, J.N., Mutlu, A.S., Xia, F., Wang, M.C., 2015. Olfaction Modulates Reproductive Plasticity through Neuroendocrine Signaling in Caenorhabditis elegans. Curr. Biol. 25, 2284–2289. https://doi.org/10.1016/j.cub.2015.07.023

Starich, T.A., Hall, D.H., Greenstein, D., 2014. Two Classes of Gap Junction Channels Mediate Soma-Germline Interactions Essential for Germline Proliferation and Gametogenesis in Caenorhabditis elegans. Genetics 198, 1127–1153. https://doi.org/10.1534/genetics.114.168815

Steiner, C.C., Weber, J.N., Hoekstra, H.E., 2007. Adaptive Variation in Beach Mice Produced by Two Interacting Pigmentation Genes. PLOS Biol. 5, e219. https://doi.org/10.1371/journal.pbio.0050219

Stiernagle, T., 2006. Maintenance of C. elegans. WormBook. https://doi.org/10.1895/wormbook.1.101.1

Tax, F.E., Yeargers, J.J., Thomas, J.H., 1994. Sequence of C. elegans lag-2 reveals a cell-signalling domain shared with Delta and Serrate of Drosophila. Nature 368, 150–154. https://doi.org/10.1038/368150a0

Thondamal, M., Witting, M., Schmitt-Kopplin, P., Aguilaniu, H., 2014. Steroid hormone signalling links reproduction to lifespan in dietary-restricted Caenorhabditis elegans. Nat. Commun. 5, 4879. https://doi.org/10.1038/ncomms5879

Tolkin, T., Hubbard, E.J.A., 2021. Germline Stem and Progenitor Cell Aging in C. elegans. Front. Cell Dev. Biol. 9, 699671. https://doi.org/10.3389/fcell.2021.699671

Toomey, M.E., Frydman, H.M., 2014. Extreme Divergence of Wolbachia Tropism for the Stem-Cell-Niche in the Drosophila Testis. PLOS Pathog. 10, e1004577. https://doi.org/10.1371/journal.ppat.1004577

Vanhaeren, H., Gonzalez, N., Coppens, F., De Milde, L., Van Daele, T., Vermeersch, M., Eloy, N.B., Storme, V., Inzé, D., 2014. Combining growth-promoting genes leads to positive epistasis in Arabidopsis thaliana. eLife 3, e02252. https://doi.org/10.7554/eLife.02252

Venables, W.N., Ripley, B.D., Venables, W.N., 2002. Modern applied statistics with S.

Vigne, P., Gimond, C., Ferrari, C., Vielle, A., Hallin, J., Pino-Querido, A., Mouridi, S.E., Mignerot, L., Frøkjær-Jensen, C., Boulin, T., Teotónio, H., Braendle, C., 2021a. A single-nucleotide change underlies the genetic assimilation of a plastic trait. Sci. Adv. 7. https://doi.org/10.1126/SCIADV.ABD9941

Vigne, P., Gimond, C., Ferrari, C., Vielle, A., Hallin, J., Pino-Querido, A., Mouridi, S.E., Mignerot, L., Frøkjær-Jensen, C., Boulin, T., Teotónio, H., Braendle, C., 2021b. A single-nucleotide change underlies the genetic assimilation of a plastic trait. Sci. Adv. 7. https://doi.org/10.1126/SCIADV.ABD9941

Vonesch, S.C., Lamparter, D., Mackay, T.F.C., Bergmann, S., Hafen, E., 2016. Genome-Wide Analysis Reveals Novel Regulators of Growth in Drosophila melanogaster. PLOS Genet. 12, e1005616. https://doi.org/10.1371/journal.pgen.1005616

Weinreich, D.M., Lan, Y., Wylie, C.S., Heckendorn, R.B., 2013. Should evolutionary geneticists worry about higher-order epistasis? Curr. Opin. Genet. Dev., Genetics of system biology 23, 700–707. https://doi.org/10.1016/j.gde.2013.10.007

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L.D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T.L., Miller, E., Bache, S.M., Müller, K., Ooms, J., Robinson, D., Seidel, D.P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., Yutani, H., 2019. Welcome to the Tidyverse. J. Open Source Softw. 4, 1686. https://doi.org/10.21105/joss.01686

Wood, W.B., 1988. Determination of Pattern and Fate in Early Embryos of Caenorhabditis elegans, in: Browder, L.W. (Ed.), The Molecular Biology of Cell Determination and Cell Differentiation, Developmental Biology. Springer US, Boston, MA, pp. 57–78. https://doi.org/10.1007/978-1-4615-6817-9_2

Yochem, J., Greenwald, I., 1989. glp-1 and lin-12, genes implicated in distinct cell-cell interactions in C. elegans, encode similar transmembrane proteins. Cell 58, 553–563. https://doi.org/10.1016/0092-8674(89)90436-4