Widespread exposure to SARS-CoV-2 in wildlife communities

Amanda R. Goldberg[1], Kate E. Langwig[1], Katherine L. Brown[2,3,7], Jeffrey Marano[4,5], Pallavi Rai[4], Kelsie M. King[6], Amanda K. Sharp[6], Alessandro Ceci[7], Christopher D. Kailing[1], Macy J. Kailing[1], Russell Briggs[7], Matthew G. Urbano[7], Clinton Roby[7], Anne M. Brown[6,8,9,10,11], James Weger-Lucarelli[3,4], Carla V. Finkielstein[1,2,3,7,10,11,12]*†, Joseph R. Hoyt[1]*†

[1]*Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia, USA*
[2]*Virginia Tech Carilion School of Medicine, Virginia Tech, Roanoke, VA, USA*
[3]*Center for Emerging, Zoonotic, and Arthropod-borne Pathogens, Virginia Tech, Blacksburg, VA, USA*
[4]*Department of Biomedical Sciences and Pathobiology, Virginia Tech, Blacksburg, Virginia, USA*
[5]*Translational Biology, Medicine, and Health Graduate Program, Virginia Tech, Roanoke, VA, USA*
[6]*Program in Genetics, Bioinformatics, and Computational Biology, Virginia Tech, Blacksburg, VA, USA*
[7]*Molecular Diagnostics Laboratory, Fralin Biomedical Research Institute, Virginia Tech, Roanoke, VA, USA*
[8]*Department of Biochemistry, Virginia Tech, Blacksburg VA, USA*
[9]*Data Services, University Libraries, Virginia Tech, Blacksburg, VA, USA*
[10]*Virginia Tech Center for Drug Discovery, Virginia Tech, Blacksburg, VA, USA*
[11]*Academy of Integrated Science, Virginia Tech, Blacksburg, VA, USA*
[12] *Virginia Tech Carilion School of Medicine, Virginia Tech, Roanoke, VA, USA*

*Correspondence to: Carla V. Finkielstein, finkielc@vt.edu; Joseph R. Hoyt, hoytjosephr@gmail.com

†These authors contributed equally: Carla V. Finkielstein, Joseph R. Hoyt

Keywords: COVID wildlife, cross-species transmission, Omicron variant, SARS-CoV-2, spillover

Abstract

Pervasive SARS-CoV-2 infections in humans have led to multiple transmission events to

captive animals. While SARS-CoV-2 has a potential broad wildlife host range, most

documented infections to date are found in a single species, the white-tailed deer. The

extent of SARS-CoV-2 exposure among wildlife species and the factors that influence

wildlife transmission risk remain unknown. We sampled 23 wildlife species for SARS-

CoV-2 and examined the effects of urbanization and human use on seropositivity. Here,

we document positive detections of SARS-CoV-2 RNA in six species, including the deer

mouse, Virginia opossum, raccoon, groundhog, Eastern cottontail, and Eastern red bat. In

addition, we found that sites with high human activity had three times higher

seroprevalence than low human-use areas. We detected SARS-CoV-2 genomic sequences

from nine individuals of six species which were assigned to seven Pango lineages of the

Omicron variant. The close match to variants circulating in humans at the time suggests

at least seven recent human-to-animal transmission events. Our data support that

exposure to SARS-CoV-2 has been widespread in wildlife communities and suggests that

areas with high human activity may serve as points of contact for cross-species

transmission.

INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the causative agent

of coronavirus disease 2019 (COVID-19), has resulted in over 771 million human cases

and over six million deaths worldwide[1]. As SARS-CoV-2 becomes endemic in humans,

one of the greatest threats to public health is the resurgence of more virulent and

transmissible variants. The considerable pathogen pressure imposed by the pandemic has

caused concern as to whether SARS-CoV-2 will spill into wildlife populations, establish

a sylvatic cycle, and potentially serve as a source for new variants.

Transmission of SARS-CoV-2 to captive animals has been well documented[2-4], but

detections in free-ranging wildlife are currently limited to only a few species including

white-tailed deer (*Odocoileus virginianus*[5-7]), feral mink (*Neovison vison*[8]), and Eurasian

river otters (*Lutra lutra*[9]). Experimental infections and modeling of the functional

receptor for SARS-CoV-2 (angiotensin-converting enzyme 2: ACE2) have shown that

numerous wildlife species may be competent hosts[10-15]. However, it remains unexplored

whether a diversity of wildlife species are infected in natural settings, where exposure to

SARS-CoV-2 is likely to be indirect and at a lower exposure dose.

Since the emergence of SARS-CoV-2 in 2019, numerous variants have been detected

in humans and animals. Many variants that have become dominant in human populations

have also shown mutations that increase their infectivity in humans[16]. These mutations

and new variants may also impact the virus's ability to infect new wildlife species. SARS-

CoV-2 collected from white-tailed deer have included lineages found in human samples,

caused by human-to-deer transmission[5], as well as unique mutations suggestive of deer-

to-deer transmission[17]. This implies that only minimal adaptation may be needed for

transmission to occur among deer following initial human-to-animal transmission

events[18]. Other human peridomestic species, such as deer mice (*Peromyscus*

maniculatus)[12,13] and skunks (*Mephitis mephitis*)[14] have been shown to be capable of viral

shedding in laboratory settings[11]. Collectively, these studies raise important questions

about the extent of human-to-wildlife transmission and the ability of other wildlife

species to sustain transmission.

Establishment of SARS-CoV-2 infections in wildlife communities could result in

novel mutations that increase virulence, transmissibility, or confer immune escape,

negatively impacting both human and wildlife populations. Furthermore, as SARS-CoV-

2 adapts to not only human hosts, but potentially a wide diversity of wildlife species,

SARS-CoV-2 evolution may become more unpredictable[19]. This could present several

challenges for human health, including concerns related to vaccine development targeting

human-specific lineages, and novel impacts to pathogenicity and transmissibility of the

virus.

Here, we examine how widespread SARS-CoV-2 exposure has been in wildlife

communities. We used quantitative reverse transcription polymerase chain reaction (RT-

qPCR) to examine 789 nasopharyngeal/oropharyngeal samples from 23 species and

documented the presence of SARS-CoV-2 RNA in six of these species. In addition, we

analyzed 126 serum samples from six species collected before and after the arrival of

SARS-CoV-2 and detected neutralizing antibody titers in five of the six species. Finally,

we detected an effect of urbanization and human use on seropositivity in animals, and

examined genomic data associated with positive samples.

RESULTS

SARS-CoV-2 RNA detections in multiple species.

We amplified SARS-CoV-2 RNA from nasopharyngeal/oropharyngeal samples by RT-qPCR from six of the 23 species examined (26.1% of species sampled) and had a total of 23 unique individual animals that were positive (2.9% of samples tested; Fig. 1a-b, Supplementary Tables 1-4). This included eight deer mice (*P. maniculatus*; 4.7%, N=172), four Virginia opossums (*Didelphis virginiana*; 2.9%, N=140), four raccoons (*Procyon lotor*; 4.8%, N=84), three Eastern cottontail rabbits (*Sylvilagus floridanus;* 2.5%, N=118), three groundhogs (*Marmota monax;* 9.7%, N=31), and one Eastern red bat (*Lasiurus borealis*; 8.3%, N=12) (Fig. 1b). We had slightly higher positivity rates in field collected samples compared to samples collected at wildlife rehabilitation centers (4.04% compared to 2.24%), which may reflect repeated sampling of wildlife at a site during a SARS-CoV-2 outbreak.

Figure 1. SARS-CoV-2 RNA and neutralizing antibody prevalence in wildlife communities. Panel (**a**) represents counties where swabs were collected (both wildlife

rehabilitation centers and study sites). Grey scale indicates sample sizes. Circles represent

SARS-CoV-2 positive samples and color indicates species in panel b. Prevalence of (**b**)

RT-qPCR detections collected between May 2022 and September 2023. Detections

include species with >10 individuals sampled, and black points indicate prevalence and

95% confidence intervals (See Supplementary Table 1 for full dataset that includes all 23

species). Panel (**c**) shows seroprevalence measured using plaque reduction neutralization

tests for species sampled both prior to SARS-CoV-2 arrival in 2020 and samples

collected in our study between June and July 2022 (Supplementary Table 5). The size of

each point in panel (**c**) is in relation to the percent neutralization. Samples with a 60% or

greater percent neutralization were considered positive samples. Error bars represent 95%

confidence intervals. Panel (**d**) represents percent neutralization values of serum samples

to SARS-CoV-2 sampled both prior and post SARS-CoV-2 arrival. The black dotted

horizontal line represents a 60% neutralization cut-off. The purple diamond represents

neutralizing antibody titers for a Virginia opossum who was found conclusively RT-

qPCR positive one month prior. Grey boxes represent median and interquartile range of

samples with less than a 60% neutralization value, and black boxes represent samples

with greater than a 60% neutralization value. Organism silhouettes were sourced from

PhyloPic[20].

SARS-CoV-2 neutralizing antibodies in mammal communities

We detected SARS-CoV-2 antibodies in five of the six species we collected serum

samples from in 2022 (60% neutralization cutoff, Fig. 1c-d and Supplementary Table 5),

including the Virginia opossum (37.5%, N=8), raccoon (36.4%, N=11), Eastern grey

squirrel (*Sciurus carolinensis*; 57.1%, N=7), white-footed mouse (*Peromyscus leucopus*;

16.7%, N=6) and the deer mouse (7.1%, N=14). Percent neutralization values from

samples prior to the arrival of SARS-CoV-2 (pre-2020) were significantly lower than

those collected after SARS-CoV-2 arrival (t=-10.774, $p < 0.001$, Fig. 1d). Furthermore,

four samples (two raccoons, one opossum, and one white-footed mouse) had a percent

neutralization above a more conservative 80% cut-off, further supporting that previous

SARS-CoV-2 exposure is likely in these species (Fig. 1d). In 2022, an opossum with a

positive detection was trapped one month later and collected serum revealed a percent

neutralization value of 51.1% which was below our 60% cutoff, suggesting this cutoff

may be too conservative for some species.

Figure 2. Examination of SARS-CoV-2 exposure at the human-wildlife interface.

Panel (**a**) shows imperviousness with darker shades representing more urban areas.

Circles represent sites sampled. We placed multiple traps at eight different sites (see

Supplementary Methods for trap locations). Sites PF and NRT were sampled in both

2022 and 2023, sites BB, BM, ML, and RK were only sampled in 2022, and sites PP and

CF were only sampled in 2023. Serology samples were collected at all of the study sites

in 2022 except PF. Panel (**b**) shows the relationship between urbanization

(imperviousness) and seroprevalence collected from five different sites in VA, USA.

Panel (**c**) shows the effects of average monthly human presence (population density

collected from the 2020 U.S. Census within the trapping area of each site (Blacksburg) or

use estimates from trail counters or landowners) on seroprevalence. Grey ribbons (shaded

areas around the line) represent 95% confidence intervals.

We found a positive relationship between urbanization (imperviousness) and

wildlife seroprevalence (intercept: -1.665 ± 0.48 SE, urbanization slope 0.039 ± 0.02 SE,

$p = 0.031$, Fig. 2a and 2b). However, antibody detections were highest (80%, N=5) at one

of the least urban sites, which is a highly visited state park (average imperviousness

1.5%), and more closely matched seroprevalence at our more urbanized sites (50%, N=10

and 33%, N=12). Human visitation at the state park was similar to human activity in

urbanized sites (Fig. 2c), and we found a positive relationship between human presence

and seroprevalence of SARS-CoV-2 (univariate linear mixed model with species as a

random effect; intercept: -1.132 ± 0.36 SE, human presence coeff: 0.705 ± 0.35 SE, $p =$

0.044; Fig. 2c, Supplementary Fig. 1). This relationship was maintained across different

neutralization cutoffs (40-65%; Supplementary Tables 6 and 7).

Whole genome sequencing from nasal and oropharyngeal swabs

SARS-CoV-2 sequences were obtained for 12 of the 23 RNA positive samples

(Fig. 3, Supplementary Figs. 2-9, Supplementary Tables 3 and 4) and Pango lineages

were determined for nine total individuals (39.1%; Supplementary Table 3). The SARS-

CoV-2 sequence from the opossum trapped in 2022 was assigned to BA.2.10.1 (opossum,

N=1) and shared defining mutations in *ORF1a/b*, *S*, *E*, and *M* genes found in the BA.2

Omicron lineage (Fig. 3b and Supplementary Fig. 2). All eight sequences collected in

2023 were assigned to the XBB* Pango lineages (Supplementary Figs. 3-9,

Supplementary Table 3). These lineages were circulating among humans in Virginia

during the time of collection (Fig. 3a). The lineages of the sequences obtained from

wildlife included XBB (deer mouse, N=1; Supplementary Fig. 3), XBB.1.5 (raccoon,

N=1; Supplementary Fig. 4), XBB.1.5.10 (opossum N=1; Supplementary Fig. 5),

XBB.1.16 (Eastern cottontail, N=1; Supplementary Fig. 6), XBB.1.5.45 (groundhog,

N=1; Supplementary Fig. 7), EG.5.1.1 (deer mouse, N=2; Supplementary Fig. 8), and

JD.1 (deer mouse, N=1; Supplementary Fig. 9). The remaining three sequences included

two from Eastern red bats and one from a deer mouse, which were generated using

amplicon sequencing of the S gene. These matched the SARS-CoV-2 reference sequence

(Supplementary Table 4) with 95-100% identity but were at insufficient length for

phylogenetic analysis and lineage assignment.

The relative similarity of the sequences obtained from wild animals compared to

the closest sequences obtained from humans (Supplementary Table 3) suggest recent

introductions of these SARS-CoV-2 lineages into wildlife with at least seven independent

introductions of SARS-CoV-2 over a several month period. Interestingly, two sequences

from the same lineage (EG.5.1.1) were collected from deer mice on the same day and

location (site=PP; Fig. 3 & Supplementary Fig. 8). These two sequences clustered

together along with a sequence collected from an infected human in Vienna, Virginia

(~330km distance) three days prior (Fig. 3 & Supplementary Fig. 8). This clustering

suggests the deer mice might have been exposed to the same source of the virus or could

represent a mouse-to-mouse transmission event (Fig. 3c). Sequence from a third mouse

collected from the same day and site (site=PP) aligned with another lineage (JD.1, Fig. 3b

and Supplementary Table 3), and suggests a separate human-to-mouse introduction (Fig.

3c). At a nearby site (site = NRT; located ~52 km away) sequences produced for two

additional individuals (one deer mouse and one raccoon, Fig. 3b and Supplementary

Table 3) were also from separate lineages (XBB and XBB.1.5, respectively), further

suggesting independent human-to-animal introductions (Fig. 3c).

All sequences from wildlife were aligned to their closest related human sequence

based on phylogeny, and then assessed for any unique amino acid substitutions. Whole

genome sequencing (WGS) of the SARS-CoV-2 variant collected from the positive

opossum in July 2022 revealed several mutations shared within the Omicron clade as

well as a unique amino acid substitution not previously identified in other SARS-CoV-2

virus samples collected from humans at the time of sampling (Table 1).

The Omicron sub-lineage, BA.2.10.1(Fig. 3), includes the $G^{798}D$ mutation found in the Spike (S) protein. The nearest neighbor (EPI_ISL_14334179), which was collected in New York state 11 days prior to our sampling, contains all the nucleotide and amino acid mismatches found in the opossum sequence, compared to the reference

Table 1. Summary of mutations identified in the Virginia opossum (*Didelphis virginianus*)

SNP	Gene	Mutation
C7749T	*ORF1ab/NSP3*	Missense $T^{2495}I$
T16342C	*ORFab/Helicase*	Missense $S^{5390}P$
A20304G	*ORFab/EndoRNAse*	Synonymous
A22974T	*S*	[1]Missense $E^{471}V$
G23955A	*S*	[2]Missense $G^{798}D$

[1,2]Fig. 4a, Supplementary Figs. 10 and 11

190 sequence from Wuhan, China, except for A22974T (S:$E^{471}V$) which appears to be unique

191 to the opossum. Only one mutation was found among the eight other SARS- CoV-2

192 sequences from 2023 as compared to closely related human sequences. However, this

193   mutation (S:H146Q), in a positive detection from a groundhog, is a common mutation

194   found in other Pango XBB* lineages circulating in 2023.

Figure 3. SARS-CoV-2 collected from wildlife in comparison to human samples. (a)

Color ribbons show the proportion of each lineage circulating in humans in sampling

region at the time of data collection[21]. Circles represent each individual animal sampled.

A positive sample has a value of 1 and a negative sample has a value of 0. Circles

outlined in red represents a positive sample with lineage assignment (Supplementary

Figs. 2-9; Table 3). The fill color within each red circle represents the lineage it was

assigned. (**b**) A phylogenetic tree was generated using maximum likelihood analysis and

GTR+G4 substitution models to visualize the relatedness of the nine whole genome

sequences obtained from wild animals to 90 SARS-CoV-2 sequences derived from

human hosts representing different PANGO lineages. Additional phylogenetic analyses

show the placement of the SARS-CoV-2 sequences from each individual wild animal

within individual lineages in more detail (Supplementary Figs. 2-9). (**c**) Suspected

transmission origin for each individual is represented by a human silhouette or a question

mark ("?") if the origin is suspected to be either a human-to-animal or animal-to-animal

transmission. Inferences about transmission were determined by phylogenetic analysis

using related human host sequences and our sequences collected from wild animals as

shown in Supplementary Fig. 2-9 and in Supplementary Table 3. We also considered any

unique mutations not described from a human host, as those mutations may have come

from the originating hosts' virus. Arrow color indicates the lineage assignment and

silhouettes indicate the species from which the sample was collected.

Unique SARS-CoV-2 mutations collected from wildlife

We investigated the unique $E^{471}V$ mutation detected in the opossum sequence

from 2022, which was located in the receptor-binding motif (RBM, residues 437 to 508)

within the receptor-binding domain (RBD, residues 319 to 541) of the S protein. The

RBD is necessary for binding the viral S protein to ACE2 in both human and animal cells

and is therapeutically targeted by neutralizing monoclonal antibodies[22]. We conducted *in*

silico studies to examine the local impact of the $E^{471}V$ and $G^{798}D$ mutations on the S

protein. We employed molecular modeling and molecular mechanics/generalized borne

surface area (MM/GBSA) free energy calculations to predict the free energy of binding

interaction (ΔG_{bind}). Our focus was on examining the local impact of $E^{471}V$ and $G^{798}D$ on

the favorability of S protein-hACE2 interaction and the modifications in the S2 site as a

hypothesis for the experimentally observed traits found in SARS-CoV-2 wildlife

reservoir. Here, the open conformation structure of the S protein trimer from the

glycosylated BA.2 Omicron variant was used for residue variant mapping [PDB 7XO8][23].

Because the crystal structure shows the trimer bound to the human ACE2 (hACE2)

receptor, it is possible to investigate the local impact of mutations in the RBD on

interactions between SARS-CoV-2 S protein and the human ACE2, as a prediction for

mechanistic impact on S protein-ACE2 interaction favorability as previously reported and

validated experimentally[24] (Fig. 4). Our free energy studies predict that ΔG_{bind} between

the $S:E^{471}V$-carrying mutation and hACE2 was improved relative to that of BA.2 with

hACE2 (-76.5 and -60.2 kcal/mol for $E^{471}V$ vs. BA.2, respectively). Large-scale low-

mode conformational sampling was performed on the RBD of BA.2 and $E^{471}V$ Spike

(residues 329-531) with hACE2 to probe ΔG_{bind} on multiple potential conformations[25,26].

Consistent with the results observed in full-length Spike-hACE2 interactions, the mean

ΔG_{bind} for the top five most favorable $E^{471}V_{RBD}$ conformations (-151.7 ± 19.2 kcal/mol)

was more favorable than the mean for $BA.2_{RBD}$ conformations (125.4 ± 4.1 kcal/mol,

$p \leq 0.05$). The sampled conformations of $E^{471}V_{RBD}$ exhibited a larger range of ΔG_{bind}

values relative to BA.2$_{RBD}$, reflecting not only increased interaction flexibility but also

increased flexibility in the RBD due to the E^{471}V mutation. Residue 471 is predicted to be

part of a flexible, hydrophobic loop that interacts with the N-terminal domain of ACE2

(Fig. 4). Several point mutations (A^{475}V, S^{477}G, V^{483}A, F^{490}L) within this loop region

have been experimentally shown to improve S protein – hACE2 binding affinity and

increased resistance to neutralizing antibodies[27,28]. Our analysis using normalized Kyte-

Doolittle hydropathy scores and surface maps show a more hydrophobic interaction

surface at not just position 471, but also among flanking residues 469-474 (0.8) compared

to BA.2 [-2.6][29]. The increased favorability of the E^{471}V mutation for hACE2 interaction

aligns with previously reported trends of residues in the RBD influencing stabilization

and enhanced affinity[30]. Any differences may be attributed to entropic effects,

emphasizing the potential enthalpic contribution of this mutation to improve hACE2

interaction at the RBD interface.

Figure 4. Molecular modeling of the unique S mutation of SARS-CoV-2 collected

from a Virginia opossum. (**a**) Representation of the structure of the BA.2 S protein

(gray, PDB: 7XO8) in its open conformation bound to the human ACE2 (teal). Residues

471 (red) and 798 (orange) are shown as spheres. Glycans are displayed as sticks colored

purple. Top inset: Overlay of $E^{471}V$ (red) and BA.2 (grey). Bottom inset: Overlay of

$G^{798}D$ (orange) and BA.2 (grey). (**b**) Surface map of the BA.2 S protein (left) and the

region surrounding $E^{47}1V$ and $D^{798}G$ (right) residues. Residue side chain properties are

colored: green for hydrophobic, blue for positively charged, red for negatively charged,

teal for polar uncharged, and gray for neutral.

We mapped the second missense mutation, $G^{798}D$, in the S2 subunit of the S protein

(residues 686 to 1,273) and within the fusion peptide domain (residues 788 to 806) near

the N^{801} glycosylation site. When glycosylated, the N^{801} site significantly enhances viral

entry, with very low mutation rates observed in adjacent residues (e.g., within 2-3

residues from position 801)[31]. Interestingly, D^{798} creates a small, charged pocket on this

solvent-accessible S2 loop, decreasing the probability of glycosylation at N^{801} (0.48)

compared to BA.2 [0.61][32]. It is hypothesized that $G^{798}D$ could impact structural stability

and membrane interaction[33] (Fig. 4 and Supplementary Fig. 10) and have a negative

impact on Spike maturation and infectivity. While this approach limits the investigation

of these mutations and their role in the dynamic conformational states of the S protein, it

provides a foundation for understanding the mechanistic impact and connection of

structure to experimental observables.

DISCUSSION

Our combined results suggest that a broad diversity of mammal species have been

exposed to SARS-CoV-2 in the wild (Supplementary Table 8). While species like the

white-tailed deer have been shown to be important hosts for SARS-CoV-2, our results

highlight that evaluating the importance of each species in the context of a broader

community of hosts will be critical for controlling future zoonotic disease risk[34].

For some species, SARS-CoV-2 RNA detections and seroprevalence aligned well

with both predicted susceptibilities based on ACE2 modeling and experimental infection

studies (e.g., whether the species seroconverted, was capable of viral shedding, or

transmitting; Supplementary Table 9). Most notable is the deer mouse, which has been

predicted to have high susceptibility[35], shown experimentally to be capable of mouse-to-

mouse transmission[12,13], and produces antibodies in the lab[12,13,36]. We had the highest

number of positive RNA detections (N=8) in this species, as well as detected neutralizing

antibodies (Fig. 1, Supplementary Table 1 and 8). Raccoons have also been shown

experimentally to be capable of seroconverting following SARS-CoV-2 exposure[14,36],

and we found high seroprevalence (63.6%) in this species. However, experimental

infections did not find evidence of viral shedding or transmission in raccoons when

infected with the USA-WA1/2020[14] or the WA1/2020WY96 SARS-CoV-2 isolates[36],

whereas we had four positive RT-qPCR detections in this species (Supplementary Table

1). We also had positive RNA detections in cottontail rabbits (N=3), which, when

experimentally infected in the lab, also did not produce any clinical signs of infection, or

appear to be capable of viral shedding when infected with the WA1/2020WY96 isolate[36].

Several factors may contribute to the apparent discordance between experimental

infections and positive detections in the wild. First, experimental infection studies with

these species used earlier SARS-CoV-2 variants and most positive detections in this

study occurred recently (July-September 2023, Fig. 3a) with newly emerged lineages

(e.g., XBB). The rapid evolutionary change in SARS-CoV-2 infectivity, virulence, and

immune escape abilities may have changed species competence for SARS-CoV-2.

Supporting this, recent research has shown that Omicron's (B.1.1.529) ACE2 binding

affinity is greater in a number of wildlife species compared to humans, including white-

footed mice, red fox, two marmot species, and five marsupial species[37]. Another

possibility is that some RNA detections do not reflect active infections, but rather recent

exposure to another infected animal. Other species with positive detections (groundhog,

Virginia opossum, and Eastern red bat) have not currently been evaluated in laboratory

experiments (Supplementary Table 9). Experimental infection studies and updated

molecular modeling with new viral lineages will be essential for assessing the current risk

to other mammal species.

Many of the species in this study are considered human peri-domestics, have

broad ranges across North America, and live in and around human settlements. In

addition, some species have been introduced into Europe (gray squirrels and raccoons)

and Asia (raccoons) and have close relatives across the globe. Deer mice are of particular

importance given that they are known to be reservoirs for other pulmonary viruses[38], and

are in close direct and indirect contact with humans. It is unknown whether a sylvatic

cycle has been established in any of the species we evaluated; however, their close

connection to humans and broad spatial distribution means they are likely to experience

continuous exposure to a diversity of SARS-CoV-2 lineages in the future.

We found support for a relationship between human presence and seroprevalence,

suggesting that areas with high human activity may serve as potential hotspots for cross-

species transmission. This suggests that wildlife in areas with more human activity likely

have higher risk of exposure to SARS-CoV-2. Hence, recreational and highly urbanized

sites may be important points of contact between humans and wildlife and could be

targeted for surveillance and control. Humans and wildlife rarely come into direct

contact, but numerous indirect links likely exist. Wastewater has been proposed as a

potential source for indirect exposure to SARS-CoV-2[39], however, in rural areas where

septic tanks are a dominant form of wastewater management, this is unlikely to be the

only source. Instead, other forms of human waste, like trash receptacles, may be

important sources of indirect SARS-CoV-2 exposure in wildlife[14,40]. Urban wildlife are

regularly exposed to human refuse and have developed positive associations with

discarded food, which could serve as a bridge for transmission between humans,

companion animals, and wild species[41-43].

We obtained eight RNA sequences in 2023 from multiple Pango lineages, likely

indicating multiple introductions of the virus from humans into wildlife. This highlights

that wildlife are continually exposed to a wide diversity of SARS-CoV-2 lineages, as has

been observed in white-tailed deer[5,18]. We can further assume that these are recent

introductions, and not due to an established animal host sylvatic cycle, as they are close

matches (99.1-100.0%) to published human sequences circulating at the same time of

collection. It is important to note that in two instances, multiple animals tested positive at

the same site during the same collection period (~4 days), and animal-to-animal

transmission following human introduction cannot be ruled out. It is difficult to fully

resolve whether transmission occurred via human-animal or animal-to-animal in the

summer of 2023 because of a lack of human collected samples from that time, as the

reporting of human samples in 2023 had significantly decreased compared to previous

years. Hence, there is a much smaller sample size of known circulating human variants at

a fine resolution. Determining how wildlife are being infected (pathways such as human

refuse, wastewater, contact with infected pets, etc.) is a critical next step in disease

control and management. It will be important to continue to sequence variants from

wildlife as well as humans to assess if SARS-CoV-2 is adapting to new wildlife hosts, if

a sylvatic cycle develops, and whether there is a risk of transmission back to humans.

Analysis of sequences collected from an opossum in 2022 revealed a novel

mutation that has not been described in any other variant, which might suggest animal-to-

animal transmission or possible adaptation within the opossum. All others mutations were

found circulating in humans in other regions. Some of these mutations likely increase

binding affinity to the hACE2 receptor or confer some antibody resistance compared to

ancestral lineages. Specifically, SARS-CoV-2 collected from the opossum had mutations

in the RBM of the RBD of the S protein. The previously uncharacterized amino acid

change is predicted to improve S-hACE2 binding compared to Omicron BA.2, potentially

providing a fitness advantage by either increasing the affinity of S for the ACE2 receptor

or, alternatively, by evading the neutralizing activity of antibodies. Whether these

mutations developed in humans not captured in surveillance at the time or in wildlife

communities and were transmitted back to humans, which has been previously suggested

during the emergence of the Omicron variant[5,7], remains unknown (Fig. 3). Future work

using molecular dynamics (MD) simulations to evaluate the impact of these mutations on

structural stability, membrane interaction, and conformational states will help contribute

to our understanding of evolutionary pressures and cross-species transmission.

While single gene amplification (*S, N, or E*) below our cutoff did not meet our

criteria for assigning a positive (*i.e.,* at least two SARS-CoV-2 genes need to amplify

below the cutoff for a reference sample to be reported as positive based on our EUA-

FDA and CLIA protocols), the analytical specificity (cross-reactivity) of the RT-qPCR

(99-100% single gene specificity) suggests that many of these may be positive detections

with low levels of RNA[44]. Single-gene positive detections overlapped with all six species

confirmed in this study (Supplementary Table 1), but also encompassed an additional

seven species (Supplementary Tables 1 and 2), including the American beaver (*Castor*

canadensis; 1/2), bobcat (*Lynx rufus;* 2/3), American black bear (*Ursus americanus*; 1/7),

red fox (*Vulpes vulpes*; 2/17), white-tailed deer (3/20), skunk (1/25), and Eastern gray

squirrel (4/105). We also obtained sequence data that was a 100% match for SARS-CoV-

2 from single gene detections, further suggesting that in some cases, these may indicate

true positives (Supplementary Tables 2-4). While confirming new species requires a more

conservative and confirmatory approach, given the large number of single-gene positives,

future studies should carefully consider whether these are unconfirmed detections,

particularly in small wildlife species, which may produce lower viral loads than humans

or larger mammals like deer. Importantly, classifying unconfirmed samples as negative

could substantially alter our understanding of the ecology and community dynamics of

SARS-CoV-2 in wildlife going forward. Furthermore, our use of neutralization tests to

detect evidence of prior SARS-CoV-2 exposure may have resulted in reduced sensitivity

over approaches like ELISA that identify only binding antibodies. However, binding-

antibody-based approaches also suffer from false-positivity to other coronaviruses[45],

other infectious diseases[46], and even auto-immune diseases[47]. Thus, PRNT increases the

confidence that animals tested here were truly exposed to SARS-CoV-2 and not to

another related viruses.

Our results have greatly expanded the known range of hosts exposed to SARS-

CoV-2 in the wild, which now includes nine species (six from this study) that have been

documented with SARS-CoV-2. Several of the species that tested positive (mice, rabbits,

and opossums) have traits that may make them more suitable for establishing a SARS-

CoV-2 wildlife reservoir than previously described species. These include fast-paced life

history strategies whereby individuals reproduce frequently at a young age, and in some

cases are known reservoirs for other respiratory viruses[48,49]. Continued broad surveillance

and more detailed ecological research will be needed to fully determine the role of

wildlife communities in SARS-CoV-2 transmission and evolution.

MATERIALS AND METHODS

Study Sites

We collected nasopharyngeal or oropharyngeal samples from wildlife across 43

counties in Virginia and Washington D.C., U.S.A. We collected samples from three

wildlife rehabilitation centers in Boyce, Roanoke, and Waynesboro, VA. In addition, we

actively captured wildlife from eight sites between 2022 and 2023 that spanned a rural to

urban gradient in Giles, Montgomery, Roanoke, and Wythe counties (Fig. 2a,

Supplementary Table 10, and see Supplemental Methods for site descriptions.). At six

sites we actively trapped wildlife using live-traps from 9-May through 1-July 2022 and at

four sites (two sites were the same as 2022) we trapped from 27-June through 11-

September 2023.

Trapping and processing

We trapped each of six sites for a 2-4 day session between 9-May and 1-July 2022

(two sites were trapped for two sessions with at least 27 days between them).

Additionally, we trapped four sites for a 4-day session between 27-June and 11-

September 2023 (Fig. 2a). We used three different sized live traps to capture animals:

Tomahawk non-folding traps (Tomahawk Live Trap Co., Tomahawk, WI, U.S.A; 46 x 15

x 15cm, model 103.5), Tomahawk folding traps (66 x 23 x 23cm, model 205 or 81 x 25 x

31, model 207), and Sherman folding traps for smaller animals (Sherman Traps,

Tallahassee, FL, U.S.A; 8 x 9 x 23cm, model LFG). We set and baited traps in the

evening just before dark and checked them the following morning before 10:00 AM to

ensure animals did not get overheated.

We processed all animals in the area where they were trapped. We anesthetized

larger animals in either a bucket chamber[50] or in a 50L clear box chamber (58 x 38 x 56

cm, Supplementary Fig. 12). We used a tec-4 funnel fill with a cage mount manifold

vaporizer (Ohmeda, West Yorkshire, UK) connected to a portable medical oxygen

cylinder (size E) with a built-in regulator (Walk-02-Bout+[TM], Airgas Healthcare, Radnor,

PA). We used a 2 L/min oxygen mix to 3-4% isoflurane in the chambers. Smaller animals

were placed in a small plastic canister and anesthetized with 0.5 – 1.2mL isoflurane

(dosage depended on species and body size), placed onto a cotton ball held within a

perforated canister to prevent direct contact with the animal (Supplementary Fig. 12).

Once anesthetized, we removed animals from the chamber, and then masked animals

(except for mice; Supplementary Fig. 12) with a dose of 2 L/min oxygen and 2-3%

isoflurane.

We marked individuals with aluminum ear tags in each ear (National Band and Tag

Company, model 1005-1 or 1005-3 depending on body size). We collected

morphometric, sex and reproductive status from each animal. For RT-qPCR detections,

we swabbed each individual with a polyester swab (25-800 1PD swab for larger

mammals and 25-1000 1PD swab for smaller animals e.g., juveniles, skunks, squirrels,

bats, shrews, and mice; Puritan Medical Products, Guilford, ME, U.S.A). Given the small

swab size compared to humans and larger mammals and the reduction in material that

could be collected from these swabs, we collected two swabs per animal after July 2023.

We used an oropharyngeal swab for animals that were too small for nasopharyngeal

swabbing (mice, shrews, bats, etc.). All collection swabs were placed in tubes with

transport media[44] and kept on ice until they were transported back to the laboratory

following field work each day. Transport media consisted of a 1:1 volume of Dulbecco's

Modified Eagle's Medium (DMEM) containing low glucose (1g/l), sodium pyruvate and

L-glutamine (Corning) media and 2x DNA/RNA Shield[TM] (Zymo Research), which can

preserve samples up to 10 days at room temperature[44]. Field collected samples were all

stored in the refrigerator at 4°C and processed within five days following field collection.

We collected blood for antibody screening from the submandibular vein for mice and

from the nail quick of one of the rear toenails for all other species. Blood was stored in

microvette 500 z-gel tubes for later processing. We put all blood samples into a cooler

immediately following field collection and stored at 4°C. Three wildlife rehabilitation

centers in Virginia provided wildlife oral and nasal swabs following methods described

above. We analyzed all samples from rehabilitation centers within 45 days of collection.

Protocols were approved under Virginia Tech IACUC protocol #22-061.

Personal Protective Equipment and Contamination Control

All personnel collecting and working with the samples wore fit-tested N95 respirators

and gloves. We sterilized all equipment between each animal. We also sterilized all traps

and equipment between each trapping session, and individuals collecting the samples

were tested using the assay described below 1-2 times per week to confirm they were

negative throughout the duration of sampling. During fall of 2023, when most positive

samples were collected, all personnel were tested daily.

RNA Extraction and detection of SARS-CoV-2 viral RNA

The RT-qPCR based test we used for the identification of SARS-CoV-2 RNA in

wildlife has been thoroughly described in Ref[40]. This SARS-CoV-2 assay, authorized for

use in humans by the FDA during the pandemic and adhering to all regulatory standards,

detects three targets including the nucleocapsid (*N*), envelope (*E*), and spike (*S*) genes of

the SARS-CoV-2 virus, has high analytical specificity, and across all three targets has a

100% probability of detection at 10.0 copies per 10 µL.

We processed swab-containing media shortly after collection and total RNA was

purified using 96-well spin columns, assessed for quality control, and subjected to

synthesis and amplification using the Power SYBR^{TM} Green RNA-to-CT^{TM} 1-Step kit

(Applied Biosystems). The cytochrome c (Cytochrome C Oxidase Subunit 1; *Cox1*)

housekeeping gene, for which species-specific primers were designed (Supplementary

Table 11), and HRPP30 housekeeping gene for human samples was used as a control for

whether sufficient sample was collected from each individual. RT-qPCR reactions were

performed in a CFX384 Touch Real-Time PCR detection system (Bio-Rad), and we

included positive controls using inactive SARS-CoV-2 virus (ATCC®), no template

controls, and standard curves on each plate as previously described[44]. Samples were

reported conclusively positive when two or more SARS-CoV-2 targets (*N*, *E*, and *S*

genes) and the housekeeping gene (Supplementary Table 2) amplified below the

threshold established by corresponding standard curves[44], which is similar to the criteria

established for the Applied Biosystems TaqPath COVID-19 used for confirmation in

other wildlife studies[7]. The use of multiple targets is particularly relevant when

considering genetic variations of SARS-CoV-2 among circulating variants for which the

sole amplification of a single gene could result in a false negative result if the mutation

was in a region of the genome assessed by the test. Thus, a molecular diagnostic test

developed to detect multiple genetic targets of SARS-CoV-2 is likely less susceptible to

novel genetic variation. For the positive samples from the opossum collected in 2022, the

remaining sample was sent to the USDA National Veterinary Services

Laboratories (NVSL) in Ames, Iowa for confirmation of results. However, while

insufficient sample remained for NVSL to conduct confirmatory testing, raw sequence

data from this positive was independently analyzed and confirmed to be from an

opossum.

The Virginia Division of Consolidated Lab Services (DCLS) confirmed the

performance of the developed assay using a comparator assay that followed the CDC

EUA IFU (CDC DOC 006-0099 rev.03) "CDC 2019-Novel Coronavirus (2019-nCoV)

Real-Time RT-PCR Diagnostic Panel" that includes the same probes and N primers.

Results of all blindly tested samples were incorporated into the Emergency Use

Authorization which was approved by the Food and Drug Administration. To date, this

assay has been used for sample analysis by the Virginia Department of Health in over

230,000 tests in Southwest Virginia. Lastly, the Molecular Diagnostics Laboratory that

tested the samples is CLIA certified and participates in a proficiency testing program

from the American Proficiency Institute.

Serology data

We obtained additional samples (Supplementary Table 5) to compare seropositivity

prior to the emergence of SARS-CoV-2 to those samples we collected in 2022[51]. The

samples were transferred to gold-top microtainer tubes (BD, Franklin Lakes, NJ, USA)

and serum was separated via centrifugation at 5,000 x g for 10 minutes and transferred to

1.7 mL tubes. Serum samples were then heat inactivated at 55°C for 1 hour. Plaque

reduction neutralization assays (PRNTs) were performed similarly to prior reports[52,53].

Briefly, samples were diluted 1:10 in RPMI-1640 with 10 mM HEPES and 2% FetalPure

bovine serum (Genesee Scientific 25-525H). Diluted serum samples were mixed in a 1:1

v/v ratio with a solution containing 1300 plaque-forming units per mL (PFU/mL) of

SARS-CoV-2 Delta virus strain USA-GNL-1205/2021 (a generous gift from the World

Reference Center for Emerging Viruses and Arboviruses), or approximately 40 PFU's per

well. Our serum samples were collected in June and July of 2022, following the Delta

wave in fall of 2021, and given the life expectancy of the animal species (Supplementary

Table 12) we were sampling, we assumed Delta would have been the most likely recent

variant they were exposed to[54]. Each sample was tested in triplicate. The mixture was

then incubated at 37 °C for one hour, and then 50 µL of the virus-serum mixture was

used to inoculate wells in a confluent 24-well plate of VeroE6 hACE2-TMPRSS2 cells

and again incubated at 37 °C for one hour. After the one-hour adsorption period, we

added 500 µL of overlay media to each well and incubated the plates at 37°C, as

previously described[55]. After two days, the plates were fixed with 10% formalin and

stained with crystal violet. The level of neutralization is based on the reduction of plaque

counts compared to virus mixed with diluent. Based on previously published sero-

surveillance studies[56-59], we used the following testing criteria:

1. ≥ 90 % Neutralization at a 1:20 Dilution – Strong Positive

2. ≥ 60 % Neutralization at a 1:20 Dilution – Weak Positive

3. < 60 % Neutralization at 1:20 Dilution – Negative

Any individual with a 60% neutralization or higher was counted as positive in analyses

analyzing the effects of urbanization and human presence on SARS-CoV-2 exposure.

Urbanization and human presence variables

We obtained estimates of urban imperviousness from the National Land Cover

Database 2019[60,61]. We obtained estimates of population density from 2020 census at the

1km spatial scale[62]. We created buffers of variable sizes based on species home range

size to calculate the mean estimates of imperviousness and population density for each

individual capture location (Supplementary Table 13).

We obtained estimates of human presence at the five sites where we collected

serology data (Supplementary Table 10) and all visitation data was standardized to

number of visits per month. New River Trails State Park and Roanoke Parks and

Recreation provided monthly human visitation estimates between March and August

2022. Mountain Lake Biological Research Station provided estimates for how many

people used the facilities at the station between March and August 2022; they averaged

about 50 people at the station per month with each person staying 7 days on average.

Brush Mountain recreational area was closed to the public from March through August

2022, however, there was some trail maintenance work and the occasional trespasser so

we assumed approximately 10 people used the area per month.). For the town of

Blacksburg, we used population estimates from the 2020 census (TIGER/Line, U.S.

Census Bureau, 2020 Blocks). We created a minimum convex polygon (mcp) in the

adehabitatHR package[63] in program R v 4.1.1[64] around the different trapping locations.

We then added a 1km buffer area around each census assuming both animals and

residents move in and out of the trapping mcp area. We used ArcGIS Pro (ArcGIS Pro

2.9.0, Esri, Redlands, CA, USA) to calculate the population within each mcp area. We

used the intersect tool field calculator, and summary statistics to calculate the population

for each census block by multiplying the percent of the block within the mcp by the

population count. We then summed the population within the mcp to estimate the total

population. We then multiplied the population count by 30 (approximately days in each

month) to compare the daily estimates of occupancy to the monthly estimates from the

other sites.

Clinical specimen analysis

Human (13,221) and wildlife (789) clinical samples were analyzed for the

presence of SARS-CoV-2 by RT-qPCR. We performed whole genome and amplicon

sequencing for positive samples. We limited the description of variants in the human

population to the window at which wild sample collection took place (May 1, 2022 to

September 8, 2023 Fig. 3a and Supplementary Fig. 13).

Human clinical samples were collected as part of ongoing SARS-CoV-2

surveillance efforts in the region and collected under approval through Virginia Tech

Institutional Review Board (IRB 20-852) and the Virginia Department of Health (VDH

IRB 70046). The analysis of human clinical specimens identified 4,123 as positive, 8,350

as negative, 162 as inconclusive and 586 as invalid (no housekeeping gene detected),

respectively. (Supplementary Fig. 13a). A few samples failed sequencing or were not

assigned due to their quality/quantity and low sequence depth. As shown in

Supplementary Fig. 13b, to further examine the sequence data from the positive opossum

from 2022, various SARS-CoV-2 lineages expanded the window of our study with BA.1

(0.17%), BA.2 (15.85%), BA.2.12.1 (25.80%), BA.4 (14.83%), BA.5 (37.91%), BE

(1.57%), BF (1.45%), BG.2 (0.05%), AY.103 (0.02%), not assigned (N/A, 2.24%) being

the most prevalent among humans in the region.

Whole genome sequencing and Sanger amplification

Sanger sequencing was carried out using ARTIC primers (v3/v4.1) *. Samples

were treated with ExoSAP-IT™ Express (Applied Biosystems) and submitted to Eurofins

Genomics (Louisville, KY) for Sanger sequencing. To amplify the entire SARS-CoV-2

genome of the 2022 Virginia opossum, positive samples were reverse transcribed and

amplified using SuperScript IV Reverse Transcriptase (Thermofisher) and ARTIC nCoV-

2019 Amplicon Panel V4.1 primers (Integrated DNA technologies). Purified PCR

products were barcoded using the plexWell™ 384 Library Preparation Kit (seqWell,

MA) and the pooled library sequenced using a MiSeq System (Illumina, CA) following

manufacturer's instructions. SARS-CoV-2 sequencing analysis was accomplished using

an established pipeline and optimized workflow for validation that combines tools to

assess quality, sequence alignment, variation calling, and variant assignment. We

processed lineage assignments using Pangolin software suite (https://cov-lineages.org/)[65].

All remaining positive SARS-CoV-2 detections in 2023 by qRT-PCR were whole

genome sequenced using a GridION X5 Sequencer (Oxford Nanopore Technologies,

ONT) separately from the 2022 Virginia opossum. Amplifications were performed using

the Lunascript RT Supermix (New England BioLabs) and Midnight RT PCR Expansion

(Oxford Nanopore Technologies). Barcoding was performed using the Rapid Barcoding

Kit 96 (Oxford Nanopore Technologies) before the cDNA pool was purified. The pure

cDNA pool was quantified using either a Nanodrop 2000/2000c or Qubit Flex (Thermo

Fisher Scientific) before loading into a flow cell (R.9.4.0 or R.9.4.1 Flow Cell Versions;

Oxford Nanopore Technologies). Sequencing was performed using a GridION X5

Sequencer (Oxford Nanopore Technologies). Raw sequence data was basecalled and

demultiplexed using ONT's Guppy software, and analysis performed using ONT's

EPI2ME Labs software package running the wf-artic analysis protocol using scheme

Midnight-ONT/V3 (Oxford Nanopore Technologies and ARTIC Network). The wf-artic

analysis protocol utilizes Medaka, Nextflow, Nextclade, and Pangolin to produce a

comprehensive analysis report that assesses sequence quality and alignment, as well as

variant calling and assignment. Whole genome sequences of all human samples were

deposited in the GISAID database (https://www.gisaid.org/) and NCBI

(https://ncbi.nlm.nih.gov) SARS-CoV-2 sequence repositories.

Phylogenetic analysis

Consensus FASTA files containing available sequences from shotgun whole

genome sequencing and targeted amplicon sequencing were assembled using DECIPHER

software package[66]. These sequences were uploaded into Ultrafast Sample placement on

Exiting tRee [UShER; https://genome.ucsc.edu/cgi-bin/hgPhyloPlace][67,68] for global

phylogenetic tree analysis using 12,397,869 SARS-CoV-2 genomes obtained from

GISAID, GenBank, COG-UK and CNCB that were available as of 2022-10-03 for the

original opossum sequence and 16,284,433 SARS-CoV-2 genomes that were available as

of 2023-10-12 for the other eight SARS-CoV-2 sequences obtained from wild animals.

Sequences were also uploaded into NextClade [https://clades.nextstrain.org/][70] for clade

assignment and mutation calling.

For visualization of these nine consensus whole genome sequences in comparison

to Pango lineages, a phylogenetic tree was generated using a randomized subset of 90

SARS-CoV-2 sequences from NCBI Virus[71] that represent 15 different Pango lineages (6

sequences per lineage). The sequence alignment and a Maximum Likelihood (ML) tree

was generated using DECIPHER's TreeLine function[66] and viewed using iTOL v6[72].

Each of these SARS-CoV-2 sequences collected from wildlife were further

analyzed for similarity to published SARS-CoV-2 sequences obtained from humans

within our geographical region. We assembled separate phylogenetic trees for each Pango

lineage using human SARS-CoV-2 sequences collected from Virginia and the four states

that neighbor Southwest Virginia, including West Virginia, Kentucky, Tennessee and

North Carolina (Supplementary Tables 14 and 15). To limit the size of some of the trees,

we only included sequences collected around the same time as the dates for the wildlife

sample collection. For lineages XBB.1.4.45 and JD.1, an insufficient number of

sequences were available from these five states so all available sequences from North

America were included for generating the phylogenetic trees. These human SARS-CoV-2

sequences were downloaded from GISAID[73] and NCBI Virus. Duplicate sequences were

removed prior to tree assembly. The trees were rooted using the NCBI Reference

Sequence (NC_045512.2) for SARS-CoV-2. Sequence alignments and ML trees were

generated as described above.

We used Nextclade's mutation calling feature to identify novel amino acid

substitutions that were unique to our wildlife samples as they were not also found in their

closest neighboring sequence.

Molecular modeling

We carried out computational studies of the structure of the glycosylated Omicron

BA.2 variant bound to the human ACE2 receptor [Protein Data Bank 7XO8][23]. We

incorporated point mutations in BA.2 using PyMOL (The PyMOL Molecular Graphics

System v. 2.4.0, Schrödinger, LLC) to generate a new structure containing both E471V

and G798D mutations. We then performed energy minimization using Schrödinger-

Maestro (v. 2020.4) and OPLS3e force field[74,75]. We used Schrödinger Bioluminate to

carry out molecular mechanics/generalized borne surface area (MM/GBSA) energy

calculations[76,77] as well as surface mapping [v. 2020.4][29]. In brief, the S protein and

hACE complex underwent a second round of minimization using the local optimization

feature, the OPLS3e force field, and adopting the Variable Dielectric Surface Generalized

Born (VSGB) continuum solvation model. Prime MM-GBSA calculates the energy of

each individual protein and the complex as a unit. The total predicted binding free energy

for the complexes was then calculated using:

$\Delta G_{binding} = G_{complex} - (G_{S\ protein} + G_{hACE2})$

$\Delta G_{binding} = \Delta E_{MM} + \Delta G_{GB} + \Delta G_{SA}$

$\Delta E_{MM} = \Delta E_{Electrostatic} + \Delta E_{internal} + \Delta E_{VdW}$

These calculations involve various energy terms, including molecular mechanics (MM),

electrostatic, van der Waals, and solvation energies. ΔE_{MM} represents the total gas phase

energy in OPLS3e. $\Delta E_{internal}$ includes bond, angle, and dihedral terms. These calculations

incorporated contributions from gas phase energy, electrostatic interactions, van der

Waals forces, and solvation effects via the generalized Born (GB) method and a nonpolar

contribution. We conducted large-scale low-mode conformational sampling using

MacroModel[25,26,78]. We employed the OPLS4[79] force field, a convergence threshold of

1.0, and an energy window for saving structures of 21.0 kJ/mol, along with a maximum

atom deviation cutoff of 0.5 Å. We used the NetNGlyc 1.0 server[80] to calculate the

glycosylation propensity. The structure files and MM/GBSA energy components, such as

Coulomb, vDW, solvent, complex, are included in the CSV files on our Open Science

Framework Page (https://osf.io/82n73/).

Statistical Analyses

We used the sf[81] and exactextractr[82] packages in program R to calculate the mean

of the urbanization variables within each of our trap buffers. We calculated prevalence

and 95% confidence intervals using the "PropCIs" package[83] with the Agresti-Coull

method due to small sample sizes. We used the glmmTMB package[84] in program R to run

univariate generalized linear mixed models to assess whether seroprevalence rates were

impacted by the following covariates: urbanization (imperviousness), human presence,

and resident population and included species as a random effect. All P-values presented

represent a 2-tailed test.

DATA AVAILABILITY

All study data are included in the article and supporting information. Sequence data from

SARS-CoV-2 viruses from Virginia opossum sequenced in this study are available in the

GISAID database (https://www.gisaid.org/) and NCBI (https://ncbi.nlm.nih.gov) SARS-

CoV-2 sequence repositories.

References

1   World Health Organization *WHO COVID-19 Dashboard*, <

https://covid19.who.int/ > (2020).

2   Drozdz, M. *et al.* Current State of Knowledge about Role of Pets in Zoonotic

Transmission of SARS-CoV-2. *Viruses-Basel* **13**, 1149 (2021).

https://doi.org:10.3390/v13061149

3   Fritz, M. *et al.* High prevalence of SARS-CoV-2 antibodies in pets from COVID-

19+ households. *One Health* **11**, 100192 (2020).

https://doi.org:10.1016/j.onehlt.2020.100192

4   Murphy, H. L. & Ly, H. Understanding the prevalence of SARS-CoV-2 (COVID-

19) exposure in companion, captive, wild, and farmed animals. *Virulence* **12**,

2777-2786 (2021). https://doi.org:10.1080/21505594.2021.1996519

5   Hale, V. L. *et al.* SARS-CoV-2 infection in free-ranging white-tailed deer. *Nature*

602, 481-486 (2022). https://doi.org:10.1038/s41586-021-04353-x

6   Chandler, J. C. *et al.* SARS-CoV-2 exposure in wild white-tailed deer

(*Odocoileus virginianus*). *P Natl Acad Sci USA* **118** (2021).

https://doi.org:10.1073/pnas.2114828118

7   Kuchipudi, S. V. *et al.* Multiple spillovers from humans and onward transmission

of SARS-CoV-2 in white-tailed deer. *P Natl Acad Sci USA* **119** (2022).

https://doi.org:10.1073/pnas.2121644119

8   Aguiló-Gisbert, J. *et al.* First description of SARS-CoV-2 infection in two feral

American mink (*Neovison vison*) caught in the wild. *Animals* **11**, 1422 (2021).

https://doi.org:10.3390/ani11051422

9   Padilla-Blanco, M. *et al.* The finding of the severe acute respiratory syndrome

coronavirus (SARS-CoV-2) in a Wild Eurasian River Otter (*Lutra lutra*)

highlights the need for viral surveillance in wild mustelids. *Front Vet Sci* **9**

(2022). https://doi.org:10.3389/fvets.2022.826991

10   Martínez-Hernández, F. *et al.* Assessing the SARS-CoV-2 threat to wildlife:

Potential risk to a broad range of mammals. *Perspectives in ecology and*

conservation **18**, 223-234 (2020). https://doi.org:10.1016/j.pecon.2020.09.008

11   Meekins, D. A., Gaudreault, N. N. & Richt, J. A. Natural and Experimental

SARS-CoV-2 Infection in Domestic and Wild Animals. *Viruses-Basel* **13**, 1993

(2021). https://doi.org:10.3390/v13101993

12   Griffin, B. D. *et al.* SARS-CoV-2 infection and transmission in the North

American deer mouse. *Nature Communications* **12**, 10 (2021).

https://doi.org:10.1038/s41467-021-23848-9

13   Fagre, A. *et al.* SARS-CoV-2 infection, neuropathogenesis and transmission

among deer mice: Implications for spillback to New World rodents. *PLoS Pathog.*

17, 23 (2021). https://doi.org:10.1371/journal.ppat.1009585

14   Francisco, R. *et al.* Experimental Susceptibility of North American Raccoons

(*Procyon lotor*) and Striped Skunks (*Mephitis mephitis*) to SARS-CoV-2. *Front*

Vet Sci **8**, 10 (2022). https://doi.org:10.3389/fvets.2021.715307

15   Caserta, L. C. *et al.* White-tailed deer (Odocoileus virginianus) may serve as a

wildlife reservoir for nearly extinct SARS-CoV-2 variants of concern. *P Natl*

Acad Sci USA **120**, 12 (2023). https://doi.org:10.1073/pnas.2215067120

16   Thorne, L. G. *et al.* Evolution of enhanced innate immune evasion by SARS-

CoV-2 (vol 602, pg 487, 2022). *Nature* **604**, E14-E14 (2022).

https://doi.org:10.1038/s41586-022-04653-w

17   Pickering, B. *et al.* Divergent SARS-CoV-2 variant emerges in white-tailed deer

with deer-to-human transmission. *Nat. Microbiol*, 2011-2024

https://doi.org:10.1038/s41564-022-01268-9

18   Tan, C. C. S. *et al.* Transmission of SARS-CoV-2 from humans to animals and

potential host adaptation. *Nature Communications* **13**, 13 (2022).

https://doi.org:10.1038/s41467-022-30698-6

19   Markov, P. V. *et al.* The evolution of SARS-CoV-2. *Nature Reviews*

Microbiology **21**, 361-379 (2023). https://doi.org:10.1038/s41579-023-00878-2

20   PhyloPic. *Permalink: Image Collection*,

<https://www.phylopic.org/permalinks/5711d2ef512cd5fef54cd101c52ccf839ad2

a5e92d60323f8cbfddba5003df3a> (2023).

21   Centers for Disease Control and Prevention.   (U.S. Department of Health and

Human Services, CDC, Atlanta, GA, 2023).

22   Taylor, P. C. *et al.* Neutralizing monoclonal antibodies for treatment of COVID-

19. *Nat. Rev. Immunol.* **21**, 382-393 (2021). https://doi.org:10.1038/s41577-021-

00542-x

23   Xu, Z. *et al.* Integrated strategy of RNA-sequencing and network pharmacology

for exploring the protective mechanism of Shen-Shi-Jiang-Zhuo formula in rat

with non-alcoholic fatty liver disease. *Pharm Biol* **60**, 1819-1838 (2022).

https://doi.org:10.1080/13880209.2022.2106250

24   Kang, Y. F. *et al.* Rapid Development of SARS-CoV-2 Spike Protein Receptor-

Binding Domain Self-Assembled Nanoparticle Vaccine Candidates. *Acs Nano* **15**,

2738-2752 (2021). https://doi.org:10.1021/acsnano.0c08379

25   Watts, K. S., Dalal, P., Tebben, A. J., Cheney, D. L. & Shelley, J. C. Macrocycle

Conformational Sampling with MacroModel. *Journal of Chemical Information*

and Modeling **54**, 2680-2696 (2014). https://doi.org:10.1021/ci5001696

26   Kolossvary, I. & Guida, W. C. Low Mode Search. An Efficient, Automated

Computational Method for Conformational Analysis:□ Application to Cyclic and

Acyclic Alkanes and Cyclic Peptides. *Journal of the American Chemical Society*

118, 5011-5019 (1996). https://doi.org:doi.org/10.1021/ja952478m

27   Singh, V. *et al.* Stroke increases the expression of ACE2, the SARS-CoV-2

binding receptor, in murine lungs. *Brain Behav Immun* **94**, 458-462 (2021).

https://doi.org:10.1016/j.bbi.2021.01.039

28   Li, H. D. *et al.* Greener liquid-phase synthesis and the ACE inhibitory structure-

activity relationship of an anti-SARS octapeptide. *Org Biomol Chem* **18**, 8433-

8442 (2020). https://doi.org:10.1039/d0ob01948h

29   Beard, H., Cholleti, A., Pearlman, D., Sherman, W. & Loving, K. A. Applying

Physics-Based Scoring to Calculate Free Energies of Binding for Single Amino

Acid Mutations in Protein-Protein Complexes. *Plos One* **8**, e82849 (2013).

https://doi.org:10.1371/journal.pone.0082849

30   Koehler, M. *et al.* Molecular insights into receptor binding energetics and

neutralization of SARS-CoV-2 variants. *Nature Communications* **12**, 6977

(2021). https://doi.org:10.1038/s41467-021-27325-1

31   Yang, Q. *et al.* Role for N-glycans and calnexin-calreticulin chaperones in SARS-

CoV-2 Spike maturation and viral infectivity. *Science Advances* **8**, eabq8678

(2022). https://doi.org:10.1126/sciadv.abq8678

32   Huang, M. *et al.* Atlas of currently available human neutralizing antibodies

against SARS-CoV-2 and escape by Omicron sub-variants

BA.1/BA.1.1/BA.2/BA.3. *Immunity* **55**, 1501-1514 (2022).

https://doi.org:10.1016/j.immuni.2022.06.005

33   Shajahan, A., Pepi, L. E., Rouhani, D. S., Heiss, C. & Azadi, P. Glycosylation of

SARS-CoV-2: structural and functional insights. *Analytical and Bioanalytical*

Chemistry **413**, 7179-7193 (2021). https://doi.org:10.1007/s00216-021-03499-x

34   Fenton, A. & Pedersen, A. B. Community epidemiology framework for

classifying disease threats. *Emerg Infect Dis* **11**, 1815-1821 (2005).

35   Fischhoff, I. R., Castellanos, A. A., Rodrigues, J. P., Varsani, A. & Han, B. A.

Predicting the zoonotic capacity of mammals to transmit SARS-CoV-2.

Proceedings of the Royal Society B **288**, 20211651 (2021).

https://doi.org:10.1098/rspb.2021.1651

36   Bosco-Lauth, A. M. *et al.* Peridomestic mammal susceptibility to severe acute

respiratory syndrome coronavirus 2 infection. *Emerg Infect Dis* **27**, 2073 (2021).

https://doi.org:10.3201/eid2708.210180

37   Samanta, A., Alam, S. S. M., Ali, S. & Hoque, M. Evaluating the transmission

feasibility of SARS-CoV-2 Omicron (B.1.1.529) variant to 143 mammalian hosts:

insights from S protein RBD and host ACE2 interaction studies. *Functional &*

Integrative Genomics **23** (2023). https://doi.org:10.1007/s10142-023-00962-z

38   Botten, J. *et al.* Persistent Sin Nombre virus infection in the deer mouse

(*Peromyscus maniculatus)* model: Sites of replication and strand-specific

expression. *Journal of Virology* **77**, 1540-1550 (2003).

https://doi.org:10.1128/jvi.77.2.1540-1550.2002

39   Franklin, A. B. & Bevins, S. N. Spillover of SARS-CoV-2 into novel wild hosts

in North America: A con- ceptual model for perpetuation of the pathogen. *Sci.*

Total Environ. **733** (2020). https://doi.org:10.1016/j.scitotenv.2020.139358

40   Han, J. *et al.* Municipal solid waste, an overlooked route of transmission for the

severe acute respiratory syndrome coronavirus 2: a review. *Environ. Chem. Lett.*,

1-15 (2022). https://doi.org:10.1007/s10311-022-01512-y

41   Aberle, M. A., Langwig, K. E., Adelman, J. S. & Hawley, D. M. Effects of bird

feeder density on the foraging behaviors of a backyard songbird (the House Finch,

Haemorhous mexicanus) subject to seasonal disease outbreaks. *Canadian Journal*

of Zoology **98**, 611-621 (2020). https://doi.org:10.1139/cjz-2019-0282

42   Civitello, D. J., Allman, B. E., Morozumi, C. & Rohr, J. R. Assessing the direct

and indirect effects of food provisioning and nutrient enrichment on wildlife

infectious disease dynamics. *Philos T R Soc B* **373** (2018).

https://doi.org:10.1098/rstb.2017.0101

43   Longcore, T., Rich, C. & Sullivan, L. M. Critical Assessment of Claims

Regarding Management of Feral Cats by Trap-Neuter-Return. *Conserv Biol* **23**,

887-894 (2009). https://doi.org:10.1111/j.1523-1739.2009.01174.x

44   Ceci, A. *et al.* Development and implementation of a scalable and versatile test

for COVID-19 diagnostics in rural communities. *Nature Communications* **12**, 1-

14 (2021). https://doi.org:10.1038/s41467-021-24552-4

45   Galipeau, Y. *et al.* Relative Ratios of Human Seasonal Coronavirus Antibodies

Predict the Efficiency of Cross-Neutralization of SARS-CoV-2 Spike Binding to

ACE2. *EBioMedicine* **74**, 14 (2021). https://doi.org:10.1016/j.ebiom.2021.103700

46   Lapidus, S. *et al. Plasmodium* infection is associated with cross-reactive

antibodies to carbohydrate epitopes on the SARS-CoV-2 Spike protein. *Sci Rep-*

Uk **12** (2022). https://doi.org:10.1038/s41598-022-26709-7

47   Kharlamova, N. *et al.* False Positive Results in SARS-CoV-2 Serological Tests

for Samples From Patients With Chronic Inflammatory Diseases. *Frontiers in*

Immunology **12** (2021). https://doi.org:10.3389/fimmu.2021.666114

48   Olival, K. J. *et al.* Host and viral traits predict zoonotic spillover from mammals.

Nature **546**, 646-+ (2017). https://doi.org:10.1038/nature22975

49   Han, B. A., Schmidt, J. P., Bowden, S. E. & Drake, J. M. Rodent reservoirs of

future zoonotic diseases. *P Natl Acad Sci USA* **112**, 7039-7044 (2015).

https://doi.org:10.1073/pnas.1501598112

50   Bentler, K. T., Gossett, D. N. & Root, J. J. A novel isoflurane anesthesia

induction system for raccoons. *Wildlife Society Bulletin* **36**, 807-812 (2012).

https://doi.org:10.1002/wsb.193

51   Gomez, A. *et al.* Land use and West Nile Virus seroprevalence in wild mammals.

Emerg Infect Dis **14**, 962-965 (2008). https://doi.org:10.3201/eid1406.070352

52   Muruato, A. E. *et al.* A high-throughput neutralizing antibody assay for COVID-

19 diagnosis and vaccine evaluation. *Nature communications* **11**, 4059 (2020).

53   Marano, J. M. & Weger-Lucarelli, J. Replication in the presence of dengue

convalescent serum impacts Zika virus neutralization sensitivity and fitness.

Frontiers in Cellular and Infection Microbiology **13**, 1130749 (2023).

54   Feng, A. J. *et al.* Transmission of SARS-CoV-2 in free-ranging white-tailed deer

in the United States. *Nature Communications* **14** (2023).

https://doi.org:10.1038/s41467-023-39782-x

55   Liu, J. L. *et al.* Stabilization of a Broadly Neutralizing Anti-Chikungunya Virus

Single Domain Antibody. *Front Med-Lausanne* **8** (2021).

https://doi.org:10.3389/fmed.2021.626028

56   Bosco-Lauth, A. M. *et al.* Experimental infection of domestic dogs and cats with

SARS-CoV-2: Pathogenesis, transmission, and response to reexposure in cats. *P*

Natl Acad Sci USA **117**, 26382-26388 (2020).

https://doi.org:10.1073/pnas.2013102117

57   Bosco-Lauth, A. M. *et al.* Susceptibility of livestock to SARS-CoV-2 infection.

Emerg Microbes Infec **10**, 2199-2201 (2021).

https://doi.org:10.1080/22221751.2021.2003724

58   Shriner, S. A. *et al.* SARS-CoV-2 Exposure in Escaped Mink, Utah, USA. *Emerg*

Infect Dis **27**, 988-990 (2021). https://doi.org:10.3201/eid2703.204444

59   Palermo, P. M., Orbegozo, J., Watts, D. M. & Morrill, J. C. SARS-CoV-2

Neutralizing Antibodies in White-Tailed Deer from Texas. *Vector-Borne Zoonot*,

62-64 (2021). https://doi.org:10.1089/vbz.2021.0094

60   Dewitz, J. & U.S. Geological Survey in *US Geological Survey*   (2021).

61   Wickham, J., Stehman, S. V., Sorenson, D. G., Gass, L. & Dewitz, J. A. Thematic

accuracy assessment of the NLCD 2016 land cover for the conterminous United

States. *Remote Sens Environ* **257**, 112357 (2021).

https://doi.org:10.1016/j.rse.2021.112357

62   Center for International Earth Science Information Network (CIESIN).   (ed

Columbia University) (NASA Socioeconomic Data and Applications Center

(SEDAC), Palisades, New York, 2018).

63   Calenge, C. The package "adehabitat" for the R software: A tool for the analysis

of space and habitat use by animals. *Ecol Model* **197**, 516-519 (2006).

https://doi.org:10.1016/j.ecolmodel.2006.03.017

64   R: A Language and Environment for Statistical Computing (R Foundation for

Statistical Computing, Vienna, Austria, 2021).

65   O'Toole, A. *et al.* Assignment of epidemiological lineages in an emerging

pandemic using the pangolin tool. *Virus Evol* **7** (2021).

https://doi.org:10.1093/ve/veab064

66   Wright, E. S. RNAconTest: comparing tools for noncoding RNA multiple

sequence alignment based on structural consistency. *Rna* **26**, 531-540 (2020).

https://doi.org:10.1261/rna.073015.119

67   Sagulenko, P., Puller, V. & Neher, R. A. TreeTime: Maximum-likelihood

phylodynamic analysis. *Virus Evol* **4** (2018). https://doi.org:10.1093/ve/vex042

68   Turakhia, Y. *et al.* Ultrafast Sample placement on Existing tRees (UShER)

enables real-time phylogenetics for the SARS-CoV-2 pandemic. *Nat Genet* **53**,

809-816 (2021). https://doi.org:10.1038/s41588-021-00862-7

69   Hadfield, J. *et al.* Nextstrain: real-time tracking of pathogen evolution.

Bioinformatics **34**, 4121-4123 (2018).

https://doi.org:10.1093/bioinformatics/bty407

70   Aksamentov, I., Roemer, C., Hodcroft, E. B. & Neher, R. A. Nextclade: clade

assignment, mutation calling and quality control for viral genomes. *The Journal*

of Open Source Software **6**, 3773 (2021). https://doi.org:10.21105/joss.03773

71   (2024).

72   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for

phylogenetic tree display and annotation. *Nucleic Acids Research* **49**, W293-

W296 (2021). https://doi.org:10.1093/nar/gkab301

73   Elbe, S. & Buckland-Merrett, G. Data, disease and diplomacy: GISAID's

innovative contribution to global health. *Global Challenges* **1**, 33-46 (2017).

https://doi.org:10.1002/gch2.1018

74   Harder, E. *et al.* OPLS3: A Force Field Providing Broad Coverage of Drug-like

Small Molecules and Proteins. *J Chem Theory Comput* **12**, 281-296 (2016).

https://doi.org:10.1021/acs.jctc.5b00864

75   Shivakumar, D. *et al.* Prediction of Absolute Solvation Free Energies using

Molecular Dynamics Free Energy Perturbation and the OPLS Force Field. *J Chem*

Theory Comput **6**, 1509-1519 (2010). https://doi.org:10.1021/ct900587b

76   Jacobson, M. P. *et al.* A hierarchical approach to all-atom protein loop prediction.

Proteins **55**, 351-367 (2004). https://doi.org:10.1002/prot.10613

77   Jacobson, M. P., Friesner, R. A., Xiang, Z. X. & Honig, B. On the role of the

crystal environment in determining protein side-chain conformations. *J Mol Biol*

320, 597-608 (2002). https://doi.org:10.1016/S0022-2836(02)00470-9

78   Mohamadi, F. *et al.* Macromodel—an integrated software system for modeling

organic and bioorganic molecules using molecular mechanics. *Journal of*

Computational Chemistry **11**, 440-467 (1990).

79   Lu, C. *et al.* OPLS4: Improving Force Field Accuracy on Challenging Regimes of

Chemical Space. *J Chem Theory Comput* **17**, 4291-4300 (2021).

https://doi.org:10.1021/acs.jctc.1c00302

80   Julenius, K. NetCGlyc 1.0: prediction of mammalian C-mannosylation sites.

Glycobiology **17**, 868-876 (2007). https://doi.org:10.1093/glycob/cwm050

81   Pebesma, E. Simple Features for R: Standardized Support for Spatial Vector Data.

The R journal **10**, 439-446 (2018).

82   exactextractr: Fast extraction from raster datasets using polygons (2020).

83   PropCIs: Various Confidence Interval Methods for Proportions v. 0.3-0 (2018).

84   Brooks, M. E. *et al.* glmmTMB Balances Speed and Flexibility Among Packages

for Zero-inflated Generalized Linear Mixed Modeling. *R J* **9**, 378-400 (2017).

https://doi.org:Doi 10.32614/Rj-2017-066

ACKNOWLEDGMENTS

We thank the following for land access: J. Eustis and New River Land Trust; B. Brodie,

J. Jones and Mountain Lake Biological Research Station; R. Powers, K. Slusher, A.

McGee and Roanoke Parks and Recreation; V. Malzone, G. Gorecki, S. Sweeney, J.

Elliott, and Virginia State Parks; G. O'Malley, A. Lewis, H. Wander, K. Malewicz, V.

Corbin, R. Cohen, S. Whitehead, N. Laggan, A. Grimaudo, and G. Blanvillain. We thank

the following for help with study design and collection: M. Fisher, J. Ivan, K. Bentler, S.

Yamada, C. Cereghino, K. Pierce and the Wildlife Center of Virginia, J. Riley and the

Blue Ridge Wildlife Center, and H. Olsen-Hodges and the Southwest Virginia Wildlife

Center. Additionally, we thank G. Eastwood, A. Gomez, A. M. Kilpatrick, and NEON

Biorepository for sharing serum samples with us. We thank the following for help with

field work: M. Hellier, J. White, and M. Dwyer. This project was supported by funds

from NSF DEB-1911853 to K.E.L. and J.R.H., the Virginia Tech FBRI to C.V.F., One

Health award from VCOM to J.W, and the American Rescue Plan Act through USDA

APHIS. The findings and conclusions in this publication are those of the authors and

should not be construed to represent any official USDA or U.S. government

determination or policy. We gratefully acknowledge all data contributors, including

authors and their originating laboratories responsible for obtaining the specimens, and the

submitting laboratories for generating the genetic sequence and metadata and sharing via

the GISAID Initiative, on which this research is based. All genome sequences and

associated metadata in this dataset are published in GISAID's EpiCoV database. To view

the contributors of each individual sequence with details such as accession number, Virus

name, Collection date, Originating Lab and Submitting Lab and the list of Authors, visit

10.55876/gis8.231109hc

AUTHOR CONTRIBUTIONS: A.R.G., K.E.L., and J.R.H. designed field research;

A.R.G., C.D.K., M.J.K, K.E.L, and J.R.H. performed field research; A.R.G., J.M., P.R.

K.E.L., J.W, K.L.B., C.R, A.C., R.B., C.V.F., and J.R.H. contributed to field data

analysis; C.R., R.B., K.M.K., A.C., M.U., A.B., K.L.B., and C.V.F contributed to

molecular and genomic analysis; K.E.L., J.W., C.V.F., and J.R.H. provided project

administration; Members of the Molecular Diagnostics Laboratory provided technical

support; and all authors aided in interpreting results and to writing the paper.

DECLARATION OF INTERESTS

The authors declare no competing interests.