African eggplant-associated virus: characterization of a novel

tobamovirus identified from *Solanum macrocarpon* and assessment

of its potential impact on tomato and pepper crops

Anne K.J. Giesbers[1,2]*, Annelien Roenhorst[1,2], Martijn F. Schenk[2], Marcel Westenberg[1,2], Marleen

Botermans[1,2]

[1] Netherlands Institute for Vectors, Invasive plants and Plant health, Wageningen, The Netherlands

[2] National Plant Protection Organization (NPPO-NL), Netherlands Food and Consumer Product Safety

Authority (NVWA), Wageningen, The Netherlands

* Corresponding author: a.k.j.giesbers@nvwa.nl

Abstract

A novel tobamovirus was identified in a fruit of *Solanum macrocarpon* imported into the Netherlands in 2018. This virus was further characterized in terms of host range, pathotype and genomic properties, because many tobamoviruses have the potential to cause severe damage in important crops. In the original fruit, two different genotypes of the novel virus were present. The virus was able to infect multiple plant species from the *Solanaceae* family after mechanical inoculation, as well as a member of the *Apiaceae* family. These species included economically important crops such as tomato and pepper, as well as eggplant and petunia. Both tomato and pepper germplasm were shown to harbor resistance against the novel virus. Since most commercial tomato and pepper varieties grown in European greenhouses harbor these relevant resistances, the risk of infection and subsequent impact on these crops is likely to be low in Europe. Assessment of the potential threat to eggplant, petunia, and other susceptible species needs further work. In conclusion, this study provides a first assessment of the potential phytosanitary risks of a newly discovered tobamovirus, which was tentatively named African eggplant-associated virus.

Introduction

The International Plant Protection Convention aims to safeguard agriculture and natural resources against the entry, establishment and spread of economically and environmentally significant pests (1). In this respect, import inspections are an important tool to prevent the introduction of regulated plant pests and may serve as an early warning system on emerging pests. In 2018, a phytosanitary inspector in the Netherlands intercepted a fruit of *Solanum macrocarpon* from Mexico, which showed diffuse chlorotic mottling (Fig 1). *Solanum macrocarpon*, also known as African eggplant, is a small tropical perennial originating from Africa that is nowadays cultivated in many countries for food, medicinal purposes and as an ornamental (2, 3). Since the observed symptoms on the fruit indicated a potential viral infection, samples of this fruit were analyzed by bioassay, serology and

high-throughput sequencing analysis. The results indicated the presence of a so far unknown

tobamovirus.

Fig 1. *Solanum macrocarpon* fruit with diffuse chlorotic mottling.

The genus *Tobamovirus* (family *Virgaviridae*) currently contains 37 accepted species according to the

2021 release of the International Committee on Taxonomy of Viruses (ICTV)

(https://talk.ictvonline.org/taxonomy/), including some devastating plant viruses. Tobamoviruses are

highly stable, contagious and difficult to control. Tobamoviruses can rapidly spread locally by

mechanical contact, while transmission via seeds enables them to spread over longer distances, as

exemplified by the recent rapid spread of tomato brown rugose fruit virus (ToBRFV) (4-6) and tomato

mottle mosaic virus (ToMMV) (7). Other tobamoviruses that can cause significant economic losses in

crops if not controlled effectively, include cucumber green mottle mosaic virus (8), tobacco mosaic

virus (TMV) and tomato mosaic virus (ToMV) (9).

The potential of tobamoviruses to spread rapidly and cause severe damage has prompted us to

prioritize this novel virus for further characterization in terms of genomic properties, host range and

pathotype. Tobamoviruses can be divided into at least eight subgroups based on their complete

nucleotide sequence, the amino acid sequences of each individual viral protein and their host plant

family (10). These subgroups consist of species that can infect plant species within the families

Apocynaceae, Brassicaceae, Cactaceae, Cucurbitaceae, Leguminosae, Malvaceae, Passifloraceae, and

Solanaceae.

To assess the biological properties and potential impact of the novel tobamovirus from *S.*

macrocarpon, host range studies focused on solanaceous crops, in particular tomato (*Solanum*

lycopersicum) and pepper (*Capscium* spp.). Both tomato and pepper germplasm are known to harbor

tobamovirus resistance genes. In tomato, the *Tm-1* gene, and the *Tm-2* and *Tm2^2* alleles, originating

from wild tomato species, have been used extensively to protect against tobacco mosaic virus and

tomato mosaic virus (11). In pepper, the four *L* genes or alleles L^1, L^2, L^3 and L^4, each provide

resistance to an increasing number of tobamovirus pathotypes (12). Therefore, tomato and pepper

accessions harboring these resistances were tested to assess whether the novel tobamovirus poses a

threat to these crops.

The results of the molecular and biological characterization of this novel tobamovirus from *S.*

macrocarpon, which we tentatively named ''African eggplant-associated virus'' contribute to the

assessment of its potential risks, so that authorities can determine the need for preventive control

measures.

Materials and Methods

Virus source, propagation and maintenance

80   A novel tobamovirus was identified from a fruit of *S. macrocarpon* imported from Mexico. The isolate

81   was propagated and maintained by successive mechanical passages in *Nicotiana occidentalis* P1.

82   Young leaves were harvested just after systemic symptoms appeared and stored at -80°C for further

83   experiments as well as long-term preservation.

Mechanical inoculation and growing conditions of test plants

Symptomatic tissue (fruit or leaf material) was used for mechanical inoculation of test plants according to the procedure and conditions as described by Verhoeven and Roenhorst (2000) (13). All test plants were grown from seed, except for *Solanum tuberosum* which was grown from tubers. Tested plant species and varieties are specified in the Results section.

Host range and pathotype analysis

Nineteen species from six plant families were inoculated with the virus from *S. macrocarpon*. Per species, four plants were inoculated with the virus and four plants with inoculation buffer only (mock treatment). All plants were assessed for local and systemic disease symptoms twice a week for a total duration of four weeks and tested by DAS-ELISA 10-25 days after inoculation. Symptoms in *S. macrocarpon* were assessed until the fruits had matured. In addition, tomato and pepper accessions with different tobamovirus resistance genes (tomato: *Tm1*, *Tm2* and *Tm2²*; pepper: *L¹*, *L²*, *L³* and *L⁴*) were tested in an experiment with the same set-up. These accessions and information on their resistance genes were kindly provided by the Centre for Genetic Resources, the Netherlands (CGN).

Double Antibody Sandwich Enzyme-Linked Immunosorbent Assay (DAS-ELISA)

DAS-ELISA (14) was performed using polyclonal antisera raised against bell pepper mottle virus (BPMoV), pepper mild mottle virus (PMMoV), tobacco mosaic virus (TMV), and tomato mosaic virus (ToMV-D) from Prime Diagnostics, the Netherlands. Host range and pathotype experiments were conducted with PMMoV antiserum. Antisera were diluted at 1:1000 from a 1 mg/ml stock. Buffers used were: coating buffer (15 mM Na_2CO_3, 34.9 mM $NaHCO_3$, pH 9.6), homogenization buffer (2.9 mM KH_2PO_4, 2.7 mM KCl, 16 mM $Na_2HPO_4 \cdot 12H_2O$, 136.9 mM NaCl, containing Tween 20 at 0.5 ml/l, polyvinyl pyrrolidone at 20 g/l and albumin from bovine serum at 2.0 g/l, pH 7.4), and substrate buffer (p-nitrophenyl phosphate at 0.75 mg/ml in diethanolamine at 97.5 ml/l, pH 9.8). A volume of 200 µl was added to each well. Each sample was tested in duplicate and contained a pool of tissue from four plants of the same combination of plant species and inoculum. Samples were taken from

non-inoculated young leaves or fruits and homogenized at an approximate ratio of 1:10 (w:v), using

extraction bags and a hand-held homogenizer (Bioreba). Coating and test sample incubations were

overnight at 4°C, substrate incubations for 1-2 hours at 37°C. Optical Density (OD) at 405 nm was

measured with a microplate reader (BioRad iMark) in duplicate wells from which an average OD

value was calculated. Samples were considered positive when the OD value was at least twice the

value of the negative controls (mock-inoculated healthy plants).

Illumina sequencing and analysis

Total RNA was extracted using the RNeasy Plant Mini kit (Qiagen, Germany) following the

manufacturer's instructions. The RNA extract was sent to GenomeScan (The Netherlands) for

generation of 2 Gb of Illumina RNAseq 150PE (paired-end) data per sample. Sequencing was

performed on an Illumina NovaSeq6000. RNAseq data were analyzed in CLC Genomics workbench

v21.0.4 (Qiagen, Germany) and run in a custom workflow built for detection of *de novo* assembled

viral contigs (15). Consensus sequences (>100 nt; ARC>10) from *de novo* assembled contigs were

analyzed using megaBLAST and DIAMOND (16) with a local installation of the NCBI nr/nt database

downloaded from http://ftp.ncbi.nlm.nih.gov/. BLAST results were visualized in Krona (bitscore

threshold=25) (17). Putative viral sequences were further analyzed using Geneious Prime 2021.1.1

(Biomatters, New Zealand). Genomes of *Tobamovirus* species were aligned using MAFFT alignment

with automatic settings in Geneious, for the species of which a complete reference nucleotide

sequence was available in GenBank. A Neighbor-Joining tree was constructed using the Geneious

tree builder based on Tamura-Nei distances, bootstrap resampling, random seed 925,045, number of

replicates 1,000, and a support threshold at 70%.

Results

To determine whether the symptoms on the *S. macrocarpon* fruit could result from a viral infection,

a homogenate of symptomatic tissue was inoculated on a range of test plants routinely used as

indicators in virus diagnosis. The reaction on the inoculated test plants, six days post inoculation,

indicated the presence of a tobamovirus (Table 1). Therefore, symptomatic leaves of *N. occidentalis*

P1 were tested for different tobamoviruses by DAS-ELISA. Reactions were clearly positive for antisera

raised against BPMoV and PMMoV, but negative for TMV and ToMV-D (Table 2). Negative controls

(mock-inoculated plants) and homologous (positive) controls showed their expected outcomes.

Based on these results, the virus from *S. macrocarpon* appeared to be a tobamovirus that is

serologically related to BPMoV and PMMoV. The PMMoV antiserum was used to detect the virus in

all subsequent experiments.

Table 1. Symptomatology of indicator plants inoculated with the homogenate from the intercepted
S. macrocarpon **fruit.**

Indicator species	Symptoms	
	Local	Systemic
Datura stramonium	NL (2)	-
Nicotiana glutinosa	NL (2)	-
N. occicentalis "P1"	-	~CV, G, LN (2)
N. tabacum "White Burley"	CL, NL (2)	-
Solanum lycopersicum "Moneymaker"	-	-

143
144 -: no symptoms, ~: mild symptoms, CV: chlorotic vein banding, CL: chlorotic lesions, G: growth inhibition, LN:
145 leaf narrowing, NL: necrotic lesions. The first symptoms appeared six days post inoculation. Per species two
146 plants were inoculated. The number of plants showing specific symptoms is indicated in brackets.

147 **Table 2. Reaction of inoculated N. occidentalis P1 plants by DAS-ELISA.**

Antiserum	OD$_{405}$*
BPMoV	3.050
PMMoV	1.799
TMV	0.228
ToMV-D	0.269

*Average OD values after 90 minutes of substrate incubation. OD values were considered positive (red) when ≥
2x the OD value of mock-inoculated controls, otherwise OD values were considered negative (green).

For further characterization, the viral genome sequence of the putative tobamovirus from *S.*

macrocarpon was determined by Illumina sequencing analysis of the original fruit sample. *De novo*

assembly resulted in a near-complete genome sequence of two genotypes, each consisting of a single

stranded RNA with four predicted open readings frames (ORFs), corresponding to the genome

organization of tobamoviruses (18). These respective ORFs encode a small replicase subunit, an RNA-

dependent RNA polymerase (translated through ribosomal readthrough at the stop codon of the

small replicase subunit), a movement protein, and a coat protein. Genotype 1 (GenBank accession

number OP729507) and genotype 2 (GenBank accession number OP729508) contained 6228 and

6356 nucleotides respectively, with 96.1% nucleotide similarity. No other viruses were detected.

NCBI BLAST searches (https://blast.ncbi.nlm.nih.gov/Blast.cgi) indicated that both genotypes of the

virus showed most resemblance to isolates of PMMoV, with a maximum query coverage of 97% and

a nucleotide identity of 75%. Since the nucleotide identity with PMMoV is considerably lower than

the tobamovirus species demarcation threshold of 90% (18), the virus should be considered as a

member of a new species, for which we propose the name "African eggplant-associated virus". A

phylogenetic analysis including genome sequences of other tobamoviruses indicated that African

eggplant-associated virus is most closely related to members of species infecting solanaceous plants,

in particular PMMoV and tropical soda apple mosaic virus (Fig 2).

Fig 2. Unrooted neighbor-joining tree with (near) complete nucleotide sequences of the novel virus
(red) and other members of the genus *Tobamovirus* including their GenBank accession numbers.
The bar represents 0.1 substitutions per site. This tree was constructed by the neighbor-joining
algorithm in Geneious software, using the genetic distance model Tamura-Nei, with 1,000 bootstrap

replicates and the support threshold set at 70%. Support values (%) that were less than 100% are
indicated next to the supported internal branches.

BLAST searches also indicated a close resemblance to a bacteriophage isolate (GenBank accession

number BK018885), which aligned with the first 3,874 nucleotides of the novel virus, corresponding

to a query coverage of 60%. The nucleotide identity was 95.7% and 99.5% for genotype 1 and

genotype 2, respectively. Most probably, this sequence was incorrectly assigned to a bacteriophage.

To gain more insight into the potential host range of the novel virus, it was mechanically transferred

from *N. occidentalis* P1 to test plants (Table 3). Nine out of 19 tested species, including *S.*

macrocarpon, were susceptible based on symptomatology (i.e. systemic symptoms) and/or detection

by DAS-ELISA, of which *Ammi majus* was the only non-solanaceous plant. *A. majus*, *S. lycopersicum*

"Moneymaker" and *Solanum melongena* "Violetta Lunga 2" showed asymptomatic or latent

infections. The tested *Amaranthaceae, Brassicaceae*, *Cucurtitaceae*, and *Fabaceae* species remained

uninfected. Negative and positive controls showed their expected outcomes. Figure 3 illustrates the

most obvious symptoms per species.

Family	Species	symptoms		OD$_{405}$
		local	systemic	PMMoV
Amaranthaceae	*Chenopodium album*	CL (4)	-	0.022
	Chenopodium quinoa	CL (4), ~CR (2)	-	0.135
Apiaceae	*Ammi majus*	- / -*	-; -*	1.169 / 0.681*
Brassicaceae	*Brassica rapa* subsp. *pekinensis* "Granaat"	-	-	0.105
Cucurbitaceae	*Cucumis sativus* "Chinese Slangen"	-	-	0.000
Fabaceae	*Pisum sativum* "Kelvedon Wonder"	-	-	0.052
	Vicia faba "Witkiem"	-	-	0.071
Solanaceae	*Capsicum annuum* "Westlandse Grote Zoete"	-	C, M, Ms (4)	1.353
	Datura stramonium	NL (4)	-	0.149
	Nicotiana glutinosa	NL (4)	-	0.048
	Nicotiana hesperis "67A"	-	C, G, NLE, R (4)	0.768
	Nicotiana occidentalis "P1"	-	C, G, LC, NLE, R (4)	1.653
	Nicotiana tabacum "White Burley"	CL (4), NL (1)	-	0.101
	Petunia hybrida	NL, NV (1)	LC (3) G, M, R (4)	0.989
	Solanum lycopersicum "Moneymaker"	-	-	0.631
	Solanum macrocarpon	-	CV, ~M (2)	2.340
	Solanum melongena "Violetta Lunga 2"	- / -*	- / -*	0.763 / 1.788*
	Solanum nigrum	-	CL (3) R (2)	0.647
	Solanum tuberosum "Nicola"	-	-	0.090

Per species four plants were inoculated; the number of plants showing specific symptoms is indicated in

brackets. Average OD values after two hours of substrate incubation were considered positive (red) when ≥ 2x

the value of mock-inoculated controls, otherwise values were considered negative (green). -: no symptoms, ~:

mild symptoms, C: chlorosis, CL: chlorotic lesions, CR: chlorotic rings, CV: chlorotic vein banding, G: growth

inhibition, LC: leaf curling, M: mottling, Ms: mosaic, NL: necrotic lesions, NLE: necrotic leaf edge, NV: necrotic

vein banding, R: rugosity. * results of a repeated experiment on four new plants.

204
Fig 3. Symptoms on test plants upon inoculation with the novel tobamovirus.
a) *Chenopodium album*: local chlorotic lesions, b) *Capsicum annuum* "Westlandse Grote Zoete":
systemic chlorosis, c) *Datura stramonium*: local necrotic lesions, d) *Nicotiona glutinosa*: local necrotic
lesions, e) *Nicotiana hesperis* "67A": systemic chlorosis, growth inhibition, rugosity, f) *Nicotiana*
occidentalis "P1": systemic chlorosis, growth inhibition, leaf curling, rugosity, g) *Nicotiana tabacum*
"White Burley": local chlorotic lesions, h) *Petunia hybrida*: local necrotic lesions, systemic leaf curling,
growth inhibition, rugosity, i) *Solanum macrocarpon*: systemic chlorotic vein banding, j) *Solanum*
nigrum: systemic chlorotic lesions, rugosity.

To fulfill Koch's postulates, *S. macrocarpon* plants were observed until fruits had matured, 147 days

after inoculation. The initial virus symptoms on the leaves had disappeared completely ten days after

their first observation. For mature plants and fruits, the virus was still detected by DAS-ELISA, but

there were no phenotypic differences compared to mock-inoculated plants and fruits. Therefore,

associating the novel virus with the symptoms observed on the original fruit (Fig 1), according to

Koch's postulates, was not possible.

Furthermore, in the inoculated *S. macrocarpon* plants only genotype 2 was detected by Illumina

sequence analysis. To trace the disappearance of genotype 1, subsequent passages in *N. occidentalis*

P1 used for virus propagation were analyzed. Both genotypes were still detected in all passages and

their nucleotide sequence remained exactly the same, though the relative frequency of genotype 1

decreased with each passage. Based on sequence depth, the approximate ratios of genotype 1 to

genotype 2 were 2:3 for the original *S. macrocarpon* fruit, 1:4 for *N. occidentalis* P1 (passage 1), and

1:14 for *N. occidentalis* P1 (passage 2) (S1 Fig).

To assess whether the novel tobamovirus poses a threat to tomato and pepper crops, accessions

harboring different tobamovirus resistances were inoculated (Table 4). These included (combinations

of) *Tm1*, *Tm2*, and *Tm2^2*, and *L^1*, *L^2*, *L^3* and *L^4* resistance genes/alleles for tomato and pepper,

respectively. Even though all tomato accessions remained asymptomatic, only the four accessions

harboring the combination of *Tm1* and *Tm2/Tm2^2* tested negative by DAS-ELISA. Interestingly, the

accessions that harbored *Tm1*, *Tm2* or *Tm2^2* separately tested positive by DAS-ELISA. Therefore, a

combination of these *Tm* resistance genes seems to account for resistance against the novel

tobamovirus.

Table 4. Pathotype analysis: symptomatology and DAS-ELISA results of tomato (S. lycopersicum) and pepper (Capsicum spp.) accessions harboring different (combinations of) tobamovirus resistance genes after inoculation with the novel tobamovirus.

| Family | Species | Tobamovirus resistance | symptoms | | DAS-ELISA |
			local	systemic	PMMoV
Solanaceae	*Capsicum annuum* "Westlandse Grote Zoete"		CL, R(2)*	~C (2)*	0.371 (local), 2.156 (systemic)
	Capsicum annuum "CGN19226"	L1	CL, NL, R (4)	-	0.109
	Capsicum frutescens "CGN21546"	L2	CL, NL, R (4)	-	0.240
	Capsicum chinense "CGN21557"	L3	NL (4)	-	0.102
	Capsicum chacoense "CGN21477"	L4	LA, NL, NR (4)	-	0.165
	Solanum lycopersicum "Moneymaker"		-	-	2.118
	Solanum lycopersicum "CGN24212"	Tm1	-	-	1.975
	Solanum lycopersicum "CGN24213"	Tm2	-	-	2.131
	Solanum lycopersicum "CGN24214"	Tm2^2	-	-	2.047
	Solanum lycopersicum "CGN24218"	Tm1, Tm2/Tm2^2	-	-	0.084
	Solanum lycopersicum "Arbason"	Tm1, Tm2/Tm2^2	-	-	0.109
	Solanum lycopersicum "Cindel"	Tm1, Tm2/Tm2^2	-	-	0.096
	Solanum lycopersicum "Halay"	Tm1, Tm2/Tm2^2	-	-	0.069

Per species four plants were inoculated; the number of plants showing specific symptoms is indicated in brackets. Average OD_{405} values after 2 hours of substrate incubation were considered positive (red) when ≥ 2x the value of mock-inoculated controls, otherwise values were considered negative (green). -: no symptoms, ~: mild symptoms, C: chlorosis, CL: chlorotic lesions, LA: leaf abscission, NL: necrotic lesions, NR: necrotic rings, R: rugosity. *two plants showed local symptoms only, two other plants showed systemic symptoms only, DAS-ELISA was performed separately on pooled samples of young leaves of the two plants with local symptoms and systemic symptoms, respectively.

In contrast to the previous experiment (Table 3) in which all plants of the pepper cultivar

"Westlandse Grote Zoete" showed systemic symptoms, only two out of four inoculated plants

showed local symptoms, whereas the other two plants showed systemic symptoms only. However,

since cultivar "Westlandse Grote Zoete" does not harbor any tobamovirus resistance genes the

presence of the virus was expected in all plants, which was indeed confirmed by DAS-ELISA (Table 4).

Furthermore, all pepper accessions harboring a tobamovirus resistance gene/allele showed local but

no systemic symptoms (S2 Fig), and tested negative by DAS-ELISA (Table 4). Therefore, it appears

that each individual L^1, L^2, L^3 and L^4 gene/allele provided resistance to the novel tobamovirus. Overall,

these results showed that resistance against the novel virus is present in both the tomato and pepper

germplasm.

Finally, to gain more insight into the geographic distribution and the host range of the novel virus,

both symptomatic and asymptomatic fruits of *S. macrocarpon* and other *Solanaceae* imported into

the Netherlands in 2021, were subjected to Illumina sequencing analysis. This analysis included 19

samples from four countries, but no further material from Mexico: *Solanum betaceum* from

Colombia (2 samples), *S. macrocarpon* from Colombia (1 sample), *S. macrocarpon* from Surinam (10

samples), *S. melongena* from Malaysia (1 sample), and *Solanum torvum* from Thailand (5 samples).

The novel virus was not detected in any of these samples.

Discussion

This study assessed the potential risks of a newly discovered tobamovirus, tentatively named

"African eggplant-associated virus", from *S. macrocarpon*. The virus was easily transmitted

mechanically, and showed the ability to infect economically important crops, such as eggplant,

pepper, petunia, tomato, and other species including *Ammi majus, Nicotiana hesperis, N. occidentalis*

and *Solanum nigrum*. Overall, it was able to infect representatives of two out of six tested plant

families.

Sequence analysis indicated that the novel virus had a genome organization characteristic for

tobamoviruses, with less than 75% nucleotide identity to other tobamoviruses, which is significantly

lower than the tobamovirus species demarcation threshold of 90% nucleotide identity (18).

Therefore, it should be considered as a novel species within the genus *Tobamovirus*. Two genotypes

of the novel virus were present in the intercepted *S. macrocarpon* fruit. Although passages in *N.*

occidentalis "P1" of the novel virus did not change the nucleotide sequence of either genotype, a

decrease in relative frequency of genotype 1 was observed which led to genotype 2 taking over

completely in a subsequent passage to *S. macrocarpon.* This might be explained by adaptation to the

new host, as fitness cannot be maximized simultaneously in all potential hosts (19, 20).

Remarkably, an NCBI BLAST with the novel virus sequences resulted in a close hit with an alleged

bacteriophage isolate, which we expect to be incorrectly assigned and to actually belong to the same

novel tobamovirus. This "bacteriophage" sequence was part of a study that analyzed read data from

human metagenomic samples (21). Even though the query cover of the alignment with the novel

virus was only 60%, the high nucleotide identity with genotype 1 and genotype 2, 95.7% and 99.5%

respectively, suggests that the same virus was partially present in a human metagenomic sample.

Possibly, food or plant material contaminated with the virus had been consumed or touched by a

human.

A central paradigm in plant pathology is to demonstrate disease causation from a candidate

pathogen based on Koch's postulates. The novel virus was consistently detected in inoculated *S.*

macrocarpon plants, but we were not able to replicate the symptoms found on the original fruit.

Difficulties in demonstrating causal association are not uncommon in plant virology and the

limitations of Koch's postulates have been widely discussed (22). A potential explanation for the lack

of fruit symptoms could be the absence of genotype 1 in the inoculated *S. macrocarpon* plants.

Additionally, the development of symptoms after infection is influenced by environmental

circumstances and genotype of the plant, or a combination of these factors. Symptom variability as a

result of environmental or cultivar effects can be considered as a bottleneck for associating a novel

virus with a disease (23). Alternatively, the diffuse chlorotic mottling observed on the intercepted

fruit may have had a non-viral cause and we may have accidentally discovered an asymptomatic

infection.

A host range and pathotype study was conducted to assess the potential impact of the novel virus.

The infection of multiple solanaceous species was not unexpected, because the novel tobamovirus

was discovered in a host from this family. Additionally, *Ammi majus* from the *Apiaceae* family was

identified as a host. Although host ranges of tobamoviruses are usually quite narrow in nature, they

can be moderate to wide under experimental conditions (18). Wild species like *Ammi majus* and *S.*

nigrum that were infected experimentally might act as virus reservoirs, in which a virus can evolve

and spread to other plants and crops (24), for instance to one of the domesticated solanaceous

hosts. Such virus reservoirs complicate disease management. Vice versa, infected crops could pose a

risk to natural vegetation (25, 26).

For tomato four resistant accessions were identified which all harbored the combination of *Tm1* and

Tm2/*Tm2²*, whereas accessions with only one of these genes were tolerant, *i.e*. no symptoms were

observed while infection was detected by DAS-ELISA. This suggests that a combination of these *Tm*

genes is required for resistance. Most commercial tomato cultivars on the European market that are

grown in greenhouses harbor this combination of resistances. For open field crops, the use of these

resistances is less common, but tobamovirus infections are less likely to spread (27). Therefore, the

impact of the novel virus on tomato cultivation in Europe is likely to be low. It cannot be excluded

that another factor in the genetic background of the resistant accessions is involved. Allele dosage

might play a role as well, as it is not clear whether the resistances were homozygous or heterozygous

in the tested accessions. Even though *Tm2²* has historically been the most durable tobamovirus

resistance, it has previously been broken by ToBRFV (5) and partially broken by ToMMV (28). In a

screening of 160 tomato genotypes for ToBRFV resistance, only a single resistant genotype was

identified. A locus controlling tolerance mapped to chromosome 11 and was found to interact with a

locus on chromosome 2, at the *Tm1* region, to confer resistance (29). It would be interesting to

identify the genetics behind tolerance and resistance to African eggplant-associated virus in tomato

as well.

The breaking of *Tm2²* resistance by ToBRFV has been attributed to a replacement of the amino acid

cysteine (C68) by histidine (H67) in its movement protein. C68 is conserved among all other

Solanaceae-infecting tobamoviruses, and is involved in *Tm2²* activation and viral movement (30). The

novel tobamovirus contains C68, but surprisingly also infected a tomato accession harboring *Tm2²*.

The breaking of this resistance is therefore not due to the replacement of C68 as was shown for

ToBRFV.

For pepper, accessions harboring either of the L^1, L^2, L^3 or L^4 genes/alleles were resistant, whereas a

cultivar without any of these resistance genes was susceptible. Since most commercial pepper

varieties grown in the EPPO region harbor L resistance genes/alleles (27), the impact on pepper in

Europe is also likely to be low. However, it cannot be excluded that pepper plants harboring these

resistances are susceptible to African eggplant-associated virus under different conditions. For

example, an isolate of ToBRFV was able to infect pepper plants harboring $L^{1,3,4}$ at high temperatures,

whereas they were otherwise resistant (5).

Due to the advance of high throughput sequencing, novel virus species are being identified at a rapid

pace, but biological characterization is often lacking due to resource constraints (23, 31). However,

such experiments are crucial to determine the biological significance and predict the potential impact

of a novel virus. The current study identified a novel tobamovirus, tentatively named African

eggplant-associated virus. This virus experimentally infected multiple plant species, mainly

Solanaceae, thus being potentially harmful to several economically important crops including

eggplant, pepper, petunia and tomato. This study indicated that potato is not a host, while resistance

is present in both tomato and pepper germplasm. As most commercial tomato and pepper varieties

grown in European greenhouses harbor the relevant resistances, the risk of infection of these crops

in Europe is likely to be low. The non-resistant tomato accessions showed no symptoms under the

applied experimental conditions, indicating that the economic impact of the novel virus may be low

in this crop, even in the absence of resistance. However, the durability of these resistances against

the novel virus and their effectiveness under different environmental conditions remain unknown.

In conclusion, the present distribution of the novel virus and its prevalence are unknown. The virus

was not detected in a limited survey of solanaceous fruits imported into the Netherlands and did not

include further samples from Mexico. At present, we cannot predict the likelihood of transmission of

African eggplant-associated virus to other plant species. In general, hygiene measures and the use of

genetic resistance are the main strategies to prevent transmission of tobamoviruses. In this respect,

the observed latent infections in multiple experimental host species underline the merit of screening

symptomless plants for viral infections (32). Therefore, additional monitoring of *S. macrocarpon* and

other *Solanaceae* around the world, including non-symptomatic material, is needed to get a better

understanding of the incidence, distribution and phytosanitary risk of African eggplant-associated

virus and other potentially harmful viruses.

Acknowledgements

We would like to thank Tim Buysman (Kwaliteits-Controle-Bureau, The Netherlands) for

intercepting the *Solanum macrocarpon* fruit from which the novel virus was identified. We would

also like to thank our colleagues Robert Dees and Joanieke van Oorspronk for performing DAS-ELISA,

Jerom van Gemert, Christel de Krom and Carla Oplaat for symptom observations, Pier de Koning for

assistance with Illumina sequencing analysis, Dirk Jan van der Gaag for feedback on an early version

of the manuscript, and greenhouse staff for technical support.

References

1.   IPPC Secretariat. Strategic framework for the International Plant Protection Convention

(IPPC) 2020–2030. Protecting global plant resources and facilitating safe trade. Rome: FAO on behalf

of the Secretariat of the International Plant Protection Convention; 2021.

2.   Oboh G, Ekperigin M, Kazeem M. Nutritional and haemolytic properties of eggplants

(Solanum macrocarpon) leaves. Journal of Food Composition and Analysis. 2005;18(2-3):153-60.

3.   Arogundade O, Aderonmu OI, Matthew JO, Ayo-John EI. First Report of Tomato mosaic virus

Isolated From Solanum macrocarpon in Nigeria. Plant Disease. 2018;102(2):458-.

4.   Salem N, Mansour A, Ciuffo M, Falk BW, Turina M. A new tobamovirus infecting tomato crops

in Jordan. Arch Virol. 2016;161(2):503-6.

5.   Luria N, Smith E, Reingold V, Bekelman I, Lapidot M, Levin I, et al. A New Israeli Tobamovirus

Isolate Infects Tomato Plants Harboring Tm-22 Resistance Genes. PLoS One. 2017;12(1):e0170429.

6.   Van de Vossenberg BTLH, Dawood T, Woźny M, Botermans M. First Expansion of the Public

Tomato Brown Rugose Fruit Virus (ToBRFV) Nextstrain Build; Inclusion of New Genomic and

Epidemiological Data. PhytoFrontiers™. 2021;1(4):359-63.

7.   Ambros S, Martinez F, Ivars P, Hernandez C, de la Iglesia F, Elena SF. Molecular and biological

characterization of an isolate of Tomato mottle mosaic virus (ToMMV) infecting tomato and other

experimental hosts in eastern Spain. European Journal of Plant Pathology. 2017;149(2):261-8.

8.   Dombrovsky A, Tran-Nguyen LTT, Jones RAC. Cucumber green mottle mosaic virus: Rapidly

Increasing Global Distribution, Etiology, Epidemiology, and Management. Annu Rev Phytopathol.

2017;55:231-56.

9.   Broadbent L. Epidemiology and control of tomato mosaic virus. Annual review of

Phytopathology. 1976;14(1):75-96.

10.   Li Y, Wang Y, Hu J, Xiao L, Tan G, Lan P, et al. The complete genome sequence, occurrence

and host range of Tomato mottle mosaic virus Chinese isolate. Virology journal. 2017;14(1):1-9.

11.   Hall T. Resistance at the Tm-2 locus in the tomato to tomato mosaic virus. Euphytica.

1980;29(1):189-97.

12.   Tomita R, Sekine KT, Mizumoto H, Sakamoto M, Murai J, Kiba A, et al. Genetic basis for the

hierarchical interaction between Tobamovirus spp. and L resistance gene alleles from different

pepper species. Mol Plant Microbe Interact. 2011;24(1):108-17.

13.   Verhoeven JTJ, Roenhorst JW. Herbaceous test plants for the detection of quarantine viruses

of potato. EPPO Bulletin. 2000;30(3-4):463-7.

14.   Clark MF, Adams A. Characteristics of the microplate method of enzyme-linked

immunosorbent assay for the detection of plant viruses. Journal of general virology. 1977;34(3):475-

83.

15.   Hammond J, Adams IP, Fowkes AR, McGreig S, Botermans M, van Oorspronk JJ, et al.

Sequence analysis of 43-year old samples of Plantago lanceolata show that Plantain virus X is

synonymous with Actinidia virus X and is widely distributed. Plant Pathology. 2021;70(2):249-58.

16.   Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. Nature

methods. 2015;12(1):59-60.

17.   Ondov BD, Bergman NH, Phillippy AM. Interactive metagenomic visualization in a Web

browser. BMC bioinformatics. 2011;12(1):1-10.

18.   Adams MJ, Adkins S, Bragard C, Gilmer D, Li D, MacFarlane SA, et al. ICTV virus taxonomy

profile: Virgaviridae. The Journal of General Virology. 2017;98(8):1999.

19.   McLeish MJ, Fraile A, García-Arenal F. Evolution of plant–virus interactions: host range and

virus emergence. Current Opinion in Virology. 2019;34:50-5.

20.   García-Arenal F, Fraile A. Trade-offs in host range evolution of plant viruses. Plant Pathology.

2013;62:2-9.

21.   Tisza MJ, Buck CB. A catalog of tens of thousands of viruses from human metagenomes

reveals hidden associations with chronic diseases. Proceedings of the National Academy of Sciences.

2021;118(23).

22.   Fox A. Reconsidering causal association in plant virology. Plant Pathology. 2020;69(6):956-61.

23.   Hou W, Li S, Massart S. Is there a "biological desert" with the discovery of new plant viruses?

A retrospective analysis for new fruit tree viruses. Frontiers in Microbiology. 2020:2953.

24.   Power AG, Mitchell CE. Pathogen spillover in disease epidemics. The American Naturalist.

2004;164(S5):S79-S89.

25.   Bernardo P, Charles-Dominique T, Barakat M, Ortet P, Fernandez E, Filloux D, et al.

Geometagenomics illuminates the impact of agriculture on the distribution and prevalence of plant

viruses at the ecosystem scale. The ISME Journal. 2018;12(1):173-84.

26.   Malmstrom CM, Alexander HM. Effects of crop viruses on wild plants. Current Opinion in

Virology. 2016;19:30-6.

27.   EPPO. Pest risk analysis for tomato brown rugose fruit virus. EPPO, Paris. Available at

https://gd.eppo.int/taxon/TOBRFV/documents. 2020.

28.   Sui X, Zheng Y, Li R, Padmanabhan C, Tian T, Groth-Helms D, et al. Molecular and Biological

Characterization of Tomato mottle mosaic virus and Development of RT-PCR Detection. Plant

disease. 2017;101(5):704-11.

29.   Zinger A, Lapidot M, Harel A, Doron-Faigenboim A, Gelbart D, Levin I. Identification and

mapping of tomato genome loci controlling tolerance and resistance to tomato brown rugose fruit

virus. Plants. 2021;10(1):179.

30.   Hak H, Raanan H, Schwarz S, Sherman Y, Dinesh-Kumar SP, Spiegelman Z. Tm-2-2 resistance

targets a conserved cysteine essential for tobacco mosaic virus (TMV) movement. bioRxiv. 2022.

31.   Massart S, Candresse T, Gil J, Lacomme C, Predajna L, Ravnikar M, et al. A framework for the

evaluation of biosecurity, commercial, regulatory, and scientific impacts of plant viruses and viroids

identified by NGS technologies. Frontiers in microbiology. 2017;8:45.

32.   Ilyas R, Rohde MJ, Richert-Pöggeler KR, Ziebell H. To Be Seen or Not to Be Seen: Latent

Infection by Tobamoviruses. Plants. 2022;11(16):2166.

Supporting information

S1 Fig. Sequence depth of both virus genotypes in the original sample and after passages in *N. occidentalis "*P1" and *S. macrocarpon.*

S2 Fig. Symptoms on pepper accessions (*Capsicum spp.*) inoculated with the novel tobamovirus. a) *Capsicum annuum* "Westlandse Grote Zoete": chlorotic lesions, b) *C. annuum* "CGN19226": chlorotic lesions, leaf bulging, c) *C. chacoense* "CGN21477": necrotic lesions, leaf abscission, d) *C. chinense* "CGN21557": necrotic lesions, e) *C. frutescens* "CGN21546": chlorotic and necrotic lesions.

S3 Appendix. Raw DAS-ELISA values of Table 2-4.