**G Protein-Coupled Receptor Kinase-2 (GRK-2) controls exploration through neuropeptide**

**signaling in *Caenorhabditis elegans***

Kristen Davis[1,2,3], Christo Mitchell[4,+], Olivia Weissenfels[4,+], Jihong Bai[5], David M. Raizen[1,6],

Michael Ailion[4,*], and Irini Topalidou[4,5,*]

[1]Department of Neurology, Perelman School of Medicine, University of Pennsylvania,

Philadelphia, PA, USA

[2]Center for Excellence in Environmental Toxicology (CEET), Perelman School of Medicine,

University of Pennsylvania, Philadelphia, PA, USA

[3]Department of Neuroscience, Vickie and Jack Farber Institute of Neuroscience, Thomas

Jefferson University, Philadelphia, PA, USA

[4]Department of Biochemistry, University of Washington, Seattle, WA, USA

[5]Fred Hutchinson Cancer Center, Seattle, WA, USA

[6]Chronobiology and Sleep Institute, Perelman School of Medicine, University of Pennsylvania,

Philadelphia, PA, USA.

[+]These authors contributed equally to this work

[*]Corresponding authors

Email: etopalid@fredhutch.org

Email: mailion@uw.edu

Short title: GRK-2 controls exploration and movement quiescence

**Abstract**

Animals alter their behavior in manners that depend on environmental conditions as well

as their developmental and metabolic states. For example, *C. elegans* is quiescent during larval

molts or during conditions of satiety. By contrast, worms enter an exploration state when

removed from food. Sensory perception influences movement quiescence (defined as a lack of

body movement), as well as the expression of additional locomotor states in *C. elegans* that are

associated with increased or reduced locomotion activity, such as roaming (exploration

behavior) and dwelling (local search). Here we find that movement quiescence is enhanced,

and exploration behavior is reduced in G protein-coupled receptor kinase *grk-2* mutant animals.

*grk-2* was previously shown to act in chemosensation, locomotion, and egg-laying behaviors.

Using neuron-specific rescuing experiments, we show that GRK-2 acts in multiple ciliated

chemosensory neurons to control exploration behavior. *grk-2* acts in opposite ways from the

cGMP-dependent protein kinase gene *egl-4* to control movement quiescence and exploration

behavior. Analysis of mutants with defects in ciliated sensory neurons indicates that *grk-2* and

the cilium-structure mutants act in the same pathway to control exploration behavior. We find

that GRK-2 controls exploration behavior in an opposite manner from the neuropeptide receptor

NPR-1 and the neuropeptides FLP-1 and FLP-18. Finally, we show that secretion of the FLP-1

neuropeptide is negatively regulated by GRK-2 and that overexpression of FLP-1 reduces

exploration behavior. These results define neurons and molecular pathways that modulate

movement quiescence and exploration behavior.

**Author summary**

Many modulatory neurotransmitters affect behavior by binding to G protein-coupled

receptors (GPCRs) and initiating signals that modify neuronal activity. GPCRs are regulated by

G protein-coupled receptor kinases (GRKs). GRKs phosphorylate and promote the inactivation

of GPCRs. Here we identify GRK-2 as a regulator of distinct locomotor states in *C. elegans*. We

find that GRK-2 acts in olfactory sensory neurons to promote exploration and suppress

movement quiescence. Additionally, we show that GRK-2 acts in opposition to a neuropeptide

signaling pathway that acts in interneurons. Thus, this study demonstrates critical roles for

GRK-2 in regulating neuromodulatory signaling and locomotor behavior.

## Introduction

Animals modify their behaviors in response to changes in their environment or metabolic state. Sensory and neuromodulatory signals play a major role in adapting behavior to environmental or metabolic changes [1-6]. For example, animals experience periods of quiescence and arousal that depend on sensory and neuromodulatory cues [3,5,7-9]. Arousal is a state of increased responsiveness that can be associated with fear or starvation, whereas quiescence is characterized by decreased responsiveness and is associated with sleep and satiety. Even though sensory and neuromodulatory signals are recognized as major modulators of behavior, the precise mechanisms that mediate behavioral strategies under different inputs remain largely unknown. Here we describe how a locomotor behavior in the nematode *Caenorhabditis elegans* is altered through the action of the G protein-coupled receptor kinase-2 (GRK-2) in sensory neurons and neuromodulator signaling in interneurons.

Several examples of movement quiescence in response to changes in the external environment or the internal metabolic state of the animal have been reported in *C. elegans. C. elegans* enters an exploration state when it is removed from food  [10,11] and becomes quiescent during molting or under conditions of prolonged starvation, satiety, or cellular stress [3,12-17]. Sensory perception affects quiescence, as well as additional locomotor states in *C. elegans* that are characterized by increased or reduced locomotion activity, such as roaming/exploration and dwelling [2]. The animal covers a large area during roaming (exploring), while it restricts its activity to a small region during the dwelling state. The mechanism by which sensory perception modulates this locomotor strategy is unknown.

Quiescence, roaming, and dwelling are controlled by the function of the cGMP-dependent protein kinase (PKG) gene *egl-4* in sensory neurons*. egl-4* loss-of-function mutant animals have chemosensory defects [18] and attenuated sensory adaptation [19]. EGL-4 is a regulator of lethargus quiescence, a sleep-like behavior that takes place during molts [3], and of a satiety-induced sleep-like behavior [12]. EGL-4 also defines the time the animals spend

roaming and dwelling since *egl-4* loss of function mutant animals show increased exploration

[2,20]. Here, we describe a role for GRK-2 in the sensory neurons of *C. elegans* to regulate

movement quiescence and exploration in an opposite manner to EGL-4.

GRKs play key roles in attenuating the strength of neuromodulatory pathways by

phosphorylating G protein-coupled receptors (GPCRs) and promoting the inactivation of GPCR

signaling [21]. Vertebrates have seven GRKs that are classified into three subfamilies: the visual

GRKs (GRK1/7), the GRK2/3 subfamily, and the GRK4/5/6 subfamily [22]. *C. elegans,* which

lacks a visual system, has GRK-1 and GRK-2, which are orthologs of the mammalian GRK4/5/6

and GRK2/3 families, respectively [23,24]. *grk-2* is expressed broadly in the nervous system

and has a broad impact on animal physiology including chemosensation [23], gustatory

plasticity [25], egg-laying [26], crawling [27], swimming [27,28], and oxidative and heat stress

responses [29,30]. Elimination of *grk-2* function renders animals unable to respond to a wide

range of chemical stimuli, suggesting that GRK-2 plays a positive role in chemosensation [23].

Similarly, *grk-2* mutant animals show reduced crawling rates and are unable to swim, again

suggesting that GRK-2 plays a positive role in regulating crawling and swimming [27,28]. GRK-2

was proposed to promote crawling and swimming by negatively regulating the D2-like dopamine

receptor DOP-3 [27,28]. GRK-2 regulates chemosensation by acting in sensory neurons [23],

but it controls crawling and swimming by acting in premotor interneurons [27,28], suggesting

that GRK-2 acts in different neurons to modulate distinct behavioral outcomes.

Besides sensory perception, neuromodulation has also been shown to play a critical role

in defining arousal and quiescence as well as roaming and dwelling in *C. elegans*. For example,

mutants lacking the neuropeptide receptor NPR-1 have been used as a model for studying

arousal since both lethargus and movement quiescence are reduced in *npr-1* mutants [8,31-33].

Specifically, increased RMG interneuron activity in *npr-1* mutant animals during lethargus

induces secretion of the neuropeptide pigment dispersing factor PDF-1, resulting in increased

animal motility and reduced quiescence [31]. PDF-1 also positively regulates roaming, whereas

serotonin plays an opposite role by increasing dwelling [4]. Here we find that GRK-2 controls

exploration in an opposite manner to NPR-1 and the neuropeptides FLP-1 and FLP-18. We also

show that GRK-2 acts in different pathways than PDF and serotonergic signaling. Finally, we

demonstrate that secretion of FLP-1, a neuropeptide that regulates locomotion [34-37], depends

on GRK-2 and that overexpression of *flp-1* inhibits exploration behavior.

## Results

### grk-2 mutants have exploration and dispersal defects

Our previous work suggested that GRK-2 positively regulates crawling and swimming by negatively regulating the D2-like dopamine receptor DOP-3 [27,28]. *grk-2(gk268)* null mutant animals exhibit slow crawling and impaired swimming, and mutations in *dop-3* suppress both these phenotypes. *grk-2* mutant animals also have a decreased exploration behavior: they restrict their movements to a limited region of the bacterial lawn, whereas wild-type animals explore the entire lawn [27]. This exploration behavior defect could be due to decreased roaming, increased dwelling, or increased quiescence. We used simple radial locomotion and exploration assays ([4,38]; see also Methods) to quantify this phenotype by observing the radial distance that individual animals traveled over 1 h (Fig 1A) or the tracks that individual animals left on a lawn of *E. coli* over an 18-20 hr period (Fig 1D). We found that *grk-2(gk268)* mutant animals disperse and explore less compared to wild-type (Fig 1B and 1E). This phenotype is independent of sex since *grk-2* males also explore less than wild type males (Fig 1F). To examine whether the *grk-2* dispersal defect depends on the presence of food we performed the dispersal assay on plates without *E. coli.* We found that *grk-2(gk268)* mutant animals disperse less compared to wild type even in the absence of the bacteria lawn (Fig 1C), suggesting that the dispersal defect of *grk-2* mutant is independent of the presence or absence of food.

Although the *dop-3(vs106)* mutation suppresses the slow locomotion and swimming defects of *grk-2(gk268)* mutant animals [27,28], it did not significantly alter the dispersal or exploration behavior of *grk-2(gk268)* (Fig 1B,C and E), suggesting that GRK-2 affects the exploration behavior independently of DOP-3, and that the mechanism by which GRK-2 influences exploration behavior is different than its role in regulating crawling and swimming.

### GRK-2 acts in a different pathway than serotonergic or PDF signaling to affect exploration

Worm exploration behavior is regulated by serotonergic and PDF signaling: serotonin

inhibits exploration through the MOD-1 serotonin-gated chloride channel, while PDF-1 promotes

exploration through the Gαs-coupled PDF receptor PDFR-1 [4]. To examine whether GRK-2

acts in the PDF-1 pathway to affect the exploration behavior we compared animals that were

mutant for both *grk-2* and *pdfr-1* to single mutants. We found that *pdfr-1(ok3425)* mutant

animals display decreased exploration, as previously reported [4] and that the phenotype of the

double mutant *grk-2(gk268); pdfr-1(ok3425)* is stronger than the phenotype of the single

mutants (Fig S1A), suggesting that GRK-2 and PDFR-1 act in different pathways to control

exploration. Similarly, to examine whether GRK-2 acts in the serotonin pathway to control

exploration, we built double mutants between *grk-2* and *tph-1* mutants (*tph-1* encodes the rate-

limiting enzyme for serotonin synthesis). We verified that *tph-1(mg280)* mutant animals display

increased exploration [4] and further showed that *grk-2(gk268); tph-1(mg280)* double mutant

animals have an intermediate phenotype between that of either *grk-2* or *tph-1* single mutants

(Fig S1B), suggesting that *grk-2* and *tph-1* act in separate pathways to control exploration. Our

results imply that GRK-2 acts in a different pathway than serotonergic or PDF signaling to affect

the exploration behavior.

**GRK-2 functions in the ciliated neurons to affect the exploration behavior**

GRK-2 is expressed broadly in the nervous system [23] and acts in premotor

interneurons to control crawling and swimming [27,28]. To determine where GRK-2 acts to

affect the exploration behavior, we expressed the *grk-2* cDNA under the control of neuron-

specific promoters. Expression of *grk-2* cDNA under the pan-neuronal promoter (*rab-3p*) fully

rescued the *grk-2(gk268)* mutant exploration defect but expression in ventral cord acetylcholine

motor neurons alone (using the *acr-2* promoter) or in command interneurons alone (using the

*nmr-1* promoter) did not rescue this defect (Fig 2A). Expression of *grk-2* cDNA in the same set

of command interneurons rescued the crawling defects of *grk-2* mutant animals (Fig S2),

supporting the notion that the role that GRK-2 plays to affect the exploration behavior is different

from its role in crawling.

*grk-2* mutants fail to respond to a wide range of chemical stimuli, suggesting that GRK-2

is a positive regulator of chemosensation [23]. To test the hypothesis that the exploration defect

in *grk-2(gk268)* mutant animals is explained by its role in chemosensation, we expressed the

*grk-2* cDNA under the ciliated sensory neuron promoters *xbx-1p* and *osm-6p*. Expression of *grk-*

*2* cDNA in ciliated sensory neurons fully rescued the exploration defect of *grk-2* mutant animals

both in the presence and absence of food (Fig 2B,C). Our result that transgenic expression of

*grk-2* under *rab-3, xbx-1*, or *osm-6* heterologous promoters fully rescued the *grk-2* exploration

defect (Fig 2A-C) suggests that the precise level of GRK-2 is not critical to its function.

To further narrow down the site of action of GRK-2 in exploration behavior we expressed

the *grk-2* cDNA under the control of the *odr-3* promoter (*odr-3p*), which promotes expression in

olfactory and nociceptive neurons: AWA, AWB, AWC, ASH, and ADF [39]. Expression of *grk-2*

under the *odr-3* promoter partially rescued the exploration defect of *grk-2* mutant animals,

suggesting that *grk-2* acts in multiple ciliated neurons to affect exploration behavior (Fig 2D). An

operon GFP was included in the expression construct downstream of the 3'UTR allowing

confirmation of expression (see Methods). To confirm that *grk-2* is expressed in *odr-3*-

expressing neurons, we expressed the cDNA encoding GRK-2 protein fused to tagRFP (*grk-*

*2*::tagRFP) under the *grk-2* promoter as well as mNeon::NLS driven by the *odr-3* promoter (*odr-*

*3p*::mNeon::NLS). We confirmed that *grk-2* is indeed expressed in *odr-3*-expressing neurons

(shown with arrows in Fig 2G).

To dissect further the subset of chemosensory neurons where GRK-2 acts to control

exploration behavior we expressed the *grk-2* cDNA under the control of the *sra-6* promoter,

which is primarily expressed in the ASH and ASI chemosensory neurons (*sra-6p*; [40]).

Expression of *grk-2* under the *sra-6* promoter partially rescued the *grk-2* exploration defect,

suggesting that a part of GRK-2's role in exploration behavior is due to its action in the ASH or

ASI neurons (Fig 2E). Similarly, *grk-2* expression in the AWA, AWB, or AWC neurons (*odr-10p*:

expression in AWA [41]; *str-1p*: expression in AWB [42]; *str-2p*: expression in AWC-ON [43])

only slightly rescued the *grk-2* exploration defect (Fig 2F). Even *grk-2* expression under the

combined control of the *odr-1, str-1,* and *str-2* promoters only partially rescued the *grk-2*

exploration defect (Fig 2E), again suggesting that *grk-2* acts in more than a single type of

ciliated neuron to control exploration behavior.  To address the possibility that expression of *grk-*

*2* in any ciliated sensory neuron might rescue the exploration defect of *grk-2* mutant animals, we

expressed *grk-2* under the *gcy-8* promoter, which is solely expressed in the AFD thermosensory

neurons (*gcy-8p;* [44]). Expression of *grk-2* under the *gcy-8* promoter failed to rescue the *grk-2*

exploration defect (Fig 2H). Similarly, expression of *grk-2* under the dopaminergic neuron

promoter *dat-1* [45] failed to rescue the *grk-2* exploration defect (*dat-1p*; Fig 2I).

To examine whether GRK-2 plays a developmental role to affect exploration we placed

*grk-2* cDNA under the control of a heat shock-inducible promoter and introduced it in *grk-2*

mutant animals. Heat shock in adult animals partially but significantly restored the exploration

behavior of *grk-2* (Fig 2J). These results together suggest that GRK-2 has a functional role in

mature ciliated sensory neurons that affect the exploration behavior.

**Mutants in cilia-related genes have exploration defects similar to *grk-2***

Mutants with defects in sensory perception exhibit exploration defects on food like the

ones described here for *grk-2* [2]. Specifically, mutants with cilium-structure defects (such as

*che-2*) restrict their movements to a limited region of the bacterial lawn, while sensory signal

transduction mutants like *tax-2* and *tax-4* have a weaker defect than cilium-structure mutants

[2]. *grk-2* mutant animals are not defective in their ability to take up a fluorescent dye like the

cilium-structure mutants (Dye-filling assay; [23]), an assay indicative of the structural integrity of

the sensory cilia. To confirm that cilium structure mutants have exploration defects similar to

*grk-2*, we performed the exploration assay on *che-2(e1033)* and *daf-10(e1387)* mutant animals

in the presence or absence of food. We found that *che-2(e1033)* and *daf-10(e1387)* mutant

animals are exploration defective identical to *grk-2(gk268)* (Fig 3A-C). Additionally, the *che-*

*2(e1033)* and *daf-10(e1387)* mutations do not enhance the phenotype of *grk-2* mutant (Fig 3A-

C). The lack of an obvious enhancement is not due to a floor effect since *grk-2; pdfr-1* double

mutant animals have stronger exploration phenotype than the single mutants (Fig S1A). These

findings suggest that *che-2, daf-10*, and *grk-2* act in the same pathway to control exploration

(Fig 3A-C).

To further investigate which kinds of sensory perception affect exploration, we used the

exploration assay to test mutants with different sensory defects.  Most chemosensory and

osmosensory- defective mutants that possess defective cilia based on their Dye filling

phenotype [46] had defects in exploration (Fig 3D). Specifically, mutants with strong cilium-

structure defects (shown in blue in Fig 3D) had strong exploration defects and most mutants

with weak cilium defects (shown in green in Fig 3D) had weaker exploration defects. Other

chemosensory, osmosensory, and thermosensory defective mutants (*odr-1, odr-3, odr-8, odr-*

*10, osm-7, osm-10, tax-2, tax-4, ttx-1(p767)*), also had exploration defects but not as severe as

*grk-2*. Interestingly, chemosensory and osmosensory mutants *che-1, che-6, che-7, odr-2, odr-4,*

*odr-5, odr-7, osm-9,* and *osm-11*, as well as thermosensory *ttx-1(oy26* and *oy29)* mutants were

not significantly defective. Surprisingly, the *ocr-2* canonical null mutant *ak47* (VM396 strain)

showed an exploration defect that was not observed with the *ocr-2(ok1711)* allele (VC1233

strain).  Given that *ok1711* contains a deletion of at least three exons it is presumably a null

allele, suggesting that either the VC1233 *ocr-2(ok1711)* strain has a suppressor mutation that

rescues the exploration defect or that the VM396 *ocr-2(ak47)* strain has an exploration defect

due to a background mutation and not because of the *ocr-2(ak47)* deletion. To distinguish

between these possibilities, we outcrossed strains VC1233 and VM396 and tested the

exploration behavior of outcrossed strains BJH2762 *ocr-2(ok1711)* and BJH2763 *ocr-2(ak47)*.

We found that strain BJH2763 *ocr-2(ak47)* did not have an exploration defect, suggesting that

the *ocr-2(ak47)* mutation does not cause exploration defects and that strain VM396 has a

background mutation that makes the animals exploration defective (Fig S3A). Notably, oxygen-

sensing mutant *gcy-35* and mechanosensory-defective mutants *mec-3, mec-4,* and *glr-1* did not

show exploration defects. We conclude that ciliated neurons that are involved in

chemosensation and osmosensation play a significant role in the exploration behavior.

**GRK-2 controls movement quiescence in an opposite way to EGL-4**

*C. elegans* locomotory behavior in the presence of food alternates between two discrete

states called roaming and dwelling, where the animal either covers long distances or explores

locally, respectively [2,4]. The cGMP-dependent protein kinase EGL-4 acts in the sensory

neurons to promote dwelling [2]. *egl-4* loss of function mutant animals explore more than wild

type animals [2,4]. Also, *egl-4* mutations suppress the reduced exploration defect of *che-2*

mutants [2]. Thus, we examined whether *egl-4(n479)* mutant animals suppress the *grk-2*

exploration defect. We noticed that when the exploration assay was performed on 6 cm plates,

*egl-4* mutant animals didn't seem to explore significantly more than wild type animals, probably

due to a ceiling effect (Fig 4A). In contrast, when the assay was performed on 10 cm plates, *egl-*

*4* mutant animals explored significantly more than wild type animals (Fig 4B). In both cases, the

*egl-4(n479)* mutation significantly suppressed the *grk-2* exploration defect (Fig 4A,B).

Expression of EGL-4 in ciliated neurons reversed the *egl-4* suppression of the *grk-2* exploration

defect (Fig 4C), suggesting that GRK-2 and EGL-4 act in opposite ways in the ciliated neurons

to affect the exploration behavior.

EGL-4 also functions in sensory neurons to promote a *C. elegans* sleep state during

lethargus (lethargus quiescence; [3]). The described role of EGL-4 in quiescence prompted us

to examine whether the exploration defect of *grk-2* is the result of movement quiescence. We

measured activity and quiescence in *C. elegans* larval stage 4 (L4) animals by imaging

individual animals cultivated in a WorMotel, which is a micro-fabricated device with multiple

agar-filled wells [47,48]. Single L4 animals were placed in WorMotel, imaged for 8 h, and their

quiescence and activity were quantified. *grk-2(gk268)* mutant animals showed reduced body

movement activity and increased quiescence in this 8 h period and the *egl-4(n479)* mutation

suppressed these phenotypes (Fig 4D-F). We conclude that *grk-2* mutant animals have

enhanced movement quiescence that is suppressed by *egl-4*, indicating that *grk-2* and *egl-4* act

in opposite directions to regulate movement quiescence. These results also suggest that part of

the exploration defect of *grk-2* mutant might be the result of increased movement quiescence.

In addition to cGMP signaling through EGL-4, cAMP signaling pathways act through the

salt-inducible kinase KIN-29 in sensory neurons to regulate sensory behaviors [49]. *kin-29* plays

an important role in the metabolic regulation of sleep [50] and *kin-29* mutants, like *egl-4*

mutants, have reduced sleep. However, in contrast to *egl-4* mutant animals, *kin-29(oy38)*

mutant animals did not explore more than wild-type animals and the *kin-29(oy38)* mutation did

not rescue the exploration defect of *grk-2(gk268)* mutant (Fig S3B). We conclude that KIN-29

does not play a role in the GRK-2 pathway to control exploration and movement quiescence.

**APTF-1 promotes GRK-2-modulated movement quiescence and exploration**

To further explore the role of sleep in the reduced exploration of *grk-2* mutant, we

manipulated sleep in *grk-2* mutant animals. Sleep behavior is controlled by the AP2 transcription

factor APTF-1, which is required for the development of the sleep-promoting RIS interneuron

[51]. Although the exploration behavior of *aptf-1(gk794)* mutant animals was similar to that of

wild type animals, the *aptf-1* mutation partially suppressed the decreased exploration phenotype

of *grk-2* mutant animals (Fig 5A). Similarly, *aptf-1* mutant partially suppressed both the

enhanced quiescence and reduced activity phenotypes of *grk-2* mutant (Fig 5B-D). We

conclude that increased sleep is partially responsible for the reduced exploration phenotype of

*grk-2* mutant animals.

APTF-1 promotes sleep by turning on the expression of the sleep-inducing neuropeptide

FLP-11 in the RIS neuron. At sleep onset, increases in RIS calcium activity cause the release of

FLP-11, which induces quiescence [9]. *flp-11(tm2706)* mutant did not significantly affect the

exploration or quiescence phenotypes of *grk-2* mutant (Fig S4A-D), suggesting that GRK-2 acts

independently of FLP-11 in regulating quiescence, and that the role of APTF-1 in adult

movement quiescence is partially independent of FLP-11. Alternatively, since the effect of *flp-11*

on sleep is weaker than that of *aptf-1* [9], our assays might not be sensitive enough to detect a

phenotype for *flp-11*.

*C. elegans* displays an additional sleep-like quiescent behavior which occurs in

response to exposure to cellular stressors (stress-induced quiescence). Mutants lacking the

transcription factor CEH-17 lack a functional ALA neuron, a neuron which promotes stress-

induced quiescenc [13,14,52,53]. In contrast to the *aptf-1* mutation, the *ceh-17(np1)* mutation

did not affect the exploration or quiescence phenotypes of *grk-2* (Fig S4E-H), suggesting that

GRK-2 acts independently of the ALA neuron to control movement quiescence.

**NPR-1 negatively controls GRK-2- modulated exploration behavior**

The *C. elegans* neuropeptide receptor NPR-1 negatively regulates arousal by inhibiting

the activity of a sensory circuit that is connected by gap junctions to the RMG interneurons [54].

*npr-1* loss-of-function mutant animals have increased locomotion both in lethargus and in the

adult stage due to heightened activity of the RMG circuit [8,30-32,54]. *npr-1(ky13)* mutant

animals explored more than wild type animals and *npr-1(ky13)* suppressed the exploration

defect of *grk-2(gk268)* (Fig 6A,B). Moreover, *npr-1(ky13)* also suppressed the *grk-2* dispersal

defect in the absence of food (Fig 6C). These results suggest that *npr-1* and *grk-2* act in

opposite ways to affect the exploration behavior.

The FMRFamide-related neuropeptides (FaRPs) FLP-18 and FLP-21 are known NPR-1

ligands [55,56]. While *flp-18(db99)* and *flp-21(ok889)* single mutants did not have an exploration

defect, a *flp-18* mutation partially suppressed the defective exploration phenotype of *grk-2* (Fig

6D). The *flp-21* mutation did not affect the *grk-2* exploration defect and *flp-21; flp-18* mutants

were not stronger suppressors than *flp-18* mutants, suggesting that the *flp-21(ok889)* mutation

does not suppress the *grk-2* exploration defect (Fig 6E). The NPR-1 ligand FLP-18 only having

a partial effect suggests that it is not the only ligand for NPR-1 in this pathway. Alternatively,

NPR-1 may have ligand-independent activity.

NPR-1 is expressed in several neurons including neurons of the RMG circuit (ASH,

URX, RMG) [57] and expression in the RMG circuit completely rescues the increased

locomotion of *npr-1* mutants during lethargus [8,31]. Moreover, expression solely in the RMG

interneuron rescues the enhanced locomotion speed of *npr-1* mutants [54] while NPR-1 acts in

the ASH sensory neurons to inhibit sensory responses [8]. To examine whether ASH-specific

expression of *npr-1* is sufficient to reverse the *npr-1* suppression of the exploration defect of *grk-*

*2* mutants we expressed the *npr-1* cDNA under the control of the ASH (and ASI) - specific *sra-6*

promoter (*sra-6p*; [40]). Expression of *npr-1* under the *sra-6* promoter did not reverse the *npr-*

*1(ky13)* suppression of the exploration defect of *grk-2* (Fig 6F).

We next addressed whether RMG- specific expression of *npr-1* is sufficient to reverse

the *npr-1-* suppression of the exploration defect of *grk-2*. Since RMG- specific promoters are not

known, we used a described intersection strategy to drive *npr-1* expression only in the RMG

neuron [54]. In this approach, we used the *ncs-1* promoter to drive the Cre recombinase and the

*flp-21* promoter to drive *npr-1* preceded by a floxed STOP site. [54]. Expression of *npr-1* by this

method in *grk-2(gk268); npr-1(ky13)* mutant animals fully reversed the *npr-1* suppression of the

exploration defect of *grk-2* (Fig 6F). *flp-21* and *ncs-1* are coexpressed in several neurons

(including the RMG neurons), but of these neurons, *npr-1* is present only in ASE, ASG, and

RMG. Since ASE and ASG are gustatory neurons that have not been implicated in NPR-1

function in locomotion, our experiments suggest that NPR-1 functions in RMG to negatively

control the GRK-2 pathway that controls the exploration behavior.

**FLP-1 negatively controls GRK-2- mediated exploration behavior**

In a literature search for additional neuropeptides that could play a role in GRK-2-

modulated exploration, we came across FLP-1. *flp-1(ok2811)* null mutant animals are

hyperactive and have a loopy locomotion posture characterized by deep body bends [34-37].

Interestingly, although *flp-1(ok2811)* mutant animals explored as well as WT animals in the

presence of food, the *flp-1(ok2811)* mutation significantly suppressed the *grk-2(gk268)*

exploration defect (Fig 7A). Moreover, *grk-2(gk268); flp-1(ok2811); flp-18(db99)* triple mutant

animals behaved like the *grk-2(gk268); flp-1(ok2811)* double mutant, suggesting that FLP-1 and

FLP-18 act in the same pathway to negatively modulate GRK-2-mediated exploration behavior

(Fig 7B). Expression of FLP-1 under its own promoter and under the AVK neuron-specific *twk-*

*47* promoter reversed the *flp-1* suppression of the *grk-2* exploration phenotype, suggesting that

expression of FLP-1 in AVK interneurons is sufficient to restore its function (Fig 7C). To assess

whether *npr-1* and *flp-1* act in the same pathway, we built the triple mutant *grk-2; flp-1; npr-1*

and compared its phenotype with that of the *grk-2; npr-1* double mutant. We found that *grk-2;*

*flp-1; npr-1* behaves like *grk-2; npr-1* mutant animals, suggesting that either NPR-1 and FLP-1

act in the same pathway or that we cannot detect further suppression due to a ceiling effect (Fig

7D,E). Previous reports suggest that NPR-6 and FRPR-7 are low- and high-affinity FLP-1

receptors, respectively [36]. We tested the exploration behavior of *grk-2(gk268); npr-6(tm1497)*

and *grk-2(gk268); frpr-7(gk463846)* double mutants. *npr-6(tm1497)* did not affect the exploration

behavior of *grk-2(gk268),* but *frpr-7(gk463846)* partially but significantly suppressed the

exploration defect of *grk-2(gk268)* (Fig 7F)*.* These results suggest that *frpr-7* but not *npr-6* could

be playing a role in the exploration behavior.

**GRK-2 negatively regulates FLP-1 secretion from the AVK neuron**

To examine the possibility of a functional connection between GRK-2 and FLP-1, we

assessed whether GRK-2 affects the levels of FLP-1 secretion using the coelomocyte uptake

assay [58-60]. Proteins that are secreted into the *C. elegans* body cavity are endocytosed by

scavenger cells named coelomocytes (Fig 8A). We expressed the fusion protein FLP-

1::mCherry in AVK neurons of *flp-1* single mutant animals and of *grk-2; flp-1* double mutant

animals and quantified the intensity of fluorescence in coelomocytes as a readout of secreted

FLP-1::mCherry (see Methods; [36]). Previous work confirmed that FLP-1::mCherry follows the

dense core vesicle route for neuropeptide secretion and used the FLP-1::mCherry secretion

assay as a way of evaluating FLP-1 secretion [36]. Notably, FLP-1::mCherry expression in *grk-*

*2; flp-1* reversed the suppression of the exploration defect of *grk-2* by *flp-1,* suggesting that FLP-

1 in the FLP-1::mCherry reporter retains its function in exploration (Fig S5). *grk-2; flp-1* mutants

expressing FLP-1::mCherry had significantly increased levels of mCherry in coelomocytes in

comparison to *flp-1* mutant animals (Fig 8A,B). To examine whether background fluorescence in

the coelomocytes of *grk-2; flp-1* mutants also changed, together with FLP-1::mCherry we co-

expressed a reporter expressing GFP driven by the coelomocyte-specific promoter *unc-122* and

tested the levels of GFP fluorescence in the coelomocytes of *flp-1* and *grk-2; flp-1* mutants. We

found that GFP fluorescence remained unchanged in *grk-2; flp-1* mutants suggesting that the

increase in FLP-1::mCherry fluorescence in *grk-2; flp-1* coelomocytes is specific to FLP-

1::mCherry (Fig 8C). These results suggest that GRK-2 affects the level of secreted FLP-1,

which could explain at least in part the exploration defect observed in mutants lacking GRK-2.

Consistent with this idea, multi-copy expression of FLP-1::mCherry under the *flp-1* promoter

reduced the exploration behavior of wild type animals (Fig 8D). These results altogether indicate

that GRK-2 negatively regulates the amount of FLP-1 that is secreted by AVK neurons, resulting

in positive regulation of the exploration behavior.

## Discussion

Locomotion is the result of dynamic interactions between a genetically determined central program and feedback mechanisms that modulate the activity of the central program [61]. Feedback is mainly of sensory and neuromodulatory nature and it plays a major role in adapting the pattern of locomotion to environmental or metabolic changes [1]. Here we have demonstrated that locomotor behavioral strategies in the nematode *Caenorhabditis elegans* are altered through the action of the G protein-coupled receptor kinase-2 (GRK-2) in ciliated sensory neurons and neuropeptide signaling in interneurons. We propose that GRK-2 acts in ciliated sensory neurons to control the exploration behavior and the arousal state of the animal by controlling the level of secretion of the neuropeptide FLP-1.

### GRK-2 signaling in ciliated sensory neurons controls exploration

The sensory system influences the expression of different locomotor states in *C. elegans*, like roaming and dwelling [2]. Roaming is a behavioral exploration strategy where the animal covers long distances, while dwelling is a local search behavior strategy where animals restrict their activity to a small region. Previous studies have shown that mutants with diminished sensory responses to soluble and volatile chemicals are defective in the time they spend roaming or dwelling, suggesting that sensory perception is critical for this behavior [2]. Specifically, a class of mutants that lack a normal sensory cilium structure, including *che-2* [62,63], are unable to explore long distances. In this study we show that *grk-2* mutant animals have similarly severe phenotypes as the cilium structure mutants in that they also show reduced exploration. Our findings suggest that the positive role of GRK-2 in exploration behavior is due to its positive role in chemosensation. We show that expression of GRK-2 in the ciliated sensory neurons of *grk-2* mutant animals rescues their exploration defect, suggesting that GRK-2 acts specifically in these neurons to affect exploration. Our effort to narrow down the specific neurons where GRK-2 acts to mediate this behavior revealed that GRK-2 acts in multiple

ciliated neurons to control exploration. Interestingly, although *grk-2* mutant animals have strong

chemosensory defects, they do not have severe cilia structural defects like the cilium structure

mutants [23,64]. Also, expression of GRK-2 in adult *grk-2* mutant animals can rescue the *grk-2*

exploration defect suggesting that GRK-2 plays a non-developmental role in ciliated neurons to

promote exploration. Our study of mutants with defects in different kinds of sensory perception

showed that the activity of the ciliated neurons that are involved in chemosensation and

osmosensation play important roles in the exploration behavior, but neurons involved in

mechanosensation or oxygen-sensation do not. These results together suggest that impaired

sensory neuron function per se can reduce exploration.

Our results show that *grk-2* and the cilium structure mutant *che-2* act in the same

genetic pathway to control exploration, again suggesting that impaired ciliated sensory neuron

function disrupts exploration. In agreement, we found that mutations in the gene encoding the

cGMP-dependent protein kinase EGL-4 that suppress the altered exploration behavior of *che-2*

mutants [2], also suppress the exploration defect of *grk-2* mutant animals. Given that expression

of EGL-4 in the ciliated neurons reverses this suppression, we propose that both GRK-2 and

EGL-4 act in the same ciliated neurons to control exploration but in opposite ways. Given that

both GRK-2 and EGL-4 act in multiple ciliated neurons to control exploration behavior and that

other chemosensory and osmosensory mutants have similar exploration defects as *grk-2*, we

support a model in which GRK-2 acts in ciliated neurons to positively control sensory perception

and affect the exploration state of the animal by regulating EGL-4. We propose that in *grk-2* and

cilium structure mutants, impaired sensory perception leads to an increase in EGL-4 activity and

decreased exploration. A similar model was supported by previous studies, which showed that

EGL-4 in sensory neurons controls body size and locomotory states [2].

In previous work we have shown that GRK-2 regulates crawling and swimming [27,28]

by negatively regulating the D2-like dopamine receptor DOP-3 in premotor interneurons (Fig

9A). In the case of chemosensation and exploration, the target(s) of GRK-2, which are located

in chemosensory neurons and not interneurons, are currently unknown (Fig 9B). Similarly, we

do not yet know the identity of the phosphorylation target(s) for EGL-4.

**GRK-2 controls movement quiescence**

Animals undergo periods of quiescence and arousal in response to changes in their

environment or metabolic state. For example, *C. elegans* increases its exploration behavior

when it is removed from food [10,11] and shows quiescence during larval molting (lethargus)

and under starvation, satiety, and stress [3,12-17]. The cGMP-dependent protein kinase EGL-4

acts in sensory neurons to promote *C. elegans* sleep where the animals become quiescent

during lethargus [3]. This result, together with the observed opposite functionality between EGL-

4 and GRK-2 in ciliated neurons prompted us to examine whether *grk-2* mutants have

movement quiescence defects. Indeed, we found that *grk-2* mutant animals have enhanced

movement quiescence, which is suppressed by *egl-4*. This result suggests that GRK-2 and

EGL-4 act antagonistically to control movement quiescence outside lethargus. We propose that

GRK-2 acts in ciliated neurons to control sensory perception and affect movement quiescence

by negatively regulating EGL-4. Our findings imply that the exploration defect of *grk-2* and other

chemosensory mutants besides *grk-2* might be due in part to enhanced movement quiescence.

This series of data also suggests that *C. elegans* locomotion strategies that have been

described as exploration, roaming, and arousal might reflect the same behavior.

In addition to EGL-4, the AP2 transcription factor APTF-1 also plays an important role in

regulating sleep, largely by regulating expression of the sleep-inducing neuropeptide FLP-11 in

the RIS interneuron [51]. Indeed, we found that *aptf-1* mutants partially suppress the decreased

exploration phenotype, enhanced quiescence, and reduced activity of *grk-2*. This result

suggests that increased sleep is partially responsible for the reduced exploration phenotype of

*grk-2* mutant animals. Surprisingly, we found that *flp-11* mutations do not suppress the

enhanced quiescence of *grk-2* mutant animals. One possible explanation is that APTF-1 acts

through additional targets besides FLP-11, which is also supported by previous studies [9].

Alternatively, FLP-11 might be playing a smaller role than APTF-1 in adult quiescence and our

assays might not be sensitive enough to detect it.

**Neuromodulation through FLP-1 negatively regulates exploration**

Mutants lacking the neuropeptide receptor NPR-1 have been described as a model for

arousal both during lethargus and in adults [8,31-33]. NPR-1 is proposed to act in the RMG

interneuron to inhibit the activity of a sensory circuit that is connected to RMG via gap junctions

(54). In *npr-1* mutants, enhanced sensory activity causes aroused locomotion [8,31-33,65]. Our

results show that *npr-1* mutants suppress the exploration defects of *grk-2* mutant animals, and

that NPR-1 acts in RMG neurons to mediate this effect. The fact that *npr-1* is epistatic to *grk-2*

suggests that GRK-2 negatively regulates NPR-1 – in a *grk-2* mutant, increase in NPR-1 activity

inhibits exploration (Fig 9B). Since GRK-2 acts in ciliated neurons and NPR-1 acts in RMG

neurons to control exploration, this suggests that GRK-2 activity regulates NPR-1 in a cell-

nonautonomous way. Of the reported NPR-1 ligands FLP-18 and FLP-21, we found that only

FLP-18 acts in the GRK-2 pathway that mediates exploration, since *flp-18* mutants partially

suppress the exploration defect of *grk-2* [55,56]. Interestingly, we also found that mutants in the

gene encoding the neuropeptide FLP-1 partially suppress the *grk-2* exploration defect and that

FLP-1 acts in the same genetic pathway as FLP-18.  FLP-1 has not been reported to be a

ligand for *C. elegans* NPR-1, but it was shown to activate the *Girardia tigrina* neuropeptide

receptor 1 GtNPR-1 when expressed in Chinese hamster ovary cells [66]. This finding together

with our genetic observations could suggest that FLP-1 directly acts through NPR-1 to affect

exploration. Alternatively, and since *flp-1* and *flp-18* mutants do not act in an additive way to

suppress the *grk-2* exploration defect, FLP-1 could be acting in the GRK-2 pathway through a

different receptor than NPR-1. The partial suppression of the *grk-2* exploration defect by *frpr-7*,

suggests that the high-affinity FLP-1 receptor FRPR-7 could be playing a role.

Notably, Oranth et al (2018) [36] suggested that after >10 min off food, AVK releases

FLP-1 peptides to promote dispersal. Our genetic experiments show that mutants in *flp-1*

suppress the exploration defect of *grk-2* mutant animals and that overexpression of FLP-1

reduces exploration in the presence of food. This apparent discrepancy might be attributed to

different FLP-1 functionality in the presence or absence of food or when chemosensation is

impaired. Our genetic results together with our finding that GRK-2 modulates the secretion

levels of FLP-1 suggest that GRK-2 negatively regulates FLP-1 secretion, either directly or

indirectly (Fig 9B). Since GRK-2 acts in ciliated neurons to affect exploration and GRK-2

modulates FLP-1 secretion from AVK neurons we propose that GRK-2 acts in ciliated neurons

to mediate FLP-1 secretion from AVKs in a cell-nonautonomous way.  Given that there is no

known synaptic input from chemosensory neurons to AVKs [67] we suggest that GRK-2-driven

regulation of sensory neuron activity affects a neuromodulatory pathway that drives FLP-1

secretion from AVKs (Fig 9B). AVKs were previously reported to play important roles in

regulating locomotion by modulating motor neuron activity [36]. Alternatively, GRK-2 action in

ciliated neurons may lead to modulation of the neuronal circuit downstream of the ciliated

neurons that eventually affects secretion of FLP-1. According to our model, GRK-2 activity

positively regulates chemosensory neuron activity, which in turn inhibits EGL-4 activity,

modulating exploration and the arousal state of *C. elegans* through neuromodulatory signaling

acting in interneurons (Fig 9B). Thus, an active chemosensory system can guide *C. elegans*

towards exploration, while a defective chemosensory system inhibits dispersal and puts the

animal in a state of quiescence.

**Methods**

**Strains**

Strains were maintained at room temperature or 20° on the OP50 strain of *E. coli* [68].

The Supplementary Information contains a list of strains used, including their full genotypes

(Table S1; List of strains).

**Constructs and transgenes**

The Supplementary Information contains a list of constructs used (Table S2; List of

plasmids). All constructs made in this study were constructed using the Gateway system

(Invitrogen): a promoter region, a gene region (cDNA) or fluorescent tag, and a N- or C-terminal

3'UTR or fluorescent tag fused to a 3'UTR were cloned into the destination vector pCFJ150 [69].

For the cell-specific rescuing experiments, an operon GFP was included in the expression

constructs downstream of the 3'UTR [70]. This resulted in expression of untagged protein, but

also allowed for confirmation of proper promoter expression by monitoring GFP expression.

Extrachromosomal arrays were made by standard injection and transformation methods

[71]. We injected 5-10 ng/ul of the expression plasmid for *grk-2* and *flp-1* rescue experiments

and 20 ng/ul of the plasmids for other experiments. For the *yakEx261* FLP-1 overexpression

experiment in Fig 8D we injected 60 ng/ul. We injected 50 ng/ul of the *unc-122*::GFP co-

injection marker, 1 ng/ul of *myo-2p*::mCherry, 5 ng/ul of *myo-3p*::mCherry, and 5 ng/ul of *rab-*

*3p*::mCherry. We isolated two to three independent lines. The results shown in the figures are

for two of the lines tested.

**Exploration assays**

The exploration assay is an established assay that measures the ability of the animals to

explore (wander) during long periods of time [4]. Exploration assays were performed on 6 or 10

cm nematode growth medium (NGM) plates uniformly seeded with 200 or 500 ul of the OP50

strain of *E. coli*, respectively, and spread with a glass spreader. 10 cm plates were used in the

case of mutants that explore more than wild type animals (e.g. *egl-4, npr-1*) in order to avoid a

ceiling effect because wild type animals usually explore most of the area of a 6 cm plate in a

20h period. Bacteria were grown on the NGM agar surface at room temperature for

approximately 48 h and the plates were stored at $4°C$ until needed. On the day of the

experiment, individual first-day adult animals were transferred to the center of each plate and

were allowed to explore for 18-20 h. Then the plates were superimposed on a 50- or 150-

square grid and we counted the number of squares entered by the worm tracks. Approximately

10-40 animals were tested per strain. Every strain was tested in two or three independent trials.

The number "n" in figure legends indicates the total number of animals sampled (animals from

two or three different trials are combined).

**Dispersal assays**

The dispersal (or radial locomotion) assay is an established assay that measures the

distance animals travel from the middle of the plate in a period [38]. Since the dispersal assay

measures distance, it is only performed for short periods of time (1h on food or 20 min on plates

without food). Dispersal assays were performed on 10 cm nematode growth medium (NGM)

plates that were uniformly seeded with 500 µl of the OP50 strain of *E. coli* and spread with a

glass spreader. Bacteria were grown at room temperature for approximately 48 h and the plates

were stored at $4°C$ until needed. Approximately 20 first day adult animals were transferred to the

center of a plate and were allowed to explore for 1 h. Then the positions of the worms were

marked and the distances of the worms from the starting point were measured. Dispersal

assays without food were performed on NGM plates that were not spread with OP50 and the

animals were allowed to move freely for 20 min. Animals were first transferred to an

intermediate plate without food before being transferred to the assay plate. Approximately 20-60

animals were tested per strain. Every strain was tested in two or three independent trials. The

number "n" in figure legends indicates the total number of animals sampled (animals from two to

three different trials are combined).

**Heat shock assays**

Animals were raised to young adulthood at room temperature (23°C), then shifted to

34°C for 2 hr. They were allowed to recover at room temperature (23°C) for 4 hr and tested.

**Movement quiescence assays**

Movement quiescence assays were performed as previously described [48]. Briefly,

animals were selected as fourth larval stage L4s the day prior to the assay. They were loaded

individually onto the surface of NGM agar-filled wells on a PDMS WorMotel chip. The agar

surface in each well was covered in a thin lawn of DA837 *E. coli*. The loaded chip was placed in

a 10 cm petri dish lined with damp kimwipes to prevent desiccation and placed under a camera.

Images were acquired every 10 seconds for eight hours. The images were then analyzed using

a custom written program in Matlab. For more details on this method see [48]. 8-12 animals

were tested per condition and compared to 8-12 control animals imaged in the same WorMotel.

The "n" in figure legends indicates the total number of animals sampled.

**Body bend locomotion assays**

First-day adults were picked to a three-day-old lawn of OP50 and stimulated by poking

the tail of the animal with a worm pick. Body bends were then immediately counted for one

minute. A body bend was defined as the movement of the worm from maximum to minimum

amplitude of the sine wave. Approximately 10 animals were tested per strain. Every strain was

tested on two independent trials. The number "n" in figure legends indicates the total number of

animals sampled (animals from the different trials are combined).

**Imaging**

For fluorescence imaging, first-day adult animals were mounted on 2% agarose pads

and anesthetized with 50 mM sodium azide for ten minutes before placing the cover slip. The

animals were imaged using an Olympus FLUOVIEW FV1200 confocal microscope with a 60×

UPlanSApo oil objective (NA = 1.35) or a NikonTi2-E Crest X-light V2 spinning disk confocal

with a 60x oil objective (NA = 1.4).

**Coelomocyte assays**

For coelomocyte imaging we used FLP-1::mCherry (a gift from Alexander Gottschalk;

[36]). We imaged the anterior coelomocytes in day 1 adults using a NikonTi2-E Crest X-light V2

spinning disk confocal with a 60x oil 1.4 NA objective. Maximum intensity projections of image

stacks were obtained using FluoView software (Nikon). For quantification, the maximum

fluorescence intensity was calculated using Fiji and the background fluorescence intensity was

subtracted to obtain coelomocyte fluorescence.

**Statistics**

Statistical analysis and graphing were carried out using Prism 8 or Prism 9 (GraphPad

Software). Normally distributed data sets requiring multiple comparisons were analyzed by a

one-way ANOVA or by a one-way ANOVA followed by a Bonferroni test. Normally distributed

pairwise data comparisons were analyzed by two-tailed unpaired t tests. $p < 0.05$ (following

Bonferroni correction, when appropriate) was considered to be statistically significant (*$p < 0.05$,

**$p < 0.01$, ***$p < 0.001$, ****$p<0.0001$). Data are presented as the mean ± standard error

(SEM).

**Raw data**

Table S3 contains all the data points used for the main and supplementary figures.

**Acknowledgements**

We dedicate this paper to the memory of Giannis Topalidis, whose support was

instrumental for the completion of this manuscript. We thank Alexander Gottschalk for the FLP-

1::mCherry plasmid, Ithai Rabinowitch for the *str-1, str-2,* and *odr-10* promoter plasmids, and

Cori Bargmann for the pEM01 and pEM03 plasmids. Some strains were provided by the CGC,

which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440). This

work was funded by R01 NS109476 to MA and JB, R01 GM121481 to MA, R01 NS122779 and

R01 NS107969 to DMR. KT was supported by a T32Act ES019851 Training grant (PI: Trever

Penning).

**Author contributions**

637      KD and IT conceived the experiments. KD and IT performed the experiments and

638 analyzed the data. CM and OP performed the experiments described in Fig 3D. MA, DMR, and

639 JB assisted with providing ideas. IT wrote the paper with help from MA, DMR, and JB.

**References**

1.    MacKay-Lyons M. Central pattern generation of locomotion: a review of the evidence. Phys

Ther. 2002 Jan;82(1):69–83.

2.    Fujiwara M, Sengupta P, McIntire SL. Regulation of body size and behavioral state of *C.*

*elegans* by sensory perception and the EGL-4 cGMP-dependent protein kinase. Neuron.

2002 Dec 19;36(6):1091–102.

3.    Raizen DM, Zimmerman JE, Maycock MH, Ta UD, You Y jai, Sundaram MV, et al.

Lethargus is a *Caenorhabditis elegans* sleep-like state. Nature. 2008 Jan

31;451(7178):569–72.

4.    Flavell SW, Pokala N, Macosko EZ, Albrecht DR, Larsch J, Bargmann CI. Serotonin and

the neuropeptide PDF initiate and extend opposing behavioral states in *C. elegans*. Cell.

2013 Aug 29;154(5):1023–35.

5.    Flavell SW, Raizen DM, You YJ. Behavioral States. Genetics. 2020 Oct;216(2):315–32.

6.    Flavell SW, Gordus A. Dynamic functional connectivity in the static connectome of

*Caenorhabditis elegans*. Curr Opin Neurobiol. 2022 Feb 17;73:102515.

7.    Taghert PH, Nitabach MN. Peptide neuromodulation in invertebrate model systems.

Neuron. 2012 Oct 4;76(1):82–97.

8.    Choi S, Taylor KP, Chatzigeorgiou M, Hu Z, Schafer WR, Kaplan JM. Sensory Neurons

Arouse *C. elegans* Locomotion via Both Glutamate and Neuropeptide Release. PLoS

Genet. 2015 Jul;11(7):e1005359.

9.   Turek M, Besseling J, Spies JP, König S, Bringmann H. Sleep-active neuron specification

and sleep induction require FLP-11 neuropeptides to systemically induce sleep. eLife.

2016 Mar 7;5:e12499.

10.   Gray JM, Hill JJ, Bargmann CI. A circuit for navigation in *Caenorhabditis elegans.* Proc

Natl Acad Sci U S A. 2005 Mar 1;102(9):3184–91.

11. Hills T, Brockie PJ, Maricq AV. Dopamine and glutamate control area-restricted search

behavior in *Caenorhabditis elegans.* J Neurosci. 2004 Feb 4;24(5):1217-25.

12.   You Y jai, Kim J, Raizen DM, Avery L. Insulin, cGMP, and TGF-beta signals regulate food

intake and quiescence in *C. elegans*: a model for satiety. Cell Metab. 2008 Mar;7(3):249–

57.

13.   Hill AJ, Mansfield R, Lopez JMNG, Raizen DM, Van Buskirk C. Cellular stress induces a

protective sleep-like state in *C. elegans*. Curr Biol CB. 2014 Oct 20;24(20):2399–405.

14.   Nelson MD, Lee KH, Churgin MA, Hill AJ, Van Buskirk C, Fang-Yen C, et al. FMRFamide-

like FLP-13 neuropeptides promote quiescence following heat stress in *Caenorhabditis*

*elegans*. Curr Biol CB. 2014 Oct 20;24(20):2406–10.

15.   DeBardeleben HK, Lopes LE, Nessel MP, Raizen DM. Stress-Induced Sleep After

Exposure to Ultraviolet Light Is Promoted by p53 in *Caenorhabditis elegans*. Genetics.

2017 Oct;207(2):571–82.

16.   Skora S, Mende F, Zimmer M. Energy Scarcity Promotes a Brain-wide Sleep State

Modulated by Insulin Signaling in *C. elegans*. Cell Rep. 2018 Jan 23;22(4):953–66.

17. Wu Y, Masurat F, Preis J, Bringmann H. Sleep Counteracts Aging Phenotypes to Survive

Starvation-Induced Developmental Arrest in *C. elegans.* Curr Biol CB. 2018 Nov

19;28(22):3610-3624.e8.

18. Daniels SA, Ailion M, Thomas JH, Sengupta P. *egl-4* acts through a transforming growth

factor-beta/SMAD pathway in *Caenorhabditis elegans* to regulate multiple neuronal circuits

in response to sensory cues. Genetics. 2000 Sep;156(1):123–41.

19. L'Etoile ND, Coburn CM, Eastham J, Kistler A, Gallegos G, Bargmann CI. The cyclic GMP-

dependent protein kinase EGL-4 regulates olfactory adaptation in *C. elegans.* Neuron.

2002 Dec 19;36(6):1079–89.

20. McCloskey RJ, Fouad AD, Churgin MA, Fang-Yen C. Food responsiveness regulates

episodic behavioral states in *Caenorhabditis elegans.* J Neurophysiol. 2017 May

1;117(5):1911–34.

21. Gurevich VV, Gurevich EV. GPCR Signaling Regulation: The Role of GRKs and Arrestins.

Front Pharmacol. 2019;10:125.

22. Gurevich EV, Tesmer JJG, Mushegian A, Gurevich VV. G protein-coupled receptor

kinases: more than just kinases and not only for GPCRs. Pharmacol Ther. 2012

Jan;133(1):40–69.

23. Fukuto HS, Ferkey DM, Apicella AJ, Lans H, Sharmeen T, Chen W, et al. G protein-

coupled receptor kinase function is essential for chemosensation in *C. elegans.* Neuron.

2004 May 27;42(4):581–93.

24.  Wood JF, Wang J, Benovic JL, Ferkey DM. Structural domains required for *Caenorhabditis*

*elegans* G protein-coupled receptor kinase 2 (GRK-2) function in vivo. J Biol Chem. 2012

Apr 13;287(16):12634–44.

25.  Hukema RK, Rademakers S, Dekkers MPJ, Burghoorn J, Jansen G. Antagonistic sensory

cues generate gustatory plasticity in *Caenorhabditis elegans*. EMBO J. 2006 Jan

25;25(2):312–22.

26.  Wang J, Luo J, Aryal DK, Wetsel WC, Nass R, Benovic JL. G protein-coupled receptor

kinase-2 (GRK-2) regulates serotonin metabolism through the monoamine oxidase AMX-2

in *Caenorhabditis elegans.* J Biol Chem. 2017 Apr 7;292(14):5943–56.

27.  Topalidou I, Cooper K, Pereira L, Ailion M. Dopamine negatively modulates the NCA ion

channels in *C. elegans*. PLoS Genet. 2017 Oct;13(10):e1007032.

28.  Xu Y, Zhang L, Liu Y, Topalidou I, Hassinan C, Ailion M, et al. Dopamine receptor DOP-1

engages a sleep pathway to modulate swimming in *C. elegans.* iScience. 2021 Apr

23;24(4):102247.

29.  Heilman C, Ibarreta J, Salonga GA, Hon MC, Caracci AM, Badial T, et al. G protein

coupled receptor kinases modulate *Caenorhabditis elegans* reactions to heat stresses.

Biochem Biophys Res Commun. 2020 Oct 1;530(4):692–8.

30.  Henry SA, Crivello S, Nguyen TM, Cybulska M, Hoang NS, Nguyen M, et al. G protein-

coupled receptor kinase 2 modifies the ability of *Caenorhabditis elegans* to survive

oxidative stress. Cell Stress Chaperones. 2021 Jan;26(1):187–97.

31.  Choi S, Chatzigeorgiou M, Taylor KP, Schafer WR, Kaplan JM. Analysis of NPR-1 reveals

a circuit mechanism for behavioral quiescence in *C. elegans*. Neuron. 2013 Jun

5;78(5):869–80.

32.  Nichols ALA, Eichler T, Latham R, Zimmer M. A global brain state underlies *C. elegans*

sleep behavior. Science. 2017 Jun 23;356(6344):eaam6851.

33.  Soto R, Goetting DL, Van Buskirk C. NPR-1 Modulates Plasticity in *C. elegans* Stress-

Induced Sleep. iScience. 2019 Sep 27;19:1037–47.

34.  Hums I, Riedl J, Mende F, Kato S, Kaplan HS, Latham R, et al. Regulation of two motor

patterns enables the gradual adjustment of locomotion strategy in *Caenorhabditis elegans.*

eLife. 2016 May 25;5:e14116.

35.  Buntschuh I, Raps DA, Joseph I, Reid C, Chait A, Totanes R, et al. FLP-1 neuropeptides

modulate sensory and motor circuits in the nematode *Caenorhabditis elegans.* PloS One.

2018;13(1):e0189320.

36.  Oranth A, Schultheis C, Tolstenkov O, Erbguth K, Nagpal J, Hain D, et al. Food Sensation

Modulates Locomotion by Dopamine and Neuropeptide Signaling in a Distributed Neuronal

Network. Neuron. 2018 Dec 19;100(6):1414-1428.e10.

37.  Topalidou I. Background mutation in strain RB2126 affects the locomotion behavior of *flp-1*

mutants. MicroPublication Biol. 2021 May 9;2021.

38.  Coleman B, Topalidou I, Ailion M. Modulation of Gq-Rho Signaling by the ERK MAPK

Pathway Controls Locomotion in *Caenorhabditis elegans.* Genetics. 2018 Jun;209(2):523–

35.

39.  Roayaie K, Crump JG, Sagasti A, Bargmann CI. The G alpha protein ODR-3 mediates

olfactory and nociceptive function and controls cilium morphogenesis in *C. elegans*

olfactory neurons. Neuron. 1998 Jan;20(1):55–67.

40.  Troemel ER, Chou JH, Dwyer ND, Colbert HA, Bargmann CI. Divergent seven

transmembrane receptors are candidate chemosensory receptors in *C. elegans.* Cell. 1995

Oct 20;83(2):207–18.

41.  Sengupta P, Chou JH, Bargmann CI. *odr-10* encodes a seven transmembrane domain

olfactory receptor required for responses to the odorant diacetyl. Cell. 1996 Mar

22;84(6):899–909.

42.  Troemel ER, Kimmel BE, Bargmann CI. Reprogramming chemotaxis responses: sensory

neurons define olfactory preferences in *C. elegans.* Cell. 1997 Oct 17;91(2):161–9.

43.  Troemel ER, Sagasti A, Bargmann CI. Lateral signaling mediated by axon contact and

calcium entry regulates asymmetric odorant receptor expression in *C. elegans.* Cell. 1999

Nov 12;99(4):387–98.

44.  Yu S, Avery L, Baude E, Garbers DL. Guanylyl Cyclase expression in specific sensory

neurons: a new family of chemosensory receptors. Proc Natl Acad Sci U S A. 1997 Apr

1;94(7):3384–7.

45.  Flames N, Hobert O. Gene regulatory logic of dopamine neuron differentiation. Nature.

2009 Apr 1;458(7240):885–9.

46.  Inglis PN, Ou G, Leroux MR, Scholey JM. The sensory cilia of *Caenorhabditis elegans.*

2007 Mar 8; WormBook, ed. The *C. elegans* Research Community, WormBook. Available

from: http://www.wormbook.org

47.    Churgin MA, Jung SK, Yu CC, Chen X, Raizen DM, Fang-Yen C. Longitudinal imaging of

765           *Caenorhabditis elegans* in a microfabricated device reveals variation in behavioral decline

766           during aging. eLife. 2017 May 31;6:e26652.

767    48.    Churgin MA, Szuperak M, Davis KC, Raizen DM, Fang-Yen C, Kayser MS. Quantitative

768           imaging of sleep behavior in *Caenorhabditis elegans* and larval Drosophila melanogaster.

769           Nat Protoc. 2019 May;14(5):1455–88.

770    49.    van der Linden AM, Wiener S, You Y jai, Kim K, Avery L, Sengupta P. The EGL-4 PKG

771           acts with KIN-29 salt-inducible kinase and protein kinase A to regulate chemoreceptor

772           gene expression and sensory behaviors in *Caenorhabditis elegans.* Genetics. 2008

773           Nov;180(3):1475–91.

774    50.    Grubbs JJ, Lopes LE, van der Linden AM, Raizen DM. A salt-induced kinase is required for

775           the metabolic regulation of sleep. PLoS Biol. 2020 Apr;18(4):e3000220.

776    51.    Turek M, Lewandrowski I, Bringmann H. An AP2 transcription factor is required for a sleep-

777           active neuron to induce sleep-like quiescence in *C. elegans.* Curr Biol CB. 2013 Nov

778           18;23(22):2215–23.

779    52.    Pujol N, Torregrossa P, Ewbank JJ, Brunet JF. The homeodomain protein

780           CePHOX2/CEH-17 controls antero-posterior axonal growth in *C. elegans.* Dev Camb Engl.

781           2000 Aug;127(15):3361–71.

782    53.    Trojanowski NF, Nelson MD, Flavell SW, Fang-Yen C, Raizen DM. Distinct Mechanisms

783           Underlie Quiescence during Two *Caenorhabditis elegans* Sleep-Like States. J Neurosci

784           Off J Soc Neurosci. 2015 Oct 28;35(43):14571–84.

54.   Macosko EZ, Pokala N, Feinberg EH, Chalasani SH, Butcher RA, Clardy J, et al. A hub-

and-spoke circuit drives pheromone attraction and social behaviour in *C. elegans.* Nature.

2009 Apr 30;458(7242):1171–5.

55.   Kubiak TM, Larsen MJ, Nulf SC, Zantello MR, Burton KJ, Bowman JW, et al. Differential

activation of "social" and "solitary" variants of the *Caenorhabditis elegans* G protein-

coupled receptor NPR-1 by its cognate ligand AF9. J Biol Chem. 2003 Sep

5;278(36):33724–9.

56.   Rogers C, Reale V, Kim K, Chatwin H, Li C, Evans P, et al. Inhibition of *Caenorhabditis*

*elegans* social feeding by FMRFamide-related peptide activation of NPR-1. Nat Neurosci.

2003 Nov;6(11):1178–85.

57.   Coates JC, de Bono M. Antagonistic pathways in neurons exposed to body fluid regulate

social feeding in *Caenorhabditis elegans*. Nature. 2002 Oct 31;419(6910):925–9.

58.   Fares H, Greenwald I. Genetic analysis of endocytosis in *Caenorhabditis elegans:*

coelomocyte uptake defective mutants. Genetics. 2001 Sep;159(1):133–45.

59.   Sieburth D, Madison JM, Kaplan JM. PKC-1 regulates secretion of neuropeptides. Nat

Neurosci. 2007 Jan;10(1):49–57.

60.   Speese S, Petrie M, Schuske K, Ailion M, Ann K, Iwasaki K, et al. UNC-31 (CAPS) Is

Required for Dense-Core Vesicle But Not Synaptic Vesicle Exocytosis in *Caenorhabditis*

*elegans.* J Neurosci. 2007 Jun 6;27(23):6150–62.

61.   Rossignol S, Dubuc R, Gossard JP. Dynamic sensorimotor interactions in locomotion.

Physiol Rev. 2006 Jan;86(1):89–154.

62.  Lewis JA, Hodgkin JA. Specific neuroanatomical changes in chemosensory mutants of the

nematode *Caenorhabditis elegans.* J Comp Neurol. 1977 Apr 1;172(3):489–510.

63.  Perkins LA, Hedgecock EM, Thomson JN, Culotti JG. Mutant sensory cilia in the nematode

*Caenorhabditis elegans.* Dev Biol. 1986 Oct;117(2):456–87.

64.  Mukhopadhyay S, Lu Y, Shaham S, Sengupta P. Sensory signaling-dependent remodeling

of olfactory cilia architecture in *C. elegans.* Dev Cell. 2008 May;14(5):762–74.

65.  Nagy S, Tramm N, Sanders J, Iwanir S, Shirley IA, Levine E, et al. Homeostasis in *C.*

*elegans* sleep is characterized by two behaviorally and genetically distinct mechanisms.

eLife. 2014 Dec 4;3:e04380.

66.  Omar HH, Humphries JE, Larsen MJ, Kubiak TM, Geary TG, Maule AG, et al. Identification

of a platyhelminth neuropeptide receptor. Int J Parasitol. 2007 Jun;37(7):725–33.

67.  White JG, Southgate E, Thomson JN, Brenner S. The structure of the nervous system of

the nematode *Caenorhabditis elegans.* Philos Trans R Soc Lond B Biol Sci. 1986 Nov

12;314(1165):1–340.

68.  Brenner S. The genetics of *Caenorhabditis elegans.* Genetics. 1974 May;77(1):71–94.

69.  Frøkjær-Jensen C, Davis MW, Sarov M, Taylor J, Flibotte S, LaBella M, et al. Random and

targeted transgene insertion in *Caenorhabditis elegans* using a modified Mos1 transposon.

Nat Methods. 2014 May;11(5):529–34.

70.  Frøkjær-Jensen C, Davis MW, Ailion M, Jorgensen EM. Improved Mos1-mediated

transgenesis in *C. elegans.* Nat Methods. 2012 Jan 30;9(2):117–8.

71.   Mello CC, Kramer JM, Stinchcomb D, Ambros V. Efficient gene transfer in *C.elegans*:

extrachromosomal maintenance and integration of transforming sequences. EMBO J. 1991

Dec;10(12):3959–70.

**Figure captions**

**Fig 1. *grk-2* mutant animals have dispersal and exploration defects.**

(A) Schematic representation of the assay used to measure dispersal (see Methods).

(B) *grk-2(gk268)* mutant animals have a dispersal defect in the presence of food and *dop-3(vs106)* does not rescue this phenotype. Shown is the distance (cm) animals traveled from the center of the plate in the presence of food in 1 h. (****, $P<0.0001$. ns, $P>0.05$. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 30–53).

(C) *grk-2(gk268)* mutant animals have a dispersal defect in the absence of food and *dop-3(vs106)* does not rescue this phenotype. Shown is the distance (cm) animals traveled from the center of the plate without food in 20 min. (****, $P<0.0001$. ns, $P>0.05$. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 47-60).

(D) Schematic representation of the assay used to measure exploration (see Methods).

(E) *grk-2(gk268)* mutant hermaphrodites have an exploration defect and *dop-3(vs106)* does not rescue this defect. Shown is the number of squares (out of 50) single animals covered in approximately 20 h. (****, $P<0.0001$. ns, $P>0.05$. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 30-32).

(F) *grk-2(gk268)* mutant males have an exploration defect. (****, $P<0.0001$. Student's t-test. Error bars = SEM; n = 30).

**Fig 2. *grk-2* acts in the ciliated neurons to control exploration behavior.**

(A) Expression of *grk-2* cDNA under a pan-neuronal promoter rescues the exploration defect of *grk-2* mutant animals in the presence of food. The *grk-2* cDNA was expressed in *grk-2(gk268)* mutant animals under a pan-neuronal promoter (*rab-3p*, transgene *yakEx44*), ventral cord acetylcholine motor neuron promoter (*acr-2p*, transgene *yakEx47*), and interneuron promoter (*nmr-1p*, transgene *yakSi32*). Shown is the number of squares single animals covered in ~20 h.

(****, P<0.0001. ns, P>0.05. One way ANOVA followed by a Bonferroni test. Error bars = SEM;

n = 20-25).

(B) Expression of *grk-2* cDNA under either of two ciliated sensory neuron promoters (*xbx-1p*,

transgene *yakEx71* and *osm-6p*, transgene *yakEx53*) rescues the exploration defect of *grk-*

*2(gk268)* mutant animals in the presence of food. (****, P<0.001. ns, P>0.05. One way ANOVA

followed by a Bonferroni test. Error bars = SEM; n = 20-30).

(C) Expression of *grk-2* cDNA under a ciliated sensory neuron promoter (*xbx-1p::grk-2*,

transgene *yakEx71*) rescues the dispersal defect of *grk-2(gk268)* in the absence of food. (****,

P<0.0001. ns, P>0.05. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n =

49-55). Please note the high variability in the off-food response in comparison to the on-food

response (Fig. 2B). This could be because worms are more mobile in the absence of food and

constantly searching for a food source, which would increase the variability of the distances they

cover.

(D) Expression of *grk-2* cDNA under the *odr-3* promoter (expression in AWA, AWB, AWC, ASH,

and ADF neurons; *odr-3p::grk-2*, transgene *yakEx189*) partially rescues the exploration defect

of *grk-2(gk268)*. (****, P<0.0001. One way ANOVA followed by a Bonferroni test. Error bars =

SEM; n = 27-44).

(E) Expression of *grk-2* cDNA under the *sra-6* promoter (expression in ASH and ASI neurons;

*sra-6p::grk-2*, transgene *yakEx191*) or under the combined control of the *odr-10* (AWA

neurons), *str-1* (AWB neurons) and *str-2* (AWC ON neuron) promoters (*str-1p, str-2p, odr-*

*10p::grk-2*, transgene *yakEx194*) partially rescues the exploration defect of *grk-2(gk268)*. (****,

P<0.0001. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 25-44).

(F) Expression of *grk-2* cDNA under the *odr-10p* (AWA), *str-1p* (AWB), or *str-2p* (AWC ON)

promoters (transgenes *pekEx267, pekEx265* and *pekEx266,* respectively) partially rescues the

*grk-2* exploration defect. (***, P<0.001. **, P<0.01. *, P<0.05. One way ANOVA followed by a

Bonferroni test. Error bars = SEM; n = 20-33). Twenty of the data points shown for WT and *grk-*

*2* mutant are the same as in Figure 3B (these experiments were run in parallel).

(G) *grk-2* is expressed in many neurons in the head of the worm including *odr-3-* expressing

neurons. Representative images of the z-stack projections of the head of an animal co-

expressing tagRFP fused to GRK-2 ORF driven by the *grk-2* promoter (*grk-2*::tagRFP,

transgene *yakIs19*) and NLS-tagged mNeon driven by the *odr-3* promoter (*odr-3p*::mNeon::NLS;

transgene *yakEx204*). Arrows indicate *grk-2*::tagRFP-expressing cells that also express *odr-*

*3p*::mNeon::NLS. Scale bar: 10 µm.

(H) Expression of *grk-2* cDNA under the *gcy-8* promoter (expression in AFD neurons; *gcy-*

*8p::grk-2*, transgene *yakEx202*) fails to rescue the exploration defect of *grk-2(gk268)*.

(ns, P>0.05. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 20-40).

(I) Expression of *grk-2* cDNA under the *dat-1* promoter (expression in dopaminergic neurons;

*dat-1p::grk-2*, transgene *yakEx259*) fails to rescue the exploration defect of *grk-2(gk268)*. (ns,

P>0.05. Student's t-test. Error bars = SEM; n = 38-44).

(J) Expression of *grk-2* cDNA under the heat shock promoter *hsp-16.41* (*hsp-16.41p::grk-2*,

transgene *yakEx253*) in adult *grk-2(gk268)* partially but significantly rescues their exploration

defect. (****, P<0.0001. Student's t-test. Error bars = SEM; n = 34-44).

**Fig 3. Mutants in cilia-related genes have exploration defects.**

(A) The *che-2(e1033)* mutant has an exploration defect like *grk-2(gk268)* that is not enhanced

by *grk-2*. Shown is the number of squares single animals covered in ~20 h. (****, P<0.0001. ns,

P>0.05. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 22).

(B) The *daf-10(e1387)* mutant has an exploration defect like *grk-2(gk268)* that is not enhanced

by *grk-2*. Shown is the number of squares single animals covered in ~20 h. (****, P<0.0001. ns,

P>0.05. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 20-25). The data

points shown for WT and *grk-2* mutant are the same as in Figure 2F (these experiments were

run in parallel).

(C) The *che-2(e1033)* and *daf-10(e1387)* mutants have a dispersal defect independently of food

that is not enhanced by *grk-2(gk268).* Shown is the distance (cm) animals traveled from the

center of the plate without food in 20 min. (****, $P<0.0001$. One way ANOVA followed by a

Bonferroni test. Error bars = SEM; n = 30-40).

(D) Most chemosensory and osmosensory-defective mutants that possess strong or weak

cilium-structure defects based on a dye-filling assay ([46]; shown in blue and green,

respectively) have exploration defects. Chemosensory-defective mutants *odr-1, odr-3, odr-8,*

*odr-10, tax-2,* and *tax-4*; osmosensory-defective mutants *osm-7* and *osm-10;* and

thermosensory-defective mutant *ttx-1(p767),* also have exploration defects. In contrast,

chemosensory mutants *che-1, che-6, che-7, odr-2, odr-4, odr-5, odr-7,* osmosensory mutants

*osm-9* and *osm-11*, oxygen-sensing mutant *gcy-35,* and mechanosensory-defective mutants

*mec-3, mec-4,* and *glr-1* do not have exploration defects. The canonical null allele *ocr-2(ak47)*

(strain VM396; indicated with an asterisk) showed a significant exploration defect that was not

observed with the presumably null allele *ocr-2(ok1711)*. Outcrossed strain BJH2763 *ocr-2(ak47)*

did not show a similar exploration defect, suggesting that the *ocr-2(ak47)* mutation per se does

not cause exploration defects (Fig S3A; see text for details). Shown is the number of squares

(out of 50) single animals covered in ~20 h, normalized to WT. (***, $P<0.001$. **, $P<0.01$. *,

$P<0.05$. ns, $P>0.05$. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 10-

35).

**Fig 4. *egl-4* suppresses the exploration defect, movement quiescence, and reduced**

**activity of *grk-2* mutant animals.**

(A), (B) The *egl-4(n479)* mutation partially suppresses the exploration defect of *grk-2(gk268)*

mutant as assayed on 6 cm (A) and 10 cm (B) plates. Shown is the number of squares (out of

50, (A) or 150, (B)) single animals covered in ~20 h. Please note that *egl-4* mutant animals

explore more than wild type animals as assayed on 10 cm plates. (****, P<0.0001. ns, P>0.05.

One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 22-26).

(C) Expression of *egl-4* cDNA in ciliated sensory neurons under the *osm-6* promoter reverses

the *egl-4* suppression of the exploration defect of *grk-2* mutant (*osm-6p::egl-4*, transgene

*yakEx221*). Shown is the number of squares single animals covered in ~20 h (****, P<0.0001.

ns, P>0.05. Student's t-test. Error bars = SEM; n = 31-33).

(D) *grk-2(gk268)* mutant animals show enhanced movement quiescence, and the *egl-4(n479)*

mutation suppresses this phenotype. The graph shows the average minutes animals spent in

quiescence per hour in the WorMotel. Quiescence is the time when zero worm pixels moved

between two frames separated by 10 sec (See Methods).  One WT animal was considered an

outlier and was not included in the graph since its values were more than 3 standard deviations

from the mean.

(E) The graph shows the total minutes individual animals spent in quiescence in 8 h in the

WorMotel. (****, P<0.0001. ns, P>0.05. One way ANOVA. Error bars = SEM; n = 11-12).

(F) *grk-2(gk268)* mutant animals show reduced activity, and the *egl-4(n479)* mutation partially

suppresses this phenotype. The graph shows the activity in pixels of individual animals in 8 h in

the WorMotel. Activity is the number of pixels moved between frames separated by 10 sec.

(****, P<0.0001. One way ANOVA. Error bars = SEM; n = 11-12).

**Fig 5. *aptf-1* partially suppresses the exploration defect, movement quiescence, and**

**reduced activity of *grk-2*.**

(A) The *aptf-1(gk794)* mutation partially suppresses the exploration defect of *grk-2(gk268)*

mutant. Shown is the number of squares single animals covered in ~20 h. (****, P<0.0001. ns,

P>0.05. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 21-62).

(B) *grk-2(gk268)* mutant animals show enhanced movement quiescence, and the *aptf-1(gk794)*

mutation partially suppresses this phenotype. The graph shows the average minutes animals

spent in quiescence per hour in the WorMotel. Quiescence is the time when zero pixels moved

between two frames separated by 10 sec (See Methods).

(C) *grk-2(gk268)* mutant animals show enhanced movement quiescence, and the *aptf-1(gk794)*

mutation partially suppresses this phenotype. The graph shows the total minutes individual

animals spent in quiescence in 8 h in the WorMotel. (****, P<0.0001. ns, P>0.05. One way

ANOVA. Error bars = SEM; n = 10-12).

(D) *grk-2(gk268)* mutant animals show reduced activity, and the *aptf-1(gk794)* mutation partially

suppresses this phenotype. The graph shows the activity in pixels of individual animals in 8 h in

the WorMotel. Activity is the number of pixels moved between frames separated by 10 sec.

(****, P<0.05. **, P<0.01. One way ANOVA. Error bars = SEM; n = 10-12).

**Fig 6. *npr-1* suppresses the exploration defect of *grk-2* mutant animals.**

(A) The *npr-1(ky13)* mutation suppresses the exploration defect of *grk-2(gk268)* mutants as

assayed on 6 cm plates. Shown is the number of squares (out of 50) single animals covered in

~20 h. (****, P<0.0001. *, P<0.05. ns, P>0.05. One way ANOVA followed by a Bonferroni test.

Error bars = SEM; n = 20-42).

(B) The *npr-1(ky13)* mutation suppresses the exploration defect of *grk-2(gk268)* mutant as

assayed on 10 cm plates. Shown is the number of squares (out of 150) single animals covered

in ~20 h. (****, P<0.0001. *, P<0.05. One way ANOVA followed by a Bonferroni test. Error bars

= SEM; n = 23-25).

(C) The *npr-1(ky13)* mutation suppresses the dispersal defect of *grk-2(gk268)* in the absence of

food. (****, P<0.0001.  ***, P<0.001.  ns, P>0.05. One way ANOVA followed by a Bonferroni

test. Error bars = SEM; n = 39-51).

(D) The *flp-18(db99)* mutation partially but significantly suppresses the exploration defect of *grk-*

*2(gk268)*. (****, P<0.0001. ns, P>0.05. One way ANOVA followed by a Bonferroni test. Error

bars = SEM; n = 20-30). The data points shown for WT and *grk-2* mutant are the same as in

Figure S4A (these experiments were run in parallel).

(E) The *flp-21(ok889)* mutation does not suppress the exploration defect of *grk-2(gk268)*. (****,

P<0.0001. **, P<0.01. ns, P>0.05. One way ANOVA followed by a Bonferroni test. Error bars =

SEM; n = 20-26).

(F) Expression of *ncs-1p::nCre* and *flp-21p*::LoxStopLox::NPR-1 (expression in RMG neurons

((54); *RMGp::npr-1*, transgene *yakEx206*) reverses the *npr-1* suppression of the exploration

defect of *grk-2(gk268)* mutant. Expression of *npr-1* cDNA under the *sra-6* promoter (expression

in ASH and ASI neurons; *sra-6p::npr-1*, transgene *yakEx209*) fails to reverse the *npr-1*

suppression of the exploration defect of *grk-2(gk268)* (****, P<0.0001. ns, P>0.05. One way

ANOVA followed by a Bonferroni test. Error bars = SEM; n = 21-33).

**Fig 7. *flp-1* significantly suppresses the exploration defect of *grk-2*.**

(A) *grk-2(gk268)* mutant animals have an exploration defect and *flp-1(ok2811)* significantly

suppresses this defect. Shown is the number of squares (out of 50) single animals covered in

~20 h. (****, P<0.0001. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n =

20-26).

(B) *flp-1* and *flp-18* act in the same pathway to regulate exploration. *flp-1(ok2811)* and *flp-*

*18(db99)* partially suppress the exploration defect of *grk-2(gk268)* mutant and the *grk-2(gk268);*

*flp-1(ok2811); flp-18(db99)* triple mutant behaves like the *grk-2(gk268); flp-1(ok2811)* double

mutant. (****, P<0.0001. ***, P<0.001. ns, P>0.05. One way ANOVA followed by a Bonferroni

test. Error bars = SEM; n = 20-28).

(C) Expression of *flp-1* under its own promoter (*flp-1p::flp-1*, transgene *yakEx231*) or under the

*twk-47* promoter (expression in AVK neurons; *twk-47p::flp-1*, transgene *yakEx216),* reverses the

*flp-1* suppression of the exploration defect of *grk-2(gk268)* mutant. (****, P<0.0001. Student's t-

test. Error bars = SEM; n = 23-30).

(D) *npr-1(ky13)* suppresses the dispersal defect of *grk-2(gk268)* mutant and the *grk-2(gk268);*

*flp-1(ok2811); npr-1(ky13)* triple mutant behaves like the *grk-2(gk268); npr-1(ky13)* double

mutant. Shown is the distance (cm) animals traveled from the center of the plate in the presence

of food in 1 h. (ns, P>0.05. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n

= 51-67).

(E) *npr-1(ky13)* suppresses the exploration defect of *grk-2(gk268)* mutant and the *grk-2(gk268);*

*flp-1(ok2811); npr-1(ky13)* triple mutant behaves like the *grk-2(gk268); npr-1(ky13)* double

mutant. Shown is the number of squares (out of 50) single animals covered in ~20 h (ns,

P>0.05. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 20-23).

(F) *frpr-7(gk463846)* partially but significantly suppresses the exploration defect of *grk-2(gk268)*

mutant. *npr-6(tm1497)* has no effect on *grk-2(gk268)* exploration. (***, P<0.001. ns, P>0.05.

One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 23-39).

**Fig 8. *grk-2* mutant animals expressing FLP-1::mCherry show increased fluorescence in**

**their coelomocytes.**

(A) (Up) Schematic diagram of the coelomocyte uptake assay used to quantify FLP-1 secretion

from AVK interneurons. (Down) Representative images of coelomocytes of *flp-1(ok2811)* and

*grk-2(gk268); flp-1(ok2811)* mutant animals expressing a rescuing construct of FLP-1::mCherry

(FLP-1::mCherry; transgene *yakEx256*). Scale bars = 10 µm.

(B) mCherry fluorescence intensity of coelomocytes of *flp-1(ok2811)* and *grk-2(gk268); flp-*

*1(ok2811)* mutant animals expressing FLP-1::mCherry and a coelomocyte marker *unc-*

*122*::GFP. (FLP-1::mCherry; transgene *yakEx256).* (\*\*, P<0.01. Student's t-test. Error bars =

SEM; n = 57-59).

(C) GFP fluorescence intensity of coelomocytes of *flp-1(ok2811)* and *grk-2(gk268); flp-*

*1(ok2811)* mutant animals expressing FLP-1::mCherry and a coelomocyte marker *unc-*

*122*::GFP (*unc-122::GFP*; transgene *yakEx256*). (ns, P>0.05. Student's t-test. Error bars =

SEM; n = 29-30).

(D) Overexpression of FLP-1 reduces the exploration behavior of wild type animals. Shown is

the number of squares covered in ~20 h by single wild type animals, *flp-1* mutant expressing

low levels of FLP-1::mCherry (transgene *yakEx256;* 5 ng/ul injected), and wild type animals

overexpressing FLP-1 (transgene *yakEx261;* 60 ng/ul injected). (\*\*\*\*, P<0.0001. \*\*\*, P<0.001.

One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 29-47).

1042

**Fig 9. Model for GRK-2 modulation of locomotion behaviors in *C. elegans.***

GRK-2 positively controls (A) crawling and swimming by acting in premotor interneurons in a

manner that depends on the D2-like dopamine receptor DOP-3 [27,28] and (B) exploration by

acting in ciliated neurons in an opposite way from EGL-4. We propose a model in which GRK-2

acts in multiple ciliated sensory neurons to positively control sensory perception and (1) inhibit

the activity of the EGL-4 kinase in the same ciliated neurons, (2) negatively regulate the

neuropeptide receptor NPR-1 in RMG interneurons in a cell-nonautonomous way (e.g. by

regulating the level of secretion of an NPR-1 ligand), and (3) negatively control the levels of

FLP-1 secretion from AVK interneurons. AVKs play critical roles in regulating locomotion by

modulating motor neuron activity [36]. Since the ciliated neurons and AVK interneurons do not

have direct connections, we propose that GRK-2 controls FLP-1 secretion from AVK

interneurons in a cell-nonautonomous way through a neuromodulatory pathway. Proteins that

positively regulate crawling, swimming, or exploration are shown in green and proteins that

negatively regulate crawling, swimming, or exploration are shown in red. Dashed lines show

indirect interactions. Created with BioRender.com.

1058

**Fig S1. *grk-2* acts in parallel to *pdfr-1* and *tph-1* to control exploration.**

(A) *grk-2(gk268)* and *pdfr-1(ok3425)* mutants have exploration defects. The double mutant *grk-*

*2(gk268); pdfr-1(ok3425)* has a stronger phenotype than the single mutants suggesting that

these genes act in parallel to control exploration behavior. Shown is the number of squares (out

of 50) single animals covered in ~20 h. (****, $P<0.0001$. **, $P<0.01$. One way ANOVA followed

by a Bonferroni test. Error bars = SEM; n = 23-28).

(B) *grk-2(gk268)* mutant animals explore less and *tph-1(mg280)* more than WT animals. The

double mutant *tph-1(mg280); grk-2(gk268)* has an intermediate phenotype suggesting that

these genes act in parallel pathways to control exploration behavior. (****, $P<0.0001$. ***,

$P<0.001$. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 23-41).

1069

1070    **Fig S2. *grk-2* acts in command interneurons to regulate crawling.**

1071    *grk-2* cDNA expression in command interneurons (*Pnmr-1*, transgene *yakSi32*) is sufficient to

1072    rescue the slow locomotion of *grk-2(gk268)* mutant animals. (****, $P<0.0001$. ns, $P>0.05$. One

1073    way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 20).

1074

**Fig S3. *ocr-2(ak47), ocr-2(ok1711,)* and *kin-29(oy38)* mutants do not have exploration**

**defects.**

(A) *ocr-2(ak47), ocr-2(ok1717)* (strains BJH2763 and BJH2762, respectively) and *kin-29(oy38)*

mutants do not have exploration defects. Shown is the number of squares (out of 50) single

animals covered in ~20 h. (ns, $P>0.05$. One way ANOVA followed by a Bonferroni test. Error

bars = SEM; n = 20-24).

(B) *kin-29(oy38)* mutant animals do not have an exploration defect and the double mutant *grk-*

*2(gk268); kin-29(oy38)*) has an exploration defect similar to *grk-2(gk268).* (ns, P>0.05. One way

ANOVA followed by a Bonferroni test. Error bars = SEM; n = 20-21).

1084

**Fig S4. Mutations in *flp-11* and *ceh-17* do not affect the exploration defect and movement**

**quiescence of *grk-2*.**

(A) The *flp-11(tm2706)* mutation does not affect the exploration defect of *grk-2(gk268)* mutant

animals. Shown is the number of squares (out of 50) single animals covered in ~20 h. (****,

P<0.0001. ns, P>0.05. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n =

20-28). The data points shown for WT and *grk-2* mutant are the same as in Figure 6D (these

experiments were run in parallel).

(B) *grk-2(gk268)* mutant animals show enhanced movement quiescence and the *flp-11(tm2706)*

mutation does not affect this phenotype. The graph shows the average minutes animals spent in

quiescence per hour in the WorMotel. Quiescence is the time when there

were zero pixels moved between two frames separated by 10 sec (See Methods).

(C) The graph shows the total minutes individual animals spent in quiescence in 8h in the

WorMotel. (****, P<0.0001. ns, P>0.05. One way ANOVA. Error bars = SEM; n = 11-12).

(D) *grk-2(gk268)* mutant animals show reduced activity and the *flp-11(tm2706)* mutation does

not affect this phenotype. The graph shows the activity in pixels of individual animals in 8 h in

the WorMotel. Activity is the number of pixels moved between frames separated by 10 sec.

(****, P<0.0001. ns, P>0.05. One way ANOVA. Error bars = SEM; n = 11-12).

(E) The *ceh-17(np1)* mutation does not affect the exploration defect of *grk-2(gk268)* mutant

animals. Shown is the number of squares (out of 50) single animals covered in ~20 h. (ns,

P>0.05. One way ANOVA. Error bars = SEM; n = 20-24).

(F) *grk-2(gk268)* mutant animals show enhanced movement quiescence, and the *ceh-17(np1)*

mutation does not affect this phenotype.

(G) The graph shows the total minutes individual animals spent in quiescence in 8 h in the

WorMotel. (*, P<0.05. ***, P<0.001. ns, P>0.05. One way ANOVA. Error bars = SEM; n = 8-12).

(H) The graph shows the activity in pixels of individual animals in 8 h in the WorMotel.

(****, P<0.0001. ns, P>0.05. Error bars = SEM; n = 8-12).

1111

1112    **Fig S5. Expression of FLP-1::mCherry in *grk-2; flp-1* mutant animals reverses the**

1113    **suppression of the *grk-2* exploration defect by *flp-1*.**

1114    Shown is the number of squares (out of 50) single animals covered in ~20 h. (****, P<0.0001.

1115    ns, P>0.05. One way ANOVA followed by a Bonferroni test. Error bars = SEM; n = 34-39).

1116

1117    **Table S1. List of strains.**

1118    **Table S2. List of plasmids.**

1119    **Table S3. Raw data.**

1120

**Figure 1**

# Figure 2

**A** ON FOOD

*rab-3*: All neurons
*acr-2*: Ventral Cord motor neurons
*nmr-1*: Command interneurons

**B** ON FOOD

*xbx-1, osm-6*: Ciliated neurons

**C** OFF FOOD

*xbx-1*: Ciliated neurons

**D** ON FOOD

*odr-3*: AWA, AWB, AWC, ASH, ADF

**E** ON FOOD

*sra-6*: ASH, ASI
*odr-10*: AWA
*str-1*: AWB
*str-2*: AWC ON

**F** ON FOOD

**G** *grk-2*::tagRFP    *odr-3p*::mNeon::H2B    Merge

**H** ON FOOD

*gcy-8*: AFD

**I** ON FOOD

*dat-1*: dopamine neurons

**J** ON FOOD

# Figure 3

# Figure 4

# Figure 5

**Figure 7**

**Figure 8**

A

AVK interneurons

FLP-1::mCherry secretion

Coelomocyte uptake

Coelomocytes

*flp-1*    *grk-2; flp-1*

B  FLP-1::mCherry

**

20000
15000
10000
5000
0

Fluorescence

*flp-1*    *grk-2; flp-1*

C  *unc-122p*::GFP

ns

80000
60000
40000
20000
0

Fluorescence

*flp-1*    *grk-2; flp-1*

D  FLP-1 overexpression

****
***

60
40
20
0

# of squares covered

WT; FLP-1    *flp-1*; FLP-1    WT; FLP-1

**Figure 9**