Eccentric cycling involves greater mental demand and cortical activation of the frontoparietal network

Lénaic Borot[1,2], Benjamin Pageaux[3,4,5], Davy Laroche[6,7], Grégoire Vergotte[1], Romuald Lepers[6], Stéphane Perrey[1]

1. *EuroMov Digital Health in Motion, Univ Montpellier, IMT Mines Ales, Montpellier, France*
2. *Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, United Kingdom*
3. *École de kinésiologie et des sciences de l'activité physique (EKSAP), Faculté de médecine, Montréal, Canada*
4. *Centre de recherche de l'Institut universitaire de gériatrie de Montréal (CRIUGM), Montréal, Canada*
5. *Centre interdisciplinaire de recherche sur le cerveau et l'apprentissage (CIRCA), Montréal, Canada.*
6. *INSERM UMR 1093-CAPS, Univ. Bourgogne Franche-Comté, UFR des Sciences du Sport, Dijon, France*
7. *INSERM, CIC 1432, Module Plurithematique, Plateforme d'Investigation Technologique, Dijon, France ; CHU Dijon-Bourgogne, Centre d'Investigation Clinique, Module Plurithematique, Plateforme d'Investigation Technologique Dijon, France*

Short title: Cortical control of eccentric cycling

Corresponding author:

Prof. Stephane Perrey (PhD)

EuroMov Digital Health in Motion

700 avenue du Pic Saint Loup

34090 Montpellier France

Tel. (+33) 4 34 43 26 23

Email: stephane.perrey@umontpellier.fr

Abstract

Eccentric, compared to concentric muscle contraction, is thought to be attributed to different strategies of neuro-motor processing and a higher level of mental demand. This study aimed to evaluate the mental demand and related-cortical activation patterns to eccentric and concentric cycling at matched perceived effort and torque. Nineteen healthy men (30±6 yrs) performed four different 5-min cycling exercise conditions on a semi-recumbent isokinetic cycle ergometer: 1) concentric at a moderate perceived effort (23 on the CR100® scale) without torque feedback; 2) concentric and 3) eccentric at the same average torque produced in the first condition; and 4) eccentric at the same moderate perceived effort than the first concentric condition. The order of conditions 2-4 was randomised. After each condition, mental demand was monitored using the NASA-TLX scale. Changes in oxy- (O_2Hb) and deoxy- (HHb) haemoglobin during cycling exercise were measured over the two prefrontal cortices and the right parietal lobe from a 15-probe layout using a continuous-wave NIRS system. Mental demand was significantly higher when performing eccentric compared to concentric cycling ($p = .012$) and when the intensity was fixed by the torque rather than the perceived effort ($p < .001$). For both torque- or perceived effort-matched exercises, O_2Hb was significantly greater ($p < .001$) in both prefrontal cortices and right parietal lobe, and HHb decreased in the left and right prefrontal cortices during eccentric compared to concentric cycling. The current study supports that acute eccentric cycling involves a higher mental demand and frontoparietal network activation compared to concentric cycling.

1. INTRODUCTION

Most human movements involve both eccentric and concentric muscle contractions (e.g., walking, running, jumping) and a large proportion of our daily living activities require accurate control of movements employing eccentric contractions (e.g., walking downstairs, sitting, lowering weight). During eccentric contractions, the external force on the muscle is greater than the force that the muscle can generate, producing a lengthening contraction (vice versa for concentric contractions, Duchateau & Enoka, 2008). In the last decades, eccentric cycling at moderate intensity has been shown to promote functional (knee-extensor strength) and structural (hypertrophy) muscle adaptations in different populations due to its capability to achieve high loads on muscle tissue at low cardiorespiratory stress (athletic: Douglas et al., 2017; Lastayo et al., 1999; and chronic diseases: Pageaux et al., 2020).

Over more than 60 years, increasing research have been undertaken to understand mechanical, peripheral and neural features, as well as the metabolic cost of eccentric contractions. Eccentric cycling was initially introduced to investigate the physiology of concentric and eccentric muscle work (Abbott, Bigland, & Ritchie, 1952). Eccentric cycling requires substantially less oxygen uptake than concentric cycling to develop the same absolute external mechanical power output (Abbott, Bigland, & Ritchie, 1952; Perrey et al., 2001). In addition, corticospinal excitability, electromyographic activity and rating of perceived effort are lower during eccentric compared to concentric cycling at the same absolute mechanical workload (Laroche et al., 2013; Peñailillo, Mackay & Abbiss, 2018; Hollander et al., 2003; Perrey et al., 2001; Clos & Lepers, 2020; Clos et al., 2022). Noteworthy that at a given metabolic workload, heart rate and rating of perceived effort were found higher in eccentric compared to concentric cycling (Perrey et al., 2001; Clos et al., 2019; Clos et al., 2022).

Eccentric cycling is associated with substantial effort perception and sustained attention when subjects are asked to match a given target mechanical power output (Kan, Speelman, & Nosaka, 2019; Peñailillo, Blazevich, & Nosaka, 2017). However, little is known about cortical activity patterns of attentional resources during eccentric cycling. Current literature contains several lines of evidence indicating that neurobehavioral relationships in eccentric contractions are associated with higher processing demands by various brain areas for controlling motor output (see for a review Perrey, 2018).

At the behavioural level, several studies have reported greater variability in the motor output during eccentric compared to concentric contractions (Christou & Carlton, 2002; Fang et al., 2001, 2004; Semmler et al., 2007). This increased motor variability may indicate less practice and that eccentric contractions might be more cognitively demanding than concentric contractions (Perrey, 2018). This hypothesis is supported by the recent findings from Kan, Speelman, and Nosaka (2019) during submaximal eccentric and concentric cycling. First, an overall higher perception of the workload, evaluated by NASA task load index (NASA-TLX) questionnaire, was found during eccentric compared to concentric cycling. This difference is specifically related to two items of the questionnaire: mental demand and frustration. Second, the average choice reaction time was slower and the accuracy score was lower during a sustained attention to response task after eccentric compared to concentric cycling. Altogether, these findings suggest that eccentric cycling is taxing more the brain than concentric cycling does.

A neuroimaging modality used in the study of Kan, Speelman, and Nosaka (2019) was functional near-infrared spectroscopy (fNIRS). fNIRS is an optical neuroimaging technique based on the theory of neurovascular coupling (Liao et al., 2013; Nippert, Biesecker, & Newman, 2018) and is relatively robust to motion artifacts (Perrey, 2008). Since the local supply of oxygen is greater than its consumption in activated brain regions,

a higher concentration of oxy-haemoglobin (O_2Hb) and a decreased concentration of deoxy-haemoglobin (HHb) are to be observed (Perrey, 2008; Ferrari & Quaresima, 2012). Changes in O_2Hb and HHb recorded with 2 channels over the prefrontal cortex in Kan, Speelman, and Nosaka (2019) were found to be smaller during two eccentric cycling compared to a concentric cycling, while there were higher scores for the two eccentric cycling trials regarding the mental demand subscale of NASA-TLX. These preliminary findings suggest that a reduced supply of oxygen compared to relative mental demand occurs during eccentric cycling. However, they seem not to be supported by previous studies comparing the influence of the mode of muscle contractions from isolated and small muscle groups on brain activity (Perrey, 2018). Using electroencephalography during elbow flexion at matched torque, greater amplitude of movement-related cortical potentials in eccentric compared to concentric contractions were shown for electrodes located over the motor areas, prefrontal cortex and parietal lobe (Yue et al., 2000; Fang et al., 2001-2004). Using functional magnetic resonance imagery (fMRI), Yao et al. (2014) observed an increased blood-oxygen-level dependent signal in similar brain areas during an eccentric task of the first dorsal interosseous muscle as compared to a concentric task. Yet, none of the neuroimaging studies mentioned above directly investigated cortical activity over multiple brain areas during whole-body eccentric and concentric exercise, such as cycling.

According to the previous studies, the control of eccentric contraction would require important activation of the prefrontal regions, the right parietal lobe as well as of the motor regions. These different areas are part of the central executive network known to be involved in cognitive-motor tasks modulating attentional demand (Derosière et al., 2015) as well as in the modulation of mental demand during continuous cycling exercise (Radel, Brisswalter, & Perrey, 2017). As attention-demanding tasks exhibit increased activity in the prefrontal cortex and posterior parietal cortex, we hypothesized that the greater mental demand of eccentric cycling compared to concentric cycling would be associated with cortical activity

increases in bilateral prefrontal cortices and right posterior parietal cortex. Finally, most of the aforementioned studies investigating eccentric cycling used an experimental design imposing an absolute mechanical power output. Using effort perception for matching exercise intensity between eccentric and concentric cycling can be an alternative to circumvent the comparison of neurophysiological responses, either at a same mechanical or at the same metabolic workload (Besson et al., 2013; Clos et al., 2021). An effort perception approach to prescribe eccentric cycling has been proposed for the tailoring of rehabilitation program in deconditioned populations, such as patients living with cardio-respiratory chronic diseases leading to muscle weakness (Pageaux et al., 2020). Considering the gaps in the literature, the main objective of the present study was to quantify the acute changes of attention-related cortical activity as well as those of relevant brain motor areas from fNIRS during submaximal eccentric and concentric cycling exercise. As the method to prescribe exercise is likely to influence cortical activity and mental demand, eccentric and concentric cycling was performed at both matched perceived effort and torque.

2. METHODS

2.1. Participants

Nineteen healthy male adults (mean age: 30 ± 6 years) volunteered in the study. All participants reported no sign of neurological, respiratory and cardiovascular disease or medication, which might affect brain and muscle functions. Participants provided written informed consent prior to participation in the study. The study was approved by the local ethics committee (ID: RCB_2016-A01722-49) and was conducted in accordance with the ethical standards specified by the Declaration of Helsinki. This study is belonging to the ExoMode project registered on ClinicalTrials.gov (NCT03280875). The sample size required for the present study was calculated using G*Power (3.1.9.6). The fNIRS results of previous studies (Borot,

Vergotte, & Perrey, 2018; Kan, Speelman, & Nosaka, 2019) were used as group parameters. The power analysis indicated that a total of at least 17 participants was required for the brain activation (f = .85; α = .05; 1-β = .90). Accordingly, a sample size of 17 participants was recruited. Two additional participants were included in case of experimental outliers.

2.2. Experimental procedures

Participants were requested to come to the laboratory twice. The first visit consisted of a familiarization session and the second visit was the experimental session. Data collection occurs solely in the experimental session.

2.2.1 Familiarisation session

The first visit to the laboratory was a familiarisation session with a double objective. First, participants were familiarised with the CR100® scale (Borg & Borg, 2001) to evaluate their perception of effort and quadriceps pain, as well as to self-regulate the cycling exercise intensity based on a fixed perceived effort intensity. Second, the participants experimented with pedalling, driven by a motor either clockwise (concentric cycling) or anticlockwise (eccentric cycling), on an isokinetic semi-recumbent prototype cycle ergometer (Tech Med TMS, 89290 Champs-sur- Yonne, France). The seat of the semi-recumbent cycle ergometer allows for greater torso stabilization compared with a normal upright cycling position. This familiarisation cycling session consisted of participants understanding that they need to resist the pedals moving anticlockwise while they would have to accompany the pedals moving clockwise by exerting a suitable force to match a target line appearing on a screen in front of them. Participants were asked to perform eccentric and concentric cycling in a well-coordinated manner such that the knee extensors of each leg are activated at the right timing and intensity to ensure smooth pedal rotation and torque production. This familiarization session was carried out with the participant reaching submaximal torque values indicated

by visual feedback on the control screen. Finally, the participants performed four submaximal cycling exercise at a moderate intensity for three minutes. The mode of muscle contraction (eccentric vs concentric) and the prescription of the exercise (fixed perceived effort or torque) were manipulated as follow: 1) eccentric and 2) concentric cycling at a moderate effort intensity (23 on the CR100® scale); 3) eccentric and 4) concentric cycling at a fixed torque value. The familiarization session was carried out at least 48 hours before the experimental session.

2.2.2 Recording session

In this second session, participants performed four cycling exercise conditions for a duration of 5 min (see Figure 1), as experienced in the familiarisation session. In the first condition, participants performed concentric cycling at a fixed perception of effort intensity corresponding to 23 on the CR100® scale (moderate intensity). No force feedback was provided to the participants who were instructed to self-regulate their force during each pedal revolution solely on their effort perception intensity, and to keep it moderate during the 5-min of exercise. The maximal torque values produced at each pedal revolution were extracted from the cycle ergometer every 15 s from minutes 2 to 5 of the exercise. These values were then averaged to individually determine the target torque used in the subsequent conditions where exercise intensity was prescribed with a fixed torque. Then, the other three conditions were performed in a randomised order: (i) concentric cycling at an intensity corresponding to the average torque produced in the first condition, (ii) eccentric cycling at an intensity corresponding to the average torque produced in the first condition, (iii) eccentric cycling at a fixed perception of effort intensity corresponding to 23 on the CR100® scale (moderate, same perception of effort intensity than the first condition). Five minutes of rest were allowed between each exercise condition so that the physiological responses (heart rate and fNIRS signals) return to baseline levels. Heart rate was recorded

with a chest belt and heart rate monitor (Polar V800, Finland) during rest and exercise periods for each condition. The average heart rate was obtained during the resting period before the onset of exercise and over the last minute of each exercise condition.

A control experiment was performed on 5 subjects to ensure that there was no effect from the repetition of the four exercise sessions on muscle fatigue occurrence, defined as a decreased maximal force/torque production capacity of the working muscles (Gandevia, 2001). This control experiment consisted of duplicating the experimental session without measuring neurophysiological variables. Before the test session, the maximal voluntary isometric contraction (MVIC) of the dominant knee extensors was assessed (under verbal encouragement) as the best performance in three 5-s trials separated by a 30-s resting period. Then, one MVIC was performed in less than 20-s after the completion of each exercise condition. MVIC were performed on an isokinetic dynamometer (System Pro 4, Biodex Medical System, New York, NY) at a knee angle of 90°. The MVIC values did not significantly change throughout the test session for all the participants (F (4,16) = .71, p = .60; η^2_p = .15).

** please insert Figure 1 here **

2.3. Psychological variables

Perception of effort, also known as perceived exertion or sense of effort, can be described as a cognitive feeling of work associated with voluntary actions (Pageaux, 2016). Participants were asked to rate their perception of effort and quadriceps pain (CR100® scale) and affective valence during the last 20 seconds of each condition. In-task affective valence (i.e., the pleasure/displeasure one feels) was measured using the Feeling scale (Hardy & Rejeski, 1989), an 11-point bipolar scale ranging from -5 to +5. Seven anchors are provided ranging from "very good" (+5) to "very bad" (-5). Finally, participants were asked

to complete at the end of each condition two subscales (mental demand and physical demand) of a commonly used subjective workload level questionnaire, the NASA-TLX (Hart & Staveland, 1988). For each subscale, participants had to answer from 0 (very low or high) to 100 (very high or low).

2.4. Cortical activity

2.4.1 Recording

A two-wavelength (765 and 856 nm) continuous-wave 15-channels fNIRS system (Oxymon MkIII, Artinis Medical Systems, The Netherlands) was utilised to measure the changes in O_2Hb and HHb over regions of interests (ROI): left and right dorsolateral posterior parietal cortex, right parietal lobe and motor regions on both hemispheres as a whole (cycling task). The O_2Hb and HHb data were acquired during rest and exercise periods at a sampling rate of 10 Hz. O_2Hb and HHb concentration changes (expressed in $\Delta\mu M$) obtained by using the modified Beer-Lambert law were displayed on a PC monitor in real-time. The optodes (5 receivers and 13 transmitters resulting in a total of 15 channels, Figure 2a) were placed on the participants using a fit neoprene fNIRS head cap to get a stable optical contact with the scalp by generating an equal amount of pressure. The pressure was adequate to induce a partial transient blockage of the skin circulation during the fNIRS study (Takahashi et al., 2011) as witnessed by the presence of the well-defined circles over the skin that disappears about 15 min after the test session. In the 15 channels, the measurement points were defined as the midpoint of the corresponding illuminator-detector pairs interspaced of 3 cm. The Brodmann's areas covered by the different channels were extracted via the NFRI function (Singh et al., 2005) from the MNI (Montreal Neurological Institute) coordinates (Figure 2b). Cartesian coordinates were obtained with a Fastrak® 3D Digitizer (Polhemus Inc., USA) allowing the localization of each optode and channel from four anatomical reference points: nasion, right and left mastoids and vertex.

** please insert Figure 2 here **

2.4.2 Data analysis

Raw data (optical intensity) extracted from Oxysoft software (v3.0.75, Artinis Medical Systems) was converted in optical density (OD) using Matlab® (MATLAB 2014b, The MathWorks, USA). Then, to reduce possible head motion artifacts two methods implemented in the Homer2 toolbox (Huppert et al., 2009) were used. First, the moving standard deviation and spline interpolation methods were applied (Scholkmann et al., 2010) using parameters: SDTresh = 20, AMPTresh = 0.5, tMotion = 0.5s, tMask = 2s and p = 0.99. Second, we used wavelet-based signal decomposition (Molavi & Dumont, 2012) with parameter: iqr = 0.1, as recommended in Cooper et al. (2012) and Di Lorenzo et al. (2019). Then OD time series were converted into relative concentrations of O_2Hb and HHb using the modified Beer-Lambert law (Kocsis, Herman, & Eke, 2006) corrected by a differential pathlength factor depending on the age of the participant (Duncan et al., 1996). A low pass Butterworth zero-phase digital filter (order 4) with a cut-off frequency at 0.08 Hz (Yücel et al., 2016) was applied to limit physiological artifacts (heart rate: ± 1 Hz, respiration: ± 0.4 Hz and Mayer wave: ± 0.1 Hz). For each participant, channels not having a sufficient signal-to-noise ratio were excluded from the analyses after observation of the power spectrum density of the O_2Hb signals where the presence of a cardiac rhythm in the signal (peak around 1 Hz) indicates good contact between the optodes and the scalp (Themelis et al., 2007). After this preprocessing step, 14% of the channels were excluded.

Following a splitting of the fNIRS signals from events indicating the beginning of each condition, the average between -30 seconds and -10 seconds of the minute of rest preceding each condition was subtracted from the subsequent data during the exercise period. The following dependent variables attesting to exercise-related cortical activation magnitude

were calculated per channel for the period (representative of the steady state level) between 2 minutes and 4 minutes 30 seconds of exercise for each condition: area under the curve of O_2Hb and HHb and average values. The average of the channels positioned on 4 ROIs (the right and the left posterior parietal cortex and primary motor, and right parietal lobe) was extracted for each participant and each condition.

2.5 Statistical analysis

Statistical analyses were performed using Statistica (7.1, Statsoft Inc., France). Data were examined for normality and sphericity using Shapiro-Wilk and Mauchly tests, respectively. Where Mauchly's tests indicated violations of the sphericity assumption, Greenhouse–Geisser corrections were applied. A one-way repeated-measure ANOVA was used to assess the differences between the four exercise conditions for heart rate recorded during the rest period. A fully repeated measures two-way ANOVA with the factors *mode of muscle contraction* (eccentric vs. concentric) and *mode of exercise prescription* (torque vs. perception of effort) were used to compare the NASA-TLX scores and fNIRS responses. In case of significant effect, post hoc comparisons were adjusted with the Bonferroni correction.

Wilcoxon signed rank tests were performed for quadriceps pain and the Feeling scale values when comparing the mode of muscle contraction and the mode of exercise prescription. Student paired *t*-test was carried out between the concentric and eccentric torque conditions for the perceived effort. For the cycling torque data, Wilcoxon signed rank test was performed between the first condition (concentric at fixed perceived effort) and the two conditions (eccentric and concentric) where the intensity was fixed by a torque value.

Effect sizes were reported with the partial eta square (η^2_p) for the effects of the ANOVAs (Lakens, 2013) and with the Cohen's *d* for pairwise comparison (Cohen, 1988). The significance level for all analyses was established at $p = .05$ (two-tailed). All the fNIRS *p*-

values were corrected using false discovery rate to control for type 1 error due to the spatial dependency between brains areas (Benjamini & Yekutieli, 2001).

3. RESULTS

3.1. Torque and heart rate

No differences were observed between the torque value to match assessed from the first condition (concentric cycling at a fixed perception of 23; 41.12 ± 14.11 Nm) and either concentric (42.73 ± 13.80 Nm; W (17) = 47.5; p = 0.17; r_{rb} = 0.33) or eccentric (40.27 ± 13.15 Nm; W (17) = 40; p =0.08; r_{rb} = 0.42) cycling at an intensity corresponding to the average torque produced in the first condition.

No difference in resting heart rate was observed before the start of each cycling exercise condition (F (3, 45) = 1.25; p = .30; η^2_p = .08). During the cycling exercises, a main effect of the mode of muscle contraction revealed a higher heart rate during concentric cycling compared to eccentric cycling (F (1, 18) = 40.3; p < .001; η^2_p = .69, 101 ± 13 vs. 87 ± 13 bpm). No main effect of the mode of exercise prescription (F (1, 18) = .09; p = .77; η^2_p = .005; perception of effort: 94 ± 14 bpm; torque: 94 ± 15 bpm) or mode of muscle contraction × mode of exercise prescription interaction (F (1, 18) = .88; p = .36; η^2_p = .05; observed power = .15) was observed.

Taken together, these results confirm that participants performed properly the four cycling conditions as expected.

3.2. Psychological variables

When the exercise was prescribed with a fixed torque feedback, the *perception of effort intensity* was higher (t (17) = 3.39; p < .01; d = 1.075) during concentric cycling compared to eccentric cycling (23.2 ± 3.9 and 17.4 ± 6.8).

No main effects of the mode of muscle contraction (W (38) = 96.5; p = .75; r_{rb} = .05) and prescription of intensity (W (38) = 53; p = .44; r_{rb} = .13) were found for perceived quadriceps pain during the exercise as well as for the affect (Feeling scale) (W (38) = 94; p = .68; r_{rb} = .07 and W (38) = 79; p = .12; r_{rb} = .25, respectively).

For the perceived measures with the NASA-TLX subscales, one participant did not complete the questionnaire at the end of the experimental session. Therefore, data are available for 18 participants. For the *mental demand scores*, the main effect of the mode of muscle contraction revealed a higher mental demand during eccentric cycling compared to concentric cycling (F (1, 17) = 7.86, p = .01; η^2_p = .32; Figure 3). The main effect of the mode of exercise prescription revealed a higher mental demand when the exercise was prescribed with a fixed torque compared to a fixed perception effort (F (1, 17) = 26.22; p < .001; η^2_p = .61; Figure 3). No mode of muscle contraction × mode of exercise prescription interaction was observed (F (1, 17) = .45; p = .51; η^2_p = .02). For the *physical demand scores*, the main effect of the mode of muscle contraction revealed a higher physical demand during concentric cycling compared to eccentric cycling (F (1, 17) = 6.66; p = .02; η^2_p = .28; Figure 3). No main effect of the mode of exercise prescription was observed (F (1, 17) = .54; p = .47; η^2_p = .03; observed power = .11). Follow up tests on the mode of muscle contraction × mode of exercise prescription interaction (F (1, 17) = 6.99; p = .02, η^2p = .29) are presented in Figure 3. The physical demand was greater for the torque condition in concentric cycling compared to eccentric cycling (p = .005, d = 0.315).

** please insert Figure 3 here **

3.3. Cortical responses

For the *left prefrontal cortex* (Figure 4A) and O_2Hb, a main effect of the mode of muscle contraction revealed a higher area under the curve during eccentric cycling

compared to concentric cycling (F (1, 18) = 16.35; $p < .01$, η^2_p = .48). Neither a main effect of the mode of exercise prescription method (F (1, 18) = .1, $p = .75$; η^2_p = .01) nor a mode of muscle contraction × mode of exercise prescription interaction (F (1, 18) = 1.48; $p = .24$; η^2_p = .08) was observed. Regarding HHb, a main effect of the mode of muscle contraction revealed a lower area under the curve during eccentric cycling compared to concentric cycling (F (1, 18) = 35.77, $p < .001$, η^2_p = .67). Neither a main effect of the mode of exercise prescription (F (1, 18) = .49, $p = .49$; η^2_p = .03) nor a mode of muscle contraction × mode of exercise prescription interaction (F (1, 18) = .07; $p = .80$; η^2_p = .003) was observed.

For the *right prefrontal cortex* (Figure 4B) and O$_2$Hb, a main effect of the mode of muscle contraction revealed a higher area under the curve during eccentric cycling compared to concentric cycling (F (1, 18) = 6.96; $p < .05$; η^2_p = .28). Neither a main effect of the mode of exercise prescription (F (1,18) = .3; $p = .59$; η^2_p = .02) nor a mode of muscle contraction × mode of exercise prescription interaction (F (1, 18) = 2.06; $p = .17$; η^2_p = .10) was observed. Regarding HHb, a main effect of the mode of muscle contraction revealed a lower area under the curve during eccentric cycling compared to concentric cycling (F (1, 18) = 22.26, $p < .001$, η^2_p = .55). Neither a main effect of the mode of exercise prescription (F (1,18) = .63, $p = .44$; η^2_p = .03) nor a mode of muscle contraction × mode of exercise prescription interaction (F (1, 18) = 3.15, $p = .09$; η^2_p = .15) was observed. For bilateral *motor cortex* and O$_2$Hb (Figure 4C), no main effect of the mode of muscle contraction was observed (F (1,18) = 1.01, $p = .33$; η^2_p = .05). Neither a main effect of the mode of exercise prescription (F (1,18) = .18, $p = .68$; η^2_p = .01) nor a mode of muscle contraction × mode of exercise prescription interaction (F (1, 18) = .37, $p = .55$; η^2_p = .02) was observed. Regarding HHb, a main effect of the mode of muscle contraction revealed a lower area under the curve during eccentric cycling compared to concentric cycling (F (1, 18) = 38.26, $p < .001$, η^2_p = .68). Neither a main effect of the mode of exercise prescription (F (1, 18) = .73, $p = .40$; η^2_p

= .04) nor a mode of muscle contraction × mode of exercise prescription interaction (F = (1, 18) = 3.42, p = .08; η^2_p = .16) was observed.

For the *right parietal lobe* (Figure 4D) and O_2Hb, a main effect of the mode of muscle contraction revealed a higher area under the curve during eccentric cycling compared to concentric cycling (F (1, 13) = 11.12, p = .02, η^2_p = .46). Neither a main effect of the mode of exercise prescription method (F (1,13) = 1.09, p = .32; η^2_p = .08; Figure 4D) nor a mode of muscle contraction × mode of exercise prescription interaction (F (1, 13) = .20, p = .66; η^2_p = .01) was observed. Regarding HHb, no main effect of the mode of muscle contraction was observed (F (1, 13) = .38, p = .55; η^2_p = .03). Neither a main effect of the mode of exercise prescription (F (1, 13) = 2.2, p = .16; η^2_p = .14 nor a mode of muscle contraction × mode of exercise prescription interaction (F (1, 13) = 1.33, p = .27; η^2_p = .09) was observed.

** please insert Figure 4 here **

4. DISCUSSION

The main objective of the present study was to quantify the acute changes of attention-related cortical activity as well as those of relevant brain motor areas from fNIRS during eccentric and concentric cycling exercises performed at both matched perceived effort (23 on CR100® scale) and torque. The main results showed a greater mental demand and cortical activation in the right and left prefrontal cortices and right parietal lobe during eccentric cycling compared to concentric cycling. These findings confirm previous results suggesting that eccentric cycling is more cognitively demanding that concentric cycling (Kan, Speelman, & Nosaka, 2019) and extend them by identifying specific brain activation patterns underlying this greater mental demand.

4.1. Mental demand during eccentric cycling

Mental demand was subjectively measured with the NASA-TLX subscale. The mental demand was reported to be 45% greater during eccentric cycling compared to concentric cycling, while physical demand was 38% greater during concentric cycling. These findings support the results of Kan, Speelman, and Nosaka, (2019). Nevertheless, mental and physical demand values were 2 to 3 times lower in the present study. The higher mental and physical demand found in the Kan, Speelman & Nosaka (2019) is likely due to a greater exercise intensity compared to the present study, as confirmed by the higher mean heart rate values reported in their study as compared to ours (eccentric cycling: ~+29 bpm; concentric cycling: +56 bpm). While mental demand refers to the mental and perceptual activity imposed by a task and refers to related information processing (Hart and Staveland, 1988), effort refers to the voluntary engagement of mental and physical resources to perform, or attempt to perform, in a task (Inzlicht, Shenhav, & Olivola, 2018; Pageaux, 2016; Preston & Wegner, 2009). In the present study, coordinating properly the leg movements during eccentric and concentric cycling required dedicated motor skills and so adequate demand for attention. As described in the methods, participants were asked to perform eccentric and concentric cycling in a well-coordinated manner to ensure smooth pedal rotation and torque production. This was similar to the cycling tasks as proposed in Kan, Speelman, and Nosaka, (2019), where the authors observed that attention and vigilance were maintained after eccentric and concentric cycling. However, higher cognitive load (based on both the NASA TLX score and choice reaction time) was found in the study of Kan, Speelman, and Nosaka (2019) during eccentric cycling. Altogether, it appears that eccentric cycling is more cognitively demanding than concentric cycling and that the less cognitive demand of eccentric and concentric cycling in the present study did not likely change cognitive functions post-exercise.

Regarding the mode of exercise prescription, the mental demand was significantly higher when the intensity was set by a target torque value compared to an exercise

prescribed by a fixed perceived effort. To the best of our knowledge, this study is the first to observe this difference in mental demand. A lower mental demand during self-regulated exercise based on effort perception is not surprising as in this specific condition, participants did not have to visually focus on a torque target to reach at each pedal revolution. Contrary to the mental demand, the mode of exercise prescription did not alter the reported physical demand of the cycling exercise. This observation is in line with no significant difference of heart rate between the two modes of exercise prescription.

4.2. Cortical activity during eccentric cycling

Neuroimaging studies using various methods and exercise paradigms reveal inconsistencies in the role of the primary motor cortex in the production of eccentric muscle contractions (see for review, Perrey, 2018). Yao et al. (2014) during unilateral movements of the first dorsal interosseous muscle in MRI scanner, as Fang et al. (2001, 2004) in an EEG study for the elbow flexors, showed a greater activation of the primary motor cortex contralateral to the active muscles when performing eccentric compared to concentric contractions. Conversely, Olsson et al. (2012) and Kwon and Park (2011) reported a reduction in the primary motor cortex blood-oxygen-level dependent signal amplitude during eccentric compared to concentric contractions. The present study is the first to collect, during an eccentric cycling exercise (i.e., a bilateral task), fNIRS data over the precentral gyrus (BA 4/6) and primary motor areas (BA 4). The present study did not observe significant differences in motor cortices activity between eccentric and concentric cycling. Our results can be explained because the tasks performed are fundamentally different from the literature. First, the different studies cited above performed unilateral distal movements of the upper limb, unlike our study where the participants performed a bilateral task mobilizing larger muscle masses of the lower limbs. In addition, in the present study, the cycling task was performed at low intensity (23 on the CR100® scale or using an associated torque).

Studies showing a greater activity of the contralateral primary motor during eccentric muscle contractions involved exercises performed either at maximum intensity (Fang et al., 2004) or at a sub-maximum intensity fixed relative to maximal voluntary contraction force or body weight (Fang et al., 2001; Yao et al., 2014). Noteworthy, in the study of Olsson et al. (2012) a continuous imagery task paradigm was used by alternating concentric (lifting a weight) and eccentric (slowing down a weight) muscle contractions. For other cortical areas involved in the motor network (e.g. primary and secondary motor areas not recorded here), a consensus seems to exist on the greater cortical activity observed during eccentric contractions (Perrey, 2018).

The main results of the present study revealed a greater activity of the left and right prefrontal cortices, as well as the right parietal lobe, when performing eccentric cycling compared to concentric cycling, regardless of the mode of exercise prescription. These observations likely unveil underlying neurophysiological mechanisms sustaining the increased mental demand reported by the participants during eccentric cycling. The left and right prefrontal cortices as well as the right parietal lobe are known for their roles in the modulation of mental demand during prolonged exercise (Radel, Brisswalter, & Perrey, 2017) as well as during sustained attentional cognitive tasks (Mandrick et al., 2013, Derosière et al., 2013). When focusing on a specific movement to perform, several brain areas, constituting the central executive network, are engaged. More specifically, these regions would be part of the fronto-parietal network that consists mainly of the dorsolateral prefrontal cortex, the frontal cortex and the parietal lobe (Vincent et al., 2008). An increase in eccentric force variability has been proposed as a sign of an increased motor control difficulty during eccentric contractions compared to concentric contractions (Kwon & Park, 2011; Olsson et al., 2012). Controlling the variability in motor output (not assessed in the present study) during eccentric contractions would likely require greater attentional demand, thus explaining the observed increase in the activity of the prefrontal cortices and the right parietal lobe. Some

authors have highlighted the complementarity of regional brain activity magnitude measures and those linked to the dynamic activity of brain networks (Esfahlani et al., 2020). On this basis, the fMRI study of Yao et al. (2016) examined the strength of functional connectivity between the primary motor cortex and other cortical regions in the motor control network (e.g. supplementary motor area, cerebellum) involved in eccentric and concentric contractions. The authors observed a weaker functional connectivity during eccentric compared to concentric contractions of the right first dorsal interosseous muscle. A weaker statistical dependence between two-time series from different brain areas may reflect more involvement of specialised regions (segregation) and thus less functional integration (Friston, 2011). This means that higher cortical activation may be required during eccentric contraction to compensate for a weaker functional connectivity. Yao et al. (2016) hypothesized that the brain control of a concentric contraction will be more efficient than an eccentric contraction. This first result needs to be supported by more evidence, notably for a bilateral task mobilizing larger muscle masses of the lower limbs. The analysis of functional connectivity could be a relevant and complementary approach to better understand the functional organisation of the brain when performing eccentric cycling over time and its possible implications for rehabilitation and training.

4.3. Methodological considerations

In the present study, we controlled for the induction of muscle fatigue by providing sufficient recovery period between cycling exercise conditions, thus allowing us to ensure that our results are not confounded by a decreased in force production capacity of the working muscles with changes in the corticospinal excitability (Taylor et al., 1996; Gandevia et al., 1996). Exercise-induced muscle fatigue plays an important role in the modulation of cortical activity during exercise. In an fMRI study involving a maximal-effort handgrip contraction, Liu et al. (2002) observed an early increase followed by a decrease in brain activity in the

primary motor cortex and in particular in the prefrontal cortex. The exercise intensity in our experimental design was low as indicated by perceived effort and heart rate values. In addition, our control experiment showed no significant decrease in the maximal force production capacity of the knee extensors after the whole exercise protocol.

The present study faces one limitation, which might be taken into account in follow-up studies. The fNIRS signal can be contaminated by the physiological responses to the task performed (e.g. heart rate, blood pressure). This signal contamination related to non-neural activity can lead to false positives or negatives in our analyses (Tachtsidis & Scholkmann, 2016). To overcome these issues, the use of short-distance channels (approximately 1 cm between the transmitters and the receivers) was proposed (Brigadoi & Cooper, 2015). In this study, we did not use such short-distance channels for separating signals that stem from blood flow changes in the extracerebral layers (scalp blood flow) as well as the cerebral tissue layer (systemic blood flow). As indicated in the methods, we minimized this potential effect by inducing a partial transient blockade of the skin blood flow during the fNIRS data collection (Takahashi et al., 2011). Interestingly, heart rate was ~13% higher in concentric as reported before (e.g. Overend et al., 2000; Hollander et al., 2003), whereas the greatest cortical activation was observed during eccentric cycling. These elements allow us to think that the results might not be influenced by systemic interference but future investigations are necessary to confirm this hypothesis.

5. CONCLUSION

The present study showed that the higher mental demand observed during eccentric cycling compared to concentric cycling was associated to a greater activation of the left and right prefrontal cortices as well as the right parietal lobe. These data suggest a specific involvement of a frontoparietal network in the control of eccentric cycling. Based on the findings of this study, future studies should investigate the potential use of wearable fNIRS

for clinical and sports application via an online neurofeedback approach, for example, to decrease the mental demand occurring when using eccentric cycling. Another important application is that self-regulated eccentric exercise based on perceived effort is less mentally demanding, suggesting that it may be more appropriate if one session needs to minimize the overall workload for example.

ACKNOWLEDGEMENT

This research work was supported by the French National Research Agency (EXOMODE–ANR–15–CE19-0023).

AUTHOR CONTRIBUTIONS

Lenaic Borot: Conceptualization; data curation; formal analysis; investigation; methodology; validation; visualization; writing –original draft; writing –review and editing. Benjamin Pageaux: Conceptualization; investigation; methodology; visualization; validation; writing –review and editing. Davy Laroche: Conceptualization; funding acquisition; methodology; resources; supervision; writing –review and editing. Gregoire Vergotte: methodology; formal analysis; writing –review and editing. Romuald Lepers and Stephane Perrey: Conceptualization; investigation; funding acquisition; methodology; project administration; resources; supervision; validation; writing –review and editing.

CONFLICT OF INTEREST

All authors declare that they have no conflicts of interest.

REFERENCES

Abbott, B. C., Bigland, B., & Ritchie, J. M. (1952). The physiological cost of negative work. *The Journal of physiology*, *117*(3), 380. https://doi.org/10.1113/jphysiol.1952.sp004755

Benjamini, Y., & Yekutieli, D. (2001). The control of the false discovery rate in multiple testing under dependency. *Annals of statistics*, 1165-1188. http://doi.org/10.1214/aos/1013699998

Besson, D., Joussain, C., Gremeaux, V., Morisset, C., Laurent, Y., Casillas, J. M., & Laroche, D. (2013). Eccentric training in chronic heart failure: feasibility and functional effects. Results of a comparative study. *Annals of physical and rehabilitation medicine*, *56*(1), 30-40. https://doi.org/10.1016/j.rehab.2013.01.003

Borg, G., & Borg, E. (2001). A new generation of scaling methods: Level-anchored ratio scaling. *Psychologica*, 28(1), 15-45.

Borot, L., Vergotte, G., & Perrey, S. (2018). Different hemodynamic responses of the primary motor cortex accompanying eccentric and concentric movements: A functional NIRS study. *Brain Sciences*, *8*(5), 75. https://doi:10.3390/brainsci8050075

Brigadoi, S., & Cooper, R. J. (2015). How short is short? Optimum source–detector distance for short-separation channels in functional near-infrared spectroscopy. *Neurophotonics*, *2*(2), 025005. https://doi.org/10.1117/1.NPh.2.2.025005

Christou, E. A., & Carlton, L. G. (2002). Motor output is more variable during eccentric compared with concentric contractions. *Medicine and science in sports and exercise*, *34*(11), 1773-1778. http://doi.org/10.1097/00005768-200211000-00013

Clos, P., Laroche, D., Stapley, P. J., & Lepers, R. (2019). Neuromuscular and perceptual responses to sub-maximal eccentric cycling. *Frontiers in Physiology*, *10*, 354. https://doi.org/10.3389/fphys.2019.00354

Clos, P., & Lepers, R. (2020). Leg Muscle Activity and Perception of Effort before and after Four Short Sessions of Submaximal Eccentric Cycling. *International Journal of Environmental Research and Public Health*, *17*(21), 7702. https://doi.org/10.3390/ijerph17217702

Clos, P., Mater, A., Laroche, D., & Lepers, R. (2022). Concentric versus eccentric cycling at equal power output or effort perception: Neuromuscular alterations and muscle pain. *Scandinavian Journal of Medicine & Science in Sports*, *32*(1), 45-59. https://doi.org/10.1111/sms.14053

Clos, P., Mater, A., Legrand, H., Poirier, G., Ballay, Y., Martin, A., & Lepers, R. (2022). Corticospinal excitability is lower during eccentric than concentric cycling. *Frontiers in Physiology*, 392. https://doi.org/10.3389/fphys.2022.854824

Cohen, J. (1988). Statistical Power Analysis for the Behavioral Sciences (2nd ed.). Routledge. https://doi.org/10.4324/9780203771587

Cooper, R., Selb, J., Gagnon, L., Phillip, D., Schytz, H. W., Iversen, H. K., ... & Boas, D. A. (2012). A systematic comparison of motion artifact correction techniques for functional near-infrared spectroscopy. *Frontiers in neuroscience*, *6*, 147. https://doi.org/10.3389/fnins.2012.00147

Derosière, G., Billot, M., Ward, E. T., & Perrey, S. (2015). Adaptations of motor neural structures' activity to lapses in attention. *Cerebral cortex*, *25*(1), 66-74. https://doi.org/10.1093/cercor/bht206

Derosière, G., Mandrick, K., Dray, G., Ward, T. E., & Perrey, S. (2013). NIRS-measured prefrontal cortex activity in neuroergonomics: strengths and weaknesses. *Frontiers in human neuroscience*, *7*, 583 https://doi.org/10.3389/fnhum.2013.00583

Di Lorenzo, R., Pirazzoli, L., Blasi, A., Bulgarelli, C., Hakuno, Y., Minagawa, Y., & Brigadoi, S. (2019). Recommendations for motion correction of infant fNIRS data applicable to multiple data sets and acquisition systems. *NeuroImage*, *200*, 511-527. https://doi.org/10.1016/j.neuroimage.2019.06.056

Douglas, J., Pearson, S., Ross, A., & McGuigan, M. (2017). Chronic adaptations to eccentric training: a systematic review. *Sports Medicine*, *47*(5), 917-941. https://doi.org/10.1007/s40279-016-0628-4

Duchateau, J., & Enoka, R. M. (2008). Neural control of shortening and lengthening contractions: influence of task constraints. *The Journal of physiology*, *586*(24), 5853-5864. https://doi.org/10.1113/jphysiol.2008.160747

Duncan, A., Meek, J. H., Clemence, M., Elwell, C. E., Fallon, P., Tyszczuk, L., ... & Delpy, D. T. (1996). Measurement of cranial optical path length as a function of age using phase resolved near infrared spectroscopy. *Pediatric research*, *39*(5), 889-894. https://doi.org/10.1203/00006450-199605000-00025

Esfahlani, F. Z., Jo, Y., Faskowitz, J., Byrge, L., Kennedy, D. P., Sporns, O., & Betzel, R. F. (2020). High-amplitude cofluctuations in cortical activity drive functional connectivity. *Proceedings of the National Academy of Sciences*, *117*(45), 28393-28401. https://doi.org/10.1073/pnas.2005531117

Fang, Y., Siemionow, V., Sahgal, V., Xiong, F., & Yue, G. H. (2001). Greater movement-related cortical potential during human eccentric versus concentric muscle contractions. *Journal of Neurophysiology*, *86*(4), 1764-1772. https://doi.org/10.1152/jn.2001.86.4.1764

Fang, Y., Siemionow, V., Sahgal, V., Xiong, F., & Yue, G. H. (2004). Distinct brain activation patterns for human maximal voluntary eccentric and concentric muscle actions. *Brain research*, *1023*(2), 200-212. https://doi.org/10.1016/j.brainres.2004.07.035

Ferrari, M., & Quaresima, V. (2012). A brief review on the history of human functional near-infrared spectroscopy (fNIRS) development and fields of application. *Neuroimage*, *63*(2), 921-935. https://doi.org/10.1016/j.neuroimage.2012.03.049

Friston, K. J. (2011). Functional and effective connectivity: a review. *Brain connectivity*, *1*(1), 13-36. https://doi.org/10.1089/brain.2011.0008

Gandevia, S. C. (2001). Spinal and supraspinal factors in human muscle fatigue. *Physiological reviews.* https://doi.org/10.1152/physrev.2001.81.4.1725

Gandevia, S. C., Allen, G. M., Butler, J. E., & Taylor, J. L. (1996). Supraspinal factors in human muscle fatigue: evidence for suboptimal output from the motor cortex. *The Journal of physiology*, *490*(2), 529-536. https://doi.org/10.1113/jphysiol.1996.sp021164

Hardy, C.J., & Rejeski, W.J. (1989). Not What, but How One Feels: The Measurement of Affect during Exercise. *Journal of Sport & Exercise Psychology, 11*, 304-317. https://doi.org/10.1123/JSEP.11.3.304

Hart, S. G., & Staveland, L. E. (1988). Development of NASA-TLX (Task Load Index): Results of empirical and theoretical research. In *Advances in psychology* (Vol. 52, pp. 139-183). North-Holland. https://doi.org/10.1016/S0166-4115(08)62386-9

Hollander, D. B., Durand, R. J., Trynicki, J. L., Larock, D., Castracane, V. D., Hebert, E. P., & Kraemer, R. R. (2003). RPE, pain, and physiological adjustment to concentric and eccentric contractions. *Medicine & Science in Sports & Exercise*, *35*(6), 1017-1025. http://doi.org/10.1249/01.MSS.0000069749.13258.4E

Huppert, T. J., Diamond, S. G., Franceschini, M. A., & Boas, D. A. (2009). HomER: a review of time-series analysis methods for near-infrared spectroscopy of the brain. *Applied optics*, *48*(10), D280-D298. https://doi.org/10.1364/AO.48.00D280

Inzlicht, M., Shenhav, A., & Olivola, C. Y. (2018). The effort paradox: Effort is both costly and valued. *Trends in cognitive sciences*, *22*(4), 337-349. https://doi.org/10.1016/j.tics.2018.01.007

Kan, B., Speelman, C., & Nosaka, K. (2019). Cognitive demand of eccentric versus concentric cycling and its effects on post-exercise attention and vigilance. *European Journal of Applied Physiology*, *119*(7), 1599-1610. https://doi.org/10.1007/s00421-019-04149-y

Kocsis, L., Herman, P., & Eke, A. (2006). The modified Beer–Lambert law revisited. *Physics in Medicine & Biology*, *51*(5), N91. https://doi.org/10.1088/0031-9155/51/5/N02

Kwon, Y. H., & Park, J. W. (2011). Different cortical activation patterns during voluntary eccentric and concentric muscle contractions: an fMRI study. *NeuroRehabilitation*, *29*(3), 253-259. http://doi.org/10.3233/NRE-2011-0701

Lakens, D. (2013). Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANOVAs. *Frontiers in psychology*, 863. https://doi.org/10.3389/fpsyg.2013.00863

Laroche, D., Joussain, C., Espagnac, C., Morisset, C., Tordi, N., Gremeaux, V., & Casillas, J. M. (2013). Is it possible to individualize intensity of eccentric cycling exercise from perceived exertion on concentric test? *Archives of physical medicine and rehabilitation*, *94*(8), 1621-1627. https://doi.org/10.1016/j.apmr.2012.12.012

Lastayo, P. C., Reich, T. E., Urquhart, M., Hoppeler, H., & Lindstedt, S. L. (1999). Chronic eccentric exercise: improvements in muscle strength can occur with little demand for oxygen. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, *276*(2), R611-R615. https://doi.org/10.1152/ajpregu.1999.276.2.R611

Liao, L. D., Tsytsarev, V., Delgado-Martínez, I., Li, M. L., Erzurumlu, R., Vipin, A., ... & Thakor, N. V. (2013). Neurovascular coupling: in vivo optical techniques for functional brain imaging. *Biomedical engineering online*, *12*(1), 1-20. https://doi.org/10.1186/1475-925X-12-38

Liu, J. Z., Dai, T. H., Sahgal, V., Brown, R. W., & Yue, G. H. (2002). Nonlinear cortical modulation of muscle fatigue: a functional MRI study. *Brain research*, *957*(2), 320-329. https://doi.org/10.1016/S0006-8993(02)03665-X

Mandrick, K., Derosiere, G., Dray, G., Coulon, D., Micallef, J. P., & Perrey, S. (2013). Prefrontal cortex activity during motor tasks with additional mental load requiring attentional demand: a near-infrared spectroscopy study. *Neuroscience research*, *76*(3), 156-162. https://doi.org/10.1016/j.neures.2013.04.006

Molavi, B., & Dumont, G. A. (2012). Wavelet-based motion artifact removal for functional near-infrared spectroscopy. *Physiological measurement*, *33*(2), 259. https://doi.org/10.1088/0967-3334/33/2/259

Nippert, A. R., Biesecker, K. R., & Newman, E. A. (2018). Mechanisms mediating functional hyperemia in the brain. *The Neuroscientist*, *24*(1), 73-83. https://doi.org/10.1177/1073858417703033

Olsson, C. J., Hedlund, M., Sojka, P., Lundström, R., & Lindström, B. (2012). Increased prefrontal activity and reduced motor cortex activity during imagined eccentric compared to concentric muscle actions. *Frontiers in Human Neuroscience*, *6*, 255. https://doi.org/10.3389/fnhum.2012.00255

Overend, T. J., Versteegh, T. H., Thompson, E., Birmingham, T. B., & Vandervoort, A. A. (2000). Cardiovascular stress associated with concentric and eccentric isokinetic exercise in young and older adults. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, *55*(4), B177-B182. https://doi.org/10.1093/gerona/55.4.B177

Pageaux, B., Besson, D., Casillas, J. M., Lepers, R., Gremeaux, V., Ornetti, P., ... & Laroche, D. (2020). Progressively increasing the intensity of eccentric cycling over four training sessions: A feasibility study in coronary heart disease patients. *Annals of physical and rehabilitation medicine*, *63*(3), 241-244. https://doi.org/10.1016/j.rehab.2019.09.007

Pageaux, B. (2016). Perception of effort in Exercise Science: Definition, measurement and perspectives. *European Journal of Sport Science*, *16*(8), 885-894. https://doi.org/10.1080/17461391.2016.1188992

Peñailillo, L., Blazevich, A. J., & Nosaka, K. (2017). Factors contributing to lower metabolic demand of eccentric compared with concentric cycling. *Journal of Applied Physiology*, *123*(4), 884-893. https://doi.org/10.1152/japplphysiol.00536.2016

Peñailillo, L., Mackay, K., & Abbiss, C. R. (2018). Rating of perceived exertion during concentric and eccentric cycling: are we measuring effort or exertion? *International Journal of Sports Physiology and Performance*, *13*(4), 517-523. https://doi.org/10.1123/ijspp.2017-0171

Perrey, S. (2008). Non-invasive NIR spectroscopy of human brain function during exercise. *Methods*, *45*(4), 289-299. https://doi.org/10.1016/j.ymeth.2008.04.005

Perrey, S. (2018). Brain activation associated with eccentric movement: A narrative review of the literature. *European Journal of Sport Science*, *18*(1), 75-82. https://doi.org/10.1080/17461391.2017.1391334

Perrey, S., Betik, A., Candau, R., Rouillon, J. D., & Hughson, R. L. (2001). Comparison of oxygen uptake kinetics during concentric and eccentric cycle exercise. *Journal of Applied Physiology*, *91*(5), 2135-2142. https://doi.org/10.1152/jappl.2001.91.5.2135

Preston, J., & Wegner, D. M. (2009). Elbow grease: When action feels like work. In E. Morsella, J. A. Bargh, & P. M. Gollwitzer (Eds.), *Oxford handbook of human action* (pp. 569–586). Oxford University Press

Radel, R., Brisswalter, J., & Perrey, S. (2017). Saving mental effort to maintain physical effort: a shift of activity within the prefrontal cortex in anticipation of prolonged exercise. *Cognitive, Affective, & Behavioral Neuroscience*, *17*(2), 305-314. https://doi.org/10.3758/s13415-016-0480-x

Scholkmann, F., Spichtig, S., Muehlemann, T., & Wolf, M. (2010). How to detect and reduce movement artifacts in near-infrared imaging using moving standard deviation and spline interpolation. *Physiological measurement*, *31*(5), 649. https://doi.org/10.1088/0967-3334/31/5/004

Semmler, J. G., Tucker, K. J., Allen, T. J., & Proske, U. (2007). Eccentric exercise increases EMG amplitude and force fluctuations during submaximal contractions of elbow flexor muscles. *Journal of Applied Physiology*, *103*(3), 979-989. https://doi.org/10.1152/japplphysiol.01310.2006

Singh, A. K., Okamoto, M., Dan, H., Jurcak, V., & Dan, I. (2005). Spatial registration of multichannel multi-subject fNIRS data to MNI space without MRI. *Neuroimage*, *27*(4), 842-851. https://doi.org/10.1016/j.neuroimage.2005.05.019

Takahashi, T., Takikawa, Y., Kawagoe, R., Shibuya, S., Iwano, T., & Kitazawa, S. (2011). Influence of skin blood flow on near-infrared spectroscopy signals measured on the forehead during a verbal fluency task. *Neuroimage*, *57*(3), 991-1002. https://doi.org/10.1016/j.neuroimage.2011.05.012

Tachtsidis, I., & Scholkmann, F. (2016). False positives and false negatives in functional near-infrared spectroscopy: issues, challenges, and the way forward. *Neurophotonics*, *3*(3), 031405. https://doi.org/10.1117/1.NPh.3.3.031405

Taylor, J. L., Butler, J. E., Allen, G. M., & Gandevia, S. C. (1996). Changes in motor cortical excitability during human muscle fatigue. *The Journal of physiology*, *490*(2), 519-528. https://doi.org/10.1113/jphysiol.1996.sp021163

Themelis, G., D'Arceuil, H. E., Diamond, S. G., Thaker, S., Huppert, T. J., Boas, D. A., & Franceschini, M. A. (2007). Near-infrared spectroscopy measurement of the pulsatile component of cerebral blood flow and volume from arterial oscillations. *Journal of biomedical optics*, *12*(1), 014033. https://doi.org/10.1117/1.2710250

Vincent, J. L., Kahn, I., Snyder, A. Z., Raichle, M. E., & Buckner, R. L. (2008). Evidence for a frontoparietal control system revealed by intrinsic functional connectivity. *Journal of neurophysiology*, *100*(6), 3328-3342. https://doi.org/10.1152/jn.90355.2008

Xia, M., Wang, J., & He, Y. (2013). BrainNet Viewer: a network visualization tool for human brain connectomics. *PloS one*, *8*(7), e68910. https://doi.org/10.1371/journal.pone.0068910

Yao, W. X., Jiang, Z., Li, J., Jiang, C., Franlin, C. G., Lancaster, J. L., ... & Yue, G. H. (2016). Brain functional connectivity is different during voluntary concentric and eccentric muscle contraction. *Frontiers in physiology*, *7*, 521. https://doi.org/10.3389/fphys.2016.00521

Yao, W. X., Li, J., Jiang, Z., Gao, J. H., Franklin, C. G., Huang, Y., ... & Yue, G. H. (2014). Aging interferes central control mechanism for eccentric muscle contraction. *Frontiers in Aging Neuroscience*, *6*, 86. https://doi.org/10.3389/fnagi.2014.00086

Yücel, M. A., Selb, J., Aasted, C. M., Lin, P. Y., Borsook, D., Becerra, L., & Boas, D. A. (2016). Mayer waves reduce the accuracy of estimated hemodynamic response functions in functional near-infrared spectroscopy. *Biomedical optics express*, *7*(8), 3078-3088. https://doi.org/10.1364/BOE.7.003078

Yue, G. H., Liu, J. Z., Siemionow, V., Ranganathan, V. K., Ng, T. C., & Sahgal, V. (2000). Brain activation during human finger extension and flexion movements. *Brain Research*, *856*(1-2), 291-300. https://doi.org/10.1016/S0006-8993(99)02385-9

Figure 1 Overview of the experimental protocol. Four different 5-min cycling exercise conditions were realised in one session. In the first condition, participants performed concentric cycling at a fixed perception of effort (23 on the CR100® scale) without torque feedback. Then, the other three conditions were performed in a randomised order: eccentric cycling at a fixed perception of effort (23 on the CR100®), concentric or eccentric cycling at an intensity corresponding to the average torque produced in the first condition. Five minutes of rest were allowed between each exercise condition.

ROI	MNI			BA
	x	y	z	
PFC Left	-47	48	18	45-46
	-40	47	30	45-46
	-28	65	18	10-46
PFC Right	39	64	7	10-46
	54	44	7	45-46
	52	42	21	45-46
Motor Cortex	-38	-21	71	4-6
	-14	-2	75	6
	-1	-10	73	6
	12	0	75	6
	22	-10	75	6
	36	-20	73	4-6
PL Right	28	-77	56	7
	30	-85	44	7-19
	49	-70	48	7-39

Figure 2 The functional near infrared spectroscopy (fNIRS) multichannel layout with 15 channels covering four regions of interest (ROI): motor cortex (yellow), left (dark blue) and right (light blue) prefrontal cortex (PFC) and parietal lobe (PL red). On the left, spatial organization of different optodes (BrainNet viewer toolbox (Xia, Wang & He, 2013)); in black the receivers, in colour the emitters, and pink edge the channels. On the right, MNI coordinates to each fNIRS channel (ROI) with corresponding Brodmann areas (BA).

Figure 3 Mean (± standard deviation) and individual scatter for NASA-TLX mental demand (top) and physical demand (bottom) scores. There were main effects of the mode of muscle contraction (eccentric vs. concentric) and of the prescription of the exercise (fixed perceived effort -PoE- or torque) for the mental demand. There was a main effect of the mode of muscle contraction (eccentric vs. concentric) and an interaction between the mode of muscle contraction and the prescription of the exercise for the physical demand; the physical demand was greater for the torque condition in concentric cycling compared to eccentric cycling. * *p* < .05, ** *p* < .01, *** *p* < .001.

Figure 4 Mean (± standard deviation) and individual scatter for the area under the

curve (AUC) changes in oxyhaemoglobin (O₂Hb, red) and deoxyhaemoglobin (HHb,

blue) during eccentric (Ecc) and concentric (Conc) cycling. Left (A) and right (B) pre-frontal cortex (PFC), Motor cortex (C) and parietal lobe (PL, D). Main effect of the mode of muscle contraction (eccentric vs. concentric) at * $p < .05$, ** $p < .01$, *** $p < .001$.