# Eccentric cycling involves greater mental demand and cortical activation of the frontoparietal network

**Lénaic Borot[1,2], Benjamin Pageaux[3,4,5], Davy Laroche[6,7], Grégoire Vergotte[1], Romuald Lepers[6], Stéphane Perrey[1*]**

1.  *EuroMov Digital Health in Motion, Univ Montpellier, IMT Mines Ales, Montpellier, France*
2.  *Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, United Kingdom*
3.  *École de kinésiologie et des sciences de l'activité physique (EKSAP), Faculté de médecine, Montréal, Canada*
4.  *Centre de recherche de l'Institut universitaire de gériatrie de Montréal (CRIUGM), Montréal, Canada*
5.  *Centre interdisciplinaire de recherche sur le cerveau et l'apprentissage (CIRCA), Montréal, Canada.*
6.  *INSERM UMR 1093-CAPS, Univ. Bourgogne, UFR des Sciences du Sport, Dijon, France*
7.  *INSERM, CHU Dijon-Bourgogne, Centre d'Investigation Clinique CIC 1432, Module Plurithématique, Plateforme d'Investigation Technologique, Dijon, France*

**Short title:** Cortical control of eccentric cycling

**\*Corresponding author**:

Prof. Stephane Perrey (PhD)
EuroMov Digital Health in Motion
700 avenue du Pic Saint Loup
34090 Montpellier France
Tel. (+33) 4 34 43 26 23

Email: stephane.perrey@umontpellier.fr

**Abstract**

Eccentric, compared to concentric exercise, is proposed to involve different neuro-motor processing strategies and a higher level of mental demand. This study compared eccentric and concentric cycling at matched perceived effort and torque for the mental demand and related-cortical activation patterns. Nineteen men (30 ± 6 yrs) performed four different 5-min cycling conditions at 30 RPM on a semi-recumbent isokinetic cycle ergometer: 1) concentric at a moderate perceived effort (23 on the CR100® scale) without torque feedback; 2) concentric and 3) eccentric at the same average torque produced in the first condition; and 4) eccentric at the same moderate perceived effort than the first concentric condition. The conditions 2-4 were randomised. After each condition, mental demand was monitored using the NASA Task Load Index scale. Changes in oxy-($O_2$Hb) and deoxy-(HHb) haemoglobin during exercise were measured over both prefrontal cortices and the right parietal lobe from a 15-probe layout using a continuous-wave NIRS system. Mental demand was significantly higher during eccentric compared to concentric cycling (+52%, $p$ = .012) and when the exercise intensity was fixed by the torque rather than the perceived effort (+70%, $p$ < .001). For both torque- or perceived effort-matched exercises, $O_2$Hb increased significantly ($p$ < .001) in the left  and right prefrontal cortices, and right parietal lobe, and HHb decreased in the left, and right, prefrontal cortices during eccentric compared to concentric cycling. This study supports that acute eccentric cycling, compared to concentric cycling, involves a higher mental demand and frontoparietal network activation.

**Keywords:** muscle contraction, bilateral motor task, perceived effort, optical imaging, brain, cognitive demand

**INTRODUCTION**

Most human movements involve both eccentric and concentric muscle contractions (e.g., walking, running, jumping) and a large proportion of our daily living activities require accurate control of movements employing eccentric contractions (e.g., walking downstairs, sitting, lowering weight). During eccentric contractions, the external force on the muscle is greater than the force that the muscle can generate, producing a lengthening contraction; and vice versa for concentric contractions.[1] In the last decades, eccentric cycling at moderate intensity has been shown to promote functional (knee-extensor strength) and structural (hypertrophy) muscle adaptations in different populations due to its capability to achieve high loads on muscle tissue at low cardiorespiratory stress (athletic[2] and chronic diseases[3]).

Over the last 60 years, increasing research has been undertaken to understand mechanical, peripheral and neural features, as well as the metabolic cost of eccentric contractions. Eccentric cycling was initially introduced to investigate the physiology of concentric and eccentric muscle work[4]. Eccentric cycling requires substantially less oxygen uptake than concentric cycling to develop the same absolute external mechanical power output.[4,5] In addition, corticospinal excitability, electromyographic activity and rating of perceived effort are lower during eccentric compared to concentric cycling at the same absolute mechanical workload.[5-7] Notably, at a given metabolic workload, heart rate and rating of perceived effort were found higher in eccentric compared to concentric cycling.[5,6]

Eccentric cycling is associated with substantial effort perception and sustained attention when subjects are asked to match a given target mechanical power output.[8,9] However, little is known about cortical activity patterns of attentional resources involved

during eccentric cycling. Current literature contains several lines of evidence indicating

that neurobehavioral relationships in eccentric contractions are associated with higher

processing demands by various brain areas for controlling motor output.[10]

At the behavioural level, several studies have reported greater variability in the

motor output during eccentric compared to concentric contractions.[11] This increased

motor variability may indicate less practice and that eccentric contractions might be

more cognitively demanding than concentric contractions.[10] This hypothesis is

supported by the recent findings from Kan et al.[9] during submaximal eccentric and

concentric cycling. First, an overall higher perception of the workload, evaluated by the

NASA task load index (NASA-TLX) questionnaire, was reported during eccentric

compared to concentric cycling. This difference is specifically related to two items of

the questionnaire: mental demand and frustration. Second, the average dual-task

reaction time in assessing cognitive load was 15% slower, and the accuracy score was

7% lower, during eccentric compared to concentric cycling. Altogether, these findings

suggest that eccentric cycling is more taxing the brain than concentric cycling does.

A neuroimaging modality used in the study of Kan et al.[9] was functional near-

infrared spectroscopy (fNIRS). Functional NIRS is an optical neuroimaging technique

based on the theory of neurovascular coupling[12] and is relatively robust to motion

artifacts.[13] Since the local supply of oxygen is greater than its consumption in activated

brain regions, a higher concentration of oxy-haemoglobin ($O_2Hb$) and a decreased

concentration of deoxy-haemoglobin (HHb) are to be observed.[13] Changes in $O_2Hb$

and HHb recorded with 2 channels over the prefrontal cortex in Kan et al.[9] were found

to be smaller during two eccentric cycling exercises compared to a concentric cycling

exercise, while there were higher scores for the two eccentric cycling exercise

regarding the mental demand subscale of NASA-TLX. These preliminary findings

suggest that a reduced supply of oxygen compared to relative mental demand occurs

during eccentric cycling. However, they seem not to be supported by previous studies

comparing the influence of the mode of muscle contractions from isolated and small

muscle groups on brain activity.[10] Using electroencephalography during elbow flexion

at matched torque, a greater amplitude of movement-related cortical potentials in

eccentric compared to concentric contractions was shown for electrodes located over

the motor areas, prefrontal cortex and parietal lobe.[14] Using functional magnetic

resonance imagery (fMRI), Yao et al.[15] observed an increased blood-oxygen-level

dependent signal in similar brain areas during an eccentric task of the first dorsal

interosseous muscle as compared to a concentric task. Yet, none of the neuroimaging

studies mentioned above directly investigated cortical activity over multiple brain areas

during whole-body eccentric and concentric exercise, such as cycling.

According to the aforementioned studies, the control of eccentric contraction

would require important activation of the prefrontal regions, the right parietal lobe as

well as the motor regions. These different areas are part of the central executive

network known to be involved in cognitive-motor tasks modulating attentional

demand[16] as well as in the modulation of mental demand during continuous cycling

exercise.[17] As attention-demanding tasks exhibit increased activity in the prefrontal

cortex and posterior parietal cortex, we hypothesized that the greater mental demand

of eccentric cycling compared to concentric cycling would be associated with cortical

activity increases in bilateral prefrontal cortices and right posterior parietal cortex.

Finally, most of the aforementioned studies investigating eccentric cycling used an

experimental design imposing an absolute mechanical power output. Using effort

perception for matching the exercise intensity between eccentric and concentric

cycling can be an alternative to circumvent the comparison of neurophysiological responses, either at the same mechanical or at the same metabolic workload.[18] An effort perception approach to prescribe eccentric cycling has been proposed for the tailoring of rehabilitation programs in deconditioned populations, such as patients living with cardio-respiratory chronic diseases leading to muscle weakness.[3] Considering the gaps in the literature, the main objective of the present study was to quantify the acute changes of attention-related cortical activity as well as those of relevant brain motor areas from fNIRS during submaximal eccentric and concentric cycling exercise. As the method to prescribe exercise is likely to influence cortical activity and mental demand, eccentric and concentric cycling was performed at both matched perceived effort and torque.

**METHODS**

**Participants**

Nineteen healthy male adults (mean age: 30 ± 6 years) volunteered in the study. All participants reported no sign of neurological, respiratory and cardiovascular disease or medication, which might affect brain and muscle functions. Participants provided written informed consent prior to participation in the study. The study was approved by the local ethics committee (ID: RCB_2016-A01722-49) and was conducted in accordance with the ethical standards specified by the Declaration of Helsinki. This study is belonging to the ExoMode project registered on ClinicalTrials.gov (NCT03280875). The sample size required for the present study was calculated using G*Power (3.1.9.6). The fNIRS results between the concentric and eccentric cycling conditions in one previous study[9] were used as group parameters. The power analysis indicated

that a total of at least 17 participants was required for the brain activation (f = .85 – $np2= \sim.42$ , corresponding to large effect – ; α = .05; 1-β = .90). Accordingly, a sample size of 17 participants was recruited. Two additional participants were included in case of experimental outliers.

**Experimental procedures**

Participants were requested to visit the laboratory twice. The first visit consisted of a familiarization session and the second visit was the experimental session. Data collection occurred solely in the experimental session.

### *Familiarisation session*

The first visit to the laboratory was a familiarisation session with a double objective. First, participants were familiarised with the CR100® scale to evaluate their perception of effort and quadriceps pain, as well as to self-regulate the cycling exercise intensity based on a fixed perceived effort intensity. Second, the participants experimented with pedalling, driven by a motor either clockwise (concentric cycling) or anticlockwise (eccentric cycling), on an isokinetic semi-recumbent prototype cycle ergometer (Tech Med TMS, 89290 Champs-sur- Yonne, France). The seat of the semi-recumbent cycle ergometer allows for greater torso stabilization compared with a normal upright cycling position. This familiarisation cycling session consisted of participants understanding that they need to resist the pedals moving anticlockwise while they would have to accompany the pedals moving clockwise by exerting a suitable force to match a target line appearing on a screen in front of them. Participants were asked to perform eccentric and concentric cycling in a well-coordinated manner such that the knee extensors of each leg are activated at the right timing and intensity to ensure smooth pedal

rotation and torque production. This familiarization session was carried out with the

participant reaching submaximal torque values indicated by visual feedback on the

control screen. The visual feedback was adjusted to produce 75W. To ensure precise

execution of the concentric and eccentric isokinetic exercise, the researcher employed

manual palpation of the quadriceps as a mean of verification. Specifically, this involved

the researcher using manual palpation of the quadriceps to confirm that participants

were adequately familiarized with the task and successfully contracted their quadriceps

at the appropriate timing during pedal rotations. Finally, the participants performed four

submaximal cycling exercises at a moderate intensity for three minutes at 30 RPM.

The mode of muscle contraction (eccentric vs concentric) and the prescription of the

exercise (fixed perceived effort or torque) were manipulated as follow: 1) eccentric and

2) concentric cycling at a moderate effort intensity (corresponding to 23 on the CR100®

scale); 3) eccentric and 4) concentric cycling at a fixed torque value. The familiarization

session was carried out at least 48 hours before the experimental session.

***Recording session***

In this second session, participants performed four cycling exercise conditions for 5

min at 30 RPM (see Figure 1), as experienced in the familiarisation session. The choice

of the exercise – rest ratio of 5 min – 5 min is based on pilot testing as well as the

literature. First, as the literature demonstrated that 5 min of concentric or eccentric

cycling is sufficient to obtain a steady state in physiological responses[4,5], we decided

to use this duration for the cycling bouts. Second, we tested this ratio during a pilot

experiment in 5 participants to ensure that completion of our experimental protocol did

not induce a marked decrease in maximal voluntary isometric contraction peak torque

(MVIC) of the knee extensor (i.e., neuromuscular fatigue), a confound factor known to

increase perception of effort.[19,20] This pilot experiment consisted of duplicating the

experimental session without measuring neurophysiological variables. Before the test

session, MVIC of the dominant knee extensors was assessed (under verbal

encouragement) as the best performance in three 5-s trials separated by a 30-s resting

period. Then, one MVIC was performed in less than 20-s after the completion of the

test. MVIC was performed on an isokinetic dynamometer (System Pro 4, Biodex

Medical System, New York, NY) at a knee angle of 90°. We observed no marked

decrease in knee extensor MVIC before starting the protocol (pre) vs after (post)

completion of the protocol [MVC pre = 318 ± 77 Nm; MVC post = 318 ± 80 Nm; $p$ =

0.982. d (95%CI) = -0.011 (-0.887, 0.867). Finally, to control for potential muscle

damage at the onset of the experimental visit, we asked the participants at their arrival

to the laboratory for the presence of muscle pain in their lower leg at rest or while

walking, and no pain was reported.

In the first condition, participants performed concentric cycling at a fixed perception of

effort intensity corresponding to 23 on the CR100® scale (moderate intensity). No force

feedback was provided to the participants who were instructed to self-regulate their

force during each pedal revolution solely on their effort perception intensity, and to

keep it moderate during the 5-min of exercise. The maximal torque values produced

at each pedal revolution were extracted from the cycle ergometer every 15 s from

minutes 2 to 5 of the exercise. These values were then averaged over the last 3

minutes, excluding the last 15 s (i.e., when the participants were reporting their

perception), to individually determine the target torque used in the subsequent

conditions where exercise intensity was prescribed with a fixed torque. The same data

points were considered for analysis in all conditions. Then, the other three conditions

were performed in a randomised order: (i) concentric cycling at an intensity

corresponding to the average torque produced in the first condition, (ii) eccentric cycling at an intensity corresponding to the average torque produced in the first condition, (iii) eccentric cycling at a fixed perception of effort intensity corresponding to 23 on the CR100® scale (moderate, same perception of effort intensity than the first condition). Five minutes of rest were allowed between each exercise condition so that the physiological responses (heart rate and fNIRS signals) return to baseline levels. Heart rate was recorded with a chest belt and heart rate monitor (Polar V800, Finland) during rest and exercise periods for each condition. The average heart rate was obtained during the resting period before the onset of exercise and over the last minute of each exercise condition.

**Figure 1.** Overview of the experimental protocol. Four different 5-min cycling exercise conditions were realised in one session. In the first condition, participants performed concentric cycling at a fixed perception of effort (23 on the CR100® scale) without torque feedback. Then, the other three conditions were performed in a randomised order: eccentric cycling at a fixed perception of effort (23 on the CR100®), concentric or eccentric cycling at an intensity corresponding to the average torque produced in the first condition. Five minutes of rest were allowed between each exercise condition.

**Psychological variables**

Perception of effort, also known as perceived exertion or sense of effort, can be

described as a cognitive feeling of work associated with voluntary actions.[20]

Participants were asked to rate their perception of effort and quadriceps pain (CR100®

scale) and affective valence during the last 20 seconds of each condition. At the

beginning of the first sessions, participants were provided with validated instructions

to dissociate between perceptions of effort and muscle pain[21] and had the possibility

to ask questions to the researchers if clarification needed. In-task affective valence

(i.e., the pleasure/displeasure one feels) was measured using the Feeling scale, an

11-point bipolar scale ranging from -5 to +5. Seven anchors are provided ranging from

"very good" (+5) to "very bad" (-5). Finally, participants were asked to complete at the

end of each condition two subscales (mental demand and physical demand) of a

commonly used subjective workload level questionnaire, the NASA-TLX scale. For

each subscale, participants had to answer from 0 (very low or high) to 100 (very high

or low). The NASA TLX scale was provided to the participants immediately after

cessation of the 5-min cycling bouts. The NASA TLX scale was completed within 1 min

post cycling bout for all participants.

**Cortical activity**

*Recording*

A two-wavelength (765 and 856 nm) continuous-wave 15-channels fNIRS system

(Oxymon MkIII, Artinis Medical Systems, The Netherlands) was utilised to measure the

changes in $O_2Hb$ and HHb over regions of interest (ROI): left and right dorsolateral

posterior parietal cortex, right parietal lobe and motor regions on both hemispheres as

a whole. The $O_2Hb$ and HHb data were acquired during rest and exercise periods at a

sampling rate of 10 Hz. $O_2Hb$ and HHb concentration changes (expressed in $\Delta\mu M$)

obtained by using the modified Beer-Lambert law were displayed on a PC monitor in

real time. The optodes (5 receivers and 13 transmitters resulting in a total of 15

channels, Figure 2) were placed on the participants using a fit neoprene fNIRS head

cap to get a stable optical contact with the scalp by generating an equal amount of

pressure. The pressure was adequate to induce a partial transient blockage of the skin

circulation during the fNIRS study[22] as witnessed by the presence of well-defined

circles over the skin that disappears about 15 min after the test session. In the 15

channels, the measurement points were defined as the midpoint of the corresponding

illuminator-detector pairs interspaced by 3 cm. The Brodmann's areas covered by the

different channels were extracted via the NFRI function[23] from the MNI (Montreal

Neurological Institute) coordinates (Figure 2). Cartesian coordinates were obtained

with a Fastrak® 3D Digitizer (Polhemus Inc., USA)  allowing the localization of each

optode and channel from four anatomical reference points: nasion, right and left

mastoids and vertex.

| ROI | MNI | | | BA |
|---|---|---|---|---|
| | x | y | z | |
| | -47 | 48 | 18 | 45-46 |
| PFC Left | -40 | 47 | 30 | 45-46 |
| | -28 | 65 | 18 | 10-46 |
| | 39 | 64 | 7 | 10-46 |
| PFC Right | 54 | 44 | 7 | 45-46 |
| | 52 | 42 | 21 | 45-46 |
| | -38 | -21 | 71 | 4-6 |
| | -14 | -2 | 75 | 6 |
| Motor | -1 | -10 | 73 | 6 |
| Cortex | 12 | 0 | 75 | 6 |
| | 22 | -10 | 75 | 6 |
| | 36 | -20 | 73 | 4-6 |
| | 28 | -77 | 56 | 7 |
| PL Right | 30 | -85 | 44 | 7-19 |
| | 49 | -70 | 48 | 7-39 |

**Figure 2.** The functional near infrared spectroscopy (fNIRS) multichannel layout with

15 channels covering four regions of interest (ROI): motor cortex (yellow), left (dark

blue) and right (light blue) prefrontal cortex (PFC) and parietal lobe (PL red). On the

left, spatial organization of different optodes (BrainNet viewer toolbox[24]); in black the

receivers, in colour the emitters, and pink edge the channels. On the right, MNI

coordinates to each fNIRS channel (ROI) with corresponding Brodmann areas (BA).

***Data analysis***

Raw data (optical intensity) extracted from Oxysoft software (v3.0.75, Artinis Medical

Systems) was converted in optical density (OD) using Matlab® (MATLAB 2014b, The

MathWorks, USA). Then, to reduce possible head motion artifacts two methods

implemented in the Homer2 toolbox[25] were used. First, the moving standard deviation

and spline interpolation methods were applied using parameters: SDTresh = 20,

AMPTresh = 0.5, tMotion = 0.5s, tMask = 2s and p = 0.99. Second, we used wavelet-

based signal decomposition with the parameter: iqr = 0.1, as recommended.[9] Then OD

time series were converted into relative concentrations of $O_2Hb$ and HHb using the

modified Beer-Lambert law corrected by a differential pathlength factor depending on

the age of the participant.[26] A low-pass Butterworth zero-phase digital filter (order 4)

with a cut-off frequency of 0.08 Hz was applied to limit physiological artifacts (heart

rate: ± 1 Hz, respiration: ± 0.4 Hz and Mayer wave: ± 0.1 Hz). For each participant,

channels not having a sufficient signal-to-noise ratio were excluded from the analyses

after observation of the power spectrum density of the $O_2Hb$ signals where the

presence of a cardiac rhythm in the signal (peak around 1 Hz) indicates good contact

between the optodes and the scalp.[27] After this preprocessing step, 14% of the

channels were excluded.

Following a splitting of the fNIRS signals from events indicating the beginning of each

condition, the average between -30 seconds and -10 seconds of the minute of rest

preceding each condition was subtracted from the subsequent data during the exercise

period. The following dependent variables attesting to exercise-related cortical

activation magnitude were calculated per channel for the period (representative of the

steady-state level) between 2 minutes and 4 minutes 30 seconds of exercise for each

condition: area under the curve of $O_2Hb$ and HHb and average values. The average of

the channels positioned on 4 ROIs (the right and the left posterior parietal cortex and

primary motor, and right parietal lobe) was extracted for each participant and each

condition.

**Statistical analysis**

Statistical analyses were performed using Statistica (7.1, Statsoft Inc., France). Data

were examined for normality and sphericity using Shapiro-Wilk and Mauchly tests,

respectively. Where Mauchly's tests indicated violations of the sphericity assumption,

Greenhouse–Geisser corrections were applied. A one-way repeated-measure ANOVA

was used to assess the differences between the four exercise conditions for heart rate

recorded during the rest period. A fully repeated measures two-way ANOVA with the

factors *mode of muscle contraction* (eccentric vs. concentric) and *mode of exercise*

*prescription* (torque vs. perception of effort) were used to compare the NASA-TLX

scores and fNIRS responses. In case of significant effect, post hoc comparisons were

adjusted with the Bonferroni correction.

Wilcoxon signed-rank tests were performed for quadriceps pain and the Feeling scale

values when comparing the mode of muscle contraction and the mode of exercise

prescription. Student paired *t*-tests were carried out between the concentric and

eccentric torque conditions for the perceived effort. For the cycling torque data,

Wilcoxon signed-rank test was performed between the first condition (concentric at

fixed perceived effort) and the two conditions (eccentric and concentric) where the

intensity was fixed by a torque value.

Effect sizes were reported with the partial eta square ($\eta^2_p$) for the effects of the

ANOVAs[28] and with Cohen's *d* and rank-biserial correlation ($r_{rb}$) for pairwise

comparison. The significance level for all analyses was established at $p = .05$ (two-

tailed). All the fNIRS *p*-values were corrected using false discovery rate to control for

type 1 error due to the spatial dependency between brain areas.

**RESULTS**

**Torque and heart rate**

The average torque during concentric cycling at a moderate perceived effort was 41.1

± 14.1 Nm. No difference was observed between the average torque produced when

performing concentric cycling at a moderate perceived effort, and the average torque

produced when performing concentric (42.7 ± 13.8 Nm; W (17) = 47.5; p = .17; rrb =

.33) or eccentric cycling at a fixed torque (40.3 ± 13.12 Nm; W (17) = 40; p = .08; rrb =

.42).

No difference in resting heart rate was observed before the start of each cycling exer-

cise condition ($F_{(3, 45)} = 1.25$; $p = .30$; $\eta^2_p = .08$). During the cycling exercises, the

main effect of the mode of muscle contraction revealed a higher heart rate during con-

centric cycling compared to eccentric cycling ($F_{(1, 18)} = 40.3$; $p < .001$; $\eta^2_p = .69$, 101

± 13 vs. 87 ± 13 bpm). No main effect of the mode of exercise prescription ($F_{(1, 18)} =$

.09; $p = .77$; $\eta^2_p = .00$; perception of effort: 94 ± 14 bpm; torque: 94 ± 15 bpm) or mode

of muscle contraction × mode of exercise prescription interaction ($F$ (1, 18) = .88; $p$ =

.36; $\eta^2_p$ = .05) was observed.

**Psychological variables**

When the exercise was prescribed with a fixed torque feedback, the *perception of effort*

*intensity* was higher ($t$ (17) = 3.39; $p$ < .01; $d$ = 1.07) during concentric cycling com-

pared to eccentric cycling (23.2 ± 3.9 and 17.4 ± 6.8).

No main effects of the mode of muscle contraction ($W$ (38) = 96.5; $p$ = .75; $r_{rb}$ = .05)

and prescription of intensity ($W$ (38) = 53; $p$ = .44; $r_{rb}$ = .13) were found for perceived

quadriceps pain during the exercise as well as for the affects (Feeling scale) ($W$ (38)

= 94; $p$ = .68; $r_{rb}$ = .07 and $W$ (38) = 79; $p$ = .12; $r_{rb}$ = .25, respectively).

For the perceived measures with the NASA-TLX subscales, one participant did not

complete the questionnaire at the end of the experimental session. Therefore, data are

available for 18 participants. For the *mental demand scores*, the main effect of the

mode of muscle contraction revealed a higher mental demand during eccentric cycling

compared to concentric cycling ($F$ (1, 17) = 7.86, $p$ = .01; $\eta^2_p$ = .32; Figure 3a). The

main effect of the mode of exercise prescription revealed a higher mental demand

when the exercise was prescribed with a fixed torque compared to a fixed perception

effort ($F$ (1, 17) = 26.22; $p$ < .001; $\eta^2_p$ = .61; Figure 3b). No mode of muscle contraction

× mode of exercise prescription interaction was observed ($F$ (1, 17) = .45; $p$ = .51; $\eta^2_p$

= .02). For the *physical demand scores*, the main effect of the mode of muscle

contraction revealed a higher physical demand during concentric cycling compared to

eccentric cycling ($F$ (1, 17) = 6.66; $p$ = .02; $\eta^2_p$ = .28; Figure 3c). No main effect of the

mode of exercise prescription was observed ($F$ (1, 17) = .54; $p$ = .47; $\eta^2_p$ = .03;

observed power = .11). Follow-up tests on the mode of muscle contraction × mode of

exercise prescription interaction ($F$ (1, 17) = 6.99; $p$ = .02, $\eta^2_p$ = .29) are presented in Figure 3d. The physical demand was greater for the torque condition in concentric cycling compared to eccentric cycling ($p$ = .005, $d$ = 0.315).

**Figure 3.** Perceived mental and physical demand scores measured with the NASA-TLX scale. The mental demand was perceived higher during eccentric cycling compared to concentric cycling (main effect of the muscle contraction mode, panel a), and during the exercise prescribed with the torque feedback compared to the perception of effort (main effect of the exercise prescription mode, panel b). The physical demand was perceived lower during eccentric cycling compared to concentric cycling (main effect the muscle contraction mode, panel c), and the interaction effect revealed a

specific lower perceived physical demand during the eccentric cycling compared to

concentric cycling when exercise was prescribed with the torque feedback. Concentric

cycling and eccentric cycling are represented with circle and square markers, respec-

tively. Individual (n=18) data are represented with empty markers and mean (±95% CI)

as filled markers. * $p < .05$, ** $p < .01$, *** $p < .001$.

**Cortical responses**

For the *left prefrontal cortex* (Figure 4A) and $O_2Hb$, the main effect of the mode

of muscle contraction revealed a higher area under the curve during eccentric cycling

compared to concentric cycling ($F (1, 18) = 16.35$; $p < .01$, $\eta^2_p = .48$). Neither the main

effect of the mode of exercise prescription method ($F (1, 18) = .1$; $p = .75$; $\eta^2_p = .01$),

nor a mode of muscle contraction × mode of exercise prescription interaction ($F (1, 18)$

$= 1.48$; $p = .24$; $\eta^2_p = .08$) was observed. Regarding HHb, the main effect of the mode

of muscle contraction revealed a lower area under the curve during eccentric cycling

compared to concentric cycling ($F (1, 18) = 35.77$, $p < .001$, $\eta^2_p = .67$). Neither the main

effect of the mode of exercise prescription ($F (1, 18) = .49$, $p = .49$; $\eta^2_p = .03$) nor a

mode of muscle contraction × mode of exercise prescription interaction ($F (1, 18) =$

$.07$; $p = .80$; $\eta^2_p = .003$) was observed.

For the *right prefrontal cortex* (Figure 4B) and $O_2Hb$, the main effect of the mode

of muscle contraction revealed a higher area under the curve during eccentric cycling

compared to concentric cycling ($F (1, 18) = 6.96$; $p < .05$; $\eta^2_p = .28$). Neither the main

effect of the mode of exercise prescription ($F (1,18) = .30$; $p = .59$; $\eta^2_p = .02$) nor a

mode of muscle contraction × mode of exercise prescription interaction ($F (1, 18) =$

$2.06$; $p = .17$; $\eta^2_p = .10$) was observed. Regarding HHb, the main effect of the mode of

muscle contraction revealed a lower area under the curve during eccentric cycling

compared to concentric cycling ($F$ (1, 18) = 22.26, $p$ < .001, $\eta^2_p$ = .55). Neither the main

effect of the mode of exercise prescription ($F$ (1,18) = .63, $p$ = .44; $\eta^2_p$ = .03) nor a

mode of muscle contraction × mode of exercise prescription interaction ($F$ (1, 18) =

3.15, $p$ = .09; $\eta^2_p$ = .15) was observed. For bilateral *motor cortex* and $O_2Hb$ (Figure

4C), no main effect of the mode of muscle contraction was observed ($F$ (1,18) = 1.01,

$p$ = .33; $\eta^2_p$ = .05). Neither the main effect of the mode of exercise prescription ($F$ (1,18)

= .18, $p$ = .68; $\eta^2_p$ = .01) nor a mode of muscle contraction × mode of exercise

prescription interaction ($F$ (1, 18) = .37, $p$ = .55; $\eta^2_p$ = .02) was observed. Regarding

HHb, the main effect of the mode of muscle contraction revealed a lower area under

the curve during eccentric cycling compared to concentric cycling ($F$ (1, 18) = 38.26, $p$

< .001, $\eta^2_p$ = .68). Neither the main effect of the mode of exercise prescription ($F$ (1,

18) = .73, $p$ = .40; $\eta^2_p$ = .04) nor a mode of muscle contraction × mode of exercise

prescription interaction ($F$ = (1, 18) = 3.42, $p$ = .08; $\eta^2_p$ = .16) was observed.

For the *right parietal lobe* (Figure 4D) and $O_2Hb$, the main effect of the mode of

muscle contraction revealed a higher area under the curve during eccentric cycling

compared to concentric cycling ($F$ (1, 13) = 11.12, $p$ = .02, $\eta^2_p$ = .46). Neither the main

effect of the mode of exercise prescription method ($F$ (1,13) = 1.09, $p$ = .32; $\eta^2_p$ = .08;

Figure 4D) nor a mode of muscle contraction × mode of exercise prescription

interaction ($F$ (1, 13) = .20, $p$ = .66; $\eta^2_p$ = .01) was observed. Regarding HHb, no main

effect of the mode of muscle contraction was observed ($F$ (1, 13) = .38, $p$ = .55; $\eta^2_p$ =

.03). Neither the main effect of the mode of exercise prescription ($F$ (1, 13) = 2.2, $p$ =

.16; $\eta^2_p$ = .14 nor a mode of muscle contraction × mode of exercise prescription

interaction ($F$ (1, 13) = 1.33, $p$ = .27; $\eta^2_p$ = .09) was observed.

**Figure 4.** Changes in the area under the curve (AUC) for the oxyhaemoglobin (O2Hb, red) and deoxyhaemoglobin (HHb, blue). Data are presented for the main effect of the muscle contraction mode. The four regions of interest were the left prefrontal cortex (N=19, panel a), the right prefrontal cortex (N=19, panel b), the motor cortex (N=19, panel c) and the right parietal lobe (n=14, panel d). Data missing for the right parietal lobe is due to a damaged optical fiber occurring during one day of the data collection. Concentric cycling and eccentric cycling are represented with circle and square markers, respectively. Individual data are represented with empty markers and mean (±95% CI) as filled markers. * *p* < .05, ** *p* < .01, *** *p* < .001.

## DISCUSSION

The main objective of the present study was to quantify the acute changes in attention-related cortical activity as well as those relevant to brain motor areas from fNIRS during eccentric and concentric cycling exercises performed at both matched perceived effort (23 on CR100® scale) and torque. The main results showed a greater mental demand

and cortical activation in the right and left prefrontal cortices and right parietal lobe during eccentric cycling compared to concentric cycling. These findings confirm previous results suggesting that eccentric cycling is more cognitively demanding than concentric cycling[9] and extend them by identifying specific brain activation patterns underlying this greater mental demand.

**Mental demand during eccentric cycling**

Mental demand was subjectively measured with the NASA-TLX subscale. The mental demand was reported to be 45% greater during eccentric cycling compared to concentric cycling, while physical demand was 38% greater during concentric cycling. These findings support the results of Kan et al.[9] Nevertheless, mental and physical demand values were 2 to 3 times lower in the present study. The higher mental and physical demand found in Kan et al.[9] is likely due to a greater exercise intensity compared to the present study, as confirmed by the higher mean heart rate values reported in their study as compared to ours (eccentric cycling: ~+29 bpm; concentric cycling: +56 bpm). While mental demand refers to the mental and perceptual activity imposed by a task and refers to related information processing, effort refers to the voluntary engagement of mental and physical resources to perform, or attempt to perform, in a task. [20,29] In the present study, coordinating the leg movements properly during eccentric and concentric cycling required dedicated motor skills and so adequate demand for attention. As described in the methods, participants were asked to perform eccentric and concentric cycling in a well-coordinated manner to ensure smooth pedal rotation and torque production. This was similar to the cycling exercises proposed by Kan et al.[9], where the authors observed that attention and vigilance were maintained after eccentric and concentric cycling. However, higher cognitive load

(based on both the NASA TLX score and choice reaction time) was found in the study

of Kan et al.[9] during eccentric cycling. Altogether, it appears that eccentric cycling is

more cognitively demanding than concentric cycling and that the less mental demand

of eccentric and concentric cycling in the present study did not likely change cognitive

functions post-exercise.

Regarding the mode of exercise prescription, the mental demand was

significantly higher when the intensity was set by a target torque value compared to an

exercise prescribed by a fixed perceived effort. To the best of our knowledge, this study

is the first to observe this difference in mental demand. A lower mental demand during

self-regulated exercise based on effort perception is not surprising as, in this specific

condition, participants did not have at each pedal revolution to visually focus on a

torque target to reach. Contrary to the mental demand, the mode of exercise

prescription did not alter the reported physical demand of the cycling exercise. This

observation is in line with no significant difference in heart rate between the two modes

of exercise prescription.

**Cortical activity during eccentric cycling**

Neuroimaging studies using various methods and exercise paradigms reveal

inconsistencies in the role of the primary motor cortex in the production of eccentric

muscle contractions.[10] Yao et al.[15] during unilateral movements of the first dorsal

interosseous muscle in an MRI scanner, as Fang et al.[14] in an EEG study for the elbow

flexors, showed a greater activation of the primary motor cortex contralateral to the

active muscles when performing eccentric compared to concentric contractions.

Conversely, a reduction was reported in the primary motor cortex blood-oxygen-level

dependent signal amplitude during eccentric compared to concentric contractions.[30]

The present study is the first to collect, during an eccentric cycling exercise (i.e., a bilateral task), fNIRS data over the precentral gyrus (BA 4/6) and primary motor areas (BA 4). The present study did not observe significant differences in motor cortices activity between eccentric and concentric cycling. Our results can be explained because the tasks performed are fundamentally different from the literature. First, the different studies cited above performed unilateral distal movements of the upper limb, unlike our study where the participants performed a bilateral task mobilizing larger muscle masses of the lower limbs. In addition, in the present study, the cycling exercise was performed at moderate intensity (23 on the CR100® scale or using an associated torque). Studies showing a greater activity of the contralateral primary motor during eccentric muscle contractions involved exercises performed either at maximum intensity[14] or sub-maximum intensity fixed relative to maximal voluntary contraction force or body weight.[14,15] Noteworthy, in the study of Olsson et al.[30] a continuous imagery task paradigm was used by alternating concentric (lifting a weight) and eccentric (slowing down a weight) muscle contractions. For other cortical areas involved in the motor network (e.g. primary and secondary motor areas not recorded here), a consensus seems to exist on the greater cortical activity observed during eccentric contractions.[10]

The main results of the present study revealed a greater activity of the left and right prefrontal cortices, as well as the right parietal lobe, when performing eccentric cycling compared to concentric cycling, regardless of the mode of exercise prescription. These observations likely unveil underlying neurophysiological mechanisms sustaining the increased mental demand reported by the participants during eccentric cycling. The left and right prefrontal cortices as well as the right parietal lobe are known for their roles in the modulation of mental demand during prolonged exercise[17] as well

as during sustained attentional cognitive tasks.[16] When focusing on a specific move-

ment to perform, several brain areas, constituting the central executive network, are

engaged. More specifically, these regions would be part of the fronto-parietal network

that consists mainly of the dorsolateral prefrontal cortex, the frontal cortex and the pa-

rietal lobe.[31] An increase in eccentric force variability has been proposed as a sign of

an increased motor control difficulty during eccentric contractions compared to con-

centric contractions.[11] Controlling the variability in motor output (not assessed in the

present study) during eccentric contractions would likely require greater attentional de-

mand, thus explaining the observed increase in the activity of the prefrontal cortices

and the right parietal lobe. Some authors have highlighted the complementarity of re-

gional brain activity magnitude measures and those linked to the dynamic activity of

brain networks.[32] On this basis, the fMRI study of Yao et al.[33] examined the strength of

functional connectivity between the primary motor cortex and other cortical regions in

the motor control network (e.g. supplementary motor area, cerebellum) involved in ec-

centric and concentric contractions. The authors observed a weaker functional con-

nectivity during eccentric compared to concentric contractions of the right first dorsal

interosseous muscle. A weaker statistical dependence between two-time series from

different brain areas may reflect more involvement of specialised regions (segregation)

and thus less functional integration.[34] This means that higher cortical activation may

be required during eccentric contraction to compensate for a weaker functional con-

nectivity. Yao et al.[33] hypothesized that the brain control of a concentric contraction will

be more efficient than an eccentric contraction. This first result needs to be supported

by more evidence, notably for a bilateral task mobilizing larger muscle masses of the

lower limbs. The analysis of functional connectivity could be a relevant and comple-

mentary approach to better understand the functional organisation of the brain when

performing eccentric cycling over time and its possible implications for rehabilitation

and training.

**Methodological considerations**

In the present study, we controlled for the induction of muscle fatigue by providing a

sufficient recovery period between cycling exercise conditions, thus allowing us to

ensure that our results are not confounded by a decrease in force production capacity

of the working muscles with changes in the corticospinal excitability.[35] Exercise-

induced muscle fatigue plays an important role in the modulation of cortical activity

during exercise. In an fMRI study involving a maximal-effort handgrip contraction, Liu

et al.[36] observed an early increase followed by a decrease in brain activity in the

primary motor cortex and in particular in the prefrontal cortex. The exercise intensity in

our experimental design was low as indicated by perceived effort and heart rate values.

In addition, our control experiment showed no significant decrease in the maximal force

production capacity of the knee extensors after the whole exercise protocol.

The present study faces one limitation, which might be taken into account in follow-up

studies. The fNIRS signal can be contaminated by the physiological responses to the

task performed (e.g. heart rate, blood pressure). This signal contamination related to

non-neural activity can lead to false positives or negatives in our analyses.[37] To over-

come these issues, the use of short-distance channels (approximately 1 cm between

the transmitters and the receivers) was proposed.[38] In this study, we did not use such

short-distance channels for separating signals that stem from blood flow changes in

the extracerebral layers (scalp blood flow) as well as the cerebral tissue layer (systemic

blood flow). As indicated in the methods, we minimized this potential effect by inducing

a partial transient blockade of the skin blood flow during the fNIRS data collection.[22]

Interestingly, heart rate was ~13% higher in concentric as reported before[39], whereas

the greatest cortical activation was observed during eccentric cycling. These elements

allow us to think that the results might not be influenced by systemic interference but

future investigations are necessary to confirm this hypothesis.

Finally, it is important to note that 2-3 participants did not report a higher mental de-

mand during eccentric cycling compared to concentric cycling. These interindividual

variability responses may be due to the differences in the degree of expertise of our

participants with eccentric cycling. Therefore, future studies should explore the link

between the repeated and chronic use of eccentric cycling and a potential decrease in

its mental demand. Regarding the variability in fNIRS responses, 3-5 participants pre-

sented results contradictory to the group. In line with this observed variability, we

strongly encourage future researchers to increase their sample size beyond the one

used in our present study to replicate and confirm our observations.

**CONCLUSION**

In the present study we took advantage of using functional near infrared spectroscopy

covering four regions of interest to investigate how mental demand induced by

eccentric exercise modulated the central executive network. The present study showed

that the higher mental demand observed during eccentric cycling compared to

concentric cycling was associated with a greater activation of the left and right

prefrontal cortices as well as the right parietal lobe. These findings provide evidence

that heightened activation of the frontoparietal network may support eccentric

movement control.

**PERSPECTIVES**

A large proportion of our daily living activities require accurate control of movements employing eccentric contractions (e.g. walking downstairs, sitting). We encourage future studies to replicate and extend our results using different locomotion modalities such as uphill or downhill running, which are known to predominantly involve concentric and eccentric contractions of the knee extensors[40], respectively. As physiological responses to physical exercise are well known to be linked to the exercise intensity, it could also be of interest to explore the interaction between the exercise intensity and mental demand during various exercise modalities. Based on the findings of this study, future studies should investigate the potential use of wearable fNIRS for clinical and sports applications via an online neurofeedback approach to modulate the mental demand. Further investigations are also warranted to examine the impact of task familiarization on mental demand and the related brain activation during eccentric cycling. As learning is associated with increased automaticity and decreased resources engaged in a task, it is plausible that participants will experience lower mental demand when they become acquainted with the task. Another application is that self-regulated eccentric exercise based on perceived effort is less mentally demanding, suggesting that it may be more appropriate if one session needs to minimize the overall workload. Overall, eccentric cycling workload can be prescribed based on the individual's perceived rate of exertion, but also at a brain activation workload similar to that elicited by concentric cycling to improve both neuro-motor processing strategies and mental demand.

**ACKNOWLEDGEMENT**

This research work was supported by the French National Research Agency (EXOMODE–ANR–15–CE19-0023). We thank the anonymous reviewers for their careful reading of our manuscript and their many insightful comments and suggestions.

**AUTHOR CONTRIBUTIONS**

Lenaic Borot: Conceptualization; data curation; formal analysis; investigation; methodology; validation; visualization; writing –original draft; writing –review and editing. Benjamin Pageaux: Conceptualization; investigation; methodology; visualization; validation; writing –review and editing. Davy Laroche: Conceptualization; funding acquisition; methodology; resources; supervision; writing –review and editing. Gregoire Vergotte: methodology; formal analysis; writing –review and editing. Romuald Lepers and Stephane Perrey: Conceptualization; investigation; funding acquisition; methodology; project administration; resources; supervision; validation; writing – review and editing.

**CONFLICT OF INTEREST**

All authors declare that they have no conflicts of interest.

# REFERENCES

1. Duchateau J, Enoka RM. Neural control of shortening and lengthening contractions: influence of task constraints. *J Physiol (Lond)*. 2008;586:5853-5864.

2. Douglas J, Pearson S, Ross A, McGuigan M. Chronic adaptations to eccentric training: a systematic review. *Sports Med.* 2017;47:917-941.

3. Pageaux B, Besson D, Casillas JM, Lepers R, Gremeaux V, Ornetti P, Laroche D. Progressively increasing the intensity of eccentric cycling over four training sessions: A feasibility study in coronary heart disease patients. *Ann Phys Rehab Med*. 2020;63:241-244.

4. Abbott BC, Bigland B, Ritchie JM. The physiological cost of negative work. *J Physiol (Lond).* 1952;117:380.

5. Perrey S, Betik A, Candau R, Rouillon JD, Hughson RL. Comparison of oxygen uptake kinetics during concentric and eccentric cycle exercise. *J Appl Physiol.* 2001;91:2135-2142

6. Clos P, Mater A, Laroche D, Lepers R. Concentric versus eccentric cycling at equal power output or effort perception: Neuromuscular alterations and muscle pain. *Scand J Med Sci Sports.* 2022;32:45-59.

7. Peñailillo L, Mackay K, Abbiss CR. Rating of perceived exertion during concentric and eccentric cycling: are we measuring effort or exertion? *Int J Sports Physiol Perf.* 2018;13:517-523.

8. Peñailillo L, Blazevich AJ, Nosaka K. Factors contributing to lower metabolic demand of eccentric compared with concentric cycling. *J Appl Physiol*. 2017;123:884-893.

9. Kan B, Speelman C, Nosaka, K. Cognitive demand of eccentric versus concentric cycling and its effects on post-exercise attention and vigilance. *Eur J Appl Physiol.* 2019;119:1599-1610.

10. Perrey S. Brain activation associated with eccentric movement: A narrative review of the literature. *Eur J Sport Sci.* 2018;18:75-82.

11. Christou EA, Carlton LG. Motor output is more variable during eccentric compared with concentric contractions. *Med Sci Sports Exerc.* 2002;34:1773-1778.

12. Liao LD, Tsytsarev V, Delgado-Martínez I, Li ML, Erzurumlu R, Vipin A, Thakor NV. Neurovascular coupling: in vivo optical techniques for functional brain imaging. *Biomed Eng Online.* 2013;*2*:1-20

13. Perrey S. Non-invasive NIR spectroscopy of human brain function during exercise. *Methods*. 2008;45: 289-299.

14. Fang Y, Siemionow V, Sahga, V, Xiong, F, Yue GH. Distinct brain activation patterns for human maximal voluntary eccentric and concentric muscle actions. *Brain Res*. 2004;1023: 200-212.

15. Yao WX, Li J, Jiang Z, Gao JH, Franklin CG, Huang Y, Yue GH. Aging interferes central control mechanism for eccentric muscle contraction. *Front Aging Neurosci*. 2014;6:86.

16. Derosière G, Billot M, Ward ET, Perrey S. (2015). Adaptations of motor neural structures' activity to lapses in attention. *Cereb Cortex*. 2015;25:66-74.

17. Radel R, Brisswalter J, Perrey S. Saving mental effort to maintain physical effort: a shift of activity within the prefrontal cortex in anticipation of prolonged exercise. *Cogn Affect Behav Neurosci*. 2017;17(2):305-314

18. Besson D, Joussain C, Gremeaux V, Morisset C, Laurent Y, Casillas JM, Laroche D. Eccentric training in chronic heart failure: feasibility and functional effects. Results of a comparative study. *Ann Phys Rehab Med*. 2013;56:30-40.

19. Pageaux B, Lepers R. Fatigue Induced by Physical and Mental Exertion Increases Perception of Effort and Impairs Subsequent Endurance Performance. *Front Physiol*. 2016;7:587.

20. Pageaux B (2016). Perception of effort in Exercise Science: Definition, measurement and perspectives. *Eur J Sport Sci*. 2016;16:885-894.

21. Pageaux B, Clos P, Impellizzeri F, Audiffren M, Stapley P, Grémeaux V, Perrey, S, O'Connor P, Marcora S, Lepers R, Laroche D. (2020). Towards Standardized instructions for measuring perception of effort and muscle pain during physical exercise. *Med Sci Sports Exerc*. 2020;52:499

22. Takahashi T, Takikawa Y, Kawagoe R, Shibuya S, Iwano T, Kitazawa S. Influence of skin blood flow on near-infrared spectroscopy signals measured on the forehead during a verbal fluency task. *Neuroimage*. 2011;57:991-1002.

23. Singh AK, Okamoto M, Dan H, Jurcak V, Dan I. Spatial registration of multichannel multi-subject fNIRS data to MNI space without MRI. *Neuroimage*. 2005;27:842-851.

24. Xia M, Wang J, He Y. BrainNet Viewer: a network visualization tool for human brain connectomics. *PloS one*. 2013;8:e68910.

25. Huppert TJ, Diamond SG, Franceschini MA, Boas D.A. HomER: a review of time-series analysis methods for near-infrared spectroscopy of the brain. *Appl Opt.* 2009;48:D280-D298

26. Duncan A, Meek JH, Clemence M, Elwell CE, Fallon P, Tyszczuk L, Delpy DT. Measurement of cranial optical path length as a function of age using phase resolved near infrared spectroscopy. *Pediatric Res.* 1996;39:889-894.

27. Themelis G, D'Arceuil HE, Diamond SG, Thaker S, Huppert TJ, Boas DA, Franceschini MA. Near-infrared spectroscopy measurement of the pulsatile component of cerebral blood flow and volume from arterial oscillations. *J Biomed Opt.* 2007;12:014033.

28. Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANOVAs. *Front Psychol.* 2013;4:863.

29. Inzlicht M, Shenhav A, Olivola CY. The effort paradox: Effort is both costly and valued. *Trends Cogn Sci.* 2018;22: 337-349.

30. Olsson CJ, Hedlund M, Sojka P, Lundström R, Lindström B. Increased prefrontal activity and reduced motor cortex activity during imagined eccentric compared to concentric muscle actions. *Front Hum Neurosci.* 2012;6:255.

31. Vincent JL, Kahn I, Snyder AZ, Raichle ME, Buckner RL. Evidence for a frontoparietal control system revealed by intrinsic functional connectivity. *J Neurophysiol.* 2008;100: 3328-3342.

32. Esfahlani FZ, Jo Y, Faskowitz J, Byrge L, Kennedy DP, Sporns O, Betzel RF. High-amplitude cofluctuations in cortical activity drive functional connectivity. *Proc Natl Acad Sci.* 2020;117:28393-28401.

33. Yao WX, Jiang Z, Li J, Jiang C, Franlin CG, Lancaster JL, Yue GH. Brain functional connectivity is different during voluntary concentric and eccentric muscle contraction. *Front Physiol.* 2016;7: 521.

34. Friston KJ (2011). Functional and effective connectivity: a review. *Brain Connect.* 2011;1:13-36.

35. Gandevia SC, Allen GM, Butler JE, Taylor JL. Supraspinal factors in human muscle fatigue: evidence for suboptimal output from the motor cortex. *J Physiol (Lond).* 1996;490: 529-536

36. Liu JZ, Dai TH, Sahgal V, Brown RW, Yue GH. Nonlinear cortical modulation of muscle fatigue: a functional MRI study. *Brain Res.* 2002;957:320-329.

37. Tachtsidis I, Scholkmann F False positives and false negatives in functional near-infrared spectroscopy: issues, challenges, and the way forward. *Neurophotonics*, 2016;3:031405.

38. Brigadoi S, Cooper RJ. How short is short? Optimum source–detector distance for short-separation channels in functional near-infrared spectroscopy. *Neurophotonics*, 2015;2:025005.

39. Hollander DB, Durand RJ, Trynicki JL, Larock D, Castracane VD, Hebert EP, Kraemer RR. RPE, pain, and physiological adjustment to concentric and eccentric contractions. *Med Sci Sports Exerc*. 2003;35:1017-1025.

40. Garnier YM, Lepers R, Dubau Q, Pageaux B, Paizis C. Neuromuscular and perceptual responses to moderate-intensity incline, level and decline treadmill exercise. *Eur J Appl Physiol*, 2018;118:2039-2053.