Microprism-based two-photon imaging of the mouse inferior colliculus reveals novel

organizational principles of the auditory midbrain.

Baher A. Ibrahim[1,2], Yoshitaka Shinagawa[1, 2], Austin Douglas[4], Gang Xiao[1,2,3], Alexander R.

Asilador[2,3,4], & Daniel A. Llano[1,2,3,4,5]

1- Department of Molecular & Integrative Physiology, University of Illinois at Urbana-

Champaign, IL, USA.

2- Beckman Institute for Advanced Science & Technology, University of Illinois at Urbana-

Champaign, IL, USA.

3- Neuroscience Program, University of Illinois at Urbana-Champaign, IL, USA.

4- School of Molecular & Cell Biology, University of Illinois at Urbana-Champaign, IL, USA.

5- Carle Illinois College of Medicine, University of Illinois at Urbana-Champaign, United States,

IL, USA.

Abstract:

To navigate real-world listening conditions, the auditory system relies on the integration of

multiple sources of information. However, to avoid inappropriate cross-talk between inputs,

highly connected neural systems need to strike a balance between integration and segregation.

Here, we develop a novel approach to examine how repeated neurochemical modules in the

mouse inferior colliculus lateral cortex (LC) allow controlled integration of its multimodal inputs.

The LC had been impossible to study via imaging because it is buried in a sulcus. Therefore we

coupled two-photon microscopy with the use of a microprism to reveal the first-ever sagittal

views of the LC to examine neuronal responses with respect to its neurochemical motifs under

anesthetized and awake conditions. This approach revealed marked differences in the acoustic

response properties of LC and neighboring nonlemniscal portions of the inferior colliculus. In

addition, we observed that the module and matrix cellular motifs of the LC displayed distinct

somatosensory and auditory responses. Specifically, neurons in modules demonstrated

primarily offset responses to acoustic stimuli with enhancement in responses to bimodal stimuli,

whereas matrix neurons showed onset response to acoustic stimuli and suppressed responses

to bimodal stimulation. Thus, this new approach revealed that the repeated structural motifs of

the LC permit functional integration of multimodal inputs while retaining distinct response

properties.

Introduction:

Real-world listening occurs in environments cluttered by noise. To navigate these conditions, the auditory system relies on the integration of multiple sources of input. Broad integration of neural inputs, however, comes at a potential cost. Dense convergence can lead to inappropriate cross-talk between input streams, generating a processing "soup" whereby individual neural pathways cannot be independently modulated. Therefore, highly connected neural systems need to preserve the specificity of inputs in the face of massive integration (Fair et al., 2007; Sporns, Tononi, & Edelman, 2000; Tononi, Sporns, & Edelman, 1994)

Several brain systems have solved the integration/segregation problem by creating repeated structural motifs that allow independent modulation of different pathways. For example, in the basal ganglia, neurochemical periodicity of the striosome and matrix regions permits the retention of functional independence of input streams in the face of massive convergence. As a result, basal ganglia direct and indirect pathways may be independently modulated, while allowing controlled crosstalk between channels (Alexander, Crutcher, & DeLong, 1990; Brimblecombe & Cragg, 2017; Draganski et al., 2008; Gerfen, 1985)

In the auditory system, the integration/segregation challenge is particularly formidable in the inferior colliculus (IC). The IC serves as a specialized integration hub because of its heavy innervation by higher auditory centers as well as by nonauditory regions. Thus, the IC stands at the crossroads between multiple brain regions and is thus important for linking acoustic stimuli with non-acoustic inputs. The IC comprises a lemniscal division, the central nucleus (ICC), that receives primarily ascending auditory projections (Oliver, 2005; Willard & Ryugo, 1983; Willott, 2001) and two non-lemniscal divisions known as the dorsal (DC) and lateral (LC) cortices that receive massive descending auditory as well as non-auditory input (Bajo & Moore, 2005; Coleman & Clerici, 1987; Loftus, Malmierca, Bishop, & Oliver, 2008; Saldaña, Feliciano, &

Mugnaini, 1996; Schreiner & Winer, 2005; Winer, Larue, Diehl, & Hefti, 1998) as reviewed in (A.

M. H. Lesicko & Geffen, 2022).

The LC and DC appear to differ in their strategy for integrating these inputs. The DC does not

contain known structural heterogeneities but has a tonotopic arrangement of neuronal

responses on its surface (Barnstedt, Keating, Weissenberger, King, & Dahmen, 2015; A. B.

Wong & Borst, 2019). In contrast, the LC contains repeated neurochemical motifs, here referred

to as modules, that express the inhibitory neurotransmitter GABA as well as other metabolic

markers (Chernock, Larue, & Winer, 2004). Areas outside the modules, referred to as matrix,

stain strongly for calretinin (Dillingham, Gay, Behrooz, & Gabriele, 2017). We and others have

determined that nearly all of the long- and short-range connectivity of the LC is governed by the

module/matrix organization (Dillingham et al., 2017; Lamb-Echegaray, Noftz, Stinson, &

Gabriele, 2019; A. M. H. Lesicko, Sons, & Llano, 2020; A. M. Lesicko, Hristova, Maigler, &

Llano, 2016).

Unfortunately, how the LC integrates these inputs while maintaining distinct paths of information

flow is not known. For example, LC neurons respond to multisensory stimuli (Jain & Shore,

2006; Zhou & Shore, 2006), but information from different modalities enters the LC in

segregated channels (A. M. Lesicko et al., 2016). Thus, integration across the module/matrix

border must occur, but the mechanisms of this integration are unknown. Traditional methods of

studying the LC, such as in vivo electrophysiology, pose significant challenges in

unambiguously assigning individually recorded neurons to residence in matrix or modules.

Imaging approaches, such as two-photon (2P) microscopy, have proven to be enormously

valuable in understanding the fine spatial structure of neural response properties in the auditory

system (Bandyopadhyay, Shamma, & Kanold, 2010; Romero et al., 2020; Rothschild, Nelken, &

Mizrahi, 2010). However, because of its superficial location on the dorsal surface of the mouse

midbrain, the DC is the only part of the IC that has been characterized by 2P imaging (Barnstedt

et al., 2015; Ito, Hirose, Murase, & Ikeda, 2014; A. B. Wong & Borst, 2019). Optically, it has

been impossible to characterize the functions of the LC, which is laterally embedded deep in a

brain sulcus.

Here, we describe a novel approach using a 45^o microprism inserted into the sulcus that

separates the LC from the cerebellum to investigate the fine-scale functional organization of the

LC from its lateral surface through 2P imaging. The 2P images obtained from the microprism

revealed the first-ever images of the LC in vivo showing its characteristic GABAergic modules.

Compared to the DC imaged from the surface, the acoustically and somatosensory-evoked

calcium signals obtained by the 2P imaging of the LC through the microprism revealed

functional distinctions between the different cellular motifs of the LC. For instance, while

modules were more responsive to somatosensory stimulation than matrix, the latter was more

acoustically-responsive and had a lower detection threshold for complex sounds. Additionally,

modules and matrix were found to differently process the spectral and temporal features of the

acoustic information. These findings suggest that the non-lemniscal divisions of the IC contain

rich but distinct platforms for the processing of different features of auditory and non-auditory

information.

Results:

The insertion of 45^o microprism for imaging the LC from the side.

GAD67-GFP knock-in mice on a Swiss Webster background were crossed with Thy1-

jRGECO1a mice to produce a GAD67-GFPx jRGECO1a hybrid mouse, which was used to

visualize GABAergic modules, to identify the dense network of GABAergic neurons and

terminals in modules of the LC in the green channel (Tamamaki et al., 2003), and the calcium

indicator, jRGECO1a, in the red channel (Dana et al., 2018). A standard craniotomy was

performed (Goldey et al., 2014) with modifications to suit the specific characteristics of the IC in

terms of its location, depth, and high density of the surrounding vasculature (Xiong et al., 2017).

The craniotomy (~ 3 x 4 mm) did not drill directly into the sagittal and lambdoid sutures which

allowed minimal disruption of the superficial sinuses below the skull (Figure 1A) as indicated by

the minimal bleeding near the bone flap (Figure 1B). During drilling, pressurized air and a cold

stream of saline were used to remove the bone debris and dissipate heat. Because the IC sits

deeper in the brain compared to the surrounding structures, the direct insertion of the 45°

microprism was found to push the IC ventrally. Therefore, a fine diamond knife was used to

separate the lateral surface of the IC from the surrounding tissues without damaging the

transverse and sigmoid sinuses (Figures 1C and D). This step was followed by a gentle

retraction of the knife medially to generate a pocket to secure the microprism (Figure 1E). The

successful insertion of the microprism was assessed by examining the reflection of the diamond

knife from the microprism's top glass surface (Figures 1F and G). Holding the microprism down

and retracting the knife upward simultaneously resulted in securing the microprism in its pocket

(Figure 1H and I). Locking the microprism down in its pocket for 10 minutes with the diamond

knife (Figure 1J) is critical to ensure that the microprism is stabilized and to prevent bleeding.

Filling the gap between the skull surface and the microprism with saline was found to generate

motion artifact due to the heart pulsation, so 1% agarose was used to fill the gap to reduce the

motion artifact and add more stability to the microprism (Figure 1K). To compare response

properties between the LC and DC, direct imaging (i.e., without a microprism) of the dorsal

surface of the IC, where the DC is located, was also conducted by scanning the DC surface

from the most medial to lateral regions following the convex curvature of the surface (Figures 1L

and M).

The 2P characterization of GABAergic and non-GABAergic neuronal distribution on the

LC and DC surfaces with histological validation.

Following the surgery, the anesthetized animals were taken immediately for imaging.

Examination of the distribution of GABAergic modules in the LC was critical to validate the

target area for functional calcium imaging. Therefore, the 920 nm laser beam was initially used

to scan for GFP, which is expressed in the GABAergic cells of GAD67-GFP knock-in mice. The

insertion of the microprism was initially viewed by low magnification fluorescence imaging of the

GFP signals from the whole surface of the IC along with the inserted microprism (Figure 2A).

This step validated the location of the microprism relative to the IC under the microscope and

ensured there was no bleeding that could block the field of view. The 2P images of the GFP

signals from the LC imaged via the microprism, or for simplicity LC$_{(microprism)}$, across different

animals revealed the LC characteristic GABAergic modules (Figures 2C and E, the irregular

white lines), which were consistent with previous 3D reconstructions of modules from the

coronal sections (Chernock et al., 2004; A. M. Lesicko et al., 2016). To further validate the

microprism location and the ability to image the LC from the lateral surface, a laser lesion (see

Methods) was created via the microprism (Figure 2G), which was validated histologically (Figure

2H). For comparison, the whole dorsal surface of the IC was scanned from medial to lateral

borders (Figure 2B, the dotted white lines). Unlike the LC, most of the dorsal surface of the IC

showed a non-clustered distribution of the GABAergic cells (Figures 2D and F, blue boxes).

However, the lateral horizon of the IC dorsal surface showed clusters of GABAergic cells and

terminals that were confirmed to be modules, which comprise the prominent feature of the LC

(Figures 2D and F, red boxes). To further validate the presence of GABAergic modules in the

dorsolateral surface of the IC, a laser lesion was created in a GABAergic module located in that

area (Figures 2I and J, dotted red circle), which was confirmed histologically (Figure 2K),

demonstrating that it is possible to image a portion of the LC directly from its dorsal surface.

Using the 1040 nm laser, the GAD67-GFPx jRGECO1a hybrid mouse showed a non-clustered

cellular expression of jRGECO1a on the surface of the DC (Figure 3A) or on the surface of the

LC$_{(microprism)}$ (Figure 3B), as expected. The IC of anesthetized animals in each experimental

setup was then imaged to measure calcium signals over the course of pseudorandom

presentations of either 500 ms pure tones of different combinations of frequencies (5-40 kHz,

half octave intervals) and sound levels (40-80 dB SPL, 10 dB intervals) or broadband noise

without or with amplitude modulation (AM-noise) of different combinations of AM frequencies (2-

256 Hz, one-octave intervals) and sound levels (40-80 dB SPL, 10 dB intervals). As shown by

two representative cells from the DC imaged from the dorsal surface or LC imaged via

microprism, these acoustic stimulation paradigms successfully elicited transient cellular calcium

signals that varied depending on the stimulus type (Figures 3C and D). The expression

distributions of jRGECO1a were histologically examined for the 2P imaged areas of the DC and

LC (Figure 3-figure supplement 1A). The DC and LC, have a similar fraction of cells expressing

jRGECO1a (paired sample t-Test, $t_{(9)}$ = 0.98, p = 0.34, n = 10 coronal sections from 2 animals)

(Figure 3-figure supplement 1B and C). Also, the cells expressing jRGECO1a in all regions of

the IC were only nonGABAergic neurons as they did not express GFP which is expressed

exclusively in the GABAergic neurons (Figure 3-figure supplement 1D).

Acoustic responsiveness differs in the LC and DC:

As indicated by the GFP signals and consistent with Figure 2, imaging the LC of the hybrid mice

via microprism showed clusters of GABAergic modules, which take different irregular shapes on

the surface of the LC (Figure 4A - irregular white lines). Based on the best pure tone frequency

(BTF) of each responsive cell (Barnstedt et al., 2015), an inconsistent and fragmentary

tonotopic organization was seen over the LC$_{(microprism)}$ (Figure 4B) as indicated by a low R^2 of the

best linear and quadratic regressions fit made between the BTFs of cells and their locations

(Figure 4 - figure supplement 1A). Additionally, the best fit of these regressions was found to be

along different axes across the tested animals (Figure 4 - figure supplement 1A). Despite this

inconsistent topographical organization, it was found that the cells in the modules were tuned to

relatively higher frequencies than those in the matrix (Mann-Whitney Test: z = 4.03, p = 5.6x10[-]

[5], a median of best-tuned frequency = 10 and 7.1 Hz for modules (215 cells from 7 animals) and

matrix, (230 cells from 7 animals) respectively).

For comparison, the surface of the DC was imaged directly from the dorsal surface using the

routine imaging setup. To view the majority of the DC surface, imaging was done by scanning

the DC from its medial to the lateral horizon to generate a composite map (see Methods). As

indicated by the GFP signals, it was noted that the most lateral region of the surface of the IC

showed clusters of the GABAergic cells and terminals (i.e., modules, Figure 4C – irregular white

lines) and are therefore considered part of the LC. Based on the BTF of each responsive cell

(Barnstedt et al., 2015), the DC showed a tonotopic organization of cells along the rostromedial

to caudolateral axis tuned to lower or higher frequencies, respectively, (Figure 4D) as indicated

by a high R^2 of the regression fits made between BTFs of cells and their locations along the

rostromedial to caudolateral axis. (Figure 4- figure supplement 1B). It was noted that the R^2 of

the nonlinear quadratic regression was higher than that of the linear fit, which allowed that

model to include the cells tuned to higher frequencies at the rostromedial part of the DC, which

is consistent with the tonotopic organization that was previously described (Barnstedt et al.,

2015; A. B. Wong & Borst, 2019). It was also noticed that a large area of the DC was tuned to

the lowest tested frequency (5kHz), suggesting a low-frequency bias on the surface of the DC.

To quantify tuning differences between DC and LC, the cumulative distribution function based

on the cells' BTFs showed a low-frequency bias in the DC, similar to previous work (Ito et al.,

2014; Romand & Ehret, 1990; Shen, Xu, & Shen, 2003; Stiebler & Ehret, 1985). Although

around 50% of the cells of $LC_{(microprism)}$ showed also a low-frequency bias to 5 and 7.1 kHz,

collectively, most of the LC cells were tuned to relatively higher frequencies than those of the

DC (Figure 5A, Mann-Whitney Test: z = -11.34, p < 0.001).

Detection of AM is critical to identify natural sounds as well as to aggregate sound sources

during auditory scene analysis, which is important for communication (Joris, Schreiner, & Rees,

2004; Rees & Langner, 2005; Singh & Theunissen, 2003). The IC also is the first site in the

ascending auditory pathway where a substantial number of neurons are rate-tuned to AM

(Frisina, 2001; Joris et al., 2004; Langner, 1992; Nelson & Carney, 2007; Rabang et al., 2012).

Even though calcium indicators do not provide a spike-by-spike readout of neural activity, their

integrated signal does correlate with the number of spikes produced by a neuron over short time

windows (Vogelstein et al., 2010; Yaksi & Friedrich, 2006). Thus, this integrated signal can be

used to measure rate responses to AM. Using this integrated signal, we tested the ability of the

cells of the LC $_{(microprism)}$ to respond to broadband noise at different AM rates within each cellular

motif. Within the fraction of responsive cells, the LC$_{(microprism)}$ showed cells of different response

profiles. While some cells responded only to pure tones or tone-selective (Tone-sel), other cells

responded only to AM-noise or noise-selective (Noise-sel). However, some cells responded to

both pure tones and AM-noise (Non-sel), when the two stimuli were individually presented.

While the LC$_{(microprism)}$ had similar fractions of Noise-sel and Non-sel cells, both fractions were

significantly higher than that of the Tone-sel cells (Figure 5B, one-way-ANOVA, p = 1.6×10^{-4}).

We also observed that modules and matrix differentially process the spectral information. For

instance, the matrix had a significantly greater fraction of Noise-sel cells compared to the

modules (Figure 5C, Mann-Whitney Test, z = -2.32, p = 0.02). In contrast, both cellular motifs

had similar fractions of Tone-sel and Non-sel cells (Figure 5C, Mann-Whitney Test: z = 0.48, p =

0.62 for Tone-sel and z = 1.36, p = 0.17 for Non-sel). These findings suggest that neurons in the

matrix are more responsive to sounds of greater spectral and temporal complexity than those in

the modules. Similar to the LC, the DC showed significantly higher fractions of Noise-sel and

Non-sel cells than Tone-sel cells, but the cellular fraction of Non-sel cells was higher than that of

Noise-sel cells (Figure 5D, one way-ANOVA, $f_{(2,9)}$ = 86.9, p = 1.2×10^{-6}).

By pooling the cells based on their response to either pure tones or AM-noise, animals showed

a lower fraction of responsive neurons to the acoustic stimulation in the LC $_{(microprism)}$ compared

to the DC regardless of the sound type (two-way ANOVA, $f_{(1,1,18)}$ = 9.1, 7.8, and 0.1, p = 0.007,

0.01, and 0.75 for region (LC vs DC), stimulus (pure tones vs AM-noise), and interaction,

respectively, n = 7 animals for LC and 4 animals for DC, the % of responsive cells ± SEM =

48±7% for LC and 75±2% for DC). To evaluate if any optical artifacts due to imaging through a

prism could be responsible for this finding, the number of responsive cells imaged from the

lateral-most dorsal surface of the IC, which we consider part of the LC because of the presence

of GABAergic modules (LC imaged from the dorsal surface or LC$_{(dorsal surface)}$), was quantified and

compared with that obtained from the LC$_{(microprism)}$. We observed that there were no significant

differences between the number of tone or noise-responsive neurons between the LC$_{(microprism)}$ or

LC$_{(dorsal surface)}$ (two-way ANOVA, $f_{(1,1, 18)}$ = 0.07, 14.1, and 2.11, p = 0.79, 0.001, and 0.16 for the

region, stimulus (pure tones vs AM-noise), and interaction, respectively, n = 7 animals for

LC$_{(microprism)}$ and 4 animals for LC$_{(dorsal surface)}$, the % of responsive cells ± SEM = 48±7% for

LC$_{(microprism)}$ and 45±11% for LC$_{(dorsal surface)}$).

LC and DC respond differently to spectral and temporal modulation of acoustic inputs:

Given that more cells from LC $_{(microprism)}$ and DC had responses to AM-noise compared to pure

tones, we further quantified the selectivity to spectral or temporal modulation by computing the

spectral modulation index (SMI) or the temporal modulation index (TMI), respectively, as well as

the number of modulated cells for all the responsive cells (see Methods) across the tested

sound levels.

For the LC$_{(microprism)}$, the matrix has a greater proportion of spectrally modulated cells (cells

responding only to pure tones and unmodulated noise) than modules. The higher fractions of

modulated cells in matrix were most prominently at sound levels below 70 dB (Figure 5E, two-

way ANOVA: p = $7.1x10^{-7}$, 1.0, and 0.01 for the region, sound level, and interaction,

respectively) suggesting that the cells in the matrix have a lower threshold to detect the spectral

features of sound. Similarly, the matrix had more cells responsive to temporal modulation (cells

responding to AM-noise) as a main effect (Figure 5F, two-way ANOVA, p = 2.7×10^{-4}, 1.0, and

0.32 for the region, sound level, and interaction, respectively), which was consistent with the

above results (Figure 5C).

The DC and LC imaged via microprism were found to have different profiles of modulation of

responses to spectral and temporal complexity. Although there is a general trend for SMI and

TMI to decrease with increasing sound-pressure level, corresponding to a decrease in the

response to pure-tone and AM-noise, respectively, overall there is a statistically-significant

higher level of average SMI and TMI in the LC compared to the DC (Figures 5G and H, SMI:

two-way ANOVA, p < 0.001 for all region, sound level, and interaction, TMI: two-way ANOVA, p

< 0.001 for all regions, sound level, and interactions). These results suggested that neurons in

the LC are more sensitive to spectral and temporal modulation than those in the DC.

LC modules and matrix differentially respond to acoustic and somatosensory stimuli:

Somatosensory inputs from many cortical and subcortical structures have been identified in the

non-lemniscal division of the IC, particularly in the LC (A. M. Lesicko et al., 2016; Lohse,

Zimmer-Harwood, Dahmen, & King, 2022; Zhou & Shore, 2006) Previous work has shown that

while somatosensory projections mostly target the modules, the auditory projections tend to

target the matrix (A. M. Lesicko et al., 2016), which indicates that the LC is a multisensory

integration site. Therefore, we examined the somatosensory and auditory cellular responses on

the surface of the LC by coupling the 2P imaging with the microprism with a whisker or acoustic

stimulation (see Methods). Acoustic stimulation was done with 500 ms of unmodulated

broadband noise at 80 dB SPL. After the acoustic or somatosensory stimulation, evoked cellular

responses were observed from the neurons of the LC$_{(microprism)}$ as well as the DC imaged directly

from the dorsal surface. According to these cellular responses, different types of neurons were

found in modules and matrix of the $LC_{(microprism)}$ as well as the DC (Figure 6A-D). Also, the

auditory response evoked by auditory stimulation only was found to be modulated (suppressed

vs enhanced) when the auditory stimulation was simultaneously presented with somatosensory

stimulation (bimodal stimulation) (Figure 6E and F). The cells of every structure were then

mapped according to their responses. Auditory and somatosensory responses were evoked in

the neurons of both matrix and modules imaged via microprism (Figures 7A, B, and D). Within

the pool of responsive cells, the cells were categorized into cells that respond only to auditory

stimulation [auditory selective cells (Aud-sel)], only to somatosensory stimulation

[somatosensory selective cells (Som-sel)], or nonselectively to both stimuli (Aud/Som-nonsel)]

(Figure 7F). Both modules and matrix showed a higher fraction of Aud-sel cells than that of

Som-sel and Aud/Som-nonsel cells (Figure 7H, Chi-Square test, χ^2 = 33.1, p = 3.5x10^{-6}).

Pooling the cells based on their auditory vs somatosensory responses revealed that more

auditory-responsive cells (Aud-sel + Aud/Som-nonsel) were found in the matrix, while more

somatosensory-responsive cells (Som-sel + Aud/Som-nonsel) were clustered in the modules

(Figures 7B, D, F, and I, Chi-Square test, χ^2 = 26.0, p = 9.3x10^{-5}), which was consistent with the

reported difference in inputs to modules compared to matrix (A. M. Lesicko et al., 2016).

Additionally, while most of the acoustically-responsive cells in the matrix showed onset

responses, the cells showing acoustically-driven offset responses were clustered in the modules

(Figures 7C and J, Chi-Square test, χ^2 = 120.5, p <0.001). Moreover, a higher fraction of

Aud/Som-nonsel cells in the modules had offset, not onset, auditory response than those

located in the matrix (Chi-Square test, χ^2 = 13.2, p = 0.004, % of Aud/Som-nonsel cells = 57%,

15%, 8%, and 20%, for auditory offset cells in modules, auditory onset cells in modules, auditory

offset cells in matrix, and auditory onset cells in the matrix, respectively), which suggests a

multisensory integrative function by those cells as discussed later. Given the specific spatial

location of the somatosensory-responsive cells in the modules which contain clusters of

GABAergic cells, we asked if somatosensory stimulation could suppress the auditory response

when both stimuli are presented simultaneously (bimodal stimulation). To test this hypothesis,

we calculated and visualized the response index (RI), which was calculated as the ratio of the

cellular response to bimodal stimulation to the cellular response to acoustic response alone for

auditory-responsive cells. Accordingly, some cells showed suppressed (RI < 1) or enhanced

(RI > 1) auditory responses in both modules and matrix following bimodal stimulation. We

observed that matrix and modules have more cells with suppressed auditory responses than

those having enhanced auditory responses. However, matrix had a higher fraction of cells with

suppressed responses than modules, while modules had a higher fraction of cells with

enhanced responses (Figures 7G and K, Chi-Square test, χ^2 = 12.9, p = 0.04).

Consistent with the above data, the LC$_{(dorsal\ surface)}$ (Figures 8A-D, dorsolateral white rectangle)

shared common response profiles with the LC$_{(microprism)}$. For instance, both regions had a higher

percentage of somatosensory responsive cells than the DC (Figures 8A-D, dorsal yellow and

dorsomedial blue rectangles) as quantitatively shown (Figure 8H, one-way ANOVA, p = 0.02),

which suggests that the LC is the main region of the IC shell that processes somatosensory

information, consistent with the anatomical reports (Aitkin, Dickhaus, Schult, & Zimmermann,

1978; Jain & Shore, 2006; A. M. Lesicko et al., 2016; Liu et al., 2023). Similar to the LC$_{(microprism)}$,

the matrix and modules imaged from the dorsal surface had more percentage of Aud-sel cells

than Som-sel and Aud/Som-nonsel cells (Figures 8E and I, Chi-Square test, χ^2 = 86.1, p

<0.001). Collectively, the matrix had more auditory responsive cells than modules, while

modules had more somatosensory responsive cells than matrix (Figure 8J, Chi-Square test, χ^2 =

66.5, p <0.001). Additionally, more cells with onset auditory response were found in the matrix,

while the cells with offset auditory responses were clustered in the modules (Figures 8F and K,

Chi-Square test, χ^2 = 500.2, p <0.001).

Consistent with the LC$_{(microprism)}$, some cells had suppressed or enhanced auditory responses

following bimodal stimulation (Figure 8-figure supplement 1). The matrix imaged from the dorsal

surface showed also more cells with suppressed auditory responses than those having

enhancing auditory responses induced by bimodal stimulation. Additionally, the matrix had a

higher fraction of cells with suppressed responses than modules, while modules had a higher

fraction of cells with enhanced responses than matrix. In contrast to the LC$_{(microprism)}$, the modules

imaged directly from the dorsal surface had a higher fraction of cells with enhanced responses

than those with suppressed responses, which may indicate that the ratio between suppressed

and enhanced responses in modules could be altered based on the dorsoventral location

(Figures 8G and L, Chi-Square test, χ^2 = 18.5, p = 1.6X10^{-5}). In general, these data confirmed

the findings obtained from the LC imaged via microprism and emphasized the functional

difference between the LC and DC.

LC and DC differ in their acoustic and multimodal response properties in awake animals:

To determine if the presence of ketamine/xylazine anesthesia influenced the response

properties detailed above, additional experiments were performed in awake animals (n = 6 total,

3 for the LC$_{(microprism)}$ and 3 for the DC). Similar to what was observed under anesthesia, the cells

of the LC$_{(microprism)}$ did not show clear evidence of tonotopy (Figure 9B), which was indicated by a

low R^2 of the regressions fit between the BTFs of the cells and their locations over different

anatomical axes (Figure 9-figure supplement 1A). Similarly, the cells of the modules were tuned

to relatively higher frequencies than those in matrix (Mann-Whitney Test, z = 3.8, p = 1.0 x 10^{-4},

median of best frequency = 10 vs 7.1 kHz for modules (148 cells from 3 animals) vs matrix (287

cells from 3 animals)). Additionally, the data obtained from imaging the DC under awake

preparations showed a similar tonotopic map (Figure 9D) as that observed under anesthesia

indicated by a high R^2 of the regression fit between the BTFs of the cells and their locations over

the rostromedial and caudolateral axis (Figure 9-figure supplement 1B). Yet, a greater

proportion of high-frequency cells were visually observed in the mediodorsal portion of the DC,

which was indicated by a higher R^2 of the nonlinear quadratic fit than that of the linear fit,

leading to the observation of a narrow low-frequency zone similar to the previous work under

awake preparations (A. B. Wong & Borst, 2019). The $LC_{(microprism)}$ had also more cells tuned to

higher frequencies compared to the DC (Figure 10A, Mann-Whitney test, $p=1.6 \times 10^{-6}$). We also

examined the width of the receptive field (RF) of the DC vs LC neurons at each tested sound

level under anesthetized and awake preparations using the binarized receptive field sum (RFS)

method (Bowen, Winkowski, & Kanold, 2020). Both DC and LC of the anesthetized animals had

higher RFS mean and median than those of awake animals given that ketamine was known to

broaden the RF (Guo et al., 2012). In both preparations (anesthetized and awake), the DC had

a higher RFS mean than that of the LC (Figure 10-figure supplement 1, Kruskal-Wallis ANOVA,

$p < 0.001$), which could be consistent with the finding that the DC had a lower SMI than LC.

Similar to what was shown in the anesthetized animals, the fraction of Noise-sel cells was

higher than that of Tone-sel cells in both $LC_{(micrprism)}$ and DC imaged directly from the top in

awake animals (Figure 10B and D). In contrast to the anesthetized animals, the fraction of Non-

sel cells had different profiles. For instance, while the percentage of Non-sel cells was higher

than Noise-sel and Tone-sel cells in the $LC_{(microprism)}$ (Figure 10B, one way ANOVA, $f_{(2,6)} = 41.6$, p

$= 3.0 \times 10^{-4}$), they were significantly lower than that of Noise-sel cells and similar to that of Tone-

sel cells in the DC (Figure 10D, one way ANOVA, $f_{(2,6)} = 12.8$, $p = 0.006$). Despite that modules

and matrix had similar fractions of Noise-sel, Tone-sel, and Non-sel cells (Figure 10C, Mann-

Whitney Test (Modules vs matrix), z = -0.46, -0.46, and -1.85, p = 0.64, 0.64, and 0.06 for

Noise-sel, Tone-sel, and Non-sel, respectively), matrix had more spectrally (Figure 10E, two-

way ANOVA: $p = 5.3 \times 10^{-10}$, 1.0, and 0.17 for the region, sound level, and interaction,

respectively) and temporally (Figure 10F, two-way ANOVA: $p = 4.6 \times 10^{-7}$, 1.0, and 0.11 for the

region, sound level, and interaction, respectively) modulated cells than modules, which was

similar to anesthetized animals. Additionally, the $LC_{(microprism)}$ and DC showed different profiles

than those found by anesthetized animals. While LC and DC had similar SMI (Figure 10G, two-

way ANOVA: p = 0.22, $2.1x10^{-12}$, and $1.1x10^{-6}$ for the region, sound level, and interaction,

respectively), the $LC_{(microprism)}$ had a lower average of TMIs compared to the DC (Figure 10H,

two-way ANOVA: p = $1.9x10^{-0}$, $6.6x10^{-24}$, and 0.01 for the region, sound level, and interaction,

respectively).

Similar to the anesthetized preparations, the auditory and somatosensory responsive cells were

topographically organized across modules and matrix (Figures 11A, C, and E). However, it was

noticed that the somatosensory and auditory responsive cells were clustered within a pool of

nonresponsive cells (Figure 11E). Some of these nonresponsive cells to unmodulated

broadband noise (Figure 11-supplementary figures 1A) were found to respond to pure tones

(Figure 11-supplementary figures 1B, blue arrows) indicating that these nonresponsive cells

may be sensitive to a specific feature or type of sensory stimulations. Additionally, some of the

cells that did not respond to any of the tested stimulations showed spontaneous activity (Figure

11-supplementary figure 1C) indicating the viability of the nonresponsive cells. Within the pool of

responsive cells, the matrix and modules imaged via microprism had higher fractions of Aud-sel

cells than those of Som-sel and Aud/Som-nonsel cells (Figures 11E and G, Chi-Square test, χ^2

= 42.5, p = $4.4x10^{-8}$), which was similar to the data obtained from the anaethetized animals.

Additionally, matrix had a higher fraction of auditory responsive cells than modules, while

modules had a higher fraction of somatosensory responsive cells than matrix (Figures 11A, D,

and H, Chi-Square test, χ^2 = 30.2, p = $1.2X10^{-6}$). In contrast to the anesthetized animals, the

offset cells were very rare (only 6 cells/3 animals), and were evenly distributed between

modules and matrix resulting in no significance difference between matrix and modules (Figures

11C and I, Chi-Square test, χ^2 = 2.8, p = 0.41). Similar to anesthetized data, we found that

matrix and modules had higher fractions of cells with auditory suppressed responses than those

with auditory enhanced responses induced by bimodal stimulation indicating an overall

suppression in the cellular auditory response. In contrast to the anesthetized animals, modules

had a higher fraction of cells with suppressed response than matrix and a lower fraction of cells

with enhanced responses compared to matrix (Figures 11G and J, χ^2 test = 31.9, p = 5.2x10^{-7})

indicating the possibility that the bimodal interactions in modules could be affected by the

arousal state of the animal.

Discussion:

Summary:

As a new imaging approach, a microprism was used to image the LC surface, to provide the

first-ever sagittal image of the LC in vivo and to examine its functional organization, which

revealed the distinction between strategies used by LC and DC for processing acoustic and

non-acoustic information. Imaging the LC either via microprism or from the dorsal surface

revealed that it contains GABAergic modules that were embedded in a non-GABAergic matrix.

Although modules and matrix were found to be activated by somatosensory and auditory stimuli,

they were functionally distinct. The matrix had more auditory-responsive cells with a lower

threshold for complex sound detection and had more cells responsive to complex spectral and

temporal auditory features than modules. In contrast, modules had more somatosensory

responsive cells, consistent with anatomical differences in input to the modules compared to

matrix (A. M. Lesicko et al., 2016). Somatosensory stimulation suppressed auditory responses

in the matrix, suggesting that multisensory interactions exist between the matrix and modules.

The novelty of the imaging setup and technical considerations

Microprism-based 2P- microscopy has been successfully used to image the cortical layers in the

somatosensory and visual cortices of mice (Andermann et al., 2013) and to image other

inaccessible cortical surfaces like those folded in the brain sulci such as the medial prefrontal

cortex (Low, Gu, & Tank, 2014) or entorhinal cortex (Beckmann et al., 2019; Heys, Rangarajan,

& Dombeck, 2014; Low et al., 2014). In those studies, the glass window glued to the microprism

was used, which requires significant skill and potential removal of brain regions to allow prism

access (Heys et al., 2014). Another approach is to use a micro metal or diamond-made knife to

create a pocket with minimal tissue damage (Andermann et al., 2013; Marks & Goard, 2021).

Accordingly, we combined and modified some of these procedures to improve the microprism

implantation to image the sagittal surface of the LC, which is deeply embedded into the brain

tissue in a highly vascular area, with minimum damage to the IC or the surrounding vasculature.

Previous work using microprisms has been done chronically for up to 60 days (Heys et al.,

2014; Low et al., 2014; Wenzel, Hamm, Peterka, & Yuste, 2017), with minimal tissue

inflammation (Andermann et al., 2013). The current study was initially done on anesthetized

animals to examine and validate the new approach, which later revealed similar organizational

differences between LC and DC in awake animals. Although imaging through a microprism and

1-mm cover glass appears to have minimal impact on image quality compared to direct 2P

imaging (Chia & Levene, 2009), it was important to compare LC imaging via microprism with the

direct 2P imaging of the dorsal surface to validate the new imaging method. We found no

significant differences between LC response properties when imaging via microprism or imaging

directly from the dorsal surface. However, we did observe significant functional differences

between the LC and the DC.

Responsiveness to spectral and temporal features.

While the LC showed inconsistent and fragmented tonotopy, the DC showed a consistent one,

as previously reported (Barnstedt et al., 2015; A. B. Wong & Borst, 2019). These data were

supported by a lower R^2 of the best regression fit between BTFs of the LC cells and their

locations compared to a higher R^2 obtained for the DC. The direction of the best regression fit

shown by the LC was inconsistent across animals, which could be related to the fact that the

microprism did not permit a full view of the LC surface. Given that microprism permits imaging of

only a single area per animal, it is expected that each animal may have a different profile of

cellular organization based on the location of the imaged area on the LC.

The surface of the DC was biased to lower frequencies compared to LC, consistent with

previous imaging and recording studies (Ito et al., 2014; Romand & Ehret, 1990; Shen et al.,

2003; Stiebler & Ehret, 1985). Despite their response to pure tones, the DC and LC, particularly

in the matrix, were more responsive to AM-noise, which fits their function as nonlemniscal

divisions (Chen, Cheng, Ito, & Song, 2018; Faye-Lund & Osen, 1985; Loftus et al., 2008).

The spectral and temporal features of animal vocalization and speech are not independent

(Singh & Theunissen, 2003). We observed that LC neurons, particularly in the matrix, had

greater responsiveness to spectral and temporal modulation than the DC. Given the role of

inhibition in spectral and temporal processing (Cai et al., 2018; Chang, Bao, Imaizumi,

Schreiner, & Merzenich, 2005; Wehr & Zador, 2003), GABAergic cells of the modules could play

an important role in increasing the spectral and temporal modulation indices, especially at lower

sound levels, which could be critical for localizing objects in noisy environments given the

importance of spectral and temporal cues for sound localization (Davis, Ramachandran, & May,

2003; Goodman, Benichoux, & Brette, 2013; Sinex, 2005). Therefore, future work is critical to

image the functional responses of the GABAergic cells of the modules and compare them with

the non-GABAergic cells inside and outside the modules.

Somatosensory vs. auditory responses:

Consistent with the previous anatomical findings (A. M. Lesicko et al., 2016), modules showed

more somatosensory responsive cells than matrix. Since modules receive somatosensory

inputs from different brain regions (A. M. Lesicko et al., 2016; Lohse et al., 2022; Zhou & Shore,

2006), the somatosensory response shown by modules could be modulated by different types of

somatosensory stimulations, which is an open area for future studies. In contrast to modules,

we observed that most auditory-responsive neurons, particularly onset-responsive neurons,

were found in the matrix, which receives mostly auditory inputs (A. M. Lesicko et al., 2016).

However, a subpopulation of acoustically-driven offset-responsive neurons preferably resides in

the modules compared to the matrix. Given that the offset response could be generated as a

rebound to the termination of the ongoing inhibition during the sound presentation (Xu, 2014),

GABAergic cells in the modules could play an important role in the coding of sound termination

as found before in the cortex (Solyga & Barkat, 2021). Because some of the offset acoustically-

driven cells inside the modules were also activated by the somatosensory stimulation, this

fraction of cells could then encode the termination of somatosensory-associated sounds through

the local inhibitory circuits of the modules. Under in-vivo recording of the mouse IC, the offset

responses were previously reported from cells that were non-GABAergic and located in the

caudal and shallower parts of the IC (Kasai, Ono, & Ohmori, 2012). These data suggest that the

module/matrix system permits the preservation of distinct multimodal response properties,

which is consistent with their anatomical didtinction, despite the massive integration of inputs in

the LC.

Consistent with previous reports (Aitkin et al., 1978; Aitkin, Kenyon, & Philpott, 1981; Jain &

Shore, 2006), we observed that combined auditory-somatosensory stimulation generally

suppressed neural responses to auditory stimuli and that this suppression was the most

prominent in the LC matrix of anesthetized animals. Mechanistically, it is not known if such

suppressive responses are created in the LC or are inherited from other brain regions since

auditory-somatosensory interactions occur as early as in the cochlear nucleus, and are also

found in the IC, thalamus, and cortex (Aitkin et al., 1981; Haenggeli, Pongstaporn, Doucet, &

Ryugo, 2005; Kimura, Imbe, Donishi, & Tamai, 2007; Li & Mizuno, 1997; Lohse, Dahmen, Bajo,

& King, 2021; Shore, 2005; Shore, Vass, Wys, & Altschuler, 2000; C. Wong, Chabot, Kok, &

Lomber, 2015), reviewed in (Brunelle & Llano, 2023; Lohse et al., 2022). We note that a

systematic exploration of the stimulus space was not conducted here such that other acoustic or

somatosensory stimuli, different relative amplitudes, or timings could have produced

enhancement rather than suppression. Future work will be needed to further explore this

stimulus space to more fully understand multisensory processing in the LC.

Effect of ketamine anesthesia:

The data obtained from anesthetized animals were similar to those obtained from awake

animals in the findings related to the tonotopy, the preference of the cells to respond to AM-

noise in the LC and DC, the increased sensitivity of matrix to detect the spectral and temporal

features of sound, and the bimodal response properties of the matrix vs modules. However,

anesthetized and awake animals showed some differences that could be related to the effect of

ketamine. For instance, ketamine was found to broaden the RF of the auditory-responsive

neurons (Guo et al., 2012). Consistently, the width of the RF of all DC and LC neurons was

higher under ketamine anesthesia compared to awake preparation. However, the DC neurons

had a wider RF than those in the LC. The relatively narrower RF of the LC could be consistent

with their higher SMI (discussed below) compared to that of the DC. Both findings could be

related to the degree of inhibition that may shape the inhibitory bands of the RF in the LC

neurons given that the LC has prominent GABAergic modules. Also, while the average of the

SMIs was negative with no difference between the $LC_{(microprism)}$ and the DC of awake animals, it

was positive with a higher value in the $LC_{(microprism)}$ compared to the DC of the anesthetized

animals indicating the preference of cells to respond more to pure tones under anesthesia.

Given that ketamine was reported to increase the duration of the tone-evoked response in the

auditory cortex (Guo et al., 2012), the same effect could increase the AUC for the tone-evoked

response that might result in a positive average of SMI. Additionally, while LC had a higher

average of TMIs than that of the DC of anesthetized animals, the LC had a lower mean of TMIs

than that of the DC of awake animals indicating that ketamine may specifically modulate the

response properties of the LC via its effects on GABAergic cells (Behrens et al., 2007; Deane et

al., 2020; Weckmann et al., 2019), given that GABAergic inhibition reportedly contripute to the

processing of the temporally modulated signals (Alluri et al., 2021; Burger & Pollak, 1998; Cai et

al., 2018; Gourévitch, Mahrt, Bakay, Elde, & Portfors, 2020). In both anesthetized and awake

animals, matrix neurons had always a consistent ratio between the cells with suppressed (~

70%) and enhanced (~30%) auditory responses induced by bimodal stimulation. However,

modules had variable ratios across anesthetized and awake animals. While modules had a

lower fraction of cells with enhanced responses than matrix in awake animals, they had a higher

fraction of cells with enhanced responses in anesthetized animals, which could be consistent

with the direct inhibitory effect of ketamine on GABAergic cells (Behrens et al., 2007; Deane et

al., 2020; Weckmann et al., 2019). Given that modules are clusters of GABAerigic cells and

terminals, modules could be specifically modulated by ketamine anesthesia.

Conclusions:

Here, using a novel microprism approach to image the LC in anesthetized and awake mice, we

observed that 1) DC displayed tonotopic organization which was not consistently present in the

LC, 2) DC neurons were generally more sound-responsive than LC neurons but that neurons in

both regions had preferential responses to spectrally- and temporally-complex stimuli, 3) that

matrix regions of the LC were more responsive to spectrally- and temporally-complex stimuli

than modules and that modules were more responsive to somatosensory stimuli and 4) that

bimodal stimulation led to suppressive responses that were more prominent in the matrix. In

general, these data highlight significant organizational differences between the LC and DC and

between modules and matrix in the LC. Future work using microprisms in the LC will help to

reveal the potential differential role of matrix vs. module neurons in multisensory integration

tasks in behaving animals.

Materials and methods:

Animal subjects:

Male mice of 8-12 weeks of age were used. Males were used as our preliminary data suggest

male/female differences in DC organization (Ibrahim, Shinagawa, Xiao, Asilador, & Llano, 2022)

and will be published separately. GAD67-GFP knock-in mice (developed and shared with

permission from Dr. Yuchio Yanagawa at Gunma University and obtained from Dr. Douglas

Oliver at the University of Connecticut), where GFP is exclusively expressed in GABAergic cells

(Tamamaki et al., 2003), were used to visualize the GABAergic cells and LC modules. To

simultaneously monitor calcium signals and visualize the GFP+ cells, Tg(Thy1-

jRGECO1a)GP8.20Dkim/J mice (Jackson Laboratory, Stock# 030525) were crossed with

GAD67-GFP knock-in mice to generate GAD67-GFPxThy1-jRGECO1a hybrid mice. The mice

were housed and bred at the animal research facility of The Beckman Institute for Advanced

Science and Technology. The animal care facilities are approved by the Association for

Assessment and Accreditation of Laboratory Animal Care (AAALAC). To phenotype the GFP-

positive animals, a single-photon fluorescence microscope was used for the transcranial

examination of the pups at postnatal day 4 using excitation: 472/30 nm and emission 520/35 nm

filters along with dichroic 505 nm long pass at low power magnification (2.5x/0.08, Olympus

objective, MPlanFL N, Japan). The positive pups exhibited a green fluorescence in the

cerebellum, the cerebral cortex, and the olfactory bulb and were only kept with their parents until

weaning. The genotyping for jRGECO1a was done using a probe composed of a forward

primer: GCCGCCGAGGTCAAGA & a reverse primer: TCCAACTTGATGTCGACGATGTAG) by

Transnetyx™ (Transnetyx, USA). Samples of the mice's tail snips around one month of age

were used for genotyping. All applicable guidelines for the care and use of animals were

followed. All procedures were approved by the Institutional Animal Care and Use Committee

(IACUC).

Surgery

Surgery

The general procedures of craniotomy were previously described (Goldey et al., 2014), with

modifications to place craniotomy over the IC. Before surgery, mice were anesthetized with a

mixture of ketamine, xylazine, and acepromazine (100 mg/kg, 3 mg/kg, and 3mg/kg,

respectively) delivered intraperitoneally. The anesthesia was maintained during the surgery and

imaging using only ketamine (100 mg/kg). To prevent neural edema during or after the

craniotomy, an intramuscular injection of dexamethasone sodium (4.8 mg/kg) was given just

before the surgery using an insulin syringe. After placing the animal in the stereotaxic apparatus

(David Kopf Instruments, USA), both eyes were protected by applying Optixcare lubricant eye

gel (Aventix Animal Health, Canada). The hair on the scalp was then removed by massaging

the scalp with a depilatory cream (Nair) using a cotton-tipped applicator and leaving the cream

on the scalp for 4-5 minutes. The cream was then removed by a thin plastic sheet (flexible ruler)

to leave a hair-free area on the scalp**.** The remaining tiny hairs were then removed by alcohol

swab and the area was then sterilized by applying 10% Povidone Iodine (Dynarex, USA) using

a sterile cotton-tipped applicator. The medial incision was made with a scalpel blade #10, and

0.2 ml of 0.5% lidocaine was injected intradermally into the scalp. The skin extending from the

medial line to the temporalis muscle was completely removed using a pair of microscissors to

produce a wide skinless area above the skull. Number 5/45 forceps were used to remove any

remaining periosteum. The remaining dried or firmly attached pieces of periosteum were

removed with a scalpel blade #10. The skull was cleaned with sterile saline and dried with

gently pressurized air. Using the stereotaxic apparatus, a wide area of ~3 x 4 mm above the left

IC was made. A micro drill bit (size #80, Grainger, USA) was used to drill through the skull

starting from the rostrolateral (Figure 1A, yellow circle) region to lambda (Figure 1A, black circle)

following the border in a clockwise direction (Figure 1A, dotted black rectangle). To prevent

overheating of the superficial brain tissues and to mitigate the occasional spurts of skull

bleeding during the drilling, ice-cold sterile saline was used to intermittently irrigate the surface.

A stream of pressurized air was also applied during the drilling procedures to prevent

overheating and remove the debris produced by the drilling. Caution was taken not to pierce the

dura when performing the craniotomy while crossing the sagittal or the lambdoid sutures to

avoid damaging the underlying sinuses. After drilling, the skull was irrigated in sterile saline and

the bone flap (the undrilled bone over the craniotomy area) was gently examined for complete

separation from the rest of the skull (Figure 1B). Using a pair of no. 5/45 forceps, the bone flap

was gently removed. To control the bleeding if it occurred, a piece of sterile hemostatic gel

(Adsorbable Gelatin Sponge USP, Haemosponge, GBI, India) which was pre-soaked in ice-cold

saline, was applied to the bleeding spot. Once the bleeding ceased, the brain was kept covered

in sterile saline. In some surgeries, the dura was peeled off from the surface of the IC while

removing the bone flap. In the surgeries where the dura remained intact, a Bonn microprobe

(F.S.T, Germany, Item # 10032-13) was used to pierce the dura in an area that is not above the

IC and devoid of cortical vasculature (e.g., a part of exposed cerebellum). After piercing, the

dura was carefully and gently lifted, and a pair of no. 5/45 forceps were used to grab the dura to

gently tear it to the extent of the transverse sinus to avoid bleeding of the major venous

structures.

Using the stereotaxic apparatus, a 30^{o} diamond knife (F.S.T, Germany, Item # 10100-30) was

placed in its vertical position. From the medial line, the knife was placed at the lateral horizon of

the IC. The knife was then inserted 1.5 mm ventrally. Caution was taken to avoid hitting any

major blood vessels to avoid bleeding. From its position, the knife was retracted 0.3-0.5 mm

medially to create an initial pocket for the insertion of the microprism (Figures 1C-E). The

microprism (1.5 mm-silver coated, Tower Optical Corporation) was then inserted, so one of its

sides was parallel to the lateral surface of the diamond blade, the other side was facing up, and

its hypotenuse was facing ventrally (Figures 1F and G). The wooden piece of a cotton swab was

carefully driven down to push the microprism until its top surface was at the same level as the

IC dorsal surface. Keeping the microprism in its place with the wooden end of a swab, the

diamond knife was carefully retracted upward leaving the microprism behind (Figure 1H). If the

retraction of the diamond knife pulled the prism up, the wooden piece would be used to push it

down simultaneously with the knife movement. To control the bleeding, a piece of sterile

hemostatic gel was used as before. The tip of the diamond knife was used to secure the

microprism in its position for at least 10 minutes (Figure J). During that time, the solution of the

1% agarose was prepared for use. A pair of forceps were then used to gently place a sterile 5

mm cover glass #1 (Thomas Scientific, USA) over the skull covering the opened area above the

IC. The cover glass was secured by a wooden trimmed piece of sterile cotton swab by gently

pressing the cover glass from the top. Since the surface of the IC was located below the

surfaces of both the cerebellum and cerebral cortex, the cover glass usually traps about a 0.7-

1.0 mm thick sheet of sterile saline. The movement of the liquid medium caused by heart

pulsation was found to be a source of motion artifact during imaging, so the gap between the

surface of the IC and the cover glass was filled with 1% agarose gel that was made in saline at

a temperature range of 33-35^{o}C (Figure 1K). The extra agarose gel pieces were cut off and

removed by the scalpel blade and the wet skull was then dried out using pressurized air. A

titanium headpost as described before (Goldey et al., 2014) was glued carefully on the top of

the skull to be at the same level as the cover glass. Following the manufacturer's instructions,

the C&B Metabond (Parkell, Japan) was used to secure the headpost in its place. Excluding the

steps of the microprism insertion, the same procedures were followed for DC imaging from the

surface (Figure 1L). For awake preparations, the same procedures were followed under

isoflurane anesthesia starting with 4% of isoflurane as an induction dose with 1-2% isoflurane

during the surgery. The animal was transferred under the microscope objective and left to

recover from isoflurane anesthesia for 1 hour before imaging. To ensure the animal's recovery,

the animal was visually inspected for its breathing rate (Ewald, Werb, & Egeblad, 2011) and

ongoing limb movements.

Acoustic stimulation:

Using a custom-made MATLAB (The MathWorks, Natick, MA, USA) code, either 500 ms pure

tone or AM-noise was generated. Thirty-five (5x7) pure tones with a series of sound pressure

levels (40, 50, 60, 70, 80 dB SPL) and a series of carrier frequencies (5000-40000 Hz with a

half-octave gap) were presented with a cosine window. Forty-five (5x9) 100% AM white noise

bursts with a series of sound pressure levels (40, 50, 60, 70, 80 dB SPL) and modulation

frequencies (0, 2, 4, 8, 16, 32, 64, 128, 256 Hz) were generated. The stimuli of either pure tone

or AM-noise combinations were played in random sequence to the mice with a 600 ms

interstimulus interval (ISI) by a TDT RP2.1 processor (Tucker-Davis Technologies, US) and

delivered by a TDT ES1 speaker (Tucker-Davis Technologies, US). In other sets of

experiments, 500 ms of unmodulated broadband noise was presented 5 times with ISI of 600

ms at 80 dB SPL to examine the acoustic activity between modules and matrix compared to

somatosensory stimulation.

The output of the TDT ES1 speaker was calibrated using a PCB 377A06 microphone, which

feeds a SigCal tool to generate a calibration file for all tested frequencies (5–40 kHz). To enable

the custom-made MATLAB code to read this calibration file, the values were first processed by

MATLAB signal processing toolbox (sptool) to generate a 256-tap FIR filter to apply the

calibration using the following parameters [arbitrary magnitudes, least square, order: 256,

sampling rate: 97,656.25, frequency vector (5–40 kHz), amplitude vector (40–80 dB SPL), and

weight vector ones (1128)].

Somatosensory stimulation:

The somatosensory stimulation was done by deflecting the right whiskers of the animals at an

average excursion of 1.2 mm with a 50 Hz rate over 500 ms to match the acoustic stimulus. A

total of five stimuli were presented with 600 ms ISI. The whisker deflection was performed using

a Brüel & Kjær vibrator (V203 10/32 UNF- CE, United Kingdom) that was driven by a wave

generator (4063, BK precision, USA) after amplifying the signals with an SLA1 100 W power

amplifier (ART, USA).

Two-Photon imaging:

Immediately after surgery, the anesthetized animal was taken and secured under the

microscope objective by clamping the arms of the head post to two perpendicular metal posts

mounted on the microscope stage. A custom-built 2P microscope was used. The optical and the

controlling components were supplied from Bruker, Olympus, and Thorlabs. The imaging of the

DC was made using a 20x water-immersion objective (LUMPlanFI/IR, 20X, NA: 0.95, WD: 2

mm; Olympus Corporation, Tokyo, Japan), while the lateral surface of the IC was imaged by the

long working distance 16x water-immersion objective (N16XLWD-PF - CFI LWD Plan Fluorite

Objective, 16X, NA: 0.8, WD: 3.0 mm; Nikon, Tokyo, Japan) to be able to reach the focal point

of the microprism at its hypotenuse. For imaging both the GFP or jRGECO1a signals, the

excitation light was generated by InSight X3 laser (Spectra-Physics Lasers, Mountain View, CA,

USA) tuned to a wavelength of 920 or 1040 nm, respectively. A layer of a 1:1 mixture of

wavelength multipurpose ultrasound gel (National Therapy, Canada) with double-deionized

water was used to immerse the objective. This gel was able to trap the water and reduce its

evaporation during imaging. The emitted signals were detected by a photomultiplier tube

(Hamamatsu H7422PA-4, Japan) following a t565lp dichroic and a Chroma barrier et525/70m

filter for GFP and et595/50m filter for jRGECO1a signals. Images (512x512 pixels) were

collected at a frame rate of 29.9 Hz in the resonant galvo mode. Data were collected from the

dorsal surface of the IC by scanning the surface of the IC based on the GFP and jRGECO1a

signals through the medial and lateral horizons of the IC. Generally, the scanning was started by

moving the 20x objective to the most ventromedial position, where there is no ability to see any

cells (Figure 2B, medial horizon). Then, the objective was driven to slowly move dorsolaterally

to get cells in focus (Figure 1M), where the average depth of the first field of view was 169 μm.

The lateral movement of the objective was intended to be aligned and parallel to the curvature

of the DC surface indicated by getting the superficial cells in focus to avoid any possibility of

imaging the upper layers of the central nucleus of the IC, so the average depth of the most

superficial point of the imaging was 92 μm. The scanning was ended by moving the objective to

the most ventrolateral position of the IC, where there is no ability to see GFP laterally (Figure

2B, lateral horizon), where the last field of view was collected at an average depth of 225 μm.

Generally, each field of view was selected based on the expression of GFP signals as an

indicator for the cells in focus and being acoustically active by using a search stimulus that was

500 ms broadband noise with zero modulation at 80 dB SPL. The z-axis of the objective was

determined relative to the pia surface. The relative position of each region was tracked through

the micromanipulator (MP-285, Sutter Instruments) that controlled the microscope objective.

The positions of imaged areas were further aligned across animals to a common coordinate

using the midline, lateral extremes, and the vasculature landmarks taken via imaging of the GFP

signals of the cranial window with low magnification (4x Olympus objective, 4x4 binning, 50 ms

frame time) using a Lumen 200 bulb (Prior Scientific Inc, USA) and CoolSNAP MYO camera

(Photometrics, USA, excitation: 488 nm and emission: 515–550 nm). Imaging of the lateral

surface of the IC via microprism was done similarly. In both cases, the frame timing of the

scanner and the sound stimuli were both digitized and time-locked using a Digidata 1440A

(Molecular Devices, Sunnyvale, CA, USA) with Clampex v. 10.3 (Molecular Devices, Sunnyvale,

CA, USA).

Laser lesion:

The laser lesion was produced using a 720 or 820 nm laser beam at 74-80 mW power at the

level of the objective for 100 to 250 seconds depending on the size of the lesioned area, the

speed of laser scanning (frame rate), and the depth of the lesion (z-axis length). In brief, the

area of interest and the dimension of the laser lesion were initially determined using 920 nm as

indicated by the GFP signals through scanning one of the GABAergic modules. The depth of the

GABAergic modules was then determined by counting the optical slices starting from their

superficial surface. The tunable laser was then switched from 920 nm to the shorter wavelength

(720 or 820 nm) to scan the same area and depth at a rate (1-2 frames/ second).

Data processing:

Data collection: The data were collected as separate movies (512x512 pixels) for each pure

tone or AM-noise runs in a resonant galvo mode. Depending on the amplitude and frequency

combinations for each type of acoustic stimulus, 40- or 52-second periods were assigned as a

movie's length for pure tone (35 stimulus combinations), or AM-noise (45 stimulus

combinations), respectively. Using ImageJ software, the z-projection was used to compute one

single image representing either the sum, the standard deviation, or the median of all the image

sequences in the movie. Based on these single images, the region of interest (ROI) was

manually drawn around each detectable cell body. Figures 3A and B represent the sum of the

image sequences of the movie collected from the dorsal surface, or the lateral surface of the IC

via microprism, respectively. For the subsequent processing steps, Python open-source libraries

were used (listed below).

Motion Correction and Filtering: The imread function from the OpenCV library was used in

grayscale mode to import images into a numpy array from a single folder containing TIFF

images. The array was then exported as a single TIFF stack into a temporary folder using the

mimwrite function from the imageio library, and the process was repeated for each folder. The

NoRMCorre algorithm (Pnevmatikakis & Giovannucci, 2017) embedded in the CaImAn library

(Giovannucci et al., 2019) was used to apply motion correction to each of the TIFF files. The

data were then imported into a numpy array, rounded, and converted to 16-bit integers. The

images were filtered using a 2D Gaussian filter with a sigma value of 1 (surface View) / 2 (prism

View) in each direction, then a 1D Gaussian temporal filter with a sigma value of 2 was applied

using the ndimage.gaussian_filter and ndimage.gaussian_filter1d function from the scipy library,

respectively.

Data Extraction: The ROI sets, which were manually created using ImageJ, were imported using

the read_roi_zip function from the read_roi library. The sets were then used to create two

masks; one mask was used as a replica of the ROIs and the second mask was made around

the original ROI (roughly four times larger in area). The smaller mask was applied to find the

average pixel value within each ROI, while the larger mask was applied to find the average pixel

value of the neuropil. The neuropil correction was applied using the following equation

(Akerboom et al., 2012; Kerlin, Andermann, Berezovskii, & Reid, 2010);

$$\text{Corrected value} = \text{Date value} - (0.4 \text{ X Neuropil value})$$

777   To identify the calcium signals, Δf/f was calculated by using the following equation.

$$\frac{\Delta f}{f} = \frac{(\text{Date value} - \text{Background value})}{\text{Background value}}$$

where the background value is the slope estimating the background levels with fluorescence

bleaching factored in. The data were then reorganized so that all segments with the same

stimulus frequency and stimulus amplitude were grouped. The area under the curve (AUC) of

the calcium signals of the excitatory responses only was then used as a metric to determine the

magnitude of the response and for subsequent analysis (Berens et al., 2018; Overk,

Rockenstein, Florio, Cheng, & Masliah, 2015).

Cell Flagging: The correlation coefficient between each of the trials was calculated using the

stats. Pearson function from the scipy library. The average correlation coefficient was calculated

for each stimulus frequency and stimulus amplitude. Similar to previous work (A. B. Wong &

Borst, 2019), if the average correlation coefficient was above the threshold of 0.6, the cell was

flagged as being responsive to that stimulus combination of frequency and amplitude, and the

best tone and modulation frequencies were calculated for every cell (Barnstedt et al., 2015).

Knowing BTF and the best modulation frequency (BMF) for each neuron enabled us to calculate

the SMI and TMI at each sound level following these equations;

792   SMI

$$= \frac{\text{Response to BTF} - \text{Response to unmodulated noise}}{\text{Response to BTF} + \text{Response to unmodulated noise}}$$

793   The SMI has values ranging from (-1 to 1). While neurons with values closer to 1 are more

794   responsive to pure tone, neurons with values closer to -1 are more responsive to noise with no

795   modulation.

796   TMI

$$= \frac{\text{Response to BMF} - \text{Response to unmodulated noise}}{\text{Response of BMF} + \text{Response to unmodulated noise}}$$

Similarly, the TMI has values ranging from (-1 to 1). While neurons with values closer to 1 are

more responsive to AM-noise, neurons with values closer to -1 are more responsive to noise

with no modulation.

For the comparison between somatosensory and auditory responses, the threshold of 0.4 was

used as a cutoff threshold to determine the responsive cells to each stimulation. The average

response of each somatosensory, acoustic (500 ms of unmodulated noise), or simultaneous

somatosensory and acoustic (bimodal) stimulation was then calculated for each responsive cell.

The correlation method was validated as a tool to determine the responsive cells to

somatosensory stimulation by generating an ROC curve comparing the automated method to a

blinded human interpretation. The AUC of the ROC curve was 0.88. This high AUC value

indicates that the correlation method can rank the random responsive cells than the random

nonresponsive cells. At the correlation coefficient (0.4), which was the cutoff value to determine

the responsive cells for somatosensory stimulation, the specificity was 87% and the sensitivity

72%, the positive predictive value was 73%, and the negative predictive value was 86%.

Additionally, all the false-responsive cells were excluded from the analysis. To examine the

effect of somatosensory stimulation on auditory responses, the response index (RI) was

calculated for the auditory-responsive cells using the following equation;

814   RI

$$= \frac{\text{Auditory response to bimodal stimulation}}{\text{Auditory response to auditory stimulation only}}$$

Reurons with RI values > 1 have enhanced or higher responses induced by simultaneous

auditory and somatosensory stimulation, while neurons with RI values < 1 have downregulated

or suppressed auditory responses induced by auditory and simultaneous somatosensory

stimulation.

Map Generation: The average radius for all ROIs was calculated to ensure that all cells on the

tonotopic map had uniform radii. A color key was also generated for every corresponding map.

A blank canvas was generated using the Image module from the pillow library, and a circle was

created for each cell and filled with a corresponding color to its value.

Receptive field sum (RFS) analysis: The binary RFS was calculated for every cell by

determining whether the cell was responsive to a particular combination of frequency and

amplitude (Bowen et al., 2020). A value of 1 was given to the responsive cell, if it responds to a

single frequency/amplitude combination. In contrast, a value of 1 was added to the tally for each

responsive combination if the cell was broadly responsive. Accordingly, the higher and lower

values of RFS indicate a broad and narrow RF, respectively. To examine the differences

between the RFS of the cells at different areas under different preparations, the cumulative

distribution function was used.

Assessment of tonotopic organization.

The tonotopy was assessed by examining the best fit of linear or nonlinear quadratic polynomial

regressions between the BTFs of cells and their locations along different anatomical axes. For

the $LC_{(microprism)}$, the regression fit was examined along the main four axes such as dorsal to

ventral, rostral to caudal, dorsocaudal to ventrorostral, and dorsorostral to ventrocoudal. Based

on the previous reports showing the tonotopic organization over the DC (Barnstedt et al., 2015;

A. B. Wong & Borst, 2019), the tonotopy was examined along the rostromedial to caudolateral

axis. The x_2 and y_2 coordinates of each cell were determined. At the starting point of each axis,

the x_1 and y_1 coordinates were assigned values as (0, 0). Then the distance (d) between each

cell and the starting point at each axis was calculated based on the following equation;

$$d = \sqrt{(x2 - x1)^2 + (y2 - y1)^2}$$

For each 50 um (bin size), the geometric average of the BTFs of all cells as well as their

distances from the starting point at each axis were computed. The linear or quadratic polynomial

regressions were then examined between the geometric averages of BTFs and the geometric

averages of distances of cells for each animal.

Detecting the spontaneous activity from the nonresponsive cells

Movies have been recorded from the nonresponsive cells of the $LC_{(microprism)}$ when no stimulation

was presented. The time trace of jRGECO1a signals was extracted from each cell. A polynomial

fitting of the time trace was used to correct for photobleaching and to adjust for the baseline for

the whole trace. The sliding average of the baseline was then calculated for 1 second (~29

frames), and the z score was calculated for the subsequent frames. The spontaneous activity

was detected when the z-score was > 3.0.

Statistics:

All statistical analyses and graphs were made using Origin Pro software. The normality of the

distributions of the data was examined using a Kolmogorov-Smirnov test. The Mann-Whitney

test as well as two-sample or paired t-test were used to examine the difference in data

distribution or the difference between samples of non-normal and normal distribution,

respectively. Either one-way or two-way ANOVA as well as Fisher as a post hoc test were used

for the mean comparison between different groups of normal distribution. The Chi-square (χ^2)

test or Kruskal-Wallis ANOVA tests were used to compare the samples based on cell numbers

of their cellular population. The cumulative frequency function was computed to calculate the

fraction of cells tuned at each frequency based on their best frequency. Differences were

deemed significant when p-value <0.05. The upper and lower borders of each box plot are at

the 25th and 75th percentile, with whiskers at 95% spread.

Histology:

After the 2P imaging, the mouse was transcardially perfused with 4% paraformaldehyde (PFA)

solution in PBS solution. The isolated brain was then placed overnight in PFA for a complete

fixation. The brain was then moved from the PFA to a graded sucrose solution (10, 20, and

30%) until it sunk. 50 µm coronal sections were taken at the level of the IC using a Leica

cryostat. The sections were then mounted on a slide, coverslipped, and imaged using a

confocal microscope for imaging the GFP signals as an indicator for the GABAergic cells in the

GAD67-GFP knock-in mouse (excitation: 488 nm and emissions: 515–550 nm).

For the examination of jrgeco1a expression in the IC, the mouse was transcardially perfused

with saline with a high concentration of potassium and a high ratio of calcium to magnesium [(in

mM) 3.5 KCl, 25 glucose, 125 NaCl, 1.2 KH2PO4, 2 CaCl2, 0.5 MgSO4] under ketamine

anesthesia. The brain was then fixed by a subsequent perfusion of 4% PFA. Selected 40 μm

coronal sections at the level of the IC were immunostained for NeuN using (NeuN (D4G40) XP®

Rabbit mAB, Cell Signaling, Cat# 62994, 1:100) primary antibody and (goat anti-rabbit IgG

conjugated to Alexa 405, 1:200) secondary antibody following the immunostaining procedures

that were previously described (Issa, Sekaran, & Llano, 2023; Vaithiyalingam Chandra Sekaran

et al., 2021). Confocal images were obtained using a Confocal Zeiss LSM 710 Microscope

using 10 and 40x objectives and tile scan functions. For NeuN, Alexa 405 signals were obtained

by a 405 nm laser and emission at 415-466 nm. The GFP signals were obtained by a 488 nm

laser and emission at 501-551 nm. The jRGECO1a signals were obtained by 561 nm laser and

emission at 600-700 nm. All cells with somas expressing a red fluorescence of jrgeco1a were

quantified in 2P-imaged areas of the DC and LC. The density of the cells expressing jrgeco1a

was then calculated based on the cells expressing NeuN.

Artwork:

The cartoon of the mouse brain in Figures 1 C, F, and I was obtained from Brain Explorer 3D

viewer Allen Mouse Brain (https://connectivity.brain-map.org/3d-viewer?v=1) and modified to be

suitable for illustration. All figures were designed and made using Adobe Illustrator (Adobe, San

Jose, CA). To keep working within the Adobe environment to avoid losing the resolution of the

figures, Adobe Photoshop (Adobe, San Jose, CA) was used to crop the borders of some images

to save space. The color pallets were chosen to avoid the combination of the green and red

colors for all maps, so they will be suitable for color-blind readers. To induce the same effect in

the microscope images combining the red (jRGECO1a) and green (GFP) fluorescence, we kept

the green signals of the GFP unchanged, but we substituted the red fluorescence of jRGECO1a

with magenta as a pseudocolor.

Acknowledgment

We thank Dr. Kush Paul (Bruker) for initial training on the craniotomy surgery, Dr. Jason

Maclean (The Department of Neurobiology, University of Chicago) for assistance with 2P

imaging, Dr. Michael Jacobs Goard (Neuroscience Research Institute, The University of

California- Santa Barbara) for his guidance in the use of microprisms, Dr. Taher Saif

(Department of Mechanical Science and Engineering, The University of Illinois Urbana-

Champaign) for measuring the deflecting distance of the somatosensory stimulating probe using

a high-speed camera, Dr. Pengfei Song (Department of Electric and Computer Engineering,

The University of Illinois Urbana-Champaign) for his assistance with the somatosensory

stimulating probe, and Dr. Nathiya Vaithiyalingam Chandra Sekaran (Department of Molecular

and Intergartive physiology, The University of Illinois Urbana-Champaign) for her assistance in

immunohistochemistry and confocal imaging. We also thank the NIDCD for funding this work

under grants # R01DC013073 and R01DC016599.

Competing interests

On behalf of all authors, there are no financial or non-financial competing interests to be

declared.

References

Aitkin, L. M., Dickhaus, H., Schult, W., & Zimmermann, M. (1978). External nucleus of inferior colliculus:
auditory and spinal somatosensory afferents and their interactions. *J Neurophysiol, 41*(4), 837-
847. doi:10.1152/jn.1978.41.4.837
Aitkin, L. M., Kenyon, C. E., & Philpott, P. (1981). The representation of the auditory and somatosensory
systems in the external nucleus of the cat inferior colliculus. *J Comp Neurol, 196*(1), 25-40.
doi:10.1002/cne.901960104
Akerboom, J., Chen, T. W., Wardill, T. J., Tian, L., Marvin, J. S., Mutlu, S., . . . Looger, L. L. (2012).
Optimization of a GCaMP calcium indicator for neural activity imaging. *J Neurosci, 32*(40), 13819-
13840. doi:10.1523/JNEUROSCI.2601-12.2012
Alexander, G. E., Crutcher, M. D., & DeLong, M. R. (1990). Basal ganglia-thalamocortical circuits: parallel
substrates for motor, oculomotor, "prefrontal" and "limbic" functions. *Prog Brain Res, 85*, 119-
146.
Alluri, R. K., Rose, G. J., Leary, C. J., Palaparthi, A., Hanson, J. L., Vasquez-Opazo, G. A., . . . Luong, K.
(2021). How auditory selectivity for sound timing arises: The diverse roles of GABAergic
inhibition in shaping the excitation to interval-selective midbrain neurons. *Prog Neurobiol, 199*,
101962. doi:10.1016/j.pneurobio.2020.101962

Andermann, M. L., Gilfoy, N. B., Goldey, G. J., Sachdev, R. N., Wölfel, M., McCormick, D. A., . . . Levene,
M. J. (2013). Chronic cellular imaging of entire cortical columns in awake mice using
microprisms. *Neuron, 80*(4), 900-913. doi:10.1016/j.neuron.2013.07.052
Bajo, V. M., & Moore, D. R. (2005). Descending projections from the auditory cortex to the inferior
colliculus in the gerbil, Meriones unguiculatus. *J Comp Neurol, 486*(2), 101-116.
doi:10.1002/cne.20542
Bandyopadhyay, S., Shamma, S. A., & Kanold, P. O. (2010). Dichotomy of functional organization in the
mouse auditory cortex. *Nat Neurosci, 13*(3), 361-368. doi:10.1038/nn.2490
Barnstedt, O., Keating, P., Weissenberger, Y., King, A. J., & Dahmen, J. C. (2015). Functional
Microarchitecture of the Mouse Dorsal Inferior Colliculus Revealed through In Vivo Two-Photon
Calcium Imaging. *J Neurosci, 35*(31), 10927-10939. doi:10.1523/JNEUROSCI.0103-15.2015
Beckmann, L., Zhang, X., Nadkarni, N. A., Cai, Z., Batra, A., Sullivan, D. P., . . . Zhang, H. F. (2019).
Longitudinal deep-brain imaging in mouse using visible-light optical coherence tomography
through chronic microprism cranial window. *Biomed Opt Express, 10*(10), 5235-5250.
doi:10.1364/BOE.10.005235
Behrens, M. M., Ali, S. S., Dao, D. N., Lucero, J., Shekhtman, G., Quick, K. L., & Dugan, L. L. (2007).
Ketamine-induced loss of phenotype of fast-spiking interneurons is mediated by NADPH-
oxidase. *Science, 318*(5856), 1645-1647. doi:10.1126/science.1148045
Berens, P., Freeman, J., Deneux, T., Chenkov, N., McColgan, T., Speiser, A., . . . Bethge, M. (2018).
Community-based benchmarking improves spike rate inference from two-photon calcium
imaging data. *PLoS Comput Biol, 14*(5), e1006157. doi:10.1371/journal.pcbi.1006157
Bowen, Z., Winkowski, D. E., & Kanold, P. O. (2020). Functional organization of mouse primary auditory
cortex in adult C57BL/6 and F1 (CBAxC57) mice. *Sci Rep, 10*(1), 10905. doi:10.1038/s41598-020-
67819-4
Brimblecombe, K. R., & Cragg, S. J. (2017). The Striosome and Matrix Compartments of the Striatum: A
Path through the Labyrinth from Neurochemistry toward Function. *ACS Chem Neurosci, 8*(2),
235-242. doi:10.1021/acschemneuro.6b00333
Brunelle, D. L., & Llano, D. A. (2023). Role of auditory-somatosensory corticothalamic circuit integration
in analgesia. *Cell Calcium, 111*, 102717. doi:10.1016/j.ceca.2023.102717
Burger, R. M., & Pollak, G. D. (1998). Analysis of the role of inhibition in shaping responses to
sinusoidally amplitude-modulated signals in the inferior colliculus. *J Neurophysiol, 80*(4), 1686-
1701. doi:10.1152/jn.1998.80.4.1686
Cai, D., Han, R., Liu, M., Xie, F., You, L., Zheng, Y., . . . Yuan, K. (2018). A Critical Role of Inhibition in
Temporal Processing Maturation in the Primary Auditory Cortex. *Cereb Cortex, 28*(5), 1610-
1624. doi:10.1093/cercor/bhx057
Chang, E. F., Bao, S., Imaizumi, K., Schreiner, C. E., & Merzenich, M. M. (2005). Development of spectral
and temporal response selectivity in the auditory cortex. *Proc Natl Acad Sci U S A, 102*(45),
16460-16465. doi:10.1073/pnas.0508239102
Chen, C., Cheng, M., Ito, T., & Song, S. (2018). Neuronal Organization in the Inferior Colliculus Revisited
with Cell-Type-Dependent Monosynaptic Tracing. *J Neurosci, 38*(13), 3318-3332.
doi:10.1523/JNEUROSCI.2173-17.2018
Chernock, M. L., Larue, D. T., & Winer, J. A. (2004). A periodic network of neurochemical modules in the
inferior colliculus. *Hear Res, 188*(1-2), 12-20. doi:10.1016/S0378-5955(03)00340-X
Chia, T. H., & Levene, M. J. (2009). Microprisms for in vivo multilayer cortical imaging. *J Neurophysiol,
102*(2), 1310-1314. doi:10.1152/jn.91208.2008
Coleman, J. R., & Clerici, W. J. (1987). Sources of projections to subdivisions of the inferior colliculus in
the rat. *J Comp Neurol, 262*(2), 215-226. doi:10.1002/cne.902620204

Dana, H., Novak, O., Guardado-Montesino, M., Fransen, J. W., Hu, A., Borghuis, B. G., . . . Svoboda, K.
(2018). Thy1 transgenic mice expressing the red fluorescent calcium indicator jRGECO1a for
neuronal population imaging in vivo. *PLoS One, 13*(10), e0205444.
doi:10.1371/journal.pone.0205444
Davis, K. A., Ramachandran, R., & May, B. J. (2003). Auditory processing of spectral cues for sound
localization in the inferior colliculus. *J Assoc Res Otolaryngol, 4*(2), 148-163. doi:10.1007/s10162-
002-2002-5
Deane, K. E., Brunk, M. G. K., Curran, A. W., Zempeltzi, M. M., Ma, J., Lin, X., . . . Happel, M. F. K. (2020).
Ketamine anaesthesia induces gain enhancement via recurrent excitation in granular input
layers of the auditory cortex. *J Physiol, 598*(13), 2741-2755. doi:10.1113/JP279705
Dillingham, C. H., Gay, S. M., Behrooz, R., & Gabriele, M. L. (2017). Modular-extramodular organization
in developing multisensory shell regions of the mouse inferior colliculus. *J Comp Neurol, 525*(17),
3742-3756. doi:10.1002/cne.24300
Draganski, B., Kherif, F., Klöppel, S., Cook, P. A., Alexander, D. C., Parker, G. J., . . . Frackowiak, R. S.
(2008). Evidence for segregated and integrative connectivity patterns in the human Basal
Ganglia. *J Neurosci, 28*(28), 7143-7152. doi:10.1523/JNEUROSCI.1486-08.2008
Ewald, A. J., Werb, Z., & Egeblad, M. (2011). Monitoring of vital signs for long-term survival of mice
under anesthesia. *Cold Spring Harb Protoc, 2011*(2), pdb.prot5563. doi:10.1101/pdb.prot5563
Fair, D. A., Dosenbach, N. U., Church, J. A., Cohen, A. L., Brahmbhatt, S., Miezin, F. M., . . . Schlaggar, B. L.
(2007). Development of distinct control networks through segregation and integration. *Proc Natl
Acad Sci U S A, 104*(33), 13507-13512. doi:10.1073/pnas.0705843104
Faye-Lund, H., & Osen, K. K. (1985). Anatomy of the inferior colliculus in rat. *Anat Embryol (Berl), 171*(1),
1-20. doi:10.1007/BF00319050
Frisina, R. D. (2001). Subcortical neural coding mechanisms for auditory temporal processing. *Hear Res,
158*(1-2), 1-27. doi:10.1016/s0378-5955(01)00296-9
Gerfen, C. R. (1985). The neostriatal mosaic. I. Compartmental organization of projections from the
striatum to the substantia nigra in the rat. *J Comp Neurol, 236*(4), 454-476.
doi:10.1002/cne.902360404
Giovannucci, A., Friedrich, J., Gunn, P., Kalfon, J., Brown, B. L., Koay, S. A., . . . Pnevmatikakis, E. A.
(2019). CaImAn an open source tool for scalable calcium imaging data analysis. *Elife, 8*.
doi:10.7554/eLife.38173
Goldey, G. J., Roumis, D. K., Glickfeld, L. L., Kerlin, A. M., Reid, R. C., Bonin, V., . . . Andermann, M. L.
(2014). Removable cranial windows for long-term imaging in awake mice. *Nat Protoc, 9*(11),
2515-2538. doi:10.1038/nprot.2014.165
Goodman, D. F., Benichoux, V., & Brette, R. (2013). Decoding neural responses to temporal cues for
sound localization. *Elife, 2*, e01312. doi:10.7554/eLife.01312
Gourévitch, B., Mahrt, E. J., Bakay, W., Elde, C., & Portfors, C. V. (2020). GABA. *J Neurophysiol, 123*(1),
134-148. doi:10.1152/jn.00377.2019
Guo, W., Chambers, A. R., Darrow, K. N., Hancock, K. E., Shinn-Cunningham, B. G., & Polley, D. B. (2012).
Robustness of cortical topography across fields, laminae, anesthetic states, and
neurophysiological signal types. *J Neurosci, 32*(27), 9159-9172. doi:10.1523/JNEUROSCI.0065-
12.2012
Haenggeli, C. A., Pongstaporn, T., Doucet, J. R., & Ryugo, D. K. (2005). Projections from the spinal
trigeminal nucleus to the cochlear nucleus in the rat. *J Comp Neurol, 484*(2), 191-205.
doi:10.1002/cne.20466
Heys, J. G., Rangarajan, K. V., & Dombeck, D. A. (2014). The functional micro-organization of grid cells
revealed by cellular-resolution imaging. *Neuron, 84*(5), 1079-1090.
doi:10.1016/j.neuron.2014.10.048

Ibrahim, Baher, Shinagawa, Yoshitaka, Xiao, Gang, Asilador, Alexander, & Llano, Daniel. (2022). *Using*
Microprisms, the 2P Calcium Imaging Revealed a Functional Distinction between the Dorsal &
Lateral Cortices of the Mouse Inferior Colliculus. Paper presented at the Association for Research
in Otolaryngology, Virtual.
Issa, L. K., Sekaran, N. V. C., & Llano, D. A. (2023). Highly branched and complementary distributions of
layer 5 and layer 6 auditory corticofugal axons in mouse. *Cereb Cortex, 33*(16), 9566-9582.
doi:10.1093/cercor/bhad227
Ito, T., Hirose, J., Murase, K., & Ikeda, H. (2014). Determining auditory-evoked activities from multiple
cells in layer 1 of the dorsal cortex of the inferior colliculus of mice by in vivo calcium imaging.
Brain Res, 1590, 45-55. doi:10.1016/j.brainres.2014.09.049
Jain, R., & Shore, S. (2006). External inferior colliculus integrates trigeminal and acoustic information:
unit responses to trigeminal nucleus and acoustic stimulation in the guinea pig. *Neurosci Lett,*
395(1), 71-75. doi:10.1016/j.neulet.2005.10.077
Joris, P. X., Schreiner, C. E., & Rees, A. (2004). Neural processing of amplitude-modulated sounds. *Physiol*
Rev, 84(2), 541-577. doi:10.1152/physrev.00029.2003
Kasai, M., Ono, M., & Ohmori, H. (2012). Distinct neural firing mechanisms to tonal stimuli offset in the
inferior colliculus of mice in vivo. *Neurosci Res, 73*(3), 224-237.
doi:10.1016/j.neures.2012.04.009
Kerlin, A. M., Andermann, M. L., Berezovskii, V. K., & Reid, R. C. (2010). Broadly tuned response
properties of diverse inhibitory neuron subtypes in mouse visual cortex. *Neuron, 67*(5), 858-871.
doi:10.1016/j.neuron.2010.08.002
Kimura, A., Imbe, H., Donishi, T., & Tamai, Y. (2007). Axonal projections of single auditory neurons in the
thalamic reticular nucleus: implications for tonotopy-related gating function and cross-modal
modulation. *Eur J Neurosci, 26*(12), 3524-3535. doi:10.1111/j.1460-9568.2007.05925.x
Lamb-Echegaray, I. D., Noftz, W. A., Stinson, J. P. C., & Gabriele, M. L. (2019). Shaping of discrete
auditory inputs to extramodular zones of the lateral cortex of the inferior colliculus. *Brain Struct*
Funct, 224(9), 3353-3371. doi:10.1007/s00429-019-01979-6
Langner, G. (1992). Periodicity coding in the auditory system. *Hear Res, 60*(2), 115-142.
doi:10.1016/0378-5955(92)90015-f
Lesicko, A. M. H., & Geffen, M. N. (2022). Diverse functions of the auditory cortico-collicular pathway.
Hear Res, 425, 108488. doi:10.1016/j.heares.2022.108488
Lesicko, A. M. H., Sons, S. K., & Llano, D. A. (2020). Circuit Mechanisms Underlying the Segregation and
Integration of Parallel Processing Streams in the Inferior Colliculus. *J Neurosci, 40*(33), 6328-
6344. doi:10.1523/JNEUROSCI.0646-20.2020
Lesicko, A. M., Hristova, T. S., Maigler, K. C., & Llano, D. A. (2016). Connectional Modularity of Top-Down
and Bottom-Up Multimodal Inputs to the Lateral Cortex of the Mouse Inferior Colliculus. *J*
Neurosci, 36(43), 11037-11050. doi:10.1523/JNEUROSCI.4134-15.2016
Li, H., & Mizuno, N. (1997). Single neurons in the spinal trigeminal and dorsal column nuclei project to
both the cochlear nucleus and the inferior colliculus by way of axon collaterals: a fluorescent
retrograde double-labeling study in the rat. *Neurosci Res, 29*(2), 135-142. doi:10.1016/s0168-
0102(97)00082-5
Liu, M., Xie, F., Dai, J., Zhang, J., Yuan, K., & Wang, N. (2023). Brain-wide inputs to the non-lemniscal
inferior colliculus in mice. *Neurosci Lett, 793*, 136976. doi:10.1016/j.neulet.2022.136976
Loftus, W. C., Malmierca, M. S., Bishop, D. C., & Oliver, D. L. (2008). The cytoarchitecture of the inferior
colliculus revisited: a common organization of the lateral cortex in rat and cat. *Neuroscience,*
154(1), 196-205. doi:10.1016/j.neuroscience.2008.01.019

Lohse, M., Dahmen, J. C., Bajo, V. M., & King, A. J. (2021). Subcortical circuits mediate communication between primary sensory cortical areas in mice. *Nat Commun, 12*(1), 3916. doi:10.1038/s41467-021-24200-x

Lohse, M., Zimmer-Harwood, P., Dahmen, J. C., & King, A. J. (2022). Integration of somatosensory and motor-related information in the auditory system. *Front Neurosci, 16*, 1010211. doi:10.3389/fnins.2022.1010211

Low, R. J., Gu, Y., & Tank, D. W. (2014). Cellular resolution optical access to brain regions in fissures: imaging medial prefrontal cortex and grid cells in entorhinal cortex. *Proc Natl Acad Sci U S A, 111*(52), 18739-18744. doi:10.1073/pnas.1421753111

Marks, T. D., & Goard, M. J. (2021). Stimulus-dependent representational drift in primary visual cortex. *Nat Commun, 12*(1), 5169. doi:10.1038/s41467-021-25436-3

Nelson, P. C., & Carney, L. H. (2007). Neural rate and timing cues for detection and discrimination of amplitude-modulated tones in the awake rabbit inferior colliculus. *J Neurophysiol, 97*(1), 522-539. doi:10.1152/jn.00776.2006

Oliver, Douglas L. (2005). Neuronal Organization in the Inferior Colliculus. In J.A. Winer & C.E. Schreiner (Eds.), *The Inferior Colliculus.* New York, NY.: Springer.

Overk, C. R., Rockenstein, E., Florio, J., Cheng, Q., & Masliah, E. (2015). Differential calcium alterations in animal models of neurodegenerative disease: Reversal by FK506. *Neuroscience, 310*, 549-560. doi:10.1016/j.neuroscience.2015.08.068

Pnevmatikakis, E. A., & Giovannucci, A. (2017). NoRMCorre: An online algorithm for piecewise rigid motion correction of calcium imaging data. *J Neurosci Methods, 291*, 83-94. doi:10.1016/j.jneumeth.2017.07.031

Rabang, C. F., Parthasarathy, A., Venkataraman, Y., Fisher, Z. L., Gardner, S. M., & Bartlett, E. L. (2012). A computational model of inferior colliculus responses to amplitude modulated sounds in young and aged rats. *Front Neural Circuits, 6*, 77. doi:10.3389/fncir.2012.00077

Rees, A., & Langner, G. (2005). Temporal Coding in the Auditory Midbrain. In J.A. Winer & C.E. Schreiner (Eds.), *The Inferior Colliculus.* . New York, NY.: Springer.

Romand, R., & Ehret, G. (1990). Development of tonotopy in the inferior colliculus. I. Electrophysiological mapping in house mice. *Brain Res Dev Brain Res, 54*(2), 221-234. doi:10.1016/0165-3806(90)90145-o

Romero, S., Hight, A. E., Clayton, K. K., Resnik, J., Williamson, R. S., Hancock, K. E., & Polley, D. B. (2020). Cellular and Widefield Imaging of Sound Frequency Organization in Primary and Higher Order Fields of the Mouse Auditory Cortex. *Cereb Cortex, 30*(3), 1603-1622. doi:10.1093/cercor/bhz190

Rothschild, G., Nelken, I., & Mizrahi, A. (2010). Functional organization and population dynamics in the mouse primary auditory cortex. *Nat Neurosci, 13*(3), 353-360. doi:10.1038/nn.2484

Saldaña, E., Feliciano, M., & Mugnaini, E. (1996). Distribution of descending projections from primary auditory neocortex to inferior colliculus mimics the topography of intracollicular projections. *J Comp Neurol, 371*(1), 15-40. doi:10.1002/(SICI)1096-9861(19960715)371:1<15::AID-CNE2>3.0.CO;2-O

Schreiner, Christoph E, & Winer, Jeffery A. (2005). *The inferior colliculus*: Springer Science+ Business Media, Incorporated.

Shen, Junxian, Xu, Zhimin, & Shen, Lijian. (2003). Topography of acoustic response characteristics in the midbrain inferior colliculus of Kunming mouse. *<i data-test="journal-title" style="margin: 0px; box-sizing: inherit;">Chinese Science Bulletin   volume, 48*, 161–165. doi:https://doi.org/10.1360/03tb9033

Shore, S. E. (2005). Multisensory integration in the dorsal cochlear nucleus: unit responses to acoustic
and trigeminal ganglion stimulation. *Eur J Neurosci, 21*(12), 3334-3348. doi:10.1111/j.1460-
9568.2005.04142.x
Shore, S. E., Vass, Z., Wys, N. L., & Altschuler, R. A. (2000). Trigeminal ganglion innervates the auditory
brainstem. *J Comp Neurol, 419*(3), 271-285. doi:10.1002/(sici)1096-
9861(20000410)419:3<271::aid-cne1>3.0.co;2-m
Sinex, D. G. (2005). Spectral processing and sound source determination. *Int Rev Neurobiol, 70*, 371-398.
doi:10.1016/S0074-7742(05)70011-8
Singh, N. C., & Theunissen, F. E. (2003). Modulation spectra of natural sounds and ethological theories of
auditory processing. *J Acoust Soc Am, 114*(6 Pt 1), 3394-3411. doi:10.1121/1.1624067
Solyga, M., & Barkat, T. R. (2021). Emergence and function of cortical offset responses in sound
termination detection. *Elife, 10*. doi:10.7554/eLife.72240
Sporns, O., Tononi, G., & Edelman, G. M. (2000). Connectivity and complexity: the relationship between
neuroanatomy and brain dynamics. *Neural Netw, 13*(8-9), 909-922. doi:10.1016/s0893-
6080(00)00053-8
Stiebler, I., & Ehret, G. (1985). Inferior colliculus of the house mouse. I. A quantitative study of tonotopic
organization, frequency representation, and tone-threshold distribution. *J Comp Neurol, 238*(1),
65-76. doi:10.1002/cne.902380106
Tamamaki, N., Yanagawa, Y., Tomioka, R., Miyazaki, J., Obata, K., & Kaneko, T. (2003). Green fluorescent
protein expression and colocalization with calretinin, parvalbumin, and somatostatin in the
GAD67-GFP knock-in mouse. *J Comp Neurol, 467*(1), 60-79. doi:10.1002/cne.10905
Tononi, G., Sporns, O., & Edelman, G. M. (1994). A measure for brain complexity: relating functional
segregation and integration in the nervous system. *Proc Natl Acad Sci U S A, 91*(11), 5033-5037.
doi:10.1073/pnas.91.11.5033
Vaithiyalingam Chandra Sekaran, N., Deshpande, M. S., Ibrahim, B. A., Xiao, G., Shinagawa, Y., & Llano,
D. A. (2021). Patterns of Unilateral and Bilateral Projections From Layers 5 and 6 of the Auditory
Cortex to the Inferior Colliculus in Mouse. *Front Syst Neurosci, 15*, 674098.
doi:10.3389/fnsys.2021.674098
Vogelstein, J. T., Packer, A. M., Machado, T. A., Sippy, T., Babadi, B., Yuste, R., & Paninski, L. (2010). Fast
nonnegative deconvolution for spike train inference from population calcium imaging. *J
Neurophysiol, 104*(6), 3691-3704. doi:10.1152/jn.01073.2009
Weckmann, K., Deery, M. J., Howard, J. A., Feret, R., Asara, J. M., Dethloff, F., . . . Turck, C. W. (2019).
Ketamine's Effects on the Glutamatergic and GABAergic Systems: A Proteomics and
Metabolomics Study in Mice. *Mol Neuropsychiatry, 5*(1), 42-51. doi:10.1159/000493425
Wehr, M., & Zador, A. M. (2003). Balanced inhibition underlies tuning and sharpens spike timing in
auditory cortex. *Nature, 426*(6965), 442-446. doi:10.1038/nature02116
Wenzel, M., Hamm, J. P., Peterka, D. S., & Yuste, R. (2017). Reliable and Elastic Propagation of Cortical
Seizures In Vivo. *Cell Rep, 19*(13), 2681-2693. doi:10.1016/j.celrep.2017.05.090
Willard, F. H., & Ryugo, D. K. (1983). Anatomy of the central auditory system. In J. F. Willot (Ed.), *The
auditory psychobiology of the mouse.* (pp. 201-304). Springfield, IL: Thomas.
Willott, James F. (2001). *Handbook of Mouse Auditory Research From Behavior to Molecular Biology* (1st
Edition ed.). Boca Raton: CRC Press.
Winer, J. A., Larue, D. T., Diehl, J. J., & Hefti, B. J. (1998). Auditory cortical projections to the cat inferior
colliculus. *J Comp Neurol, 400*(2), 147-174.
Wong, A. B., & Borst, J. G. G. (2019). Tonotopic and non-auditory organization of the mouse dorsal
inferior colliculus revealed by two-photon imaging. *Elife, 8*. doi:10.7554/eLife.49091

Wong, C., Chabot, N., Kok, M. A., & Lomber, S. G. (2015). Amplified somatosensory and visual cortical
projections to a core auditory area, the anterior auditory field, following early- and late-onset
deafness. *J Comp Neurol, 523*(13), 1925-1947. doi:10.1002/cne.23771
Xiong, B., Li, A., Lou, Y., Chen, S., Long, B., Peng, J., . . . Gong, H. (2017). Precise Cerebral Vascular Atlas in
Stereotaxic Coordinates of Whole Mouse Brain. *Front Neuroanat, 11*, 128.
doi:10.3389/fnana.2017.00128
Xu, Na   Fu, Zi-Ying   Chen,   Qi-Cai. (2014). The function of offset neurons in auditory information
processing. *Translational Neuroscience, 5*(4), 275-285. doi:https://doi.org/10.2478/s13380-014-
0235-5
Yaksi, E., & Friedrich, R. W. (2006). Reconstruction of firing rate changes across neuronal populations by
temporally deconvolved Ca2+ imaging. *Nat Methods, 3*(5), 377-383. doi:10.1038/nmeth874
Zhou, J., & Shore, S. (2006). Convergence of spinal trigeminal and cochlear nucleus projections in the
inferior colliculus of the guinea pig. *J Comp Neurol, 495*(1), 100-112. doi:10.1002/cne.20863

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194   **Figure 1:**

1195

Figure 1: A brief Illustration of the surgical procedures.

(A) A surface view of the mouse's skull showing the landmarks of the craniotomy over the IC.

(B) An image of the bone flap after finishing the drilling. (C-E) A cartoon and two images of the

brain surface showing the insertion and the pocket made by the diamond knife at the lateral

surface of the left IC. (F-H) A cartoon and two images of the brain surface show the insertion of

the microprism into its pocket to directly face the LC after displacing the blade of the diamond

knife. (I-K) A cartoon and two images of the brain surface show the final setup of the microprism

for two different animals. (L) An image of the brain surface shows the placement of a glass

window for the surface imaging of the DC. (M) A cartoon of the coronal section of the IC

demonstrating the different fields of views taken along the mediolateral axis parallel to the

curvature of the IC surface at different depths (1-5) from the pial surface. Cereb: Cerebellum,

Ctx: Cerebral cortex, DC: Dorsal cortex of the IC, IC: Inferior colliculus, ICC: Central nucleus of

the IC, Lam. Lambdoid, LC: Lateral cortex of the IC, Sag. Sagittal, SS: Sigmoid sinus, Surf:

Surface, TS: Transverse sinus.

1210

1211   **Figure 2:**

1212

Figure 2: The distribution of GABAergic cells on the LC(microprism), and DC imaged from the

dorsal surface.

All images were taken for the GFP signals exclusively expressed in GABAergic cells. (A-B) Low

magnification of fluorescence images for the GFP signals obtained from the IC surface with or

without the microprism, respectively. The dotted white line in B represented the medial and

lateral horizons of the IC. (C and E) The 2P images of the GABAergic cells on the LC$_{(microprism)}$

show the GABAergic modules across multiple animals. (D and F) The 2P images of the

GABAergic cells on either the dorsolateral surface of the DC imaged directly from its dorsal

surface showing the GABAergic modules across multiple animals (red boxes) or its dorsomedial

surface showing a homogenous distribution of GABAergic cells of different sizes with no signs

of the GABAergic modules across multiple animals (blue boxes). (G) A low magnification

fluorescence image of the IC surface with the microprism showing a laser lesion spot made

through the microprism to validate the position of the microprism relative to the IC. (H) A

histological coronal section at the level of the IC shows that the laser lesion made by the

microprism was located at the sagittal surface of the LC$_{(microprism)}$. (I-J) Low-magnification

fluorescence images of the IC surface from the dorsal surface show a laser lesion spot made on

the dorsolateral surface of the LC on the GABAergic module. The dotted white lines represent

the alignment of the laser lesion (dotted red circle) to the location of the GABAergic module at

the dorsolateral surface. (K) A histological coronal section at the level of the IC shows that the

laser lesion made on the IC surface targeted the GABAergic module at the dorsolateral portion

of the LC. All solid irregular white lines were made at the border of the GABAergic modules; D:

Dorsal, IC: Inferior colliculus, L: Lateral, R: Rostral. Cereb: Cerebellum, TS: Transverse sinus.

1235

1236

1237

1238

1239 **Figure 3:**

1240

Figure 3: 2P expression of jRGECO1a along with evoked signals in the DC imaged from

the dorsal surface and LC(microprism).

(A-B) The 2P images of the jRGECO1a expression on the DC imaged from the dorsal surface

and the LC(microprism), respectively. (C-D) The time traces of the evoked calcium signals by either

different frequencies and amplitudes of pure tones or different overall amplitudes and rates of

AM of broadband noise, respectively, which were obtained from two different cells on the DC

(top traces) or the LC(microprism) (bottom traces); The length of the colored bars (top) indicates the

intensity of the stimulus, while the color indicates the tone or AM frequency depending on the

color scale (bottom). AM-noise: Amplitude modulated broadband noise, D: Dorsal, DC: Dorsal

cortex, L: Lateral, LC: Lateral cortex, R: Rostral.

1251

1252

1253

1254

Figure 3-figure supplement 1:

 Figure 3-figure supplement 1:

 (A) Image of the IC showing the immunostaining to NueN (blue-column 1), GAD67-GFP (green-

 column 3), jRGECO1a (red-column 3), merged image for the three proteins (column 4), and

 merged image for jRGECO1a (magenta) and GFP (green) – dotted lines represents the central

1261   nucleus of the IC in all channels as well as the modules in the green channel. (B) A magnified

1262   view of the DC in A – dotted line represents the central nucleus of the IC (Icc). (C) A magnified

1263   view of the LC in A – dotted line represents one of the GABAergic modules of the LC. (D) A

1264   magnified view of the white square in C shows some GABAergic cells (blue arrows) and some

1265   nonGABAergic cells expressing jRGECO1a (white arrows). D: dorsal, L: lateral.

1266

1267

1268

1269

1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

Figure 4:

Tone freq (kHz) 5.0 ▬▬ ▬▬ 40.0

Figure 4: The auditory cellular organization of the LC(microprism) vs. DC imaged from the

dorsal surface.

(A and C) The 2P images of GFP (green) and jRGECO1a expression (red) on either the

LC(microprism) or the DC imaged from the dorsal surface, respectively. (B and D) The pseudocolor

images show the responsive cells to the pure tone of different combinations of frequencies and

levels on the surface of the LC(microprism) or the DC imaged from its dorsal surface, respectively.

D: Dorsal, L: Lateral, R: Rostral.

1295

1296   **Figure 4-figure supplement 1**

1297

1298   **Figure 4-figure supplement 1:**

1299   (A-B) Linear and nonlinear quadratic regression fit between the BTFs and the locations of the

1300   cells of the $LC_{(microprism)}$ and the DC, respectively.

1301

1302

1303

1304

1305

1306

1307

1308

1309   **Figure 5:**

1310

Figure 5: The cellular response to pure tones and AM-noise of the LC$_{(microprism)}$ vs. DC

imaged from the dorsal surface.

(A) The cumulative distribution function of the best tone frequencies of all cells collected either

from the LC imaged via microprism (blue line) or the DC imaged directly from its dorsal surface

(red line) (Mann-Whitney Test: z = -11.34, p < 0.001, a median of best-tuned frequency = 5 and

7.1 Hz for the DC (2484 cells from 4 animals) and the LC$_{(microprism)}$ (445 cells from 7 animals),

respectively, *p<0.05, DC vs LC$_{(microprism)}$). (B) A bar graph showing the fractions of cells

responding to only pure tone (Tone-sel), only AM-noise (Noise-sel), or to both nonselectively

(Non-sel) on the LC$_{(microprism)}$ (7 animals) (one way-ANOVA, $f_{(2,18)}$ = 14.9, p = 1.6x10^{-4}, Fisher

posthoc test: p = 2.0x10^{-4} and 1.3x10^{-4} for Noise-sel and Non-sel vs Tone-sel, respectively, p =

0.84 for Noise-sel vs Non-sel, % of responsive cells ± SEM = 5±1%, 46±7%, and 48±7% for

Tone-sel, Noise-sel, and Non-sel, respectively, #p<0.05 vs Tone-sel). (C) Box graphs showing

the fractions of Tone-sel, Noise-sel, and Non-sel cells within modules and matrix of the

LC$_{(microprism)}$. (D) A bar graph showing the fractions of Tone-sel, Noise-sel, or Non-sel cells on

the DC imaged directly from the dorsal surface (4 animals) (one way-ANOVA, $f_{(2,9)}$ = 86.9, p =

1.2x10^{-6}, Fisher posthoc test: p = 1.5x10^{-4} and 3.4x10^{-7} for Noise-sel and Non-sel vs Tone-sel,

respectively, p = 7.0X10^{-5} for Noise-sel vs Non-sel, % of responsive cells ± SEM = 0.6±0.2%,

32±4%, and 67±4% for Tone-sel, Noise-sel, and Non-sel, respectively, #p<0.05 vs Tone-sel and

$p<0.05 vs Noise-sel). (E and F) Bar graphs showing the percentage of spectrally (two-way

ANOVA: $f_{(1, 4, 20)}$ = 50.3, 0.0, and 4.0 – p = 7.1x10^{-7}, 1.0, and 0.01 for the region, sound level,

and interaction, respectively, Fisher post hoc, p = 5.1x10^{-6}, 4.9x10^{-4}, 0.02, 0.058, and 0.14 for

modules vs. matrix at 40, 50, 60, 70, and 80 dB, respectively, % of spectral modulated cells ±

SEM (modules vs matrix) = 9±1 vs 90±1% at 40 dB, 22±4 vs 77±4% at 50 dB, 33±10 vs

66±10% at 60 dB, 36±13 vs 63±13% at 70 dB, and 43±11 vs 56±11% at 80 dB, n = 3 animals)

or temporally (two-way ANOVA: $f_{(1, 4, 20)}$ = 19.3, 0.0, and 1.2 – p = 7.1x10^{-7}, 1.0, and 0.32 for the

region, sound level, and interaction, respectively, % of temporally modulated cells ± SEM = 38±

3 and 61± 3% for modules vs matrix, n = 3 animals) modulated cells, respectively, across

different sound levels in modules (green bars) vs. matrix (blue bars) imaged via microprism,

(**p<0.05, matrix vs. modules). (G and H) Box plots showing the mean (black lines) and the

distribution (colored dots) of the SMI (two-way ANOVA: $f_{(1, 4, 14583)}$ = 48.4, 210.4, and 21.3 – p <

0.001 for the region, sound level, and interaction, respectively, Fisher post hoc test, p = 6.2x10^{-}

^{4}, 4.2x10^{-5}, <0.001, and < 0.001 for LC vs. DC at 40, 60, 70, and 80 dB respectively, mean of

SMI ± SEM: 0.16±0.01 and 0.08±0.003 for LC vs DC, n = 1020 cells from 7 animals

(LC$_{(microprism)}$), and 13574 cells from 4 animals (DC)) and TMI (two way ANOVA: $f_{(1, 4, 16139)}$ =

431.4, 59.1, and 8.3 – p < 0.001 for the region, sound level, and interaction, respectively, Fisher

post hoc test, p = 0.002, <0.001, < 0.001, <0.001, and <0.001 for LC vs. DC at 40, 50, 60, 70,

and 80 dB, respectively, mean of TMI ± SEM: 0.35±0.006 and 0.21±0.001 for LC vs DC, n =

1441 cells from 7 animals (LC$_{(microprism)}$), and 14708 cells from 4 animals (DC)), respectively,

across different sound levels in the LC$_{(microprism)}$ (blue) vs. the DC (red), *p<0.05, LC vs. DC). DC:

Dorsal cortex, LC: Lateral cortex. SMI: spectral modulation index, TMI: temporal modulation

index.

1352

1353

1354

1355

1356

1357

1358

1359 **Figure 6:**

1360

1361 **Figure 6: The waveforms of the somatosensory vs. acoustic responses of the LC(microprism)**

1362 **and the DC. imaged via microprism.**

1363 Examples of different cell types based on their responses to auditory or somatosensory

1364 stimulations as well as their response timing to the auditory stimulation as indicated by

1365 waveforms of the evoked calcium signals.

1366

1367

1368

1369

1370

1371

1372

1373

1374

1375

1376

1377

1378

1379

1380

Figure 7:

 Figure 7: Types of somatosensory and auditory responses from the LC neurons imaged

 via microprism.

 (A) The 2P images of GFP (green) and jRGECO1a expression (magenta) in the LC$_{(microprism)}$

 show the modules within white irregular lines. (B, D, and F) The pseudocolor images show the

auditory (white circles) or somatosensory (blue circles) responsive cells of the $LC_{(microprism)}$ as

well as the merged response (green circles), respectively. (C and E) The pseudocolor images

show onset (ON) (blue circles), offset (OFF) (green circles), or onset/offset (ON/OFF) (white

circles) responsive cells of the $LC_{(microprism)}$ to auditory and somatosensory stimuli, respectively.

(G) The pseudocolor images show enhanced (white circles) and suppressive response (blue

circles) of the auditory responsive cells of the $LC_{(microprism)}$ following the bimodal stimulation

based on their response index (RI). (H) A bar graph showing the percentage of responsive cells

to auditory stimulation only or auditory selective cells (Aud-Sel, yellow bars), to somatosensory

stimulation only or somatosensory selective cells (Som-sel, blue bars), or nonselectively to both

auditory and somatosensory stimulations (Aud/Som-nonsel, black bars) in modules vs matrix

(Chi-Square test, χ^2 = 33.1, p = $3.5x10^{-6}$, % of responsive cells = 74%, 8%, and 18% for Aud-

sel, Som-sel, and Aud/Som-nonsel, respectively at (Modules, 235 cells) vs 92%, 2%, and 6% for

Aud-sel, Som-sel, and Aud/Som-sel, respectively at (matrix, 297 cells) – 4 animals, **p<0.05 vs

modules). (I) A bar graph showing the percentage of all cells having auditory responses (Aud.

resp) or those having somatosensory responses (Som. resp) in modules vs matrix (Chi-Square

test, χ^2 = 26.0, p = $9.3x10^{-5}$, % of responsive cells (Modules vs matrix) = 43% vs 57% for (Aud-

resp, 509 cells) and 73% vs 27% for (Som. resp, 84 cells) – 4 animals, **p<0.05 vs modules).

(J) A bar graph showing the percentage of auditory responsive cells with onset (Aud-ON) vs

those with offset (Aud-OFF) responses in modules vs matrix (Chi-Square test, χ^2 = 120.5, p

<0.001, % of responsive cells (Modules vs matrix) = 28% vs 72% for (Aud-ON, 390 cells) and

90% vs 10% for (Aud-OFF, 93 cells) – 4 animals, **p<0.05 vs modules). (K) A bar graph

showing the percentage of auditory responsive cells with suppressed vs those with enhanced

responses following bimodal stimulation in modules vs matrix (Chi-Square test, χ^2 = 12.9, p =

0.04, % of responsive cells = 54% vs 46% for suppressed vs enhanced (modules, 217 cells)

and 70% vs 30% for suppressed vs enhanced (matrix, 292 cells) – 4 animals, **$p<0.05$ vs

modules). D: Dorsal, R: Rostral.

1413

1414

1415

1416

1417

1418

1419

1420

1421

1422

1423

1424

1425

1426

1427

1428

1429

1430

Figure 8:

Figure 8: The somatosensory vs. auditory responses over the DC neurons.

1433

(A) A low magnification of the image obtained from the IC surface. (B-D) The top row: The 2P

images of GFP (green) and jRGECO1a expression (magenta) of three main regions

(dorsolateral, dorsal, and dorsomedial) of the DC surface imaged from the dorsal surface

showing the modules within irregular white lines in the dorsolateral region, the bottom row: The

pseudocolor images show the somatosensory responsive cells (blue circles) after

somatosensory stimulations. (E) The pseudocolor images show the responsive cells of the

LC$_{(dorsal\ surface)}$ to auditory (white circles), somatosensory (blue circles), or both stimulations

(green circles). (F) The pseudocolor images show auditory onset (Aud-ON – blue circles), offset

(Aud-OFF, green circles), or onset/offset (Aud-ON/OFF, white circles) responsive cells of the LC

imaged from the dorsal surface to auditory stimulation. (G) The pseudocolor images show the

enhanced (white circles) and suppressive (blue circles) response of the auditory responsive

cells of the LC imaged from the dorsal surface following the bimodal stimulation based on their

response index (RI). (H) A box plot shows the percentage of somatosensory responsive cells in

the LC$_{(microprism)}$, the LC$_{(dorsal\ surface)}$ or the DC (one-way ANOVA, $p = 0.02$, the percentage of

somatosensory responsive cells mean ± SEM = 1.3 ± 0.83% (DC, 4 animals), 8.2 ± 2.4%

(LC$_{(dorsal\ surface)}$, 4 animals), and 8.1 ± 0.64% (LC$_{(microprism)}$, 4 animals, *$p<0.05$, vs. DC). (I) A bar

graph showing the percentage of responsive cells to auditory stimulation only or auditory

selective cells (Aud-Sel, yellow bars), to somatosensory stimulation only or somatosensory

selective cells (Som-sel, blue bars), or nonselectively to both auditory and somatosensory

stimulations (Aud/Som-nonsel, black bars) in modules vs matrix (Chi-Square test, $\chi^2 = 86.1$, p

<0.001, % of responsive cells = 69%, 9%, and 22% for Aud-sel, Som-sel, and Aud/Som-nonsel

(modules, 152 cells) and 95%, 1%, and 4% % for Aud-sel, Som-sel, and Aud/Som-nonsel

(matrix, 573 cell–) - 4 animals, **$p<0.05$ vs modules). (J) A bar graph showing the percentage of

all cells having auditory responses (Aud. resp) or those having somatosensory responses (Som.

resp) in modules vs matrix (Chi-Square test, $\chi^2 = 66.5$, p <0.001, % of responsive cells

(Modules vs matrix) = 20% vs 80% for (Aud. resp, 705 cells) and 61% vs 39% for (Som. resp,

79 cells) - 4 animals, **$p<0.05$ vs modules). (K) A bar graph showing the percentage of auditory

responsive cells with onset (Aud-ON) vs those with offset (Aud-OFF) responses in modules vs

matrix (Chi-Square test, χ^2 = 500.2, p <0.001, % of responsive cells (Modules vs matrix) = 5%

vs 95% for (Aud-ON, 584 cells) and 95% vs 5% for (Aud-OFF, 116 cells) - 4 animals, **p<0.05

vs modules). (L) A bar graph showing the percentage of auditory responsive cells with

suppressed vs those with enhanced responses following bimodal stimulation in modules vs

matrix (Chi-Square test, χ^2 = 18.5, p = 1.6×10^{-5}, % of responsive cells = 47% vs 53% for

suppressed vs enhanced (modules, 138 cells) and 67% vs 33% for suppressed vs enhanced

(matrix, 567 cells) - 4 animals, **p<0.05 vs modules). Cereb: Cerebellum, Ctx: Cerebral cortex,

M: medial, R: Rostral, TS: Transverse sinus.

1470

1471

1472

1473

1474

1475

1476

1477

1478

1479

1480

1481

1482

1483　　**Figure 9:**

1484

Figure 9: The acoustic response from LC(microprism) and DC imaged from the dorsal surface

in awake animals.

(A and C) The 2P images of GFP (green) and jRGECO1a expression (magenta) on either the

$LC_{(microprism)}$ surface imaged via the microprism or the DC imaged directly from the dorsal

surface, respectively showing the modules within irregular white lines. (B and D) The

pseudocolor images show the responsive cells to the pure tone of different combinations of

frequencies and levels of the $LC_{(microprism)}$ or the DC imaged directly from the dorsal surface,

respectively. D: Dorsal, L: Lateral, R: Rostral.

1493

1494

1495

1496

1497

1498

1499

1500

1501

1502

1503

1504

1505

1506

1507

1508 **Figure 9-figure supplement 1:**

1509

1510 **Figure 9-figure supplement 1:**

1511 (A-B) Linear and nonlinear quadratic regression fit between the BTFs and the locations of the

1512 cells of the LC$_{(microprism)}$ and the DC, respectively.

1513

1514

1515

1516

1517

1518

1519   **Figure 10:**

1520

Figure 10: The cellular response to pure tones and AM-noise of the LC$_{(microprism)}$ vs. DC

imaged from the dorsal surface in awake animals.

(A) The cumulative distribution function of the best tone frequencies of all cells collected either

from the LC imaged via microprism (blue line) or the DC imaged directly from its dorsal surface

(red line) (Mann-Whitney test, z = -4.8, p=1.6x10^{-6}, a median of best-tuned frequency = 5 and

10 kHz for the DC (767 cells from 3 animals) and the LC$_{(microprism)}$ (371 cells from 3 animals),

respectively, *p<0.05 vs DC) (B) A bar graph showing the fractions of cells responding to only

pure tone (Tone-sel), only AM-noise (Noise-sel), or to both nonselectively (Nonsel) on the

LC$_{(microprism)}$ (3 animals) (one way ANOVA, f$_{(2,6)}$ = 41.6, p = 3.0x10^{-4}, Fisher posthoc test: p = 0.04

and 1.1x10^{-4} for Noise-sel and Non-sel vs Tone-sel, respectively, p = 7.3x10^{-4} for Noise-sel vs

Non-sel, % of responsive cells ± SEM = 6±3%, 24±6%, and 68±5 for Tone-sel, Noise-sel, and

Non-sel, respectively, #p<0.05 vs Tone-sel and $p<0.05 vs Noise-sel). (C) Box graphs showing

the fractions of Tone-sel, Noise-sel, and Non-sel cells within modules and matrix of the

LC$_{(microprism)}$. (D) A bar graph showing the fractions of Tone-sel, Noise-sel, or Non-sel cells on

the DC imaged directly from the dorsal surface (3 animals) (one way ANOVA, f$_{(2,6)}$ = 12.8, p =

0.006, Fisher posthoc test: p = 0.002 and 0.12 for Noise-sel and Non-sel vs Tone-sel,

respectively, p = 0.01 for Noise-sel vs Non-sel, % of responsive cells ± SEM = 9±1%, 61±9%,

and 28±8 for Tone-sel, Noise-sel, and Non-sel, respectively, #p<0.05 vs Tone-sel and $p<0.05

vs Noise-sel). (E and F) Bar graphs showing the percentage of spectrally (two-way ANOVA:

f$_{(1,4,16)}$ = 173.0, 0, and 1.8 - p = 5.3x10^{-10}, 1.0, and 0.17 for the region, sound level, and

interaction, respectively, % of spectrally modulated cells ± SEM = 23±2% vs 76±2% for modules

vs matrix – 3 animals, **p<0.05 vs modules) or temporally (two-way ANOVA: f$_{(1,4,16)}$ = 65.9, 0,

and 2.2 - p = 4.6x10^{-7}, 1.0, and 0.11 for the region, sound level, and interaction, respectively, %

of temporally modulated cells ± SEM = 23±2% vs 76±2% for modules vs matrix, n = 3 animals,

**p<0.05 vs modules) modulated cells, respectively, across different sound levels in modules

(green bars) vs. matrix (blue bars) imaged via microprism, (**p<0.05, matrix vs. modules).

(G and H) Box plots showing the mean (black lines) and the distribution (colored dots) of the

SMI (two-way ANOVA: $f_{(1,4,2411)}$ = 1.4, 15.3, and 8.3 - p = 0.22, 2.1×10^{-12}, and 1.1×10^{-6} for the

region, sound level, and interaction, respectively, SMI ± SEM = -0.04±0.02 vs -0.06±0.01 for

$LC_{(microprism)}$ vs DC – n = 619 cells from 3 animals ($LC_{(microprism)}$), and 1826 cells from 3 animals

(DC)) and TMI (two-way ANOVA: $f_{(1,4,2869)}$ = 36.2, 29.3, and 3.0 - p = 1.9×10^{-9}, 6.6×10^{-24}, and

0.01 for the region, sound level, and interaction, respectively, SMI ± SEM = 0.2±0.01 vs

0.32±0.007 for $LC_{(microprism)}$ vs DC – n = 781 cells from 3 animals ($LC_{(microprism)}$), and 2098 cells

from 3 animals (DC), *p<0.05 vs DC). DC: Dorsal cortex, LC: Lateral cortex. SMI: spectral

modulation index, TMI: temporal modulation index.

1556

1557

1558

1559

1560

1561

1562

1563

1564

1565

1566

1567   **Figure 10-figure supplement 1:**

1568

1569

Figure 10-figure supplement 1: The RFS of DC and LC neurons at different experimental

preparations:

The cumulative distribution function of the RFS of all cells collected from DC or LC under

different experimental preparations (Kruskal-Wallis ANOVA, χ^2 = 346.9, p < 0.001, posthoc

Dunn's test; *(DC vs LC): z = -6.4 and -4.47 with p <0.001 and 4.6x10^{-5} under anesthesia and

awake preparations, respectively; #(anesthesia vs awake): z = 13.6 and 7.8 with p <0.001 for

DC and LC, respectively; mean ± SEM = 5.1±0.08, 4.5±0.2, 3.07±0.1, and 2.0±0.08 for

(DC/anesthesia, median = 4, 2487 cells / 4 animals), (LC/anesthesia, median = 3, 472 cells / 7

animals), (DC/awake, median = 2, 647 cells / 3 animals), (LC/awake, median = 2, 249 cells / 3

animals).

1580

1581

1582

1583

1584

1585

1586

1587

1588

1589

1590

1591

1592

1593

1594

1595

1596

1597

Figure 11:

Figure 11: The somatosensory response from LC$_{(microprism)}$ in awake animals.

(A, C, and E) The pseudocolor images show the auditory (white circles) or somatosensory (blue

circles) responsive cells of the LC$_{(microprism)}$ as well as the merged response (green circles),

respectively. (B and D) The pseudocolor images show onset (ON) (blue circles), offset (OFF)

(green circles), or onset/offset (ON/OFF) (white circles) responsive cells of the LC$_{(microprism)}$ to

auditory and somatosensory stimuli, respectively. (F) The pseudocolor images show enhanced

(white circles) and suppressive response (blue circles) of the auditory responsive cells of the

LC$_{(microprism)}$ following the bimodal stimulation based on their response index (RI). (G) A bar graph

showing the percentage of responsive cells to auditory stimulation only or auditory selective

cells (Aud-Sel, yellow bars), to somatosensory stimulation only or somatosensory selective cells

(Som-sel, blue bars), or nonselectively to both auditory and somatosensory stimulations

(Aud/Som-nonsel, black bars) in modules vs matrix (Chi-Square test, χ^2 = 42.5, p = 4.4x10^{-8}, %

of responsive cells = 49%, 29%, and 22% for Aud-sel, Som-sel, and Aud/Som-nonsel (Modules,

55 cells) and 89%, 3%, and 8% for aud-sel, som-sel, and aud/som-nonsel (matrix, 147 cells) - 3

animals, **p<0.05 vs modules). (H) A bar graph showing the percentage of all cells having

auditory responses (Aud. resp) or those having somatosensory responses (Som. resp) in

modules vs matrix (Chi-Square test, χ^2 = 30.2, p = 1.2X10^{-6}, % of responsive cells (Modules vs

matrix) = 21% vs 79% for (aud-resp, 182 cells), 64% vs 36% for (som-resp, 44 cells) - 3

animals, **p<0.05 vs modules). (I) A bar graph showing the percentage of auditory responsive

cells with onset (Aud-ON) vs those with offset (Aud-OFF) responses in modules vs matrix (Chi-

Square test, χ^2 = 2.8, p = 0.41, % of responsive cells (Modules vs matrix) = 21% vs 79% for

(aud-ON, 176 cells), 50% vs 50% for (aud-OFF, 6 cells) - 3 animals, **p<0.05 vs modules). (J) A

bar graph showing the percentage of auditory responsive cells with suppressed vs those with

enhanced responses following bimodal stimulation in modules vs matrix (Chi-Square test =

31.9, p = 5.2x10^{-7}, matrix: The percentage of responding cells = 95% vs. 5% for suppressed vs

enhanced (Modules, 40 cells) and 70% vs 30% for suppressed vs enhanced (matrix, 142 cells)

– 3 animals, **p<0.05 vs modules). D: Dorsal, R: Rostral.

1627

1628

1629

1630

1631

1632

1633

1634

1635

1636

1637

1638

1639

1640

1641

1642

1643

1644

1645 **Figure 11-figure supplement 1:**

1646

Figure 11-figure supplement 1: The activity of the nonresponsive cells of the LC(microprism)

in awake animals.

(A) The pseudocolor image shows the nonresponsive (red circles) vs responsive (white circles)

cells to the auditory stimulation with unmodulated broadband noise at the LC(microprism). (B) The

pseudocolor image shows the nonresponsive (red circles) vs responsive (white circles) cells to

the auditory stimulation with unmodulated broadband noise or pure tones (blue arrows) at the

LC(microprism). (C) Time traces of the spontaneous activity of four nonresponsive cells (yellow

numbers in B).