Complex third-party effects in the *Dictyostelium-Paraburkholderia* symbiosis:

prey bacteria that are eaten, carried, or left behind

Trey J. Scott*

David C. Queller

Joan E. Strassmann

Department of Biology, Washington University in St. Louis

*Corresponding Author; E-mail: tjscott@wustl.edu

Acknowledgements

This material is based upon work supported by the National Science Foundation under grants DEB-1753743, IOS-1656756, and DEB-2237266. We thank Tyler Larsen for comments on an early draft of the manuscript. We also thank Debbie Brock and the rest of the Strassmann/Queller lab group for feedback during the development and execution of this project.

Conflict of interest

We have no conflicts of interest to declare.

Keywords: Symbiosis, cooperation, conflict, Dictyostelium discoideum, Paraburkholderia

Abstract

Symbiotic interactions may change depending on the abundance of third parties

like predators, prey, or pathogens. Third-party interactions with prey bacteria are central

to the symbiosis between *Dictyostelium discoideum* social amoeba hosts and

Paraburkholderia facultative bacterial symbionts. Symbiosis with inedible

Paraburkholderia allows host *D. discoideum* to carry prey bacteria through the dispersal

stage where host amoebae aggregate and develop into fruiting bodies that disperse spores.

Carrying prey bacteria benefits hosts when prey bacteria are scarce but harms hosts when

prey bacteria are plentiful. Symbiont-carrying hosts leave some bacteria behind; this

could explain the harm to hosts if left-behind bacteria include uneaten prey bacteria.

Thus, understanding both benefits and costs in this symbiosis requires measuring

how many prey bacteria are eaten, carried, and left behind by infected hosts. We

found that *Paraburkholderia* infection makes hosts leave behind both symbionts and prey

bacteria. However, the number of prey bacteria left uneaten was small and did not explain

why infected hosts produced fewer spores than uninfected hosts. Turning to the bacteria

that are carried, we found that hosts carry more prey bacteria after developing in prey-

poor environments than in prey-rich ones. This suggests that carriage is actively modified

to ensure hosts have prey in the harshest conditions. Our results show that multifaceted

interactions with third parties shape the evolution of symbioses in complex ways.

Introduction

The fitness effects of symbiotic interactions can change depending on the environment (Bronstein, 1994; Chamberlain *et al.*, 2014; Horas *et al.*, 2022; Scott *et al.*, 2022). One crucial component of the environment that often selects for symbiotic interactions is a third species that interacts with hosts or symbionts (Palmer *et al.*, 2008; Wendling *et al.*, 2017; Wood *et al.*, 2018; Hafer-Hahmann & Vorburger, 2020; Cassidy *et al.*, 2022). For example, in the symbiosis between ants and *Acacia* plants, ants benefit *Acacia* by fending off herbivores. However, when herbivores were prevented from accessing *Acacias*, ordinarily mutualistic interactions between ants and *Acacias* shifted towards antagonism because ants no longer provided a benefit to host *Acacias* (Palmer *et al.*, 2008). Such shifts can influence whether host and symbiont fitness interests are aligned or in conflict (Keeling & McCutcheon, 2017; Iwai *et al.*, 2019). Fitness alignment or conflict then shapes the evolution of these interactions (Scott & Queller, 2019; O'Brien *et al.*, 2021). However, the details of how third parties affect the fitness interests of symbiotic partners are not well understood for many kinds of symbioses (Chamberlain *et al.*, 2014).

An important model of microbial symbiosis is the social amoeba *Dictyostelium discoideum*, that is famous for its complex and well-studied multicellular life cycle (Kessin, 2001; Jahan *et al.*, 2021). *D. discoideum* is a soil amoeba that feeds on many bacteria (Raper, 1937; Brock *et al.*, 2018). Upon starvation, amoebae aggregate to form a multicellular fruiting body that disperses spores to new soil patches (Kessin, 2001). Development of the fruiting body progresses through well-studied stages starting with aggregation of individual amoebae which then become motile slugs (Medina *et al.*,

2019). Slugs further develop into a fruiting body. During fruiting body development,

about 20% of the cells die and become stalk while the remaining cells develop into spores

that sit atop the stalk in a structure called a sorus (Strassmann & Queller, 2011).

D. discoideum has symbioses with certain facultatively intracellular

Paraburkholderia bacteria (DiSalvo *et al.*, 2015; Brock *et al.*, 2020). In natural isolates,

around 25% of wild-collected *D. discoideum* are infected (Haselkorn *et al.*, 2019). There

are three symbiont species – *P. agricolaris*, *P. hayleyella*, and *P. bonniae* – and we will

refer to them collectively in this paper as either symbionts or simply *Paraburkholderia*

(though readers should understand that there are many species of *Paraburkholderia* not

involved in this symbiosis).

These symbionts are adapted for interacting with amoebae (Shu *et al.*, 2018a; b;

Brock *et al.*, 2020). Symbionts move towards hosts (Shu *et al.*, 2018b) and make host

phagosomes less acidic so that the symbionts are not digested (Tian *et al.*, 2022). *P.*

hayleyella and *P. bonniea* also have reduced genomes relative to *P. agricolaris* and most

other *Paraburkholderia* (Brock *et al.*, 2020; Noh *et al.*, 2022) and such reduction is often

found in endosymbionts and pathogens that have been associated closely with their hosts

during a long evolutionary history (McCutcheon & Moran, 2012). Some *D. discoideum*

hosts may also be adapted to their symbionts. Hosts infected by *P. hayleyella* appear to

suffer less from *P. hayleyella* toxicity than hosts that are naturally infected by *P.*

agricolaris (Shu *et al.*, 2018a).

Fitness of *D. discoideum* hosts and *Paraburkholderia* symbionts is affected by

interactions with a third set of organisms, the various prey bacteria that are eaten by host

amoebae (Brock *et al.*, 2011, 2018; DiSalvo *et al.*, 2015; Scott *et al.*, 2022).

Paraburkholderia symbionts are largely inedible by hosts but they cause host carriage of

additional, edible bacteria (Brock *et al.*, 2011; DiSalvo *et al.*, 2015; Khojandi *et al.*,

2019). Edible prey bacteria are mostly carried extracellularly in the sorus while

Paraburkholderia symbionts are mostly carried inside host spores (Khojandi *et al.*,

2019). When host spores disperse to locations where prey bacteria are rare, they benefit

from *Paraburkholderia* infection because they are able to seed new prey bacteria

populations (Brock *et al.*, 2011; DiSalvo *et al.*, 2015; Scott *et al.*, 2022). When hosts

disperse to prey-rich conditions, infected hosts do not gain from carrying prey bacteria

and instead experience a cost producing fewer spores relative to uninfected *D.*

discoideum (Brock *et al.*, 2011; DiSalvo *et al.*, 2015). Higher *Paraburkholderia* densities

impose higher costs on hosts but again this is affected by third-party prey, with the effect

being stronger in pre-rich conditions (Scott *et al.*, 2022).

The third-party bacteria can also affect *Paraburkholderia* fitness due to

competition (Medina *et al.*, 2023). One consequence is that symbionts reach higher

densities when they are carried to locations with fewer prey bacteria (Scott *et al.*, 2022).

Thus prey-poor conditions for the host are low-competition conditions for the symbiont

(though for consistency, we will stick to calling them prey-poor and prey-rich conditions

rather than low-competition and high-competition conditions).

While the importance of prey bacteria has been well established for this symbiosis

(DiSalvo *et al.*, 2015; Khojandi *et al.*, 2019; Scott *et al.*, 2022), there are still many

questions about the effects of prey bacteria during development of *D. discoideum*. In this

study, it is the disposition of prey bacteria during host development that is of interest –

should they be eaten, carried, or left behind?

Leaving prey bacteria behind would not seem to be adaptive, but this appears to

happen when hosts are infected by *Paraburkholderia* (Brock *et al.*, 2011, 2016b).

Initially, before the distinct roles of inedible symbionts and edible prey were

appreciated, bacteria left behind on plates was taken as a possible sign of "prudent

predation" by host amoebae — if some bacteria were going to be saved for carriage, hosts

cannot eat all the available prey and may stop feeding and start developing earlier (Brock

et al., 2011). It was hypothesized that hosts leaving prey bacteria uneaten might explain

why carriage is costly in some conditions (Brock *et al.*, 2011). Hosts that were prudent

predators left prey uneaten and missed out on potential growth and proliferation.

The essential role of *Paraburkholderia* in the symbiosis, including causing hosts

to leave behind bacteria is now known (DiSalvo *et al.*, 2015; Brock *et al.*, 2016b). At

least some of the bacteria left behind are not prey bacteria but inedible *Paraburkholderia*

symbionts (Scott *et al.*, 2022). Because of this new information, the prudent predation

hypothesis should be re-evaluated by determining the relative abundances of left behind

prey bacteria and *Paraburkholderia* symbionts. The fitness consequences of leaving prey

behind, in terms of spore production and development time, should also be evaluated.

While infected hosts may leave some prey bacteria uneaten, they also gain the

ability to carry prey bacteria along with the dispersing host spores in sori (Brock *et al.*,

2011; DiSalvo *et al.*, 2015; Khojandi *et al.*, 2019). It is unknown whether the number of

carried prey bacteria changes in different environments. If carriage is passive, we expect

hosts to carry more prey after developing in a prey-rich environment and fewer prey

bacteria after developing in a prey-poor environment.

Alternatively, the ability to carry prey bacteria could reflect the evolutionary

interests of hosts and symbionts. Since soil environments tend to be spatially and

temporally structured (Sun *et al.*, 2003; Vos *et al.*, 2013), developing in a prey-poor

environment may be associated with an increased probability that the next environment

will also tend to be prey-poor. If this is the case, it would be more adaptive for hosts if

they carried more prey bacteria after developing in a prey-poor environment than in a

prey-rich environment. This may also be adaptive for the *Paraburkholderia* symbionts

since it allows better survival of their hosts.

We investigate three questions about the complex role of prey bacteria in the

symbiosis between *D. discoideum* and two commonly studied *Paraburkholderia*

symbionts, *P. agricolaris* and *P. hayleyella* (Shu *et al.*, 2018a; Garcia *et al.*, 2019; Scott

et al., 2022; Medina *et al.*, 2023). We first re-evaluate some of the ideas behind the

prudent predation hypothesis. Second, we ask whether any prey bacteria that are left

uneaten results in costs due to delayed development or reduced spore production. Lastly,

we turn to whether the number of prey bacteria carried inside sori changes between prey-

poor and prey-rich environments.

Methods

Clones and Culturing Methods

In our experiments, we used several types of infected and uninfected hosts (Table

1). We used two types of infected host, naturally infected hosts (n=7 for each symbiont

species) and, to better control for infection levels, cured-and-reinfected hosts (n=4 for

each symbiont species). The naturally infected hosts were compared to naturally

uninfected controls (n=6) and the cured-and-reinfected host were compared to a different

set of naturally uninfected controls (n=4). To better control for genetics, cured-and-

reinfected hosts were also compared to their cured clonemates (n=4 for each symbiont

species). To generate cured-and-reinfected clones, naturally infected clones were cleared

of their natural infections with antibiotics and then reinfected at 1:1000 ratios of

Paraburkholderia to prey bacteria to control infection densities (Scott *et al.*, 2022). Data

reported here for cured-and-reinfected hosts and their uninfected controls come from

experiments performed in Scott et al. (2022) but here we include additional measures not

included in the original study. Naturally uninfected controls in this comparison were also

treated with antibiotics to control for antibiotic treatment. We stored all prepared clones

in a -80 C freezer for repeated use.

To remove any effects of being in the freezer and to ensure that infected amoebae

carried *K. pneumoniae* prey bacteria, we grew amoebae through one round of feeding and

fruiting body formation and then collected spores to initiate our experiments. This step

may allow infection densities time to equilibrate (Miller *et al.*, 2020). Amoebae were

grown from frozen spores at room temperature on 100 x 15mm plates filled with 30mL

SM/5 agar (2 g glucose (Fisher Scientific), 2 g Bacto Peptone (Oxoid), 2 g yeast extract

(Oxoid), 0.2 g $MgSO_4 * 7H_2O$ (Fisher Scientific), 1.9 g KH_2PO_4 (Sigma-Aldrich), 1 g

K_2HPO_4 (Fisher Scientific), and 15 g agar (Fisher Scientific) per liter) with 200 μL of

1.5 OD_{600} fluorescently labeled *K. pneumoniae* suspended in KK2 buffer (2.25 g

KH_2PO_4 (Sigma-Aldrich) and 0.67 g K_2HPO_4 (Fisher Scientific) per liter) spread over

the entirety of the agar. *K. pneumoniae* used in this study expressed green-fluorescent

protein (GFP) and were provided by dictyBase (Fey et al. 2019).

Dispersal experiment setup

To determine the consequences of hosts leaving behind bacteria, we performed a

dispersal experiment to seed new plates with sorus contents. To do this, we transferred

200 µL of $2x10^5$ spores/mL suspension obtained from the equilibration step to plates with

or without 200 µL of 1.5 OD_{600} GFP-expressing *K. pneumonia*e. Spore counts to make

spore suspensions were obtained using a hemocytometer. OD_{600} for plating bacteria was

measured with a BioPhotometer (Eppendorf, New York). We let bacteria and amoebae on

plates proliferate at room temperature for six days, enough time for amoebae to form

fruiting bodies.

We replicated these dispersal experiments with cured-and-reinfected hosts along

with uninfected controls on two separate dates. To understand if our results held with

natural infections, we performed an additional dispersal experiment with naturally

infected clones and uninfected controls. To rule out that leaving uneaten prey bacteria

was a host trait, we also performed a separate set of dispersal experiments comparing

cured clones to their reinfected counterparts.

Measurement of bacteria and host spore production

After dispersed amoebae and bacteria were allowed to proliferate for six days (to

allow hosts to complete development into fruiting bodies), we measured the density of

uneaten prey bacteria, *Paraburkholderia*, and host spore production. To measure the

density of uneaten prey bacteria, we first collected host spores and bacteria from after the

dispersal step. We collected host spores and bacteria by washing plates with 15 mL of

KK2 buffer after six days of growth. We counted host spores from this washed solution

using a hemocytometer. To measure bacterial density, we first removed host spores by

centrifuging the wash solution for three minutes at 360 rcf and reserving the supernatant.

We measured the optical density of the supernatant at 600 nm (OD_{600}) as well as

fluorescence with an excitation wavelength of 485 and emission wavelength of 515 nm in

a 96 well plate with a Tecan Infinite 200 Pro microplate reader. Removing host spores by

manually removing sori with a pipette tip resulted in similar densities of uneaten prey

bacteria consistent earlier findings (Khojandi *et al.*, 2019) that the number of prey

bacteria inside spores and sori is minimal relative to that left on the plate (data not

shown).

Since the supernatant from infected clones contains GFP-expressing *K.*

pneumoniae and unlabeled *Paraburkholderia*, the total OD_{600} is due to both kinds of

bacteria but fluorescence comes only from *K. pneumoniae*. To calculate the amount of

OD_{600} due to fluorescing *K. pneumoniae*, we generated a standard curve relating

fluorescence to OD_{600} using serial dilutions of GFP-expressing *K. pneumoniae* in KK2

and used this curve to predict the OD_{600} of *K. pneumoniae* in our samples. The remaining

OD_{600} is the amount due to *Paraburkholderia* symbionts, after also subtracting off the

background OD_{600} of the KK2 buffer. An OD_{600} of 0.1 translates to around 5×10^7 *K.*

pneumoniae cells or 1 x 10^8 *Paraburkholderia* cells according to the validation dataset in

Scott et al. (2022).

Development assays for cured and reinfected hosts

207   To determine how symbionts and their interactions with prey affected host

development time, we took time-lapse images of cured hosts and reinfected counterparts

developing in six-well plates filled with 10 mL SM/5. Each 6-well plate contained three

wells with cured hosts and three wells with reinfected hosts of either *P. hayleyella* clones

(QS23, QS38, and QS45) or *P. agricolaris* clones (NC21, QS159, and QS161). We

repeated development experiments on six-well plates three times for each

Paraburkholderia species. We first grew clones from the freezer on SM/5 plates for six

days in case there were freezer effects, as above. We then collected host spores and

plated 30 µL of $2x10^5$ spores per mL and 30 µL of 1.5 OD_{600} *K. pneumoniae* in each

well. Photos were taken every hour until fruiting bodies developed using a Canon EOS

Mark IV. We inspected photos to determine time points for when aggregates, slugs, and

fruiting bodies first appeared in each well.

Measurement of carried bacteria

221   To determine how prey context affected the number of prey bacteria carried in

sori, we used cured-and-reinfected hosts that were grown on prey-poor and prey-rich

plates after the dispersal step. We haphazardly sampled a single sorus from each of our

experimental plates from reinfection experiments (replicated on two different dates). To

count the number of carried prey, we suspended single-sorus contents in KK2 buffer and

plated out serial dilutions from no dilution to 1:1000 dilutions. Since *K. pneumoniae* were

labeled with GFP, we could differentiate colonies of *K. pneumoniae* and

*Paraburkholderi*a. We counted colonies from the dilution plate that appeared closest to

having 100 colonies and then back-calculated to get the total count for a single sorus.

This experiment was completely replicated on two separate dates.

Statistical Methods

To understand how *Paraburkholderia* infection affected the density of *K.*

pneumoniae bacteria left uneaten, we fit linear models in R (version 3.6.3). Because

experiments with cured-and-reinfected (and uninfected control) hosts were performed on

two different dates, we used linear mixed models (LMM) with date as a random effect.

We fit LMMs using the nlme4 package (Pinheiro & Bates, 2006) in R. For other

comparisons that involved only single measures, we used linear models fit by ordinary

linear regression (LM) with the lm function in R. We performed Tukey post-hoc tests for

pairwise comparisons using the *emmeans* package (Lenth *et al.*, 2019).

To determine whether the timing of development was affected by

Paraburkholderia infection, we used generalized linear mixed models (GLMMs) with

Poisson errors. We included the date that experiments were performed as a random

effect to capture variation within plates.

To test how prudent predation affects host spore production relative to other

explanations, we again used LMMs and LMs for cured-and-reinfected and naturally

infected infections, respectively, along with respective uninfected controls. If prudent

predation reduces host spore production, we expected a decrease in spore production with

increasing uneaten prey bacteria. Differences in host fitness could also be affected by the

density of *Paraburkholderia* left on plates (Scott *et al.*, 2022) or by infection category

(uninfected vs infected) (Brock *et al.*, 2011; DiSalvo *et al.*, 2015; Haselkorn *et al.*, 2019).

To account for these possibilities, we also fit models with these variables along with a

model that includes both uneaten prey density and *Paraburkholderia* density. In total, we

compared five models of spore production: (1) uneaten prey density (Prey model), (2)

Paraburkholderia density (Para model), (3) uneaten prey density + *Paraburkholderia*

density (Prey + Para model), (4) categorical infection status (Infection model), and (5) a

null model fit with only the intercept for each *Paraburkholderia* species.

To compare effects more easily across the different models, we scaled all

variables by subtracting the mean and dividing by twice the standard deviation. This

scaling procedure is useful for comparing models with multiple continuous predictors

(density of prey bacteria and symbionts) and with categorical factors (infected and

uninfected) because all measures end up on roughly the same scale (Gelman *et al.*, 2020).

To compare the fit of the five models described above, we calculated AICc for

each model. AICc is a measure of model fit that accounts for model complexity and that

corrects for small sample sizes (Burnham & Anderson, 2004). Models that fit the data

best have lower AICc values. For our model selection analysis we show results with 85%

confidence intervals instead of the standard 95% confidence intervals because 85%

confidence intervals are more consistent with model selection with AIC (Arnold, 2010).

To measure potential collinearity in our models that included prey and *Paraburkholderia*

density, we calculated variance inflation factors (VIF) using the check_collinearity

function in the performance package (Lüdecke *et al.*, 2021). VIFs lower than 5 indicate

low collinearity, values between 5 and 10 indicate moderate collinearity, and values

above 10 indicate high collinearity.

To compare the ability of hosts to carry *K. pneumoniae* inside fruiting bodies in

prey-rich and prey-poor environments, we used a zero-inflated negative binomial model

(ZINB) fit with the glmTMB package (Brooks *et al.*, 2017) in R. This zero-inflated

model incorporated two processes: (1) zero-inflation for whether fruiting bodies

contained or did not contain prey bacteria and (2) negative binomial for the counts of

carried prey bacteria. The zero-inflation parameter captures whether hosts are more likely

to carry while the negative binomial parameter captures whether hosts carry more prey

bacteria when they carry. Using this ZINB, we compared different hypotheses about how

prey-rich and prey-poor environments affected the ability to carry and the number carried

with AICc. We fit models to carriage data from both species and asked whether models

including prey environment, *Paraburkholderia* species, or the interaction between prey

environment and species had low AICc values. We did this model selection for both the

zero-inflation part of the model and for the negative binomial part of the model. We also

compared models with the random effects of clone and date included in either the zero-

inflation or negative binomial parts of the model.

Results:

D. discoideum hosts infected with Paraburkholderia leave some prey bacteria uneaten

We investigated the prey bacteria left behind after hosts formed fruiting bodies by

measuring the density of leftover *K. pneumoniae* from cured-and-reinfected hosts

compared to naturally uninfected controls. These prey densities are estimated from

fluorescence measurements since the *K. pneumoniae* used in this study expresses green

fluorescent protein (GFP; see methods). First, we wanted to confirm that some of the

bacteria hosts leave behind are prey bacteria and not just *Paraburkholderia* symbionts.

Leaving prey bacteria uneaten so that hosts miss out on potential growth and proliferation

was a key component of the prudent predation hypothesis, but previous studies did not

differentiate what kind of bacteria were left behind (Brock et al. 2011, 2016*b*).

Cured-and-reinfected hosts with *P. agricolaris* left 4.4 times as much prey

bacteria as naturally uninfected hosts (LMM; estimate = 0.009, se = 0.003, df = 20, P <

0.005; Figure 1A). Cured-and-reinfected hosts that were infected with *P. hayleyella* left

7.4 times as much prey bacteria than naturally uninfected hosts (Figure 1A; LMM;

estimate = 0.017, se = 0.003, df = 20, P < 0.001).

This comparison supports the view that symbionts cause prey bacteria to be left

uneaten. To control for host clone genotypes, we next compared cured-and-reinfected

hosts to the same host genotypes that were cured of their native *Paraburkholderia*

symbionts but not reinfected. Cured-and-reinfected hosts left more prey bacteria than

their cured counterparts (*P. agricolaris*: LM; estimate = 0.004, se = 0.002, df = 6, P =

0.041, *P. hayleyella*: LM; estimate = 0.004, se = 0.001, df = 6, P = 0.019; Figure 1B).

Host development time is not affected by Paraburkholderia infection

We hypothesized that leaving prey bacteria uneaten would result in faster

starvation and faster development by infected hosts. To test this, we took time-lapse

photos of amoebae aggregating into fruiting bodies, comparing cured-and-reinfected

hosts to the same host genotypes that were cured of their *Paraburkholderia* symbionts.

We determined time courses for when hosts aggregated, formed slugs, and formed

fruiting bodies. We were unable to detect that development times were slower for cured-

and-reinfected hosts relative to cured hosts for *P. agricolaris* (GLMM; estimate across all

developmental timepoints = 0.015, se = 0.037, P =   0.682) or *P. hayleyella* (GLMM;

estimate across all developmental timepoints = 0.044, se = 0.037, P = 0.231; Figure

2A&B). Individual developmental stages also did not differ due to *Paraburkholderia*

infection (P > 0.05).

Prey bacteria left on plates are not associated with reduced spore production in hosts

Leaving more prey uneaten is suspected to lower host fitness and could therefore

explain the cost of *Paraburkholderia* infection relative to uninfected hosts (Brock *et al.*,

2011). Alternately, the cost of infection may be more directly due to *Paraburkholderia*,

either infection itself or the density of *Paraburkholderia* symbionts. In addition to these

first three models (food bacteria density, *Paraburkholderia* infection status, and

Paraburkholderia density), we also fit a model that included both uneaten prey bacteria

density and *Paraburkholderia* density. To compare the relative support for these different

hypotheses, we fit a model for each hypothesis (see Methods) and compared fitted

models using AICc.

We first tested the role of uneaten prey bacteria and *Paraburkholderia* left on

plates on host spore production using our cured-and-reinfected hosts and naturally

uninfected controls (Table 1). For infections with either species of *Paraburkholderia*,

null models fit to only the intercept fit the data best (Figure 3A & S1; the model that

includes uneaten prey bacteria and *Paraburkholderia* density for *P. agricolaris* cured-

and-reinfected hosts — Prey + Para — did show confidence intervals that did not overlap

zero, but this was the worst model in terms of AICc). We thus found no evidence that

leaving prey explained the cost of infection for cured-and-reinfected hosts relative to

uninfected controls.

Naturally infected hosts and their uninfected controls also showed no support for

a cost due to leaving prey bacteria behind. Models of host spore production that included

uneaten prey (Prey model in Figure 3B) were poor predictors of host spore production.

This poor performance of prey as a predictor was also true when *Paraburkholderia* was

included as a covariate (Prey + Para model). This lack of fit for the *P. hayleyella* Para +

Prey model may be due to correlations between *Paraburkholderia* density and uneaten

food bacteria (VIF = 7.62).

These results show no support for a cost due to leaving prey bacteria behind (even

when accounting for *Paraburkholderia* density). We next turned to the direct role of

Paraburkholderia. Because models fit to only the intercept were the best for cured-and-

reinfected hosts relative to their controls, there was no evidence that *Paraburkholderia*

was costly in these clones, the same result as previously shown (Scott *et al.*, 2022). In

contrast, naturally infected hosts were afflicted by a cost of *Paraburkholderia* infection

(Figure 3B & S2). The best models for both *P. agricolaris* and *P. hayleyella* showed that

infected hosts had lower spore production than uninfected hosts (*P. agricolaris*: LM;

estimate = -0.508, se = 0.247, df = 11; *P. hayleyella*: LM; estimate = -0.804, se = 0.160,

df = 11). *Paraburkholderia* density predicted host spore production less well than

infection status. However, models that included *Paraburkholderia* density on its own

were better than models with prey bacteria density (Figure 3B & S2). These models thus

show little evidence for a cost of uneaten food bacteria and instead show that

Paraburkholderia infection is responsible for reduced host spore production in naturally

infected hosts.

Uneaten prey bacteria are a minority of left-behind bacteria

Prey bacteria left behind did not have the predicted effects on either development

time or spore production costs. One reason may be that the amount of left behind prey is

not as great as formerly thought (Brock *et al.*, 2011). Hosts may leave too few prey

bacteria to noticeably affect spore production. In fact, uneaten prey bacteria make up a

minority of left-behind bacteria. Uneaten prey bacteria are about 9% and 14% of bacteria

on plates for *P. agricolaris* and *P. hayleyella* cured-and-reinfected hosts, respectively,

and 21% and 19% for *P. agricolaris* and *P. hayleyella* naturally infected hosts,

respectively (Figure 4).

More prey bacteria are carried after hosts develop in prey poor conditions

To determine whether the number of prey bacteria carried in sori simply reflects

the number of prey bacteria in the previous environment or is modified according to the

interests of hosts or symbionts, we measured the number of fluorescent prey bacteria

inside sori after growth of cured-and-reinfected hosts on prey-rich and prey-poor plates

(Figure 5A). We found that for both *Paraburkholderia* species (Figure 5B&C), sori

produced in prey-rich environments were less likely to contain prey bacteria than those in

prey-poor environments (ZINB , zero-inflation estimate = 1.950, se = 0.987). This shows

that the ability to carry prey bacteria is affected by prey context, but in the opposite

direction to that expected if prey was carried according to the density of prey bacteria in

the environment. We did not detect a difference in the number of prey bacteria that were

caried between prey-rich and prey-poor environments (Figure 5D&E). Thus, the ability to

carry prey bacteria may depend on the fitness interests of hosts and symbionts that benefit

from carrying prey bacteria when harsh conditions are expected.

Discussion

394   Third parties that interact with symbiotic partners commonly affect the outcome

of symbiotic interactions (Palmer *et al.*, 2008; Chamberlain *et al.*, 2014; Hafer-Hahmann

& Vorburger, 2020; Cassidy *et al.*, 2022) and alter selection for cooperation and conflict

between symbiotic partners (Keeling & McCutcheon, 2017; Wood *et al.*, 2018). Often the

details of how third parties affect fitness effects are unknown. In the symbiosis between

D. discoideum social amoebae and *Paraburkholderia* bacteria, the third-party prey

bacteria are normally eaten by the host but can also be carried or left behind. We studied

how many prey bacteria are carried and left behind, and their impacts, by tracking

fluorescently labeled prey bacteria (*K. pneumoniae*) during fruiting body formation by

hosts.

404   Our most surprising finding was that infected hosts were more likely to carry prey

bacteria after growing in prey-poor environments (Figure 5). This result is surprising

because it means that hosts or symbionts actively change the number of carried prey

bacteria depending on environmental conditions. This result provides possible evidence

for mutual benefit between *D. discoideum* hosts and *Paraburkholderia* symbionts. Hosts

and symbionts are likely to benefit from modifying carriage in this way if future soil

environments tend to resemble past soil environments. Repeated prey-poor environments

are where hosts should most benefit from carrying and seeding out prey bacteria.

Paraburkholderia symbionts could also benefit from carriage of prey because keeping

hosts alive means there are hosts to further disperse *Paraburkholderia*. However, this

dispersal benefit to the symbionts must be balanced against the costs of increased

competition with the prey bacteria.

We speculate that carrying more prey in prey-poor contexts may represent

cooperation between hosts and symbionts that allows the symbiosis to persist over

repeated harsh environments. Such harsh conditions are potentially an important force

shaping cooperation in this (Scott *et al.*, 2022, 2023b) and other symbioses (Henry *et al.*,

2021; Veresoglou *et al.*, 2021; Cornwallis *et al.*, 2023). These studies in symbiotic

systems complement research on the role of harsh environments in the evolution of

cooperation that has been focused on interactions between members of the same species

(Kennedy *et al.*, 2018), most commonly in cooperatively breeding birds (Cornwallis *et*

al., 2017; Griesser *et al.*, 2017; Capilla-Lasheras *et al.*, 2021).

Although seeding out prey bacteria has been shown to be important when we move

spores to a prey-poor environment in the lab, we have little idea how frequently this

occurs in nature. The finding that carriage of prey bacteria changes in an apparently

adaptive fashion suggests that it is indeed important in the field. The importance of third-

parties in this case may be an interesting exception to the relatively small effects from

third-parties that have been observed across other studies (Chamberlain *et al.*, 2014).

Questions remain about how hosts and symbionts affect the carriage of prey

bacteria. *Paraburkholderia* are more often carried inside spores while prey bacteria are

more often carried outside spores (Khojandi *et al.*, 2019). Differences in carriage of

bacteria could potentially result from differences in lectins (Dinh *et al.*, 2018) or

polyphosphates (Rijal *et al.*, 2020) that affect the ability of bacteria to invade and survive

inside hosts. Hosts in the absence of their *Paraburkholderia* symbionts are able to modify

the contents of fruiting bodies through their immune cells that protect against toxins and

bacteria by collecting potential threats and dropping off during the slug stage (Chen *et al.*,

2007). However, the role of these immune cells in the symbiosis (Brock *et al.*, 2016a;

Scott *et al.*, 2023a) needs to be further explored. Manipulation of phagosomes could also

play a role in determining the contents of fruiting bodies. *Paraburkholderia* symbionts

increase the pH of phagosomes, presumably to prevent host digestion of symbionts (Tian

et al., 2022). Similar modification of lysosomes is used by pathogens to evade human

immune cells during infection (Isberg *et al.*, 2009; Leseigneur *et al.*, 2020). More work is

needed to understand how symbionts and food bacteria get into fruiting bodies and how

both hosts and symbionts contribute to bacterial carriage.

We also found that *Paraburkholderia* infection prevents hosts from eating all the

prey bacteria in an environment (Figure 1). A prior study suggested that leaving prey

bacteria uneaten because of prudent predation may explain the cost of infection relative

to uninfected hosts (Brock *et al.*, 2011). Instead, we found that the quantity of left-behind

prey bacteria may be too small to noticeably affect host spore production since we

observed that only a minority of left-behind bacteria was food bacteria, with the majority

being indigestible *Paraburkholderia* (Figure 4).

To get a better sense of how important left-behind prey bacteria could be for

hosts, we can use the density vs bacteria count curve in Figure S1B of Scott et al. 2022

and the average left-behind prey bacteria in this study to estimate the number of

individual left-behind prey bacteria. Using this estimation, *P. agricolaris* infected hosts

left 7.060×10^6 individual prey bacteria and *P. hayleyella* infected hosts left 1.206×10^7

individual prey bacteria. Using Kessin's (2001, p. 21) rough estimate that an amoebae

needs to eat 1,000 bacteria to divide, we calculated that the number of uneaten prey

bacteria is only enough to produce 0.002% and 0.004% more spores than what we

recovered from plates (collected spores were around 3×10^8) for *P. agricolaris* or *P.*

hayleyella infected hosts, respectively. These percentages are a rough and conservative

approximation as they assume that amoebae devote all consumed prey bacteria to

proliferation. However, the calculations show that it is unlikely that uneaten prey bacteria

can affect host fitness in a detectable manner.

Instead of hosts paying a cost in potential growth because they leave prey bacteria

uneaten, we suspect that *Paraburkholderia* infection causes both observations: that hosts

leave prey uneaten and that infected hosts pay a cost. Tentative support for this idea

comes from our findings that *Paraburkholderia* and uneaten prey bacteria densities were

correlated. This correlation could result from higher *Paraburkholderia* densities

interfering with the ability of hosts to eat prey bacteria.

An interesting remaining question in the *D. discoideum-Paraburkholderia*

symbiosis and in other symbioses is the mechanisms that result in reduced fitness when

hosts are infected. One potential explanation is that *Paraburkholderia* symbionts directly

feed on host cells or otherwise extract nutrients from hosts. Measures of

Paraburkholderia density inside sori have so far not been found to be correlated with

host fitness within species (Miller *et al.*, 2020), though the more toxic *P. hayleyella* does

appear to infect a higher percentage of spores than less toxic *P. agricolaris* (Shu *et al.*,

2018a; Khojandi *et al.*, 2019). Another promising hypothesis that deserves further study

is that hosts have lower spore production because of "collateral damage" from

competitive interactions between *Paraburkholderia* and prey bacteria (Scott *et al.*, 2022).

Competition between bacteria is often mediated by chemical warfare (Granato *et al.*,

2019) that could reduce host *D. discoideum* fitness as a side-effect.

The role of *Paraburkholderia* in reducing host spore production is more evidence

in support of there being some evolutionary conflict in this symbiosis (DiSalvo *et al.*,

2015; Scott *et al.*, 2022). However, the magnitude of conflict may differ across

Paraburkholderia symbiont species. This conflict may be more pronounced between *P.*

hayleyella and its hosts than between *P. agricolaris* and its hosts. We found that *P.*

hayleyella density could explain some of the decrease in naturally infected host spore

production but we did not find the same for *P. agricolaris* density (Figure 3). This

difference between *P. agricolaris* and *P. hayleyella* is consistent with prior studies that

found that *P. hayleyella* is more harmful than *P. agricolaris* (Shu *et al.*, 2018a;

Haselkorn *et al.*, 2019; Khojandi *et al.*, 2019). In addition to being more harmful, *P.*

hayleyella also has a reduced genome relative to *P. agricolaris* (Brock *et al.*, 2020; Noh

et al., 2022). Reduced genomes are a common result of persistent host association in

beneficial symbionts and pathogens (McCutcheon & Moran, 2012). While *P. hayleyella*

has maintained its ability to harm hosts over this persistent association, it is important to

remember that infection also comes with the ability to carry prey bacteria (DiSalvo *et al.*,

2015; Khojandi *et al.*, 2019). Thus, this symbiosis involves a balance between harm and

benefits to hosts.

Whether a specific symbiosis involves fitness alignment or conflict may depend

on a third party that affects the costs and benefits of symbiosis. Our results show that

third parties can have complex effects on conflict and fitness alignment in symbioses; the

symbiosis between *D. discoideum* and *Paraburkholderia* appears to involve elements of

conflict and cooperation that depends on how third-party bacteria are eaten, carried, and

left behind.

References

Arnold, T.W. 2010. Uninformative Parameters and Model Selection Using Akaike's
Information Criterion. *J. Wildl. Manag.* **74**: 1175–1178.

Brock, D.A., Callison, W.É., Strassmann, J.E. & Queller, D.C. 2016a. Sentinel cells,
symbiotic bacteria and toxin resistance in the social amoeba *Dictyostelium*
discoideum. *Proc. R. Soc. B Biol. Sci.* **283**: 20152727.

Brock, D.A., Douglas, T.E., Queller, D.C. & Strassmann, J.E. 2011. Primitive agriculture
in a social amoeba. *Nature* **469**: 393–396.

Brock, D.A., Haselkorn, T.S., Garcia, J.R., Bashir, U., Douglas, T.E., Galloway, J., *et al.*
2018. Diversity of Free-Living Environmental Bacteria and Their Interactions
With a Bactivorous Amoeba. *Front. Cell. Infect. Microbiol.* **8**: 411.

Brock, D.A., Jones, K., Queller, D.C. & Strassmann, J.E. 2016b. Which phenotypic traits
of Dictyostelium discoideum farmers are conferred by their bacterial
symbionts? *Symbiosis* **68**: 39–48.

Brock, D.A., Noh, S., Hubert, A.N.M., Haselkorn, T.S., DiSalvo, S., Suess, M.K., *et al.*
2020. Endosymbiotic adaptations in three new bacterial species associated
with *Dictyostelium discoideum* : *Paraburkholderia agricolaris* sp. nov.,
Paraburkholderia hayleyella sp. nov., and *Paraburkholderia bonniea* sp. nov.
PeerJ **8**: e9151.

Bronstein, J.L. 1994. Conditional outcomes in mutualistic interactions. *Trends Ecol.*
Evol. **9**: 214–217.

Brooks, M., E., Kristensen, K., Benthem, K., J. ,van, Magnusson, A., Berg, C., W., Nielsen,
A., *et al.* 2017. glmmTMB Balances Speed and Flexibility Among Packages for
Zero-inflated Generalized Linear Mixed Modeling. *R J.* **9**: 378.

Burnham, K.P. & Anderson, D.R. 2004. Multimodel Inference: Understanding AIC and
BIC in Model Selection. *Sociol. Methods Res.* **33**: 261–304.

Capilla-Lasheras, P., Harrison, X., Wood, E.M., Wilson, A.J. & Young, A.J. 2021.
Altruistic bet-hedging and the evolution of cooperation in a Kalahari bird. *Sci.*
Adv. **7**: eabe8980.

Cassidy, S.T., Markalanda, S., McFadden, C.J. & Wood, C.W. 2022. Herbivory modifies
plant symbiont number and impact on host plant performance in the field.
Evolution evo.14641.

Chamberlain, S.A., Bronstein, J.L. & Rudgers, J.A. 2014. How context dependent are
species interactions? *Ecol. Lett.* **17**: 881–890.

Chen, G., Zhuchenko, O. & Kuspa, A. 2007. Immune-like Phagocyte Activity in the
Social Amoeba. *Science* **317**: 678–681.

Cornwallis, C.K., Botero, C.A., Rubenstein, D.R., Downing, P.A., West, S.A. & Griffin,
A.S. 2017. Cooperation facilitates the colonization of harsh environments.
Nat. Ecol. Evol. **1**: 0057.

Cornwallis, C.K., Van 'T Padje, A., Ellers, J., Klein, M., Jackson, R., Kiers, E.T., *et al.*
2023. Symbioses shape feeding niches and diversification across insects. *Nat.*
Ecol. Evol., doi: 10.1038/s41559-023-02058-0.

Dinh, C., Farinholt, T., Hirose, S., Zhuchenko, O. & Kuspa, A. 2018. Lectins modulate
the microbiota of social amoebae. *Science* **361**: 402–406.

DiSalvo, S., Haselkorn, T.S., Bashir, U., Jimenez, D., Brock, D.A., Queller, D.C., *et al.*
2015. *Burkholderia* bacteria infectiously induce the proto-farming symbiosis
of *Dictyostelium* amoebae and food bacteria. *Proc. Natl. Acad. Sci.* **112**:
E5029–E5037.

Garcia, J.R., Larsen, T.J., Queller, D.C. & Strassmann, J.E. 2019. Fitness costs and
benefits vary for two facultative *Burkholderia* symbionts of the social
amoeba, *Dictyostelium discoideum*. *Ecol. Evol.* **9**: 9878–9890.

Gelman, A., Hill, J. & Vehtari, A. 2020. *Regression and other stories*. Cambridge
University Press.

Granato, E.T., Meiller-Legrand, T.A. & Foster, K.R. 2019. The Evolution and Ecology
of Bacterial Warfare. *Curr. Biol.* **29**: R521–R537.

Griesser, M., Drobniak, S.M., Nakagawa, S. & Botero, C.A. 2017. Family living sets the
stage for cooperative breeding and ecological resilience in birds. *PLOS Biol.*
15: e2000483.

Hafer-Hahmann, N. & Vorburger, C. 2020. Parasitoids as drivers of symbiont
diversity in an insect host. *Ecol. Lett.* **23**: 1232–1241.

Haselkorn, T.S., DiSalvo, S., Miller, J.W., Bashir, U., Brock, D.A., Queller, D.C., *et al.*
2019. The specificity of *Burkholderia* symbionts in the social amoeba farming
symbiosis: Prevalence, species, genetic and phenotypic diversity. *Mol. Ecol.*
28: 847–862.

Henry, L.P., Bruijning, M., Forsberg, S.K.G. & Ayroles, J.F. 2021. The microbiome
extends host evolutionary potential. *Nat. Commun.* **12**: 5141.

Horas, E.L., Metzger, S.M., Platzer, B., Kelly, J.B. & Becks, L. 2022. Context-dependent
costs and benefits of endosymbiotic interactions in a ciliate–algae system.
Environ. Microbiol. 1462-2920.16112.

Isberg, R.R., O'Connor, T.J. & Heidtman, M. 2009. The Legionella pneumophila
replication vacuole: making a cosy niche inside host cells. *Nat. Rev. Microbiol.*
7: 13–24.

Iwai, S., Fujita, K., Takanishi, Y. & Fukushi, K. 2019. Photosynthetic Endosymbionts
Benefit from Host's Phagotrophy, Including Predation on Potential
Competitors. *Curr. Biol.* **29**: 3114-3119.e3.

Jahan, I., Larsen, T., Strassmann, J.E. & Queller, D.C. 2021. Group maintenance in
aggregative multicellularity. In: *The Evolution of Multicellularity*, pp. 111–
134. CRC Press.

Keeling, P.J. & McCutcheon, J.P. 2017. Endosymbiosis: The feeling is not mutual. *J.*
Theor. Biol. **434**: 75–79.

Kennedy, P., Higginson, A.D., Radford, A.N. & Sumner, S. 2018. Altruism in a volatile
world. *Nature* **555**: 359–362.

Kessin, R.H. 2001. *Dictyostelium: evolution, cell biology, and the development of*
multicellularity. Cambridge University Press.

Khojandi, N., Haselkorn, T.S., Eschbach, M.N., Naser, R.A. & DiSalvo, S. 2019.
Intracellular Burkholderia Symbionts induce extracellular secondary
infections; driving diverse host outcomes that vary by genotype and
environment. *ISME J.* **13**: 2068–2081.

Lenth, R., Singmann, H., Love, J., Buerkner, P. & Herve, M. 2019. Package 'emmeans.'

Leseigneur, C., Lê-Bury, P., Pizarro-Cerdá, J. & Dussurget, O. 2020. Emerging Evasion
Mechanisms of Macrophage Defenses by Pathogenic Bacteria. *Front. Cell.*
Infect. Microbiol. **10**: 577559.

Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P. & Makowski, D. 2021.
performance: An R Package for Assessment, Comparison and Testing of
Statistical Models. *J. Open Source Softw.* **6**: 3139.

McCutcheon, J.P. & Moran, N.A. 2012. Extreme genome reduction in symbiotic
bacteria. *Nat. Rev. Microbiol.* **10**: 13–26.

Medina, J.M., Queller, D.C., Strassmann, J.E. & Garcia, J.R. 2023. The social amoeba
Dictyostelium discoideum rescues Paraburkholderia hayleyella, but not P.
agricolaris, from interspecific competition. *FEMS Microbiol. Ecol.* **99**: fiad055.
Oxford University Press.

Medina, J.M., Shreenidhi, P.M., Larsen, T.J., Queller, D.C. & Strassmann, J.E. 2019.
Cooperation and conflict in the social amoeba Dictyostelium discoideum. *Int.*
J. Dev. Biol. **63**: 371–382.

Miller, J.W., Bocke, C.R., Tresslar, A.R., Schniepp, E.M. & DiSalvo, S. 2020.
Paraburkholderia Symbionts Display Variable Infection Patterns That Are
Not Predictive of Amoeba Host Outcomes. *Genes* **11**: 674.

Noh, S., Capodanno, B.J., Xu, S., Hamilton, M.C., Strassmann, J.E. & Queller, D.C. 2022.
Reduced and Nonreduced Genomes in *Paraburkholderia* Symbionts of Social
Amoebas. *mSystems* e00562-22.

O'Brien, A.M., Jack, C.N., Friesen, M.L. & Frederickson, M.E. 2021. Whose trait is it
anyways? Coevolution of joint phenotypes and genetic architecture in
mutualisms. *Proc. R. Soc. B Biol. Sci.* **288**: 20202483.

Palmer, T.M., Stanton, M.L., Young, T.P., Goheen, J.R., Pringle, R.M. & Karban, R. 2008.
Breakdown of an Ant-Plant Mutualism Follows the Loss of Large Herbivores
from an African Savanna. *Science* **319**: 192–195.

Pinheiro, J. & Bates, D. 2006. *Mixed-effects models in S and S-PLUS*. Springer science &
business media.

Raper, K.B. 1937. Growth and development of Dictyostelium discoideum with
different bacterial associates.

Rijal, R., Cadena, L.A., Smith, M.R., Carr, J.F. & Gomer, R.H. 2020. Polyphosphate is an
extracellular signal that can facilitate bacterial survival in eukaryotic cells.
Proc. Natl. Acad. Sci. **117**: 31923–31934.

Scott, T.J., Larsen, T.J., Brock, D.A., Uhm, S.Y.S., Queller, D.C. & Strassmann, J.E. 2023a.
Symbiotic bacteria, immune-like sentinel cells, and the response to
pathogens in a social amoeba. *R. Soc. Open Sci.* **10**: 2023–05. Cold Spring
Harbor Laboratory.

Scott, T.J. & Queller, D.C. 2019. Long-term evolutionary conflict, Sisyphean arms
races, and power in Fisher's geometric model. *Ecol. Evol.* **9**: 11243–11253.

Scott, T.J., Queller, D.C. & Strassmann, J.E. 2022. Context dependence in the
symbiosis between Dictyostelium discoideum and Paraburkholderia. *Evol.*
Lett. **6**: 245–254.

Scott, T.J., Stephenson, C.J., Rao, S., Queller, D. & Strassmann, J.E. 2023b.
Unpredictable soil conditions affect the prevalence of a microbial symbiosis.
bioRxiv.

Shu, L., Brock, D.A., Geist, K.S., Miller, J.W., Queller, D.C., Strassmann, J.E., *et al.* 2018a.
Symbiont location, host fitness, and possible coadaptation in a symbiosis
between social amoebae and bacteria. *eLife* **7**: e42660.

Shu, L., Zhang, B., Queller, D.C. & Strassmann, J.E. 2018b. Burkholderia bacteria use
chemotaxis to find social amoeba Dictyostelium discoideum hosts. *ISME J.* **12**:
1977–1993.

Strassmann, J.E. & Queller, D.C. 2011. Evolution of cooperation and control of
cheating in a social microbe. *Proc. Natl. Acad. Sci.* **108**: 10855–10862.

Sun, B., Zhou, S. & Zhao, Q. 2003. Evaluation of spatial and temporal changes of soil
quality based on geostatistical analysis in the hill region of subtropical China.
Geoderma **115**: 85–99.

Tian, Y., Peng, T., He, Z., Wang, L., Zhang, X., He, Z., *et al.* 2022. Symbiont-Induced
Phagosome Changes Rather than Extracellular Discrimination Contribute to
the Formation of Social Amoeba Farming Symbiosis. *Microbiol. Spectr.*
e01727-21. Am Soc Microbiol.

Veresoglou, S.D., Johnson, D., Mola, M., Yang, G. & Rillig, M.C. 2021. Evolutionary bet-
hedging in arbuscular mycorrhiza-associating angiosperms. *New Phytol.*
nph.17852.

Vos, M., Wolf, A.B., Jennings, S.J. & Kowalchuk, G.A. 2013. Micro-scale determinants
of bacterial diversity in soil. *FEMS Microbiol. Rev.* **37**: 936–954.

Wendling, C.C., Piecyk, A., Refardt, D., Chibani, C., Hertel, R., Liesegang, H., *et al.* 2017.
Tripartite species interaction: eukaryotic hosts suffer more from phage
susceptible than from phage resistant bacteria. *BMC Evol. Biol.* **17**: 98.

Wood, C.W., Pilkington, B.L., Vaidya, P., Biel, C. & Stinchcombe, J.R. 2018. Genetic
669   conflict with a parasitic nematode disrupts the legume-rhizobia mutualism.
670   *Evol. Lett.* **2**: 233–245.

693 **Figure and Table Legends**

694 Table 1: Clones used in experiments.

Infection Type	Description	*Paraburkholderia* Treatment	Clones
Cured-and-reinfected	Clones that were cured of any native symbionts with tetracycline and reinfected in the lab with 0.1% of their native symbiont	*P. agricolaris* infected	QS159, QS161, QS606, NC21
		P. hayleyella infected	QS395, QS45, QS38, QS23
Naturally uninfected controls (for comparison with cured-and-reinfected)	Control clones that were treated with tetracycline.	Naturally uninfected control	QS6, QS138, QS472, QS527
Cured (for same-clone comparison	Clones that were cured of any native symbionts with	Cured of *P. agricolaris* infection	QS159, QS161, QS606, NC21

with cured-and-reinfected)	tetracycline, but not reinfected.	Cured of *P. hayleyella* infection	QS395, QS45, QS38, QS23
Naturally infected hosts	Clones with natural infections that have not been treated with antibiotics.	*P. agricolaris* infected	QS494, QS756, QS788, QS113, QS453, QS70, QS606
		P. hayleyella infected	QS45, QS101, QS46, QS2, QS23, QS529, QS38
Naturally uninfected controls (for comparison with naturally infected)	Clones without natural infections that have not been treated with antibiotics.	Naturally uninfected control	QS4, QS6, QS14, QS18, QS9, QS8

Figure 1: Symbionts cause hosts to leave prey uneaten. (A) Density left on plate of
uneaten *K. pneumoniae* prey bacteria (measured by OD_{600}) for naturally uninfected hosts,
P. agricolaris hosts that have been cured and then reinfected, and *P. hayleyella* hosts that
have been cured and then reinfected. (B) Density left on plate of uneaten prey bacteria for
cured *P. agricolaris* and *P. hayleyella* hosts compared to their cured-and-reinfected
counterparts. Dot and lines show mean and standard deviation, respectively. Asterisks
indicate significant differences (in panel A comparisons are between infected and
uninfected). Point shapes of small points indicate different dates on which experiments
were replicated.

Figure 2: Symbionts do not affect developmental time. Developmental time points at

different stages of development for cured hosts and their counterparts that have been

cured and then reinfected with (A) *P. agricolaris* and (B) *P. hayleyella* hosts. Dot and

lines show mean and standard deviation, respectively. Point shapes of small points

indicate different dates on which experiments were replicated.

Figure 3: Host spore production as a function of different bacteria densities on plates and

infection status in (A) cured-and-reinfected (B) and naturally infected hosts relative to

their respective uninfected counterparts (see Table 1). We compared models of host spore

production (shown in different colors), predicted by *Paraburkholderia* density only,

uneaten prey bacteria only, both *Paraburkholderia* and uneaten prey bacteria, or a

categorical variable for infection status (infected with *Paraburkholderia* or not).

Estimated effects are shown as points and 85% confidence intervals are shown as lines

(null models with only the intercept are not shown). ΔAICc values are shown in

increasing order from top to bottom and colored according to the kind of model.

Intercepts are not shown.

Figure 4: Most of the bacteria left behind by infected hosts were not prey bacteria.

Percent of the total left-behind bacteria (includes prey bacteria and *Paraburkholderia*

symbionts) that was prey bacteria from cured-and-reinfected (A) and naturally infected

(B) hosts. Dot and lines show mean and standard deviation, respectively. Point shapes for

small points indicate different dates on which experiments were replicated

Figure 5: Hosts are less likely to carry prey bacteria in sori after developing in prey-rich

environments. (A) Photo of fluorescent *K. pneumoniae* prey bacteria colonies plated from

an individual sorus. (B&C) Sori with carried food bacteria for cured-and-reinfected *P.*

agricolaris (B) or *P. hayleyella* (C) hosts from prey-poor and prey-rich contexts. Carriage

outcomes for individual fruiting bodies are shown as small points that indicate carried

prey (1) or did not carry prey (0). Large points and lines show the proportion of sori with

prey and the standard deviation. (D&E) Number of prey bacteria of sori that contained

prey (1s in B&C), for hosts carrying *P. agricolaris* (D) or *P. hayleyella* (E) from prey-

poor and prey-rich contexts. Large points and lines in D&E show the average and

standard deviation. Carried prey bacteria were quantified by plating serial dilutions (from

748   undiluted to 1:1000) of single fruiting bodies. Point shapes for individual fruiting bodies

749   (small points) indicate different dates on which experiments were replicated.