Third-party effects in the *Dictyostelium-Paraburkholderia* symbiosis: food bacteria that are eaten, carried, or left behind

Trey J. Scott*

David C. Queller

Joan E. Strassmann

Department of Biology, Washington University in St. Louis

*Corresponding Author; E-mail: tjscott@wustl.edu

Acknowledgements

This material is based upon work funded by the National Science Foundation under grants DEB-1753743 and IOS-1656756. We thank Tyler Larsen for comments on an early draft of the manuscript. We also thank Debbie Brock and the rest of the Strassmann/Queller lab group for feedback during the development and execution of this project.

Conflict of interest

We declare no conflicts of interest.

Keywords: Symbiosis, cooperation, conflict, Dictyostelium discoideum, Paraburkholderia

Abstract

Symbiotic interactions change depending on the abundance of third parties like

predators, prey, or pathogens. Third-party interactions with food bacteria are central to

the symbiosis between *Dictyostelium discoideum* social amoeba hosts and inedible

Paraburkholderia bacterial symbionts. Symbiosis with *Paraburkholderia* allows host *D.*

discoideum to carry food bacteria through the dispersal stage where host amoebae

aggregate and develop into fruiting bodies that disperse spores. Carrying bacteria benefits

hosts when food bacteria are scarce but harms hosts when food bacteria are plentiful. The

nature of this cost is unknown, but hosts leave bacteria behind when they carry

symbionts. If this left-behind bacteria includes uneaten food bacteria, infected hosts may

lose potential growth. Thus, decisions about how many food bacteria to eat, to carry,

and to leave behind are crucial for understanding both benefits and costs in this

symbiosis. We investigated how many food bacteria are uneaten and carried in this

symbiosis by measuring fluorescently labeled food bacteria after fruiting body

development. We found that *Paraburkholderia* infection makes hosts leave both

symbionts and uneaten food bacteria but leaving food bacteria uneaten did not explain

costs to hosts. Counts of food bacteria in fruiting bodies showed that hosts carry more

food bacteria after developing in food-poor environments than in food-rich. This

indicates that hosts, and possibly *Paraburkholderia* symbionts, actively modify how

many food bacteria are carried to ensure hosts have food in the harshest conditions.

Decisions about how many third-party bacteria to eat, carry, or leave may thus have

important effects on this symbiosis.

Introduction

The fitness effects of symbiotic interactions can change depending on the

environment (Bronstein, 1994; Chamberlain *et al.*, 2014; Horas *et al.*, 2022; Scott *et al.*,

2022). One crucial component of the environment can be a third species that interacts

with hosts and symbionts (Palmer *et al.*, 2008; Wendling *et al.*, 2017; Wood *et al.*, 2018;

Hafer-Hahmann & Vorburger, 2020; Cassidy *et al.*, 2022). For example, in the symbiosis

between ants and *Acacia* plants, ants benefit *Acacia* by fending off herbivores. However,

when herbivores were prevented from accessing *Acacias*, ordinarily mutualistic

interactions between ants and *Acacias* shifted towards antagonism because ants no longer

provided a benefit to host *Acacias* (Palmer *et al.*, 2008). Such shifts can influence

whether host and symbiont fitness interests are aligned or in conflict (Keeling &

McCutcheon, 2017; Iwai *et al.*, 2019). However, the details of how third parties affect

symbioses are not well understood for many kinds of symbioses (Chamberlain *et al.*,

2014).

The social amoeba *Dictyostelium discoideum* is an important model of microbial

symbiosis. *D. discoideum* is a soil amoeba that feeds on bacteria (Raper, 1937). Upon

starvation, amoebae aggregate to form a multicellular fruiting body that disperses spores

to new soil patches (Kessin, 2001). Development of the fruiting body progresses through

well-studied stages starting with aggregation of individual amoebae which then become

motile slugs (Bozzaro, 2019). Slugs can move to locations with food or find a location to

develop into a fruiting body. During fruiting body development, about 20% of the cells

die and become stalk while the remaining cells develop into spores that sit atop the stalk

in a structure called a sorus (Strassmann & Queller, 2011).

D. discoideum has symbioses with three species of facultatively intracellular

Paraburkholderia bacteria (Brock *et al.*, 2020). In natural isolates, around 25% of wild-

collected *D. discoideum* are infected by one, and occasionally two, of the three species of

symbionts (Haselkorn *et al.*, 2019). When not referring specifically to one of the three

species – *P. agricolaris*, *P. hayleyella*, or *P. bonniae* – we will refer to them collectively

in this paper as either *Paraburkholderia* or symbionts. These symbionts are specialized

for interacting with amoebae (Shu *et al.*, 2018a; b; Brock *et al.*, 2020). Symbionts move

towards hosts (Shu *et al.*, 2018b) and make host phagosomes less acidic so that the

symbionts are not digested (Tian *et al.*, 2022). One species of symbiont, *P. hayleyella,*

appears to have led to host adaptation in response to the presence of symbionts (Shu *et*

al., 2018a). *P. hayleyella* and *P. bonniae* also have reduced genomes relative to *P.*

agricolaris and most other *Paraburkholderia* (Brock *et al.*, 2020; Noh *et al.*, 2022),

which indicates that *P. hayleyella* and *P. bonniea* have been associated closely with their

hosts during a long evolutionary history (McCutcheon & Moran, 2012).

Fitness of *D. discoideum* hosts and *Paraburkholderia* symbionts is affected by

interactions with a third set of organisms, the various food bacteria in the environment

that are eaten by host amoebae (Brock *et al.*, 2011; DiSalvo *et al.*, 2015; Scott *et al.*,

2022). *Paraburkholderia* symbionts are largely inedible by hosts, and are carried

undigested inside the sorus along with additional edible bacteria (Brock *et al.*, 2011;

DiSalvo *et al.*, 2015; Khojandi *et al.*, 2019). When hosts disperse to food-poor

environments where food bacteria are rare, hosts benefit from *Paraburkholderia* infection

because hosts are able to carry food bacteria to seed new populations (Brock *et al.*, 2011;

DiSalvo *et al.*, 2015; Scott *et al.*, 2022). When infected hosts disperse to food-rich

environments, hosts pay a cost of producing fewer spores relative to uninfected *D.*

discoideum (Brock *et al.*, 2011; DiSalvo *et al.*, 2015). Presence of more

Paraburkholderia is also associated with lower host spore production, but this effect

depends on whether food bacteria are present (Scott *et al.*, 2022). Thus, host fitness

depends on the abundance of food bacteria in the environment prior to fruiting body

formation.

Fitness of inedible *Paraburkholderia* symbionts also changes depending on the

density of edible bacteria that are eaten by hosts, probably because they compete for

resources. We could thus refer to food-rich and food-poor host environments as being

high and low competition environments for *Paraburkholderia* symbionts. To be

consistent between host and symbiont environments, we will use food-rich and food-poor

but we note that this affects competition for symbionts. After symbionts are carried to

food-poor environments, both *P. agricolaris* and *P. hayleyella* reach higher densities

relative to food-rich environments where they face more competition (Scott *et al.*, 2022).

The effect of food bacteria on hosts and symbionts highlights how food bacteria

play an important role in this symbiosis. However, there are still many questions about

how food bacteria affect this symbiosis throughout the development of *D. discoideum*. Of

special interest is the period when fruiting bodies are formed and food bacteria are

transferred into the sorus for carriage. This carriage period is important because the

abundance of food bacteria impacts both parties (Brock *et al.*, 2011; DiSalvo *et al.*, 2015;

Scott *et al.*, 2022). It is important to note that carrying food bacteria may be a joint

phenotype (Queller, 2014; Queller & Strassmann, 2018) that both hosts and symbionts

affect together rather than being solely controlled by hosts. Joint phenotypes are

interesting to study in symbioses because of the potential for cooperation and conflict

over the value of the joint phenotype (Scott & Queller, 2019; Quides *et al.*, 2021). Here,

it is the disposition of food bacteria that is of interest – should they be eaten, carried, or

left behind?

Leaving food bacteria behind would not seem to be adaptive, but this appears to

happen when hosts are infected by *Paraburkholderia* but not when they are uninfected

(Brock *et al.*, 2011, 2016b). Initially, the key role of *Paraburkholderia* in causing hosts to

leave behind bacteria was not known. It was thought that the bacteria left behind were all

food bacteria (Brock *et al.*, 2011). This was taken as a possible sign of prudent predation

by host amoebae — if some bacteria were going to be saved for carriage, hosts cannot eat

all the available food and may stop feeding and start developing earlier (Brock *et al.*,

2011). It was hypothesized that hosts leaving food bacteria uneaten might explain why

carriage is costly in some environments (Brock *et al.*, 2011).

The essential role of *Paraburkholderia* in the symbiosis, including causing hosts

to leave behind bacteria is now known (DiSalvo *et al.*, 2015; Brock *et al.*, 2016b).

Moreover, at least some of the bacteria left behind are not food bacteria but

Paraburkholderia symbionts (Scott *et al.*, 2022). Because of this new information, the

prudent predation hypothesis that hosts are leaving behind bacteria in order to carry

should be re-evaluated. Several questions need to be clarified. What fraction of the

bacteria left behind are food bacteria – how much real food is left on the table? Does

Paraburkholderia infection cause hosts to leave the food bacteria uneaten? Are food

bacteria left because the hosts cease eating and start developing earlier? What costs are

there for leaving food bacteria uneaten, or are costs instead due to the number of

Paraburkholderia symbionts?

While infected hosts may leave some food bacteria uneaten, they also gain the

ability to carry food bacteria along with dispersing spores in sori (Brock *et al.*, 2011;

DiSalvo *et al.*, 2015; Khojandi *et al.*, 2019). It is unknown whether the number of carried

food bacteria changes in different environments. Changes in the number of carried food

bacteria could depend on the abundance of food bacteria in the environment or on the

fitness interests of *D. discoideum* hosts and *Paraburkholderia* symbionts.

The number of carried food bacteria could reflect a passive process that mirrors

the densities of food bacteria in the environment. In this case, hosts that can carry (those

that are infected with *Paraburkholderia*) would carry food bacteria proportional to the

density of food bacteria in the environment. If this is the case, we expect hosts to carry

more food after developing in a food-rich environment and fewer food bacteria after

developing in a food-poor environment.

Alternatively, the number of carried food bacteria could reflect the evolutionary

interests of hosts and symbionts. Since soil environments tend to be spatially and

temporally structured (Sun *et al.*, 2003; Vos *et al.*, 2013), developing in a food-poor

environment may be associated with an increased probability that future environments

will also tend to be food-poor. If this is the case, hosts may carry more food bacteria

after developing in a food-poor environment than in a food-rich environment because

hosts that seed the next food-poor environment with food bacteria will have an

advantage. From the *Paraburkholderia* symbiont's perspective, allowing more carriage

of food bacteria would keep their hosts alive and allow further dispersal of symbionts by

hosts. Altering carriage in response to environmental conditions could therefore be

adaptive for both hosts and symbionts.

We investigate three questions about the role of food bacteria in the symbiosis

between *D. discoideum* and two commonly studied *Paraburkholderia* symbionts, *P.*

agricolaris and *P. hayleyella* (Shu *et al.*, 2018a; Garcia *et al.*, 2019; Scott *et al.*, 2022).

We first re-evaluate some of the ideas behind the prudent predation hypothesis by

measuring the density of any food bacteria that are left uneaten. We then test the fitness

consequences of prudent predation on hosts and ask if hosts that leave more food bacteria

uneaten pay a cost in spore production. Lastly, we turn to whether the number of food

bacteria carried inside sori changes between food-poor and food-rich environments.

Methods

Clones and Culturing Methods

To compare between infected and uninfected hosts, we used 12 host *D.*

discoideum clones, 4 clones uninfected, 4 infected with *P. agricolaris*, and 4 infected

with *P. hayleyella*. We refer to these infected clones as "reinfected" (Table 1) because

they were cured with tetracycline and reinfected with 0.1% of the *Paraburkholderia*

strain with which they were isolated. This reinfection scheme ensures that initial infection

densities were the same across clones.

We also used a set of uncured clones that have not been treated with antibiotics to

modify their native infection levels. For these clones, we used 7 clones naturally infected

with *P. agricolaris*, 7 clones naturally infected with *P. hayleyella*, and 6 clones that were

not infected with any *Paraburkholderia* symbionts (Table 1). One naturally uninfected

clone (QS1) was contaminated so we excluded it from the analysis leaving 6 clones in

our last infection category. These uncured clones were useful for checking whether our

results applied to unmanipulated infection densities.

To remove any effects of being in the freezer and to ensure that infected amoebae

carried fluorescently labeled *Klebsiella pneumoniae* food bacteria, we grew amoebae

through one round of feeding and fruiting body formation and then collected spores to

initiate our experiments. This step may allow infection densities time to equilibrate

(Miller *et al.*, 2020). Amoebae were grown from frozen spores at room temperature on

SM/5 plates (2 g glucose (Fisher Scientific), 2 g Bacto Peptone (Oxoid), 2 g yeast extract

(Oxoid), 0.2 g $MgSO_4$ * $7H_2O$ (Fisher Scientific), 1.9 g KH_2PO_4 (Sigma-Aldrich), 1 g

K_2HPO_4 (Fisher Scientific), and 15 g agar (Fisher Scientific) per liter) with 200 µL of

1.5 OD_{600} fluorescently labeled *K. pneumoniae* suspending in KK2 buffer (2.25 g

KH_2PO_4 (Sigma-Aldrich) and 0.67 g K_2HPO_4 (Fisher Scientific) per liter). *K.*

pneumoniae used in this study expressed green-fluorescent protein (GFP) and were

provided by dictyBase (Fey et al. 2019). To mimic dispersal to food-poor and food-rich

environments, we transferred 200uL of $2x10^5$/mL spore solution to plates with or without

200uL of 1.5 OD_{600} GFP-expressing *K. pneumonia*e. We let bacteria and amoebae on

plates proliferate at room temperature for six days, enough time for amoebae to form

fruiting bodies.

Measurement of bacteria and host spore production

To measure the density of uneaten food bacteria after *D. discoideum* fruiting body

formation, we first collected host spores and bacteria from plates after six days of growth

by washing plates with 15 mL of KK2 buffer. We counted host spores from this washed

solution using a hemocytometer. To measure bacterial density, we first removed host

spores by centrifuging the wash solution for three minutes at 1300 rpm and collecting the

supernatant. Removing host spores by manually removing sori with a pipette tip resulted

in similar densities of uneaten food bacteria suggesting that the number of bacteria inside

spores and sori is minimal relative to that left on the plate. After removing host material

by centrifugation, we measured the optical density of the supernatant at 600 nm (OD_{600})

as well as fluorescence with an excitation wavelength of 485 and emission wavelength of

515 nm in a 96 well plate with a Tecan Infinite 200 Pro microplate reader.

Since the washed plate solution from infected clones contains GFP-expressing *K.*

pneumoniae and unlabeled *Paraburkholderia*, the total OD_{600} is due to both kinds of

bacteria but fluorescence only comes from the *K. pneumoniae*. To calculate the amount

of OD_{600} due to fluorescing *K. pneumoniae*, we generated a standard curve relating

fluorescence to OD_{600} using serial dilutions of GFP-expressing *K. pneumoniae* in KK2

and used this curve to predict the OD_{600} of *K. pneumoniae* in our samples. The remaining

OD_{600} is the amount due to *Paraburkholderia* symbionts after subtracting off the

background OD_{600} of the KK2 buffer. An OD_{600} of 0.1 translates to around 5×10^7 *K.*

pneumoniae cells and 1×10^8 *Paraburkholderia* cells according to the validation dataset

in Scott et al. (2022).

To measure how many *K. pneumoniae* bacteria were carried in sori, we

haphazardly sampled a single sorus from each of our experimental plates from re-

infection experiments (n = 24). We suspended single sorus contents in KK2 buffer and

plated out serial dilutions. Since *K. pneumoniae* are labeled with GFP, we could

differentiate colonies of *K. pneumoniae* and *Paraburkholderia* and get counts of colony

forming units (CFU). We then used these CFU counts to back-calculate the number of *K.*

pneumoniae bacteria inside fruiting bodies.

Development assays for cured and reinfected hosts

To determine how symbionts affected host development, we took time-lapse

images of cured and reinfected hosts growing in six well plates. We grew clones from the

freezer on SM/5 plates for six days in case there were freezer effects as above. We then

collected host spores and plated 30 uL of 2×10^5 spores per mL and 30 uL of 1.5 OD *K.*

pneumoniae in each well. Photos were taken every hour until fruiting bodies developed

using a Canon EOS Mark IV. We inspected photos to determine time points for when

aggregates, slugs, and fruiting bodies first appeared in each well. We included three

replicates for each species and performed each individual replicate on a separate date.

Calculating effect of uneaten food on host fitness

To estimate how much host proliferation was lost by leaving food bacteria

uneaten, we used Kessin's (2001, p. 21) estimate that an amoeba is roughly 1,000 times

larger than a bacteria and would thus require around 1,000 bacteria to divide. To estimate

the number of food bacteria cells left uneaten, we used the validation data (Figure S1) in

Scott et al. (2022) that predicted *K. pneumoniae* colony forming units from OD_{600}. The

number of additional amoebae that would result from eating the uneaten food bacteria is

then the estimated number of food bacteria divided by 1,000.

Statistical Methods

233   To compare the density of *K. pneumoniae* bacteria left uneaten on the plate for

different infection categories (uninfected vs *P. agricolaris* infected vs *P. hayleyella*

infected and cured vs infected), we fit linear models in R (version 3.6.3). For uncured

infection comparisons that involved single measures of different clones, we used linear

models fit by generalized least squares (GLS) in the *nlme* package (Pinheiro & Bates,

2006). For reinfection experiments, which were performed on two separate dates with the

same clones, we fit linear mixed models (LMM) and included clone identity or date as a

random effect depending on which random effect resulted in a better fit. To compare

cured and infected hosts, we used ordinary linear regression with the *lm* function in R.

We performed Tukey post-hoc tests for pairwise comparisons using the *emmeans*

package (Lenth *et al.*, 2019).

244   To determine whether the timing of development was affected by

245   *Paraburkholderia* infection, we used generalized linear mixed models (GLMMs) with a

246   Poisson link function. We included the date that experiments were performed as a

247   random effect to capture variation within plates.

248   To determine how leaving food bacteria uneaten and *Paraburkholderia* infection

affect host spore production, we again used LMM and GLS for reinfections and uncured

infections, respectively. If prudent predation results in a cost for hosts, we expected a

decrease in fitness with increasing uneaten food bacteria. To test this, we fit linear models

of host spore production for infections with each species. Uninfected hosts' spore

production was included in each model to act as a baseline fitness when hosts are not

affected by infection. Differences in host fitness could also be affected by the density of

Paraburkholderia left on plates (Scott *et al.*, 2022) or by infection category (uninfected

vs infected) (Brock *et al.*, 2011; DiSalvo *et al.*, 2015; Haselkorn *et al.*, 2019). To account

for these possibilities, we also fit models with these variables along with a model that

includes both uneaten food density and *Paraburkholderia* density. In total, we compared

five models of spore production: (1) uneaten food density (Food model), (2)

Paraburkholderia density (Para model), (3) uneaten food density + Paraburkholderia

density (Food + Para model), (4) categorical infection status (Infection model), and (5) a

null model fit with only the intercept. To more easily compare effects across the different

models, we scaled all variables by subtracting the mean and dividing by the standard

deviation. We selected the best models among these five using AICc, a measure of model

fit that corrects for small sample sizes (Burnham & Anderson, 2004). Models that fit the

data best have lower AICc values.

To compare the number of *K. pneumoniae* inside fruiting bodies in food-rich and

food-poor environments, we used generalized mixed models with a logistic link function

to test whether the food environment affected the probability of having food in a sorus.

We transformed our response variable, the number of *K. pneumoniae* inside a fruiting

body, to a presence absence variable for the logistic regression. We used a logistic

function in this case because of the number of zeros in our data, which could cause

problems for linear mixed models and for generalized mixed models with Poisson fits.

We fit our model to data from both species but determined that species identity was not a

useful predictor using AICc.

Results:

D. discoideum hosts infected with Paraburkholderia leave food bacteria uneaten

We investigated the food bacteria left behind after hosts formed fruiting bodies by

measuring the density of leftover *K. pneumoniae* from cured and reinfected hosts and

naturally uninfected controls (Table 1). These densities are estimated from fluorescence

measurements since the *K. pneumoniae* used in this study expresses green fluorescent

protein (GFP; see methods). First, we wanted to confirm that some of the bacteria hosts

leave behind are food bacteria and not just *Paraburkholderia* symbionts. Leaving food

bacteria uneaten so that hosts miss out on potential growth and proliferation was a key

component of the prudent predation hypothesis, but previous studies did not differentiate

what kind of bacteria were left behind (Brock et al. 2011, 2016*b*). Hosts that were cured

and reinfected with *P. agricolaris* and hosts that were cured and reinfected with *P.*

hayleyella leave more food bacteria on the plate (Figure 1A) than naturally uninfected *D.*

discoideum (Tukey Post-Hoc test (TPH), p ≤ 0.001 for both *Paraburkholderia* species).

This comparison supports the view that symbionts cause food bacteria to be left

uneaten but does not exclude the possibility that the differences are due to the different *D.*

discoideum clones that we used. Therefore, we next compared cured and reinfected hosts

to the same host genotypes that were cured of their native *Paraburkholderia* symbionts

but not reinfected. Both *P. agricolaris* (p = 0.0139) and *hayleyella* (p = 0.0162) re-

infections resulted in more food bacteria being left uneaten relative to cured but not

reinfected *P. agricolaris* and *hayleyella* hosts (Figure 1B).

Host development is not affected by Paraburkholderia infection

We hypothesized that leaving food bacteria uneaten would be associated with

faster development by infected hosts. To test this, we took time-lapse photos of amoebae

aggregating into fruiting bodies that either did or did not also carry *Paraburkholderia.*

We determined time courses for when they aggregated, formed slugs, and formed fruiting

bodies. We found that development times were not affected by *Paraburkholderia*

infection for either species (GLMMs; $p > 0.05$; Figure 2A&B).

Food bacteria left on plates is not associated with reduced spore production in hosts;

Paraburkholderia infection is

Leaving more food uneaten is suspected to lower host fitness and could therefore

explain the cost of *Paraburkholderia* infection relative to uninfected hosts (Brock *et al.*,

2011). Alternately, the cost of infection may be due to *Paraburkholderia* — this could be

measured as a categorical variable (infected vs uninfected) or as a continuous variable

(the density of *Paraburkholderia* symbionts on plates). Costs could also result from the

density of both uneaten food bacteria and *Paraburkholderia* symbionts.

We first tested the role of uneaten food bacteria and *Paraburkholderia* on host

spore production using our cured and reinfected hosts. For infections with both species of

Paraburkholderia, null models fit to only the intercept fit the data best (Figure 3A;

Supplemental File 1; the model that includes uneaten food bacteria and *Paraburkholderia*

density for *P. agricolaris* infected hosts — Food + Para — did show confidence intervals

that did not overlap zero, but this was the worst model in terms of AICc).

Our experiment with cured and reinfected hosts showed no effects of any of these

variables (Figure 3A) on host spore production (the null model without any variables was

the best fit). This includes no significant cost of infection, contrary to earlier results

(Brock *et al.*, 2011; DiSalvo *et al.*, 2015; Haselkorn *et al.*, 2019). The lack of cost may be

because hosts were infected with an extremely low density of *Paraburkholderia*

symbionts (Scott *et al.*, 2022). To test whether unmanipulated infection densities changed

our results, we performed a separate experiment with uncured hosts that have not been

treated with antibiotics (Table 1). Similar uncured clones have been shown to exhibit

costs of infection (Brock *et al.*, 2011).

Uncured hosts were afflicted by a cost of infection, but this cost was due to

Paraburkholderia infection instead of lost growth and proliferation from leaving uneaten

food bacteria. For uncured hosts with *P. agricolaris*, models of host spore production that

included uneaten food (Food model in Figure 3B), *Paraburkholderia* density (Para

model), and both variables together (Food + Para) were poor predictors of host spore

production (Supplemental File 1). The model with infection status was the best predictor

of host spore production for *P. agricolaris* uncured hosts (Supplemental File 1; note that

it is within 2 AICc units of the null model). These *P. agricolaris* results identify a cost of

infection like that found in prior studies (Brock *et al.*, 2011; DiSalvo *et al.*, 2015;

Haselkorn *et al.*, 2019).

For uncured hosts with *P. hayleyella*, we also found that *Paraburkholderia*

infection was responsible for costs. Uneaten food bacteria was associated with lower

spore production only if uneaten food was the only variable included in the model (Figure

3B). Including *Paraburkholderia* density changed the effect of uneaten food to slightly

positive with confidence intervals that overlap zero. *Paraburkholderia* density was a

better predictor of hosts spore production than uneaten food (Supplemental File 1) and

was highly correlated with uneaten food density (> 0.9). For uncured hosts infected with

P. hayleyella, infection status was again the best predictor (Supplemental File 1). Our

model comparisons for *P. hayleyella* also demonstrated a cost of infection. These models

of host spore production in uncured hosts point to a more direct role for

Paraburkholderia symbionts in reducing host spore production rather than a cost from

prudent predation.

Uneaten food bacteria are a minority of left-behind bacteria

Food bacteria left behind did not have the predicted effects on either development

time or spore production costs. One reason may be that the amount of left behind food is

not as great as formerly thought. In fact, uneaten food bacteria make up a minority of left-

behind bacteria, with the majority being *Paraburkholderia* (Figure 4). Using Kessin's

(2001, p. 21) rough estimate that an amoebae needs to eat 1,000 bacteria to divide, we

calculate that the number of uneaten food bacteria is only enough to produce 0.001-

0.004% more spores than what we recovered from plates. It is thus unlikely that uneaten

food bacteria can noticeably affect host fitness.

More food bacteria are carried after hosts develop in food poor conditions

To determine whether the number of food bacteria carried in sori simply reflects

the number of food bacteria in the previous environment or the interests of hosts and

symbionts, we measured the number of fluorescent food bacteria inside sori after growth

on food-rich and food-poor plates (Figure 5A). Counter-intuitively, we found that for

both *Paraburkholderia* species (Figure 5B&C), sori produced in food-rich environments

had a lower proportion of containing food bacteria (GLMM, Log odds = -1.894, se =

0.942, p = 0.044). This shows that the number of carried food bacteria is affected by food

context, but in the opposite direction to that expected if food was carried in proportion to

the density of food bacteria in the environment. Thus, the amount of carried food bacteria

may depend on the fitness interests of hosts and symbionts that benefit from carrying

more food bacteria when harsh conditions are expected.

Discussion

Third parties that interact with symbiotic partners can affect the fitness effects of

interactions (Palmer *et al.*, 2008; Chamberlain *et al.*, 2014; Wood *et al.*, 2018; Hafer-

Hahmann & Vorburger, 2020; Cassidy *et al.*, 2022). Often the details of how third parties

affect fitness effects are unknown. In the symbiosis between *D. discoideum* social

amoebae and *Paraburkholderia* bacteria, the third-party food bacteria is normally eaten

by the host but can also be carried or left behind. We studied how many food bacteria are

carried and left behind, and their impacts, by tracking fluorescently labeled food bacteria

(*K. pneumoniae*) during fruiting body formation by hosts.

Our most surprising finding was that hosts were more likely to carry food bacteria

after growing in food-poor environments (Figure 5). This result is surprising because it

means that hosts actively change the number of carried food bacteria depending on

environmental conditions. Since bacterial carriage is a joint phenotype affected by both

hosts and symbionts, *Paraburkholderia* symbionts may also play a role in modifying

carriage depending on the environment.

Our observation that food bacteria carriage depended on the environment (Figure

5) is possible evidence for mutual benefit between *D. discoideum* hosts and

Paraburkholderia symbionts. Host and symbionts are likely to benefit from modifying

carriage if future soil environments tend to resemble past soil environments Repeated

food-poor environments are where hosts should most benefit from carrying extra food

bacteria. *Paraburkholderia* symbionts could benefit from a stable association with hosts

that allows for continued dispersal, though more work is needed on the costs and benefits

of symbiosis for *Paraburkholderia* symbionts. We thus speculate that carrying more food

in these contexts may represent cooperation between hosts and symbionts that allows the

symbiosis to persist over repeated harsh environments. Such harsh conditions are

potentially an important force shaping this symbiosis (Scott *et al.*, 2022).

Questions remain about how hosts and symbionts affect the joint phenotype of

carrying food bacteria. *Paraburkholderia* are more often carried inside spores while food

bacteria are more often carried outside spores (Khojandi *et al.*, 2019), but the mechanism

for this is unknown. Hosts are able to normally modify the contents of fruiting bodies

through their immune cells that protect against toxins and bacteria by collecting potential

threats and dropping off during the slug stage (Chen *et al.*, 2007) but the role of these

immune cells play in the symbiosis (Brock *et al.*, 2016a) needs to be further explored.

Manipulation of phagosomes could also play a role in determining the contents of fruiting

bodies. *Paraburkholderia* symbionts increase the pH of phagosomes, presumably to

prevent host digestion of symbionts (Tian *et al.*, 2022). Similar modification of

lysosomes is used by pathogens to evade human immune cells during infection (Isberg *et*

al., 2009; Leseigneur *et al.*, 2020). More work is needed to understand how symbionts

and food bacteria get into fruiting bodies and how both hosts and symbionts contribute to

bacterial carriage.

We also found that *Paraburkholderia* infection prevents hosts from eating all the

food bacteria in an environment (Figure 1). We expected this result given the role of

Paraburkholderia in causing hosts to carry food bacteria (DiSalvo *et al.*, 2015) and leave

behind bacteria (Brock *et al.*, 2016b; Scott *et al.*, 2022). Infection with *Paraburkholderia*

likely interferes with host digestion through modification of phagosomes (Tian *et al.*,

2022). This indigestion hypothesis is supported by our finding that *Paraburkholderia*

infection did not affect development time (Figure 2). Development time should be faster

for infected hosts since they appear to eat less food and should starve sooner. One

possible explanation is that *Paraburkholderia* symbionts are making hosts sick and

delaying development enough to result in no difference between uninfected and infected

hosts.

A prior study suggested that leaving food bacteria uneaten because of prudent

predation may explain the cost of infection relative to uninfected hosts (Brock *et al.*,

2011). We instead found that these costs to hosts were better explained by

Paraburkholderia infection than by the amount of food bacteria left behind (Figure 3).

The quantity of left behind food bacteria may be too small to noticeably affect host spore

production since we observed that only a minority of left-behind bacteria was food

bacteria, with the majority being indigestible *Paraburkholderia* (Figure 4). A rough

calculation of lost growth also showed that uneaten food bacteria was not enough to

substantially increase host fitness. We thus conclude that there is little support that

prudent predation explains the cost of infection in this symbiosis.

Instead of hosts paying a cost in potential growth because they leave food bacteria

uneaten, we suspect that *Paraburkholderia* infection causes both observations: that hosts

leave food uneaten and infected hosts pay a cost. Support for this idea comes from our

findings that *Paraburkholderia* density was correlated with uneaten food bacteria and

Paraburkholderia density was a better predictor of host fitness in *P. hayleyella* uncured

hosts (Figure 3). The role of *Paraburkholderia* in reducing host spore production is more

evidence in support of there being some conflict in this symbiosis (DiSalvo *et al.*, 2015;

Scott *et al.*, 2022).

Conflict may be more pronounced between *P. hayleyella* and its hosts than

between *P. agricolaris* and its hosts. We found that *P. hayleyella* density could explain

some of the decrease in uncured host spore production but did not find the same for *P.*

agricolaris density (Figure 3; Supplemental File 1). This difference between *P.*

agricolaris and *P. hayleyella* may result from *P. hayleyella* being more toxic than *P.*

agricolaris (Shu *et al.*, 2018a; Haselkorn *et al.*, 2019; Khojandi *et al.*, 2019). In addition

to being more toxic, *P. hayleyella* also has a reduced genome relative to *P. agricolaris*

(Brock *et al.*, 2020; Noh *et al.*, 2022). Reduced genomes are a common result of

persistent host association in beneficial symbionts and pathogens (McCutcheon & Moran,

2012). Since *P. hayleyella* maintains toxicity, it likely retains some pathogenic ability

against hosts.

456    An interesting remaining question in the *D. discoideum-Paraburkholderia*

457  symbiosis and in other symbioses is the mechanisms controlling conflict and causing

458  symbiont infection to reduce fitness. One potential explanation is that *Paraburkholderia*

459  symbionts directly feed on host cells or otherwise extract nutrients from hosts. Measures

460  of *Paraburkholderia* density inside sori have so far not been found to be correlated with

461  host fitness within species (Miller *et al.*, 2020), though the more toxic *P. hayleyella* does

462  appear to infect a higher percentage of spores than less toxic *P. agricolaris* (Shu *et al.*,

463  2018a; Khojandi *et al.*, 2019). Another promising hypothesis for the *D. discoideum-*

464  *Paraburkholderia* symbiosis that deserves further study is that hosts have lower spore

465  production because of "collateral damage" from competitive interactions between

466  *Paraburkholderia* and food bacteria (Scott *et al.*, 2022). Competition between bacteria is

467  often mediated by chemical warfare (Granato *et al.*, 2019) that could reduce host *D.*

468  *discoideum* fitness as a side-effect.

469    Whether a specific symbiosis involves fitness alignment or conflict may depend

470  on a third party that affects the costs and benefits of symbiosis. Our results show that

471  third parties can have complex effects on symbioses; the symbiosis between *D.*

472  *discoideum* and *Paraburkholderia* appears to involve elements of conflict and

473  cooperation that are affected in multiple ways by food bacteria.

476  **References**

477  Bozzaro, S. 2019. The past, present and future of Dictyostelium as a model system.
478    *Int. J. Dev. Biol.* **63**: 321–331.

Brock, D.A., Callison, W.É., Strassmann, J.E. & Queller, D.C. 2016a. Sentinel cells,
symbiotic bacteria and toxin resistance in the social amoeba *Dictyostelium*
discoideum. *Proc. R. Soc. B Biol. Sci.* **283**: 20152727.

Brock, D.A., Douglas, T.E., Queller, D.C. & Strassmann, J.E. 2011. Primitive agriculture
in a social amoeba. *Nature* **469**: 393–396.

Brock, D.A., Jones, K., Queller, D.C. & Strassmann, J.E. 2016b. Which phenotypic traits
of Dictyostelium discoideum farmers are conferred by their bacterial
symbionts? *Symbiosis* **68**: 39–48.

Brock, D.A., Noh, S., Hubert, A.N.M., Haselkorn, T.S., DiSalvo, S., Suess, M.K., *et al.*
2020. Endosymbiotic adaptations in three new bacterial species associated
with *Dictyostelium discoideum* : *Paraburkholderia agricolaris* sp. nov.,
Paraburkholderia hayleyella sp. nov., and *Paraburkholderia bonniea* sp. nov.
PeerJ **8**: e9151.

Bronstein, J.L. 1994. Conditional outcomes in mutualistic interactions. *Trends Ecol.*
Evol. **9**: 214–217.

Burnham, K.P. & Anderson, D.R. 2004. Multimodel Inference: Understanding AIC and
BIC in Model Selection. *Sociol. Methods Res.* **33**: 261–304.

Cassidy, S.T., Markalanda, S., McFadden, C.J. & Wood, C.W. 2022. Herbivory modifies
plant symbiont number and impact on host plant performance in the field.
Evolution evo.14641.

Chamberlain, S.A., Bronstein, J.L. & Rudgers, J.A. 2014. How context dependent are
species interactions? *Ecol. Lett.* **17**: 881–890.

Chen, G., Zhuchenko, O. & Kuspa, A. 2007. Immune-like Phagocyte Activity in the
Social Amoeba. *Science* **317**: 678–681.

DiSalvo, S., Haselkorn, T.S., Bashir, U., Jimenez, D., Brock, D.A., Queller, D.C., *et al.*
2015. *Burkholderia* bacteria infectiously induce the proto-farming symbiosis
of *Dictyostelium* amoebae and food bacteria. *Proc. Natl. Acad. Sci.* **112**:
E5029–E5037.

Garcia, J.R., Larsen, T.J., Queller, D.C. & Strassmann, J.E. 2019. Fitness costs and
benefits vary for two facultative *Burkholderia* symbionts of the social
amoeba, *Dictyostelium discoideum*. *Ecol. Evol.* **9**: 9878–9890.

Granato, E.T., Meiller-Legrand, T.A. & Foster, K.R. 2019. The Evolution and Ecology
of Bacterial Warfare. *Curr. Biol.* **29**: R521–R537.

Hafer-Hahmann, N. & Vorburger, C. 2020. Parasitoids as drivers of symbiont
diversity in an insect host. *Ecol. Lett.* **23**: 1232–1241.

Haselkorn, T.S., DiSalvo, S., Miller, J.W., Bashir, U., Brock, D.A., Queller, D.C., *et al.*
2019. The specificity of *Burkholderia* symbionts in the social amoeba farming
symbiosis: Prevalence, species, genetic and phenotypic diversity. *Mol. Ecol.*
28: 847–862.

Horas, E.L., Metzger, S.M., Platzer, B., Kelly, J.B. & Becks, L. 2022. Context-dependent
costs and benefits of endosymbiotic interactions in a ciliate–algae system.
Environ. Microbiol. 1462-2920.16112.

Isberg, R.R., O'Connor, T.J. & Heidtman, M. 2009. The Legionella pneumophila
replication vacuole: making a cosy niche inside host cells. *Nat. Rev. Microbiol.*
7: 13–24.

Iwai, S., Fujita, K., Takanishi, Y. & Fukushi, K. 2019. Photosynthetic Endosymbionts
Benefit from Host's Phagotrophy, Including Predation on Potential
Competitors. *Curr. Biol.* **29**: 3114-3119.e3.

Keeling, P.J. & McCutcheon, J.P. 2017. Endosymbiosis: The feeling is not mutual. *J.*
Theor. Biol. **434**: 75–79.

Kessin, R.H. 2001. *Dictyostelium: evolution, cell biology, and the development of*
multicellularity. Cambridge University Press.

Khojandi, N., Haselkorn, T.S., Eschbach, M.N., Naser, R.A. & DiSalvo, S. 2019.
Intracellular Burkholderia Symbionts induce extracellular secondary
infections; driving diverse host outcomes that vary by genotype and
environment. *ISME J.* **13**: 2068–2081.

Lenth, R., Singmann, H., Love, J., Buerkner, P. & Herve, M. 2019. Package 'emmeans.'

Leseigneur, C., Lê-Bury, P., Pizarro-Cerdá, J. & Dussurget, O. 2020. Emerging Evasion
Mechanisms of Macrophage Defenses by Pathogenic Bacteria. *Front. Cell.*
Infect. Microbiol. **10**: 577559.

McCutcheon, J.P. & Moran, N.A. 2012. Extreme genome reduction in symbiotic
bacteria. *Nat. Rev. Microbiol.* **10**: 13–26.

Miller, J.W., Bocke, C.R., Tresslar, A.R., Schniepp, E.M. & DiSalvo, S. 2020.
Paraburkholderia Symbionts Display Variable Infection Patterns That Are
Not Predictive of Amoeba Host Outcomes. *Genes* **11**: 674.

Noh, S., Capodanno, B.J., Xu, S., Hamilton, M.C., Strassmann, J.E. & Queller, D.C. 2022.
Reduced and Nonreduced Genomes in *Paraburkholderia* Symbionts of Social
Amoebas. *mSystems* e00562-22.

Palmer, T.M., Stanton, M.L., Young, T.P., Goheen, J.R., Pringle, R.M. & Karban, R. 2008.
Breakdown of an Ant-Plant Mutualism Follows the Loss of Large Herbivores
from an African Savanna. *Science* **319**: 192–195.

Pinheiro, J. & Bates, D. 2006. *Mixed-effects models in S and S-PLUS*. Springer science &
business media.

Queller, D.C. 2014. Joint phenotypes, evolutionary conflict and the fundamental
theorem of natural selection. *Philos. Trans. R. Soc. B Biol. Sci.* **369**: 20130423.

Queller, D.C. & Strassmann, J.E. 2018. Evolutionary Conflict. *Annu. Rev. Ecol. Evol.*
Syst. **49**: 73–93.

Quides, K.W., Salaheldine, F., Jariwala, R. & Sachs, J.L. 2021. Dysregulation of host-
control causes interspecific conflict over host investment into symbiotic
organs. *Evolution* **75**: 1189–1200.

Raper, K.B. 1937. Growth and development of Dictyostelium discoideum with
different bacterial associates.

Scott, T.J. & Queller, D.C. 2019. Long-term evolutionary conflict, Sisyphean arms
races, and power in Fisher's geometric model. *Ecol. Evol.* **9**: 11243–11253.

Scott, T.J., Queller, D.C. & Strassmann, J.E. 2022. Context dependence in the
symbiosis between Dictyostelium discoideum and Paraburkholderia. *Evol.*
Lett. **6**: 245–254.

Shu, L., Brock, D.A., Geist, K.S., Miller, J.W., Queller, D.C., Strassmann, J.E., *et al.* 2018a.
Symbiont location, host fitness, and possible coadaptation in a symbiosis
between social amoebae and bacteria. *eLife* **7**: e42660.

Shu, L., Zhang, B., Queller, D.C. & Strassmann, J.E. 2018b. Burkholderia bacteria use
chemotaxis to find social amoeba Dictyostelium discoideum hosts. *ISME J.* **12**:
1977–1993.

Strassmann, J.E. & Queller, D.C. 2011. Evolution of cooperation and control of
cheating in a social microbe. *Proc. Natl. Acad. Sci.* **108**: 10855–10862.

Sun, B., Zhou, S. & Zhao, Q. 2003. Evaluation of spatial and temporal changes of soil
quality based on geostatistical analysis in the hill region of subtropical China.
Geoderma **115**: 85–99.

Tian, Y., Peng, T., He, Z., Wang, L., Zhang, X., He, Z., *et al.* 2022. Symbiont-Induced
Phagosome Changes Rather than Extracellular Discrimination Contribute to
the Formation of Social Amoeba Farming Symbiosis. *Microbiol. Spectr.*
e01727-21. Am Soc Microbiol.

Vos, M., Wolf, A.B., Jennings, S.J. & Kowalchuk, G.A. 2013. Micro-scale determinants
of bacterial diversity in soil. *FEMS Microbiol. Rev.* **37**: 936–954.

Wendling, C.C., Piecyk, A., Refardt, D., Chibani, C., Hertel, R., Liesegang, H., *et al.* 2017.
Tripartite species interaction: eukaryotic hosts suffer more from phage
susceptible than from phage resistant bacteria. *BMC Evol. Biol.* **17**: 98.

Wood, C.W., Pilkington, B.L., Vaidya, P., Biel, C. & Stinchcombe, J.R. 2018. Genetic
conflict with a parasitic nematode disrupts the legume-rhizobia mutualism.
Evol. Lett. **2**: 233–245.

Figure and Table Legends

Table 1: Clones used in experiments.

Type of infection	Description	*Paraburkholderia* Treatment	Clones
Cured and reinfected	Clones that were cured of any native symbionts with tetracyline and reinfected in the lab with 0.1% of their native symbiont	*P. agricolaris* infected	QS159, QS161, QS606, NC21
		P. hayleyella infected	QS395, QS45, QS38, QS23
		Uninfected control	QS6, QS138, QS472, QS527
Uncured	Clones with or without natural infections that have	*P. agricolaris* infected	QS494, QS756, QS788, QS113, QS453, QS70, QS606

	not been treated with antibiotics	*P. hayleyella* infected	QS45, QS101, QS46, QS2, QS23, QS529, QS38
		Uninfected control	QS4, QS6, QS14, QS18, QS9, QS8

Figure 1: Symbionts cause hosts to leave food uneaten. (A) Density left on plate of

uneaten *K. pneumoniae* food bacteria (measured by OD_{600}) for naturally uninfected, *P.*

agricolaris cured and reinfected, and *P. hayleyella* cured and reinfected hosts. Reinfected

hosts were reinfected with their natural symbiont partners. (B) Density left on plate of

uneaten food bacteria for naturally infected *P. agricolaris* and *P. hayleyella* hosts that

were either cured or reinfected. Shapes indicate clones (see Figure 1A in Scott et al.

2022). Dot and lines show mean and standard deviation, respectively. Asterisks indicate

significant differences (in panel A comparisons are between infected and uninfected).

Figure 2: Symbionts do not affect development. Developmental time points for cured

and reinfected (A) *P. agricolaris* and (B) *P. hayleyella* hosts at different stages of

development. Shapes indicate clones (see Figure 1A in Scott et al. 2022). Dot and lines

show mean and standard deviation, respectively.

Figure 3: Estimated effects of bacteria left on plates and infection status on host spore production in (A) cured and reinfected (B) and uncured hosts (see Table 1). Hosts without *Paraburkholderia* are included as baselines in all models. We compared models of host spore production (shown in different colors), predicted by *Paraburkholderia* density only (OD), uneaten food bacteria only (OD), both *Paraburkholderia* and uneaten food bacteria, or a categorical variable for infection status (infected with *Paraburkholderia* or not). Estimated effects are shown as points and 85% confidence intervals are shown as lines (null models with only the intercept are not shown). Asterisk color indicates the best model according to AICc (there is no asterisk in A because the null models fit with only the intercept are the best models).

Figure 4: Most of the bacteria left behind by hosts were not food bacteria. Percent of the

total left behind bacteria (includes food bacteria and Paraburkholderia symbionts) that

was food bacteria from cured and reinfected (A) and uncured (B) hosts. Shapes in A

show host clone replicates that were measured on two separate dates (see Figure 1A in

Scott et al. 2022). Points in B are independent, so we do not show clone identity. Dot and

lines show mean and standard deviation, respectively.

Figure 5: Hosts are less likely to carry food bacteria in sori after developing in food-rich

environments. (A) Photo of fluorescent *K. pneumoniae* food bacteria colonies plated from

an individual sorus. (B) Proportion of sori with carried food bacteria for reinfected *P.*

agricolaris hosts from food-poor and food-rich contexts. (C) Proportion of sori with

carried food bacteria for reinfected *P. hayleyella* hosts from food-poor and food-rich

contexts. Points show proportions and lines show the standard deviation. Asterisks

indicate significant differences (GLMM).