Hydrodynamic and anthropogenic disturbances co-shape microbiota rhythmicity

and community assembly within intertidal groundwater-surface water

continuum

Ze Zhao[1, 2, 3, 4,5], Lu Zhang[2, 3, 4, 5], Guoqing Zhang[2, 3, 4, 5], Han Gao[2, 3], Xiaogang Chen[2,

3], Ling Li[2, 3], Feng, Ju[2, 3, 4, 5*]

[1]College of Environmental & Resources Sciences, Zhejiang University, Hangzhou

310058, China

[2]Key Laboratory of Coastal Environment and Resources of Zhejiang Province, School

of Engineering, Westlake University, Hangzhou 310030, Zhejiang Province, China

[3]Institute of Advanced Technology, Westlake Institute for Advanced Study, 18

Shilongshan Road, Hangzhou, Zhejiang 310024, China

[4]Center of Synthetic Biology and Integrated Bioengineering, Westlake University,

Hangzhou, Zhejiang 310030, China

[5]Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou, Zhejiang

Province, China

*Corresponding to:

Dr. Feng Ju (Assistant Professor)

Tel: 571-87380995

Fax: 0571-85271986

E-mail: jufeng@westlake.edu.cn

Address: E2-326, Westlake University YUNGU CAMPUS, No.600 Dunyu Road,

Sandun Town, Xihu District, 310030, Hangzhou, Zhejiang Province, China

Abstract

Tidal hydrodynamics drive the groundwater-seawater exchange and shifts in microbiota structure in the coastal zone. However, how the coastal water microbiota structure and assembly patterns respond to periodic tidal fluctuations and anthropogenic disturbance remain unexplored in the intertidal groundwater-surface water (GW-SW) continuum, although it affects biogeochemical cycles and coastal water quality therein. Here, through hourly time-series sampling in the saltmarsh tidal creek, rhythmic patterns of microbiota structure in response to daily and monthly tidal fluctuations in intertidal surface water are disentangled for the first time. The similarity in archaeal community structures between groundwater and ebb-tide surface water (R^2=0.06, p=0.2) demonstrated archaeal transport through groundwater discharge, whereas multi-source transport mechanisms led to unique bacterial biota in ebb-tide water. Homogeneous selection (58.6%-69.3%) dominated microbiota assembly in the natural intertidal GW-SW continuum and the presence of 157 rhythmic ASVs identified at ebb tide and 141 at flood tide could be attributed to different environmental selection between groundwater and seawater. For intertidal groundwater in the tidal creek affected by anthropogenically contaminated riverine inputs, higher microbial diversity and shift in community structure were primarily controlled by increased co-contribution of dispersal limitation and drift (jointly 57.8%) and enhanced microbial interactions. Overall, this study fills the knowledge gaps in the tide-driven water microbial dynamics in coastal transition zone and the response of intertidal groundwater microbiota to anthropogenic pollution of overlying waters. It also highlights the potential of microbiome analysis in enhancing coastal water quality monitoring and identifying anthropogenic pollution sources (e.g., aquaculture pathogenic *Vibrio*) through the detection of rhythmic microbial variances associated

with intertidal groundwater discharge and seawater intrusion.

1. Introduction

Coastal ecosystems (e.g., salt marshes and mangroves) play a critical role in global climate regulation [1] and biogeochemical cycles [2, 3], yet they are currently facing a multitude of local and global challenges caused by climate change and human activities (e.g., pollution). In addition to the well-known river-runoff input and atmosphere deposition, submarine groundwater discharge (SGD) or porewater exchange from coastal aquifers contributes to nutrients and microbial loadings onto the coastal ecosystems, thus affecting the microbially-driven biogeochemical cycling and water quality within the groundwater-seawater mixing system [4-6]. Most existing studies have focused on microbiota structure in only one compartment of the intertidal zone, such as coastal sediment [7], groundwater or coastal seawater [8-11]. However, microbiota dynamics in the intertidal groundwater-surface water (GW-SW) continuum, especially in the context of the high structural variability and complexity impacted by bidirectional water exchange, remain poorly explored. Moreover, although periodic disturbances (e.g., tidal fluctuation) are revealed to influence the microbiota structure in various environments [12, 13], high-resolution profiling of microbiota dynamics is lacking. Hourly time-series sampling, e.g., during daily or monthly tidal cycles, would help to resolve microbiota dynamic patterns and assembly rules in the intertidal aquatic continuum and further identify key processes and factors governing the microbiota structure.

Hydrodynamic disturbance can cause steep variations in physicochemical

conditions over short spatial or temporal scales, and thus are regarded as a critical

factor controlling microbiota assembly and biogeochemical processes in coastal

sediments [12]. Various environmental factors (e.g., salinity, pH, temperature and

nutrient [8, 14-17]) are implicated in exerting selective pressures on the intertidal

groundwater or coastal seawater microbiome. The results are broadly consistent with

the niche-based theory, which asserts that deterministic processes such as

environmental filtering and biotic interactions largely control the patterns of

microbiota composition and distribution [18]. In addition to deterministic mechanisms,

stochastic processes, for instance, dispersal and ecological drift, also contribute to

microbial assembly. According to the well-established community assembly theory

[19, 20], microbiota structure and succession are controlled by four fundamental

ecological processes: selection (homogeneous selection and heterogeneous selection),

dispersal (homogenizing dispersal and dispersal limitation), drift and diversification,

which vary in their degree of determinism and stochasticity [21] and the relative

importance of the first three can be quantified by the null model [22, 23]. Until now,

only a few studies have reported on the microbiota assembly patterns in the intertidal

sediment [17, 24, 25] or coastal seawater [26], but the relative contribution of

ecological processes greatly varied across studies. Comparatively, microbiota

assembly processes have never been quantified in the intertidal GW-SW continuum,

especially under co-effects of hydrodynamic and anthropogenic disturbances.

Deterministic processes, including environmental or habitat filtering and biotic

interactions, are treated as joint selection forces in the community assembly and

interact dynamically to drive the observed community patterns [27, 28]. However,

their respective contribution cannot be parsed out based on the quantitative approach

of the null model [23]. Instead, deterministic processes and ecological relationships

can be interactively and topologically explored in a microbial co-occurrence or

molecular ecological network, which is fundamental for predicting microbial species

interactions and niche patterns in both natural [29, 30] and engineered [31-33]

ecosystems. Previous studies have reported that enhanced seawater intrusion

weakened the complexity of microbial co-occurrence associations in intertidal

sediment [24], whereas it decreased the network size but caused higher compactness

of groundwater microbial interactions [10]. However, microbial co-occurrence

patterns in the intertidal GW-SW continuum and their dynamics under periodic tidal

fluctuations and increasing anthropogenic impacts (e.g., land-sourced pollutants) are

almost blank. Therefore, complementing null model with community-wide

co-occurrence patterns will enable more complete insights into the ecological

processes shaping microbial community diversity in the context of the

determinism-versus-stochasticity dichotomy.

To explore the tide-driven dynamics of microbiota structure and community

assembly mechanisms in an intertidal aquatic continuum, we collected intertidal

groundwater and hourly time-series surface water in a natural saltmarsh tidal creek

(NT) located in Jiangsu Province, China (Fig. 1A), where mudflats are estimated to

cover more than 90% of the 954-km Jiangsu coast [34]. Meanwhile, a geographically

adjacent but human-influenced tidal creek (HT) was chosen and compared with NT to

resolve the response of intertidal groundwater microbiota to anthropogenic pollution

of overlying waters. The objectives of this study are: (i) to investigate the temporal

dynamics of microbiota structure in intertidal surface water with daily and monthly

tidal fluctuations; (ii) to explore the processes and mechanisms involved in microbiota

assembly in intertidal GW-SW continuum under tidal fluctuations; and (iii) to

elucidate the effects of anthropogenically contaminated river input on the spatial

distribution and community assembly of intertidal groundwater microbiota. This study

will provide the first and in-depth understanding of microbiota dynamics and

assembly mechanisms of the intertidal water co-shaped by hydrodynamic and

anthropogenic disturbances in the coastal ecosystem.

127 2. Materials and Methods

128 2.1. Field sample collection and physicochemical analysis

Samples were collected from the Saltmarsh Tidal Creeks located at Dafeng Milu

National Nature Reserve, Yancheng City, Jiangsu Province, China, in August 2020

(Fig, 1A, Table S1). The reserve is located in Yancheng Coastal Wetland, which is the

largest coastal wetlands nature reserve in China as listed in Ramsar Sites Information

Service. Conducing microbial research here is representative and significant. Twelve

groundwater samples were collected at low (L), mid (M), high (H) and above high (A)

tidal marks along three sampling transects of the natural tidal creek (Natural tidal

creek groundwater, NTG-1, NTG-2 and NTG-3). Eight groundwater samples were

collected at mid (M) tidal marks of 8 sampling sites in the human-influenced tidal

creek (Human-influenced tidal creek groundwater, HTG-1 to HTG-8). All

groundwater samples were collected at the lowest surface water level during ebb tide,

at a depth of 1.0-1.5 m. The outlet of HT is connected to the estuary of Chuandong

River and the estuarine water flows into the creek at flood tide. Besides, the

Terrestrial River upstream of HT discharges water into the creek at ebb tide, which is

controlled by a floodgate. The upstream of Terrestrial River flows through human

living (Yancheng City) and aquaculture area (bottom left of the map in Fig. 1A).

Higher concentration of antibiotics for human and aquaculture use (0.66-3.07 ng/L of

Metronidazole and 7.32-24.57 of Roxithromycin) and human-use-only Metformin

(5.89-19.35) and sweetener (150.25-582.62), were detected in HT surface water

compared with in NT surface water (0.20-1.11, 0-1.42, 0.49-2.12 and 64.06-128.21

ng/L) (Table S2), indicating serious water pollution in the HT (e.g., drug and nutrient)

due to aquaculture wastewater and domestic sewage discharge. On the contrary, the

downstream mouth of NT is far away from the Chuandong River estuary (about 4.5

km) and there is no terrestrial surface water input along the tidal creek, so it is

considered a natural tidal creek under less anthropogenic influence.

To investigate the transport and dynamics of microbial communities impacted by

bidirectional exchange of groundwater and coastal water during tidal fluctuations,

surface water samples were collected hourly for 14-15 hours from the Times Series

Station downstream of the NT (Natural tidal creek surface water, NTS) to monitor a

complete phase of ebb and flood tide (Fig. 1A). This monitoring was conducted once

at neap tide (N1-N14 on August 12[th]) and once at spring tide (S1-S15 on August

6[th]-7[th]) to investigate the difference in the intensity of tidal disturbances. Each water

sample (300-1400 ml) was filtered on-site through 0.2 µm pore size polycarbonate

filters (Millipore, USA) for about 30 minutes. The filters were transported to the

laboratory with dry ice in 24 h and stored at -20 ▢ until DNA extraction.

Physiochemical properties, including water depth, temperature (temp), pH, salinity,

dissolved oxygen (DO), dissolved organic and inorganic carbon (DOC and DIC), and

^{222}Rn (groundwater isotope tracer) of water samples were measured according to our

earlier publication [35] (detailed in Supplementary Text S1 and Fig. S1).

In total, 49 water samples were grouped according to their sources and sampling

times: (i) intertidal groundwater collected from human-influenced (HTG, 8) and

natural tidal creek (NTG, 12); (ii) time-series surface water samples grouped

according to two different grouping rules, namely NTS-Neap (14) and NTS-Spring

(15) collected during neap and spring tide, or NTS-Ebb (17) and NTS-Flood (12)

collected during ebb and flood tide designated by the *in-situ* observation of water

depth.

2.2.   DNA extraction, qPCR and 16S rRNA gene sequencing

Total genomic DNA was extracted using DNeasy PowerSoil Kit (Qiagen, Germany).

The V4-V5 regions of prokaryotic 16S rRNA genes were amplified using primers

515F and 926R targeting both bacteria and archaea (recommended by the Earth

Microbiome Project) [36], and sequenced on the Illumina NovaSeq 6000 platform

with 250-bp paired-end reads at the Magigene Biotechnology Co. Ltd (Guangzhou,

China). In addition, real-time quantitative PCR (qPCR) was performed in technical

triplicates to quantify the 16S rRNA gene copies (as a proxy for prokaryotic biomass)

in water samples using primers 515F and 926R on a Jena qTOWER^{3}G (Germany)

(detailed in Supplementary Text S2).

The raw sequencing data were processed following the QIIME2 pipeline [37]

(detailed in Text S3). In brief, forward and reverse reads were trimmed to 240 and 220

base pairs and low-quality reads were removed using Trimmomatic [38]. All reads

were then subjected to de-noising using the DADA2 pipeline with default parameters

[39]. Taxonomic assignment of amplicon sequence variant (ASVs) was conducted

using the Silva v138.1 database.   All the following statistical analysis and

visualization were performed with R software (4.0.5). The outputs from QIIME2,

including table of amplicon sequence variants (ASVs), taxonomy and tree files were

processed by R package phyloseq [40], MicrobiotaProcess and microeco [41]. Briefly,

ACE and Shannon indices were used to evaluate the alpha-diversity of microbial

communities. The ASV table was transformed to relative abundance using phyloseq

function *transform_sample_counts.* Non-metric multidimensional scaling (NMDS) analysis and permutational multivariate analysis of variance (PERMANOVA) was performed based on weighted UniFrac distance. To test whether each of the ASVs in time-series surface water displayed rhythmic patterns in response to tidal fluctuations, that is, recurrent changes over time, we performed Lomb Scargle periodogram (LSP, as implemented in the lomb package) analysis for relative abundance of ASVs in NTS-Neap and NTS-Spring, respectively. The method has previously been used for testing the seasonality in marine microbial communities [42]. Briefly, we obtain density distribution of each period for each ASV, showing which is the most recurrent period for each ASV. Peak normalized power (PNmax) measures the strength of this period. Microbial ASVs are considered rhythmic ASVs only of PNmax > 0.1 and $p < 0.05$ (detailed in Supplementary Text S4). In order to better observe the dynamics of ASV abundance in response to tidal fluctuations, cluster heatmap was constructed with Centered Logarithm Ratio values (CLR, adding a pseudocount of 1) of rhythmic ASVs. Random forest analyses were performed using the 'randomForest' package with default parameters to predict physicochemical variables using rhythmic ASVs. Stratified random sampling was used to select 70% samples to be used as the training dataset and the model were then validated on the remaining 30% samples (detailed in supplementary Text S5). Furthermore, the rhythmic patterns of physicochemical factors were also characterized by LSP analysis with same parameters. The distance-decay relationships (DDRs) [43] of archaeal and bacterial communities in groundwater groups were measured based on taxonomic diversity by linear regression between log-transformed community similarity (Bray-Curtis distance) and geographic distances.

2.3. Community assembly processes of tidal creek water microbiota

A recently proposed framework 'iCAMP' [22] was used to quantify the contribution

of different ecological processes to community assembly based on phylogenetic

bin-based null model analysis. The observed taxa were first divided into different

'bins' according to a phylogenetic signal threshold ($ds = 0.2$) within the phylogenetic

tree and the relative importance of heterogeneous selection, homogeneous selection,

homogenizing dispersal, dispersal limitation, and drift in individual 'bins' was

calculated to evaluate the dominated 'bins' and dominated taxa contributing to the

ecological processes. In brief, 37294 observed ASVs were divided into 687

phylogenetic bins (renamed to Bin1, Bin2, *etc.*). Phylogenetic tree of 300 most

abundant Bins (89.6-97.9% of total contribution to community assembly) and the

relative importance of ecological processes in each Bin were displayed by an online

tool 'iTOL' [44]. Furthermore, all 687 Bins were grouped based on their top member

lineage at class level, named 'Bin-class'. Contributions of the Top 15 and other

Bin-class to dominant ecological processes among groups were visualized by R

package 'pheatmap'. In addition, Mantel tests with constrained permutations were

performed to reveal the correlations between individual processes and various

environmental factors and the scripts are publicly available from GitHub of 'iCAMP'

(https://github.com/DaliangNing/iCAMP1).

2.4. Checkerboard score and co-occurrence network analysis

Checkerboard score (C-score) model was used as the metric of species segregation

and its variance (C_{var}-score) was used as the metric of both species segregation and

aggregation in the intertidal water microbiota [45]. The higher the observed

standardized effect (SES) values of C-score, the greater degree of species segregation

relative to a random distribution. The C-score tests were performed using the scripts

of package MbioAssy1.0 (https://github.com/emblab-westlake/MbioAssy1.0) [31] and

more information can be found in Text S6. Co-occurrence network and global

network properties were generated using the Molecular Ecological Network Analysis

Pipeline [46, 47] and visualized with Gephi v0.9.2. Six networks were constructed

based on abundance profiles of ASVs that present in at least half of the samples of

each sample group. The resulting correlation matrix (Pearson correlation coefficient)

was analyzed with the random matrix theory- (RMT-) based approach to determine

the correlation threshold. Moreover, 1000 corresponding Erdös-Réyni random

networks, the small-world coefficient (σ) and the keystone nodes were generated for

each network and more details are described in Text S7.

3. Results

3.1. Microbiota diversity and structural variations in intertidal water continuum

We first investigated the diversity and structural variations of prokaryotic (i.e.,

archaeal and bacterial) communities in intertidal groundwater and surface water. The

total concentration of water prokaryotic biomass, as measured by the copy number of

16S rRNA gene, varied from 3.96×10^4 to 6.36×10^6 copies/mL (Table S3), with

significantly higher median concentration (Wilcoxon rank-sum test, $p < 0.05$, Fig. S2)

in the groundwater or porewater (1.94×10^6 copies/mL) than the surface water ($3.61 \times$

10^5 copies/mL), as well as in the surface water during the neap tide (5.68×10^5

copies/mL) than the spring tide (3.07×10^5 copies/mL). Tide-driven prokaryotic

diversity and structural variations in intertidal groundwater and surface water

microbiota were also assessed comparatively. Based on ACE and Shannon indices

(Wilcoxon rank-sum test, $p < 0.05$, Fig. 1B), archaeal alpha-diversity in the natural

tidal creek groundwater (NTG) was significantly higher than (NTS-Flood and

NTS-Spring) or equal to (NTS-Ebb and NTS-Neap) that in the surface water, whereas

bacterial alpha-diversity showed a reversed trend, i.e., lower in the NTG than the

NTS-Ebb, NTS-Neap and NTS-Spring, and equal to the NTS-Flood. Comparatively,

human-influenced creek groundwater (HTG) harbored significantly higher

prokaryotic alpha-diversity than not only the groundwater (NTG) but also the surface

water (NTS) of the natural creek.

When weighing relative abundance, archaeal communities were significantly

more abundant in the human-influenced creek groundwater (HTG: 1.76%-49.66%)

than the natural one (NTG: 0.56%-14.25%) (Wilcoxon rank-sum test, $p = 0.02$, Table

S4), while they were more enriched in the downstream than upstream groundwater of

both creeks (Fig. S3). Overall, archaeal communities in both groundwater and surface

water were mainly comprised of phylum *Nanoarchaeota* (12.0-76.4%),

Crenarchaeota (6.2-67.0%), *Thermoplasmatota* (0.3-9.3%), and *Halobacterota*

(0.3-53.0%), which showed remarkably dynamics of relative abundance in broad

ranges and across sampling groups (Fig. 1C). For example, family *Woesearchaeales*

and SCGC AAA011-D5 were the most abundant Archaea in NTG and NTS-Ebb,

whereas family *Nitrosopumilaceae* and Marine Group II (typical seawater archaea)

were most abundant in NTS-Flood (Fig. S4). Notably, up to 9.8% of unclassified

archaea were detected in groundwater, suggesting the prevalence of numerous

yet-to-be-characterized 'archaeal dark matter' in the coastal zone. Moreover, we

observed weak but statistically significant structural differences (PERMANOVA, $R^2 =$

0.14, $p = 0.02$, Table S5) in archaeal communities between HTG and NTG (Fig. 2A),

as well as higher community dissimilarities within HTG (Wilcoxon rank-sum test, $p <$

0.05). Dynamics of archaeal community structure in the surface water were strongly

related to tidal cycles, with similar archaeal structure observed among NTG and

NTS-Ebb ($R^2 = 0.06$, $p = 0.202$) and significant variations between NTS-Ebb and

NTS-Flood ($R^2 = 0.60$, $p < 0.001$), which was further confirmed by NMDS

ordinations in Fig. 2A. This result well suggested the microbial transport through

SGD: i) the transport of groundwater microbes to surface water via groundwater

discharge at ebb tide and ii) the transport of coastal microbes to intertidal surface

water via seawater intrusion at flood tide.

Echoing archaeal communities, bacterial communities showed significant

structural differences between HTG and NTG ($R^2 = 0.16$, $p = 0.002$), as well as lower

community dissimilarities within the NTG (Fig. 2B). The overall groundwater

bacterial communities were dominated by *Proteobacteria* (19.0-82.7%), *Bacteroidota*

(1.8-40.7%) and *Campilobacterota* (1.5-39.7%), while the surface water communities

were dominated by *Proteobacteria* (28.8-71.5%), *Bacteroidota* (5.5-21.4%),

Cyanobacteria (1.2-39.7%), *Actinobacteriota* (6.8-22.5%) and *Planctomycetota*

(0.8-22.5%) (Fig. 1D). Unlike archaea, bacterial communities were structurally

different among NTG, NTS-Ebb, and NTS-Flood ($R^2 = 0.46$-0.58, $p < 0.01$, Table S5),

as illustrated by their clearly separate clusters in the NMDS diagram (Fig. 2B).

Besides, PERMANOVA analysis also showed significant community structural

differences between NTS-Neap and NTS-Spring ($R^2 = 0.15$, $p < 0.01$).

3.2. Microbiota rhythm and spatial dynamic patterns in intertidal surface

water

To further resolve the high-resolution temporal dynamics of surface water microbiota,

we performed NMDS analysis within the surface water samples (n = 29) to

investigate the time-series community structural dynamics at hourly intervals during

two whole tidal cycles including neap tide (August 12[th]) and spring tide (August

6[th]-7[th]). Complying with our hypothesis that periodic hydrodynamic disturbance

should generate microbiota rhythm patterns, we observed four clusters of surface

water samples separated by ebb-flood tide and neap-spring tide with a circular

trajectory (Fig. 3A), indicating rhythm patterns of microbiota structure remarkably in

the surface water following daily (ebb-flood) and monthly (neap-spring) tidal cycles.

The intriguing circadian rhythm patterns of microbiota dynamics were superimposed

by highly synchronized and complex bacterial community dynamics and contrastingly

weak (but abrupt) and simpler archaeal community dynamics (Fig. S5).

Further, we correlated surface water microbiota (NTS-Neap, NTS-Spring, and

NTS) to eight environmental factors, i.e., water depth, temperature, salinity, pH, DO,

DIC, DOC, and ^{222}Rn, to verify the aforementioned microbiota rhythm patterns (Fig.

3B). These factors are generally regarded as indicators of tidal cycles [2], and seven

of them (except for DOC) were identified to display significant rhythm patterns with

tidal cycles (PNmax = 0.34-0.77 and P-value < 0.01, Table S6). The results showed

that seven factors, except for temperature, exhibited significant correlations with

surface water microbiota during neap tide, spring tide or both, supporting the rhythm

patterns of microbiota structure in tidal creek surface water.

In order to test if each microbial ASV exhibited significantly rhythmic pattern

(identified as rhythmic ASVs) during tidal time series, that is, their abundance

exhibited periodic fluctuations, we performed Lomb Scargle periodogram (LSP) test

in NTS-Neap and NTS-Spring, respectively (see Methods and Supplementary Text

S4). We identified 271 rhythmic ASVs in NTS-Neap that spanned rhythmic period of

1.9 to 13.0 hours (Fig. S6 and Table S7), accounting for 40.5% to 72.3% of total

relative abundance. And 128 rhythmic ASVs (2.0 to 14.0 hours of periods and 6.8% to

47.5% of abundance) were identified from 1686 ASVs in NTS-Spring. Most of the

rhythmic period of rhythmic ASVs (13 and 14 h, 335 out of 399) is aligned with the

tidal cycles (~12 h) of the sampling site. Rhythmic prokaryotic subcommunities (i.e.,

rhythmic ASVs in NTS-Neap and NTS-Spring) exhibited a highly consistent rhythm

pattern with the whole communities, whereas weaker rhythmic pattern was observed

in the non-rhythmic subcommunities (Fig. S7A), as suggested by the lower variance

explained by the first and second axes based on PCoA analysis (weighted-UniFrac

distance, 66.2% and 17.1% in rhythmic subcommunities, 36.7% and 23.5% in

non-rhythmic subcommunities) (Fig. S7B). This suggested that the rhythmic patterns

observed in whole prokaryotic communities were mainly shaped by the numerous and

abundant rhythmic subcommunities.

Based on clustering heatmap plot for centered logarithm ratio abundances of 271

rhythmic ASVs from LSP test in NTS-Neap (Fig. 3C), 115 ASVs showed higher

abundance at ebb tide (ebb cluster) and 99 ASVs exhibited high abundance at flood

tide (flood cluster), whereas the remaining 57 ASVs with low abundance were

detected in only a few time points and their rhythmic characteristics were not clear

(Table S7). For example, family *Rhodobacteraceae* (21), *Woesearchaeales* (9), and

Cyanobiaceae (7) dominated in ebb cluster and Clade I (8) and SAR86 clade (7)

dominated in flood cluster. For clustering analysis of 128 rhythmic ASVs in

NTS-Spring (Fig. 3D), 64 ASVs showed higher abundance at ebb tide. Among them,

family *Cyanobiaceae* (9), *Flavobacteriaceae* (8) and *Microbacteriaceae* (4) showed

high abundance at ebb tide in both NTS-Neap and NTS-Spring. Forty-one ASVs

displayed higher abundance at flood tide in NTS-Spring, for example, *Nitrincolaceae*

(5) and *Thiomicrospiraceae* (4), all of which exhibited higher abundance in the early

ebb tide after flood tide (Fig. 3D). Remaining 23 ASVs showed low abundance across

samples. Overall, the union set of 157 rhythmic ASVs, which obtained by combining

rhythmic ASVs of ebb clusters in NTS-Neap (115) and NTS-Spring (64), represented

unique and abundant microbes in intertidal groundwater (Table S7). In turn, the other

union set of 141 rhythmic ASVs showing higher signals at flood tide (flood cluster)

represented unique microbes in coastal seawater. For the rhythmic ASVs at genus

level with mean relative abundance greater than 0.1% in ebb-tide and flood-tide

groups, we selected 23 rhythmic genera with more than five-fold abundance

difference as microbial indicators for intertidal groundwater discharge (e.g., SCGC

AAA011-D5) and seawater intrusion (e.g., *Candidatus* Nitrosopumilus), respectively

(Table S8). To validate that these rhythmic ASVs were useful indicators of water

physiochemistry, eight random forest regression models were constructed to predict

each physicochemical variable. Linear regression models comparing the predicted to

actual variables showed stronger correlations (adjusted R^2 = 0.80-0.99, Fig. S8) and

regression slopes closer to 1 (0.56-1.15, which represents a perfect prediction), except

for temperature and DO, indicating high predictive accuracy and great differentiation

between ebb-tide and flood-tide surface water samples.

Besides temporal dynamics of tidal creek water microbiota, we also measured

distance-decay relationships (DDRs) based on Bray-Curtis dissimilarity of community

structure to understand the spatial microbiota turnovers in intertidal groundwater.

Both archaeal and bacterial communities showed significant DDR patterns (P < 0.05),

i.e., decreased community similarity with increased geographical distance, whether

for the HTG or the NTG (Fig. S9). Moreover, bacterial communities exhibited

stronger DDR patterns in the NTG (slope = -0.3443) than HTG (slope = -0.1926),

while archaeal communities showed limited differences in their decay patterns

between the creeks, revealing that human influence in the coastal zone may reduce the

spatial heterogenicity of Bacteria which are overall more sensitive to human influence

than Archaea. In particular, archaeal communities showed a far steeper slope

(-0.99/-1.09 vs. -0.19/-0.34) and higher fitness value (0.43/0.29 vs. 0.23/0.18) of

DDRs than bacterial communities, indicating an overall stronger turnover rate and

higher spatial heterogenicity of Archaea than Bacteria in the intertidal groundwater.

3.3. Microbiota assembly processes and mechanisms in intertidal groundwater

and surface water

To explore mechanisms underpinning spatiotemporal dynamics and patterns in the

intertidal GW-SW continuum, the relative contribution of five ecological processes

(see methods) to microbiota assembly was quantitatively evaluated. Overall,

homogeneous selection (38.5%-69.3%), dispersal limitation (5.2%-49.0%) and drift

(8.8%-22.5%) were the dominant processes that together explained 96.3% to 98.5%

of community assemblage (Fig. 4A and Table S9), and therefore the subsequent

analysis was focused on them. The results showed that homogeneous selection

(58.6%-69.3%) predominated in the microbiota assembly of natural GW-SW

continuum, including NTG and NTS. And the contribution of dispersal limitation

markedly decreased in NTS-Flood (5.2%) and NTS-Spring (7.1%). Compared to the

NTG, the importance of dispersal limitation dramatically increased and predominated

in the human-influenced groundwater (49.0% in HTG), accompanying decreased

homogeneous selection (38.5%) and drift (8.8%).

Further, the above-mentioned relative importance of each ecological process was

quantified down to the refined level of individual phylogenetic lineages (Bins)

defined in iCAMP (Table S11) [22]. Our results revealed that homogeneous selection

was dominated in 11.4% to 13.0% of 687 Bins (Table S10), accounting for a summed

abundance of 49.3% to 54.1%, while dispersal limitation dominated in 26.1% to 71.0%

Bins (abundance: 2.9%-39.2%) and drift dominated in 16.6% to 59.0% Bins

(abundance: 10.5%-43.5%), revealing the prevalence and dominance of homogeneous

selection in a few species with disproportionally higher abundance in the intertidal

GW-SW continuum. By exhibiting the contribution of each Bin, we found that the

contribution of a few high-abundance Bins (e.g., the sixteen Bins marked in Fig. 4B)

was remarkably higher than that of others and thus cumulatively accounted for the

major contribution in the microbiota assembly. For example, the most abundant Bin1

(7.3%) contributed up to 11.6% of homogeneous selection in NTG. The results of

grouping Bins at class level, i.e., 104 Bin-class (Table S12), also showed the major

contribution of abundant Bin-class in the natural tidal creek (Fig. 4C). For example,

the most abundant Bin-class *Alphaproteobacteria*, *Gammaproteobacteria* and

Bacteroidia contributed most to both homogeneous selection, dispersal limitation and

drift in the natural GW-SW continuum. Comparatively, the enhanced dispersal

limitation in the HTG was attributed to the jointly increased contribution of high and

low abundance Bins, which was confirmed by the higher contribution of Bin-class

Nanoarchaeia (8.5%), *Dehalococcoidia* (3.6%), and *Clostridia* (1.4%) (Fig. 4C).

Overall, these results indicated that high-abundance microbial lineages contributed

most to homogeneous selection and drift in the natural intertidal GW-SW continuum,

whereas the accumulated contributions of both abundant and rare lineages resulted in

the enhanced dispersal limitation in the human-influenced groundwater.

3.4. Environmental factors influencing the ecological processes of microbiota

assembly

Correlations between changes in environmental factors and dominant ecological

processes among paired samples in each group were further determined with Mantel

test ($p < 0.05$) to identify the major drivers of ecological processes. Overall,

homogeneous selection and dispersal limitation showed statistically similar

correlation patterns (Mantel test for p-value correlation matrix: r = 0.70, $p < 0.05$)

with various environmental factors, while weaker and less frequent correlations were detected between drift and environmental factors (Fig. 5 and Table S13). Both homogeneous selection and dispersal limitation showed significant correlations with salinity, pH, DO, DIC and DOC in NTS-Neap and NTS-Spring. Considering the fluctuating environmental factors with tidal flows and their significant inter-correlations (Fig. S10), significant correlations between these factors and dispersal limitation may indicate the influence of GW-SW mixing on the temporal microbiota turnover in intertidal surface water. Notably, the environmental factors associated with homogeneous selection were quite different in the HTG (DIC and DOC) and NTG (salinity, pH and DO), and so was dispersal limitation and drift, suggesting a potential effect of carbon nutrients on the microbiota assembly in the human-influenced groundwater.

3.5. Deterministic microbial segregation and co-occurrence patterns of tidal creek water microbiota

Once stochastic and deterministic processes are demonstrated to co-drive the microbiota assembly in the intertidal GW-SW continuum, we applied C-score and co-occurrence network analyses to explore the contribution of microbial interactions (e.g., competition and mutualism) [22] on the observed assembly patterns, which are included in selection process but could not parse out by the null model analysis. Based on positive values of standardized effect size (SES) of C-score and their ranks (Fig. 6A), the intertidal water microbiota exhibited significant and stronger segregated distributions during Neap than Spring tide (51.2 vs 21.8), during Ebb than Flood (21.8 vs 13.2), and in HTG than NTG (13.9 vs 5.3), compared with those expected under 'random' null hypothesis (Table S14). Moreover, the SES values of both C-score (5.3) and C_{var}-score (3.9) were the lowest in NTG (5.3 and 3.9), while their values were

much higher in HTG (13.9 and 11.7), revealing an overall lower degree of both

species segregation and aggregation (e.g., resulted from competition and cooperation)

in the natural than the human-influenced groundwater.

Because C-score analysis revealed significant roles of both cooperative (e.g.,

synergistic) and competitive (e.g., antagonistic) microbial interactions in microbiota

assembly [48], we further explored microbial co-occurrence patterns by establishing

six networks for intertidal groundwater or surface water microbiota (Fig. 6B).

Topological analysis showed that the networks displayed significantly different

topological structures from their corresponding random networks of identical size.

They all had highly modular structures (modularity > 0.4) and showed 'small-world'

properties (σ > 1) with coefficients between 1.93 to 6.19 (Table S15), revealing

non-random network structures and assembly patterns that were unlikely to appear

due to chance. In the natural intertidal GW-SW continuum, intriguingly, a higher

percentage of rhythmic ASVs was observed in each surface water network

(61.5%-73.7%) than in the groundwater network (44.8%, Table S15). Meanwhile,

seven of the eight keystone taxa (5 module hubs and 3 connectors) identified in the

surface water network were also rhythmic ASVs (Table S16). The key role of

rhythmic ASVs observed in the surface water networks suggested the impact of tidal

fluctuations on microbial interactions in intertidal surface water.

The anthropogenic influence on co-occurrence patterns of groundwater

microbiota was comparatively explored between HTG and NTG (Fig. 6B). The

number of nodes (358), edges (2413), graph density (0.038), and average clustering

coefficient (0.491) of HTG network were all remarkably higher than that of NTG

network (288, 713, 0.018 and 0.308, respectively), suggesting higher frequencies of

species co-occurrence associations in the human-influenced than natural groundwater.

Moreover, mantel tests showed that both temperature and DIC/DOC correlated

positively (though not always significant) with almost all modules in HTG network (P

< 0.05, Fig. S11), whereas none of these parameters correlated with modules in NTG

network, suggesting they played a potential essential role in shaping the modularized

structures in HTG network representing discrete ecological niches in the

human-influenced tide creek. Ten keystone ASVs were observed in HTG network and

were all identified as 'connector', but only two keystone ASVs (one 'module hub' and

one 'connector') were identified in NTG network (Table S16 and Fig. S12).

4. Discussion

Tidal hydrodynamics drive the groundwater-seawater exchange and shifts in

microbiota structure in the intertidal zone. However, the periodic tidal fluctuations

and complex geological environment make it difficult to accurately resolve the

temporal dynamics of microbiota structure based on a few discrete time-series

samples in this intertidal groundwater-surface water continuum. Through hourly

time-series microbiota profiling and modeling, this study provides first insights into

the microbiota rhythmic patterns in response to periodic tidal fluctuations in intertidal

surface water of Yancheng, Jiangsu Province, China, and identifies homogeneous

selection as the key process driving microbiota structuring and assembly in the

intertidal GW-SW continuum. Furthermore, by comparing the groundwater

microbiota of a tidal creek affected by anthropogenically contaminated river inputs

with the natural creek, we found that shifts in prokaryotic biodiversity and community

structure were accompanied by enhanced dispersal limitation, drift and microbial

interactions but weakened homogeneous selection.

4.1. Tide cycles drive periodic transport and rhythm patterns of microbiota

along the natural intertidal water continuum

The high-resolution temporal dynamics of microbiota structure in tidal creek surface

water were followed during whole tidal cycles. Our results revealed that tidal

fluctuations shaped the daily (i.e., ebb-flood) and monthly (i.e., neap-spring)

dynamics of microbiota structure. Unlike archaea transport mainly via groundwater

discharge during ebb tide, potential multi-source transport mechanisms other than

groundwater discharge may account for the unique bacterial community structure of

ebb-tide surface water that differs from the intertidal groundwater. For example,

bacterial biota might also originate from the release of particle-attached bacteria via

particle resuspension from surface sediments and flushing of the surficial sediment

bacteria [49]. The likelihood of such events in our study is confirmed by the unique

higher abundance of family *Cyanobiaceae* (Fig. S4) and genera *Candidatus* Aquiluna,

Roseibacillus, *NS11-12 marine group* (Fig. S13) in the ebb-tide surface water

compared to the intertidal groundwater and flood-tide surface water. The family

Cyanobiaceae have been reported to prefer to develop on the surface of saltmarsh

tidal mudflats [7, 50]. Taken together, tidal fluctuations pose a domain-level

community-specific influence on the transport mechanisms of Archaea and Bacteria.

This discrepancy may be related to the fact that archaeal prefer to live in the

subsurface sediment, while bacteria have a wider niche breadth in the intertidal zone

[51].

Especially, based on the hourly time-series sampling of intertidal surface water,

we have further found that periodic tidal fluctuations on a daily and monthly basis

lead to the rhythm patterns of microbiota structure. The rhythmicity of environmental

water microbiota has only been implicated in several long-term studies on the

seasonal or circadian reoccurring patterns of marine water community structure [42,

52, 53]. Although the periodic shift of coastal microbiota structure associated with

tidal fluctuation has been reported before through observation of a few discrete-time

samples [49], our results based on hourly time-series sampling demonstrated that tidal

cycles dominated the rhythmic recurrent of not only groundwater-related microbiota

during ebb tide but also seawater-related microbiota during flood tide. The rhythmic

ASVs considered as microbial indicator of intertidal groundwater discharge (ebb-tide)

and coastal seawater (flood-tide) shared very few at the family level (6 of 214 shared

in NTS-Neap and 13 of 105 shared in NTS-Spring, Table S7). Even for these shared

family, they are classified as different genera. The rhythmic subcommunities

suggested that different ecological niches among intertidal groundwater and coastal

seawater selected for different microbial populations, which was responsible for the

unique but abundant biota observed in ebb-tide and flood-tide, respectively. In

construct, the non-rhythmic subcommunities with similar abundance in ebb and flood

tide surface water should be adapted to both groundwater and seawater environment,

and dispersal with the bidirectional water exchange. This was confirmed by the higher

contribution of rhythmic genera to homogeneous selection relative to dispersal

limitation in intertidal groundwater (e.g., genera *Cyanobium* PCC-6307 and

Rhodobacteraceae) (Table S10). Intriguingly, the rhythm pattern during spring tide

was weaker (fewer rhythmic ASVs) than during neap tide, probably because the high

tide level inundates the saltmarsh and overlying water is discharged together with the

groundwater at ebb tide, interfering with the characterization of rhythm pattern.

Besides, change of tidal periods in the investigated tidal creek resulting from tidal

asymmetry (e.g., decrease in the duration of ebb and increase in the duration of flood)

[54], as well as the hysteresis patterns on different time scales (e.g., diel and

neap-spring) [35], may explain the occurrence of rhythmic period shorter than tidal

cycles.

Furthermore, the random forest models represented high predictive accuracy in
physicochemical variables and great differentiation between ebb-tide and flood-tide
surface water samples. Given that these variables, e.g., salinity, pH [55] and ^{222}Rn
[56], were commonly used as environmental tracers to establish intertidal SGD or
seawater intrusion in the coastal zone, our results demonstrated the potential of
microbiome analysis to enhance monitoring of coastal water quality and pollution
sources by detecting rhythmic microbial variances that signify anthropogenic
contamination from intertidal groundwater discharge and/or seawater intrusion. For
example, the presence of high abundance species such as *Vibrio mimicus* (0.47%) and
Vibrio cholerae (0.15%) in HTG-1M, which are important pathogens associated with
aquaculture [57], suggested potential aquaculture pollution in intertidal groundwater.
This finding supports the hypothesis that high abundance of such fecal or aquaculture
indicators, detected as ebb-tide rhythmic ASVs in coastal seawater, may indicate
coastal anthropogenic pollution resulting from intertidal groundwater discharge. In
addition, among the 23 rhythmic genera we selected as microbial indicators, SCGC
AAA011-D5 and *Woesearchaeales* belonging to order *Woesearchaeales* were also
reported to be predominant (45%-76% of archaea) in intertidal unconfined aquifer but
accounted for less than 1% in seawater, whereas genera *Candidatus* Nitrosopumilus
and Marine Group II exhibited the opposite trend [58]. This suggested that these
rhythmic ASVs and filtered rhythmic genera have a broad applicability in indicating
coastal groundwater discharge and seawater intrusion. Further validation can be
achieved through larger-scale multi-habitat studies in the future. We propose to
leverage the informatics of microbiota to add biologically relevant information to the
monitoring and assessment of coastal ecosystem.

4.2. Homogeneous selection and microbial interactions co-contribute to the microbiota assembly in the natural intertidal water continuum

Given the tide-driven hydrodynamic disturbance and the accompanying environmental gradients across the temporal and spatial scales in the coastal zone [59], the corresponding ecological processes, i.e., selection and dispersal, are expected to play a key role in the microbiota assembly in the natural tidal creek. Despite microbial dispersal is expected to be enhanced under periodic tidal fluctuations, high levels of dispersal can result in a homogenization of the local and regional species pool, which can lead to an overall short-time increase in stochasticity (during initial formation phases of tidal creeks) followed by an increase in deterministic selection at a regional scale that decreases species diversity [60] (Fig. 1B). Meanwhile, the enhanced homogeneous selection can lead to increased species competition, suggested by the species segregated distributions based on C-score analyses. Compared with the surface water, weakening of water mixing disturbance in intertidal groundwater resulted from the barrier effect of cohesive sediment can account for the relatively lower dispersal rates. We conclude that homogeneous selection and microbial interactions co-contributed to the microbiota assembly patterns in the natural intertidal water continuum.

4.3. Anthropogenic influences alter microbiota structure and community assembly patterns in intertidal groundwater

Intertidal zone harbors abundant and diverse microbial environments owing to the land-sea interaction and are extremely sensitive to natural changes and nearshore human activities [50, 61]. Nutrient loading derived from human activities and tidal dynamics is among the critical pressures affecting the intertidal environments [62].

Considering the high spatial heterogeneity of intertidal environment, we selected two

geographically adjacent (within 6 km) tidal creeks for comparative analysis of

intertidal groundwater microbiota, so as to maintain a basically consistent

environmental conditions except for the single variable of anthropogenic disturbance

in NT. Our study found that the human-influenced tidal creek exhibited both higher

species diversity and community structural dissimilarities of groundwater microbiota

than the natural one, whether for archaeal or bacterial communities. This contrasting

profile is expected to result from the shift in microbiota structure caused by polluted

creek surface water intrusion due to Terrestrial River water input upstream (during

ebb) and estuary-water input downstream (during flood) of the creek.

Compared with the natural groundwater, the human-influenced groundwater

showed remarkably increased (and dominant) dispersal limitation, decreased

homogeneous selection and drift, and stronger microbial co-occurrence patterns. First,

the river water input carries abundant nutrients from aquaculture and human activities,

which can create additional niches to stimulate the growth of various microorganisms,

especially less abundant ones. Such growth could increase species diversity (Fig. 1B)

and enhance stochastic processes like birth and death, the latter of which is consistent

with the enhanced dispersal limitation (the influence of dispersal limitation acting in

concert with drift [23]) and drift (the influence of drift acting alone) in the

human-influenced groundwater, as noticed in prior studies on terrestrial freshwater

[63] and groundwater [13]. Second, nutrient inputs could weaken environmental

selection by providing more resources and result in less-stressed groundwater

conditions, partly resulting in decreased homogeneous selection. Nutrient-associated

factors dominated selection process in the human-influenced groundwater, suggested

by the relationships between homogeneous selection and DIC or DOC (Fig. 5).

Higher abundance of nitrogen-metabolizing genera, e.g., nitrifying *Nitrosopumilus*

[64] and denitrifying *Hydrogenophaga* [65, 66], and carbon-metabolizing genera, e.g.,

Bathyarchaeia [67], and *Sphaerochaeta* [68] in the human-influenced than the natural

groundwater (Fig. S14) further supported the important effect of nutrient input. Third,

exogenous water inputs at ebb tide provided supplement water and therefore

weakened tidal fluctuations (i.e., weakened groundwater discharge during ebb tide),

leading to enhanced dispersal limitation.

Further, our finding on the important role of stochastic processes (dispersal

limitation in concert with drift) in controlling the human-influenced groundwater

microbiota is consistent with the prior studies indicating that stochasticity decreased,

whereas environmental selection could be stronger in a fluidic ecosystem with a

disturbance, in turn, stochasticity increases after adding exogenous nutrients [13].

Moreover, the combined contribution of dispersal and homogeneous selection have

also shaped DDR patterns of intertidal groundwater microbiota. However, it seems

that nutrient-driven selection processes dampened the effect of dispersal limitation,

resulting in decreased bacterial community dissimilarities across geographic distance

in the human-influenced groundwater compared with the natural groundwater.

Interestingly, unlike bacterial communities, archaeal communities showed little

change in DDR patterns in the human-influenced groundwater, potentially due to a

narrow niche breadth of subsurface archaea and thus less influenced by anthropogenic

activities [51].

Despite its lower importance of homogeneous selection (environmental filtering),

the human-influenced groundwater microbiota showed both a higher degree of species

segregation and higher frequencies of species co-occurrence associations, indicating

stronger microbial interactions. On the one hand, enhanced dispersal limitation and

potentially stronger competition (increased species diversity with similar ecological

niches) enhanced the species segregation distribution. On the other hand, complex

physicochemical gradients (e.g., abundant nutrients, salinity and DOC) [69] created

multiple niches that harbor various microbial groups with different metabolic

capacities (Fig. S11) and in turn created opportunities for synergistic or other

cooperative interactions (mutualistic and commensal metabolism cooperation), which

was supported by enrichment of copiotrophic genera *Woesearchaeales* [70] and

Flavobacterium [71] and more in the human-influenced groundwater (Fig. S14).

Meanwhile, more keystone 'connectors' enhanced the stability of the community

under higher level of microbial cooperation (reduce stability) [72] in the

human-influenced groundwater. Overall, the enhanced microbial interaction is one of

the essential ecological processes that cannot be ignored when exploring the

microbiota assembly in the human-influenced intertidal groundwater.

5. Conclusions

By overcoming obstacles of time-series sampling in the salt marsh, this study first

unravels microbiota rhythm patterns in response to periodic tidal fluctuations in the

intertidal groundwater-surface water continuum where microbiota assembly was

found to be dominantly driven by homogeneous selection. Discharge of

anthropogenically polluted river water into the tidal creek is demonstrated to

remarkably alter the microbiota biodiversity, structure and assembly patterns in the

intertidal groundwater via bidirectional water exchange. Our first combined use of

rhythm analysis and random forest models suggested the potential to enhance

monitoring of coastal water quality and pollution sources by detecting rhythmic

microbial variances that signify anthropogenic contamination from intertidal

groundwater discharge and/or seawater intrusion. This study also reveals how the

anthropogenically polluted river input shift intertidal groundwater microbiota via water exchange, which in turn may affect the microbially-driven biogeochemical cycling in coastal aquifers. Altogether, this study can provide valuable insights into the bio-monitoring and conservation of coastal zone ecosystems.

Data Availability

The raw sequencing data generated in this study have been deposited in the China National GeneBank DataBase (CNGBdb) with accession number CNP0003642.

Acknowledgment

This work was supported by the Zhejiang Provincial Natural Science Foundation of China under Grant No. LR22D010001 and the HRHI program 202309010 of Westlake Laboratory of Life Sciences and Biomedicine. We acknowledge the Westlake University HPC Center for computational support. Thank Mr. Duofei Hu and Ms. Shengjie Hu for assistance in field sampling, and Ms. Yisong Xu for laboratory management supports.

Author Contributions

All authors contributed intellectual input and assistance to this study. F.J. obtained funding and supervised the project. The original concepts were conceived by F.J. and Z.Z. with advice from L.L. Field sampling was carried out by Z.Z. and X.C. Data analysis were done by Z.Z. with assistance from G.Z. The manuscript was prepared by Z.Z. and F.J. and revised by L.L., L.Z. and H.G.

Competing Interests

The authors declare no competing interests.

References

1.   Heckbert S, Costanza R, Poloczanska ES, Richardson AJ. Climate Regulation as a Service from
Estuarine and Coastal Ecosystems. In: Wolanski E, McLusky D, editors. Treatise on Estuarine and
Coastal Science. Waltham: Academic Press; 2011. p. 199-216.

2.   Xin P, Wilson A, Shen CJ, Ge ZM, Moffett KB, Santos IR, et al. Surface Water and Groundwater
Interactions in Salt Marshes and Their Impact on Plant Ecology and Coastal Biogeochemistry. Rev
Geophys. 2022;60(1).

3.   Santoro AE. Microbial nitrogen cycling at the saltwater-freshwater interface. Hydrogeol J.
2010;18(1):187-202.

4.   Santos IR, Chen XG, Lecher AL, Sawyer AH, Moosdorf N, Rodellas V, et al. Submarine
groundwater discharge impacts on coastal nutrient biogeochemistry. Nature Reviews Earth &
Environment. 2021;2(5):307-23.

5.   Cho HM, Kim G, Kwon EY, Moosdorf N, Garcia-Orellana J, Santos IR. Radium tracing nutrient
inputs through submarine groundwater discharge in the global ocean. Sci Rep. 2018;8(1):2439.

6.   Ruiz-Gonzalez C, Rodellas V, Garcia-Orellana J. The microbial dimension of submarine
groundwater discharge: current challenges and future directions. FEMS Microbiol Rev. 2021;45(5).

7.   Hong Y, Wu J, Wilson S, Song B. Vertical Stratification of Sediment Microbial Communities
Along Geochemical Gradients of a Subterranean Estuary Located at the Gloucester Beach of Virginia,
United States. Front Microbiol. 2018;9:3343.

8.   Adyasari D, Hassenruck C, Oehler T, Sabdaningsih A, Moosdorf N. Microbial community
structure associated with submarine groundwater discharge in northern Java (Indonesia). Sci Total
Environ. 2019;689:590-601.

9.   Sang SL, Dai H, Hu BX, Hao YR, Zhou T, Zhang J. The Study of Hydrogeochemical
Environments and Microbial Communities along a Groundwater Salinity Gradient in the Pearl River
Delta, China. Water-Sui. 2019;11(4).

10.   Chen L, Zhang J, Dai H, Hu BX, Tong J, Gui D, et al. Comparison of the groundwater microbial
community in a salt-freshwater mixing zone during the dry and wet seasons. J Environ Manage.
2020;271:110969.

11.   Chen X, Ye Q, Sanders CJ, Du J, Zhang J. Bacterial-derived nutrient and carbon source-sink
behaviors in a sandy beach subterranean estuary. Mar Pollut Bull. 2020;160:111570.

12.   Chen YJ, Leung PM, Cook PLM, Wong WW, Hutchinson T, Eate V, et al. Hydrodynamic
disturbance controls microbial community assembly and biogeochemical processes in coastal
sediments. Isme J. 2022;16(3):750-63.

13.   Zhou J, Deng Y, Zhang P, Xue K, Liang Y, Van Nostrand JD, et al. Stochasticity, succession, and
environmental perturbations in a fluidic ecosystem. Proc Natl Acad Sci U S A. 2014;111(9):E836-45.

14.   Hanley TC, Bowen JL, Kearns PJ, Hughes AR. Short- and long-term effects of nutrient
enrichment on salt marsh plant production and microbial community structure. Journal of Ecology.
2021;109(11):3779-93.

15.   Yao Z, Du S, Liang C, Zhao Y, Dini-Andreote F, Wang K, et al. Bacterial Community Assembly
in a Typical Estuarine Marsh with Multiple Environmental Gradients. Appl Environ Microbiol.
2019;85(6).

16.   Adyasari D, Hassenruck C, Montiel D, Dimova N. Microbial community composition across a

coastal hydrological system affected by submarine groundwater discharge (SGD). Plos One.

2020;15(6):e0235235.

17.   Zhang G, Bai J, Tebbe CC, Zhao Q, Jia J, Wang W, et al. Salinity controls soil microbial

community structure and function in coastal estuarine wetlands. Environ Microbiol.

2021;23(2):1020-37.

18.   Chave J. Neutral theory and community ecology. Ecology Letters. 2004;7(3):241-53.

19.   Vellend M. Conceptual synthesis in community ecology. Q Rev Biol. 2010;85(2):183-206.

20.   Nemergut DR, Schmidt SK, Fukami T, O'Neill SP, Bilinski TM, Stanish LF, et al. Patterns and

processes of microbial community assembly. Microbiol Mol Biol Rev. 2013;77(3):342-56.

21.   Zhou J, Ning D. Stochastic Community Assembly: Does It Matter in Microbial Ecology?

Microbiol Mol Biol Rev. 2017;81(4).

22.   Ning D, Yuan M, Wu L, Zhang Y, Guo X, Zhou X, et al. A quantitative framework reveals

ecological drivers of grassland microbial community assembly in response to warming. Nat Commun.

2020;11(1):4717.

23.   Stegen JC, Lin X, Fredrickson JK, Chen X, Kennedy DW, Murray CJ, et al. Quantifying

community assembly processes and identifying features that impose them. Isme J. 2013;7(11):2069-79.

24.   Gao GF, Peng D, Zhang Y, Li Y, Fan K, Tripathi BM, et al. Dramatic change of bacterial assembly

process and co-occurrence pattern in Spartina alterniflora salt marsh along an inundation frequency

gradient. Sci Total Environ. 2021;755(Pt 1):142546.

25.   Gao GF, Peng D, Tripathi BM, Zhang Y, Chu H. Distinct Community Assembly Processes of

Abundant and Rare Soil Bacteria in Coastal Wetlands along an Inundation Gradient. mSystems.

2020;5(6).

26.   Wang K, Yan H, Peng X, Hu H, Zhang H, Hou D, et al. Community assembly of bacteria and

archaea in coastal waters governed by contrasting mechanisms: A seasonal perspective. Mol Ecol.

2020;29(19):3762-76.

27.   Kraft NJB, Adler PB, Godoy O, James EC, Fuller S, Levine JM. Community assembly,

coexistence and the environmental filtering metaphor. Funct Ecol. 2015;29(5):592-9.

28.   Faust K, Raes J. Microbial interactions: from networks to models. Nat Rev Microbiol.

2012;10(8):538-50.

29.   Berry D, Widder S. Deciphering microbial interactions and detecting keystone species with

co-occurrence networks. Front Microbiol. 2014;5:219.

30.   Yuan MM, Guo X, Wu LW, Zhang Y, Xiao NJ, Ning DL, et al. Climate warming enhances

microbial network complexity and stability. Nature Climate Change. 2021;11(4):343-U100.

31.   Ju F, Xia Y, Guo F, Wang Z, Zhang T. Taxonomic relatedness shapes bacterial assembly in

activated sludge of globally distributed wastewater treatment plants. Environ Microbiol.

2014;16(8):2421-32.

32.   Ju F, Zhang T. Bacterial assembly and temporal dynamics in activated sludge of a full-scale

municipal wastewater treatment plant. Isme J. 2015;9(3):683-95.

33.   Niederdorfer R, Fragner L, Yuan L, Hausherr D, Wei J, Magyar P, et al. Distinct growth stages

controlled by the interplay of deterministic and stochastic processes in functional anammox biofilms. Water Res. 2021;200:117225.

34.   Shen Y, Feng N-h, Zhou Q, Liu Y-m, Chen Z-y. The status and its influence of reclamation on Jiangsu coast. Marine Sciences. 2006;10:008.

35.   Chen XG, Santos IR, Hu DF, Zhan LC, Zhang Y, Zhao Z, et al. Pore-water exchange flushes blue carbon from intertidal saltmarsh sediments into the sea. Limnol Oceanogr Lett. 2022;7(4):312-20.

36.   Fadeev E, Cardozo-Mino MG, Rapp JZ, Bienhold C, Salter I, Salman-Carvalho V, et al. Comparison of Two 16S rRNA Primers (V3-V4 and V4-V5) for Studies of Arctic Microbial Communities. Frontiers in Microbiology. 2021;12.

37.   Bolyen E, Rideout JR. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nature Biotechnology. 2019;37(8):852-7.

38.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 2014;30(15):2114-20.

39.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods. 2016;13(7):581-3.

40.   McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. Plos One. 2013;8(4):e61217.

41.   Liu C, Cui Y, Li X, Yao M. microeco: an R package for data mining in microbial community ecology. FEMS Microbiol Ecol. 2021;97(2):fiaa255.

42.   Lambert S, Tragin M, Lozano JC, Ghiglione JF, Vaulot D, Bouget FY, et al. Rhythmicity of coastal marine picoeukaryotes, bacteria and archaea despite irregular environmental perturbations. Isme J. 2019;13(2):388-401.

43.   Jiao S, Yang Y, Xu Y, Zhang J, Lu Y. Balance between community assembly processes mediates species coexistence in agricultural soil microbiomes across eastern China. Isme J. 2020;14(1):202-16.

44.   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. Nucleic Acids Res. 2021;49(W1):W293-W6.

45.   Hu A, Ju F, Hou L, Li J, Yang X, Wang H, et al. Strong impact of anthropogenic contamination on the co-occurrence patterns of a riverine microbial community. Environ Microbiol. 2017;19(12):4993-5009.

46.   Deng Y, Jiang YH, Yang Y, He Z, Luo F, Zhou J. Molecular ecological network analyses. BMC Bioinformatics. 2012;13:113.

47.   Zhou J, Deng Y, Luo F, He Z, Tu Q, Zhi X. Functional molecular ecological networks. Mbio. 2010;1(4).

48.   Eglen RM, Whiting RL. Competitive and non-competitive antagonism exhibited by 'selective' antagonists at atrial and ileal muscarinic receptor subtypes. Br J Pharmacol. 1987;90(4):701-7.

49.   Lee E, Shin D, Hyun SP, Ko KS, Moon HS, Koh DC, et al. Periodic change in coastal microbial community structure associated with submarine groundwater discharge and tidal fluctuation. Limnol Oceanogr. 2017;62(2):437-51.

50.   Stal LJ. Microphytobenthos as a biogeomorphological force in intertidal sediment stabilization. Ecol Eng. 2010;36(2):236-45.

51.   Chen W, Wen D. Archaeal and bacterial communities assembly and co-occurrence networks in
subtropical mangrove sediments under Spartina alterniflora invasion. Environ Microbiome.
2021;16(1):10.

52.   Fuhrman JA, Cram JA, Needham DM. Marine microbial community dynamics and their
ecological interpretation. Nat Rev Microbiol. 2015;13(3):133-46.

53.   Auladell A, Barberan A, Logares R, Garces E, Gasol JM, Ferrera I. Seasonal niche differentiation
among closely related marine bacteria. Isme J. 2022;16(1):178-89.

54.   Blanton JO, Lin GQ, Elston SA. Tidal current asymmetry in shallow estuaries and tidal creeks.
Cont Shelf Res. 2002;22(11-13):1731-43.

55.   Werner AD, Bakker M, Post VEA, Vandenbohede A, Lu CH, Ataie-Ashtiani B, et al. Seawater
intrusion processes, investigation and management: Recent advances and future challenges. Advances
in Water Resources. 2013;51:3-26.

56.   Zhong Q, Wang X, Wang Q, Zhang F, Li L, Wang Y, et al. (222)Rn, (210)Pb and (210)Po in
coastal zone groundwater: Activities, geochemical behaviors, consideration of seawater intrusion effect,
and the potential radiation human-health risk. Appl Radiat Isot. 2020;166:109386.

57.   Nilavan E, Vaiyapuri M, Sheela GS, Nadella RK, Thandapani M, Kumar A, et al. Prevalence of
Vibrio mimicus in Fish, Fishery Products, and Environment of South West Coast of Kerala, India. J.
AOAC Int. 2021;104:790–794.

58.   Ma Z, Gao L, Sun M, Liao Y, Bai S, Wu Z, et al. Microbial Diversity in Groundwater and Its
Response to Seawater Intrusion in Beihai City, Southern China. Front. Microbiol. 2022;13, 876665.

59.   Martiny JB, Eisen JA, Penn K, Allison SD, Horner-Devine MC. Drivers of bacterial beta-diversity
depend on spatial scale. Proc Natl Acad Sci U S A. 2011;108(19):7850-4.

60.   Custer GF, Bresciani L, Dini-Andreote F. Ecological and Evolutionary Implications of Microbial
Dispersal. Front Microbiol. 2022;13:855859.

61.   McCall BD, Pennings SC. Geographic variation in salt marsh structure and function. Oecologia.
2012;170(3):777-87.

62.   Li H, Chi Z, Li J, Wu H, Yan B. Bacterial community structure and function in soils from tidal
freshwater wetlands in a Chinese delta: Potential impacts of salinity and nutrient. Sci Total Environ.
2019;696:134029.

63.   Ren L, He D, Chen Z, Jeppesen E, Lauridsen TL, Sondergaard M, et al. Warming and nutrient
enrichment in combination increase stochasticity and beta diversity of bacterioplankton assemblages
across freshwater mesocosms. Isme J. 2017;11(3):613-25.

64.   Walker CB, de la Torre JR, Klotz MG, Urakawa H, Pinel N, Arp DJ, et al. Nitrosopumilus
maritimus genome reveals unique mechanisms for nitrification and autotrophy in globally distributed
marine crenarchaea. Proc Natl Acad Sci U S A. 2010;107(19):8818-23.

65.   Willems A, Gillis M. Hydrogenophaga. Bergey's Manual of Systematics of Archaea and Bacteria.
2015:1-15.

66.   Willems A, Busse J, Goor M, Pot B, Falsen E, Jantzen E, et al. Hydrogenophaga, a New Genus of
Hydrogen-Oxidizing Bacteria That Includes Hydrogenophaga flava comb. nov. (Formerly
Pseudomonas flava), Hydrogenophaga palleronii (Formerly Pseudomonas palleronii), Hydrogenophaga

pseudoflava (Formerly Pseudomonas pseudoflava and "Pseudomonas carboxydoflava"), and

Hydrogenophaga taeniospiralis (Formerly Pseudomonas taeniospiralis). Int J Syst Bacteriol.

1989;39(3):319-33.

67.   Romano RG, Bendia AG, Moreira JCF, Franco DC, Signori CN, Yu TT, et al. Bathyarchaeia

occurrence in rich methane sediments from a Brazilian ria. Estuar Coast Shelf S. 2021;263.

68.   Oshurkova V, Troshina O, Trubitsyn V, Ryzhmanova Y, Bochkareva O, Shcherbakova V.

Characterization of Methanosarcina mazei JL01 Isolated from Holocene Arctic Permafrost and Study

of the Archaeon Cooperation with Bacterium Sphaerochaeta associata GLS2T. Proceedings.

2020;66(1):4.

69.   Veach AM, Stegen JC, Brown SP, Dodds WK, Jumpponen A. Spatial and successional dynamics

of microbial biofilm communities in a grassland stream ecosystem. Mol Ecol. 2016;25(18):4674-88.

70.   Liu X, Li M, Castelle CJ, Probst AJ, Zhou Z, Pan J, et al. Insights into the ecology, evolution, and

metabolism of the widespread Woesearchaeotal lineages. Microbiome. 2018;6(1):102.

71.   Ho A, Di Lonardo DP, Bodelier PLE. Revisiting life strategy concepts in environmental microbial

ecology. FEMS Microbiology Ecology. 2017;93(3).

72.   Ya T, Du S, Li ZY, Liu SD, Zhu MH, Liu XJ, et al. Successional Dynamics of Molecular

Ecological Network of Anammox Microbial Communities under Elevated Salinity. Water Research.

2021;188.

Figure 1. Sampling and overview of archaeal and bacterial communities in tidal creek groundwater and surface water. (A) Map of human-influenced (HT) and natural (NT) salt marsh tidal creeks sampled in this study. Red dots in the graph above show the sampling sites of intertidal groundwater and red triangle marks the times series station in NT for collecting surface water samples (NTS). Blue arrows show the water flow direction of tidal creek and Chuandong River, as well as the Terrestrial River water input upstream of HT. Groundwater samples from NT were collected at low (L), mid (M), high (H) and above high (A) tidal marks along three sampling transect of NT, whereas groundwater samples from HT were all collected at mid (M) tidal marks, as shown in the bottom-left graph. The blue and pink rectangles in the bottom-right graph mark the time periods for collecting time-series flood and ebb tide surface water, respectively, during neap tide and spring tide. (B) Alpha diversity shown as ACE and Shannon indices. The significance level in the median differences across groups is statistically checked using wilcoxon rank-sum tests. Significance is expressed as ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, NS-non significant. (C) The composition of all phylum-level archaeal communities. (D) The composition of phylum-level bacterial communities. Top 16 most abundant phylum are shown and other phylum are classified as 'Others'.

Figure 2. The structural variations of microbial communities within and between tidal creek groundwater and surface water. The non-multidimensional scaling (NMDS) ordination based on weighted UniFrac distance is employed to characterize the compositional variations of archaeal (A) and bacterial (B) communities in HTG, NTG, NTS-Ebb/Flood (left graph) or NTS-Neap/Spring (right graph). The archaeal and bacterial community distance across groups based on weighted UniFrac distance is displayed (right panel) and statistically significant differences across groups are calculated based on wilcoxon rank-sum test (****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, ns-non significant).

Figure 3. Rhythmicity of microbiota in tidal creek surface water. (A) The NMDS ordination of prokaryotic (archaeal and bacterial) communities in tidal creek surface water is displayed and trajectories show sampling order of surface water during neap tide (N1-N14 with brown lines) and spring tide (S1-S15 with gray lines). The four ellipses mark surface water samples collected during the neap-ebb, neap-flood, spring-ebb and spring-flood tide, respectively. (B) Pairwise comparisons of environmental factors are shown with a color gradient denoting Pearson's correlation. Microbiota composition in NTS-Neap, NTS-Spring and NTS groups was related to each factor by partial mantel tests. Edge width corresponds to mantel's r statistic for the corresponding distance correlations and edge color denotes the statistical significance based on 9,999 permutations. (C) Cluster heatmap plot of centered logarithm ratio abundances of 271 rhythmic ASVs identified in the NTS-Neap group. (D) Cluster heatmap plot of 128 rhythmic ASVs identified in the NTS-Spring group. The black rectangles (cluster) mark the ASVs showing higher abundance during ebb tide or flood tide and the number of ASVs classified to each Family for ebb cluster and flood cluster was annotated in the graph.

Figure 4. Community assembly processes across different phylogenetic groups. (A) Relative importance of five ecological processes across groups, including heterogeneous selection (HeS), homogeneous selection (HoS), dispersal limitation (DL), homogenizing dispersal (HD), and drift (DR). (B) Phylogenetic tree of top member lineage in each Bin was displayed at class-level. Relative abundance of each Bin was displayed in first rectangle. Dominant processes in each Bin (colored rectangle) and Bin contribution (stacked bars) to major ecological processes (HoS, DL and DR) in HTG, NTG, NTS-Neap and NTS-Spring were displayed in 2-5 annulus. Top Bins with higher contribution to ecological processes were marked by gray strip. Only the top300 Bins were shown in this figure, accounting for a relative abundance of 96.1% of all Bins. (C) Bin-class (top15 and others) contributions to the major ecological processes (HoS, DL, and DR) were displayed in the left heatmap. Sum of the relative abundance of all Bins in each Bin-class was displayed in right bar chart. The vertical numbers at the right of the heatmap showed the number of Bins classified at class-level.

Figure 5. Effects of environmental factors on major ecological processes. Heatmap plot of correlations between environmental factors and dominant ecological processes based on mantel test. The correlation was determined based on the difference (with a 'd' before the factors) or the mean (with a 'm' before the factors) of a factor between each pair of samples. Significance was expressed as **$p < 0.01$, *$p < 0.05$.

Red edges: significant positive correlations. Green edges: significant negative correlations

Figure 6. Microbial segregation and co-occurrence patterns of tidal creek water microbiota. (A) Standardized effect sizes (SES) values of C-score and C_{var}-score for intertidal groundwater and surface water microbiota over six groups. Higher observed SES values of C-score suggest greater degrees of species segregation than would be expected by chance. Higher SES values of C_{var}-score indicate greater degrees of both species segregation and aggregation. All *P*-values are < 0.0001 (detailed in Text S6 and Table S14). (B) Microbial co-occurrence networks in groundwater and surface water groups. n: number of nodes; e: number of edges; m: number of modules. Nodes in each network are colored by modularity. Red edges represent significant positive correlations and green edges represent significant negative correlations. More details are described in Text S7 and Table S15.