Mathematical characterization of fractal river networks

Author: Akira Terui[1,*]

Email: hanabi0111@gmail.com

[1]Department of Biology, University of North Carolina at Greensboro, Greensboro, NC 27410, USA

Abstract

Carraro and Altermatt[1] performed an extensive analysis to study scaling properties of river networks. They concluded that: (1) branching "probability" is better termed as branching "ratio" for real river networks because it is not a probability; (2) branching ratio is a scale-dependent quantity as the value changes across spatial resolutions at which river networks are extracted (expressed as the threshold catchment area A_T that initiates channels or pixel size l); (3) Optimal Channel Networks most accurately predicted the metapopulation stability and capacity. However, the supporting ground for these conclusions is either seriously flawed or inconclusive due to the improper use of branching probability, scale invariance, dimensions, and units.

Maintext

Rivers form complex branching networks, and the ecological implication of river network complexity has gained great interest over the past few decades[2–4]. To this end, there have been concerted efforts to construct virtual river networks to provide theoretical insights into how river network structure controls riverine ecological dynamics. Carraro and Altermatt[1] compared scaling properties and ecological dynamics in virtual river networks produced by three different simulation methods – balanced binary trees (BBTs)[5], random branching networks (RBNs)[6,7], and optimal channel networks (OCNs)[8]. In addition, the authors reported power-law scaling of a structural characteristic (the number of links per unit distance, termed "branching ratio" in their article) in real rivers. Carraro and Altermatt[1] concluded that: (1) branching "probability" is better termed as branching "ratio" for real river networks because it is not a probability; (2) branching ratio is a scale-dependent quantity as the value changes across spatial resolutions at which river networks are extracted (expressed as the threshold catchment area A_T that initiates channels or pixel size l); (3) OCNs most accurately predicted the metapopulation stability and capacity. However, the supporting ground for these conclusions is either seriously flawed or inconclusive due to the improper use of branching probability, scale invariance, dimensions, and units.

First, branching probability p is certainly a probability by definition. In previous studies, branching probability p has been defined as a cumulative probability distribution of link length L [km] in real river networks (a "*link*" represents a river segment from one confluence to another). For example, Terui et al.[6,7] fitted an exponential distribution to link length as $L_j \sim \mathrm{Exp}(\theta)$ (θ is the rate parameter and j refers to an individual link) and estimated branching probability as $p = 1 - \exp(-\theta L)$ (L was set to be 1 [km] in Terui et al.[6,7]). Thus, p [km^{-1}] in this example is the *probability* of including a confluence (or an upstream terminal) per unit river km. Therefore, the author's claim"...*it is in fact improper to refer to a probability when analyzing the properties of a realized river network*..." is simply a misunderstanding. Instead, branching ratio p_r is

identical to the rate parameter θ [km^{-1}], which is the inverse of average link length $\frac{N_L}{N}$ (N_L the number of links in a network [-]; N the total river length [km]). To show this equality ($\theta = p_r$), I delineated 50 river networks analyzed in Carraro and Altermatt[1] using MERIT Hydro[9], with which θ and p_r are independently estimated as described above. MERIT Hydro has constant pixel size of 3 arc-second (~90 m at the equator) across the globe. I used $A_T = 1$ [km^2] to capture sufficient details of real river networks. The relationship between p_r and θ fell exactly on a 1:1 line (Figure 1A), thus $\theta = p_r$. By definition, branching probability is a monotonic increasing function of p_r with a possible range of $0 - 1$ (Figure 1B); yet, they are not identical. Note that, in the following paragraphs, I use p_r for my analysis to be consistent with the original article. Since branching probability p is a simple transformation of p_r, the arguments are translatable between the two.

Figure 1: (A) Equality between rate parameter θ and branching ratio p_r. Dots indicate estimated values at 50 rivers analyzed in Carraro and Altermatt[1]. The gray line denotes a 1:1 relationship. (B) Relationship between branching probability ($p = 1 - \exp(-\theta)$) and ratio ($p_r = \frac{N_L}{N}$). Dots are estimated values of p and p_r at the 50 rivers. Note that p and p_r were estimated independently; nevertheless, the data points fell exactly on the theoretical relationship between p and θ (the gray line), confirming the equality between θ and p_r. Inlet zooms into the observed range of p_r, which is denoted by the vertical broken lines.

Second, the term "scale invariance" was falsely used in their article. To explain this, let me consider an object y whose structural property is assessed under observation scale x. For example, Mandelbrot[10] studied the length of a coastline as multiples of a ruler with a unit length x (the "observation scale"). In this practice, the observed object is expressed as a function of scale x ($y = f(x)$), and the function $f(x)$ is said to be scale invariant if it suffices the following equation[11,12]:

$$f(\lambda x) = \lambda^z f(x)$$

where λ is an arbitrary constant and z is the scaling exponent. The above equation is interpreted as the sign of "scale invariance" because the multiplicative extension/shrink of observation scale x by factor λ results in the same shape of the original object y but with a *different scale*[12] (i.e., the structural property is retained). The power-law function is the most famous example that suffices this definition of scale invariance[11,13]:

$$y = f(x) = cx^z$$

where c is the scaling constant. The scale invariance can be easily proved by multiplying the scaling factor λ:

$$y' = f(\lambda x) = c\lambda^z x^z = \lambda^z f(x) = \lambda^z y$$

Carraro and Altermatt[1] provided evidence that branching ratio p_r follows a power law along the axes of observation scale A_T and pixel size l (i.e., length on a side) using OCNs (Equation 1 in the original article). Also, the authors visually showed that the relationship between p_r and A_T follows a power-law using the data from 50 real river networks (Figure 3a in the original article). Nevertheless, the authors claimed that "*Here we show that an alleged property of such random networks (branching probability) is a scale-dependent quantity that does not reflect any recognized metric of rivers' fractal character…*" (Abstract) This interpretation is the complete opposite as it has been defined in the literature of scale-invariance[11–14], including the author's previous publication[15]. Instead, by definition, the conclusion derived from their analysis must be "*branching ratio is a scale invariant feature that reflects the fractal nature of rivers.*"

Third, dimensions and units are improperly treated in their article. A dimension is the power of an axis along which a physical quantity is measured, and a unit is a way to assign a number to a particular dimension to make it relative. For a simple straight line, length is a dimension (dimension = 1), and a meter is a unit of length. Throughout the article, the authors used the number of pixels to measure the total river length N, the total catchment area A, and the threshold catchment area A_T. There is no issue with using the number of pixels as a unit. However, a problem in their article is that they obscured the dimensions of pixels. For example, they made it very unclear that the river length N has a dimension of pixel "length," while the total catchment area A and the threshold catchment area A_T have a dimension of pixel "area." In particular, the authors incorrectly defined p_r as a dimensionless quantity (Methods) even though its unit is [pixel length^{-1}].

The improper use of units caused more serious problems in the analysis of "real" river networks; the authors used different pixel sizes (length on a side) among watersheds (103 m to 1268 m; Supplementary Table 2 in the original article), meaning that the same number of pixels translates into very different lengths and areas. For example, in Figure 3 in the original article, the observation scale A_T ranges $20 - 500$ pixels. This pixel range translates into $0.2 - 5.3$ km^2 for the Toss river (with smallest pixel size) and $32.2 - 803.9$ km^2 for the Stikine river (with largest pixel size). Further, the branching ratio p_r is also affected by this variation in pixel size as its unit is [pixel length^{-1}]. Once the pixel length is converted to a unit of km, the branching ratio represents the number of links per 0.1 km in Toss, whereas it represents the number of links per 1.3 km in Stikine. Hence, the authors compared incomparable values, and their results reported in their article are not reliable.

To explore the consequences of the improper use of pixels, I re-analyzed their data with MERIT Hydro[9]; as noted above, this layer has constant pixel size, unlike the author's analysis. I extracted river networks with 20 values of A_T [km^2] ($A_T = 1, ..., 1000$ with an equal interval at a $\log_{10}$ scale, but confined to $A_T < A$ for small rivers), at which I estimated branching ratio as $p_r = \frac{N_L}{N}$ [km^{-1}]. In the original article, the authors did not perform any statistical analysis even though $p_r (= \theta)$ is a statistical parameter characterizing a link length distribution (see above); thus, its estimation accuracy is affected by the number of links, i.e., the sample size. To statistically substantiate the power-law of p_r along the axis of A_T, one must account for this heterogeneity of estimation accuracy. To this end, I fitted the following log-linear models with robust regression (i represents an individual data point of p_r estimated in a given watershed at a given scale A_T).

$$\log_{10} p_{r,i} = \log_{10} c_{w(i)} + z \log_{10} A_{T,i} + \varepsilon_i \qquad \text{(M0)}$$
$$\log_{10} p_{r,i} = \log_{10} c_{w(i)} + z_{w(i)} \log_{10} A_{T,i} + \varepsilon_i \quad \text{(M1)}$$

where ε_i is the error term that is properly weighted by Huber's function. Robust regression analysis is appropriate because it is robust to outliers caused by the small sample size (typical for large A_T values). The first model (M0) assumes the "universal" scaling with the single exponent z across watersheds; i.e., the branching ratio at all the 50 watersheds follows the same power law with the watershed-specific constant $\log_{10} c_{w(i)}$ ($w(i)$ is watershed w for a data point i). In contrast, the second model (M1) assumes the "localized" scaling with the watershed-specific exponent $z_{w(i)}$. I estimated the evidence ratio of the two models using the approximated Bayes Factor (BF)[16], which is defined as $\text{BF} = \exp\left(\frac{\text{BIC(M1)} - \text{BIC(M0)}}{2}\right)$. In this definition, a value of BF > 1 gives the support for M0 over M1; for example, if BF $= 2$, the model M0 is twice as likely as the model M1.

The analysis provided decisive support for M0 with BF $\approx 3.00 \times 10^{20}$. Under M0, as evident from its model formula, the rank of the expected branching ratio $\text{E}(\log_{10} p_r)$ never changed across scales, maintaining the order of the watershed-specific constant $\log_{10} c_{w(i)}$ (Figure 2; see regression lines). This result is inconsistent with the author's statement "*by extracting different river networks at various scales (i.e., various A_T values) and assessing the rivers' rank in terms of p_r, one observes that rivers that look more "branching" (i.e., have higher p_r) than others for a given A_T value can become less "branching" for a different A_T value (Fig. 3).*" I also must note that I did not find any significant correlation between watershed area A [km^2] and branching ratio p_r [km^{-1}] when extracted with a constant value of $A_T = 1$ km^2 across watersheds (Spearman's rank correlation $= -0.19$, p-value $= 0.19$), as opposed to the statement in the original article "*…if different river networks spanning different catchment areas (say, in km^2) are compared, all of them extracted from the same DEM (same l and same A_T in km^2), then the larger river network will appear more branching (i.e., have larger p_r)*[1]." The lack of correlation between branching probability p and watershed area A has also been reported in two previous studies[6,7], one of which used 184 watersheds for the analysis[7]. This result is not surprising at all because p_r is the number of links "divided" by the total river length by definition. My re-analysis revealed that the author's statements merely reflect the lack of appropriate quantitative analysis and/or a statistical artifact of inconsistent units across watersheds.

It is worthwhile to note that if scaling exponent varies by watershed (i.e., z_w; albeit not the case for my analysis), the dimensional p_r is incomparable across watersheds because its dimension ($= 2z_w$) is different (just like length and area are not comparable). To understand this argument, one must recognize that, by writing $\text{E}(p_r) = c_w A_T^{z_w}$, we equate the dimension in both sides of the equation. From this equation, the dimension of p_r is "estimated" to be $2z_w$ since the scaling exponent z_w applies to the unit of A_T as $\text{E}(p_r) = c_w A_T^{z_w}$ [km^{2z_w}]. Therefore, to compare branching ratio across watersheds, we must non-dimensionalize p_r as $\text{E}(\bar{p}_r) = \frac{\text{E}(p_r)}{A^{z_w}} = c_w$ [-] (see Figure 2B for this exercise). Evidently, the expected value of dimensionless branching ratio $\text{E}(\bar{p}_r)$ remains constant across scales (c_w) with no possibility of changes in the rank (Figure 2B). This technique of dimensional analysis has been widely used when comparing or non-dimensionalizing the self-similar structure of scale-invariant objects (see Figure 1 in Rinaldo et al.[14] and equation (2.2) in Rodriguez-Iturbe and Rinaldo[11] for examples). In fact, past empirical studies used $A_T = 1$ km^2 such that the estimated dimensional p_r approximates the rescaled $\bar{p}_r$ ($p_r = \bar{p}_r \approx c_w$ for $A_T = 1$ in a given unit, regardless of the value of z)[6,7]. Note that $\bar{p}_r$ fluctuates unpredictably at coarser observation scales (larger A_T relative to A; Figure 2B). This is simply because of the small sample size (i.e., the number of links) at coarser resolutions, which inflates the statistical uncertainty of parameter estimation for p_r. However, a solution is simple: use a sufficiently small observation scale (e.g., $A_T = 1$ km^2) to yield a large sample size N_L. The $\bar{p}_r$ will converge to the single value that represents the self-similar structure unique to each

137 river network (Figure 2B). Therefore, there is no mathematical supporting ground for the author's statement
138 "...*branching probability is a non-descriptive property of a river network, which by no means describes its*
139 *inherent branching character*..."

Figure 2: Log-log plot substantiates the power-law scaling of dimensional branching ratio (p_r; Panel A) along the axis of observation scale A_T. Once non-dimensionalized, the properly rescaled branching ratio ($\bar{p}_r$; Panel B) will converge to values unique to each river as $A_T \rightarrow 0$. Colors indicate rivers highlighted in the original article[1]. Individual data points are shown in dots, whose transparency is proportional to the number of links N_L (i.e., sample size). Lines are predicted values from the model M0 (i.e., the expected value of dimensional or rescaled branching ratio).

Lastly, the issue of units is pertinent to their metapopulation simulations, as they used pixel size l as a unit
of "population scale" and "dispersal distance." This makes their simulation results hardly interpretable. For
example, they assumed a population scale of 103 m in Toss, while assuming 1268 m in Stikine. A reasonable
interpretation of this simulation setup is that the authors simulated metapopulation dynamics of "different"
species among watersheds. This is not trivial as the population scale defines the number of local populations
within a metapopulation – one of the most influential parameters dictating metapopulation CV (see Equations
(2) and (3) in the original article). If one assumes the same population scale across watersheds (i.e., the same
species), then Stikine should be ~12 times greater in the number of local populations N_p compared to Toss.
Nevertheless, the authors used a nearly-constant number of N_p for all the 50 real rivers (e.g., $E(N_p) \approx 1088$
for $A_T = 500$ [pixel area]) by changing the population scale. Similarly, the average dispersal distance α was
measured in the unit of pixel length ($\alpha = 100$ [pixel length] in Figure 6 in the original article), meaning that it
varies from 10.3 km to 126.8 km among watersheds when converted to a unit of km. It is difficult to envision
that such a huge variation in dispersal capability exists within the same species. Again, the parameter α has
a decisive influence on metapopulation CV and capacity in the author's model (see Methods in the original
article). Critically, Figure 6 and Supplementary Figures 4 – 9 of the original article "aggregated" these
metapopulations of different species in real rivers and compared the overall summary statistics (e.g., median)
with those obtained from BBTs, RBNs, and OCNs. Therefore, the authors compared incomparable values.
A better alternative should have been to use the same pixel size (= population scale) among watersheds to
assume metapopulations of the "same" species, then pairing metapopulation metrics by watershed. In doing

so, the authors could examine how predicted values from different virtual networks correlate with values in real rivers with a reference to a 1:1 line. Although I do not know how this improvement will alter one of the main conclusions "*OCNs most accurately predicted the metapopulation stability and capacity*," it is clear that their simulation results are not ecologically interpretable.

Data availability

Codes and data are available at https://github.com/aterui/public-proj_fractal-river.

Competing interest

None declared.

Reference

1. Carraro, L. & Altermatt, F. Optimal Channel Networks accurately model ecologically-relevant geo-morphological features of branching river networks. *Communications Earth & Environment* **3**, 1–10 (2022).

2. Grant, E. H. C., Lowe, W. H. & Fagan, W. F. Living in the branches: Population dynamics and ecological processes in dendritic networks. *Ecology Letters* **10**, 165–175 (2007).

3. Carrara, F., Altermatt, F., Rodriguez-Iturbe, I. & Rinaldo, A. Dendritic connectivity controls biodiversity patterns in experimental metacommunities. *Proceedings of the National Academy of Sciences* **109**, 5761–5766 (2012).

4. Mari, L., Casagrandi, R., Bertuzzo, E., Rinaldo, A. & Gatto, M. Metapopulation persistence and species spread in river networks. *Ecology Letters* **17**, 426–34 (2014).

5. Yeakel, J. D., Moore, J. W., Guimarães, P. R. & de Aguiar, M. A. M. Synchronisation and stability in river metapopulation networks. *Ecology Letters* **17**, 273–283 (2014).

6. Terui, A. *et al.* Metapopulation stability in branching river networks. *Proceedings of the National Academy of Sciences* **115**, E5963–E5969 (2018).

7. Terui, A., Kim, S., Dolph, C. L., Kadoya, T. & Miyazaki, Y. Emergent dual scaling of riverine biodiversity. *Proceedings of the National Academy of Sciences* **118**, e2105574118 (2021).

8. Carraro, L. *et al.* Generation and application of river network analogues for use in ecology and evolution. *Ecology and Evolution* **10**, 7537–7550 (2020).

9. Yamazaki, D. *et al.* MERIT hydro: A high-resolution global hydrography map based on latest topography dataset. *Water Resources Research* **55**, 5053–5073 (2019).

10. Mandelbrot, B. How long is the coast of Britain? Statistical self-similarity and fractional dimension. *Science* **156**, 636–638 (1967).

11. Rodríguez-Iturbe, I. & Rinaldo, A. *Fractal River Basins: Chance and Self-Organization*. (Cambridge University Press, 2001).

12. Proekt, A., Banavar, J. R., Maritan, A. & Pfaff, D. W. Scale invariance in the dynamics of spontaneous behavior. *Proceedings of the National Academy of Sciences* **109**, 10564–10569 (2012).

13. Brown, J. H. *et al.* The fractal nature of nature: Power laws, ecological complexity and biodiversity. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* **357**, 619–626 (2002).

194   14.   Rinaldo, A., Rigon, R., Banavar, J. R., Maritan, A. & Rodriguez-Iturbe, I. Evolution and selection of river networks: Statics, dynamics, and complexity. *Proceedings of the National Academy of Sciences* **111**, 2417–2424 (2014).

196   15.   Giometto, A., Altermatt, F., Carrara, F., Maritan, A. & Rinaldo, A. Scaling body size fluctuations. *Proceedings of the National Academy of Sciences* **110**, 4646–4650 (2013).

198   16.   Wagenmakers, E.-J. A practical solution to the pervasive problems of p values. *Psychonomic Bulletin & Review* **14**, 779–804 (2007).