Signatures of kin selection in a natural population of the bacteria *Bacillus*

subtilis

Laurence J. Belcher[1*], Anna E. Dewar[1], Chunhui Hao[1], Melanie Ghoul[1a], Stuart A. West[1a]

[1] Department of Biology, University of Oxford, Oxford, OX1 3SZ, United Kingdom

[a] Joint last author

*Corresponding author

Email: laurence.belcher@biology.ox.ac.uk

Abstract

Laboratory experiments have suggested that bacteria perform a range of cooperative behaviours, which are favoured because they are directed towards relatives (kin selection). However, there is a lack of evidence for cooperation and kin selection in natural bacterial populations. Molecular population genetics offers a promising method to study natural populations, because theory predicts that kin selection will lead to relaxed selection, which will result in increased polymorphism and divergence at cooperative genes. Examining a natural population of *Bacillus subtilis*, we found consistent evidence that putatively cooperative traits have higher polymorphism and greater divergence than putatively private traits expressed at the same rate. In addition, we were able to eliminate alternative explanations for these patterns, and found more deleterious mutations in genes controlling putatively cooperative traits. Overall, our results suggest cooperation favoured by kin selection, with an average relatedness of $r=0.77$ between interacting individuals.

Introduction

Laboratory studies have suggested that bacteria cooperate in a diversity of ways (1, 2). Individual cells produce and secrete molecules to collectively scavenge nutrients, fight antibiotics, and move through their environment (3–5). This cooperation is however vulnerable to cheating by non-producers, who withhold their own cooperation whilst benefiting from that of others (6). A resolution to this vulnerability is kin selection, where cooperation is favoured because the benefits of cooperation go to related cells which share the gene for cooperation (7). Laboratory experiments have also supported a role of kin selection, with experimental evolution, and by showing how clonal growth makes neighbouring cells highly related, and limited diffusion keeps secreted molecules in the neighbourhood (3, 8–11).

In contrast, there is little evidence for cooperation and kin selection in natural populations of bacteria (12–15). The extent to which bacteria cooperate and interact with close relatives is likely to be highly dependent upon environmental conditions. It is hard to know whether the artificial environments and gene knockouts of lab experiments are representative of natural populations. Experiments have also shown that some traits can be cooperative in some environments but private in others (16). Across species comparative studies have shown that cooperation is more common in species where relatedness is higher (17, 18), but this doesn't

help us determine whether specific traits are evolving as cooperative public goods. For this,

we need a way to study bacteria in their natural environment.

Molecular population genetics offers a promising method to test for cooperation and kin

selection in natural populations. Theory predicts that kin selection leaves a distinct signature

(footprint) of selection in genomes (Figure 1) (19–23). If a gene encodes a trait that provides

a direct benefit to the individual performing that trait, then the cells which perform that trait

carry the gene for that trait. Similarly, in a clonal population, relatedness between cells is

$r = 1$, and so the cells which benefit from cooperation also carry the cooperative gene. If

instead, relatedness between cells is $r < 1$, the cells which benefit from cooperation won't

always carry the gene for cooperation. This dilution of the benefit of cooperation leads to

relaxed (reduced) selection, relative to directly beneficial traits or in a clonal population ($r =$

1).

The relaxed selection when $r < 1$ results in an increased probability of fixation for deleterious

mutations, and a decreased probability of fixation for beneficial mutations (14-16). The

consequence of this change in fixation probabilities, when $r<1$, is that it would lead to

increased polymorphism and divergence in cooperative genes relative to genes that have

direct fitness effects (Figure 1). Consequently, by examining patterns of polymorphism and

divergence, we can test for signatures of cooperation favoured by kin selection. This method

has been applied in microbes to the social amoeba *Dictyostelium discoideum* with mixed

results, although relatedness is close to $r = 1$ in that species, and so we might not expect to a

significant signature of kin selection (24–27). We have also previously applied this method to

the opportunistic pathogen *Pseudomonas aeruginosa*, and found evidence for cooperative

traits favoured by kin selection (28). A key advantage of this method is that it solves the

problem caused by traits that are only cooperative in certain contexts, as the signature of

selection will represent the average conditions experienced over time.

Figure 1: Population genetic theory for cooperative traits. (A) Representation of the categorisation of traits as private (yellow) or cooperative (blue). Cooperative traits involve the production and secretion of molecules whose fitness benefits are shared with other nearby cells. Private traits are those whose fitness benefits are only felt by the individuals expressing the trait (B) Prediction for relative polymorphism and divergence for private (yellow) and cooperative (blue) genes. If relatedness, $r=1$, then cooperative genes (middle blue violin) should have the same polymorphism and divergence as private genes (left yellow violin). In contrast, if $r<1$ then cooperative genes (right blue violin) should show greater polymorphism and divergence than private genes. Figure based on (19, 21).

103 However, there were two potential problems with the only previous analysis with bacteria, on

104 *P. aeruginosa* (28). Firstly, the genomes had been collected over several decades, and from

105 six different continents. The underlying population genetic theory assumes that all genomes

106 are sampled from a single population, at a single time point. Violating this assumption could

107 have led to biased or spurious results (29). Secondly, we were unable to directly control for

108 gene expression. If a gene is only occasionally expressed, in certain conditions, then this can

109 also lead to relaxed selection, with increased polymorphism and divergence (21). Belcher *et*

110 *al*. lacked data on gene expression, and so controlled for this problem by making targeted

111 comparisons between cooperative and private traits that they argued were likely to be

112 expressed at different rates (28). If their assumptions did not hold, and expression rates

113 differed, then then their patterns could alternatively be explained by gene expression rather

114 than kin selection for cooperation.

116 We were able to address these problems by taking advantage of two recently developed data

117 sets in the bacteria *Bacillus subtilis*. *B. subtilis* is found in soil and the gastrointestinal tracts

118 of several animals, including humans, and is used on an industrial scale by biotechnology

companies (30). A number of laboratory studies have suggested that *B. subtilis* is a highly
cooperative species, which secretes a number of potentially cooperative enzymes (31–34).
First, we used a natural population consisting of 31 environmental isolates collected as part of
a citizen science project in Dundee, Scotland (35) (Supplementary Table 1). All of these
strains were collected from the wild around the same time, and from similar niches. Second,
we were able to directly control for gene expression rates, by using two genome-wide studies
of gene expression across several timepoints during biofilm formation (36, 37). Taken
together, these studies allow us to satisfy the assumptions of the underlying population
genetic theory.

Results & Discussion

We compared genes controlling traits that are hypothesised to be cooperative, with traits that
are hypothesised to be private. We identified six different types of putatively cooperative
behaviour, where an appropriate comparison could be made with genes controlling private
traits, that are likely to be expressed at similar rates (Table 1). We used the first type, quorum
sensing (QS), for the main analysis, and we summarise the other five types at the end of the
results section.

Quorum Sensing

We started by examining genes induced by the ComQXPA quorum sensing (QS) signalling
system (33, 34). This system regulates gene expression in response to the density of a
diffusible signal molecule. At high cell densities, the density of the signal molecule increases,
causing ComA to activate, and the upregulation of a number of traits (38–41). We
categorized genes as cooperative or private based on a search of the literature in *B. subtilis*
(Methods). For example, the fifteen genes coding for the exopolysaccharide EPS are classed
as cooperative. EPS is the main biofilm matrix component (42), and is required for biofilm
formation (43). EPS is costly to produce, it provides benefits to non-producers, and non-
producers can exploit producers (44). This is a classic public good. Similarly, TasA, a protein
fibre that is needed for biofilm structural integrity (45, 46), has also been shown in lab
experiments to be a public good (32). Mutants lacking either EPS or TasA can also
complement each other, private further evidence that the benefits of these genes are shared
(47). Private genes include those coding for traits such as asparagine synthase (AsnB), which

controls peptidoglycan hydrolysis for cell growth and cell-wall synthesis (48).. We found that

quorum sensing controls a mixture of private and cooperative traits in *B. bacillus,*

categorizing N=25 of our quorum sensing-controlled genes as cooperative, and N=153 as

private (Supplementary Table 2).

We started by focusing on quorum sensing because it offers a number of advantages for our

purpose. First, the large size and nature of this network means that there are sufficient private

and cooperative genes for a targeted analysis (N=153 & N=25 respectively). Second, shared

control by the same signalling system means that private and cooperative genes controlled by

the quorum sensing system are likely to have similar levels of expression (49, 50).. Third,

there is data on gene expression allowing us to directly compare expression rates (36).

Fourth, the coregulation of genes acts as a control for mutations in noncoding regulatory and

promoter regions that could affect the production of cooperative public goods.

164
165 *QS: Controlling for conditional expression*

166
Differential gene expression can also influence the strength of selection and so needs to be

controlled for. Theory tells us that the fraction of generations in which a trait is expressed

can determine the extent to which selection is relaxed (21). To control for this, we need to

compare genes that are switched on or off in the same conditions. Shared control by the same

signalling system means that private and cooperative genes controlled by the quorum sensing

system are likely to have similar levels of expression. We were, however, also able to test this

assumption directly by examining two data sets on gene expression (36, 37). Gene expression

depends strongly upon environmental conditions and so there is no single rate of gene

expression for each gene.

176
Futo *et al*. measured gene expression at 11 different points in the formation of a biofilm, and

normalised their results to the median to convert expression levels to the same scale (36).

This gives us an excellent dataset to test our control for conditional expression, as we can use

simple correlations between pairs of genes to see if they are up- and down-regulated at the

same time. For our 178 private and cooperative genes, there are 15753 unique pairs of genes.

The average correlation in gene expression for a pair of these genes is 0.302 (Spearman's

correlation). We then used a bootstrap approach to see if this correlation was greater or lower

than for randomly chosen genes. We took a random set of 178 genes and calculate mean

pairwise correlation in the same way, and repeated 10000 times. We found that the mean

pairwise Spearman's correlation was 0.251 for the random (bootstrap) samples, and that the

correlation in expression of our quorum sensing-controlled genes was higher than 99.7% of

our bootstrap samples (Figure 2). Consequently, our candidate set of genes have expression

rates that are correlated significantly higher than expected by chance, supporting our choice

for their use in an analysis of signature of selection (p<0.004).

Pisithkul provided a different dataset measuring gene-expression in biofilms (37). Whereas

Futo measured expression over two months in a biofilm with a solid-air interface, Pisithkul

focussed on the initial stages of biofilm growth, measuring expression over 24 hours in a

biofilm with a liquid--air interface. We again found that quorum sensing-controlled genes

have expression rates that are correlated significantly higher than expected by chance (N=160

genes, p<0.02; Supplement S11).

Figure 2: Average correlation in gene expression between genes during biofilm growth. (**A**) Correlation across 11 timepoints of biofilm formation for randomly sampled gene sets of the same size as our quorum sensing-controlled genes (N=178). Data from Futo et al. (2021).

 (**B**) Correlation across 8 timepoints of biofilm formation for randomly sampled gene sets of the same size as our quorum sensing-controlled genes (N=160). Data from Pisithkul et al. (2019). In both panels, the orange line shows the average correlation for the quorum sensing-controlled genes, which is >99.7% of our random samples for panel A, and >98.5% of random samples for panel B.

199
200   *QS: Polymorphism*
201

We found that genes for putatively cooperative traits had significantly greater polymorphism

than genes for private traits (Figure 3; ANOVA   ; Games Howell

Test p<0.001; N=25 cooperative genes, N=153 private genes). Cooperative genes also had

significantly greater polymorphism when we only examined only non-synonymous sites

(Kruskal-Wallis   ; Dunn Test p=0.0240. Figure 4a), or only synonymous

sites (ANOVA   ; Games Howell Test p=0.007) (Figure 4b).

The trend for greater synonymous polymorphism is possibly surprising as such sites should

be under much weaker selection, and we wouldn't necessarily expect to see an effect of kin

selection. However we also found this pattern in *P. aeruginosa* (28). There is some evidence

that differences in use of preferred codons between cooperative and private genes could

explain this pattern (Supplement S1). Synonymous mutations can also have substantial fitness

effects on social traits (51–54).

The ratio between non-synonymous and polymorphism does not differ between cooperative

and private genes (Kruskal-Wallis   ; Dunn Test p=0.183.

Supplementary Figure 1), which reflects the fact that cooperative genes have elevated

diversity at both types of site, possibly due to the large fitness effects of many traits (both

cooperative and private) that are quorum sensing-controlled (55, 56) (Supplement S2).

221
222
223

Figure 3: Nucleotide diversity per site for private (yellow) and cooperative (blue) genes controlled by quorum sesnsing. Each point is a gene, and the horizontal line shows the median for each group. The grey line shows the median for background private genes

across the genome.

224
225

Figure 4: Nucleotide diversity at non-synonymous (A) and synonymous (B) sites for private (yellow) and cooperative (blue) genes controlled by quorum sensing. Each point is a gene, and the horizontal line shows the median for each group. The grey line shows the median for background private genes across the genome.

226
As an additional control, we were able to repeat all of these analyses comparing cooperative

quorum sensing genes against a different group of private genes not controlled by quorum

sensing (N=1832). This different set of N=1832 private genes, which we call 'background

genes' are those which aren't controlled by quorum sensing and whose products are found in

the cytoplasm, where they are least likely to have a cooperative function. In all cases we

found the same pattern, with cooperative quorum-sensing genes showing elevated

polymorphism compared to background genes (Supplement S2).

234
235 *QS: Divergence*
236
237 We found that cooperative genes had significantly greater divergence compared to private

238 genes at both at non-synonymous and synonymous sites (Non-synonymous: Kruskal-Wallis

239   ; Dunn Test p=0.00553; Figure 5a; Synonymous: ANOVA

240   ; Games Howell Test p=0.011; Figure 5b). We examined synonymous and non-

synonymous sites separately because we measure divergence through rates of protein

evolution. A signature of selection can be found at both synonymous and non-synonymous

sites, implying that synonymous variation is not neutral (26, 28). Because divergence is

elevated at both types of site (similar to polymorphism), there is no difference in the ratio of

non-synonymous to synonymous divergence (Kruskal-Wallis   ;

Dunn Test p=0.189. Supplementary Figure 2), although both sets of quorum sensing genes

have a higher ratio than the background genes (Supplement S2). This could reflect stronger

selection on quorum sensing traits than background traits on average.

Figure 5: Divergence at non-synonymous (A) and synonymous (B) sites for private (yellow) and cooperative (blue) genes controlled by quorum sensing. Divergence is measured by rates of protein evolution, e.g. number of synonymous substitutions per synonymous site for panel B. Each point is a gene, and the horizontal line shows the median for each group. The grey line shows the median for background private genes across the genome.

250
251
252 *Alternative hypotheses*
253
We were also able to eliminate alternative explanations for the patterns that we observed with

both polymorphism and divergence (Figures 3-5). Greater polymorphism would also have

been expected if cooperative genes were more likely to be experiencing balancing or

frequency dependent selection, while greater divergence would arise from positive/directional

selection leading to fixation of adaptive mutations (19, 20). Alternatively, the pattern we

observed could have been caused by other differences between putatively cooperative and

private genes. We assessed these alternative hypotheses by testing other predictions that they

make.

First, we found no evidence that cooperative genes were more likely to be under balancing

selection, which would lead to a deficit of rare alleles in the population (Tajima's D, Fu &

Li's F, Fu & Li's D, Supplement S3). Second, we found no evidence that cooperative genes

are more likely to be under positive selection, which would lead to an excess of non-

synonymous divergence compared to non-synonymous polymorphism, as adaptive mutations

would quickly spread and only be detected as divergence (Mcdonald-Kreitman test,
Neutrality Index, Direction of Selection statistic, Supplement S4). Third, we found no
evidence that the patterns observed are due to any differences in gene length or likelihood of
horizontal gene transfer between cooperative and private genes, or a lack of statistical power
(Supplement S5). Fourth, we found no evidence that our results could be explained by noise
due to variation in the recombination rate. The set of strains that we analysed vary in genetic
competence, the ability to take up DNA from the environment (35), which is a form of
recombination in bacteria. This variation in competency could create noise in our population
genetic measures that focus on SNPs, due to the variation in recombination. Considering the
31 strains we analysed, 18 are genetically competent (35). We conducted an analysis of
polymorphism and divergence using only the competent strains, and found the same patterns
as when we use all strains (Supplement S7).

Finally, we found no evidence that the patterns we observed could be caused by division of
labour (57–60), which is a cooperative hypothesis not mutually exclusive with kin selection.
In *B. subtilis* biofilms, some cells will produce the polysaccharide EPS, and some will
produce TasA amyloid fibres (61). Mutants lacking cannot grow alone but can grow together
(32). The extra level of conditional expression in these public goods (over that caused by
being quorum sensing-controlled), could leave a signature of relaxed selection that isn't
caused by kin selection. To investigate this possibility, we took advantage of the fact that this
heterogeneity is ultimately caused by Spo0A. Spo0A is a bistable switch that is active in only
a subset of cells (61, 62) and activates SinR anti-repressors which control the operons for
both EPS and TasA (32, 63, 64). If the extra conditionality of division of labour was causing
an effect, we would expect that genes under the control of Spo0A (N=20) would have greater
polymorphism than other quorum sensing-controlled genes (N=157), but this is not the case
(Supplement S6). We note that the cooperative genes EPS and TasA that are known to have
division of labour both stand out as highly polymorphic within the genes controlled by
Spo0A, providing support to our hypothesis that sociality causes the effect. We note that the
cooperative genes EPS and TasA that are known to have division of labour both stand out as
highly polymorphic within the genes controlled by Spo0A, providing support to our
hypothesis that sociality causes the effect.

299
300 *Other Cooperative Traits*
301

Our above analyses have provided strong evidence for relaxed selection due to kin selection
for cooperation, considering genes controlled by quorum sensing. We then tested if the same
pattern was found in five other types (groups) of traits, where we could compare genes for
putatively private and cooperative traits, that are likely to be expressed at similar rates (Table
1; Figure 6). The extent to which the distinction between private and cooperative traits can be
made varies across these other groups of traits. Consequently, we might not expect to see a
signature of kin selection in every case, and so our aim is to see if there is a relatively
consistent pattern.

First, *B. subtilis* produces and secretes a siderophore named bacillibactin, which binds to iron
in the environment (65, 66) (Figure 6b). The bound complex can be taken into the cell, but is
also available to non-producers, and is therefore a public good. We separated the genes
involved in the bacillibactin pathway into cooperative and private components, with: genes
involved in biosynthesis and export classed as cooperative, and those involved in uptake and
release of bound iron classed as private. Second, *B. subtilis* exhibits resistance to
antimicrobials by either pumping intact antibiotics outside the cell (private), or by producing
enzymes such as beta-lactamases that detoxify the environment for the entire community
(67–69) (cooperative; Figure 6c).

Third, *B. subtilis* produces proteases to break down proteins, with different proteases acting
either inside the cell (private), or secreted to act outside the cell (70–72) (cooperative; Figure
6d). Fourth, *B. subtilis* produces toxins which can either be contact-dependant (relatively
private), or diffusible throughout the community (cooperative public goods; Figure 4d).
However, this comparison is relative, and possibly weak, as killing cells with contact-
dependant toxins can also provide a cooperative benefit to other local cells, that experience
reduced competition. Fifth, *B. subtilis* has a number of antimicrobial traits which are more
defensive against predators without affecting the predators' growth (private) and those which
are more offensive against competitors (cooperative) (73, 74). This is also a weak
comparison, as bacillaene (the defensive molecule) is secreted from cells, and so likely also
has some cooperative component. However, the defensive traits provide a relatively more
private benefit in providing personal protection, whereas the removal of competitors by the
offensive traits provides a relatively more cooperative benefit. For all comparisons, we find
that the set of genes have significantly correlated expression, using the same methodology
and data as for the comparison with quorum sensing genes (Supplement S9) (36).

341 **Table 1**: Traits used for comparisons of cooperative vs. private genes. The full gene lists are
342 in supplementary tables 2-7.

Trait	Private genes	Cooperative genes
1. **Quorum sensing traits**	Genes which only affect the fitness of the producing cell (N=25)	Genes for public goods that potentially provide benefits to the local group of cells (N=153)
2. **Iron scavenging**	Genes for uptake and use of iron via the *B. subtilis* siderophore bacillibactin (N=5)	Genes for biosynthesis and secretion of bacillibactin (N=5)
3. **Antimicrobial resistance**	Genes for ABC transporters involved in multi-drug resistance. These genes transport the intact antibiotic outside the cell (N=11)	Genes for enzymes that deactivate beta-lactam and aminoglycoside antibiotics outside the cell (N=3)
4. **Proteases**	Genes for intracellular proteases, which are likely to be involved in processing and regulation of proteins within the cell (N=9)	Genes for extracellular proteases, which are likely to be involved in collective feeding and motility (N=8)
5. **Toxins**	Genes for contact-dependant LXG toxins, which are delived by a Type VII secretion system (N=6)	Genes for the secreted toxin bacilysin, which is activate against a broad range of bacterial competitors (N=8)
6. **Antimicrobial activity**	Genes for the secondary metabolite bacillaene, which inhibits predation by *M. xanthus* (N=13)	Genes for the secondary metabolite plipistatin, which is active against fungal competitors (N=4)

Figure 6: Diagram of how traits are categorised as either private (yellow) or cooperative (blue). Panel (A) shows categorisation of quorum sensing controlled traits for the main analysis, with private traits giving fitness benefits only to those expressing the gene, and cooperative traits giving fitness benefits that can potentially be shared with other cells. Panels B-F show other cooperative traits; (B) Bacillibactin (iron scavenging). (C) Antimicrobial resistance. (D) Proteases. (E) Toxins. (F) Antimicrobials. Note that for some of these comparisons it is the relative level of sociality that is different. Figure created with BioRender.com

The number of cooperative genes was too small to analyse each case separately – for

example, the iron-scavenging comparison involves only 10 genes (5 private and 5

cooperative). Consequently, we examined the data in three ways. First, we grouped all

cooperative genes together into one set (N=52) and compared them to the grouped private

genes across all comparisons (N=194). Second, we grouped all of the cooperative genes from

just the five new comparisons (i.e. not quorum sensing) (N=27), and compared them to the

private genes from these five comparisons (N=41). Third, we consider each of the six

categories of cooperative vs. private genes (Table 1) as its own data point (N=6).

Other Cooperative Traits: Polymorphism and divergence

We found that polymorphism was consistently higher in cooperative genes than in private

genes (Figure 7). This pattern is consistent across the three different ways that we can

analysed our data: all genes comparison (ANOVA $F_{2,127} = 10.59, p < 0.0001$; Games Howell

Test p=<0.001); just the genes for the social traits other than quorum sensing (ANOVA

$F_{2,47} = 5.94, p < 0.01$; Games Howell Test p<0.01); and the six categories comparison

(Wilcoxon signed-rank test V=21, p=0.031).

Polymorphism at non-synonymous sites is also significantly higher in cooperative genes than

in all private genes. This pattern is consistent across the three different ways that we can

analysed our data: all genes comparison (Kruskal-Wallis

$X^2(2) = 19.71, p < 10^{-4}, \text{Dunn Test p} < 10^{-4}$); just the genes for the social traits other than

quorum sensing (Kruskal-Wallis $X^2(2) = 9.90, p < 0.01, \text{Dunn Test p} < 0.01$); and the six

categories comparison (Wilcoxon V=21, p=0.031) (Supplementary Figure 3).

Polymorphism at synonymous sites is also significantly higher in all cooperative genes than

in all private genes when analysing all genes (ANOVA $F_{2,130} = 4.21, p = 0.016$; Games

Howell Test p=0.019). However, this trend was not significant when examining just the

genes for the social traits other than quorum sensing (ANOVA $F_{2,47} = 2.28, p = 0.113$; Games

Howell Test p=0.10), or when using categories as data points (synonymous polymorphism is

marginally higher in private genes than cooperative genes for the iron-scavenging and AMR

categories; Wilcoxon V=18, p=0.156). We would expect the pattern to be weaker with

synonymous polymorphism, as these sites are likely to be under weaker selection.

Non-synonymous divergence is significantly greater in all cooperative genes compared to all private genes. This pattern is consistent across the three different ways that we can analysed our data: all genes comparison (Kruskal-Wallis $X^2(2) = 22.9$, $p < 10^{-4}$, Dunn Test $p < 10^{-4}$); the genes for the social traits other than quorum sensing (Wilcoxon V=21, p=0.031 Supplementary Figure 5); and the six categories comparison (Kruskal-Wallis $X^2(2) = 14.7$, $p < 0.001$, Dunn Test $p < 0.001$). As would be expected, the pattern is more mixed for synonymous divergence. Whilst synonymous divergence is significantly greater in all cooperative genes compared to all private genes (ANOVA $F_{2,125} = 8.33$, $p < 0.001$; Games Howell Test p<0.01), this isn't the case when examining just the genes for the social traits other than quorum sensing (ANOVA $F_{2,45} = 2.10$, $p = 0.13$; Games Howell Test p=0.39), or the six categories comparison (Wilcoxon V=13, p=0.688) (Supplementary Figure 6). Similarly, the ratio between non-synonymous and synonymous divergence is significantly greater in all cooperative genes compared to all private genes (Kruskal-Wallis $X^2(2) = 25.0$, $< 10^{-5}$, Dunn Test $p = 0.021$), but not when we just looked at the genes for the social traits other than quorum sensing (Kruskal-Wallis $X^2(2) = 13.2$, < 0.01, Dunn Test $p = 0.11$) or the six categories comparison (Wilcoxon V=18, p=0.156) (Supplementary Figure 7). This may reflect differences in selection on synonymous variation on different traits, or the weakness of some of these comparisons due to small sample sizes. We confirmed that none of our cooperative vs. private comparisons show significant differences in balancing or positive selection (Supplement S10), suggesting that what we are seeing is a signature of kin selection, and that we may just lack power in our other comparisons.

Overall, these results incorporating other traits in addition to quorum sensing controlled genes, provide support to the main result that there is a signature of kin selection for cooperation (Figures 3-5). Nonetheless, the *a priori* distinction between private and cooperative traits is weaker for some comparisons, and we could expect exceptions within the overall pattern. The main exception in our analyses was toxin comparison, where we compared contact-dependant LXG toxins (N= 6 genes) to the secreted antimicrobial bacilycin (N=8 genes). Both of these sets of genes are involved in competition in biofilms, and both are controlled by the DegS-DegU system, so likely expressed at similar rates (75, 76). Possible confounding factors in this case include the fact that although we classified the contact-dependent toxins as private, they also provide cooperative benefits to local cells by

410   eliminating competitors. The LXG toxins also stand out because three of them are on phage
411   elements (75), and it may be that the strength or type of selection is different on these genes,
412   masking any effect of sociality.

Figure 7: Private (yellow) vs. cooperative (blue) polymorphism in genes for six traits.
Panel A shows the quorum sensing-controlled genes used in the main analysis. Panels B-F
show the other cooperative traits.

413
All traits: Deleterious mutations

As an additional robustness test for our conclusions, we analysed deleterious mutations. If
kin selection is favouring cooperation, we should also observe more deleterious mutations in
genes controlling cooperative traits compared to private traits. This is because relaxed
selection slows the rate at which deleterious mutations are purged from the population (19,
20, 22). We tested this prediction by looking for loss-of-function mutations, that generate
stop codons or frameshift mutations, and hence act as deleterious mutations, in our SNP data.
We repeated this analysis with two different data sets.

First, we used all cooperative genes from our six comparisons in Table 1. We measured how
many cooperative and private genes have deleterious mutations, and compared this to an
expectation based on their relative frequency across the genome. Cooperative genes were
significantly more likely than private genes to have deleterious mutations (X^2 (1) = 12.3, $p <$

0.001. This pattern also holds if we count total number of deleterious mutations, rather than

just presence or absence (X^2 (1) = 11.0, $p < 0.001$.

To test the robustness of this result, we repeated the analysis using the localization prediction

tool PSORTb to categorize genes for extracellular proteins as 'cooperative', and genes for

proteins that aren't secreted as 'private (77). This method has been previously used in several

studies to estimate whether genes are for cooperative (social) or private traits (78, 79). By

using PSORTb we are able to analyse all genes, which increases our sample size and

statistical power. We removed the 17% of all genes with unknown localization, leaving us

with a set of genes of known sociality. We found deleterious mutations in 293 genes, of

which 17 are cooperative (5.8%). This is significantly more than expected given that

cooperative genes only make up 2.0% of genes (binomial test P<0.001), matching our

prediction that deleterious mutations should be biased towards cooperative genes. If we count

total deleterious mutations (rather than number of genes with at least one) we see the same

pattern, with cooperative genes making up 5.5% of mutations (20 of 361).

Relatedness estimation

The genetic relatedness between interacting cells (*r*) is a key parameter for social evolution.

Relatedness can be very hard to estimate for natural populations of bacteria and other

microbes, except for extreme cases where interactions take place in some physical structure

such as a fruiting body or a filament (80, 81). Population genetic data allows relatedness to be

estimated indirectly, because the degree to which selection is relaxed, and greater

polymorphism will be observed, depends upon the relatedness between interacting cells.

Consequently, we can work backwards from the polymorphism data to obtain an indirect

estimate of relatedness (28). Examining the polymorphism data from all genes, we estimated

relatedness to be r=0.77 (95% CI 0.67-0.91) (Supplement S8). This estimate assumes that

both the magnitude of selection and the distribution of selection coefficients is the same on

average for cooperative and private genes. An advantage of this indirect method for

estimating relatedness is that it does not require knowledge about factors that would be hard

or impossible to measure in natural populations. For example, the spatial details of how

cooperative interactions play out, such as how far do public goods diffuse, and who benefits,

as well as how much these vary in different environments, and the frequency with which

different environments are encountered (28, 82). Indeed, cooperative traits in *B. subtilis* vary

in the degree to which they are shared depending on whether groups are exhibiting sliding

motility or growing in biofilms (16). In contrast, the indirect measure provided by population
genetics represents an average for the different cooperative traits, over the different
environments encountered, over evolutionary time.

Conclusions

We have found strong evidence of kin selection for cooperation in a natural population of *B.*
subtilis. Our analyses controlled for possible confounding factors, such as expression rate,
and eliminated alternative explanations for polymorphism and divergence, providing
evidence that complements the lab experiments demonstrating sociality in this species (32,
42–46, 61, 83–86). Taken together with a previous study, population genetic analyses have
now provided evidence of kin selection for cooperation in both a gram-positive (*B. subtilis*)
and a gram-negative (*P. aeruginosa*) bacteria (28). These results suggest convergent, and
potentially widespread, kin selection for cooperation, based on very different underlying
mechanisms, across bacteria.

A possible complication with studying cooperation in bacteria is that the extent to which
traits are cooperative, and their importance for fitness, can depend greatly upon
environmental conditions (8, 87–90). One advantage of the molecular population genetic
approach that we have used is that it averages across different environments over
evolutionary time. Consequently, rather than examining a specific environment, it provides
an 'average' answer. In the case of *B. subtilis,* for the traits that we have examined, we have
found evidence of kin selection for cooperation, with an estimated average relatedness of
$r=0.77$.

480
481
482

Acknowledgements

484
485 We thank: Ming Liu and Ákos Kovács for useful discussion and comments on the
486 manuscript. This work was supported by the European Research Council (834164: LJB, AED
487 & SAW; SESE: MG).

Material & Methods

Strains

We use the whole-genome sequences of 31 strains of *Bacillus subtilis* from (35). The strains
are environmental isolates, collected from a citizen science project in Dundee where people
brought soil samples from their garden. *B. subtilis* is most commonly found in soil, but is also
found living as a commensal in animal intestines (91), and in marine environments (92)..
Whilst durable spores than can disperse through the air can allow long-distance migration
(93), and strains don't phylogenetically cluster based on environment (94), several factors are
in favour of using these strains as our natural population.. Firstly, these samples were all
collected at the same time for the same project (35). Further, we know that rates of migration
between populations scales with geographic distance, most of the sequence diversity within
the species in contained in local population (93), and there is evidence for fine-scale genetic
structure in micro-scale populations (95).

We downloaded raw sequence data for each strain from the European Nucleotide Archive
(accession number PRJEB43128). The full list of strains can be found in Supplementary
Table S1.

Supplementary Table 1: List of strains used

Strain ID	Location	Assembly
NRS6096	Tayport (UK) garden soil	GCA_905311035.1
NRS6099	Tayport (UK) garden soil	GCA_905310985.1
NRS6103	Tayport (UK) community garden soil	GCA_905310995.1
NRS6105	Tayport (UK) community garden soil	GCA_905311425.1
NRS6107	Tayport (UK) garden soil	GCA_905311405.1
NRS6108	Tayport (UK) garden soil	GCA_905311395.1
NRS6110	Tayport (UK) vegetable plot	GCA_905311415.1
NRS6111	Tayport (UK) vegetable plot	GCA_905311375.1
NRS6116	Tayport (UK) garden soil	GCA_905311385.1
NRS6118	Tayport (UK) potato patch	GCA_905311435.1
NRS6120	Tayport (UK) potato patch	GCA_905312035.1
NRS6121	Tayport (UK) shrub bed	GCA_905315035.1
NRS6127	Tayport (UK) garden soil	GCA_905315045.1
NRS6128	Tayport (UK) garden soil	GCA_905315055.1
NRS6131	Tayport (UK) worm bin	GCA_905315025.1
NRS6134	Tayport (UK) vegetable patch	GCA_905315395.1
NRS6137	Tayport (UK) community garden soil	GCA_905315385.1

NRS6141	Tayport (UK) garden soil	GCA_905315375.1
NRS6145	Tayport (UK) garden soil	GCA_905315685.1
NRS6148	Tayport (UK) garden soil	GCA_905315695.1
NRS6153	Tayport (UK) garden soil	GCA_905315705.1
NRS6160	Tayport (UK) garden soil	GCA_905315715.1
NRS6167	Tayport (UK) soil from planter	GCA_905316385.1
NRS6181	Tayport (UK) garden soil	GCA_905318255.1
NRS6183	Lochee (UK) garden soil	GCA_905319155.1
NRS6186	Newport (UK) garden soil	GCA_905319135.1
NRS6190	Tayport (UK) garden soil	GCA_905319565.1
NRS6194	Tayport (UK) garden soil	GCA_905319555.1
NRS6202	Kirriemuir (UK) garden soil	GCA_905319535.1
NRS6205	Kirriemuir (UK) garden soil	GCA_905319825.1
NRS6206	Kirriemuir (UK) garden soil	GCA_905319815.1

Genes regulated by quorum sensing

521
522
For our set of quorum sensing controlled genes, we combine three published datasets: (1) 88
genes controlled by ComXAP (40); (2) 114 genes controlled by degU (96); (3) 40 genes
controlled by Spo0A (63).. We didn't use a fourth possible dataset, of 166 genes affected by
the competence transcription factor ComK, which are indirectly regulated by ComA (97)..
This is partly because our undomesticated reference strain (NCIB 3610) carries a plasmid-
encoded protein which interferes with the competence machinery (98) (Supplement S7).). In
addition, we wanted to focus on the quorum sensing systems known to produce public goods
(Figure 3 of (34)).

Identifying social genes

532
533
We used an artisan approach to identify social genes based on laboratory studies which have
demonstrated that a trait is cooperative. The gold-standard test for a cooperative gene
involves a wildtype strain which produces the traits, and a mutant strain which doesn't. If a
trait is cooperative, then the producer will outperform the non-producer when each is grown
clonally, but non-producers will outperform producers in groups (2). As an example, we look
at the first gene on the list, bslA (formely known as yuaB), which is involved in biofilm
formation, and specifically in making the biofilm hydrophobic to resist chemical attack (99).
A non-producer of bslA cannot form normal biofilms on its own, but can get into mature
biofilms when in co-culture with producers (100). Further work mixing producers and non-
producers at a range of starting ratios demonstrated that the biofilm can maintain function as
long as >50% of cells are producers (99).

545
546 The full list of cooperative genes can be found in Supplementary Table 2.

547
548 **Supplementary Table 2**: List of social genes

Gene	Function	Reference
bslA	biofilm-surface layer protein BslA	Arnaouteli 2020
bslB	biofilm-surface layer protein BslB	Morris 2017

epsA	hypothetical protein	Arnaouteli 2020
epsB	protein tyrosine kinase EpsB	Arnaouteli 2020
epsC	polysaccharide biosynthesis protein	Arnaouteli 2020
epsD	glycosyltransferase family 4 protein	Arnaouteli 2020
epsE	glycosyltransferase EpsE	Arnaouteli 2020
epsF	glycosyltransferase family 1 protein	Arnaouteli 2020
epsG	biofilm exopolysaccharide biosynthesis protein EpsG	Arnaouteli 2020
epsH	glycosyltransferase	Arnaouteli 2020
epsI	polysaccharide pyruvyl transferase family protein	Arnaouteli 2020
epsJ	lipoprotein	Arnaouteli 2020
epsK	cyclic-di-GMP receptor EpsK	Arnaouteli 2020
epsL	sugar transferase	Arnaouteli 2020
epsM	acetyltransferase	Arnaouteli 2020
epsN	aminotransferase class I/II-fold pyridoxal phosphate-dependent enzyme	Arnaouteli 2020
epsO	polysaccharide pyruvyl transferase family protein	Arnaouteli 2020
sipW	signal peptidase I	Arnaouteli 2020
srfAA	surfactin non-ribosomal peptide synthetase SrfAA	Kalamara 2018
srfAB	surfactin non-ribosomal peptide synthetase SrfAB	Kalamara 2018
srfAC	surfactin non-ribosomal peptide synthetase SrfAC	Kalamara 2018
srfAD	surfactin biosynthesis thioesterase SrfAD	Kalamara 2018
srfT	MFS transporter	Kalamara 2018
tapA	amyloid fiber anchoring/assembly protein TapA	Arnaouteli 2020
tasA	biofilm matrix protein TasA	Arnaouteli 2020

549
550 For the robustness check of whether deleterious mutations are over- or under-represented in
551 cooperative genes, we used the protein localisation tool PSORTb 3.0 (77). We categorize
552 cooperative genes as those which PSORTb predicts to be extracellular. We also follow
553 previous studies in removing genes for which PSORTb cannot make a definitive prediction
554 (78)
555

556 *Controlling for conditional expression*
557

558 Conditional expression can lead to the same signatures of relaxed selection as kin selection
559 for cooperation. We directly examined expression rates for the set of genes regulated by
560 quorum sensing, using data from (36), who measured gene expression of >4000 genes at 11
561 timepoints during biofilm formation.
562

For any pair of genes, we can calculate the correlation in gene expression across the 11
timepoints of the biofilm. For the 178 quorum sensing genes in our dataset, there are 15753
unique pairs of genes. The mean pairwise Spearman's correlation in gene expression is 0.302.
To test whether this set of genes is more or less correlated than a randomly chosen set of
genes, we use a bootstrap approach. We take a random set of 178 genes and calculate mean
pairwise correlation in the same way as before. Then we repeat 10000 times. We find that the
correlation in expression of our quorum sensing-controlled genes is higher than in 99.7% of
our bootstrap samples (Supplementary Figure 3), demonstrating that our candidate set of
genes is appropriate for our analysis of signature of selection.

Other cooperative traits

We also examined five other types (groups) of traits, where we could compare genes for putatively private and cooperative traits, that are likely to be expressed at similar rates (Table 1; Figure 4). Firstly, we used iron-scavenging via siderophores, which is a well-studied cooperative trait that is important for growth and survival of bacteria (3, 101). Specifically, we looked at the *B. subtilis* siderophore bacillibactin (65, 66). We classified the genes for biosynthesis bacillibactin as cooperative, and genes for uptake and release of bound bacillibactin as private (Supplementary Table 3).We classified the genes for biosynthesis bacillibactin as cooperative, and genes for uptake and release of bound bacillibactin as private (Supplementary Table 3).

Second, we looked at antibiotic resistance genes. There are many mechanisms of antibiotic resistance, some of which are cooperative. For example, the secretion of beta-lactamases is a cooperative trait as they detoxify the external environment, providing benefits to the local population (5, 102). We also classified aminoglycoside resistance as cooperative, as the modification of the antibiotic detoxifies the local environment (103). For the private genes, we used the eight ABC transporters that are thought to be involved in multi-drug resistance by pumping antibiotics outside the cell (69, 104) (Supplementary Table 4).

Thirdly, we looked at the range of peptidase proteases produced by *B subtilis*. The functions of these proteases are broad, covering processing, regulation, and feeding, but we can separate them in cooperative and private genes by looking at those which are secreted (i.e. are extracellular) and those which are not secreted (70). The secreted proteases are more likely to have cooperative fitness effects on other cells, through nutrition, interacting with host immune systems etc (Supplementary Table 6)..

Fourthly, we looked at toxins. For the cooperative genes, we used bacilycin, which is a secreted antimicrobial peptide that is active against a range of bacteria (105). Because bacilycin can diffuse through the environment, it likely has cooperative fitness effects on others. *B. subtilis* also has many toxins which are involved in contact-dependent inhibition, and therefore likely have private effects on fitness. For the private gene, we used six LXG toxins (75), which are delivered by a Type VII secretion system. Whilst these toxins can still have cooperative effects by removing competitors, they are by their nature less cooperative than secreted molecule such as bacilycin. Both of these sets of genes are under control of the DegS-DegU system (Supplementary Table 5).

Fifthly, we looked at antimicrobials. *B. subtilis* produce a series of antimicrobial molecules, which vary in which organisms they target, how they act, and how they are secreted. We can however distinguish between antimicrobials that have a more defensive role in traits such as predation avoidance, and those which have a more offensive role in competition with other species. Whilst both of these categories likely have some component of cooperative and private effects on fitness, the offensive ones will be relatively more cooperative. This is because defensive molecules have a stronger effect on the individuals producing them (Supplementary Table 7).

Identifying deleterious mutations

We used the variant annotation tool SnpEff (106) to look for SNPs that generate deleterious
mutations in our dataset.. Specifically, we annotate two types of mutations; (1) premature
stop codons and (2) frameshift mutations. This gives us a list of genes that have at least one
deleterious mutation. To test if a given set of genes are overrepresented for deleterious
mutations, we use two percentages; (1) the % of genes in the whole genome that are in that
set, and (2) the % of genes with deleterious mutations that are in this set. We compare these
values using a binomial test with the null hypothesis that the number of deleterious mutations
in the gene set is equal to that expected by the frequency of the gene set. For any given gene
set, we conduct a further test where we use the total number of deleterious mutations in that
set of genes, rather than just presence/absence of deleterious mutations for that gene.
631

Statistics & Figures

We conducted all statistical analysis in R (107).. For the main statistical analysis comparing
molecular population genetic parameters of cooperative and private genes we use one of two
statistical tests, depending on the variable in question. For variables that are normally
distributed, we used an ANOVA to compare the three groups of genes. Because of unequal
sample sizes we used Welch ANOVA which doesn't require equal variance. We used the
Games-Howell post-hoc test, which is similar to Tukey's HSD, but designed for Welch
ANOVA where we don't have to assume equal variance. For variables that aren't normally
distributed, we used the Kruskal-Wallis test, which compares medians. We then used the
Dunn test for post-hoc comparisons of groups.

All results figures were made using the ggplot2 package in R (108) using colour palettes from
the packages wesanderson (github.com/karthik/wesanderson) and BirdBrewer
(https://github.com/lauriebelch/BirdBrewer). Figure 4 illustrating the secondary comparisons
was made using BioRenderFigure 4 illustrating the secondary comparisons was made using
BioRender.
649

Bioinformatics

652　Raw reads for each of the 31 strains were downloaded from the European Bioinformatics
653　Institute's European Nucleotide Archive (accession number PRJEB43128). We then used a
654　SNP calling pipeline to find SNPs in each strain compared to the reference NCIB 3610
655　(accession NZ_CP020102.1).
656

Trimming and quality-control
658　We used Trimmomatic to remove adapters remove low quality reads, which we did by
659　remove leading and trailing reads if the quality score was <3 or if average quality in a four-
660　base sliding window was <20. We manually checked the output of this step using the reports
661　produced by FastQC (109).

Mapping
664　We used BWA (110) to map reads from each strain to the reference strain. We used
665　SAMtools (111) to convert the mapping files from BAM to SAM, and used Picard tools
666　(112)to remove PCR duplicates.

Variant calling
669　We used BCFtools (113) to call variants on all strains and produce a VCF file that can be
670　read by R for population genetic analysis.

671
Filtering and quality control
We conducted further filtering to remove indels, and filter for mapping quality, read depth,
and strain bias using the default setting of SAMtools vcfutils python script. We then removed
all sites which hadn't been called in at least 80% of strains. We also used the coverageBed
tool in BEDtools (114) to record the percentage of each genes length that had been mapped,
in order to adjust per-site measures to the correct length. After filtering, we had a total of
256,769 SNPs among the 31 strains.
679
680 *Outgroup*
681 We used the phylogeny in Kalamara 2019 (the source of these strains) to identify *Bacillus*
682 *subtilis subsp. spizizenii* str. W23 as an appropriate outgroup (accession NC_014479.1, raw
683 sequencing data SRR2063059). We used the same variant calling pipeline as above to
684 produce a second VCF which included the SNPs from the outgroup.
685
686 **Population genetic measures**
687
We used the PopGenome package from R (115) to conduct the main molecular population
genetic analysis. All parameters were scaled to the corresponding mapped gene length, and
any gene with mapped length <50% of their full length or lacking polymorphism data was
removed from the analysis, leaving 3817 genes for the population genetics analysis. Using
PopGenome, we calculated Nucleotide polymorphism, Tajima's D, Fu and Li's D*, the
Mcdonald-Kreitman p-value, Direction of Selection statistic, and neutrality index. We also
calculated separate measures for synonymous and non-synonymous sites where appropriate.
695
696 To calculate divergence, we measured the rate of protein evolution K_a/K_s by comparing the
697 reference strain to the outgroup. We did this by creating a pseudo-genome of the outgroup by
698 inserting the relevant SNPs into the reference sequence using the GATK suite of tools (116).
699 This pseudo-genome could then be read by R, and we used the seqinR package (117) to
700 calculate divergence.

Supplementary Material

S1: Codon usage

In the main text, we showed that the elevated polymorphism in cooperative genes relative to private genes occurs at both synonymous and non-synonymous sites. This may be due to selection on synonymous codon usage, which is common across bacteria (118). In general, some codons may be preferred due to GC or AT preference, tRNA availability, or metabolic costs (119–122). We investigated whether codon usage differences between cooperative and private genes can explain the elevated synonymous polymorphism observed in social genes.

We used the R package 'sscu' (strength of selected codon usage), which calculates several measures of codon usage bias (123).

For many measures, we compare social genes to a set of highly expressed genes, under the assumption that highly expressed genes are under the strongest selection to optimize codon usage. We followed Rocha 2004 (122) in choosing genes that code for ribosomal proteins as this set. For our social genes, we used the set of genes coding for extracellular proteins (as determined by PSORTb), as this is a measure that can be systematically applied to the whole genome (unlike our 'artisan' method, which requires manual curation of genes).

The first measure we calculated is RSCU, relative synonymous codon usage. This gives a measure of the relative use of each codon, compared to the null expectation that each codon for an amino acid is used equally (RSCU =1) (Supplementary Figure S1.1).

Supplementary Figure S1.1: RSCU for cooperative genes (blue) alongside highly expressed (ribosomal) genes (yellow). The dotted line shows RSCU=1, which is when each codon for an amino acid is equally likely to be used. If codons are used at random, we would expect all RSCU values to be close to 1. If certain codons are strongly preferred, then each amino acid should have a codon for which RSCU>>1

753
754
To determine if codon usage is more even in social genes or highly expressed genes, we
calculate Shannon entropy for each amino acid. If each codon is used evenly, then Shannon
entropy will be 0. If one codon is used more frequently than random, then Shannon entropy
will be negative. This is analogous to how entropy is used in ecology to calculate species
richness. An amino acid mostly being coded for with the same codon is equivalent to one
species dominating in a species richness measure.
761
762 On average, Shannon entropy is much less negative for cooperative genes (-0.215 compared
763 to -0.869 for highly expressed genes), suggesting that cooperative genes have much more
764 even usage of codons, suggesting that there may be relaxed selection for synonymous codon
765 usage in cooperative genes (Supplementary Figure S1.2)
766

Supplementary Figure S1.2: Shannon entropy for cooperative (blue) and highly-expressed (yellow) genes for each amino acid.

767
768
769 Cooperative genes also have higher GC3 (GC content at third positions) than our set of
770 highly expressed genes. [0.407 in cooperative compared to 0.322 in ribosomal] which may
771 relate to variable expression.
772
The effective number of codons is also substantially higher in cooperative genes than in our
set of highly expressed genes (55.1 compared to 43.4), further demonstrating how
cooperative genes are using more codons, and therefore less likely to use preferred codons.
We also conducted chi-squared tests to determine for each codon whether it was used
significantly more than expected (from random chance) in cooperative genes than highly
expressed genes, and found that only 14 codons are used significantly more often than
expected by chance in cooperative genes, compared to 22 codons that are used significantly
more often than expected for highly-expressed genes.
781
782 Overall, there is some evidence that decreased usage of preferred codons may explain the
783 increase in synonymous polymorphism. This is the third study to find the counter-intuitive
784 pattern of increased synonymous polymorphism in social genes (*P. aeruginosa* (28) ; *D.*
785 *discoideum* (26), so more research is needed in this area.

786 **S2: Cooperative vs. background genes**
787
788 We compared cooperative genes to private genes in the main analysis, and here we also
789 compare to a set of background genes. For the background genes we use those which produce
790 proteins that localise to the cytoplasm, as these are least likely to have cooperative functions.
791 This produces a set of 1832 genes.
792
793 Here, we present the results of the post-hoc comparison between cooperative genes and
794 background genes for the main set of molecular population genetic measures
795

Measure	P-value

Nucleotide polymorphism	0.004*
Non-synonymous polymorphism	0.006*
Synonymous polymorphism	0.056
Ratio of non-synonymous to synonymous polymorphism	0.103
Ratio of non-synonymous to synonymous divergence	0.017*
Non-synonymous divergence	0.004*
Synonymous divergence	0.485
Tajima's D	0.220
Neutrality Index	1.000

We find the same pattern as in the main analysis, with cooperative genes having a signature of higher non-synonymous polymorphism and divergence, without evidence for increased likelihood of positive or balancing selection.

This comparison is less well-controlled than the main analysis, but these results add further weight to our conclusion that signature of selection we observed in quorum sensing - controlled genes is a signature of kin selection.

S3: Balancing selection

A logical explanation for cooperative genes to be more polymorphic than private genes would be that cooperative genes are more likely to be under balancing selection. This could occur if both cheats and cooperators are maintained in a population because the fitness advantage of cheats declines as they become more common (124, 125). It could also occur if multiple greenbeard recognition alleles are maintained (126).

To detect balancing selection from sequence alignment data, we can use several population genetic measures such as Tajima's D and Fu & Li's F* and D* that look at allele frequencies to determine if balancing selection is occurring. Tajima's D looks at the distribution of allele frequencies, whereas Fu & Li's measures look at singletons (rare variants found in only one

strain). To interpret these measures, we use statistical tests to determine if each gene is significantly different from the neutral expectation. We then extract the list of genes with significant support, and test if they are overrepresented for social genes using binomial tests.

Tajima's D

We use the beta-distribution test with alpha=0.025 from the Pegas package in R (127, 128) to identify which genes have evidence for balancing selection. The test could be significant either due to balancing selection and a lack of rare alleles (D>>0) or a recent selective sweep (D<<0), so we exclude significant genes where D<0.

We have 242 genes with significant evidence for balancing selection. Only one of those is a cooperative gene, so cooperative genes are not significant overrepresented in those under balancing selection (binomial test, p=0.380). That gene is the aminoglycoside resistance gene aadK.

Fu & Li's D* and F*

We use the critical values from (129) for n=100 genes and alpha=0.025, which is 1.53 for D* and 1.73 for F*. The probability of D* or F* being greater than this critical value for by chance is 0.025. Although we have >100 genes, this is likely to be a good approximation as the critical value scales with the natural-log of n. Although we have >100 genes, this is likely to be a good approximation as the critical value scales with the natural-log of n.

We have 21 genes with significant evidence for balancing selection from Fu & Li's D*. None of them are cooperative. We have 13 genes with significant evidence for balancing selection from Fu & Li's D*. None of them are cooperative

Overall, these results support our conclusions that cooperative genes have higher polymorphism due to relaxed selection, rather than balancing selection.

S4: Positive selection

Cooperative genes might show greater divergence than private genes due to being more likely to be under positive or directional selection.

We use several complementary measures to look for signatures of positive selection in our genes. First, we use the Mcdonald-Kreitman (MK) test, which compares the ratio of non-synonymous to synonymous divergence with the ratio of non-synonymous to synonymous polymorphism (130). If there is lots more non-synonymous divergence than non-synonymous polymorphism, then this could indicate positive selection. The logic is that with strong positive selection, advantageous mutations don't spend much time as polymorphisms, and are mainly detected through divergence. .

We have 15 genes which have evidence for significant positive selection from the MK test.
None of them are cooperative.

Secondly, we use the neutrality index, which uses the same information as the MK test, but
rather than just a binary significance test, it uses the full information to make a continuous
variable (130).. This can be interpreted by comparing averages and distributions of different
groups of genes. We log-transformed neutrality index to normalise, meaning that positive
values indicate positive selection. There is no difference in neutrality index between
cooperative and private genes, and both also don't differ from the background set of genes
(Kruskal-Wallis test, chi-squared=0.06, df=2, p=0.974) Supplementary Figure 4.1.

Supplementary Figure 4.1: Log-transformed neutrality index of private and cooperative
genes. The dashed line shows the median for background genes.

897 We also use the direction of selection statistic, which again uses the same information as the
898 MK test to make a continuous variable, with positive values indicating positive selection.
899 Whilst there is a slight trend for cooperative genes to have a higher direction of selection
900 statistic than private genes, this is not significant (Kruskal-Wallis test, chi-squared=0.09,
901 df=2, p=0.955) Supplementary Figure 4.2.

Supplementary Figure 4.2: Direction of selection statistic for private and cooperative genes. The dashed line shows the median for background genes.

Overall, these findings support our conclusion that cooperative genes are not more divergent than private genes due to being more likely to be under positive selection.

S5: Alternative explanations

937 ***Gene length***
938
939 It is well-known that gene length can affect molecular population genetic parameters such as
940 polymorphism. Even though we calculate all measures per site (considering gene length), we
941 also check here that our results aren't an artefact of any differences in gene length between
942 cooperative and private genes.
943
944 Cooperative genes are on average 19% longer than private genes (965 base-pairs compared to
945 812).
946
947 We conduct a small analysis where we remove the smallest 25% of genes from our analysis,
948 which is those genes which are <450 base-pairs long.
949
950 Cooperative genes are still significantly more polymorphic and divergent than private genes
951 using this reduced dataset (Kruskal-Wallis test, chi-squared=8.16, df=2, p=0.017. Dunn Test
952 p=0.01)
953
954 ***Horizontal gene transfer / pangenome***
955
956 We focused our analysis on chromosomal genes, because frequent horizontal gene transfer
957 can make drawing conclusion from molecular population genetic parameters more
958 challenging. We check if cooperative genes are more likely to be horizontally transferred by
959 checking if they are overrepresented in the accessory genome compared to the core genome.
960
961 *B. subtilis* has an open pangenome, meaning that each additional strain sequenced adds many
962 new genes. This is usually indicative of a species living in multiple or variable environments,
963 which we know is true for this species. *B. subtilis* is also naturally competent, meaning they
964 can take-up DNA from the environment, and it is thought that the open pangenome occurs
965 because rare genes are acquired in this way from closely related species (94).
966
967 We use the panX database (pangenome.org) to assign genes as core or accessory genome
968 based on 80 *B. subtilis* genomes. If we count core genes as those present in 90% of genomes,
969 then 23.2% of the genes in our reference strain are in the *B. subtilis* accessory genome, and
970 76.8% are in the core genome.
971
972 10 out of 53 cooperative genes are in the accessory genome, which is 18.9%. Cooperative
973 genes are therefore not overrepresented in the accessory genome (binomal test p=0.622).
974
975 We conclude that cooperative genes are not more likely to be transferred horizontally, which
976 matches with previous results (78)..
977
978 **Power analysis**
979
980 We conducted a power analysis to see what would happen if we had fewer strains in our
981 population genetic analysis.
982
We took advantage of the vcf-tools software , which allowed us to randomly remove strains
from our analysis. This enabled us to conduct a basic analysis on polymorphism for a groups
of N strains (N = 4,6,8,10,12,14,16,18,20,22,24,26,28), with 22 iterations of each number of
strains

Within any analysis, we focussed on the 3570 genes which are present in all strains, and
calculated mean and median polymorphism (average pairwise polymorphism, relative to gene
length) (Supplementary Figure S5.1).

Supplementary Figure S5.1: (A) Median polymorphism, (B) Mean polymorphism as the number of strains uses in the analysis varies. The line is a loess regression fit

The graph on the left shows median polymorphism, which shows a threshold effect once we
get to 16 strains, and also much smaller error bars. This shows that a smaller number of
strains will miss a lot of the diversity between strains.

The graph on the right shows mean polymorphism. The main pattern is that the standard error
in mean polymorphism declines substantially as the number of strains increase. This means
that the likelihood of getting a good estimate of the population mean increases as the number
of strain increases, which makes logical sense.

1001

1002 The fact that mean polymorphism doesn't change much, but median does, implies that the
1003 distribution has changed. The overall conclusion is that the number of strains we have
1004 included is likely appropriate to capture the true variation in the population

1018 **S6: Division of labour**

1019

1020 We conducted an analysis to see if the division of labour in the production of public goods in
1021 *B. subtilis* could explain the signatures of selection that we observe. For this, we look at the
1022 subset of 40 genes which are controlled by Spo0A, which is active only in a subset of cells
1023 (63). Depending on whether Spo0A is on or off, the expression of extracellular
1024 polysaccharides and amyloid protein fibres is either repressed or not (32, 61).
1025
1026 Spo0A controlled genes have lower median polymorphism than other quorum sensing -
1027 controlled genes, whether measured as overall polymorphism (0.0045 vs. 0.0101), non-
1028 synonymous polymorphism (0.0015 vs. 0.0037), or synonymous polymorphism (0.016 vs.
1029 0.021).
1030
1031 Spo0A controlled genes also have lower non-synonymous divergence (0.013 vs. 0.023) and
1032 lower ratio between non-synonymous and synonymous divergence (0.049 vs. 0.097),
1033 although they have slightly higher synonymous divergence (0.286 vs. 0.266).
1034
1035 The direction of selection statistic is very similar between the two groups (-0.044 vs. -0.037).
1036
1037 This pattern is the opposite to what we would expect if the division of labour was responsible
1038 for the signature of selection, rather than sociality *per se*. If the lower conditional expression
1039 was having a large effect, then we would expect these genes to have higher polymorphism
1040 than the background set of quorum sensing-controlled genes.
1041
1042 Further, EPS and TasA genes stand out within this class of genes as having high
1043 polymorphism, implying that the social effect is important in causing the signature of
1044 selection that we observe.

S7: Competence

The reference strain NCIB 3610 has a plasmid-encoded gene ComI, which interferes with the competence machinery (Konkol 2013). We know that there is variation in natural competence in our strains, as they were all screened for competency by (35). 18 out of the 31 strains we used are genetically competent.

Some of the competence genes (comGF, comGE, comGG) have extremely high polymorphism. Here, we conduct a small analysis where we restrict our analysis to only competent strains, to see if this polymorphism is caused by disuse in non-competent strains. We find that the competence operon comG is still amongst the most polymorphic even when we are only looking at competent strains.

Furthermore, cooperative genes are still significantly more polymorphic than private genes in the competent strains (Kruskal-Wallis test, chi-squared=14.7, df=2, p<0.0001).

Overall, we conclude that variation in natural competence isn't responsible for the difference between cooperative and private genes that we observe.

S8: Estimation of relatedness

According to theory, the degree to which selection is relaxed in cooperative genes relative to private genes is inversely proportional to relatedness. This result emerges from a simple population genetics model, which shows that a slightly deleterious allele with a cooperative effect on fitness will reach equilibrium frequency inversely proportional to the relationship between the actor and recipient (r) (19). If we assume weak selection and a large population (and ignore higher-order terms), we can directly map this prediction to relative levels of nucleotide polymorphism between alleles with cooperative and private effects on fitness.

As we noted in our previous work on *P. aeruginosa,* we have to make further assumptions that our set of cooperative genes experience the same average strength of selection and distribution of fitness effects as private genes, but this approach has the advantage of many experimental attempts to estimate relatedness for social interactions in that we don't need to know the scale at which interactions take place, or the relative weighting of environments in which traits are more or less social.

Cooperative genes have a median polymorphism of 0.01078, and private genes have median polymorphism of 0.00918. This leads to a calculation of relatedness as r=0.77.

We note that this measure might vary depending on how you define the population, which is tricky in bacteria due to horizontal gene transfer and other complications (131). In *B. subtilis*, for example, we know that strains from the same plant root or gram of soil can vary in their production of public goods, and don't always group together phylogenies (132–134).

S9: Correlation in gene expression for secondary comparisons

Here, we use the dataset from Futo *et al*. 2020 to test if the genes we used in the secondary comparisons also tend to be co-expressed. For each set of genes we first calculate the mean pairwise correlation between all possible pairs of genes. We then use a bootstrap approach of randomly sampling 10,000 other gene sets of the same size, to see if the gene set has higher correlated expression than expected by chance.

For all gene sets, we find that the correlation is greater than >95% of randomly sampled gene-sets (Supplementary Table S9.1)

Supplementary Table S9.1: Results from pairwise correlation in gene expression of a gene set, and bootstrap iterations of randomly sampled gene sets of the same size

Gene set	Correlation for gene set	Mean correlation for 10,000 iterations	Percentile of gene set
Iron scavenging	0.880	0.329	100
Antibiotic resistance	0.429	0.306	95.7
Proteases	0.397	0.293	95.3
Toxins	0.534	0.305	99.8
Antimicrobials	0.640	0.294	100

S10: Positive and balancing selection in other cooperative traits

We checked whether the other cooperative traits that we examined differ in signatures of positive and balancing selection, as this could cloud our conclusion on the nature of selection. We use Tajima's D to detect balancing selection, and neutrality index to detect positive selection.

Cooperative genes don't differ in balancing selection, whether we use all genes from the six comparisons (ANOVA $F_{2,121} = 7.50, p < 0.001$; Games Howell Test p=0.39), just the five secondary comparisons (ANOVA $F_{2,44} = 4.72, p = 0.14$; Games Howell Test p=0.33), or consider each gene as a data point (Wilcoxon signed-rank test V=7, p=0.563).

Cooperative genes also don't differ in positive selection, whether we use all genes from the six comparisons (Kruskal-Wallis $X^2(2) = 2.11, p = 0.35,$ Dunn Test $p = 0.70$), just the five secondary comparisons (Kruskal-Wallis $X^2(2) = 5.37, p = 0.07,$ Dunn Test $p = 0.98$), or consider each gene as a data point (Wilcoxon signed-rank test V=7, p=0.563).

Overall, we can conclude that the signature of selection we see across the five other cooperative traits are most consistent with kin selection causing the effective relaxation of selection

S11: Correlations in gene expression of QS-controlled genes

To test the robustness of our result that quorum sensing-controlled genes tend to be expressed at the same time, we repeated our analysis using the dataset from Pisithkul (37).

Whilst Futo *et al*. measured gene expression in a solid-air interface biofilm over two months (N=11 timepoints), Pisithkul *et al*. used a liquid-air interface, and measured gene expression over 24 hours (N=7 timepoints), representing the initial stages of biofilm growth.

Pisithkul *et al*. provided their data in the form of reads per kilobase of transcript per million mapped reads. For each gene, we normalised the data with the following steps;

1) We calculated median expression across the four replicates of each timepoint
2) We then divided each measure by the median expression at timepoint one (eight hours). This allows for better comparison between genes
3) We then used a log2 transformation to normalize the resulting relative expression measures

The average correlation for the N=160 quorum sensing controlled genes that we were able to match to the data was 0.411. The average correlation for N=10,000 randomly sampled gene sets of the same size (N=160) was 0.385. Quorum sensing-controlled genes have a higher correlation than 98.5% of the randomly sampled gene sets.

1311

1312 **Supplementary Figures**

Supplementary Figure 1: Ratio between non-synonymous and synonymous nucleotide diversity per site for private (yellow) and cooperative (blue) genes controlled by quorum sensing. Each point is a gene, and the horizontal line shows the median for each group. The grey line shows the median for background private genes across the genome.

1313

Supplementary Figure 2: Ratio between non-synonymous and synonymous divergence for private (yellow) and cooperative (blue) genes controlled by quorum sensing. Divergence is measured by rates of protein evolution, e.g. number of synonymous substitutions per synonymous site for panel B. Each point is a gene, and the horizontal line shows the median for each group. The grey line shows the median for background private

genes across the genome.

Supplementary Figure 3: Private (yellow) vs. cooperative (blue) non-synonymous polymorphism in genes for six traits. Panel A shows the quorum sensing -controlled genes used in the main analysis. Panels B-F show the secondary comparisons

1314

Supplementary Figure 4: Private (yellow) vs. cooperative (blue) ratio between non-synonymous and synonymous polymorphism in genes for six traits. Panel A shows the quorum sensing-controlled genes used in the main analysis. Panels B-F show the secondary

comparisons

1315

Supplementary Figure 5: Private (yellow) vs. cooperative (blue) non-synonymous divergence in genes for six traits. Panel A shows the quorum sensing -controlled genes used in the main analysis. Panels B-F show the secondary comparisons

1316

Supplementary Figure 6: Private (yellow) vs. cooperative (blue) synonymous divergence in

genes for six traits. Panel A shows the quorum sensing -controlled genes used in the main analysis. Panels B-F show the secondary comparisons

Supplementary Figure 7: Private (yellow) vs. cooperative (blue) ratio between non-synonymous and synonymous divergence in genes for six traits. Panel A shows the quorum sensing-controlled genes used in the main analysis. Panels B-F show the secondary comparisons

1338   **Supplementary Tables**

1339   **Supplementary Table 1**: List of strains used

Strain ID	Location	Assembly
NRS6096	Tayport (UK) garden soil	GCA_905311035.1
NRS6099	Tayport (UK) garden soil	GCA_905310985.1
NRS6103	Tayport (UK) community garden soil	GCA_905310995.1
NRS6105	Tayport (UK) community garden soil	GCA_905311425.1
NRS6107	Tayport (UK) garden soil	GCA_905311405.1
NRS6108	Tayport (UK) garden soil	GCA_905311395.1
NRS6110	Tayport (UK) vegetable plot	GCA_905311415.1
NRS6111	Tayport (UK) vegetable plot	GCA_905311375.1
NRS6116	Tayport (UK) garden soil	GCA_905311385.1
NRS6118	Tayport (UK) potato patch	GCA_905311435.1
NRS6120	Tayport (UK) potato patch	GCA_905312035.1
NRS6121	Tayport (UK) shrub bed	GCA_905315035.1
NRS6127	Tayport (UK) garden soil	GCA_905315045.1
NRS6128	Tayport (UK) garden soil	GCA_905315055.1
NRS6131	Tayport (UK) worm bin	GCA_905315025.1
NRS6134	Tayport (UK) vegetable patch	GCA_905315395.1
NRS6137	Tayport (UK) community garden soil	GCA_905315385.1
NRS6141	Tayport (UK) garden soil	GCA_905315375.1
NRS6145	Tayport (UK) garden soil	GCA_905315685.1
NRS6148	Tayport (UK) garden soil	GCA_905315695.1
NRS6153	Tayport (UK) garden soil	GCA_905315705.1
NRS6160	Tayport (UK) garden soil	GCA_905315715.1
NRS6167	Tayport (UK) soil from planter	GCA_905316385.1
NRS6181	Tayport (UK) garden soil	GCA_905318255.1
NRS6183	Lochee (UK) garden soil	GCA_905319155.1
NRS6186	Newport (UK) garden soil	GCA_905319135.1
NRS6190	Tayport (UK) garden soil	GCA_905319565.1
NRS6194	Tayport (UK) garden soil	GCA_905319555.1
NRS6202	Kirriemuir (UK) garden soil	GCA_905319535.1
NRS6205	Kirriemuir (UK) garden soil	GCA_905319825.1
NRS6206	Kirriemuir (UK) garden soil	GCA_905319815.1

1349   **Supplementary Table 2**: List of social genes

Gene	Function	Reference

bslA	biofilm-surface layer protein BslA	(33, 34)
bslB	biofilm-surface layer protein BslB	(135)
epsA	hypothetical protein	(33, 34)
epsB	protein tyrosine kinase EpsB	(33, 34)
epsC	polysaccharide biosynthesis protein	(33, 34)
epsD	glycosyltransferase family 4 protein	(33, 34)
epsE	glycosyltransferase EpsE	(33, 34)
epsF	glycosyltransferase family 1 protein	(33, 34)
epsG	biofilm exopolysaccharide biosynthesis protein EpsG	(33, 34)
epsH	glycosyltransferase	(33, 34)
epsI	polysaccharide pyruvyl transferase family protein	(33, 34)
epsJ	lipoprotein	(33, 34)
epsK	cyclic-di-GMP receptor EpsK	(33, 34)
epsL	sugar transferase	(33, 34)
epsM	acetyltransferase	(33, 34)
epsN	aminotransferase class I/II-fold pyridoxal phosphate-dependent enzyme	(33, 34)
epsO	polysaccharide pyruvyl transferase family protein	(33, 34)
sipW	signal peptidase I	(33, 34)
srfAA	surfactin non-ribosomal peptide synthetase SrfAA	(33, 34)
srfAB	surfactin non-ribosomal peptide synthetase SrfAB	(33, 34)
srfAC	surfactin non-ribosomal peptide synthetase SrfAC	(33, 34)
srfAD	surfactin biosynthesis thioesterase SrfAD	(33, 34)
srfT	MFS transporter	(33, 34)
tapA	amyloid fiber anchoring/assembly protein TapA	(33, 34)
tasA	biofilm matrix protein TasA	(33, 34)

1350 **Supplementary Table 3:** Cooperative and private genes for iron-scavenging via bacillibactin

Gene ID	Name	Function	Sociality	Reference
IRON SCAVENGING				
17270	dhbF	Bacillibactin biosynthesis	Cooperative	(65, 66)
17275	dhbB	Bacillibactin biosynthesis	Cooperative	(65, 66)
17280	dhbE	Bacillibactin biosynthesis	Cooperative	(65, 66)
17285	dhbC	Bacillibactin biosynthesis	Cooperative	(65, 66)
17290	dhbA	Bacillibactin biosynthesis	Cooperative	(65, 66)
1040	feuC	Membrane permease	Private	(65, 66)
1045	feuB	Membrane permease	Private	(65, 66)
1050	feuA	Periplasmic binding protein	Private	(65, 66)
17785	yusV [feuV]	ABC transporter	Private	(65, 66)
17295	besA	Ferri-bacillibactin esterase	Private	(65, 66)

1351

1352 **Supplementary Table 4:** Cooperative and private genes for antibiotic resistance

Gene ID	Name	Function	Sociality	Reference
ANTIBIOTIC RESISTANCE				
1295	blaOXA	Beta-lactamase	Cooperative	(67)

Gene ID	Name	Function	Sociality	Reference
	(ybxI)			
10225	bla (penP)	Beta-lactamase	Cooperative	(67)
14460	aadk	Aminoglycoside resistance	Cooperative	(68)
5445	bmrC	Multidrug ABC transporter	Private	(69)
5550	bmrD	Multidrug ABC transporter	Private	(69)
20835	cydC	Multidrug ABC transporter	Private	(69)
20840	cydD	Multidrug ABC transporter	Private	(69)
4595	yfiB	Multidrug ABC transporter	Private	(69)
4600	yfiC	Multidrug ABC transporter	Private	(69)
7900	yknU	Multidrug ABC transporter	Private	(69)
7905	yknV	Multidrug ABC transporter	Private	(69)
20050	ywjA	Multidrug ABC transporter	Private	(69)
18775	bmrA	Multidrug ABC transporter	Private	(69)
4830	ygaD	Multidrug ABC transporter	Private	(69)
11680	sunT	Sublancin transporter	Private	(69)

1353

Supplementary Table 5: Cooperative and private genes toxin genes

Gene ID	Name	Function	Sociality	Reference
		TOXINS		
20325	ywfA	Bacilycin biosysnthesis & export	Cooperative	(105)
20320	bacA	Bacilycin biosysnthesis & export	Cooperative	(105)
20315	bacB	Bacilycin biosysnthesis & export	Cooperative	(105)
20310	bacC	Bacilycin biosysnthesis & export	Cooperative	(105)
20305	bacD	Bacilycin biosysnthesis & export	Cooperative	(105)
20300	bacE	Bacilycin biosysnthesis & export	Cooperative	(105)
20295	bacF	Bacilycin biosysnthesis & export	Cooperative	(105)
20290	bacG	Bacilycin biosysnthesis & export	Cooperative	(105)
3890	yeeF	LXG toxin	Private	(75)
10360	yobL	LXG toxin	Private	(75)
11735	yokI	LXG toxin	Private	(75)
1830	yqcG	LXG toxin	Private	(75)
19495	ywqJ (rttN)	LXG toxin	Private	(75)
21125	yxiD	LXG toxin	Private	(75)

1355

Supplementary Table 6: Cooperative and private genes protease genes

Gene ID	Name	Function	Sociality	Reference
		PROTEASES		
5755	aprE	Extracellular protease	Cooperative	(70)
8405	Bpr	Extracellular protease	Cooperative	(70)
20660	Epr	Extracellular protease	Cooperative	(70)
1375	Mpr	Extracellular protease	Cooperative	(70)
6155	nprB	Extracellular protease	Cooperative	(70)
8100	nprE	Extracellular protease	Cooperative	(70)
20500	Upr	Extracellular protease	Cooperative	(70)
5990	wprA	Extracellular protease	Cooperative	(70)
9385	aprX	Intracellular protease	Private	(70)
7560	clpE	Intracellular protease	Private	(70)
18630	clpP	Intracellular protease	Private	(70)
8830	clpQ	Intracellular protease	Private	(70)

Gene ID	Name	Function	Sociality	Reference
13140	ispA	Intracellular protease	Private	(70)
15210	lonA	Intracellular protease	Private	(70)
15215	lonB	Intracellular protease	Private	(70)
9115	mlpA	Intracellular protease	Private	(70)
12000	ypwA	Intracellular protease	Private	(70)

1357

1358 **Supplementary Table 7:** Cooperative and private genes for the production of antimicrobials

Gene ID	Name	Function	Sociality	Reference
		PROTEASES		
9975	ppsA	Plipistatin biosynthesis	Cooperative	(73)
9970	ppsB	Plipistatin biosynthesis	Cooperative	(73)
9965	ppsC	Plipistatin biosynthesis	Cooperative	(73)
9960	ppsD	Plipistatin biosynthesis	Cooperative	(73)
9955	ppsE	Plipistatin biosynthesis	Cooperative	(73)
9295	pksB	Bacillaene biosynthesis	Private	(74)
9310	pksD	Bacillaene biosynthesis	Private	(74)
9315	pksE	Bacillaene biosynthesis	Private	(74)
9325	pksF	Bacillaene biosynthesis	Private	(74)
9330	pksG	Bacillaene biosynthesis	Private	(74)
9335	pksH	Bacillaene biosynthesis	Private	(74)
9340	pksI	Bacillaene biosynthesis	Private	(74)
9345	pksJ	Bacillaene biosynthesis	Private	(74)
9350	pksL	Bacillaene biosynthesis	Private	(74)
9355	pksM	Bacillaene biosynthesis	Private	(74)
9360	pksN	Bacillaene biosynthesis	Private	(74)
9365	pksR	Bacillaene biosynthesis	Private	(74)
9370	pksS	Bacillaene biosynthesis	Private	(74)

References

1.   J. E. Strassmann, O. M. Gilbert, D. C. Queller, Kin discrimination and cooperation in
microbes. *Annu. Rev. Microbiol.* **65**, 349–367 (2011).
2.   S. A. West, A. S. Griffin, A. Gardner, S. P. Diggle, Social evolution theory for
microorganisms. *Nat Rev Microbiol* **4**, 597–607 (2006).
3.   A. S. Griffin, S. A. West, A. Buckling, Cooperation and competition in pathogenic
bacteria. *Nature* **430**, 1024–1027 (2004).
4.   L. McNally, M. Viana, S. P. Brown, Cooperative secretions facilitate host range
expansion in bacteria. *Nat. Commun.* **5** (2014).
5.   L. A. Dugatkin, M. Perlin, J. S. Lucas, R. Atlas, Group-beneficial traits, frequency-
dependent selection and genotypic diversity: An antibiotic resistance paradigm. *Proc.*
R. Soc. B Biol. Sci. **272**, 79–83 (2005).
6.   M. Ghoul, A. S. Griffin, S. A. West, Toward an evolutionary definition of cheating.
Evolution (N. Y). **68**, 318–331 (2014).
7.   W. D. W. D. Hamilton, The Genetical Evolution of Social Behaviour. II. *J. Theor.*
Biol. **7**, 17–52 (1964).
8.   R. Kümmerli, A. S. Griffin, S. A. West, A. Buckling, F. Harrison, Viscous medium
promotes cooperation in the pathogenic bacterium Pseudomonas aeruginosa. *Proc. R.*
Soc. B Biol. Sci. **276**, 3531–3538 (2009).
9.   S. P. Diggle, A. S. Griffin, G. S. Campbell, S. A. West, Cooperation and conflict in
quorum-sensing bacterial populations. *Nature* **450**, 411–414 (2007).
10.   J. E. Strassmann, Y. Zhu, D. C. Queller, Altruism and social cheating in the social
amoeba Dictyostelium discoideum. *Nature* **408**, 965–967 (2000).
11.   G. J. Velicer, L. Kroos, R. E. Lenski, Developmental cheating in the social bacterium
Myxococcus xanthus. *Nature* **404**, 598–601 (2000).
12.   M. Ghoul, S. B. Andersen, S. A. West, Sociomics: Using Omic Approaches to
Understand Social Evolution. *Trends Genet.* **33**, 408–419 (2017).
13.   O. X. Cordero, L. -a. L. A. Ventouras, E. F. DeLong, M. F. Polz, Public good
dynamics drive evolution of iron acquisition strategies in natural bacterioplankton
populations. *Proc. Natl. Acad. Sci.* **109**, 20059–20064 (2012).
14.   E. Butaite, M. Baumgartner, S. Wyder, R. Kümmerli, Siderophore cheating and
cheating resistance shape competition for iron in soil and freshwater Pseudomonas
communities. *Nat. Commun.* **8**, 414 (2017).
15.   S. B. Andersen, *et al.*, Long-term social dynamics drive loss of function in pathogenic
bacteria. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 10756–10761 (2015).
16.   T. Jautzus, J. van Gestel, Á. T. Kovács, Complex extracellular biology drives surface
competition during colony expansion in Bacillus subtilis. *ISME J.* (2022)
https:/doi.org/10.1038/s41396-022-01279-8.
17.   R. M. Fisher, C. K. Cornwallis, S. A. West, Group formation, relatedness, and the
evolution of multicellularity. *Curr. Biol.* **23**, 1120–1125 (2013).
18.   C. Simonet, L. McNally, Kin selection explains the evolution of cooperation in the gut
microbiota. *Proc. Natl. Acad. Sci.* (2021) https:/doi.org/10.1073/pnas.2016046118.
19.   T. A. Linksvayer, M. J. Wade, Genes with social effects are expected to harbor more
sequence variation within and between species. *Evolution (N. Y).* **63**, 1685–1696
(2009).
20.   T. A. Linksvayer, M. J. Wade, Theoretical predictions for sociogenomic data: The
effects of kin selection and sex-limited expression on the evolution of social insect
genomes. *Front. Ecol. Evol.* **4**, 1–10 (2016).
21.   J. D. Van Dyken, M. J. Wade, The genetic signature of conditional expression.
Genetics **184**, 557–570 (2010).
22.   J. D. van Dyken, M. J. Wade, Detecting the molecular signature of social conflict:

Theory and a test with bacterial quorum sensing genes. *Am. Nat.* **179**, 436–450 (2012).

23.   J. D. Van Dyken, T. A. Linksvayer, M. J. Wade, Kin selection-mutation balance: A
model for the origin, maintenance, and consequences of social cheating. *Am. Nat.* **177**,
288–300 (2011).

24.   E. A. Ostrowski, *et al.*, Genomic Signatures of Cooperation and Conflict in the Social
Amoeba. *Curr. Biol.* **25**, 1661–5 (2015).

25.   S. Noh, K. S. Geist, X. Tian, J. E. Strassmann, D. C. Queller, Genetic signatures of
microbial altruism and cheating in social amoebas in the wild. *Proc. Natl. Acad. Sci.*
U. S. A. **115**, 3096–3101 (2018).

26.   J. L. de Oliveira, *et al.*, Conditional expression explains molecular evolution of social
genes in a microbe. *Nat. Commun.* **10**, 3284 (2019).

27.   O. M. Gilbert, K. R. Foster, N. J. Mehdiabadi, J. E. Strassmann, D. C. Queller, High
relatedness maintains multicellular cooperation in a social amoeba by controlling
cheater mutants. *Proc. Natl. Acad. Sci.* **104**, 8913–8917 (2007).

28.   L. J. Belcher, A. E. Dewar, M. Ghoul, S. A. West, Kin selection for cooperation in
natural bacterial populations. *Proc. Natl. Acad. Sci. U. S. A.* **119** (2022).

29.   M. W. Hahn, *Molecular Population Genetics* (OUP USA, 2018).

30.   N. A. Logan, P. De Vos, "Bacillus" in *Bergey's Manual of Systematics of Archaea and*
Bacteria, F. A. R. and W. B. W. M.E. Trujillo, S. Dedysh, P. DeVos, B. Hedlund, P.
Kämpfer, Ed. (2015) https:/doi.org/10.1002/9781118960608.gbm00530.

31.   J. W. Veening, *et al.*, Bet-hedging and epigenetic inheritance in bacterial cell
development. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 4393–4398 (2008).

32.   A. Dragoš, *et al.*, Division of Labor during Biofilm Matrix Production. *Curr. Biol.* **28**,
1903-1913.e5 (2018).

33.   S. Arnaouteli, N. C. Bamford, N. R. Stanley-Wall, Á. T. Kovács, Bacillus subtilis
biofilm formation and social interactions. *Nat. Rev. Microbiol.* **19**, 600–614 (2021).

34.   M. Kalamara, M. Spacapan, I. Mandic-Mulec, N. R. Stanley-Wall, Social behaviours
by Bacillus subtilis: quorum sensing, kin discrimination and beyond. *Mol. Microbiol.*
110, 863–878 (2018).

35.   M. Kalamara, J. C. Abbott, C. E. Macphee, N. R. Stanley-Wall, Biofilm
hydrophobicity in environmental isolates of Bacillus subtilis. *Microbiol. (United*
Kingdom) **167** (2021).

36.   M. Futo, *et al.*, Embryo-Like Features in Developing Bacillus subtilis Biofilms. *Mol.*
Biol. Evol. **38**, 31–47 (2021).

37.   T. Pisithkul, *et al.*, Metabolic remodeling during biofilm development of bacillus
subtilis. *MBio* **10** (2019).

38.   M. Špacapan, *et al.*, The ComX Quorum Sensing Peptide of Bacillus subtilis Affects
Biofilm Formation Negatively and Sporulation Positively. *Microorg. 2020, Vol. 8,*
Page 1131 **8**, 1131 (2020).

39.   M. M. Nakano, L. Xia, P. Zuber, Transcription initiation region of the srfA operon,
which is controlled by the comP-comA signal transduction system in Bacillus subtilis.
J. Bacteriol. **173**, 5487 (1991).

40.   N. Comella, A. D. Grossman, Conservation of genes and processes controlled by the
quorum response in bacteria: Characterization of genes controlled by the quorum-
sensing transcription factor ComA in Bacillus subtilis. *Mol. Microbiol.* **57**, 1159–1174
(2005).

41.   D. López, R. Kolter, Extracellular signals that define distinct and coexisting cell fates
in Bacillus subtilis. *FEMS Microbiol. Rev.* **34**, 134–149 (2010).

42.   S. S. Branda, *et al.*, Fruiting body formation by Bacillus subtilis Steven. *Proc. Natl.*
Acad. Sci. **98**, 11621–11626 (2001).

43.   S. S. Branda, *et al.*, Genes Involved in Formation of Structured Multicellular
1483   Communities by Bacillus subtilis. *J. Bacteriol.* **186**, 3970 (2004).
1484   44.   J. Van Gestel, F. J. Weissing, O. P. Kuipers, Á. T. Kovács, Density of founder cells
1485   affects spatial pattern formation and cooperation in Bacillus subtilis biofilms. *ISME J.*
1486   **8**, 2069–2079 (2014).
1487   45.   D. Romero, C. Aguilar, R. Losick, R. Kolter, Amyloid fibers provide structural
1488   integrity to Bacillus subtilis biofilms. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 2230–2234
1489   (2010).
1490   46.   E. Erskine, C. E. MacPhee, N. R. Stanley-Wall, Functional Amyloid and Other Protein
1491   Fibers in the Biofilm Matrix. *J. Mol. Biol.* **430**, 3642 (2018).
1492   47.   S. S. Branda, F. Chu, D. B. Kearns, R. Losick, R. Kolter, A major protein component
1493   of the Bacillus subtilis biofilm matrix. *Mol. Microbiol.* **59**, 1229–1238 (2006).
1494   48.   K. I. Yoshida, Y. Fujita, S. D. Ehrlich, Three Asparagine Synthetase Genes of Bacillus
1495   subtilis. *J. Bacteriol.* **181**, 6081 (1999).
1496   49.   S. Azimi, A. D. Klementiev, M. Whiteley, S. P. Diggle, Bacterial Quorum Sensing
1497   during Infection. *Annu. Rev. Microbiol.* (2020) https:/doi.org/10.1146/annurev-micro-
1498   032020-093845.
1499   50.   S. T. Rutherford, B. L. Bassler, Bacterial quorum sensing: Its role in virulence and
1500   possibilities for its control. *Cold Spring Harb. Perspect. Med.* **2**, 1–25 (2012).
1501   51.   M. P. Zwart, *et al.*, Unraveling the causes of adaptive benefits of synonymous
1502   mutations in TEM-1 β-lactamase. *Heredity (Edinb).* **121**, 406–421 (2018).
1503   52.   M. Meir, *et al.*, Competition between social cheater viruses is driven by
1504   mechanistically different cheating strategies. *Sci. Adv.* **6** (2020).
1505   53.   S. Azimi, *et al.*, Allelic polymorphism shapes community function in evolving
1506   Pseudomonas aeruginosa populations. *ISME J.* (2020) https:/doi.org/10.1038/s41396-
1507   020-0652-0.
1508   54.   M. L. M. Salverda, J. A. G. M. de Visser, M. Barlow, Natural evolution of TEM-1 β-
1509   lactamase: Experimental reconstruction and clinical relevance. *FEMS Microbiol. Rev.*
1510   **34**, 1015–1036 (2010).
1511   55.   M. Whiteley, S. P. Diggle, E. P. Greenberg, Progress in and promise of bacterial
1512   quorum sensing research. *Nature* **551**, 313–320 (2017).
1513   56.   M. Schuster, D. J. Sexton1, B. A. Hense, Why quorum sensing controls private goods.
1514   *Front. Microbiol.* **8**, 1–16 (2017).
1515   57.   G. A. Cooper, S. A. West, Division of labour and the evolution of extreme
1516   specialization. *Nat. Ecol. Evol.* **2**, 1161–1167 (2018).
1517   58.   G. A. Cooper, H. Frost, M. Liu, S. A. West, The consequences of group structure and
1518   efficiency benefits for the evolution of division of labour. *Elife* **10** (2021).
1519   59.   S. A. West, G. A. Cooper, Division of labour in microorganisms: an evolutionary
1520   perspective. *Nat. Rev. Microbiol. 2016 1411* **14**, 716–723 (2016).
1521   60.   M. Liu, S. A. West, G. A. Cooper, Relatedness and the evolution of mechanisms to
1522   divide labor in microorganisms. *Ecol. Evol.* **11**, 14475–14489 (2021).
1523   61.   Y. Chai, F. Chu, R. Kolter, R. Losick, Bistability and biofilm formation in Bacillus
1524   subtilis. *Mol. Microbiol.* **67**, 254–263 (2008).
1525   62.   J. E. González-Pastor, Cannibalism: a social behavior in sporulating Bacillus subtilis.
1526   *FEMS Microbiol. Rev.* **35**, 415–424 (2011).
1527   63.   V. Molle, *et al.*, The Spo0A regulon of Bacillus subtilis. *Mol. Microbiol.* **50**, 1683–
1528   1701 (2003).
1529   64.   K. Kobayashi, SlrR/SlrA controls the initiation of biofilm formation in Bacillus
1530   subtilis. *Mol. Microbiol.* **69**, 1399–1410 (2008).
65.   M. Miethke, *et al.*, Ferri-bacillibactin uptake and hydrolysis in Bacillus subtilis. *Mol.*

Microbiol. **61**, 1413–1427 (2006).

66.   H. Pi, J. D. Helmann, Sequential induction of Fur-regulated genes in response to iron
limitation in Bacillus subtilis. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 12785–12790
(2017).

67.   T. Bucher, *et al.*, An active β-lactamase is a part of an orchestrated cell wall stress
resistance network of Bacillus subtilis and related rhizosphere species. *Environ.*
Microbiol. **21**, 1068–1085 (2019).

68.   N. Noguchi, M. Sasatsu, M. Kono, "Genetic mapping in Bacillus subtilis 168 of the
aadK gene which encodes arninoglycoside 6-adenylyltransferase" (1993).

69.   C. Torres, C. Galián, C. Freiberg, J. R. Fantino, J. M. Jault, The YheI/YheH
heterodimer from Bacillus subtilis is a multidrug ABC transporter. *Biochim. Biophys.*
Acta **1788**, 615–622 (2009).

70.   C. R. Harwood, Y. Kikuchi, The ins and outs of Bacillus proteases: Activities,
functions and commercial significance. *FEMS Microbiol. Rev.* **46**, 1–20 (2022).

71.   B. M. Koo, *et al.*, Construction and Analysis of Two Genome-Scale Deletion Libraries
for Bacillus subtilis. *Cell Syst.* **4**, 291-305.e7 (2017).

72.   S. Pohl, *et al.*, Proteomic analysis of Bacillus subtilis strains engineered for improved
production of heterologous proteins. *Proteomics* **13**, 3298–3308 (2013).

73.   D. Romero, *et al.*, The Iturin and Fengycin Families of Lipopeptides Are Key Factors
in Antagonism of Bacillus subtilis Toward Podosphaera fusca. *MPMI* **20**, 430–440
(2007).

74.   S. Müller, *et al.*, Bacillaene and Sporulation Protect Bacillus subtilis from Predation by
Myxococcus xanthus. *Appl. Environ. Microbiol.* **80**, 5603 (2014).

75.   K. Kobayashi, Diverse LXG toxin and antitoxin systems specifically mediate
intraspecies competition in Bacillus subtilis biofilms. *PLOS Genet.* **17**, e1009682
(2021).

76.   A. Mariappan, O. Makarewicz, X. H. Chen, R. Borriss, Two-Component Response
Regulator DegU Controls the Expression of Bacilysin in Plant-Growth-Promoting
Bacterium Bacillus amyloliquefaciens FZB42. *Microb. Physiol.* **22**, 114–125 (2012).

77.   N. Y. Yu, *et al.*, PSORTb 3.0: Improved protein subcellular localization prediction
with refined localization subcategories and predictive capabilities for all prokaryotes.
Bioinformatics **26**, 1608–1615 (2010).

78.   A. Dewar, *et al.*, Plasmids do not consistently stabilize cooperation across bacteria but
may promote broad pathogen host-range. *Nat. Ecol. Evol.* **5**, 1624–1636 (2021).

79.   T. Nogueira, *et al.*, Horizontal Gene Transfer of the Secretome Drives the Evolution of
Bacterial Cooperation and Virulence. *Curr. Biol.* **19**, 1683–1691 (2009).

80.   J. M. Flowers, *et al.*, Variation, Sex, and Social Cooperation: Molecular Population
Genetics of the Social Amoeba Dictyostelium discoideum. *PLoS Genet.* **6**, e1001013
(2010).

81.   O. M. Gilbert, J. E. Strassmann, D. C. Queller, High relatedness in a social amoeba:
The role of kin-discriminatory segregation. *Proc. R. Soc. B Biol. Sci.* **279**, 2619–2624
(2012).

82.   S. Nee, S. A. West, A. F. Read, Inbreeding and parasite sex ratios. *Proc. R. Soc. B*
Biol. Sci. **269**, 755–760 (2002).

83.   B. Kraigher, M. Butolen, P. Stefanic, I. Mandic Mulec, Kin discrimination drives
territorial exclusion during Bacillus subtilis swarming and restrains exploitation of
surfactin. *ISME J. 2021 163* **16**, 833–841 (2021).

84.   M. Martin, *et al.*, Cheaters shape the evolution of phenotypic heterogeneity in Bacillus
subtilis biofilms. *ISME J. 2020 149* **14**, 2302–2312 (2020).

85.   N. A. Lyons, B. Kraigher, P. Stefanic, I. Mandic-Mulec, R. Kolter, A Combinatorial

Kin Discrimination System in Bacillus subtilis. *Curr. Biol.*, 1–10 (2016).

86. P. Stefanic, B. Kraigher, N. A. Lyons, R. Kolter, I. Mandic-Mulec, Kin discrimination
between sympatric Bacillus subtilis isolates. *Proc. Natl. Acad. Sci. U. S. A.* **112**,
14042–14047 (2015).

87. S. A. West, K. Winzer, A. Gardner, S. P. Diggle, Quorum sensing and the confusion
about diffusion. *Trends Microbiol.* **20**, 586–594 (2012).

88. J. B. Xavier, W. Kim, K. R. Foster, A molecular mechanism that stabilizes cooperative
secretions in Pseudomonas aeruginosa. *Mol. Microbiol.* **79**, 166–179 (2011).

89. D. J. Sexton, M. Schuster, Nutrient limitation determines the fitness of cheaters in
bacterial siderophore cooperation. *Nat. Commun. 2017 81* **8**, 1–8 (2017).

90. B. D. Connelly, E. L. Bruger, P. K. McKinley, C. M. Waters, Resource abundance and
the critical transition to cooperation. *J. Evol. Biol.* **30**, 750–761 (2017).

91. N. K. M. Tam, *et al.*, The intestinal life cycle of Bacillus subtilis and close relatives. *J.*
Bacteriol. **188**, 2692–2700 (2006).

92. L. Fan, *et al.*, Genome sequence of Bacillus subtilis subsp. spizizenii gtP20b, isolated
from the Indian ocean. *J. Bacteriol.* **193**, 1276–1277 (2011).

93. M. Roberts, F. Cohan, Recombination and Migration Rates in Natural Populations of
Bacillus Subtilis and Bacillus Mojavensis. *Evolution (N. Y).* **49**, 1081–1094 (1995).

94. P. H. Brito, *et al.*, Genetic competence drives genome diversity in bacillus subtilis.
Genome Biol. Evol. **10**, 108–124 (2018).

95. K. E. Duncan, N. Ferguson, K. Kimura, X. Zhou, C. A. Istock, Fine-Scale Genetic and
Phenotypic Structure in Natural Populations of Bacillus subtilis and Bacillus
licheniformis: Implications for Bacterial Evolution and Speciation. *Evolution (N. Y).*
48, 2002 (1994).

96. K. Kobayashi, Gradual activation of the response regulator DegU controls serial
expression of genes for flagellum formation and biofilm formation in Bacillus subtilis.
Mol. Microbiol. **66**, 395–409 (2007).

97. R. M. Berka, *et al.*, Microarray analysis of the Bacillus subtilis K-state: genome-wide
expression changes dependent on ComK. *Mol. Microbiol.* **43**, 1331–1345 (2002).

98. M. A. Konkol, K. M. Blair, D. B. Kearns, Plasmid-encoded ComI inhibits competence
in the ancestral 3610 strain of Bacillus subtilis. *J. Bacteriol.* **195**, 4085–4093 (2013).

99. S. Arnaouteli, *et al.*, Bifunctionality of a biofilm matrix protein controlled by redox
state. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E6184–E6191 (2017).

100. A. Ostrowski, A. Mehert, A. Prescott, T. B. Kiley, N. R. Stanley-Wall, YuaB functions
synergistically with the exopolysaccharide and TasA amyloid fibers to allow biofilm
formation by Bacillus subtilis. *J. Bacteriol.* **193**, 4821–4831 (2011).

101. R. Kümmerli, *et al.*, Co-evolutionary dynamics between public good producers and
cheats in the bacterium Pseudomonas aeruginosa. *J. Evol. Biol.* **28**, 2264–2274 (2015).

102. I. Frost, *et al.*, Cooperation, competition and antibiotic resistance in bacterial colonies.
ISME J. **12**, 1582–1593 (2018).

103. K. Poole, Aminoglycoside resistance in Pseudomonas aeruginosa. *Antimicrob. Agents*
Chemother. **49**, 479–487 (2005).

104. Y. Quentin, G. Fichant, F. Denizot, Inventory, assembly and analysis of Bacillus
subtilis ABC transport systems. *J. Mol. Biol.* **287**, 467–484 (1999).

105. O. Ertekin, *et al.*, Analysis of a bac operon-silenced strain suggests pleiotropic effects
of bacilysin in Bacillus subtilis. *J. Microbiol.* **58**, 297–313 (2020).

106. P. Cingolani, *et al.*, A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff. *Fly (Austin).* (2012)
https:/doi.org/10.4161/fly.19695.

107. R, R: A language and environment for statistical computing. *R Found. Stat. Comput.*

Vienna, Austria. (2020).

108.   H. Wickham, *ggplot2 Elegant Graphics for Data Analysis* (2016).

109.   S. Andrews, FastQC. *Babraham Bioinforma.* (2010).

110.   H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* (2009) https:/doi.org/10.1093/bioinformatics/btp324.

111.   H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics*
(2009) https:/doi.org/10.1093/bioinformatics/btp352.

112.   Broad Institute, Picard toolkit. *Broad Institute, GitHub Repos.* (2019).

113.   P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *Gigascience* **10**, 1–4
(2021).

114.   A. R. Quinlan, I. M. Hall, BEDTools: A flexible suite of utilities for comparing
genomic features. *Bioinformatics* (2010) https:/doi.org/10.1093/bioinformatics/btq033.

115.   B. Pfeifer, U. Wittelsbürger, S. E. Ramos-Onsins, M. J. Lercher, PopGenome: An
efficient swiss army knife for population genomic analyses in R. *Mol. Biol. Evol.* **31**,
1929–1936 (2014).

116.   A. McKenna, *et al.*, The genome analysis toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res.* (2010)
https:/doi.org/10.1101/gr.107524.110.

117.   D. Charif, J. R. Lobry, "SeqinR 1.0-2: a contributed package to the R project for
statistical computing devoted to biological sequences retrieval and analysis." in
Structural Approaches to Sequence Evolution: Molecules, Networks, Populations, U.
Bastolla, M. Porto, H. Roman, M. Vendruscolo, Eds. (Springer Verlag, 2007)
https:/doi.org/10.1007/978-3-540-35306-5_10.

118.   F. Hildebrand, A. Meyer, A. Eyre-Walker, Evidence of selection upon genomic GC-
content in bacteria. *PLoS Genet.* **6** (2010).

119.   P. M. Sharp, E. Bailes, R. J. Grocock, J. F. Peden, R. E. Sockett, Variation in the
strength of selected codon usage bias among bacteria. *Nucleic Acids Res.* **33**, 1141–
1153 (2005).

120.   H. Akashi, T. Gojobori, Metabolic efficiency and amino acid composition in the
proteomes of Escherichia coli and Bacillus subtilis. *Proc. Natl. Acad. Sci. U. S. A.* **99**,
3695–3700 (2002).

121.   R. Raghavan, Y. D. Kelkar, H. Ochman, A selective force favoring increased G+C
content in bacterial genes. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 14504–14507 (2012).

122.   E. P. C. Rocha, Codon usage bias from tRNA's point of view: Redundancy,
specialization, and efficient decoding for translation optimization. *Genome Res.* **14**,
2279–2286 (2004).

123.   Y. Sun, S. G. E. Andersson, SSCU: an R/Bioconductor package for analyzing selective
profile in synonymous codon usage (2018).

124.   A. Ross-Gillespie, *et al.*, Frequency Dependence and Cooperation: Theory and a Test
with Bacteria. *Am. Nat.* **170**, 331–342 (2007).

125.   J. Gore, H. Youk, A. van Oudenaarden, Snowdrift game dynamics and facultative
cheating in yeast. *Nature* **459**, 253–256 (2009).

126.   P. G. Madgwick, L. J. Belcher, J. B. Wolf, Greenbeard Genes : Theory and Reality.
Trends Ecol. Evol., 1–12 (2019).

127.   F. Tajima, Statistical method for testing the neutral mutation hypothesis by DNA
polymorphism. *Genetics* (1989).

128.   E. Paradis, Pegas: An R package for population genetics with an integrated-modular
approach. *Bioinformatics* **26**, 419–420 (2010).

129.   Y. X. Fu, W. H. Li, Statistical tests of neutrality of mutations. *Genetics* (1993)
https:/doi.org/10.1093/genetics/133.3.693.

130.   N. Stoletzki, A. Eyre-Walker, Estimation of the neutrality index. *Mol. Biol. Evol.* **28**,
1683   63–70 (2011).
1684   131.   E. P. C. Rocha, Neutral theory, microbial practice: Challenges in bacterial population
1685   genetics. *Mol. Biol. Evol.* **35**, 1338–1347 (2018).
1686   132.   A. Oslizlo, *et al.*, Exploring ComQXPA quorum-sensing diversity and biocontrol
1687   potential of Bacillus spp. isolates from tomato rhizoplane (2015)
1688   https:/doi.org/10.1111/1751-7915.12258 (November 7, 2022).
1689   133.   H. T. Kiesewalter, *et al.*, Genomic and Chemical Diversity of Bacillus subtilis
1690   Secondary Metabolites against Plant Pathogenic Fungi. *mSystems* **6** (2021).
1691   134.   K. Steinke, O. S. Mohite, T. Weber, Á. T. Kovács, Phylogenetic Distribution of
1692   Secondary Metabolites in the Bacillus subtilis Species Complex. *mSystems* **6** (2021).
1693   135.   R. J. Morris, *et al.*, Natural variations in the biofilm-associated protein BslA from the
1694   genus Bacillus. *Sci. Rep.* **7** (2017).
1695