SOX9-positive pituitary stem cells differ according to their position in the gland and

maintenance of their progeny depends on context.

135 characters

Running Title: Stem cells and adaptative response in the pituitary.

Karine Rizzoti[a,1], Probir Chakravarty[b], Daniel Sheridan[a], and Robin Lovell-Badge[a,1].

[a] Laboratory of Stem Cell Biology and Developmental Genetics, The Francis Crick Institute,

London NW1 1AT, UK

[b] Bioinformatics core, The Francis Crick Institute, London NW1 1AT, UK

[1] authors for correspondence karine.rizzoti@crick.ac.uk and robin.lovell-badge@crick.ac.uk

ABSTRACT
19
20
Stem cell (SC) differentiation and maintenance of resultant progeny underlie cell-turnover in
many organs, but it is difficult to pinpoint the contribution of either process. In the pituitary, a
central regulator of endocrine axes, adult SCs undergo activation following target organ
ablation, providing a well-characterized paradigm to study an adaptive response in a multi-
organ system. Here we used single cell technologies to characterize SC heterogeneity and
mobilization together with lineage tracing. We show that SC differentiation occurs more
frequently than thought previously. In adaptive conditions, differentiation increases and is
more diverse than demonstrated by the lineage tracing experiments. Detailed examination of
SC progeny suggests that maintenance of selected nascent cells underlies SC output,
highlighting a trophic role for the microenvironment. Analyses of cell trajectories further
predict pathways and potential new regulators. Our model provides a valuable system to
study the influence of evolving states on the mechanisms of SC mobilization.
33
34
35   Teaser: Pituitary stem cells are diverse and differentiate more than thought but only selected
36   progeny persist according to need.
37
38

INTRODUCTION

The pituitary gland controls the three major endocrine axes, reproduction, stress and
metabolism, and is involved in homeostasis; all under control of the hypothalamus. Its
function requires modulation of hormonal output over short and long terms, and this is
regulated at several levels, including rates and patterns of hormone secretion, synthesis and
storage, and changes in endocrine cell numbers. Different mechanisms are known to underlie
endocrine cell emergence: division of endocrine cells, differentiation from resident SCs and
transdifferentiation, although the relevance of the latter is unclear. Under normal and stable
conditions, it is thought that division from endocrine cell types is sufficient for the typical
low cell turnover seen in the pituitary (*1*). This has been directly shown in corticotrophs using
a genetic trick where cell division induces apoptosis of POMC expressing cells. A gradual
loss of corticotrophs is observed in these pituitaries in support of the role of these endocrine
cells in their own replacement (*2*). In agreement, long term lineage tracing of adult SOX2 and
SOX9 +ve pituitary SCs in steady-state conditions suggests that SCs differentiate rarely and
thus do not significantly participate in cell turnover (*3*) (*4*). Furthermore, adult ablation of the
SCs does not appear to affect pituitary function (*5*). In contrast, when the regenerative or
adaptative potential of the gland is stimulated, either by endocrine cell ablation (*6*) or target
organ ablation (*3*) respectively, SC-derived new endocrine cells of the appropriate type
emerge. In this latter paradigm, differentiation from SCs appears to be the main process
underlying emergence of new corticotrophs after adrenalectomy (Ax) (*2*), providing a well
characterized multi-organ system to explore mechanisms of SC activation.

Here, we explore the characteristics of pituitary SC fate acquisition using single cell
approaches combined with lineage tracing analyses. We observe that SOX9 +ve pituitary SCs
comprise several functionally distinct sub-populations, located in the ventral and dorsal part
of the epithelium lining the pituitary cleft and in the anterior pituitary parenchyma. We then
explored mechanisms underlying SC mobilization following target organ ablation through
characterizing cell trajectories and gene regulatory networks (GRNs) potentially involved in
cell fate acquisition. We observed that while fate acquisition of the targeted cell type
(corticotrophs after Ax and gonadotrophs after gonadectomy) was efficiently induced,
differentiation toward all endocrine cell types unexpectedly also took place. Examination of
longer-term outcomes using lineage tracing further showed that only a proportion of the
differentiated progeny is retained, specifically that of the targeted cell type. More precisely,
we only observe SC-derived corticotrophs after Ax, and lactotrophs in females, following
Selective Estrogen Receptor Modulator (SERM) treatment. Therefore, while pituitary SC
differentiation appears both more frequent and diverse than previously thought, our results
suggest that the selective maintenance of nascent cells, according to perceived
pathophysiological need and likely sex-dependent hormonal cues, is an important regulatory
step controlling their adaptive response. Our model thus allows characterisation of this
response following pathophysiological manipulations that may well relate to normal life-
changing events, such as puberty, pregnancy and lactation.

82
83
84 RESULTS
85
86 <u>Distinct regulatory pathways govern AL and IL SC identity</u>
87
The anterior lobes of three $Sox9^{iresGFP/+}$ adult male pituitaries were dissociated and the
transcriptome of single GFP +ve cells analysed after fluorescence activated cell sorting
(FACS), as previously described (*3*). After quality checks, 1993 cells were retained for
further analysis; clustering was performed by generating UMAP plots. Three major clusters
were identified (Fig.1A). Co-expression of *Sox2* and *Sox9* in clusters 0 and 2 (Fig.1B,
Sup.Fig.1), representing 69% of the cells, suggested that these were likely to represent SCs.
The presence of two SC clusters had previously been reported in whole pituitary datasets (*7*)
(*8*). The remaining cluster (1), representing 24% of the dataset, comprised cells where *Sox*9
was detected, but where *Sox2* was greatly reduced or absent. Finally, cells each expressing a
different hormone and the corresponding lineage specifying factors, were grouped within a
single smaller cluster (3) representing 3% of the cells (Fig.1B, Sup.Fig.1). SOX9 is not
expressed in any endocrine cell in quiescent conditions (*9*). However, we observed some rare
SOX9;ACTH +ve cells, and more numerous GFP;ACTH +ve cells in $Sox9^{iresGFP/+}$ animals
exclusively after activation of the SCs by pituitary target organ ablation (*3*) (Sup. Fig.2). This
suggests that differentiating, and recently differentiated cells are present in the SOX9iresGFP
fraction due to persistence of the GFP, which correlates with what has been suggested to
happen in both human and mouse pituitaries, where hormone positive cells were included in
the SC compartment of whole pituitary single cell RNAseq datasets (*10*). In addition, a small
proportion of contaminating pituitary cells (estimated to represent 25% of cluster 3) is likely
to be present (Sup.Fig.3). Therefore, most cells present in cluster 3 are likely to represent
cells committed toward an endocrine fate, with the largest population (41% of cluster 3 cells)
expressing *Prolactin*.
110
111

Figure 1

Figure 1: Single cell analysis of the $Sox9^{iresGFP/+}$ population in male pituitaries.

A) UMAP clustering of $Sox9^{iresGFP/+}$ cells.

B) Expression levels of *Sox2* and *Sox9* were examined to identify SCs (*9*) while genes

encoding for hormones (*Gh*, *Prl*, *Tshβ*, *Lhβ*, *Fshβ*, *Pomc*) were examined to distinguish

committed cells (*Pou1f1* for somatotrophs, lactotrophs and thyrotrophs.

C) Heatmap displaying selected markers for the main clusters.

The two SC clusters were further explored. Analysis of differentially expressed genes (DEG)

(Fig.1C, Sup.Fig.1B) shows that clusters 0 and 2 are relatively similar and are mostly

distinguished by differences in levels of expression of specific genes. One notable exception

is *Aqp3*, encoding for a water channel, which is specific for cluster 0. In contrast, *Aldh1a2*,

encoding for an enzyme involved in retinoic acid (RA) synthesis, and the transcription factor

Msx1 were preferentially present in cluster 0, while *Ednrb*, encoding for the endothelin B

receptor, and the cell adhesion molecule *Claudin3* were both expressed at higher levels in

cluster 2. RA is required during pituitary morphogenesis (*11*) (*12*) and *Msx1* interacts with

SOX2 for *Prop1* regulation (*13*), while nothing is known about *Ednrb* or *Claudin3* in

pituitary SCs. Conservation of most of these markers in SCs from whole pituitary datasets (*7*)

(*8*) suggests that our two SC clusters correlate with those observed by others. In the gland,

expression of the five markers is restricted to SCs (Fig.2, Sup. Fig.4). In the epithelium lining

the cleft, cluster 0 markers ALDH1a2 and MSX1 are expressed exclusively on the AL side

(Fig.2 A,, Sup.Fig4B), while cluster 2 markers EDNRB and CLAUDIN-3 (Fig.2 C,D) are

present on both sides. In the parenchyma ALDH1a2, MSX1 and CLAUDIN-3 co-localise
with SOX2/9 while expression of EDNRB is not detectable. In situ hybridisation for *Ednrb*
confirms a clear enrichment in cleft lining cells (Sup.Fig.5). AQP3, whose transcript is
present in about half of cluster 0 cells, is exclusively detected in the parenchymal SCs,
revealing further heterogeneity between cleft-lining and parenchymal populations. In
addition, it is only expressed in a subset of SOX2;SOX9 positive parenchymal SCs (Fig.2B),
demonstrating the existence of parenchymal subpopulations. Examination of the five markers
in scRNAseq datasets from whole pituitaries (21) confirms that the strongest, and sometimes
exclusive, expression is seen in SCs (Sup. Fig.4). All together, these results suggest that
cluster 0 is enriched in anterior lobe SCs, while cluster 2 has a more generic pituitary SC
identity.
Pathway analysis in cluster 0 (Sup Table.1) highlighted activity of the NOTCH and WNT
pathways and presence of Ephrin signalling, as previously reported in pituitary SCs (*14*) (*15*).
Cluster 2 cells are characterised by expression of cytoskeleton remodelling and adhesion
genes consistent with the EMT-like process required during pituitary progenitor cell fate
acquisition (*16*). In conclusion, regionalisation of these two SC subpopulations is underlined
by the expression of genes involved in the activity of specific regulatory pathways, such as
retinoic acid signalling in AL SCs. Further investigation is required to understand what
distinguishes AQP3 positive and negative parenchymal SCs.

Folliculo-stellate (FS) cells are specialised AL glial cells (*17*). These express S100b in the
rat, but not in the mouse ((*18, 19*), Sup. Fig.6). To clarify their relationship with SCs, we
examined expression of *Aldolase C*, a marker for rodent FS cells (*19*) and of *Slc152a*, which
encodes a transporter responsible for the FS specific import of AMCA (*20*). Both are
selectively present in SCs, in clusters 0 and 2 (Sup.Fig.6A). Furthermore, in a whole pituitary
dataset, *Sox2* and *Aldoc* are exclusively present in the SC cluster (Sup.Fig.6B). All together
these results confirm, as suggested earlier (*9, 21*) (*22, 23*), that in the mouse FS and SCs are
overlapping populations.

In conclusion, single-cell analysis of the $Sox9^{iresGFP/+}$ pituitary fraction successfully resolved
heterogeneity of this population, suggesting differential activity of signalling pathways, and
clarified the relationship between SCs and FS cells.

-

Figure 2

Figure 2: Differential expression of SC cluster markers.

A) Co-immunofluorescence for ALDH1A2, GFP and SOX2 in a male SOX9iresGFP

pituitary section. ALDH1A2 is exclusively expressed in SOX2;SOX9 AL SC (yellow arrow

shows its absence from IL SC, and white arrow shows its expression in parenchymal SC).

Aldh1a2 violin plot shows enrichment in cluster 0.

B) Co-immunofluorescence for AQP3, GFP and SOX2 in a male SOX9iresGFP pituitary

section. AQP3 is expressed in 66% of SOX2 +ve parenchymal SC (white arrow shows

SOX2;SOX9iresGFP parenchymal SC not expressing AQP3; counting was performed on n=3

animals). *Aqp3* violin plot shows enrichment in cluster 0.

C) Co-immunofluorescence for CLAUDIN3, GFP and SOX2 in a male SOX9iresGFP

pituitary section. CLAUDIN3 is exclusively expressed in SOX2;SOX9 SC. *Cldn3* UMAP

shows enrichment in cluster 2.

D) Co-immunofluorescence for EDNRB and GFP in a male SOX9iresGFP pituitary section.

EDNRB is exclusively expressed in SOX9 cleft SC. *Ednrb* UMAP shows enrichment in

cluster 2. Similar expression profiles were observed in females for all markers.

The scale bar in A represents 10 μm for all panels. White arrows show absence of expression

in parenchymal SC. AL, anterior lobe, IL intermediate lobe.

<u>A novel population of parenchymal SOX9;PDGFRβ +ve cells</u>

Cluster 1, representing 25% of the cells, is characterised by expression of PDGFRβ (Fig.1C,

Fig.3E). *In situ*, PDGFRβ;SOX9 +ve cells are exclusively detected in the parenchyma.

Expression of SOX9 in these double +ve cells is weaker than in SCs (Fig.1B, Fig.3A) and

SOX2 is not detected (Fig.1B , Fig.3B). In contrast with the *in silico* analysis, only 8±1.62%

SD of parenchymal SOX9 +ve cells express PDGFRβ in males (Fig.3C). This discrepancy

could be due either to an over-representation of cluster 1 cells, because epithelial SCs are

more difficult to recover than parenchymal cells, or to the difficulty in detecting the cells by

immunofluorescence. We observe significantly fewer PDGFRβ;SOX9 +ve cells in females

than in males (1.3±0.6% SD, p= 0.029, Fig.3C). PDGFRβ +ve pituitary cells are known to

comprise a mural vascular network (*18*). While we did not attempt to fully characterize the

SOX9;PDGFRβ population, we observe some SOX9;PDGFRβ cells in close contact with

capillaries, displaying a pericyte like morphology (Fig.3D). In the embryo, vascular and

mural cells have an extra-Rathke's pouch origin, as they emerge from immigrating neural

crest cells (*24, 25*). Because cluster 1 cells do not express the generic pituitary marker *Pitx1*

(Fig.3E), we analysed a potential neural crest origin by performing WNT1-Cre lineage

tracing. 79 ± 3.1% SEM of SOX9;PDGFRβ cells (n=130 cells from 4 males) derive from

WNT1-Cre +ve cells (Fig3C, F) while in females only 47.5 ± 5.5% SEM of the cells derive

from the neural crest (n= 57 cells from 3 females, p= 0.024). Therefore, we propose that

cluster 1 comprises a sexually dimorphic mural network with a dual embryonic origin. The

pituitary capillary network, which controls endocrine output, adapts according to the

physiological context and this is particularly important for gonadotroph function (*26*). The

sex differences we observe may be related to this aspect.

<u>Figure 3: Characterisation of a SOX9;PDGFRβ +ve AL cell population.</u>

A) Co-immunofluorescence for PDGFRβ, GFP and SOX9 in a SOX9iresGFP male AL

section. Triple +ve cells are indicated (arrows). The lower panels represent the magnification

of the region indicated on the merge panel above. The dotted line indicates the pituitary cleft.

B) Co-immunofluorescence for PDGFRβ, GFP and SOX2 in AL. A SOX9;PDGFRβ +ve cell

which is negative for SOX2 (arrow).

C) Percentage of GFP;SOX9;PDGFRβ/GFP;SOX9 +ve cells in *Sox9iresGFP* male and

female AL sections (average 500 SOX9 cells counted/animal, p = 0.029). In SOX9iresGFP

pituitary sections, parenchymal SOX9; GFP were captured (Sup.Table.4). Amongst these, the

percentage of those also positive for PDGFRβ cells was determined. Percentage of

SOX9;PDGFRβ +ve cells of neural crest origin (p = 0.024; Mann Whitney test). In

Wnt1Cre;R26ReYFP pituitary sections, Sox9;PDGFRb cells were captured. Amongst these,

the percentage of those also positive for Wnt1Cre;ReYFP cells was determined

(Sup.Table.4).

D) Co-immunofluorescence for PDGFRβ, GFP and PECAM in a Sox9iresGFP AL section

showing a SOX9;PDGFRβ +ve cell resembling a pericyte bridging PECAM +ve capillaries.

E) UMAP for *Pitx1* and *Pdgfrβ* showing that *Pdgfrβ* +ve cluster 1 cells do not express *Pitx1*.

F) Co-immunofluorescence for PDGFRβ, eYFP and SOX9 in a *Wnt1-Cre;R26R^(eYFP)* AL

section. Triple +ve cells are shown (arrows).

The scale bars represent 50 μm and 20 μm for A (upper and lower panels respectively) and

10 μm for D and F.

<u>All endocrine cell types are present in the $Sox9^{iresGFP/+}$ +ve compartment after SC</u>
<u>mobilization.</u>

We then performed sc-RNAseq on activated SCs. Target organ ablation induces a transient
proliferative wave in the gland, mostly affecting SCs, which can then give rise to new
endocrine cells (*2, 3, 27*). We analysed the transcriptome of $Sox9^{iresGFP/+}$ male cells 4 days
after removing the adrenals, the testes or both. We integrated these datasets with that
obtained from unchallenged animals and obtained a similar clustering of SCs (Fig.1,
Fig.4A,B, Sup. Fig.7A). A pairwise comparison for proportion test limited to SC clusters
demonstrates a significant increase in proliferative cells after target organ ablation
(Sup.Fig.7B), as previously shown (*3*). We also observe the emergence of distinct hormone
+ve clusters representing in total 25 to 30% of all cells (Fig.4A,B) compared to 3.5% in the
unchallenged dataset, suggesting increased differentiation. Target organ ablation is associated
with increased secretion, and sometimes cell numbers, of the endocrine population regulating
the resected organ. However, regardless of the glands removed, all endocrine cell lineages
emerge in the Sox9iresGFP +ve fraction. While we expected *Pomc* +ve cells to prominently
feature after Ax, these were outnumbered by *Prl* and *LHb;FSHb* +ve cells (Sup. Fig. 7C).
Similarly, gonadectomies were expected to trigger commitment toward the gonadotroph cell
fate, but *Gh* +ve cells were mostly observed. To characterise these $Sox9^{iresGFP/+}$;hormone +ve
cells we compared them with endocrine cells by integrating our datasets with one from whole
pituitaries (*28*). $Sox9^{iresGFP/+}$;hormone +ve cells cluster with endocrine cells (Fig.4C, D).
Therefore, they represent differentiated cells, at least transcriptionally. In conclusion, target
organ ablation induces differentiation of SCs, but there is an unexpected lack of specificity in
the endocrine cell types produced, with all hormonal markers being up-regulated.

Figure 4: Target organ ablation induces differentiation toward multiple endocrine lineages.
A) UMAP clustering for integrated datasets from sorted *Sox9*^{iresGFP/+} cells from unchallenged, adrenalectomized (1915 cells), gonadectomized (1546 cells) and both adrenalectomized and gonadectomized (1880 cells) animals, 4 days after surgery. B) UMAP clustering split by experiment showing the contribution of cells originating from each dataset. Cells expressing all pituitary hormones are present (dotted lines) in all conditions. C) UMAP clustering for integrated datasets from sorted *Sox9*^{iresGFP/+} cells and a whole pituitary dataset (*28*). D) UMAP clustering split by experiment show that hormone +ve cells present in the *Sox9*^{iresGFP/+} cells cluster with endocrine cells (black arrow pointing to lactotrophs).

Analyses of cell trajectories after Ax.

We decided to focus on Ax because it represents the most efficient stimulus of SC differentiation after target organ ablation (*3*). We added a new time point at day 2 after Ax

and a sham control. These new datasets were then integrated (Fig.5A) with our initial dataset
(Fig.1) and the one performed 4 days after Ax (Fig.4). UMAP clustering indicated that SCs
were present in four distinct clusters (Sup. Fig.4). In addition to the *Ednrb* and *Aldh1a2*
clusters, two SC clusters (SC3 and SC4) emerged (Fig.5A). These were mostly but not
exclusively populated by cells from the new datasets (Sup. Fig.8). While more cells were
sequenced in these new datasets (up to 3 times more), we used a different chemistry (see
material and methods) which, despite Seurat batch correction tools, may explain detection of
more genes and cells resulting in emergence of these clusters. SC3 was characterised by
higher expression of markers also present in the *Ednrb* and *Aldh1a2* clusters. SC4, mostly
derived from the Ax+48h00 dataset, showed co-expression of SC and endocrine markers
suggesting that it represents cells in a transitioning phase (Sup. Fig.9). Finally, endocrine cell
clusters were detected (Sup. Fig.9), as observed previously (Fig.4).
In sham operated animals, 4.5% of hormone expressing cells, mostly *Prl* +ve, were present,
confirming our observations in unchallenged animals (Sup. Table 2). 48 hours after Ax, the
proportion of hormone +ve cells had increased to 11%, and this had again doubled after 4
days. Because Sox9iresGFP expression stops as cells differentiate the increasing proportion
means that commitment toward differentiation augments during the 4 days following
ablation. A pairwise proportion test between the 2 new datasets, sorted identically on the
same day, showed that a shift occurs from SC4; while the proportion of cells from each
dataset in the 3 SC clusters were comparable, from SC4 the proportion of cells from the Ax
dataset becomes significantly increased compared to the control, fitting with its transitioning
phase profile (Fig.5B, Sup. Fig.9). At this shorter time point, 48 h00 after adrenal ablation,
corticotrophs are the most significantly increased cell type, while 4 days after surgery they
were outnumbered by lactotrophs. This sharp and transient increase could represent the
specific SC response to adrenals ablation.
We then performed trajectory analyses after principal component analysis (PCA) using
Slingshot (*29*) using exclusively Ax+48h and Sham Ax+48h datasets. Trajectories finishing
at each endocrine cell type were discovered, with a common root through SC clusters *Ednrb*,
Aldh1a2, proliferative cells and SC4, then ending at each endocrine cell type (Fig.5C, Sup.
Fig.10 and 11). In the corticotroph trajectory (Fig.5D), the expression pattern of 2245 genes
significantly correlates with pseudotime (Fig.5F). The expression of known SC markers
(*Ednrb*, *Sox2*, *Sox9*, *Aldh1a2*) and signalling pathway members (*Notch2*, *Hes1*, *Hey1* and
Wnt4, *Fzd2*, *Sfrp5*) was gradually lost. Pathway analyses further suggest keratin filament
remodelling, in agreement with epithelial to mesenchymal transition during differentiation.
Ismn1 (*30*) and *Prop1*(*16*), which are involved in cell fate acquisition, are also present during
early phases. Conversely, the corticotroph specifier and marker genes *Neurod1*, *Ascl1*, *Tbx19*,
Pomc and *Creb3l2* (*31*) are upregulated as cells differentiate. A sharp transition occurs as
Prop1 is upregulated and is characterised by a transient up-regulation of *Pou1f1* lineage and
gonadotroph (*Fshb*) genes. Comparisons with trajectories from other lineages reveal similar
trajectories with a transient up-regulation of unrelated lineages markers (Sup.Fig.11). This
could reflect permissive chromatin remodelling as endocrine cell fate acquisition occurs.

SCENIC analyses were then performed to explore GRNs (*32*) and compare them between the
different endocrine lineages (Fig.5G, H, Sup.Fig.11). As expected *Sox2* and *9* regulons

featured in SC clusters while *Ascl1*, *Neurod1* and *Creb3l2* were present in corticotrophs. The
thyroid (*Thra*) and the androgen (*Ar*) hormone receptor regulons featured in corticotroph-
differentiating SCs, suggesting that SCs can respond to peripheral signals. New pituitary
GNRs were also discovered, such as those involving *Runx1* and *Irf9* whose expression was
validated in SCs (Fig.5E, G, H). RUNX1 (RUNt-related transcription tactor 1) is known to
control adult skin SC maintenance by regulating WNT signalling (*33*) and is shared between
the corticotroph and gonadotroph trajectories. IRFs (Interferon Regulatory Factors) are
modulators of cell growth and differentiation beside their anti-viral functions. IRF9
associates with phosphorylated STAT1:STAT2 (Signal Transducer and Activator of
Transcription) dimers, and all are present in our four endocrine SCENIC analyses (*34*).
Finally, several RA factors were also present in all trajectories, supporting RA involvement
during endocrine cell differentiation (Fig.5H).

In conclusion, by adding new datasets, we validated our previous observations suggesting a
low differentiation rate of SCs, mostly into lactotrophs in the intact and sham operated
animal. We refined the dynamics of the adaptative response after mobilization, showing a
transient peak of corticotroph differentiation 48 hours after Ax. Finally, we were able to infer
endocrine cell trajectories revealing common and exclusive factors during each endocrine cell
fate acquisition.
336
337

Figure 5

Figure 5: Trajectory analyses following Ax.

A) UMAP clustering for integrated datasets from sorted GFP +ve cells from unchallenged,

adrenalectomized and sham + 48 h, and + 4 days, all from *Sox9iresGFP/+* pituitaries. The data

has been filtered to remove mural and vascular cells and outliers. Cluster colour scheme is

constant throughout the figure. B) Pairwise comparison for proportion test performed on

adrenalectomized and sham + 48h. The proportion of corticotrophs after Ax is significantly
different from all other cluster distributions. The proportion of cells in cluster SC4 is
significantly different from all other SC cluster distributions (Sup. Table2). C) Representation
of endocrine cell trajectories on the PCA plot for integrated dataset. D) Proportion of
cells/cluster in the corticotroph trajectory comparing Ax and sham Ax. Differentiation is
occurring predominantly after Ax. E) Immunofluorescence on pituitary sections validates
RUNX1 and IRF9 expression in SOX2 +ve SCs. RUNX1 is also expressed in F4.80 +ve
macrophages. F) Heatmap of 2245 genes associated with the corticotroph trajectory. G)
Heatmap of SCENIC analysis on the corticotroph trajectory. H) Venn diagram of the
common and unique regulons between the different endocrine trajectories (Sup.Fig. 11).
354

<u>Selective maintenance of SC progeny</u>

To examine predictions from our transcriptomic analyses, we performed two different types
of lineage tracing experiment. First, we directly assessed expression of hormonal markers on
plated $Sox9^{iresGFP/+}$ FACSorted single cells by immunofluorescence in males and females, 4
days after Ax or sham-surgery. Second, to follow mobilized SCs further, we used $Sox9^{ires-}$
$^{CreERT2/+}$; $R26R^{EYFP/+}$ animals and examined progeny 10 to 11 days after Cre induction (a
week after surgery since induction is mostly performed for 3 days before surgery).

All hormone +ve cell types were found in the GFP +ve fraction in both experimental and
control $Sox9^{iresGFP/+}$ animals, which matches our transcriptomic analyses (Fig.6 A, B).
Following Ax, the number of POMC +ve cells increased the most, in agreement with our
earliest dataset (Ax+ 48h). However, even if less efficiently stimulated after Ax, the
proportion of most other endocrine cell types was comparable to that of corticotrophs. This
again reflected our transcriptomic data. More precisely, we also observed a slight increase in
somatotroph emergence in both sexes, while lactotrophs and to a reduced extent rare
gonadotrophs appear to increase only in females.

We then carried out lineage tracing in $Sox9^{ires-CreERT2/+}$; $R26R^{EYFP/+}$ animals (Fig.6C). We
examined the endocrine cell types predicted to be mostly induced by the single cell analysis
(Sup.Fig.3C) plus corticotrophs, since we knew these are present in the SC progeny after Ax
(3) . In contrast with our previous results (Fig.6A), examination of pituitary sections
exclusively revealed the presence of POMC;eYFP +ve corticotrophs after Ax, in comparable
numbers in both sexes (Fig.6C). In females, we additionally observed lactotrophs, and did so
even without performing target organ ablation. Quantification of lactotroph emergence on
dissociated pituitaries confirmed the sex bias (Fig.6D). Correcting for CreERT2 induction
efficiency, we estimate that in females SCs contribute to 0.6% of the total population of
lactotrophs and these SC-derived lactotrophs represent 2% of the eYFP +ve population
(Fig.5F).

The two genetic tools we used, $Sox9^{iresGFP}$ and $Sox9^{ires-CreERT2/+}$, display inherent distinctions,
such as use of high-dose of the SERM tamoxifen, and purposes, where these may impair
direct comparisons. However, while results from both transcriptomic analyses and
immunofluorescence demonstrate that endocrine cells of all types are produced in
comparable numbers in the short term, we clearly observe that corticotrophs are the only

endocrine cells remaining in long-term lineage tracing experiments in males. This indicates a
selective maintenance of nascent endocrine cells.

Alternatively, hormone positive cells could acquire expression of SOX9 after surgeries,
leading to the presence of SOX9;hormone +ve cells in the SOX9iresGFP fraction but not in
the SOX9CreERT2 animals because tamoxifen was given before SC mobilization. There is
no supporting evidence for this to happen, while we know that SCs differentiate (SOX9+ve
cells become hormone +ve). Furthermore, tamoxifen is known to persist in mice so we
should still be able to see these cells in our lineage tracing experiments even if SOX9
becomes up-regulated after the surgeries (*35*). Finally, this discrepancy in the presence of
hormone+ve cells in short-term versus long-term lineage tracing already exists in the intact
animal because we see these cells our unchallenged and sham single cell RNAseq
experiment; but ours and others' lineage tracing analyses did not reveal any significant
differentiation in normal conditions (*3, 4*). Therefore, we think that the most likely
explanation is the selective maintenance of a proportion of the SC progeny.

To further explore the sex bias in SC-derived lactotrophs, we administered tamoxifen to P0
Sox9$^{ires-CreERT2/+}$; R26R$^{EYFP/+}$ pups and quantified differentiation into lactotrophs once they
reached adulthood. There was no significant difference in the proportion of SC-derived
lactotrophs between males and females (Fig.5E), as previously observed (*3*). Therefore, male
SC-derived lactotrophs can be produced, but this is temporally restricted. Furthermore,
because the proportion of lactotrophs generated from male and female pups is similar, this
suggests that tamoxifen, which can interfere with estrogen action, induces lactotroph
emergence. If lactotroph emergence was a continuous process in females their proportion
would have been far superior to that observed in males. Our results thus reveal a temporal
window of sensitivity in males, while female SCs remain responsive to the lactotroph
promoting effect of tamoxifen until adulthood. It is unclear how E2 signalling is altered, and
whether SC differentiation is induced or maintenance of nascent lactotrophs permitted.
Sexual dimorphism in SC activity has also been reported in another endocrine organ, the
adrenals, where androgens regulate cell-turnover (*36*).
414

Figure 6

<u>Figure 6: Analyses of scRNAseq predictions by short and long-term lineage tracing.</u>

A) Immunofluorescence for hormonal markers on FACsorted Sox9iresGFP +ve cells from a
female 4 days after Ax.

B) Percentages of FACSorted Sox9iresGFP +ve cells that are hormone +ve /DAPI. Cell
counts were performed on plated cells (shown in A). Pituitaries were harvested 4 days after
surgeries. Pairwise for proportion tests were performed to assess significance of differences
between Ax and Sham (Sup. Table 3).

C) Numbers of hormone;eYFP double +ve cells counted in $Sox9^{iresCreERT2/+};Rosa26^{ReYFP}$
pituitaries a week after surgeries or 10 days after tamoxifen treatment (no surgery). Counts
were performed on comparable samples of whole pituitary sections. Corticotrophs are
observed in both sexes after Ax, as previously reported. In contrast, significant numbers of
lactotrophs are exclusively observed in females. LH;FSH cells have never been observed
after gonadectomies (3) and were thus not quantified here.

D) Immunofluorescence for eYFP and PRL on dissociated non-sorted
$Sox9^{iresCreERT2/+};Rosa26^{ReYFP}$ female pituitary cells. A PRL;eYFP +ve cell is indicated
(arrow).

E) Estimation of the percentage of lactotrophs differentiated from SCs. eYFP;PRL +ve cells
from $Sox9^{iresCreERT2};Rosa26^{ReYFP}$ dissociated pituitaries were counted 10 days after tamoxifen
treatment. Numbers were corrected for Cre induction efficiency. Induction efficiency was
calculated in each animal as the percentage of SOX9;eYFP/SOX9 +ve cells present in the
cleft, which was preserved from dissociation. There is very little proliferation in normal
conditions in the cleft SCs, and we thus assumed that the number of eYFP;SOX9 positive
cells directly reflected induction efficiency. 0.6% of lactotrophs originate from SCs 10 days
after tamoxifen induction in females while we barely observed any in males (Mann-Whitney
test performed on angular transformation of percentages, p=0.0159).

F) Percentage of PRL;eYFP/eYFP +ve cells counted 10 days after tamoxifen treatment in 8
to 10 week-old animals, and in 8 week-old animals that were treated with tamoxifen as pups
(eYFP;PRL/eYFP in $Sox9^{iresCreERT2/+};Rosa26^{ReYFP}$). The proportion of SCs becoming
lactotrophs is significantly higher in adult females compared to males (t-test, p=0.03). This
sex bias disappears when tamoxifen is administered in pups (t-test test, p=0.1825). Scale bars
represent 10 microns.

<u>HGF trophic effect on pituispheres suggests that SCs and corticotrophs interact.</u>

The selective maintenance of specific nascent SC progeny, while the rest are lost, suggests
that survival is dependent on local cues in their microenvironment. In the context of Ax, the
corticotrophs are a likely source of signals. We thus re-examined our transcriptomic datasets,
screening for signalling pathways potentially activated in SCs by corticotrophs. We found
that the receptor tyrosine kinase Met is selectively upregulated in SCs while its ligand Hgf is
present both in corticotrophs and pericytes (Fig.7A-C, Sup.Fig 12 (37)), suggesting
interactions between these cell types. Furthermore, following Ax, the number of SCs
expressing Met is increased, exclusively in those proliferating (Fig.7D). This suggests a
potential trophic effect of HGF on SCs that we explored in pituisphere assays (Fig.7E, F).
Upon HGF treatment, the number of pituispheres is on average doubled compared to the

control, demonstrating a trophic effect on SCs. Because the number of corticotrophs is
increased after Ax and levels of HGF may thus potentially increase too, this pathway may
mediate or at least participate in the increased SC proliferation observed after Ax. In
agreement with a trophic role of the pathway in the pituitary, the receptor and its ligand are
both widely expressed in human pituitary adenomas (*38*).
465
466

467
Figure 7: *Met*/MET and *Hgf* are expressed in SCs and corticotrophs respectively and the
ligand has a trophic effect on pituispheres.
A) Expression levels of *Met* are shown in clusters belonging to the corticotroph trajectory
(Fig.5D). B) Immunofluorescence for MET and SOX9 showing the receptor is expressed in
SCs. C) In situ hybridisation (RNAscope) and immunofluorescence for *Hgf* and POMC
respectively showing that MET ligand is expressed in corticotrophs. *Hgf* is also predicted to
be expressed in pericytes which could explain signal outside corticotrophs. D) More cells
express *Met* after Ax specifically in the cluster comprising proliferating cells (Chi-squared
test, p= 0.0005, Sup.Table 3). E) More pituispheres form when 100ng/ml HGF is added to
basic medium (t-test, p= 0.002, Sup. Table 3, n=4). F) MET is expressed in SOX9 +ve
pituispheres. Scale bar represents 20 microns in B and D, and 10 microns in C.
479
480
DISCUSSION

While adult pituitary SCs appear superfluous for cell-turnover in the unchallenged animal,
their differentiation potential can be stimulated, for example when the function of certain
endocrine axes is compromised, such as when adrenals or gonads are ablated (*3*) (*2, 5*). Here
we have characterized SC heterogeneity and the modalities of endocrine cell emergence, by
examining Sox9iresGFP single cells. We observe that a small percentage of pituitary SCs
differentiate in the unchallenged animal, which correlates with single-cell RNAseq analyses
performed on both human and mouse pituitaries where the presence of differentiating cells
was also detected in the SC compartment (*10*). This proportion increases when SCs are
mobilized. The paradigm we applied for SC mobilization, target organ ablation, is known to
trigger a specific response in the pituitary: adrenal ablation results in increased ACTH while
gonadectomy is associated with increased gonadotropins. However, although there was a
significant induction of corticotroph cell fate acquisition shortly after ablation, we
unexpectedly observed increased differentiation toward all AL endocrine cell types.
Nevertheless, in the longer term, lineage tracing experiments show that it is exclusively the
targeted endocrine cell type that is maintained in the SC progeny. Therefore, following
differentiation, selective maintenance of specific nascent cells is likely to be an important
process regulating pituitary SC endocrine cell output, and it seems likely that this is
controlled by the SC microenvironment. Increased sensitivity of nascent pituitary cells to
apoptosis had been suggested before (*27*), and shown to increase after target organ ablation,
in agreement with our results. Selective fitness of surviving cells in the embryo relies on
access to particular trophic factors (*39*). In our Ax paradigm, the pituitary targets of the
ablation, the corticotrophs, represent a likely source of signals, a notion re-enforced by the
trophic effect of HGF we observe on pituispheres. Similarly, estrogen is known to promote
lactotroph proliferation and their emergence in embryos (*40*). We previously examined the
effects of estrogen on adult pituitary SC differentiation, but the prior administration of
tamoxifen in our lineage tracing system may explain why we did not obtain meaningful
results((*3*). Here, our data suggest that tamoxifen treatment induces emergence of lactotrophs,
both in male and female neonates, but exclusively in adult females. This in turn implies that
this process is sensitive to E2, and that male development suppresses it. The presence of
lactotrophs in the male SOX9iresGFP fraction, but not after lineage tracing, suggests that
their maintenance is promoted in females. However,to further investigate these aspects a
tamoxifen independent lineage tracing tool is required. In parallel with the likely role of
corticotrophs on SCs after Ax, lactotrophs may be involved in the selective maintenance of
nascent PRL cells emerging after tamoxifen treatment, because they are sensitive to its effect,
given that prolactin secretion is increased by the SERM (*41*). In addition to the reliance on
trophic factors, it is likely that nascent cell maintenance involves integration into homotypic
endocrine networks (*42*). This is reminiscent of what is known for nascent neurons that are
often produced in excess and whose persistence relies on synaptic connections (*43*). Although
focusing on SCs enabled us to uncover novel regulators and unexpected modalities of the
adaptative response, our analyses point toward an important role for the microenvironment,
and, in our multi-organ system, peripheric factors which are predicted to modulate the SC
response. The thyroid (*Thra*) and the androgen (*Ar*) hormone receptors appear active in
differentiating SCs, suggesting that they can respond to peripheral signals, and may integrate

this information to regulate pituitary plasticity, as recently proposed (*44*). Further
investigations of both local and peripheral interactions are now required to better understand
how this adaptative response is regulated in detail.

Differential clustering of SOX9iresGFP +ve SCs, trajectory analyses and *in situ* validation of
new markers revealed the activity of region-specific regulatory pathways. In agreement with
what had been shown by others (*7*) (*8*), we found two main populations. We show here that
these correlate with differential localisation, anterior lobe (ALDH1a2, MSX1 positive) versus
more generic AL/IL SC (CLDN3 positive) identity. This suggests that AL and IL SCs are
regulated differently. We speculate that RA may be involved in AL SC differentiation, as
shown in the embryo (*11*). The downregulation of ALDH1a2 expression in IL SCs which do
not differentiate (*2, 45*) provides an additional support to this hypothesis. The enrichment of
EDNRb in the cells lining the cleft reveals a further level of heterogeneity between cleft
lining and parenchymal cells. In the trajectory analyses, the upstream expression pattern of
Ednrb implies that the SCs lining the cleft are in a more immature state than those in the
parenchyma, which fits with this epithelium constituting the primary pituitary SC niche.
Among the 3 endothelin ligands, *Edn3* is the most abundantly expressed in the gland, more
precisely in SCs and pericytes (*37*). This suggests that this pathway could function in an
autocrine or paracrine way in cleft SCs, because these are not in close contact with capillaries
(*42*), which would make an interaction with the latter unlikely. EDNRB signalling has been
shown to regulate melanocyte SC regeneration (*46*) and we are currently investigating
whether this role is conserved in pituitary SCs. We have furthermore found that a
subpopulation of parenchymal SCs is defined by AQP3 expression. It is unclear whether this
correlates with the existence of functionally different subpopulations, such as supporting FS
cells versus SCs for example, or whether this reflects distinct SC states, but we are now able
to identify these cells, and investigate their role. Finally, the presence of RUNX, IFR and
STAT regulons during cell differentiation is also of potential interest because of the roles of
these factors in different SC contexts, and because they have been shown to interact
(RUNX/STAT and IFR/STAT). STAT transcription factors are activated by different
cytokines and growth factors, in particular IL6 whose role in the regenerative response of
pituitary SCs was recently uncovered (*7*).

Our differentiation trajectories suggest the presence of a common root for all endocrine cell
types, which only segregate as the terminal differentiation markers are upregulated in the
nascent cells. This process is likely to be fast since cells still contain GFP expressed from
Sox9iresGFP. ~~It is now important to determine whether the same genes in this common root~~
~~are associated with the different lineages and when specificity first appears~~. It is tempting to
draw a parallel with embryonic pituitary progenitors, because these were similarly shown to
all commit at the same stages (*47*). Interestingly, it was proposed that intrinsic factors and
cell communication within the developing pituitary underlie specific endocrine cell fate
acquisition, which, in a different context and with different modalities and outcomes, is what
we hypothesize is happening in our model.

In conclusion, evaluation of single cell transcriptomic predictions with short and long-term
lineage tracing experiments has proved a robust strategy revealing novel aspects of pituitary
SC adaptative responses. In circumstances where an exceptional pathophysiological need is

perceived, nascent cells of the appropriate type are maintained. Understanding how the new
cells can survive will help us to understand how SC input is controlled in the pituitary, but
this could also be relevant in other tissues where SC activity appears largely superfluous.
Finally, it is notoriously difficult to obtain mature cell types *in vitro* for use in regenerative
medicine. Therefore, identifying trophic factors for nascent cells, which may be all that can
be obtained for transplantation, appears necessary.

575
576

Acknowledgements:

We are grateful for the support from present and past members of the Lovell-Badge lab. We
are indebted to the Francis Crick Institute platforms for their excellent support and expertise.
We especially thank Jacek Mor from the Biological Research facility for taking excellent care
of the animals, Donald Bell from the Advanced Light microscopy, Deborah Schneider-
Luftman from Bioinformatics and Biostatistics and experts from the Advanced Sequencing
and Histology platforms. We also thank Frederique Ruf-Zamojski and Zidong Zhang from
the Icahn School of Medicine at Mount Sinai (New York) for sharing their expertise.
This work was supported by the Medical Research Council, U.K. (U117512772,
U117562207 and U117570590) and the Francis Crick Institute which receives its core
funding from Cancer Research UK (FC001107), the UK Medical Research Council
(FC001107), and the Wellcome Trust (FC001107).

590
591
592

593 SUPPLEMENTARY FIGURE LEGENDS:

594
595

Sup. Figure 1: Expression of known and novel markers in SOX9iresGFP unchallenged
dataset.

A) *Sox2* and *Sox9* were used to identify SCs while genes encoding for hormones (*Gh*, *Prl*,
Tshβ, *Lhβ*, *Fshβ*, *Pomc*) and lineage specific transcription factors were examined to
distinguish committed cells (*Pou1f1* for somatotrophs, lactotrophs and thyrotrophs, *Tbx19*
and *Pax7* for corticotrophs and melanotrophs, and *Nr5a1* for gonadotrophs). B) Heatmap for
the top10cluster markers.

Sup. Figure 2: Co-localisation between SOX9 and ACTH after Ax.

Co-immunofluorescence for GFP, SOX9 and ACTH on sections of pituitaries harvested one
week after Ax. Expression of Sox9iresGFP (A) and SOX9 (B) is observed in rare ACTH +ve
cells (arrows) exclusively after Ax. The scale bar represents 10 μm for A and 5μm for B.

Sup. Figure 3: Gating strategy for SOX9iresGFP cells.

Gating for SOX9iresGFP cells was validated using wild-type animals. The SOX9iresGFP
fraction we selected typically represented 1 to 2% of all cells while 0.02% of all cells were
falsely sorted in negative controls. This contamination of likely hormonal cells represents a
small proportion of the hormonal cells we detect in our unchallenged Sox9iresGFP fraction
(3% of 1.9% representing 0.08% of the whole population).

Sup. Figure 4: Expression of novel SC cell markers in whole pituitary and SC datasets.

A) UMAP of re-analysed whole pituitary dataset (*28*). This dataset was used to examine
levels of expression of *Ednrb* (B), *Aldah1a2* (C), *Cldn3* (D) and *Msx*1(E) to show exclusive
or enriched expression in SCs. F) Co-immunofluorescence for MSX1, GFP and SOX9 in a
male SOX9iresGFP pituitary section. MSX1 is exclusively expressed in SOX2;SOX9 AL SC
(yellow arrow shows its absence from IL SC, and white arrow shows its expression in
parenchymal SC). *Msx1* violin plot shows enrichment in cluster 0 of the single cell dataset
presented in Fig.1. The scale bar in A represents 10 μm.

Sup. Figure 5: In situ hybridisation for *Ednrb*.

In situ hybridisation analysis (RNAscope) of *Ednrb* expression confirming higher levels of
expression in cleft versus parenchymal SC where low levels of signal can be detected, while
levels of *Sox9* expression appear similar in both compartments. The scale bar represents 200
μm.

Sup. Figure 6: Expression of FS cells markers in SC and whole pituitary datasets.

A-C) UMAP for *S100b* (A), *Aldolase C* (B) and *Slc15a2* (C) in our Sox9iresGFP single cell
dataset.
D-F) UMAP for *S100b* (D), *Sox2* (E) and *Aldolase C* (F) in a whole pituitary dataset (*28*).

Sup. Figure 7: Unchallenged/Challenged integrated analysis.

A) UMAP clustering and marker analysis for integrated datasets from sorted *Sox9*[iresGFP/+]
cells from unchallenged, adrenalectomized, gonadectomized and both adrenalectomized and
gonadectomized animals, 4 days after surgery.
B) Pairwise comparison for proportion test ('pairwise_prop_test' function [stats R package])
on stem cell clusters (circled on UMAP in A). The number of cells assigned to cluster 10
corresponding to proliferative cells (see *Top2a* expression in A) is significantly higher in
activated stem cells (Sup. Table2).
C) Numbers and proportion of specific hormone +ve cells/all hormone +ve cells in each
dataset. While we would expect the number of hormone +ve cells to increase after target

organ ablation, as seen here, the fact that we select fewer GFP weak cells in unchallenged animals prevents us from comparing the proportion of hormone +ve cells in unchallenged and mobilized datasets and thus make this conclusion.

Sup. Figure 8: UMAP clustering for filtered integrated analysis split by dataset.

A) UMAP clustering for integrated datasets from sorted SOX9iresGFP +ve cells from unchallenged, adrenalectomized and sham + 48 h, adrenalectomized + 4 days split by dataset. SC clusters 3 and 4 mostly originate from datasets collected 48 hours after surgery.

Sup. Figure 9: Analyses of marker expression on UMAP clustering in Ax integrated datasets.

Markers of SC and hormone-secreting cell types are shown on the UMAP clustering for integrated datasets from GFP +ve cells sorted from $Sox9^{iresGFP/+}$ pituitaries from mice that were unchallenged, adrenalectomized and sham + 48 h, adrenalectomized + 4 days.

Sup. Figure 10: PCA clustering for filtered integrated analysis split by dataset.

SC4 cluster cells located in an intermediate position between SC and endocrine cells originate mostly from Ax+48h, which is consistent with mobilization and increased commitment toward differentiation.

Sup.Fig.11: Analysis of endocrine trajectories and regulons.

Heatmaps of genes associated with trajectories and SCENIC analyses for lactotroph, somatotroph and gonadotroph lineages. Representative genes are shown. Known regulators and those in common with the corticotroph analyses (Fig.5 F-H) are displayed on the heatmaps.

Sup. Figure 12: Expression of *Met* and *Hgf* in the whole pituitary dataset.

Expression of *Hgf* was plotted along with *Pomc* and *Pdgfrb* suggesting expression in *Pomc* positive corticotrophs and *Pdgfrb* positive pericytes, while *Met* expression is restricted to *Sox2* positive SCs (from re-analysed dataset (*28*)). The same pattern is observed in a different dataset (*37*).

Sup. Table1: Pathway analysis for clusters 0 and 2.
Gene markers for WT Cluster 0 were used to determine enriched pathways with Metacore using a hypergeometric test. Pathways with an adjusted p value < 0.05 are shown in the table. Genes present in our dataset are highlighted in the top5 pathways.

Sup. Table2: Endocrine cell cluster numbers from Ax integrated dataset (Fig.5)

701   Sup. Table 3: Pairwise for proportion test results.

703   Sup. Table 4: Counting numbers for Fig. 2, 3, 5 and 6.

705   Sup. Table 5: Cell type signatures.

708   MATERIALS AND METHODS
709
710
711   **Mice**
712
All animal experiments carried out were approved under the UK Animals (Scientific
Procedures) Act 1986 and under the project licences n. 80/2405 and PP8826065.
Sox9$^{iresGFP/+}$ (*Sox9^{tm1Haak}*) (*48*), *Sox9$^{iresCreERT2/+}$* (*Sox9$^{tm1(cre/ERT2)Haak}$*) (*48*), *Wnt1Cre* ((*no
gene)$^{tg(Wnt1-GAL4)11Rth}$*) (*49*) and Rosa26$^{ReYFP/ReYFP}$ (*Gt(ROSA)26Sortm1(EYFP)Cos*) (*50*) mice
were maintained on a C57Bl6 background.
For CreERT2 induction in *Sox9$^{iresCreERT2/+}$* ;Rosa26$^{ReYFP/+}$ animals, tamoxifen was
administered for 3 consecutive days at a concentration of 0.2mg/g body weight. For
Sox9$^{iresCreERT2/+}$;Rosa26$^{ReYFP/+}$ pups induction, a single subcutaneous injection of tamoxifen
(0.25mg/g body weight) was performed. Target organ ablations were performed as previously
described (*3*).
723
724   **Pituitary SC selection**
725
The anterior lobe of freshly dissected 8 to 12 week-old pituitaries was minced and incubated
for 15mn at 37°C in 1mg/ml Papain (Sigma 10108014001), 10µg/ml DNAse I (Sigma
10104159001) and 10µg/ml Rock inhibitor (Abmole Bioscience M1817) in HBSS (Sigma
H9394). pH was adjusted by addition of NaOH. The enzymatic solution was removed, and
the cells were mechanically resuspended in HBSS supplemented with DNAse I and Rock
inhibitor, as above. Cells were FACsorted for GFP using a large nozzle (100µm) and low
pressure (20psi) for optimal cell survival. For immunofluorescence, sorted cells were plated
on Superfrost slides for 90 min in a cell culture incubator, fixed 20 min on ice in 4% PFA and
immediately processed.
735
736   **Pituisphere culture.**
737
738   Dissociated anterior lobes (see above) of male and female 7 to 12 week-old mice were seeded
739   at 50.10^3 cells/ml in pituisphere medium containing EGF and FGF (*9*). 100ng/ml of HGF
740   (2207-HG-025/CF, R&D systems) was added to this basic medium in treated cultures.
741   Medium was replaced every other day and spheres manually and blindly counted after 7 days.
742   4 independent repeats were performed.
743
744
745   **Single cell sequencing**
746
The concentration and viability of the single cell suspension was measured using Trypan
Blue and the Eve automatic cell counter (unchallenged and target organ ablation d4 samples)
or acridine orange and propidium iodide with the Luna-FX7 automatic cell counter (Ax and
Sham control d2 samples). Up to 10,000 cells were loaded on Chromium Chip and

partitioned in nanolitre scale droplets using the Chromium Controller and Chromium 10x 3
prime v2.0, Chromium Single Cell 3' Reagent Kits User Guide (v2 Chemistry) (unchallenged
and target organ ablation d4 samples) or Next GEM Single Cell reagents (3 prime v3.1) (Ax
and Sham control d2 samples). Within each droplet the cells were lysed, and the RNA was
reverse transcribed. All of the resulting cDNA within a droplet shared the same cell barcode.
Illumina compatible libraries were generated from the cDNA using Chromium Next GEM
Single Cell library reagents in accordance with the manufacturer instructions (10x
Genomics). Final libraries are quality checked using the Agilent TapeStation and sequenced
using the Illumina HiSeq4000 or HiSeq2500, read configuration 26-8-0-98 (unchallenged and
target organ ablation d4 samples) or NovaSeq 6000, read configuration 28-10-10-90.
Where expression of specific genes in the whole pituitary gland is referenced as ((*37*), we
used this published dataset on the Loupe browser to examine expression of relevant markers.
763
764 **Bioinformatic analyses**
765
Alignment, Seurat analysis, integration and cluster annotation were performed using
CellRanger and Seurat. Raw reads were initially processed by the Cell Ranger v.3.0.2
pipeline (*51*), which deconvolved reads to their cell of origin using the UMI tags, aligned
these to the mm10 transcriptome (to which we added the eGFP sequence
https://www.addgene.org/browse/sequence/305137/ to detect eGFP expressing cells) using
STAR (v.2.5.1b) (*52*) and reported cell-specific gene expression count estimates. All
subsequent analyses were performed in R v.3.6.0 (*53*) using the Seurat (v3) package
(*54*). Genes were considered to be 'expressed' if the estimated (log10) count was at least 0.1.
Primary filtering was then performed by removing from consideration cells expressing fewer
than 50 genes and cells for which mitochondrial genes made up greater than 3 standard
deviations from the mean of mitochondrial expressed genes. PCA decomposition was
performed and, after consideration of the eigenvalue 'elbow-plots', the first 20 components
were used to construct the UMAP plots per sample. Multiple samples, generated in this study
and published (*28*), were integrated using 2000 variable genes and Seurat's CCA method.
Cluster specific gene markers were identified using a Wilcoxon rank sum test and the top 20
genes ranked by logFC per cluster were used to generate a heatmap. Clusters were annotated
using label tranfer methods within Seurat using the GSE120410 dataset (*28*). Clusters were
further annotated using cell specific signatures (Sup. Table 5).
Cell trajectories were identified using the package 'Slingshot' (version 1.4.0) (*29*), using the
undifferentiated cluster as a starting point and the PCA co-ordinates. Lineages were identified
showing specific trajectories ending in specific differentiated cells. Differential expression of
genes between samples were further investigated for the corticotroph lineage with a
correlation metric against pseudotime, using the 'gam' package (version 1.20) (*55*). Heatmaps
were made using the smoothed expression of differential genes, and the cells are ranked by
increasing pseudotime.
In order to investigate transcription factor activity along pseudotime, the package SCENIC
(*32*) was used. A list of 948 putative mouse regulatory binding sites found in
promoter regions of expressed genes were used to identify shared regulatory networks.
Regulon activity score per transcription factor was calculated using the AUCell function,

where the enrichment of target genes was measured using the area under the curve of the
target gene relative to the expression-based ranking of all genes. The top 200 regulons ranked
by activity score were using to generate a binarized heatmap, where the cells were ranked by
pseudotime.
799
800

801 **Immunofluorescence and RNAscope.**
802
Pituitaries were fixed after dissection by immersion in 4% PFA at 4ºC. Immunofluorescent
stainings were performed on cryosections as described (*56*). The following primary
antibodies were used: Goat anti-SOX2 (Biotechne, AF2018), rabbit anti-SOX9 (a gift from F.
Poulat, IGH Montpellier, France), rat anti-GFP (Nacalai Tesque 04404-84), chick anti-GFP
(Invitrogen, A10262), rabbit anti-EDNRb (Abcam 117529), goat anti-MSX1 (R&D system
AF5045), rabbit anti-ALDH1A2 (Abcam 75674, after antigen retrieval), rabbit anti-Claudin3
(Abcam ab15102), rat anti-PECAM (BD Pharmigen, 550274), rat anti-
PDGFRβ (eBiosciences 14-1402-81), rabbit anti-AQP3 (Alomone Labs, AQP-003, after
antigen retrieval) rabbit anti-Runx1 (Sigma, HPA004176), rabbit anti-IRF9 (Cell Signalling,
D9I5H), goat anti-Met (R&D, AF 527) and hormone antibodies anti-LH, GH, ACTH and
PRL from the National Hormone Peptide Program (A.F. Parlow, Torrance, USA). Slides
were then incubated with Alexa Fluor secondary antibodies. Imaging of stained tissue
sections was performed on a Leica SPE microscope while imaging of plated cells was done
on an Olympus spinning disk microscope.
RNAscope was performed on cryosections following the manufacturer's instructions using
Hgf (#456511) and *Ednrb* (#473801) probes.
819
820

821 **Cell counts**
822
Counts were performed blindly. For dissociated cell counts, slides stained by
immunofluorescence were scanned; automated counts were then performed using QuPath
(https://qupath.github.io/). For *Sox9$^{iresCreERT2/+}$* ;Rosa26$^{ReYFP/+}$;hormone counting on sections,
double +ve cells were manually identified and each imaged on a slide representing 1/5 of a
pituitary. Images were then reviewed, and double +ve cells counted. All counting results and
tests are presented (Sup. Table 3 and 4).
829
830 **Statistics**
831
832 Comparisons of the percentages of Sox9;PDGFRβ (Fig.3) were performed after angular
833 transformation using Mann-Whitney test on Prism. Pairwise for proportion test (Sup. Fig4B,
834 Fig.5B, Fig.6B and Sup. Table 3), t-test (Fig.6E , Fig.7D) and Chi squared test (Fig.7) were
835 performed using the Stats R package.
836
837

Author contributions:

      Conceptualization: KR, RLB
      Methodology: KR
      Bioinformatic analyses: PC
      Investigation: KR, DS
      Visualization: KR
      Supervision: KR, RLB
      Writing—original draft: KR
      Writing—review & editing: KR, RLB

Competing interests:

Authors declare that they have no competing interests.

Data and materials availability:

All data are available in the main text or the supplementary materials.
Codes are available on request. Datasets have been submitted (Awaiting confirmation of data deposited at GEO).

BIBLIOGRAPHY

1. A. Levy, Physiological implications of pituitary trophic activity. *J Endocrinol* **174**, 147-155 (2002).
2. D. Langlais, C. Couture, M. Kmita, J. Drouin, Adult pituitary cell maintenance: lineage-specific contribution of self-duplication. *Mol Endocrinol* **27**, 1103-1112 (2013).
3. K. Rizzoti, H. Akiyama, R. Lovell-Badge, Mobilized adult pituitary stem cells contribute to endocrine regeneration in response to physiological demand. *Cell Stem Cell* **13**, 419-432 (2013).
4. C. L. Andoniadou *et al.*, Sox2(+) stem/progenitor cells in the adult mouse pituitary support organ homeostasis and have tumor-inducing potential. *Cell Stem Cell* **13**, 433-445 (2013).
5. H. Roose *et al.*, Major depletion of SOX2(+) stem cells in the adult pituitary is not restored which does not affect hormonal cell homeostasis and remodelling. *Sci Rep* **7**, 16940 (2017).
6. Q. Fu *et al.*, The adult pituitary shows stem/progenitor cell activation in response to injury and is capable of regeneration. *Endocrinology* **153**, 3224-3235 (2012).
7. A. Vennekens *et al.*, Interleukin-6 is an activator of pituitary stem cells upon local damage, a competence quenched in the aging gland. *Proc Natl Acad Sci U S A* **118**, (2021).
8. F. Ruf-Zamojski *et al.*, Single nucleus multi-omics regulatory landscape of the murine pituitary. *Nature communications* **12**, 2677 (2021).

882   9.   T. Fauquier, K. Rizzoti, M. Dattani, R. Lovell-Badge, I. C. Robinson, SOX2-expressing
883   progenitor cells generate all of the major cell types in the adult mouse pituitary
884   gland. *Proc Natl Acad Sci U S A* **105**, 2907-2912 (2008).
885   10.   Z. Zhang *et al.*, Single nucleus transcriptome and chromatin accessibility of
886   postmortem human pituitaries reveal diverse stem cell regulatory mechanisms. *Cell*
887   *reports* **38**, 110467 (2022).
888   11.   L. Y. M. Cheung, S. A. Camper, PROP1-Dependent Retinoic Acid Signaling Regulates
889   Developmental Pituitary Morphogenesis and Hormone Expression. *Endocrinology*
890   **161**, (2020).
891   12.   S. Yoshida *et al.*, Retinoic acid signalling is a candidate regulator of the expression of
892   pituitary-specific transcription factor Prop1 in the developing rodent pituitary. *J*
893   *Neuroendocrinol* **30**, e12570 (2018).
894   13.   N. Nishimura *et al.*, Search for regulatory factors of the pituitary-specific
895   transcription factor PROP1 gene. *J Reprod Dev* **62**, 93-102 (2016).
896   14.   E. Laporte, A. Vennekens, H. Vankelecom, Pituitary Remodeling Throughout Life: Are
897   Resident Stem Cells Involved? *Frontiers in endocrinology* **11**, 604519 (2020).
898   15.   S. Yoshida *et al.*, Cell type-specific localization of Ephs pairing with ephrin-B2 in the
899   rat postnatal pituitary gland. *Cell Tissue Res* **370**, 99-112 (2017).
900   16.   M. I. Perez Millan, M. L. Brinkmeier, A. H. Mortensen, S. A. Camper, PROP1 triggers
901   epithelial-mesenchymal transition-like process in pituitary stem cells. *Elife* **5**, (2016).
902   17.   P. R. Le Tissier, P. Mollard, Renewing an old interest: Pituitary folliculostellate cells. *J*
903   *Neuroendocrinol*, e13053 (2021).
904   18.   L. O'Hara, H. C. Christian, N. Jeffery, P. Le Tissier, L. B. Smith, Characterisation of a
905   mural cell network in the murine pituitary gland. *J Neuroendocrinol* **32**, e12903
906   (2020).
907   19.   K. Fujiwara *et al.*, Aldolase C is a novel molecular marker for folliculo-stellate cells in
908   rodent pituitary. *Cell Tissue Res* **381**, 273-284 (2020).
909   20.   W. Allaerts, H. Vankelecom, History and perspectives of pituitary folliculo-stellate
910   cell research. *Eur J Endocrinol* **153**, 1-12 (2005).
911   21.   E. Laporte *et al.*, Decoding the activated stem cell phenotype of the neonatally
912   maturing pituitary. *Elife* **11**, (2022).
913   22.   P. A. Fletcher *et al.*, The astroglial and stem cell functions of adult rat folliculostellate
914   cells. *Glia* **71**, 205-228 (2023).
915   23.   S. Zhang *et al.*, Single-cell transcriptomics identifies divergent developmental lineage
916   trajectories during human pituitary development. *Nature communications* **11**, 5275
917   (2020).
918   24.   S. W. Davis *et al.*, beta-catenin is required in the neural crest and mesencephalon for
919   pituitary gland organogenesis. *BMC Dev Biol* **16**, 16 (2016).
920   25.   H. C. Etchevers, C. Vincent, N. M. Le Douarin, G. F. Couly, The cephalic neural crest
921   provides pericytes and smooth muscle cells to all blood vessels of the face and
922   forebrain. *Development* **128**, 1059-1068 (2001).
923   26.   P. Le Tissier *et al.*, An updated view of hypothalamic-vascular-pituitary unit function
924   and plasticity. *Nat Rev Endocrinol* **13**, 257-267 (2017).
925   27.   L. A. Nolan, A. Levy, A population of non-luteinising hormone/non-
926   adrenocorticotrophic hormone-positive cells in the male rat anterior pituitary
927   responds mitotically to both gonadectomy and adrenalectomy. *J Neuroendocrinol*
928   **18**, 655-661 (2006).

28.   L. Y. M. Cheung *et al.*, Single-Cell RNA Sequencing Reveals Novel Markers of Male
Pituitary Stem Cells and Hormone-Producing Cell Types. *Endocrinology* **159**, 3910-
3924 (2018).
29.   K. Street *et al.*, Slingshot: cell lineage and pseudotime inference for single-cell
transcriptomics. *BMC Genomics* **19**, 477 (2018).
30.   J. E. Welcker *et al.*, Insm1 controls development of pituitary endocrine cells and
requires a SNAG domain for function and for recruitment of histone-modifying
factors. *Development* **140**, 4947-4958 (2013).
31.   J. Drouin, The corticotroph cells from early development to tumorigenesis. *J
Neuroendocrinol* **34**, e13147 (2022).
32.   S. Aibar *et al.*, SCENIC: single-cell regulatory network inference and clustering. *Nat
Methods* **14**, 1083-1086 (2017).
33.   K. M. Osorio, K. C. Lilja, T. Tumbar, Runx1 modulates adult hair follicle stem cell
emergence and maintenance from distinct embryonic skin compartments. *J Cell Biol*
193, 235-250 (2011).
34.   C. M. Horvath, G. R. Stark, I. M. Kerr, J. E. Darnell, Jr., Interactions between STAT and
non-STAT proteins in the interferon-stimulated gene factor 3 transcription complex.
Mol Cell Biol **16**, 6957-6964 (1996).
35.   M. Valny, P. Honsa, D. Kirdajova, Z. Kamenik, M. Anderova, Tamoxifen in the Mouse
Brain: Implications for Fate-Mapping Studies Using the Tamoxifen-Inducible Cre-loxP
System. *Frontiers in cellular neuroscience* **10**, 243 (2016).
36.   A. Grabek *et al.*, The Adult Adrenal Cortex Undergoes Rapid Tissue Renewal in a Sex-
Specific Manner. *Cell Stem Cell* **25**, 290-296 e292 (2019).
37.   A. Mayran *et al.*, Pioneer and nonpioneer factor cooperation drives lineage specific
chromatin opening. *Nature communications* **10**, 3807 (2019).
38.   X. Z. Hou *et al.*, Expression of hepatocyte growth factor and its receptor c-Met in
human pituitary adenomas. *Neuro Oncol* **12**, 799-803 (2010).
39.   Y. Wang *et al.*, A cell fitness selection model for neuronal survival during
development. *Nature communications* **10**, 4137 (2019).
40.   K. Ogasawara *et al.*, Hormonal regulation of prolactin cell development in the fetal
pituitary gland of the mouse. *Endocrinology* **150**, 1061-1068 (2009).
41.   D. Gonzalez *et al.*, Luteinizing hormone secretion elicited in a ligand-independent
activation of progesterone receptor manner at pituitary level in the rat: differential
effect of two selective estrogen receptor modulators. *Neurosci Lett* **289**, 111-114
(2000).
42.   P. Mollard, D. J. Hodson, C. Lafont, K. Rizzoti, J. Drouin, A tridimensional view of
pituitary development and function. *Trends Endocrinol Metab* **23**, 261-269 (2012).
43.   W. Kelsch, C. W. Lin, C. P. Mosley, C. Lois, A critical period for activity-dependent
synaptic development during olfactory bulb adult neurogenesis. *J Neurosci* **29**,
11852-11858 (2009).
44.   P. R. Le Tissier, J. F. Murray, P. Mollard, A New Perspective on Regulation of Pituitary
Plasticity: The Network of SOX2-Positive Cells May Coordinate Responses to
Challenge. *Endocrinology* **163**, (2022).
45.   V. Moncho-Amor *et al.*, SOX2 is required independently in both stem and
differentiated cells for pituitary tumorigenesis in p27-null mice. *Proc Natl Acad Sci U
S A* **118**, (2021).

46. M. Takeo *et al.*, EdnrB Governs Regenerative Response of Melanocyte Stem Cells by
Crosstalk with Wnt Signaling. *Cell reports* **15**, 1291-1302 (2016).
47. S. W. Davis, A. H. Mortensen, S. A. Camper, Birthdating studies reshape models for
pituitary gland cell specification. *Developmental biology* **352**, 215-227 (2011).
48. K. Furuyama *et al.*, Continuous cell supply from a Sox9-expressing progenitor zone in
adult liver, exocrine pancreas and intestine. *Nat Genet* **43**, 34-41 (2010).
49. P. S. Danielian, D. Muccino, D. H. Rowitch, S. K. Michael, A. P. McMahon,
Modification of gene activity in mouse embryos in utero by a tamoxifen-inducible
form of Cre recombinase. *Curr Biol* **8**, 1323-1326 (1998).
50. S. Srinivas *et al.*, Cre reporter strains produced by targeted insertion of EYFP and
ECFP into the ROSA26 locus. *BMC Dev Biol* **1**, 4 (2001).
51. G. X. Zheng *et al.*, Massively parallel digital transcriptional profiling of single cells.
Nature communications **8**, 14049 (2017).
52. A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013).
53. R. C. Team, R: A language and environment for statistical computing. *R Foundation
for Statistical Computing, Vienna, Austria.* , (2019).
54. T. Stuart *et al.*, Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902
e1821 (2019).
55. T. Hastie, gam: Generalized Additive Models. . *R package version 1.20*, (2020).
56. K. Rizzoti *et al.*, SOX3 is required during the formation of the hypothalamo-pituitary
axis. *Nat Genet* **36**, 247-255 (2004).
997
998