Steady state visual evoked potentials reveal a signature of the pitch-size crossmodal association in visual cortex

Placido Sciortino, Christoph Kayser

Department of Cognitive Neuroscience, Faculty of Biology, Bielefeld University, 33501 Bielefeld, Germany

Corresponding author: Placido Sciortino, placido.sciortino@uni-bielefeld.de

Abbreviated title: Visual cortex reflects crossmodal associations

Keywords: Crossmodal associations, steady state visual evoked potentials, crossmodal correspondences, SSVEP

Abstract

Crossmodal correspondences describe our tendency to associate sensory features from different modalities with each other, such as the pitch of a sound with a specific size of a visual object. While such crossmodal correspondences (or associations) have been described in many behavioural studies their neurophysiological correlates remain debated. Under the current working model of multisensory perception both a low- and a high-level account seem plausible. That is, the neurophysiological processes shaping these associations could commence in early sensory regions, or may rather emerge only later and as a result from high-level multisensory integration processes or in semantic and object identification networks. We implemented a paradigm based on steady-state visual evoked potentials (SSVEP) to directly test the hypothesis that the associations between acoustic pitch and the visual features of size, hue or chromatic saturation are visible in brain activity from early visual cortices. Our results reveal a pitch-size congruency effect in the SSVEP over occipital electrodes and in the behavioural data, supporting a low-level account of this crossmodal association. For the associations between pitch and chromatic features we found no significant effects. We speculate that this signature of the pitch size association in early visual cortices reflects the successful pairing of congruent visual and acoustic object properties and may contribute to establishing causal relations between multisensory objects.

Introduction

Humans often associate sensory features from different modalities with each other (Kohler, 1929, 1947; Spence, 2011; Spence and Deroy, 2013). For example, a low pitched sound is typically associated with the larger of two visual objects. Such crossmodal associations have been defined as the intuitive pairing between perceptual attributes of stimuli in different senses and are known also as ''crossmodal similarities'', "crossmodal correspondences", ''synesthetic associations'' or ''sound symbolic associations''. Such crossmodal associations have been reported for various combinations of sensory modalities and features and have been studied in a variety of experimental paradigms that range from verbal reports to speeded classification tasks (Spence, 2011). Still, these behavioural results are only partly paralleled by neuroimaging data and the cerebral origins of crossmodal associations remain unclear.

In particular, it remains debated whether crossmodal associations are mediated by low-level brain regions or whether they are the result of high-level integration processes in semantic or object identification networks (McCormick et al., 2021, 2018). Under the current working model of multisensory perception, both a low- and a high-level account seem plausible. In this model, multisensory representations emerge along multiple stages of the sensory pathways, commencing in early sensory regions with automatic interactions constrained by stimulus location or timing and continuing with higher specificity in association regions (Bizley et al., 2016; Cao et al., 2019; Noppeney, 2021; Rohe and Noppeney, 2015). For example, early sensory regions may accumulate multisensory evidence while premotor and parietal regions implement the reliability-weighted integration of the unisensory cues and temporo-frontal regions merge information about object identity or semantics.

In one previous study, participants were trained on specific sound-symbolic associations and EEG recordings revealed association-specific activations around 140 ms post-stimulus onset, letting the authors speculate about an involvement of low-level sensory regions (Kovic et al., 2010). Such an account would be in line with object-related multisensory congruency effects at even earlier latencies (Molholm et al., 2004) and the activation of auditory and visual cortices in the sound-symbolic association of the Bouba-Kiki effect (Peiffer-Smadja and Cohen, 2019) or other multisensory paradigms (Brang et al., 2022; Kayser et al., 2010; Lakatos et al., 2009; Schroeder and Foxe, 2005). However, other EEG studies on the pitch-size association pointed to effects at longer latencies, possibly arising from parietal regions (Bien et al., 2012; Stekelenburg and Keetels, 2016). Such a high-level account is also supported by fMRI studies that probed the associations of pitch with elevation and pseudo-words with shapes and reported congruency effects in a wider network of parietal, frontal and insular regions (McCormick et al., 2018, 2021). These congruency effects did not overlap with regions involved in semantic processes, magnitude estimation or multisensory object integration, suggesting a distributed neurophysiological correlate that may arise from changes in multisensory attention, memory- or imagination-related processes (McCormick et al., 2021). However, multisensory attention and imagination have neurophysiological correlates that can extend into early sensory regions (Ferrari and Noppeney, 2021; Talsma et al., 2010; Vetter et al., 2014; Zuanazzi and Noppeney, 2020), which again fuels speculations that crossmodal associations may commence or shape activity at early states of cortical processing (Peiffer-Smadja and Cohen, 2019).

We here directly tested the hypothesis that associations between auditory and visual features are reflected in activity from early visual cortices - hence that their signatures are visible in functionally defined low-level sensory regions. To this end we exploited visual steady-state evoked potentials (SSVEP) as markers of neural activity arising from early visual cortices (Norcia et al., 2015; Regan, 1966; Vialatte et al., 2010). The SSVEP is elicited by the rapid presentation of visual stimuli at a fixed rate and has been used to study the involvement of visual cortices in processes such as figure ground segregation, perceptual dominance, binocular rivalry, attention, object recognition and memory (Alpers et al., 2005; Appelbaum et al., 2008, 2006; Brown and Norcia, 1997; Kaspar et al., 2010; Müller and Hübner, 2002; Silberstein et al., 2001; Sutoyo and Srinivasan, 2009; Walter et al., 2012). We chose this paradigm as it comes with two specific advantages: first, it relies on a functionally-induced and well-studied neurophysiological marker of sensory-specific activations (Norcia et al., 2015; Vialatte et al., 2010). Second, it allows probing the stimulus-selective response of visual cortex to one of two simultaneously presented stimuli, hence emphasising the 'binding' nature of multisensory associations. This binding effect may be characteristic of crossmodal correspondences but is more difficult to assess in paradigms involving only one stimulus per modality (Parise and Spence, 2009).

In our experiment we focused on three putative crossmodal associations of pitch with visual features: size, hue and chromatic saturation. All three features reflect basic visual attributes that are encoded in visual cortices and hence their association with pitch may have a correlate in low-level sensory regions. An association between pitch and size has been reported in multiple behavioural studies, whereas the associations of pitch with hue and pitch with saturation have been found less consistently (Anikin and Johansson, 2019; Bernstein et al., 1971; Caivano, 1994; Giannakis, 2001; Hamilton-Fletcher et al., 2017; Melara, 1989; Moos et al., 2014; Orlandatou, 2012; Ward et al., 2006). The comparison of these three putative associations allowed us to probe whether they potentially all share a similar neurophysiological correlate. We tested these associations by presenting two concurrent flickering visual stimuli that differed along one specific feature dimension (size, hue or saturation) together with one sound (low or high in pitch) or in the absence of a sound. Thereby we probed both the influence of the presence of any sound on the visual cortical response and the selective influence of low or high pitched sounds on the response to specific sizes, hues or saturation levels. Our results reveal a congruency-related effect for the pitch-size association in both the SSVEP and behavioural data, but not for the other two associations. Given that the SSVEP mainly originates from early visual cortices (Di Russo et al., 2007; Lauritzen et al., 2010) this result speaks for a correlate of the pitch-size association in early sensory regions and shows that SSVEP can be exploited to study crossmodal processes.

Methods

Participants

A total of 26 participants (16 females, mean age = 27.5) participated in the experiment. The sample size is comparable to previous studies using the SSVEP procedure and testing crossmodal associations in behaviour or brain activity (Appelbaum et al., 2006; Kaspar et al., 2010; Lauritzen et al., 2010; Parise and Spence, 2012, 2009). All participants gave written informed consent prior to participation and all procedures were approved by the Ethics Committee of Bielefeld University. Participants were compensated for their time (10 Euro/hour). The experiments were performed in a darkened and electrically shielded room (E:box, Desone). Participants sat on a comfortable chair in front of a computer screen with a grey background. The data from one participant had to be excluded (see below), hence all data are for a sample of 25 participants.

General paradigm

The paradigm was designed to probe three audio-visual associations: between pitch and the size of a visual object, between pitch and hue, and between pitch and chromatic saturation. We chose the association between size and pitch as this has been reported robustly across many studies (Evans and Treisman, 2010; Gallace and Spence, 2006; Parise and Spence, 2008, 2009). The associations between pitch and the hue or saturation of visual stimuli have been reported before, but the robustness of these remains debated (Anikin and Johansson, 2019; Bernstein et al., 1971; Caivano, 1994; Hamilton-Fletcher et al., 2017; Melara, 1989; Moos et al., 2014; Simpson et al.,1956; Ward et al., 2006). Concerning hue, we focused on blue and yellow, as these are considered safer than for example red and green when presented as flickering stimuli (Drew et al., 2001; Tello et al., 2015).

Visual stimulation was based on an SSVEP procedure that elicits responses primarily from visual cortices (Di Russo et al., 2007; Lauritzen et al., 2010; Norcia et al., 2015; Vanegas et al., 2013; Vialatte et al., 2010). During each trial two visual stimuli were presented for 6.2 seconds, one flickering at 7.5 Hz and one at 12 Hz and were presented on a uniform grey background (15 cd/m^2). These were presented 11 degrees of the left and right of a central fixation point. The individual stimuli were defined by their shape (irrelevant for the specific hypotheses tested), their size, hue and saturation (Figure 1). During each trial the two visual stimuli differed by only one feature, which defied the visual feature whose association with the sound was probed in the respective trial. The specific shapes were a triangle, a circle and a square. The shape was varied in order to minimise intramodal associations between the manipulated features and other properties such as angularity, spikiness, or the number of edges (Dreksler and Spence, 2019). The small and large visual stimuli subtended about 3.5 and 6.5 degrees of visual angle in diameter. For the specific values of hue and saturation see below. During each trial, either one of two sounds was presented (high pitch: 1000 Hz tone, low pitch: 500 Hz tone) or no sound was present. Across trials, the different combinations of visual features, flicker frequency, side of presentation and auditory conditions were counterbalanced.

As a result, each trial either probed one specific crossmodal association (pitch-size, pitch-hue or pitch-saturation) or presented the visual stimuli in the absence of an auditory stimulus. When a tone was

presented, one of the two visual stimuli was congruent with the sound according to the associations tested, the other was incongruent. Congruencies were operationally defined as follows and similar to the previous literature: high pitch & small size (low pitch & large size); high pitch & yellow pitch (low pitch & blue pitch); high pitch & high saturation (low pitch & low saturation). A total of 288 trials were presented, divided into 6 blocks. Each association was tested in 96 trials each that were divided into two blocks. Each trial started with a fixation period (600 to 900 ms, uniform), followed by the 6.2 s stimulation period. Inter-trial intervals lasted 1100 to 1400ms (uniform).

At the end of the stimulation period one of the two shapes was presented for an additional 4 frames (33.3ms) and hence was visible longer than the other. Participants were asked to indicate which of the two shapes was presented longer by pressing the matching right or left arrow key on a keyboard. Participants were instructed to be as fast and as accurate as possible. Importantly, on trials with a sound present the visual stimulus remaining longer could be either the congruent or incongruent one with the concurrent pitch. We employed this task to ensure that participants maintained attention to both visual stimuli and to probe for a behavioural correlate of the crossmodal association.

Stimuli were presented using the Psychtoolbox (3.0.15) from Matlab (R2017a). Visual stimuli were presented on an Asus #PG279Q monitor running at 120Hz refresh rate, with stimuli at 7.5 Hz being presented each 16th frame and those at 12Hz each 10th frame. Each stimulus was drawn for two consecutive frames on the screen. The timing of the visual stimuli was verified for each trial and participant by inspecting the time stamps provided by Psychtoolbox and was generally verified using an oscilloscope. This revealed that the actual screen refresh rate was 116.9 ± 0.12 Hz (mean ±s.e.m. across experimental days) rather than the nominal 120Hz. Hence, the two actual flicker frequencies were effectively 7.26 (termed f.low) and 11.6 Hz (f.high). These values were very consistent across participants and used for subsequent data analysis. The auditory stimuli were presented at 62 dB SPL from two speakers positioned to the left and right of the monitor. Sound intensity was calibrated using a sound level-meter (Bruel & Kjaer 2250).

Selection of hue and saturation levels
Because the perceived hue and saturation of chromatic stimuli is subjective we used participant-wise calibrated stimuli that were subjectively perceived as 'yellow' and 'blue' and had similar perceived lightness and saturation against a uniform grey background. We considered this important as otherwise two visual stimuli may also differ in their subjective saliency. For this purpose we conducted a pre-test. In a first step participants were presented a HSV colour wheel (saturation and lightness values of 0.7) on a grey background and were instructed to select the "most pure blue" or "most pure yellow" direction respectively using a mouse. Subsequently, participants were presented with one of the selected hues (at fixed saturation) along with a coloured bar presenting the other hue in varying degrees of saturation. They were then asked to select the level of saturation of the second hue matching that of the first. In a third step they were asked to equalise the level of lightness in a similar manner. This iterative process was randomized and repeated 4 times (twice matching yellow against blue and twice blue against yellow). We then calculated the average of each HSV attribute and used these for each participant as hue, saturation and lightness. While lightness was fixed through the experiment for each hue, saturation varied between high- and low-saturation trials. High saturation

was defined as the average of the chosen saturation values, while low saturation was defined by dividing the chosen high value by two.

Figure 1. Schematic of the stimuli used to test the associations of pitch with hue, size and saturation. The two visual stimuli presented on each trial differed by one dimension (either hue, size, saturation) and could take one of three shapes (circle, square, triangle) to avoid biasing the results by using a single shape. The two visual stimuli flickered independently at 7.26 and 11.6 Hz for a period of 6.2s and were accompanied by either a high pitched tone, a low pitched tone or no sound The individual features, flicker frequencies, sound conditions and side of presentation were counterbalanced across trials.

EEG recording and pre-processing

EEG signals were acquired using a 128 channel BioSemi (BioSemi, B.V.) system with Ag-AgCl electrodes mounted on an elastic cap (BioSemi). Four additional electrodes were placed at the outer canthi and below the eyes to obtain the electro-oculogram (EOG). Data were acquired at 1028 Hz sampling rate and electrodes offset were kept below 25 mV. Data analysis was performed using MATLAB (The MathWorks Inc.; R 2017A) and the FieldTrip toolbox (ver. 22/09/2019)(Oostenveld et al., 2011).

Data pre-processing was similar to our previous work (Kayser and Kayser, 2020; Park and Kayser, 2021; Sciortino and Kayser, 2022a, 2022b). The data were low-pass filtered between 0.6 and 90 Hz and resampled to 200 Hz. Electrodes with faulty contacts were interpolated (on average 0.65 ± 0.15 per participant, mean ± s.e.m.). Subsequently the data were denoised using Independent component analysis (ICA). We identified and removed components reflecting oculomotor or muscle activity following previous literature and using a semi-automatic procedure (Campos Viola et al., 2009; Hipp and Siegel, 2013; O'Beirne and Patuzzi, 1999; Sciortino and Kayser, 2022a, 2022b). In total we removed 11.5 ± 0.10 mean ± s.e.m. components per participant. To extract the SSVEP the pre-processed data were epoched around the stimulation interval, removing the first 500 ms following stimulus onset and omitting the last frames of the stimulus that introduced the behavioural task.

Quantification of the SSVEP response

To visualise and quantify the SSVEP we computed the power spectra of the data in Fieldtrip. For this we resampled the data to 40 Hz and computed the time- and epoch-averaged power using Hanning windows. Power estimates were obtained for each electrode separately and by combining data epochs either across all trials and conditions or individually for specific conditions (see below). We visually inspected these spectra and confirmed that they reflect the overall 1/f dependency known from EEG data (Hipp and Siegel, 2013; O'Beirne and Patuzzi, 1999). The data from one participant clearly deviated from this expected pattern and the data from this participant were excluded.

Based on the stimulation paradigm we expected two local peaks in these spectra around 7.26 and 11.6Hz. Because the overall shape of the EEG spectra and their overall power differs between participants, we computed the signal to noise ratio at these two stimulation frequencies (SNR) as a participant-wise normalised measure of the SSVEP response. For this we divided the power near each stimulation frequency (7.26 ± 0.1 Hz and 11.65 ± 0.1 Hz respectively) by the average power in neighbouring bands. These neighbouring bands were defined as -0.7Hz to -0.3Hz below and +0.3Hz to +0.7Hz around each stimulation band. A group-level topographical visualisation is shown in Figure 2. This reveals the expected strong SSVEP-SNR over occipital electrodes, indicating that the stimulation frequencies adopted were successful in generating a strong signal in visual cortices (Di Russo et al., 2007; Lauritzen et al., 2010; Norcia et al., 2015; Vanegas et al., 2013; Vialatte et al., 2010).

To probe our main question, we computed this SNR for specific constellations of trials. First, we split trials according to the sound conditions (no sound, high or low pitch). In addition, by design of our paradigm, we split trials according to whether the congruent (incongruent) visual stimulus on each trial was presented at the lower or higher flicker frequency. This allowed us to selectively compute the SSVEP response for trials in which the sound was congruent (incongruent) with the stimulus at each stimulation frequency. Note that by design each trial enters this analysis twice, once via the congruent and one via the incongruent visual stimulus. We then computed the SSVEP SNR across all congruent (incongruent) trials at each frequency separately. This procedure was applied to each of the three associations and resulted for each participant in one SNR for each association, congruency condition (or no sound), stimulation frequency and electrode.

Statistical analysis

For the EEG data the comparisons of interest concerned the group-level SSVEP responses between congruent and incongruent pairs. This contrast was derived for each of the three associations and the two stimulation frequencies separately. To test for a difference while correcting for multiple tests across electrodes we relied on a cluster-based permutation procedure (Maris and Oostenveld, 2007). We computed electrode-wise paired t-tests between congruent and incongruent epochs and thresholded these at a first-level threshold of p<0.01 (two-sided). Significant electrode-clusters were aggregated using the cluster-mass (using a minimal cluster-size of 3 electrodes) and the cluster-wise statistics in the actual data were compared to a surrogate distribution obtained from 5000 randomizations. For each significant cluster we report the p-value and the cluster-mass as test-statistics. As effect size we report Cohen's D at the electrode with maximal effect sizes. We also

implemented a post-hoc analysis within the significant cluster obtained for the pitch-size association (c.f. Fig. 3). Here we probed whether the SSVEP response was affected by the general presence of a sound (vs. the no-sound condition). For this we first averaged the SNR across congruent and incongruent conditions and then contrasted this against the no-sound condition (paired t-test). We report the p-value, the t-stat and as effect size the Cohen's D.

To compare the mean response times between conditions (congruent vs. incongruent, or against no-sound), we first averaged the log-transformed response times across trials, removing RTs faster than 150 and slower than 1500 ms as outliers. We then contrasted conditions using a Wilcoxon sign rank test. Effect sizes were obtained using Point biserial correlation (denoted rrb).

Results

SSVEP reveals an effect of association-congruency

The SSVEP stimulation induced robust responses over occipital electrodes. These were quantified as the SNR of the induced power at the two stimulated frequencies relative to the power in immediately neighbouring frequency bands. Figure 2 shows the group-level SNR topographies, which reveal peaks over occipital electrodes as expected.

Figure 2. SSVEP response for each stimulation frequency. The topographies show the group-average of the SNR (mean, n=25). The SNR quantifies the power at each stimulation frequency against the average of the power in immediately neighbouring frequency bins.

We asked whether this SSVEP response differed between congruent and incongruent conditions, hence whether the visual cortical response to a given stimulus is affected by the pitch of the tone presented at the same time. This statistical comparison was implemented for each stimulation frequency and association separately, correcting for multiple comparisons using a cluster-based permutation procedure. For the association between pitch and size we found a significantly stronger SSVEP response for the congruent condition at the lower stimulation frequency (n = 25 participants; p=0.02, tsum = 12.4, Cohen's D =0.70, 4 electrodes; Fig. 3A) but no significant result for the higher stimulation frequency (at p<0.05, see Table 1). The significant cluster resided over occipital electrodes, in line with the SSVEP SNR being strongest over visual cortices. For the other two associations, pitch-hue and pitch-saturation, we found no significant differences in SSVEP responses between conditions (see Table 1 for effect sizes).

Figure 3. Differences in SSVEP responses between conditions. A) Topographies of the electrode-wise t-maps for a difference between congruent and incongruent conditions for each stimulation frequency (for the operational definition of congruent and incongruent for each association see Methods). For the pitch-size association we found a significantly higher SSVEP response for congruent conditions (low pitch & large size or high pitch & small size) over left occipital electrodes for f.low (p=0.02, cluster-based permutation test, N=25 participants). **B)** SSVEP responses for the individual conditions in the pitch-size association for the cluster of 4 occipital electrodes. Dots display the participant-wise data.

Association	f. low				f. high			
	Dmin	Dmax	Tmax	Tmin	Dmin	Dmax	Tmax	Tmin
Size	-0,2	0,7	3,5	-1,2	-0,5	0,4	2,1	-2,7
Hue	-0,5	0,5	2,4	-2,5	-0,5	0,5	2,5	-2,4
Saturation	-0,4	0,5	2,5	-1,8	-0,3	0,5	2,7	-1,5

Table 1. Statistical results for the comparison of congruent and incongruent epochs for each stimulation frequency. The table lists the minimal and maximal (across electrodes) t-value and Cohen's D.

Sounds do not facilitate visual responses in general

Given that many studies have shown generic multisensory benefits for neurophysiological responses in low-level regions we also compared the SSVEP response between trials in which a sound was present compared to trials without sound. For this we focused on the cluster of electrodes revealing a significant congruency-effect in the pitch-size association . We then contrasted the SSVEP response between both sound conditions combined (average of congruent and incongruent) and the no-sound trials (Fig. 3B). This revealed no significant difference (n=25; Size: p=0.25, t=-1.17, Cohen's D=-0.18; Hue: p=0.10, t=-1.69, Cohen's D=-0.25; Saturation: p=0.86, t=-0.17, Cohen's D=0.06). This suggests that the SSVEP response was not generally affected by the presence of a sound but rather points to a specific effect of the pitch-size congruency.

Response times

As a behavioural measure of the crossmodal correspondences we included a speeded detection of a deviant stimulus at the end of the stimulation period. Specifically, one of the two stimuli stayed on longer for a brief moment (33.3 ms) and participants had to indicate which one (left or right). We reasoned that the congruency between the sound, which was already present for the 6 seconds prior to this task, may interact with the detection of this event for the congruent visual stimulus. We hence compared congruent and incongruent trials for each association.

For the pitch-size association responses in the incongruent trials were significantly faster compared to congruent trials (n=25, Wilcoxon sign-rank test, Z=2.25, p = 0.02, rrb =0.51). No significant results were found for the other associations (Hue: Z=1.06, p = 0.29, rrb =0.24; Saturation: Z=-0.39, p = 0.69, rrb =-0.08). We also probed for a general facilitation of reaction times by the presence of any sound, contrasting response times on trials with a sound and those without sound. This revealed that participants were significantly faster when a sound was present (Size= Z=4.34, p < 0.01, rrb =0.99; Hue: Z=4.05, p < 0.01, rrb =0.93; Saturation: Z=4.37, p < 0.01, rrb =1). Hence in the behavioural data we observed a general facilitation of a sound on visual task performance as well as a congruency effect for the pitch-size association.

Figure 4. Behavioural response times. The graphs show the log-transformed response times for the three conditions and each of the association blocks. Dots indicate individual participants (n = 25).

Discussion

Humans consistently associate properties of stimuli in different sensory modalities with each other (Spence, 2011; Spence and Deroy, 2013; Spence and Sathian, 2020). Well known examples are the association between pitch and size or between phonological features and visual shapes such as in the Bouba-Kiki effect (Kohler 1929, 1947). In the latter the word Bouba is associated with the round and Kiki with the more spiky among two visual shapes. Despite considerable work on multisensory perception in general, the neurophysiological correlates of such associations remain debated (Spence and Sathian, 2020). While some studies have argued for an involvement of low-level sensory regions others have implied high-level processes in association cortices (Bien et al., 2012; Kovic et al., 2010; McCormick et al., 2021, 2018; Peiffer-Smadja and Cohen, 2019; Stekelenburg and Keetels, 2016). To address this question, we developed a paradigm based on the SSVEP procedure to test the specific hypothesis that neurophysiological correlates of audio-visual associations can be found in activity arising from functionally defined early visual cortices. Our results reveal a congruency-modulation of the SSVEP for the pitch-size association over occipital electrodes that was accompanied by a behavioural congruency effect, in accordance with a low-level correlate of this association. In contrast, the associations between pitch and hue or saturation respectively did not reveal significant neurophysiological or behavioural effects.

Robust associations of pitch with size but not hue or saturation

The association between the pitch of a sound and the size of a visual object possibly arises from the physical properties of the objects surrounding us, where larger animals or objects produce lower frequency sounds compared to smaller ones (Bee et al., 2000; Carello et al., 1998; Tecumseh Fitch and Reby, 2001). This association has been reported in numerous studies using implicit behavioural measures such as reaction times or explicit matching responses (Bien et al., 2012; Evans and Treisman, 2010; Mondloch and Maurer, 2004; Parise and Spence, 2012).

Our data support the robustness of this association both at the behavioural and neurophysiological level. In contrast, the associations between pitch and chromatic properties such as a specific hue or saturation level have been reported in some but not in other studies (Anikin and Johansson, 2019; Bernstein et al., 1971; Caivano, 1994; Hamilton-Fletcher et al., 2017; Melara, 1989; Moos et al., 2014; Simpson et. al, 1956; Ward et al., 2006). Furthermore, these studies disagree in which specific features were associated. For example, some reported the association of high pitch with yellow or green relative to blue or purple (Simpson et al., 1956; Moos et al., 2014; Giannakis 2001, Orlandatou 2012), while others suggested that the specific pitch-hue or pitch-saturation associations may depend on the precise values of acoustic pitch and visual features used (Anikin and Johansson, 2019; Hamilton-Fletcher et al., 2017; Wrembel, 2009) or the context of other sound frequencies presented in the same experiment (Ward et al., 2006). Importantly, some studies did not control for then subjectively perceived lightness when manipulating hue or saturation (de Thornley Head, 2006; Simpson et a., 1956; Ward et al., 2006; but see Anikin and Johansson, 2019), which may be crucial given that an association between pitch and lightness has been reported as well (Brunel et al., 2015; Hubbard, 1996; Marks, 1987, 1975, 1974). Hence, some of the discrepancies in the previous literature may be a result of subjective differences of the respective stimuli along multiple chromatic dimensions. To minimise

such confounding factors, we relied on stimuli that were calibrated on a participant-level for their subjective hue, saturation and lightness. Based on these stimuli, we did not find significant congruency effects for pitch-hue or pitch-saturation, neither in the behavioural nor in the neurophysiological data. Hence, our results suggest that these associations may be not as robust as the association between pitch and size, or may indeed depend on the precise nature of the specific stimuli or of the behavioural paradigm employed as some have speculated (Hamilton-Fletcher et al., 2017)

The SSVEP reveals low-level correlates of associations

We introduced an SSVEP based paradigm to probe crossmodal associations for two reasons. First, given the known origin of the SSVEP response in visual cortex it allows establishing whether or not a neurophysiological correlate from these sensory regions relates to crossmodal associations. Second, it allowed us to probe the response to simultaneously presented congruent and incongruent visual stimuli with the same sound, and therefore directly taps into the characteristic binding property of crossmodal associations (Parise and Spence, 2009).

Our results reveal a stronger SSVEP response at 7 HZ over occipital electrodes for the visual stimulus whose size was congruent with the pitch. This provides direct evidence for an early and possibly bottom-up neurophysiological correlate of crossmodal associations (Kovic et al., 2010; Molholm et al., 2004; Peiffer-Smadja and Cohen, 2019). Hence our results are in line with previous speculations suggesting a low-level or short-latency correlate of cross-modal associations and similar object-related congruency effects (Brang et al., 2022; Kayser et al., 2017; Kovic et al., 2010; Molholm et al., 2004; Peiffer-Smadja and Cohen, 2019; Sadaghiani et al., 2009).

These results are also in line with the current working model of multisensory processing. This model proposes that interactions in early sensory regions are governed by low-level congruencies of the multiple inputs, such as their temporal synchronization or spatial alignment, while multisensory integration at higher association cortices becomes more functionally and task-specific (Bizley et al., 2016; Cao et al., 2019; Noppeney, 2021; Rohe and Noppeney, 2015). Our results show that multisensory processes in early visual cortices are also sensitive to congruencies pertaining to basic features such as size and pitch. Given that these features are highly correlated in everyday life based on the physical attributes of the objects surrounding us this suggests that early crossmodal interactions may be adapted to our natural environment. Hence, the low-level signature of the pitch-size association may reflect the binding or grouping of multisensory features often occurring in a correlated manner, possibly to aid the process of multisensory causal inference. Multisensory causal inference helps the brain to decide whether or which sensory features to combine for perception and which not to (Körding et al., 2007; Noppeney, 2021). While recent studies have assigned an important role of frontal regions in this process (Cao et al., 2019; Ferrari and Noppeney, 2021; Rohe and Noppeney, 2015), consequences of this process may relate to the grouping of sensory features in low-level regions. This speculation is also supported by our finding that the congruency effect for the SSVEP response was not accompanied by a general enhancement from the presence of an acoustic stimulus per se. This suggests that the association-related auditory influence on visual cortex is distinct and more specific than the previously reported general modulation of visual responses by acoustic inputs (Brang et al., 2022; Lakatos et al., 2009; Murray et al., 2016; Petro et al., 2017; Wang et al., 2008)

Our data cannot disambiguate the contribution of feed-forward and feed-back processes given that the SSVEP response was elicited and analysed over a longer stimulation interval. Hence it is possible that the neurophysiological signature over occipital regions was generated in response to multisensory feedback, for example as a result of causal inference, but possibly also including processes such as attention or memory. In fact, in our paradigm the sound may have facilitated the perceptual binding of the auditory input with the congruent of the two visual shapes via processes the spread of attention across modalities (Talsma et al., 2010). This attention-spread may be one factor contributing to the enhanced SSVEP response, which is known to be sensitive to attention effects (Appelbaum and Norcia, 2009; Lauritzen et al., 2010; Norcia et al., 2015). An alternative account would be predictive coding, whereby auditory regions relay information that is predictive of concurrent or expected features to visual cortices, possibly via direct cortico-cortical connections (Falchier et al., 2002; Garner and Keller, 2022; Morrell, 1972; Petro et al., 2017). Hence while our results demonstrate the existence of a specific neurophysiological signature of cross-modal correspondences, future work still needs to uncover the specific functional and computational benefits.

For the SSVEP stimulation we opted for frequencies in the alpha range, which are known to maximise the involvement of early visual cortices, in particular primary visual cortex (Di Russo et al., 2007; Lauritzen et al., 2010; Vanegas et al., 2013). Still, the specific generators of the SSVEP response may depend also on the stimulation frequency, which higher stimulation frequencies emphasising responses from earlier visual regions based on the short response latencies of neurons at earlier stages of the visual pathway (Alonso-Prieto et al., 2013; Appelbaum and Norcia, 2009; Cottereau et al., 2011; Liu-Shuang et al., 2014). The occipital topographies of the SSVEP responses and the pitch-size congruency effect obtained in this study are in line with an origin in early visual cortical regions. Still, further work is required to understand why we found the strongest (and significant) congruency effect at a stimulation frequency around 7 but not at 11 Hz. This may provide evidence about the specific brain regions giving rise to this signature of crossmodal associations thought studies with higher spatial resolution or functional specificity are required to study this in more detail.

We found signatures of the pitch-size association in both the SSVEP and the behavioural data. Yet, the SSVEP response to congruent visual stimuli was enhanced, while the response times were longer when the deviant stimulus was congruent with the sound. While these two signatures seemingly have opposing directions we see reasons for why this may be plausible. First, the SSVEP response was obtained over a 6 second stimulation period. In contrast, the behavioural response was obtained only at the very end of the stimulation period, hence after prolonged exposure to the stimuli. Such prolonged stimulation may induce adaptation effect which can dissociate sensory responses and behaviour (Blakemore and Campbell, 1969; Blakemore and Sutton, 1969; Gibson and Radner, 1937; Leopold et al., 2005). Second, the stronger SSVEP responses for congruent visual stimuli may reflect the enhanced binding of these specific visual features with the sound. Binding the congruent visual object with the sound may foster its perceptual constancy or persistence, which may then be detrimental for the detection of a change in the congruent (bound) compared to the incongruent (not bound) object. For example, participants have higher thresholds for discriminating spatial or temporal

conflicts in synesthetically matched compared to mismatched audio-visual stimuli (Parise and Spence, 2009). Multisensory binding may impede or slow down the neurophysiological access to the individual audio and visual sensory components, therefore in our case reducing the behavioural detection of changes relative to the congruent pair (Ernst and Banks, 2002; Ernst, 2006). Third, other studies measuring SSVEPs also found opposing effects on SSVEP response strength and behaviour. In general, the SSVEP response provides an index of visual cortical neural responses. But since the EEG reflects pre- and post-synaptic responses this does not only include the mere processing of sensory input but reflects the aggregate effect of all intracortical processes. For example, opposing effects between SSVEP and behavioural performance have been found for several tasks such as working memory, spatial attention and object recognition (Kaspar et al., 2010; Silberstein et al., 2001; Walter et al., 2012) suggesting that an apparent mismatch between SSVEP signatures and behavioural data may be a more general phenomenon.

Unspecific multisensory benefits for behaviour

Our behavioural data also speak in favour of a general facilitation of visual task performance by the presence of a sound compared to the no-sound condition. Similar sound-induced benefits of visual perception have been reported for task-irrelevant or uninformative sounds in a range of paradigms (Fiebelkorn et al., 2011; Frassinetti et al., 2002; Huang et al., 2011; Jaekl and Soto-Faraco, 2010; Lippert et al., 2007). The present data corroborate the existence of such effects. We did not see a general sound-modulation of the SSVEP responses, which speaks for a different neurophysiological origin of these unspecific effects compared to the cross-modal association. However, unspecific multisensory effects were also attributed to early sensory cortices (Brang et al., 2022; Kayser et al., 2009; Mercier and Cappe, 2020; Pérez-Bellido et al., 2022). Hence, it is possible that both association-related and unspecific multisensory processes influence low-level sensory responses but do so by different pathways and neural mechanisms and give rise to distinct neurophysiological signatures in the EEG.

General conclusion

Despite their prominence in behavioural and psychophysical studies, the neurophysiological mechanisms underlying cross-modal associations remain unclear. Using an SSVEP stimulation protocol known to engage early visual cortices we show that the association between pitch and the size of a visual object influences activity in these visual regions. Hence our data support a low-level account of crossmodal associations.

References

Alonso-Prieto, E., Belle, G.V., Liu-Shuang, J., Norcia, A.M., Rossion, B., 2013. The 6Hz fundamental stimulation frequency rate for individual face discrimination in the right occipito-temporal cortex. Neuropsychologia 51, 2863–2875. https://doi.org/10.1016/j.neuropsychologia.2013.08.018

Alpers, G.W., Ruhleder, M., Walz, N., Mühlberger, A., Pauli, P., 2005. Binocular rivalry between emotional and neutral stimuli: A validation using fear conditioning and EEG. Int. J. Psychophysiol., Neurobiology of Fear and Disgust 57, 25–32. https://doi.org/10.1016/j.ijpsycho.2005.01.008

Anikin, A., Johansson, N., 2019. Implicit associations between individual properties of color and sound. Atten. Percept. Psychophys. 81, 764–777. https://doi.org/10.3758/s13414-018-01639-7

Appelbaum, L.G., Norcia, A.M., 2009. Attentive and pre-attentive aspects of figural processing. J. Vis. 9, 18. https://doi.org/10.1167/9.11.18

Appelbaum, L.G., Wade, A.R., Pettet, M.W., Vildavski, V.Y., Norcia, A.M., 2008. Figure–ground interaction in the human visual cortex. J. Vis. 8, 8. https://doi.org/10.1167/8.9.8

Appelbaum, L.G., Wade, A.R., Vildavski, V.Y., Pettet, M.W., Norcia, A.M., 2006. Cue-Invariant Networks for Figure and Background Processing in Human Visual Cortex. J. Neurosci. 26, 11695–11708. https://doi.org/10.1523/JNEUROSCI.2741-06.2006

Bee, M.A., Perrill, S.A., Owen, P.C., 2000. Male green frogs lower the pitch of acoustic signals in defense of territories: a possible dishonest signal of size? Behav. Ecol. 11, 169–177. https://doi.org/10.1093/beheco/11.2.169

Bernstein, I.H., Eason, T.R., Schurman, D.L., 1971. Hue-Tone Sensory Interaction: A Negative Result. Percept. Mot. Skills 33, 1327–1330. https://doi.org/10.2466/pms.1971.33.3f.1327

Bien, N., ten Oever, S., Goebel, R., Sack, A.T., 2012. The sound of size: Crossmodal binding in pitch-size synesthesia: A combined TMS, EEG and psychophysics study. NeuroImage, Neuroergonomics: The human brain in action and at work 59, 663–672. https://doi.org/10.1016/j.neuroimage.2011.06.095

Bizley, J.K., Jones, G.P., Town, S.M., 2016. Where are multisensory signals combined for perceptual decision-making? Curr. Opin. Neurobiol., Systems neuroscience 40, 31–37. https://doi.org/10.1016/j.conb.2016.06.003

Blakemore, C., Campbell, F.W., 1969. On the existence of neurones in the human visual system selectively sensitive to the orientation and size of retinal images. J. Physiol. 203, 237–260. https://doi.org/10.1113/jphysiol.1969.sp008862

Blakemore, C., Sutton, P., 1969. Size Adaptation: A New Aftereffect. Science 166, 245–247. https://doi.org/10.1126/science.166.3902.245

Brang, D., Plass, J., Sherman, A., Stacey, W.C., Wasade, V.S., Grabowecky, M., Ahn, E., Towle, V.L., Tao, J.X., Wu, S., Issa, N.P., Suzuki, S., 2022. Visual cortex responds to sound onset and offset during passive listening. J. Neurophysiol. 127, 1547–1563. https://doi.org/10.1152/jn.00164.2021

Brown, R.J., Norcia, A.M., 1997. A method for investigating binocular rivalry in real-time with the steady-state VEP. Vision Res. 37, 2401–2408. https://doi.org/10.1016/S0042-6989(97)00045-X

Brunel, L., Carvalho, P.F., Goldstone, R.L., 2015. It does belong together: cross-modal correspondences influence cross-modal integration during perceptual learning. Front. Psychol. 6.

Caivano, J.L., 1994. Color and Sound: Physical and Psychophysical Relations*. Color Res. Appl. 19, 126–133. https://doi.org/10.1111/j.1520-6378.1994.tb00072.x

Campos Viola, F., Thorne, J., Edmonds, B., Schneider, T., Eichele, T., Debener, S., 2009. Semi-automatic identification of independent components representing EEG artifact. Clin. Neurophysiol. 120, 868–877. https://doi.org/10.1016/j.clinph.2009.01.015

Cao, Y., Summerfield, C., Park, H., Giordano, B.L., Kayser, C., 2019. Causal Inference in the Multisensory Brain. Neuron 102, 1076-1087.e8. https://doi.org/10.1016/j.neuron.2019.03.043

Carello, C., Anderson, K.L., Kunkler-Peck, A.J., 1998. Perception of Object Length by Sound. Psychol. Sci. 9, 211–214. https://doi.org/10.1111/1467-9280.00040

Cottereau, B.R., McKee, S.P., Ales, J.M., Norcia, A.M., 2011. Disparity-Tuned Population Responses from Human Visual Cortex. J. Neurosci. 31, 954–965. https://doi.org/10.1523/JNEUROSCI.3795-10.2011

Di Russo, F., Pitzalis, S., Aprile, T., Spitoni, G., Patria, F., Stella, A., Spinelli, D., Hillyard, S.A., 2007. Spatiotemporal analysis of the cortical sources of the steady-state visual evoked potential. Hum. Brain Mapp. 28, 323–334. https://doi.org/10.1002/hbm.20276

Dreksler, N., Spence, C., 2019. A Critical Analysis of Colour–Shape Correspondences: Examining the Replicability of Colour–Shape Associations. -Percept. 10, 2041669519834042. https://doi.org/10.1177/2041669519834042

Drew, P., Sayres, R., Watanabe, K., Shimojo, S., 2001. Pupillary response to chromatic flicker. Exp. Brain Res. 136, 256–262. https://doi.org/10.1007/s002210000605

Evans, K.K., Treisman, A., 2010. Natural cross-modal mappings between visual and auditory features. J. Vis. 10, 6. https://doi.org/10.1167/10.1.6

Falchier, A., Clavagnier, S., Barone, P., Kennedy, H., 2002. Anatomical Evidence of Multimodal Integration in Primate Striate Cortex. J. Neurosci. 22, 5749–5759. https://doi.org/10.1523/JNEUROSCI.22-13-05749.2002

Ferrari, A., Noppeney, U., 2021. Attention controls multisensory perception via two distinct mechanisms at different levels of the cortical hierarchy. PLOS Biol. 19, e3001465. https://doi.org/10.1371/journal.pbio.3001465

Fiebelkorn, I.C., Foxe, J.J., Butler, J.S., Mercier, M.R., Snyder, A.C., Molholm, S., 2011. Ready, Set, Reset: Stimulus-Locked Periodicity in Behavioral Performance Demonstrates the Consequences of Cross-Sensory Phase Reset. J. Neurosci. 31, 9971–9981. https://doi.org/10.1523/JNEUROSCI.1338-11.2011

Frassinetti, F., Bolognini, N., Làdavas, E., 2002. Enhancement of visual perception by crossmodal visuo-auditory interaction. Exp. Brain Res. 147, 332–343. https://doi.org/10.1007/s00221-002-1262-y

Gallace, A., Spence, C., 2006. Multisensory synesthetic interactions in the speeded classification of visual size. Percept. Psychophys. 68, 1191–1203. https://doi.org/10.3758/BF03193720

Garner, A.R., Keller, G.B., 2022. A cortical circuit for audio-visual predictions. Nat. Neurosci. 25, 98–105. https://doi.org/10.1038/s41593-021-00974-7

Giannakis, K., 2001. Sound mosaics: a graphical user interface for sound synthesis based on audio-visual associations. (phd). Middlesex University.

Gibson, J.J., Radner, M., 1937. Adaptation, after-effect and contrast in the perception of tilted lines. I. Quantitative studies. J. Exp. Psychol. 20, 453–467. https://doi.org/10.1037/h0059826

Hamilton-Fletcher, G., Witzel, C., Reby, D., Ward, J., 2017. Sound Properties Associated With Equiluminant Colours. Multisensory Res. 30, 337–362. https://doi.org/10.1163/22134808-00002567

Hipp, J., Siegel, M., 2013. Dissociating neuronal gamma-band activity from cranial and ocular muscle activity in EEG. Front. Hum. Neurosci. 7.

Huang, P.-C., Yeh, S.-L., Chen, Y.-C., Spence, C., 2011. Synchronous Sounds Enhance Visual Sensitivity without Reducing Target Uncertainty. Seeing Perceiving 24, 623–638. https://doi.org/10.1163/187847611X603765

Hubbard, T.L., 1996. Synesthesia-like Mappings of Lightness, Pitch, and Melodic Interval. Am. J. Psychol. 109, 219–238. https://doi.org/10.2307/1423274

Jaekl, P.M., Soto-Faraco, S., 2010. Audiovisual contrast enhancement is articulated primarily via the M-pathway. Brain Res. 1366, 85–92. https://doi.org/10.1016/j.brainres.2010.10.012

Kaspar, K., Hassler, U., Martens, U., Trujillo-Barreto, N., Gruber, T., 2010. Steady-state visually evoked potential correlates of object recognition. Brain Res. 1343, 112–121. https://doi.org/10.1016/j.brainres.2010.04.072

Kayser, C., Logothetis, N.K., Panzeri, S., 2010. Visual Enhancement of the Information Representation in Auditory Cortex. Curr. Biol. 20, 19–24. https://doi.org/10.1016/j.cub.2009.10.068

Kayser, C., Petkov, C.I., Logothetis, N.K., 2009. Multisensory interactions in primate auditory cortex: fMRI and electrophysiology. Hear. Res., Multisensory integration in auditory and auditory-related areas of cortex 258, 80–88. https://doi.org/10.1016/j.heares.2009.02.011

Kayser, S.J., Kayser, C., 2020. Shared Physiological Correlates of Multisensory and Expectation-Based Facilitation. eNeuro 7. https://doi.org/10.1523/ENEURO.0435-19.2019

Kayser, S.J., Philiastides, M.G., Kayser, C., 2017. Sounds facilitate visual motion discrimination via the enhancement of late occipital visual representations. NeuroImage 148, 31–41. https://doi.org/10.1016/j.neuroimage.2017.01.010

Körding, K.P., Beierholm, U., Ma, W.J., Quartz, S., Tenenbaum, J.B., Shams, L., 2007. Causal inference in multisensory perception. PloS One 2, e943. https://doi.org/10.1371/journal.pone.0000943

Kovic, V., Plunkett, K., Westermann, G., 2010. The shape of words in the brain. Cognition 114, 19–28. https://doi.org/10.1016/j.cognition.2009.08.016

Lakatos, P., O'Connell, M.N., Barczak, A., Mills, A., Javitt, D.C., Schroeder, C.E., 2009. The Leading Sense: Supramodal Control of Neurophysiological Context by Attention. Neuron 64, 419–430. https://doi.org/10.1016/j.neuron.2009.10.014

Lauritzen, T.Z., Ales, J.M., Wade, A.R., 2010. The effects of visuospatial attention measured across visual cortex using source-imaged, steady-state EEG. J. Vis. 10, 39. https://doi.org/10.1167/10.14.39

Leopold, D.A., Rhodes, G., Müller, K.-M., Jeffery, L., 2005. The dynamics of visual adaptation to faces. Proc. R. Soc. B Biol. Sci. 272, 897–904. https://doi.org/10.1098/rspb.2004.3022

Lippert, M., Logothetis, N.K., Kayser, C., 2007. Improvement of visual contrast detection by a simultaneous sound. Brain Res. 1173, 102–109. https://doi.org/10.1016/j.brainres.2007.07.050

Liu-Shuang, J., Norcia, A.M., Rossion, B., 2014. An objective index of individual face discrimination in the right occipito-temporal cortex by means of fast periodic oddball stimulation. Neuropsychologia 52, 57–72. https://doi.org/10.1016/j.neuropsychologia.2013.10.022

Marks, L.E., 1987. On cross-modal similarity: Auditory–visual interactions in speeded discrimination. J. Exp. Psychol. Hum. Percept. Perform. 13, 384–394. https://doi.org/10.1037/0096-1523.13.3.384

Marks, L.E., 1975. On colored-hearing synesthesia: Cross-modal translations of sensory dimensions. Psychol. Bull. 82, 303–331. https://doi.org/10.1037/0033-2909.82.3.303

Marks, L.E., 1974. On Associations of Light and Sound: The Mediation of Brightness, Pitch, and Loudness. Am. J. Psychol. 87, 173–188. https://doi.org/10.2307/1422011

McCormick, K., Lacey, S., Stilla, R., Nygaard, L.C., Sathian, K., 2021. Neural Basis of the Sound-Symbolic Crossmodal Correspondence Between Auditory Pseudowords and Visual Shapes. Multisensory Res. 35, 29–78. https://doi.org/10.1163/22134808-bja10060

McCormick, K., Lacey, S., Stilla, R., Nygaard, L.C., Sathian, K., 2018. Neural basis of the crossmodal correspondence between auditory pitch and visuospatial elevation. Neuropsychologia 112, 19–30. https://doi.org/10.1016/j.neuropsychologia.2018.02.029

Melara, R.D., 1989. Dimensional interaction between color and pitch. J. Exp. Psychol. Hum. Percept. Perform. 15, 69–79. https://doi.org/10.1037/0096-1523.15.1.69

Mercier, M.R., Cappe, C., 2020. The interplay between multisensory integration and perceptual decision making. NeuroImage 222, 116970. https://doi.org/10.1016/j.neuroimage.2020.116970

Molholm, S., Ritter, W., Javitt, D.C., Foxe, J.J., 2004. Multisensory Visual–Auditory Object Recognition in Humans: a High-density Electrical Mapping Study. Cereb. Cortex 14, 452–465. https://doi.org/10.1093/cercor/bhh007

Mondloch, C.J., Maurer, D., 2004. Do small white balls squeak? Pitch-object correspondences in young children. Cogn. Affect. Behav. Neurosci. 4, 133–136. https://doi.org/10.3758/CABN.4.2.133

Moos, A., Smith, R., Miller, S.R., Simmons, D.R., 2014. Cross-Modal Associations in Synaesthesia: Vowel Colours in the Ear of the Beholder. -Percept. 5, 132–142. https://doi.org/10.1068/i0626

Morrell, F., 1972. Visual System's View of Acoustic Space. Nature 238, 44–46. https://doi.org/10.1038/238044a0

Müller, M.M., Hübner, R., 2002. Can the Spotlight of Attention Be Shaped Like a Doughnut? Evidence From Steady-State Visual Evoked Potentials. Psychol. Sci. 13, 119–124. https://doi.org/10.1111/1467-9280.00422

Murray, M.M., Thelen, A., Thut, G., Romei, V., Martuzzi, R., Matusz, P.J., 2016. The multisensory function of the human primary visual cortex. Neuropsychologia, Special Issue: Functional Selectivity in Perceptual and Cognitive Systems - A Tribute to Shlomo Bentin (1946-2012) 83, 161–169. https://doi.org/10.1016/j.neuropsychologia.2015.08.011

Noppeney, U., 2021. Perceptual Inference, Learning, and Attention in a Multisensory World. Annu. Rev. Neurosci. 44, 449–473. https://doi.org/10.1146/annurev-neuro-100120-085519

Norcia, A.M., Appelbaum, L.G., Ales, J.M., Cottereau, B.R., Rossion, B., 2015. The steady-state visual evoked potential in vision research: A review. J. Vis. 15, 4. https://doi.org/10.1167/15.6.4

O'Beirne, G.A., Patuzzi, R.B., 1999. Basic properties of the sound-evoked post-auricular muscle response (PAMR). Hear. Res. 138, 115–132. https://doi.org/10.1016/s0378-5955(99)00159-8

Oostenveld, R., Fries, P., Maris, E., Schoffelen, J.-M., 2011. FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput. Intell. Neurosci. 2011, 156869. https://doi.org/10.1155/2011/156869

Orlandatou, K., n.d. The Role of Pitch and Timbre in the Synaesthetic Experience 8.

Parise, C., Spence, C., 2008. Synesthetic congruency modulates the temporal ventriloquism effect. Neurosci. Lett. 442, 257–261. https://doi.org/10.1016/j.neulet.2008.07.010

Parise, C.V., Spence, C., 2012. Audiovisual crossmodal correspondences and sound symbolism: a study using the implicit association test. Exp. Brain Res. 220, 319–333. https://doi.org/10.1007/s00221-012-3140-6

Parise, C.V., Spence, C., 2009. 'When Birds of a Feather Flock Together': Synesthetic Correspondences Modulate Audiovisual Integration in Non-Synesthetes. PLOS ONE 4, e5664. https://doi.org/10.1371/journal.pone.0005664

Park, H., Kayser, C., 2021. The Neurophysiological Basis of the Trial-Wise and Cumulative Ventriloquism Aftereffects. J. Neurosci. 41, 1068–1079. https://doi.org/10.1523/JNEUROSCI.2091-20.2020

Peiffer-Smadja, N., Cohen, L., 2019. The cerebral bases of the bouba-kiki effect. NeuroImage 186, 679–689. https://doi.org/10.1016/j.neuroimage.2018.11.033

Pérez-Bellido, A., Spaak, E., Lange, F.P. de, 2022. Seeing sounds: Neural mechanisms underlying auditory contributions to visual detection. https://doi.org/10.1101/2022.01.19.476187

Petro, L.S., Paton, A.T., Muckli, L., 2017. Contextual modulation of primary visual cortex by auditory signals. Philos. Trans. R. Soc. B Biol. Sci. 372, 20160104. https://doi.org/10.1098/rstb.2016.0104

Rohe, T., Noppeney, U., 2015. Cortical Hierarchies Perform Bayesian Causal Inference in Multisensory Perception. PLOS Biol. 13, e1002073. https://doi.org/10.1371/journal.pbio.1002073

Sadaghiani, S., Maier, J.X., Noppeney, U., 2009. Natural, Metaphoric, and Linguistic Auditory Direction Signals Have Distinct Influences on Visual Motion Processing. J. Neurosci. 29, 6490–6499. https://doi.org/10.1523/JNEUROSCI.5437-08.2009

Schroeder, C.E., Foxe, J., 2005. Multisensory contributions to low-level, 'unisensory' processing. Curr. Opin. Neurobiol., Sensory systems 15, 454–458. https://doi.org/10.1016/j.conb.2005.06.008

Sciortino, P., Kayser, C., 2022a. The rubber hand illusion is accompanied by a distributed reduction of alpha and beta power in the EEG. PLOS ONE 17, e0271659. https://doi.org/10.1371/journal.pone.0271659

Sciortino, P., Kayser, C., 2022b. Multivariate Analysis of Evoked Responses during the Rubber Hand Illusion Suggests a Temporal Parcellation into Manipulation and Illusion-Specific Correlates. eNeuro 9. https://doi.org/10.1523/ENEURO.0355-21.2021

Silberstein, R.B., Nunez, P.L., Pipingas, A., Harris, P., Danieli, F., 2001. Steady state visually evoked potential (SSVEP) topography in a graded working memory task. Int. J. Psychophysiol. 42, 219–232. https://doi.org/10.1016/S0167-8760(01)00167-2

Spence, C., 2011. Crossmodal correspondences: A tutorial review. Atten. Percept. Psychophys. 73, 971–995. https://doi.org/10.3758/s13414-010-0073-7

Spence, C., Deroy, O., 2013. How automatic are crossmodal correspondences? Conscious. Cogn. 22, 245–260. https://doi.org/10.1016/j.concog.2012.12.006

Spence, C., Sathian, K., 2020. Chapter 11 - Audiovisual crossmodal correspondences: Behavioral consequences and neural underpinnings, in: Sathian, K., Ramachandran, V.S. (Eds.), Multisensory Perception. Academic Press, pp. 239–258. https://doi.org/10.1016/B978-0-12-812492-5.00011-5

Stekelenburg, J.J., Keetels, M., 2016. The effect of synesthetic associations between the visual and auditory modalities on the Colavita effect. Exp. Brain Res. 234, 1209–1219. https://doi.org/10.1007/s00221-015-4363-0

Sutoyo, D., Srinivasan, R., 2009. Nonlinear SSVEP responses are sensitive to the perceptual binding of visual hemifields during conventional 'eye' rivalry and interocular 'percept' rivalry. Brain Res. 1251, 245–255. https://doi.org/10.1016/j.brainres.2008.09.086

Talsma, D., Senkowski, D., Soto-Faraco, S., Woldorff, M.G., 2010. The multifaceted interplay between attention and multisensory integration. Trends Cogn. Sci. 14, 400–410. https://doi.org/10.1016/j.tics.2010.06.008

Tecumseh Fitch, W., Reby, D., 2001. The descended larynx is not uniquely human. Proc. R. Soc. Lond. B Biol. Sci. 268, 1669–1675. https://doi.org/10.1098/rspb.2001.1704

Tello, R.J.M.G., Müller, S.M.T., Ferreira, A., Bastos, T.F., 2015. Comparison of the influence of stimuli color on Steady-State Visual Evoked Potentials. Res. Biomed. Eng. 31, 218–231. https://doi.org/10.1590/2446-4740.0739

Vanegas, M.I., Blangero, A., Kelly, S.P., 2013. Exploiting individual primary visual cortex geometry to boost steady state visual evoked potentials. J. Neural Eng. 10, 036003. https://doi.org/10.1088/1741-2560/10/3/036003

Vetter, P., Smith, F.W., Muckli, L., 2014. Decoding sound and imagery content in early visual cortex. Curr. Biol. CB 24, 1256–1262. https://doi.org/10.1016/j.cub.2014.04.020

Vialatte, F.-B., Maurice, M., Dauwels, J., Cichocki, A., 2010. Steady-state visually evoked potentials: Focus on essential paradigms and future perspectives. Prog. Neurobiol. 90, 418–438. https://doi.org/10.1016/j.pneurobio.2009.11.005

Walter, S., Quigley, C., Andersen, S.K., Mueller, M.M., 2012. Effects of overt and covert attention on the steady-state visual evoked potential. Neurosci. Lett. 519, 37–41. https://doi.org/10.1016/j.neulet.2012.05.011

Wang, Y., Celebrini, S., Trotter, Y., Barone, P., 2008. Visuo-auditory interactions in the primary visual cortex of the behaving monkey: Electrophysiological evidence. BMC Neurosci. 9, 79. https://doi.org/10.1186/1471-2202-9-79

Ward, J., Huckstep, B., Tsakanikos, E., 2006. Sound-Colour Synaesthesia: to What Extent Does it Use Cross-Modal Mechanisms Common to us All? Cortex 42, 264–280. https://doi.org/10.1016/S0010-9452(08)70352-6

Wrembel, M., 2009. On Hearing Colours — Cross-Modal Associations in Vowel Perception in a Non-Synaesthetic Population. Pozn. Stud. Contemp. Linguist. 45, 595–612. https://doi.org/10.2478/v10010-009-0028-0

Zuanazzi, A., Noppeney, U., 2020. The Intricate Interplay of Spatial Attention and Expectation: a Multisensory Perspective. Multisensory Res. 33, 383–416. https://doi.org/10.1163/22134808-20201482