Kinesin-12 KLP-18 contributes to the kinetochore-microtubule poleward flux during the metaphase of *C. elegans* one-cell embryo.

Nina Soler[1,*], Mathis Da Silva[1,*], Christophe Tascon[1], Laurent Chesneau[1], Pauline Foliard[1,†], Hélène Bouvrais[1], Sylvain Pastezeur[1], Loïc Le Marrec[2], Jacques Pecreaux[1,‡]

[1] CNRS, Univ Rennes, IGDR (Institut de Génétique et Développement de Rennes) – UMR 6290, F-35000 Rennes, France.

[2] CNRS, Univ Rennes, IRMAR (Institut de Recherche Mathématique de Rennes) – UMR 6625, F-35000 Rennes, France.

* Equally contributing authors

[†] Present address: IBGC, Bordeaux, France

[‡] Corresponding author: jacques.pecreaux@univ-rennes.fr

Abstract:

The mitotic spindle, a key structure to partition chromosomes during cell division, connects its poles to the chromosomes through microtubules. Their plus-ends, oriented towards the chromosomes, exhibit dynamic instability crucial for kinetochores' correct attachment. Involved in this process, the poleward flux implicates the displacement of microtubules towards the spindle poles, coordinated with polymerisation at the plus ends. The mechanisms behind this are diverse. It includes treadmilling powered by microtubule depolymerisation at the spindle poles, sliding of spindle microtubules by molecular motors like Kinesin-5, and pushing microtubules away from the chromosomes by chromokinesins. Interestingly, no such flux was reported in the *Caenorhabditis elegans* zygote, although all proteins contributing to flux in mammals have homologous in the nematode.

To explore this, we fluorescently labelled microtubules and conducted photobleaching. We found no global poleward flow; the bleached zone's edges moved inward. The centrosome-side front motion was caused by dynamic instability, while the chromosome-side front exhibited faster recovery, suggesting an additional mechanism. This extra velocity was localised near chromosomes, indicating that only kinetochore microtubules may undergo flux. Consistently, this flux depended on proteins ensuring the chromosome attachment and growth of the kinetochore microtubules, notably NDC-80, CLS-2[CLASP], and ZYG-9[XMAP215]. Furthermore, this flux decreased as metaphase progressed and attachments transitioned from side- to end-on; it was reduced by SKA-1 recruitment. Traditional treadmilling was unlikely as many kinetochore microtubules do not reach spindle poles in the zygote spindle. Conversely, the depletion of kinesin-12 KLP-18[KIF15], which cross-links and focuses microtubules at meiosis, reduced the front rate. Ultimately, we propose that the sole kinetochore microtubules slide along spindle microtubules likely powered by KLP-18, contrasting with solid displacement in other systems, aligning with observations in human cells of decreasing flux with increasing chromosome distance.

INTRODUCTION

During cell division, the mitotic spindle ensures a faithful partitioning of the genetic material into two identical sets distributed to each daughter cell. This specialised structure is assembled from microtubules (MTs) and their associated proteins (MAPs), especially molecular motors and cross-linkers. The microtubules are polarised semi-flexible polymers which alternate growth and shrinkage, and their plus ends are more dynamic than their minus ends (Akhmanova and Steinmetz, 2015). In the nematode C. *elegans* zygote, there are two types of microtubules in the spindle: kinetochore microtubules (kMT) with plus-ends anchored at the chromosomes and minus-ends spread along the spindle; and spindle microtubules (sMT) with minus-ends at the centrosomes and plus-ends not contacting the kinetochore (Muller-Reichert et al., 2010; Redemann et al., 2017). We included the branched microtubules along the sMT lattice in this latter category. Microtubule dynamic instability is essential for the spindle functions, a feature largely used in cancer therapies (Steinmetz and Prota, 2018; Vicente and Wordeman, 2019; Wordeman and Vicente, 2021). Numerous MAPs ensure a precise regulation of microtubule dynamics along the various phases of mitosis and spindle zones (Lacroix et al., 2018; Roostalu et al., 2018; Srayko et al., 2005). These molecular motors and MAPs stochastically bind and unbind to the microtubules, making the spindle highly dynamic (Elting et al., 2018; Nazockdast and Redemann, 2020).

An important aspect of ensuring spindle function is the attachment of the chromosomes by the kMTs while preserving the dynamic instability of the microtubule plus-end (Bakhoum and Compton, 2012). Thus, a specialised structure assembles at the centromeric regions of the chromosomes: the kinetochore (Cheeseman, 2014; Musacchio and Desai, 2017). In particular, the outer kinetochore protein NDC-80 makes the bridge with the microtubules (Cheeseman et al., 2004; Suzuki et al., 2016; Ye et al., 2016). In the nematode, chromosomes are holocentric; thus, kinetochores are scattered all along the chromosomes (Maddox et al., 2004). Chromosome attachment happens at first laterally, along the lattice of the microtubules (side-on), before being converted into an attachment at their plus-end (end-on) (Cheerambathur et al., 2013; Magidson et al., 2011). This latter can withstand the tension from the microtubules on the kinetochore. This tension is essential for the spindle assembly checkpoint (SAC) to identify chromosome misattachment and also for the subsequent mechanisms to fix these mistakes (Kuhn and Dumont, 2019; McVey et al., 2021). When attachments are correct, they are secured by the SKA complex, and the kMT dynamicity is reduced (Cheerambathur et al., 2013; Cheerambathur et al., 2017; Hanisch et al., 2006; Schmidt et al., 2012). MAPs regulate the dynamics of the attached microtubule like the rescue factor CLS-2^{CLASP} (Al-Bassam and Chang, 2011; Cheeseman et al., 2005; Matthews et al., 1998). Some ubiquitous regulators also contribute, such as the polymerisation enhancer ZYG-9^{XMAP215} (Matthews et al., 1998). In the nematode, like in other organisms, only one-fifth of the kMTs can directly reach the spindle pole (Kiewisz et al., 2022; Petry, 2016; Redemann et al., 2017). The remaining kMTs are likely connected to sMTs since their minus ends are very close to the sMT lattices.

Capturing and attaching chromosomes are partly stochastic, making errors possible. These are usually intercepted by the SAC (Nicklas, 1997). Notably and similar to mammalian embryos, in the C. *elegans* zygote, the SAC is weak, and some of its components are dispensable (Gerhold et al., 2018; Oegema and Hyman, 2006; Pintard and Bowerman, 2019; Tarailo et al., 2007). The tension exerted by the microtubules on the kinetochore involves their polymerising (Bakhoum et al., 2009; Cimini et al., 2006; Ertych et al., 2014; Lampson and Grishchuk, 2017; Schwietert et al., 2022). Combined with minus-end dynamics, it leads to poleward flux, which is a displacement of the

microtubule network away from chromosomes (Barisic et al., 2021; Hotani and Horio, 1988; Mitchison, 1989; Steblyanko et al., 2020; Waters et al., 1996). A poleward flux was measured in various organisms, from Xenopus to mammals (Brust-Mascher et al., 2004; Brust-Mascher et al., 2009; Maddox et al., 2002; Maddox et al., 2003; Maiato et al., 2005; Matos et al., 2009; Yang et al., 2008). It plays an important role in correcting chromosome attachment errors, putatively by renewing the microtubules in contact with the kinetochore (Ertych et al., 2014; Ganem et al., 2005; Matos et al., 2009; Pereira and Maiato, 2012). During anaphase A, it also contributes to a synchronous migration of the sister chromatids toward their facing spindle poles (Ganem et al., 2005; Matos et al., 2009; Mitchison and Salmon, 1992). The most obvious flux mechanism is treadmilling, where the microtubule grows on the kinetochore side in a coordinated fashion with shortening at the spindle pole–this latter part powers the mechanism (Gaetz and Kapoor, 2004; Ganem et al., 2005). Such a mechanism was also recreated *in vitro* following a minimal system approach with only four proteins: XMAP215$^{\mathrm{ZYG\text{-}9}}$, the microtubule plus-end binding EB1$^{\mathrm{EBP\text{-}2}}$, CLASP2$^{\mathrm{CLS\text{-}2}}$, and the microtubule depolymerising kinesin MCAK$^{\mathrm{KLP\text{-}7}}$ (Arpag et al., 2020). This latter kinesin is essential to depolymerise the microtubules on the centrosome side. It can be assisted by severing enzymes like katanin, spastin and fidgetin (Zhang et al., 2007). Furthermore, treadmilling is consistent with the gel-like behaviour of the spindle reported in Xenopus egg extracts (Dalton et al., 2022) and the classic flux measurements using photobleaching/photoconversion experiments (Barisic et al., 2021; Mitchison and Salmon, 1992). Their results suggest a mostly solid motion of the spindle microtubule network at the steady state. Beyond the treadmilling mechanism, alternatives are twofold. Firstly, specialised molecular motors like kinesin-5 KIF11$^{\mathrm{BMK\text{-}1}}$ and kinesin-12 KIF15$^{\mathrm{KLP\text{-}18}}$ can slide the overlapping antiparallel microtubules carrying the kMTs through cross-linking (Miyamoto et al., 2004; Steblyanko et al., 2020; Uteng et al., 2008). Notably, kinesin-12 seems more specialised in parallel microtubules (Drechsler and McAinsh, 2016). Variants were proposed by sliding bridging fibres along each other (Jagric et al., 2021). The motion of these fibres is transmitted to other microtubules, including the kMTs and the sMTs, by cross-linking agents or motors like HSET$^{\mathrm{KLP\text{-}15/16/17}}$, NuMA$^{\mathrm{LIN\text{-}5}}$ or PRC1$^{\mathrm{SPD\text{-}1}}$ (Elting et al., 2014; Risteski et al., 2022; Steblyanko et al., 2020). Secondly, chromokinesin KIF4A$^{\mathrm{KLP\text{-}19}}$ congresses the chromosome arms and generates a reaction force that pushes the kMTs away from the chromosomes (Steblyanko et al., 2020; Wandke et al., 2012). It may play a similar role in nematode, although it would also regulate microtubule dynamics (Powers et al., 2004; Zimyanin et al., 2023). These various mechanisms can also be superimposed to ensure a robust poleward movement of the kMTs.

Surprisingly, although homologous proteins are present in the nematode C. e*legans,* no flux was reported in anaphase, while a putative but weak anti-poleward flux was suggested at metaphase (Labbe et al., 2004; Redemann et al., 2017). Notably, microtubules are highly dynamic in the nematode due to tubulin specificities (Chaaban et al., 2018). In contrast, a flux was found in zygote meiosis (Lantzsch et al., 2021). These paradoxes led us to investigate the flux during mitotic metaphase using the established fluorescence recovery after photobleaching (FRAP) (Axelrod et al., 1976; Giakoumakis et al., 2017; Matsuda and Nagai, 2014; White and Stelzer, 1999). Whereas we found no net flux for the spindle as a whole, we measured a poleward displacement of the microtubules close to the chromosomes. It depends on kinetochore proteins like CLS-2$^{\mathrm{CLASP}}$, NDC-80 and the kinetochore-attachment states. We finally suggest that the kMTs be slid along the fixed sMTs by KLP-18$^{\mathrm{KIF15}}$.

RESULTS

Two distinct microtubule-dynamics mechanisms account for recovering fluorescence after photobleaching.

We investigated the dynamics of the microtubules within the metaphasic spindle of the *C. elegans* zygote, studying recovery after photobleaching a 2.6μm wide region in the spindle (Methods § Microscopy). We used the GFP::TBB-2[β-tubulin] labelling. We previously reported no phenotype for this strain (Bouvrais et al., 2021). By analysing the average intensity in the photobleached region over time, we excluded that the diffusion of tubulin dimers accounted for the recovery inside the spindle (Supplemental Text §1 and Fig S1AB). Next, to investigate a putative flux, we set out to analyse the shape of the bleached area over time (Movie S1, Fig 1A, B). We used the centrosome on the bleaching side as a spatial reference to register images and get rid of the effects of spindle pole motion (Methods § Image processing, Fig S2). We produced the kymograph of individual embryos along the spindle axis by median projection along the transverse axis (Fig S3). The fast dynamics of the microtubules imposed imaging at 12.5 frames per second, leading to a low signal-to-noise ratio (SNR). To account for low SNR, we averaged the kymographs from several embryos after aligning them on the centrosome-side top corner of the bleached region in the kymograph and matching their intensity histograms. We then segmented the bleached region of the averaged kymograph. We fitted linearly its boundaries to quantify the displacement of the two fronts over time by their slope in the kymograph (Fig 1D, blue lines). We used a leave-one-out resampling (Jackknife) to obtain the standard errors of the slopes (Efron and Tibshirani, 1993; Quenouille, 1956). Such an approach also enabled us to visualise the front-slope distributions by displaying N-1 averages (Fig 1C, Methods § Statistics on kymograph front slopes). We finally computed the slope of the mid-curve between the edges corresponding to the bleached region solid displacement (Fig 1D, black line). It read 0.017 ± 0.028 μm/s, oriented poleward ($N = 10$, mean $\pm$ standard error (se) obtained by Jackknife resampling, $p = 0.56$ compared to 0). We concluded that the bleached area underwent no global displacement, meaning no significant global flux. This is consistent with previous reports on the lack of flux in metaphase and anaphase (Labbe et al., 2004; Redemann et al., 2017).

However, a visual inspection of the kymograph (Fig 1B, D) indicated a fluorescence recovery by closing the edges through two inward-oriented fronts. We investigated this V-shaped recovery and measured the closure velocities (Fig 1C). We read 0.117 ± 0.028 μm/s on the chromosome side, which significantly differed from 0 ($N = 10$, $p = 0.002$) and 0.082 ± 0.039 on the centrosome side ($p = 0.05$), both oriented inwards. These values of the slope were significantly different from each other ($p = 1.9 \times 10^{-5}$, $N = 10$). We reckoned that these two fronts of opposite directions might reveal distinct microtubule dynamics close and far from the chromosomes. Indeed, we expect the kMTs to be numerous enough to be detected close to chromosomes but vastly overwhelmed by sMTs at the location of the centrosome front (Redemann et al., 2017).

To safeguard against artefacts, we repeated the experiment using photoconversion (see Methods § Microscopy). We crossed strains with mEOS3.2::TBB-2[β-tubulin], labelling microtubules, and mCherry::TBG-1[γ-tubulin], revealing the centrosomes. We used this second label to register the images over time, as reported above (Methods § Image processing). We produced the kymographs, and again, to cope with the low SNR, we averaged them across embryos after aligning them on the photoconverted region. We obtained a similar V-shaped pattern (Fig S1C) as expected. We concluded that the two fronts were not artefactual.

After validating that the motion of the two closing edges depended on microtubule dynamics using *zyg-9*[XMAP215](RNAi), we investigated the front on the centrosome side (Suppl Text §2, Fig S1D, E, replica in Fig S1F). This depletion decreases the polymerisation rate at the plus ends. The most

obvious possibility of accounting for the observed fronts is a flux of the microtubules themselves. While a poleward flux of kMTs was not observed in the nematode, it is well-known in numerous other species. In contrast, an anti-poleward flux of sMTs appeared much more unlikely. The edge of the bleached region on the centrosome side was located at about half the distance between the chromosomes and the spindle pole; at this place, the sMTs are three times more numerous than the kMTs (Redemann et al., 2017). We thus focused on the sMTs and reckoned that the front displacement on the centrosome side might be accounted for by the dynamics of the sMTs (growth and shrinkage) combined with diffraction due to imaging. We modelled this phenomenon and observed a closing front similar to the experimental one (Suppl Text §3, Fig S4, S5). The reduction in the centrosome-side front velocity as the spindle got longer supported the proposed mechanism. Notably, this latter phenomenon is expected to cause two symmetric closing edges, i.e., on both the centrosome and chromosome sides, that move at the same speed. We concluded that dynamic instability is sufficient to account for the front motion on the centrosome side. However, because no correlation was visible between the chromosome side velocity and the spindle length (Fig S6), and because recovery was faster on that side, it suggested an additional mechanism.

Figure 1: Tubulin fluorescence recovery after photobleaching in the metaphasic spindle of *C. elegans*. (**A**) Schematics of the FRAP experiment to measure front displacements. Thick dark grey lines depict the spindle microtubules emanating from (red) anterior and (blue) posterior centrosomes; thin light grey ones, the one branching from other microtubule lattices. Black lines correspond to kinetochore microtubules bound to (blue bars) the condensed sister chromatids. Astral microtubules are depicted in purple colour. The dashed box corresponds to the bleached area. The blue bars represent the measurement location of the fronts on (dark blue) the chromosome and (light blue) the kinetochore sides. (**B**) Averaged kymograph over the posterior half-spindle of $N = 10$ GFP::TBB-2[β-tubulin] labelled embryos. The centrosome was located on the right-hand side. The orange line delineates the bleached region as obtained by our analysis (Movie S1, Methods § Image processing). (**C**) Front velocities by analysing the segmented bleached-region of $N = 10$ GFP::TBB-2[β-tubulin] non-treated embryos. Black dots represent the average of N-1 embryos, leaving out each embryo in turn (Methods § Statistics on kymograph front slopes). Bars correspond to means; error bars are estimated standard errors using Jackknife resampling. Light blue bars are velocities on the chromosome side, and dark blue the ones on the centrosome side. It corresponds to the kymograph analysis displayed in B. (**D**) Linear fits of the edges of the segmented kymograph averaged over the non-treated embryos reported in panels B and C. Grey dots depict the pixels in the bleached region. Boundaries are highlighted by coloured dots on (turquoise, left edge) the chromosome side and (blue, right edge) the centrosome side. Grey diamonds depict the region mid-line. Linear fits are reported as lines of corresponding colours.

A poleward flux of kinetochore microtubules may account for the chromosome-side recovery.

We reckoned that the higher front rate velocity observed on the chromosome side and the lack of correlation between the spindle length and the recovery front velocity suggested that a second mechanism might superimpose specifically on the chromosome side. The most obvious addition would involve the kMTs growing at their plus ends, i.e. at the chromosomes, while sliding towards the poles and thus along the spindle microtubules. Consistently, depletion of ZYG-9[XMAP215] reduced the front velocity on the chromosome side (Fig S1D, E). This hypothesis was consistent with Redemann and colleagues not measuring this poleward flux because being far from the chromosomes. Indeed, the kMTs are abundant compared to the sMTs, only close to the chromosomes (Redemann et al., 2017).

In other organisms, the poleward flux involves all the microtubules, although their dynamics may differ (Conway et al., 2022; Risteski et al., 2022; Zhai et al., 1995). Dalton and co-authors proposed that the spindle undergoes gelation to account for the solid poleward motion of the microtubules within spindles prepared from *Xenopus laevis* extracts (Dalton et al., 2022). In contrast, we proposed that only the kMTs would undergo a poleward flux. As their number decreases with distance from the kinetochore (Redemann et al., 2017), the flux could be seen as restricted close to the kinetochores. To challenge this idea, we performed FRAP experiments, bleaching a smaller area located either close or far from the spindle pole (Fig 2A). When looking far from the chromosomes, we obtained similar front velocities on edges facing chromosomes and centrosome (Fig 2B, maroon background). It suggested that, in this spindle region, both fronts reflected similar mechanisms. Moreover, this symmetry is incompatible with sMTs undergoing a flux in either direction. Because the sMTs are in the vast majority compared to the kMTs, the recovery was likely due solely to the microtubule dynamics combined with imaging diffraction, as detailed above. Close to the kinetochore, we measured higher front velocity on the chromosome side than the centrosome one, which is consistent with our previous measurement. However, this latter value was larger than in the bleached region far from the centrosomes. It may be related to a gradient of RAN protein around the chromosomes to promote microtubule growth (Bamba et al., 2002; Srayko et al., 2005). Overall, this experiment was consistent with our hypothesis that only the kMTs underwent poleward flux. It also accounted for the difference between our findings and previous measurements because the kMT flux could only be seen close to the chromosomes (Labbe et al., 2004; Redemann et al., 2017). Anywhere else, the non-fluxing sMTs, which are in the majority, hide the kMTs flux.

To further ascertain that fluxing kMTs could cause the front motion on the chromosomes side, we set out to genetically impair the kMTs' attachment to the kinetochore. We partially depleted NDC-80 in such a condition that chromosome segregation eventually succeeded in anaphase. We observed that the spindle in metaphase elongated during early metaphase (further termed early elongation) and plateaued before the anaphase onset, as previously reported (Fig S7A) (Cheerambathur et al., 2013). We studied the microtubule dynamics by FRAP during both phases (Fig 2C, D). During the early elongation, the front velocity on the chromosome side increased, while during the plateau, it decreased compared to the control. Such a result suggested that the kinetochore-microtubule attachment was involved in setting the corresponding front velocity. We repeated the experiment and obtained similar results (Fig S7C). Our partial depletion of NDC-80 led to a delay in microtubule attaching end-on to the kinetochores (Cheerambathur et al., 2013; Cheerambathur et al., 2017; Cheeseman et al., 2004; Lange et al., 2019). Thus, kinetochore-microtubule attachments could not bear a normal load, leading to early elongation. It might account for the observed increased front velocity under the hypothesis that the kMTs underwent a poleward flux. Later, during the plateau, we observed a velocity similar to or slightly reduced compared to

the control. Altogether, these observations were suggestive that growing kMTs at the kinetochores may be necessary for the mechanism causing the chromosome recovery front.

In the classic poleward-flux mechanism, CLS-2[CLASP] is critical to ensure that the microtubules switch to polymerisation while undergoing flux (Arpag et al., 2020; Cheeseman et al., 2005). We thus partially depleted this protein. We observed a shorter spindle during metaphase, as expected (Cheeseman et al., 2005) (Fig S7B, F). We monitored the front velocity using our FRAP assay and found an increased velocity on the chromosome side that recalled the one observed during the precocious spindle elongation upon *ndc-80(RNAi)* (Fig S8, S7BD). We interpreted it similarly: the kMTs were pulled out from the kinetochores upon *cls-2(RNAi)*, causing the increased front velocity. Consistently, upon strong depletion, the spindle breaks due to the cortical pulling forces (Cheerambathur et al., 2017; Cheeseman et al., 2005). Overall, we concluded that our results, especially the observed recovery front on the chromosome side, could be explained under the hypothesis that the kinetochore-microtubule fluxing towards the poles, requiring microtubule polymerisation at the plus-ends.

Figure 2: Front velocity on the chromosome side requires kMT polymerising at the plus-ends. (**A**) Schematics of the FRAP experiment, with bleaching localised close to either the chromosomes or the centrosome to measure front displacement during recovery. Grey lines depict the spindle microtubules emanating from (blue) the posterior centrosome. Black lines correspond to the kinetochore microtubules bound to (blue bars) the condensed sister chromatids. The grey and maroon boxes are the bleached areas close to chromosomes and the centrosome, respectively. The coloured lines depict the approximate positions of the front measurements corresponding to B. (**B**) Front velocities by segmenting the dark regions of non-treated embryos, bleached either (grey background, $N = 13$) close to chromosomes or (maroon background, $N = 14$) close to centrosomes. Microtubules were labelled using GFP::TBB-2[β-tubulin]. The colour of the bars corresponds to the schematics (A). (**C**) Averaged kymographs over the posterior half-spindle for the data presented in (D), with centrosome on the right-hand side. The orange line delineates the bleached region as obtained by our analysis (Methods § Image processing). (**D**) Front velocities by segmenting the dark region of $N = 6$ *ndc-80(RNAi)* embryos bleached during early elongation, $N = 8$ *ndc-80(RNAi)* bleached during spindle length plateau and their $N = 10$ controls (typical micrograph at Fig S7A). Microtubules were labelled using GFP::TBB-2[β-tubulin]. Light blue bars are values on the chromosome side and dark blue on the centrosome side. The experiment was replicated (Fig S7C). In panels B and D, black dots represent averages of N-1 embryos, leaving out, in turn, each embryo (Methods § Statistics on kymograph front slopes). Bars correspond to means, and error bars are estimated using Jackknife resampling.

The chromosome-side poleward flux depends on the attachment status at the kinetochore.

We reckoned that during spindle assembly, the kinetochores first attached laterally, with reduced resistance to tension on kinetochores, then end-on (Cheerambathur and Desai, 2014). This second configuration couples the dynamics of the kMT plus-end with the tension (Cheeseman et al., 2004; Suzuki et al., 2015). Next, the SKA complex is recruited during late metaphase and reduces the dynamicity of the attachments (Cheerambathur et al., 2017; Schmidt et al., 2012). We thus grouped

our non-treated and control experiments into 15s-long bins by the time the bleaching was performed and measured the front velocities. On the chromosome side, we found that it significantly decreased over time (Fig 3A). It suggested that the dynamics of the kMT plus-ends might impact the measured velocity of the chromosome-side front, which aligned with our hypothesis. We also excluded that the observed correlation could result from modest variations in the distance to the chromosomes (Fig S8B). Notably, the front of the centrosome side also displayed a decreasing velocity (Fig S8A). In the framework of the proposed model for this side (Suppl Text §3), we attributed that to either the decreasing microtubule growth rate over metaphase and anaphase, or the spindle getting slightly longer over the metaphase (Goshima and Scholey, 2010; Srayko et al., 2005). Interestingly, we observed decreased front velocities on both sides during early anaphase compared to metaphase (Fig S9A). It can be consistent with locking correct chromosome attachment at the kinetochores and the reduced microtubule growth rate at anaphase (Asbury, 2017; Srayko et al., 2005).

To exclude that a cell-scale change of microtubule dynamics over time accounted for the above results, we set out to manipulate the dynamics of the kinetochore microtubules genetically, although maintaining their attachments. We depleted the protein SKA-1 by RNAi, preventing the locking of the end-on attachments. Significantly and in contrast to mammalian cells, it did not delay anaphase onset (Schmidt et al., 2012). We measured the velocity of the fronts in the time range from 60 to 30 s before anaphase onset and observed a faster rate in depleted embryos(Fig 3B, replicated in Fig S9B). Indeed, the recruitment of this protein is expected at that moment(Cheerambathur et al., 2017). This was expected in our hypothesis of a chromosome-side front motion caused by kMT poleward flux.

We then performed the converse experiment, getting precocious and stronger recruitment of SKA complex, and thus a decrease of kinetochore-microtubule dynamics. To do so, we used the previously published mutated *ndc-80,* where four Aurora kinase phosphorylation sites were changed to alanines and termed NDC-80-4A (Cheerambathur et al., 2017). The authors made it RNAi-resistant by changing some triplets to synonymous codons. We also considered the strain, with similar scrambling of gene coding but no alteration of the phosphorylation sites as control, termed NDC-80-scr. It enabled us to target the endogenous copy by RNAi. We crossed these strains with the one carrying the GFP::TBB-2$^{\beta\text{-tubulin}}$ labelling and measured the flux in the time frame from 120 to 30 seconds before anaphase onset. We observed a decreased chromosome-side front velocity in NDC-80-4A conditions with respect to NDC-80-scr, both upon *ndc-80(RNAi)* or control RNAi. This was expected, considering that SKA decreased the kMT dynamics at the kinetochore too early or too strongly (Fig 3C). In contrast, on the centrosome side, upon depleting the endogenous NDC-80 protein and rescuing with the mutated protein NDC-80-4A, we observed an increase in front velocity, suggesting that the precocious recruitment of SKA affected differentially the front on each side (Fig S9C).

Altogether, the timely evolution of the velocity of the chromosome-side front and its dependence upon the SKA complex confirmed that the kinetochore microtubules undergoing a poleward flux likely contributed to the chromosome front motion.

Figure 3: (A) Correlation between the time of bleaching and the front velocities by segmenting the dark regions of non-treated or control embryos of experiments reported in other figures. Microtubules were labelled using GFP::TBB-2$^{\beta\text{-tubulin}}$. Black dots represent averages of N-1 embryos, leaving out, in turn, each embryo (Methods § Statistics on kymograph front slopes). Error bars are estimated using Jackknife resampling. Pearson coefficient read R = -0.82 (p = 0.0037). The grey-shaded region corresponds to the confidence interval at 0.95 on the correlation-line coefficients. **(B)** Front velocities by segmenting the dark region of N = 7 *ska-1(RNAi)* embryos bleached during late metaphase (-60 to -30 s from anaphase onset) and their N = 10 controls. The experiment was replicated (Fig S9B). **(C)** Front velocities on chromosome side by segmenting the dark region of NDC-80-scr embryos either (N = 12) treated by *ndc-80(RNAi)* or (N = 24) control and bleached; NDC-80-4A embryos either (N = 14) treated by *ndc-80(RNAi)* or (N = 11) control. Corresponding velocities on the centrosome side are in Fig S9C. In panels (B and C), microtubules were labelled using GFP::TBB-2$^{\beta\text{-tubulin}}$. Black dots represent averages of N-1 embryos, leaving out, in turn, each embryo (Methods § Statistics on kymograph front slopes). Bars correspond to means; error bars are estimated standard errors using Jackknife resampling. Light blue bars are velocity values measured on the chromosome side and dark blue on the centrosome side.

The tetrameric kinesin KLP-18 is likely sliding the kinetochore microtubules along the spindle microtubules.

Which mechanism could account for the kMT poleward flux? The most classic one is treadmilling, powered by the depolymerisation at the minus end. Electronic micrographs reported that about a quarter of the kMTs reached their corresponding pole (Redemann et al., 2017). Besides, half of the microtubules had opened minus-ends at the centrosomes (O'Toole et al., 2003). The treadmilling mechanism usually involves MCAK as the main microtubule depolymeriser (Arpag et al., 2020; Brust-Mascher et al., 2004). The MCAK homologous protein, KLP-7, is also localised at the kinetochores and spindle poles (Han et al., 2015; Sarov et al., 2012). We depleted it and measured the front velocity again (Fig S10). We did not observe a decreased front velocity on the chromosome side. The trend was instead an increase in velocity that we attributed to a role of KLP-7 at the kinetochore, putatively facilitating the turnover of the kMTs (Jaqaman et al., 2010; Kline-Smith et al., 2004; Wordeman et al., 2007). *In vitro*, a minimal treadmilling system comprises CLASP^{CLS-2}, XMAP215^{ZYG-9}, MCAK^{KLP-7} and EB1^{EBP-2}. While the former two proteins were involved in causing the front velocity that we observed on the chromosome side, the depletion of the latter two did not result in significant phenotype (Suppl Text §5, Fig S10 and S11). In conclusion, we considered treadmilling an unlikely mechanism. We rather suggested that the kMTs moved locally poleward in a mechanism reminiscent of the poleward flux despite these microtubules being too short to reach the centrosome.

As an alternative, we considered the sliding of the kMTs along the sMTs, which were not moving. KLP-18, a kinesin-12 motor, appeared as a plausible candidate. Its mammal counterpart, KIF15, was shown to slide parallel microtubules *in vitro* (Drechsler and McAinsh, 2016). Furthermore, KLP-18 is also essential to correctly align the microtubules in the acentrosomal meiotic spindle of the nematode (Cavin-Meza et al., 2022; Wolff et al., 2022). Importantly, KLP-18 is localised in the mitotic spindle (Suppl Video S2). We thus set out to deplete this protein by RNAi. Because it is required for successful meiosis, conditions were hypomorphic. We measured a modest, although significant, decrease in the chromosome-side front velocity, while no effect was observed on the centrosome side (Fig 4A). It suggested that KLP-18 could contribute to kMTs poleward fluxing.

To ascertain the role of this kinesin and get a stronger phenotype, we considered the temperature-sensitive mutant *klp-18(or447ts)* compared to OD868 denoted *klp-18(+)* (Connolly et al., 2014). After culturing the worms at 15°C, they were either imaged at 15°C (permissive condition) or incubated at 25°C for 30 min prior to dissection and imaging (restrictive). In this latter condition, we selected the embryo with no apparent pronuclei defect. We measured the velocity of the front on the chromosome side and observed a decrease by a factor of two in embryos carrying the mutated protein compared to control ones (Fig 4B). It is noteworthy that the microtubule growth-

rate increases with temperature, accounting for the observed increase in front velocity between permissive and restrictive temperatures (Chaaban et al., 2018). Notably, on the centrosome side at a given temperature, we observed no apparent difference between the control and mutated KLP-18 conditions (Fig S12). We concluded that KLP-18 specifically contributed to the front velocity on the chromosome side. We proposed that this tetrameric kinesin slid the kinetochore microtubules along the spindle ones, remaining immobile.

Figure 4: **(A)** Front velocities by segmenting the bleached region of $N = 9$ hypomorphic *klp-18(RNAi)* embryos compared to $N = 6$ non-treated embryos. **(B)** Front velocities on the chromosome side by segmenting the bleached region of *klp-18(or447ts)* embryos at ($N = 6$) permissive temperature 15ºC and ($N = 9$) restrictive temperature 25ºC; corresponding control embryos klp-18(+) at ($N = 7$) permissive temperature 15ºC and ($N = 9$) restrictive temperature 25ºC. Corresponding velocities on the centrosome side are in Fig S12. Microtubules were labelled using GFP::TBB-2[β-tubulin]. Black dots represent averages of N-1 embryos, leaving out, in turn, each embryo (Methods § Statistics on kymograph front slopes). Bars correspond to means; errors are estimated using Jackknife resampling. Light blue bars are values on the chromosome side and dark blue on the centrosome side.

DISCUSSION

We investigated the dynamics of the microtubules within the metaphase spindle in the nematode one-cell embryo. While we confirmed the lack of global poleward flux by FRAP and photoconversion experiments, we showed that diffusion of tubulin dimers from the cytoplasm could not account for the recovery. Instead, we observed two opposing fronts closing the bleached region. From imaging microtubule plus-ends, we confirmed the fast growth of the spindle microtubules (Fig S13). Modelling such behaviour shows that it causes a recovery of the bleached region uniform along the spindle. In contrast, the front on the chromosome side appears to reflect a distinct mechanism superimposed on the mechanism at work on the centrosome side. We attribute this addition to the kinetochore microtubules sliding along the spindle microtubules, the latter being fixed. Reasons are threefold: (i) While far from the chromosomes, one only sees the spindle microtubules, solely undergoing dynamic instability, and the velocities of both fronts are similar. In contrast, kinetochore microtubules are the majority over the other spindle microtubules only when looking close to kinetochores, leading to differences in the front velocities between both edges. It is plainly consistent with the lack of flux detection in previous works (Labbe et al., 2004; Redemann et al., 2017). (ii) A growth of the microtubules, while attached to the kinetochores, is needed to observe the chromosome side recovery, as found by partially depleting NDC-80, CLS-2[CLASP] (considering the plateau) or ZYG-9[XMAP215]. It is noteworthy that we also found accelerated flux upon *ndc-80(RNAi)*, *cls-2(RNAi)* and *ska-1(RNAi)*, as expected for poleward flux in other organisms (Matos et al., 2009). (iii) Furthermore, the chromosome-side velocity depends on the status of the attachment of the microtubule at the kinetochore. We probed it by looking over prometaphase and metaphase until early anaphase in unperturbed conditions. We also caused either a premature or a lack of securing of attachment by the SKA complex. In light of the flux measurement close to the chromosomes, our suggested sliding mechanism contrasts with the previous proposal by Redemann and colleagues. Indeed, we found no indication that the kMTs are detaching from the centrosomes and moving towards the kinetochore. However, we cannot exclude that the kinetochore microtubules depolymerise from their minus-ends, consistent with 38% of the ends opened in metaphase (Redemann et al., 2017).

Which mechanism could account for this sliding? A classic mechanism is treadmilling, requiring depolymerisation at the minus end. This mechanism usually involves MCAK as the main microtubule depolymeriser (Arpag et al., 2020; Brust-Mascher et al., 2004). In our case, because the depletion of the homolog of MCAK, KLP-7, does not decrease the recovery velocity, instead tending to increase it, we consider this mechanism unlikely. However, it is not impossible that other microtubule depolymerisers, like KLP-13, katanin, stathmin, and fidgetin, may contribute (Zhang et al., 2007). Notably, the depletion of KLP-13 has a reduced phenotype in the zygote (Kamath et al., 2003; Sonnichsen et al., 2005). Katanin was reported to be inactive during mitosis (Clandinin and Mains, 1993). Stathmin and fidgetin display a weak phenotype upon depletion in the one-cell embryo (Kamath et al., 2003; Lacroix et al., 2014; Lacroix et al., 2016; Sonnichsen et al., 2005). We also considered the transport of the kMTs by cross-linking to sMTs unlikely as these latter are not undergoing poleward flux. We can, therefore, exclude mechanisms involving bridging microtubules (Jagric et al., 2021) or sliding of the overlapping (Miyamoto et al., 2004; Steblyanko et al., 2020; Uteng et al., 2008). We rather propose that the sliding of kMTs over the sMTs is achieved by the kinesin-12 KLP-18. This motor contributes to focusing and organising the meiotic-spindle microtubules (Cavin-Meza et al., 2022; Wolff et al., 2022). Its mammal counterpart, KIF15, was shown to slide parallel microtubules *in vitro* (Drechsler and McAinsh, 2016). Some redundant sliding mechanisms may also exist, such as the transport of the kMT minus ends to the poles by dynein using LIN-5[NuMA] (Elting et al., 2014). In all cases, the sliding mechanisms must be coupled with the polymerisation of the microtubules at the kinetochore, like in other organisms, to maintain chromosome attachment (Barisic et al., 2021).

We wondered whether restricting the poleward flux to the kMTs is a peculiarity of nematodes. Metaphase is very brief in the nematode embryo, which prevents observing the spindle in a steady state. Furthermore, careful observation of the pole-to-pole distance during the metaphase shows a slow but constant length increase. This situation contrasts with the *Xenopus laevis* spindle prepared from extracts and at steady-state, where a gelification transition was observed (Dalton et al., 2022). A plausible cause of this difference is the strong pulling forces exerted on the spindle poles by the cortical force generators (Grill et al., 2001; Grill et al., 2003; Labbe et al., 2004). Compared to other cells, these forces are very high and participate in spindle elongation. One can speculate that sliding kMTs along the sMTs may mechanically insulate the kinetochores from the poles. In early metaphase, when pulling is weak, the sliding mechanism could generate enough force to ensure a correct function of the spindle assembly checkpoint and subsequent correcting of chromosome mis-attachments. Oppositely, in late metaphase, the strong pulling could be tempered down by letting the kMTs slide along the sMTs, leading to slow spindle elongation. Indeed, while the spindle slowly elongates in the non-treated embryos at that stage, decreasing cortical forces by depleting GPR-1 and GPR-2 reduces this phenotype (Cheeseman et al., 2005; Lewellyn et al., 2010). Overall, such a mechanism may maintain some tension at the kinetochores and be compatible with the SAC and correction mechanism function. Another important difference is the centromere organisation. Nematode chromosomes are holocentric; therefore, the kMTs do not assemble to form a K-fiber. However, a recent electron microscopy study suggests that enough space exists between microtubules in HeLa cells to accommodate molecular motors, making possible a mechanism similar to the one reported here (Kiewisz et al., 2022). Finally, it was recently reported in human cells that the dynamics of microtubules along the spindle axis are not uniform (Conway et al., 2022; Risteski et al., 2022). Furthermore, the minus-ends of the kMTs would also move towards the poles in a poleward flux. Similarly, half of the kMTs in HeLa cells do not reach the corresponding centrosomes (Kiewisz et al., 2022). This observation also suggests that a mechanism similar to the one proposed for the nematode may occur in mammalian cells.

MATERIAL AND METHODS

C. *elegans* strains

All *C. elegans* strains were maintained at 20°C and cultured using standard procedures (Brenner, 1974). *C. elegans* worms were grown on NGM plates seeded with OP50 *E. coli* strain. *C. elegans* worm strains and genotypes are listed in Table 1.

Strain	Genotype	Crossing	Origin and reference
AZ244	unc-119(ed3) III; ruIs57[pie-1::GFP::tbb-2 +unc-119(+)] V		CGC (Praitis et al., 2001)
EU3068	ebp-2(or1954[ebp-2::mKate2]) II; ruIs57[pie-1::GFP::tbb-2 +unc-119(+)] V		CGC (Sugioka et al., 2018)
JEP120	ruIs57[pie-1::GFP::tbb-2 +unc-119(+)] V; gpr-2(ok1179) III.	AZ244 X TH291	TH291 is a 10x backcross of RB1150 (C. elegans Deletion Mutant Consortium, 2012)
JEP92	unc-119(ed3) ? III; ddIs180[WRM062cF05 spd-2:: 2xTY1 GFP FRT 3xFlag;unc-119(+)] ; ltIs5 [pIC36; pie-1/GFP-TEV-STag::kbp-1; unc-119 (+)]	OD9 x TH231	OD9: CGC (Cheeseman et al., 2004)
			TH231: (Decker et al., 2011)
JEP97	unc-119(ed3) ? III; ltIs37 [pAA64; pie-1/mCHERRY::his-58; unc-119 (+)]IV;ddIs44[WRM0614cB02 GLCherry::tbg-1;unc-119(+)]	OD56 x TH169	OD56: CGC (McNally et al., 2006)
			TH169: (Woodruff et al., 2015)

JEP166	ddIs44[WRM0614cB02 GLCherry::tbg-1;unc-119(+)]; pJD734_pSW077_Mos1_Pmex-5_mEOS3.2-tbb2	JEP97 x JDU562	JDU562: (Dias Maia Henriques, 2018)
JEP172	ebp-2(gk756) II; ruIs57[pie-1::GFP::tbb-2 +unc-119(+)] V	AZ244 x VC1614	VC1614: CGC (C. elegans Deletion Mutant Consortium, 2012)
SMW36	klp-18(or447ts), dpy-20 IV; ltSi220[pOD1249/pSW077; Pmex-5::GFP::tbb-2::operon_linker::mCherry::his-11; cb-unc-119(+)] I		(Wolff et al., 2022)
OD868	ltSi220[pOD1249/pSW077; Pmex-5::GFP-tbb-2-operon-linker-mCherry-his-11; cb-unc119(+)] I.		(Hollis et al., 2020)
JEP176	ltSi120[[pDC170;Pndc-80:NDC-80 reencoded; cb-unc-119(+)]II; ruIs57[pie-1::GFP::tbb-2 +unc-119(+)] V	AZ244 x OD611	
OD611	unc-119(ed3) III; ltSi120[[pDC170;Pndc-80:ndc-80 reencoded; cb-unc-119(+)]]II #3		(Cheerambathur et al., 2017)
JEP177	ltSi122[pDC178;Pndc-80:NDC-80 (8,18,44,51AAAA) reencoded; cb-unc-119(+)]II; ruIs57[pie-1::GFP::tbb-2 +unc-119(+)] V	AZ244 x OD634	
OD634	unc-119(ed3)III; ltSi122[pDC178;Pndc-80:NDC-80 (8,18,44,51AAAA) reencoded; cb-unc-119(+)]II #1		(Cheerambathur et al., 2013)
JEP186	ebp-2(or1954[ebp-2::mKate2]) II unc-119(ed3)III; ltSi462[pNH95; Pmex-5::klp-18::GFP::tbb-2 3'UTR; cb- unc-119(+)]I	EU3068 x OD1495	EU3068: CGC (Sugioka et al., 2018) OD1495: (Hattersley et al., 2016)

Table 1: Worm strains used in this article and their genotypes.

RNAi treatment

RNAi was performed by feeding as described previously (Kamath and Ahringer, 2003; Timmons and Fire, 1998). The control embryos for the RNAi experiments were fed with bacteria carrying the empty plasmid L4440. We targeted *ndc-80* by RNAi using the same sequence as in (Cheerambathur et al., 2017) after cloning it into a vector. All other clones came from the Ahringer-Source BioScience library (Kamath et al., 2003). RNAi target, clone ID or sequence used, and feeding conditions are listed in Table 2. We set the RNAi treatment duration so that we did not notice any phenotype suggesting impaired meiosis.

Target	RNAi clone ID (Ahringer-Source Bioscience) or sequence used	IPTG concentration (mM)	Feeding time (hours)	Incubation temperature (°C)
cls-2	III-4J10	4	48	20
klp-7	III-5B24	4	48	25
zyg-9	II-6M11	4	8	25

ndc-80	Oligo 1: 5' – GGGGACAAGTTTGTACAAAAAAGCAGGCTTGGA TGACAAGTACATTCAGAGATTATACAAATGATC – 3'	4	96	20 (Fig 2, S7)
	Oligo 2: 5' – GGGGACCACTTTGTACAAGAAAGCTGGGTGGTG GTTCAAGATTCATTTGAATATTAAGTCCACTG – 3'			18 (Fig 3, S9)
ska-1	I-5A22	4	96	18
klp-18	IV3O14	4	40	18
ani-2	II-4P19	2	48	25
C27D9.1	II-4A13	4	48	25

Table 2: RNAi target, clone ID and treatment conditions used in this article.

Microscopy: acquisition conditions

Preparation of C. elegans samples for imaging

C. elegans hermaphrodite adults were dissected in M9 medium (50 mM Na2HPO4, 17 mM KH2POone, 86 mM NaCl, 1 mM MgSO4) (Brenner, 1974). The released embryos were deposited on an agarose pad (2% w/v agarose, 0.6% w/v NaCl, 4% w/v sucrose) between slide and coverslip or between two 24x60 cm coverslips. To confirm the absence of phototoxicity and photodamage, we checked for normal rates of subsequent divisions (Riddle, 1997; Tinevez et al., 2012).

Imaging condition for photobleaching and photoconversion experiments

Embryo imaging to perform FRAP was achieved on a scanning confocal microscope with deconvolution (LSM 880 airyscan, Zeiss) using an apochromatic 63x/NA1.4 oil objective. Acquisition parameters were controlled by Zen black software. We imaged at 18°C, except otherwise stated, using the CherryTemp temperature control system in all cases (CherryBiotech, Rennes, France). With all strains expressing GFP::TBB-2$^{\beta\text{-tubulin}}$, we performed photobleaching using an argon laser at wavelength 488 nm and 70 µW power for 75 iterations (photobleaching time ~ 1.9 s). We then imaged using the same laser but at a 2.30 µW power. Five images were taken before photobleaching to get the basal fluorescence of the embryo. The laser power was measured at the objective output just before each microscopy session. We set to bleach a 2.6 x 19.5 µm area within the mitotic spindles (the embryo was oriented horizontally) (Fig1A). When accommodating two bleached regions per half-spindle (Fig 2AB), their width read 1.3 µm. Images were acquired at 12.5 Hz.

In the case of photoconversion experiments, we used the strain JEP166 expressing mCherry::TBG-1$^{\gamma\text{-tubulin}}$ and mEOS3.2::TBB-2$^{\beta\text{-tubulin}}$. The mEOS3.2::TBB-2$^{\beta\text{-tubulin}}$ labelling was obtained from the JDU562 strain, a kind gift from Julien Dumont's team. The mCherry::TBG-1$^{\gamma\text{-tubulin}}$ channel enables the images' registration on the centrosomes' position (see § Image processing). Five images were taken before photoconversion to have the basal fluorescence of the embryo. Photoconversion was performed using a 405 nm UV laser at 10 µW for 35 iterations (photoconversion time ~ 0.850 s). We photoconverted a 13 x 1.3 µm area within the mitotic spindle after vertically orienting the anteroposterior axis of the embryo. Images were acquired with a 561 nm laser at 9 µW power and 3 frames per second.

To study the localisation of KLP-18, we imaged *C. elegans* one-cell embryos at the spindle plane, viewing from the nuclear envelope breakdown (NEBD) until late anaphase. We used a Leica DMi8 spinning disk microscope with Adaptive Focus Control (AFC) and an HCX Plan Apo 100x/1.4 NA oil objective. Illumination was performed using a laser with an emission wavelength of 488 nm, and we used a GFP/FITC 4 nm bandpass excitation filter and a Quad Dichroic emission filter. Images were acquired using an ultra-sensitive Roper Evolve EMCCD camera that was controlled by the acquisition device by Inscoper, SAS. During the experiments, the embryos were kept at about 20°C.

In all experiments, the anaphase onset was taken as a time reference. Images were stored using OMERO software (Li et al., 2016).

Image Processing: Automatic recognition of the bleached region edges and measuring the slopes of the fluorescence-recovery fronts

Assembling a homogenous set of embryos

Observing the kymograph (Fig 1B, S3) suggested that the recovery proceeded by two fronts moving inwards. We set out to measure the velocity of these fronts. We measured the laser power at each session to safeguard against intensity variation between sessions. It also enabled us to select a set of embryos bleached in similar conditions. Furthermore, the spindle lies 10-15 μm deep in the sample, further introducing variability. Overall, we considered only the experiments where the FRAP efficacy reached at least 30%, assessed by comparing the fluorescence intensities before and after the FRAP event. We checked that we performed bleaching during metaphase, i.e. between -120 and -30 s before the anaphase onset; it ensured a full fluorescence recovery prior to anaphase onset. For the *ska-1(RNAi)* experiments, this range was reduced to [-60 -30] s to correspond to the time period when SKA-1 is recruited in non-treated embryos (Cheerambathur et al., 2017). Finally, the distance between the metaphasic plate and the bleached region was measured to guarantee that a bright corridor separated the bleached region from the hollow due to the chromosomes, ensuring a robust measure of the chromosomal front speed (Fig 1B, S3).

Obtaining a kymograph for each embryo

We performed a rigid registration of the raw images using the slightly modified version of the so-called plugin (publicly available v. 1.4) under Icy (de Chaumont et al., 2012). This registration focused on the centrosome position on the side of the bleached half-spindle, preventing spindle displacement or rotation from contributing to the measured front motion (Fig 1, S1, S2A). We then applied a median filter under Fiji to limit the noise (Schindelin et al., 2012). We computed the kymograph by considering a region of interest of a length of 23.4 μm and a width of 3.12 μm (Fig S2B). To do so, we used Fiji and performed a median projection along a direction transverse to the spindle axis (Fig S2C).

Averaging kymographs over several embryos

To ensure that each embryo contributed equally, we performed a histogram equalisation among the individual embryos' kymographs using the histogramMatcher class in sciJava (interfaced using Beanshell) (Rueden et al., 2021). Finally, we registered the kymographs aligning on the centrosome-side edge of the bleached region at the first time after bleach, using a supervised home-designed Matlab plugin (Fig S2C, arrowhead). We then averaged the kymographs pixel-wise (Fig S2D). We then cropped the embryo-averaged kymograph to consider only the 40 first seconds and the bleached half-spindle.

Kymograph segmentation

Next, we segmented the bleached area in the kymograph. As a preprocessing, we convolved the kymograph with a 3x3 kernel reading 5 in the central element and 1 elsewhere (Gaussian blur). We

then autonomously detected the non-bleached area being conservative and considering only the pixels evidently non-bleached. To do so, we set the background class as the 60% brightest pixels. We then performed grey morphology (to remove micro/dot regions due to noise). Practically, we used two iterations of opening with a 1-pixel (130 nm) circular structural element and two closing iterations with the same element (Fig S2E, black line). Similarly, we set the bleached area as the 25% darkest pixels and performed the grey morphology (Fig S2E, orange line). We used neighbourhood information to decide about the pixels with intermediate grey levels. We trained a machine learning algorithm, namely random forests featuring 100 trees (Breiman, 2001), using the above classes for training and all kymographs in a set acquired under the same condition. We then applied the algorithm to classify the pixels within the kymograph (Fig S2E, red line) using Weka under Fiji (Beanshell script) (Arganda-Carreras et al., 2017). Finally, we performed grey morphology closing (2 pixels circle), watershed and the same closing again. It enabled us to remove artefactual dotty regions while preserving the largest one corresponding to the bleached area. We then selected the largest segmented object. Indeed, because the recovery was never complete (Fig 1B) and the contrast was low, a small artefactual region may appear in the tail of the main one. Finally, we dilated the shape using a 1-pixel-radius circle twice, making the resulting shape convex.

Kymograph boundary fitting

Having segmented the bleached area, we fitted a line along the boundaries (between 2 and 30 s) to get the mean front velocities. We also computed the mid-curve (average between the positions of the two edges at each time point) and fitted it with a line.

Statistics on kymograph front slopes

We used a resampling method to get the error bars of the front velocities since we could not perform the computation on individual embryos because the images were too noisy. We used the Jackknife method, which requires computing the slope over a set containing all embryos but one (Efron and Tibshirani, 1993; Quenouille, 1956). We thus obtained a standard error on the averaged value estimated considering all embryos and could perform t-tests against the null hypothesis that the slope is 0.

Statistics representation

To compare conditions, we used the Jackknife again (Arvesen, 1969; Schechtman and Wang, 2004) or Pearson correlation when relevant. For the sake of simplicity, we depicted confidence levels using diamonds or stars (***, $P \leq 0.0005$; **, $P \leq 0.005$; *, $P \leq 0.05$; ◇ $P \leq 0.1$; n.s., $P > 0.1$). This latter means non-significant.

Acknowledgements

The strains TH291, TH231, and TH169 are kind gifts from Prof A. A. Hyman. Dr J. Dumont kindly offered JDU562. We thank Dr Gregoire Michaux for the feeding clone library and technical support. We are also thankful to Dr Arshad Desai for the gift of strains OD868, OD611 and OD634. The strain SMW36 was a kind gift of Prof Sarah M. Wignall. We also thank Drs. Gilliane Maton, Fabrice Mahé, Anne Corlu, Christophe Heligon, Rebecca Smith, Sebastien Huet, Flora Demouchy, Gregoire Michaux, Anne Pacquelet, Stéphanie Dutertre, Xavier Pinson, Marc Tramier, Ostiane d'Augustin for discussions about the project. Some strains were provided by the Caenorhabditis Genetics Center (CGC), funded by the National Institutes of Health Office of Research Infrastructure Programs (P40 OD010440; University of Minnesota). We also acknowledge La Ligue contre le cancer (comites d'Ille-et-Vilaine et du Maine-et-Loire). Microscopy imaging was performed at the Microscopy Rennes Imaging Center (MRIC), UMS 3480 CNRS/US

18 INSERM/University of Rennes 1. MRIC is a member of the national infrastructure France-BioImaging, supported by the French National Research Agency (ANR-10-INBS-04). In particular, we acknowledge the funding of the Zeiss Airyscan confocal microscope by EU funding FEDER under reference CARE Phase 2.

Author contribution

Conceptualisation: JP, LLM; Data curation: NS, MDS, CT, LC, JP; Formal analysis: NS, MDS, CT, LC, JP; Funding acquisition: JP, HB, LLM; Investigation: NS, MDS, CT, PF, LC, SP, JP; Methodology: NS, MDS, HB, LC, JP; Project administration: JP; Software: NS, JP, HB; Supervision: LLM, JP, LC; Validation: HB, LC, JP; Visualisation: NS, JP; Writing NS, JP; Writing – review and editing: LC, NS, MDS, JP, LLM, HB.

Conflict of interest

The authors declare that they have no conflict of interest.

References

Akhmanova, A., and Steinmetz, M.O. (2015). Control of microtubule organization and dynamics: two ends in the limelight. Nat Rev Mol Cell Biol *16*, 711-726.

Al-Bassam, J., and Chang, F. (2011). Regulation of microtubule dynamics by TOG-domain proteins XMAP215/Dis1 and CLASP. Trends Cell Biol *21*, 604-614.

Arganda-Carreras, I., Kaynig, V., Rueden, C., Eliceiri, K.W., Schindelin, J., Cardona, A., and Sebastian Seung, H. (2017). Trainable Weka Segmentation: a machine learning tool for microscopy pixel classification. Bioinformatics *33*, 2424-2426.

Arpag, G., Lawrence, E.J., Farmer, V.J., Hall, S.L., and Zanic, M. (2020). Collective effects of XMAP215, EB1, CLASP2, and MCAK lead to robust microtubule treadmilling. Proc Natl Acad Sci U S A *117*, 12847-12855.

Arvesen, J.N. (1969). Jackknifing U-Statistics. The Annals of Mathematical Statistics *40*, 2076-2100.

Asbury, C.L. (2017). Anaphase A: Disassembling Microtubules Move Chromosomes toward Spindle Poles. Biology (Basel) *6*.

Axelrod, D., Koppel, D.E., Schlessinger, J., Elson, E., and Webb, W.W. (1976). Mobility measurement by analysis of fluorescence photobleaching recovery kinetics. Biophys J *16*, 1055-1069.

Bakhoum, S.F., and Compton, D.A. (2012). Kinetochores and disease: keeping microtubule dynamics in check! Curr Opin Cell Biol *24*, 64-70.

Bakhoum, S.F., Thompson, S.L., Manning, A.L., and Compton, D.A. (2009). Genome stability is ensured by temporal control of kinetochore-microtubule dynamics. Nat Cell Biol *11*, 27-35.

Bamba, C., Bobinnec, Y., Fukuda, M., and Nishida, E. (2002). The GTPase Ran regulates chromosome positioning and nuclear envelope assembly in vivo. Curr Biol *12*, 503-507.

Barisic, M., Rajendraprasad, G., and Steblyanko, Y. (2021). The metaphase spindle at steady state - Mechanism and functions of microtubule poleward flux. Seminars in cell & developmental biology *117*, 99-117.

Bouvrais, H., Chesneau, L., Le Cunff, Y., Fairbrass, D., Soler, N., Pastezeur, S., Pecot, T., Kervrann, C., and Pecreaux, J. (2021). The coordination of spindle-positioning forces during the asymmetric division of the Caenorhabditis elegans zygote. EMBO reports *22*, e50770.

Breiman, L. (2001). Random Forests. Machine Learning *45*, 5-32.

Brenner, S. (1974). The genetics of Caenorhabditis elegans. Genetics *77*, 71-94.

Brust-Mascher, I., Civelekoglu-Scholey, G., Kwon, M., Mogilner, A., and Scholey, J.M. (2004). Model for anaphase B: role of three mitotic motors in a switch from poleward flux to spindle elongation. Proc Natl Acad Sci U S A *101*, 15938-15943.

Brust-Mascher, I., Sommi, P., Cheerambathur, D.K., and Scholey, J.M. (2009). Kinesin-5-dependent poleward flux and spindle length control in Drosophila embryo mitosis. Mol Biol Cell *20*, 1749-1762.

C. elegans Deletion Mutant Consortium (2012). Large-scale screening for targeted knockouts in the Caenorhabditis elegans genome. G3 *2*, 1415-1425.

Cavin-Meza, G., Kwan, M.M., and Wignall, S.M. (2022). Multiple motors cooperate to establish and maintain acentrosomal spindle bipolarity in C. elegans oocyte meiosis. eLife *11*.

Chaaban, S., Jariwala, S., Hsu, C.T., Redemann, S., Kollman, J.M., Muller-Reichert, T., Sept, D., Bui, K.H., and Brouhard, G.J. (2018). The Structure and Dynamics of C. elegans Tubulin Reveals the Mechanistic Basis of Microtubule Growth. Dev Cell *47*, 191-204 e198.

Cheerambathur, D.K., and Desai, A. (2014). Linked in: formation and regulation of microtubule attachments during chromosome segregation. Curr Opin Cell Biol *26C*, 113-122.

Cheerambathur, D.K., Gassmann, R., Cook, B., Oegema, K., and Desai, A. (2013). Crosstalk between microtubule attachment complexes ensures accurate chromosome segregation. Science *342*, 1239-1242.

Cheerambathur, D.K., Prevo, B., Hattersley, N., Lewellyn, L., Corbett, K.D., Oegema, K., and Desai, A. (2017). Dephosphorylation of the Ndc80 Tail Stabilizes Kinetochore-Microtubule Attachments via the Ska Complex. Dev Cell *41*, 424-437 e424.

Cheeseman, I.M. (2014). The kinetochore. Cold Spring Harb Perspect Biol *6*, a015826.

Cheeseman, I.M., MacLeod, I., Yates, J.R., 3rd, Oegema, K., and Desai, A. (2005). The CENP-F-like proteins HCP-1 and HCP-2 target CLASP to kinetochores to mediate chromosome segregation. Curr Biol *15*, 771-777.

Cheeseman, I.M., Niessen, S., Anderson, S., Hyndman, F., Yates, J.R., 3rd, Oegema, K., and Desai, A. (2004). A conserved protein network controls assembly of the outer kinetochore and its ability to sustain tension. Genes Dev *18*, 2255-2268.

Cimini, D., Wan, X., Hirel, C.B., and Salmon, E.D. (2006). Aurora kinase promotes turnover of kinetochore microtubules to reduce chromosome segregation errors. Curr Biol *16*, 1711-1718.

Clandinin, T.R., and Mains, P.E. (1993). Genetic studies of mei-1 gene activity during the transition from meiosis to mitosis in Caenorhabditis elegans. Genetics *134*, 199-210.

Connolly, A.A., Osterberg, V., Christensen, S., Price, M., Lu, C., Chicas-Cruz, K., Lockery, S., Mains, P.E., and Bowerman, B. (2014). Caenorhabditis elegans oocyte meiotic spindle pole assembly requires microtubule severing and the calponin homology domain protein ASPM-1. Mol Biol Cell *25*, 1298-1311.

Conway, W., Kiewisz, R., Fabig, G., Kelleher, C.P., Wu, H.Y., Anjur-Dietrich, M., Muller-Reichert, T., and Needleman, D.J. (2022). Self-organization of kinetochore-fibers in human mitotic spindles. eLife *11*.

Dalton, B.A., Oriola, D., Decker, F., Jülicher, F., and Brugués, J. (2022). A gelation transition enables the self-organization of bipolar metaphase spindles. Nature Physics *18*, 323-331.

de Chaumont, F., Dallongeville, S., Chenouard, N., Herve, N., Pop, S., Provoost, T., Meas-Yedid, V., Pankajakshan, P., Lecomte, T., Le Montagner, Y.*, et al.* (2012). Icy: an open bioimage informatics platform for extended reproducible research. Nat Methods *9*, 690-696.

Decker, M., Jaensch, S., Pozniakovsky, A., Zinke, A., O'Connell, K.F., Zachariae, W., Myers, E., and Hyman, A.A. (2011). Limiting amounts of centrosome material set centrosome size in C. elegans embryos. Curr Biol *21*, 1259-1267.

Dias Maia Henriques, A.M. (2018). Study of the Mechanisms of Chromosome Segregation in the C. elegans Embryo.

Drechsler, H., and McAinsh, A.D. (2016). Kinesin-12 motors cooperate to suppress microtubule catastrophes and drive the formation of parallel microtubule bundles. Proc Natl Acad Sci U S A *113*, E1635-1644.

Efron, B., and Tibshirani, R.J. (1993). An introduction to the bootstrap (New York ; London: Chapman & Hall).

Elting, M.W., Hueschen, C.L., Udy, D.B., and Dumont, S. (2014). Force on spindle microtubule minus ends moves chromosomes. J Cell Biol *206*, 245-256.

Elting, M.W., Suresh, P., and Dumont, S. (2018). The Spindle: Integrating Architecture and Mechanics across Scales. Trends Cell Biol *28*, 896-910.

Ertych, N., Stolz, A., Stenzinger, A., Weichert, W., Kaulfuss, S., Burfeind, P., Aigner, A., Wordeman, L., and Bastians, H. (2014). Increased microtubule assembly rates influence chromosomal instability in colorectal cancer cells. Nat Cell Biol *16*, 779-791.

Gaetz, J., and Kapoor, T.M. (2004). Dynein/dynactin regulate metaphase spindle length by targeting depolymerizing activities to spindle poles. J Cell Biol *166*, 465-471.

Ganem, N.J., Upton, K., and Compton, D.A. (2005). Efficient mitosis in human cells lacking poleward microtubule flux. Curr Biol *15*, 1827-1832.

Gerhold, A.R., Poupart, V., Labbe, J.C., and Maddox, P.S. (2018). Spindle assembly checkpoint strength is linked to cell fate in the Caenorhabditis elegans embryo. Mol Biol Cell *29*, 1435-1448.

Giakoumakis, N.N., Rapsomaniki, M.A., and Lygerou, Z. (2017). Analysis of Protein Kinetics Using Fluorescence Recovery After Photobleaching (FRAP). Methods Mol Biol *1563*, 243-267.

Goshima, G., and Scholey, J.M. (2010). Control of mitotic spindle length. Annu Rev Cell Dev Biol *26*, 21-57.

Grill, S.W., Gonczy, P., Stelzer, E.H., and Hyman, A.A. (2001). Polarity controls forces governing asymmetric spindle positioning in the Caenorhabditis elegans embryo. Nature *409*, 630-633.

Grill, S.W., Howard, J., Schaffer, E., Stelzer, E.H., and Hyman, A.A. (2003). The distribution of active force generators controls mitotic spindle position. Science *301*, 518-521.

Han, X., Adames, K., Sykes, E.M., and Srayko, M. (2015). The KLP-7 Residue S546 Is a Putative Aurora Kinase Site Required for Microtubule Regulation at the Centrosome in C. elegans. PLoS One *10*, e0132593.

Hanisch, A., Sillje, H.H., and Nigg, E.A. (2006). Timely anaphase onset requires a novel spindle and kinetochore complex comprising Ska1 and Ska2. EMBO J *25*, 5504-5515.

Hattersley, N., Cheerambathur, D., Moyle, M., Stefanutti, M., Richardson, A., Lee, K.Y., Dumont, J., Oegema, K., and Desai, A. (2016). A Nucleoporin Docks Protein Phosphatase 1 to Direct Meiotic Chromosome Segregation and Nuclear Assembly. Dev Cell *38*, 463-477.

Hollis, J.A., Glover, M.L., Schlientz, A.J., Cahoon, C.K., Bowerman, B., Wignall, S.M., and Libuda, D.E. (2020). Excess crossovers impede faithful meiotic chromosome segregation in C. elegans. PLoS Genet *16*, e1009001.

Hotani, H., and Horio, T. (1988). Dynamics of microtubules visualized by darkfield microscopy: treadmilling and dynamic instability. Cell Motil Cytoskeleton *10*, 229-236.

Jagric, M., Risteski, P., Martincic, J., Milas, A., and Tolic, I.M. (2021). Optogenetic control of PRC1 reveals its role in chromosome alignment on the spindle by overlap length-dependent forces. eLife *10*.

Jaqaman, K., King, E.M., Amaro, A.C., Winter, J.R., Dorn, J.F., Elliott, H.L., McHedlishvili, N., McClelland, S.E., Porter, I.M., Posch, M.*, et al.* (2010). Kinetochore alignment within the metaphase plate is regulated by centromere stiffness and microtubule depolymerases. J Cell Biol *188*, 665-679.

Kamath, R.S., and Ahringer, J. (2003). Genome-wide RNAi screening in Caenorhabditis elegans. Methods *30*, 313-321.

Kamath, R.S., Fraser, A.G., Dong, Y., Poulin, G., Durbin, R., Gotta, M., Kanapin, A., Le Bot, N., Moreno, S., Sohrmann, M.*, et al.* (2003). Systematic functional analysis of the Caenorhabditis elegans genome using RNAi. Nature *421*, 231-237.

Kiewisz, R., Fabig, G., Conway, W., Baum, D., Needleman, D., and Muller-Reichert, T. (2022). Three-dimensional structure of kinetochore-fibers in human mitotic spindles. eLife *11*.

Kline-Smith, S.L., Khodjakov, A., Hergert, P., and Walczak, C.E. (2004). Depletion of centromeric MCAK leads to chromosome congression and segregation defects due to improper kinetochore attachments. Mol Biol Cell *15*, 1146-1159.

Kuhn, J., and Dumont, S. (2019). Mammalian kinetochores count attached microtubules in a sensitive and switch-like manner. J Cell Biol *218*, 3583-3596.

Labbe, J.C., McCarthy, E.K., and Goldstein, B. (2004). The forces that position a mitotic spindle asymmetrically are tethered until after the time of spindle assembly. J Cell Biol *167*, 245-256.

Lacroix, B., Bourdages, K.G., Dorn, J.F., Ihara, S., Sherwood, D.R., Maddox, P.S., and Maddox, A.S. (2014). In situ imaging in C. elegans reveals developmental regulation of microtubule dynamics. Dev Cell *29*, 203-216.

Lacroix, B., Letort, G., Pitayu, L., Salle, J., Stefanutti, M., Maton, G., Ladouceur, A.M., Canman, J.C., Maddox, P.S., Maddox, A.S.*, et al.* (2018). Microtubule Dynamics Scale with Cell Size to Set Spindle Length and Assembly Timing. Dev Cell *45*, 496-511 e496.

Lacroix, B., Ryan, J., Dumont, J., Maddox, P.S., and Maddox, A.S. (2016). Identification of microtubule growth deceleration and its regulation by conserved and novel proteins. Mol Biol Cell *27*, 1479-1487.

Lampson, M.A., and Grishchuk, E.L. (2017). Mechanisms to Avoid and Correct Erroneous Kinetochore-Microtubule Attachments. Biology (Basel) *6*.

Lange, K.I., Suleman, A., and Srayko, M. (2019). Kinetochore Recruitment of the Spindle and Kinetochore-Associated (Ska) Complex Is Regulated by Centrosomal PP2A in Caenorhabditis elegans. Genetics *212*, 509-522.

Lantzsch, I., Yu, C.H., Chen, Y.Z., Zimyanin, V., Yazdkhasti, H., Lindow, N., Szentgyoergyi, E., Pani, A.M., Prohaska, S., Srayko, M.*, et al.* (2021). Microtubule reorganization during female meiosis in C. elegans. eLife *10*.

Lewellyn, L., Dumont, J., Desai, A., and Oegema, K. (2010). Analyzing the effects of delaying aster separation on furrow formation during cytokinesis in the Caenorhabditis elegans embryo. Mol Biol Cell *21*, 50-62.

Li, S., Besson, S., Blackburn, C., Carroll, M., Ferguson, R.K., Flynn, H., Gillen, K., Leigh, R., Lindner, D., Linkert, M.*, et al.* (2016). Metadata management for high content screening in OMERO. Methods *96*, 27-32.

Maddox, P., Desai, A., Oegema, K., Mitchison, T.J., and Salmon, E.D. (2002). Poleward microtubule flux is a major component of spindle dynamics and anaphase a in mitotic Drosophila embryos. Curr Biol *12*, 1670-1674.

Maddox, P., Straight, A., Coughlin, P., Mitchison, T.J., and Salmon, E.D. (2003). Direct observation of microtubule dynamics at kinetochores in Xenopus extract spindles: implications for spindle mechanics. J Cell Biol *162*, 377-382.

Maddox, P.S., Oegema, K., Desai, A., and Cheeseman, I.M. (2004). "Holo"er than thou: chromosome segregation and kinetochore function in C. elegans. Chromosome Res *12*, 641-653.

Magidson, V., O'Connell, C.B., Loncarek, J., Paul, R., Mogilner, A., and Khodjakov, A. (2011). The spatial arrangement of chromosomes during prometaphase facilitates spindle assembly. Cell *146*, 555-567.

Maiato, H., Khodjakov, A., and Rieder, C.L. (2005). Drosophila CLASP is required for the incorporation of microtubule subunits into fluxing kinetochore fibres. Nat Cell Biol *7*, 42-47.

Matos, I., Pereira, A.J., Lince-Faria, M., Cameron, L.A., Salmon, E.D., and Maiato, H. (2009). Synchronizing chromosome segregation by flux-dependent force equalization at kinetochores. J Cell Biol *186*, 11-26.

Matsuda, T., and Nagai, T. (2014). Quantitative measurement of intracellular protein dynamics using photobleaching or photoactivation of fluorescent proteins. Microscopy (Oxf) *63*, 403-408.

Matthews, L.R., Carter, P., Thierry-Mieg, D., and Kemphues, K. (1998). ZYG-9, a Caenorhabditis elegans protein required for microtubule organization and function, is a component of meiotic and mitotic spindle poles. J Cell Biol *141*, 1159-1168.

McNally, K., Audhya, A., Oegema, K., and McNally, F.J. (2006). Katanin controls mitotic and meiotic spindle length. J Cell Biol *175*, 881-891.

McVey, S.L., Cosby, J.K., and Nannas, N.J. (2021). Aurora B Tension Sensing Mechanisms in the Kinetochore Ensure Accurate Chromosome Segregation. Int J Mol Sci *22*.

Mitchison, T.J. (1989). Polewards microtubule flux in the mitotic spindle: evidence from photoactivation of fluorescence. J Cell Biol *109*, 637-652.

Mitchison, T.J., and Salmon, E.D. (1992). Poleward kinetochore fiber movement occurs during both metaphase and anaphase-A in newt lung cell mitosis. J Cell Biol *119*, 569-582.

Miyamoto, D.T., Perlman, Z.E., Burbank, K.S., Groen, A.C., and Mitchison, T.J. (2004). The kinesin Eg5 drives poleward microtubule flux in Xenopus laevis egg extract spindles. J Cell Biol *167*, 813-818.

Muller-Reichert, T., Greenan, G., O'Toole, E., and Srayko, M. (2010). The elegans of spindle assembly. Cell Mol Life Sci *67*, 2195-2213.

Musacchio, A., and Desai, A. (2017). A Molecular View of Kinetochore Assembly and Function. Biology (Basel) *6*.

Nazockdast, E., and Redemann, S. (2020). Mechanics of the spindle apparatus. Seminars in cell & developmental biology *107*, 91-102.

Nicklas, R.B. (1997). How cells get the right chromosomes. Science *275*, 632-637.

O'Toole, E.T., McDonald, K.L., Mantler, J., McIntosh, J.R., Hyman, A.A., and Muller-Reichert, T. (2003). Morphologically distinct microtubule ends in the mitotic centrosome of Caenorhabditis elegans. J Cell Biol *163*, 451-456.

Oegema, K., and Hyman, A.A. (2006). Cell division. WormBook : the online review of C elegans biology, 1-40.

Pereira, A.J., and Maiato, H. (2012). Maturation of the kinetochore-microtubule interface and the meaning of metaphase. Chromosome Res *20*, 563-577.

Petry, S. (2016). Mechanisms of Mitotic Spindle Assembly. Annu Rev Biochem *85*, 659-683.

Pintard, L., and Bowerman, B. (2019). Mitotic Cell Division in Caenorhabditis elegans. Genetics *211*, 35-73.

Powers, J., Rose, D.J., Saunders, A., Dunkelbarger, S., Strome, S., and Saxton, W.M. (2004). Loss of KLP-19 polar ejection force causes misorientation and missegregation of holocentric chromosomes. J Cell Biol *166*, 991-1001.

Praitis, V., Casey, E., Collar, D., and Austin, J. (2001). Creation of low-copy integrated transgenic lines in Caenorhabditis elegans. Genetics *157*, 1217-1226.

Quenouille, M.H. (1956). Notes on Bias in Estimation. Biometrika *43*.

Redemann, S., Baumgart, J., Lindow, N., Shelley, M., Nazockdast, E., Kratz, A., Prohaska, S., Brugues, J., Furthauer, S., and Muller-Reichert, T. (2017). C. elegans chromosomes connect to centrosomes by anchoring into the spindle network. Nat Commun *8*, 15288.

Riddle, D.L. (1997). C. elegans II (Plainview, N.Y.: Cold Spring Harbor Laboratory Press).

Risteski, P., Bozan, D., Jagric, M., Bosilj, A., Pavin, N., and Tolic, I.M. (2022). Length-dependent poleward flux of sister kinetochore fibers promotes chromosome alignment. Cell reports *40*, 111169.

Roostalu, J., Rickman, J., Thomas, C., Nedelec, F., and Surrey, T. (2018). Determinants of Polar versus Nematic Organization in Networks of Dynamic Microtubules and Mitotic Motors. Cell *175*, 796-808 e714.

Rueden, C., Schindelin, J., Hiner, M., and Eliceiri, K. (2021). SciJava Common. https://scijava.org/.

Sarov, M., Murray, J.I., Schanze, K., Pozniakovski, A., Niu, W., Angermann, K., Hasse, S., Rupprecht, M., Vinis, E., Tinney, M.*, et al.* (2012). A genome-scale resource for in vivo tag-based protein function exploration in C. elegans. Cell *150*, 855-866.

Schechtman, E., and Wang, S. (2004). Jackknifing two-sample statistics. Journal of Statistical Planning and Inference *119*, 329-340.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B.*, et al.* (2012). Fiji: an open-source platform for biological-image analysis. Nat Methods *9*, 676-682.

Schmidt, J.C., Arthanari, H., Boeszoermenyi, A., Dashkevich, N.M., Wilson-Kubalek, E.M., Monnier, N., Markus, M., Oberer, M., Milligan, R.A., Bathe, M.*, et al.* (2012). The kinetochore-

bound Ska1 complex tracks depolymerizing microtubules and binds to curved protofilaments. Dev Cell *23*, 968-980.

Schwietert, F., Volkov, V.A., Huis In 't Veld, P.J., Dogterom, M., Musacchio, A., and Kierfeld, J. (2022). Strain stiffening of Ndc80 complexes attached to microtubule plus ends. Biophys J *121*, 4048-4062.

Sonnichsen, B., Koski, L.B., Walsh, A., Marschall, P., Neumann, B., Brehm, M., Alleaume, A.M., Artelt, J., Bettencourt, P., Cassin, E.*, et al.* (2005). Full-genome RNAi profiling of early embryogenesis in Caenorhabditis elegans. Nature *434*, 462-469.

Srayko, M., Kaya, A., Stamford, J., and Hyman, A.A. (2005). Identification and characterization of factors required for microtubule growth and nucleation in the early C. elegans embryo. Dev Cell *9*, 223-236.

Steblyanko, Y., Rajendraprasad, G., Osswald, M., Eibes, S., Jacome, A., Geley, S., Pereira, A.J., Maiato, H., and Barisic, M. (2020). Microtubule poleward flux in human cells is driven by the coordinated action of four kinesins. EMBO J, e105432.

Steinmetz, M.O., and Prota, A.E. (2018). Microtubule-Targeting Agents: Strategies To Hijack the Cytoskeleton. Trends Cell Biol *28*, 776-792.

Sugioka, K., Fielmich, L.E., Mizumoto, K., Bowerman, B., van den Heuvel, S., Kimura, A., and Sawa, H. (2018). Tumor suppressor APC is an attenuator of spindle-pulling forces during C. elegans asymmetric cell division. Proc Natl Acad Sci U S A *115*, E954-E963.

Suzuki, A., Badger, B.L., Haase, J., Ohashi, T., Erickson, H.P., Salmon, E.D., and Bloom, K. (2016). How the kinetochore couples microtubule force and centromere stretch to move chromosomes. Nat Cell Biol *18*, 382-392.

Suzuki, A., Badger, B.L., and Salmon, E.D. (2015). A quantitative description of Ndc80 complex linkage to human kinetochores. Nat Commun *6*, 8161.

Tarailo, M., Kitagawa, R., and Rose, A.M. (2007). Suppressors of spindle checkpoint defect (such) mutants identify new mdf-1/MAD1 interactors in Caenorhabditis elegans. Genetics *175*, 1665-1679.

Timmons, L., and Fire, A. (1998). Specific interference by ingested dsRNA. Nature *395*, 854.

Tinevez, J.Y., Dragavon, J., Baba-Aissa, L., Roux, P., Perret, E., Canivet, A., Galy, V., and Shorte, S. (2012). A quantitative method for measuring phototoxicity of a live cell imaging microscope. Methods Enzymol *506*, 291-309.

Uteng, M., Hentrich, C., Miura, K., Bieling, P., and Surrey, T. (2008). Poleward transport of Eg5 by dynein-dynactin in Xenopus laevis egg extract spindles. J Cell Biol *182*, 715-726.

Vicente, J.J., and Wordeman, L. (2019). The quantification and regulation of microtubule dynamics in the mitotic spindle. Curr Opin Cell Biol *60*, 36-43.

Wandke, C., Barisic, M., Sigl, R., Rauch, V., Wolf, F., Amaro, A.C., Tan, C.H., Pereira, A.J., Kutay, U., Maiato, H.*, et al.* (2012). Human chromokinesins promote chromosome congression and spindle microtubule dynamics during mitosis. J Cell Biol *198*, 847-863.

Waters, J.C., Mitchison, T.J., Rieder, C.L., and Salmon, E.D. (1996). The kinetochore microtubule minus-end disassembly associated with poleward flux produces a force that can do work. Mol Biol Cell *7*, 1547-1558.

White, J., and Stelzer, E. (1999). Photobleaching GFP reveals protein dynamics inside live cells. Trends Cell Biol *9*, 61-65.

Wolff, I.D., Hollis, J.A., and Wignall, S.M. (2022). Acentrosomal spindle assembly and maintenance in Caenorhabditis elegans oocytes requires a kinesin-12 nonmotor microtubule interaction domain. Mol Biol Cell *33*, ar71.

Woodruff, J.B., Wueseke, O., Viscardi, V., Mahamid, J., Ochoa, S.D., Bunkenborg, J., Widlund, P.O., Pozniakovsky, A., Zanin, E., Bahmanyar, S.*, et al.* (2015). Centrosomes. Regulated assembly of a supramolecular centrosome scaffold in vitro. Science *348*, 808-812.

Wordeman, L., and Vicente, J.J. (2021). Microtubule Targeting Agents in Disease: Classic Drugs, Novel Roles. Cancers (Basel) *13*.

Wordeman, L., Wagenbach, M., and von Dassow, G. (2007). MCAK facilitates chromosome movement by promoting kinetochore microtubule turnover. J Cell Biol *179*, 869-879.

Yang, G., Cameron, L.A., Maddox, P.S., Salmon, E.D., and Danuser, G. (2008). Regional variation of microtubule flux reveals microtubule organization in the metaphase meiotic spindle. J Cell Biol *182*, 631-639.

Ye, A.A., Cane, S., and Maresca, T.J. (2016). Chromosome biorientation produces hundreds of piconewtons at a metazoan kinetochore. Nat Commun *7*, 13221.

Zhai, Y., Kronebusch, P.J., and Borisy, G.G. (1995). Kinetochore microtubule dynamics and the metaphase-anaphase transition. J Cell Biol *131*, 721-734.

Zhang, D., Rogers, G.C., Buster, D.W., and Sharp, D.J. (2007). Three microtubule severing enzymes contribute to the "Pacman-flux" machinery that moves chromosomes. J Cell Biol *177*, 231-242.

Zimyanin, V., Magaj, M., M., Yu, C.-H., Gibney, T., V., Mustafa, B., Ashley, X., Siller, K., Cueff, L., Bouvrais, H., Pecreaux, J.*, et al.* (2023). Lack of chromokinesin Klp-19 creates a more rigid midzone and affects force transmission during anaphase in C. elegans. bioRxiv, 2023.2010.2026.564275.