Non-invasive stimulation of the human striatum disrupts reinforcement learning of motor skills

Pierre Vassiliadis[1,2,3], Elena Beanato[1,2], Traian Popa[1,2], Fabienne Windel[1,2], Takuya Morishita[1,2], Esra Neufeld[4], Julie Duque[3], Gerard Derosiere[3,5], Maximilian J. Wessel[1,2,6] and Friedhelm C. Hummel[1,2,7] *

[1]Defitech Chair of Clinical Neuroengineering, Neuro-X Institute (INX) and Brain Mind Institute (BMI), École Polytechnique Fédérale de Lausanne (EPFL), 1202 Geneva, Switzerland.

[2]Defitech Chair of Clinical Neuroengineering, INX and BMI, EPFL Valais, Clinique Romande de Réadaptation,1951 Sion, Switzerland.

[3]Institute of Neuroscience, Université Catholique de Louvain, 1200, Brussels, Belgium

[4]Foundation for Research on Information Technologies in Society, Zurich, Switzerland

[5]Lyon Neuroscience Research Center – Impact team - Inserm U1028 – CNRS UMR5292, Lyon 1 University, Bron, France

[6]Department of Neurology, University Hospital, Julius-Maximilians-University, Würzburg, Germany

[7]Clinical Neuroscience, University of Geneva Medical School, 1202 Geneva, Switzerland.

* CORRESPONDENCE TO:

Friedhelm C. Hummel

Neuro-X Institute (INX)

École Polytechnique Fédérale de Lausanne (EPFL)

Campus Biotech, Room H4.3.132.084

Chemin des Mines 9, 1202 Geneva, Switzerland

and

EPFL Valais,

Clinique Romande de Réadaptation

Av. Grand-Champsec 90,

CH-1951 Sion

Email: friedhelm.hummel@epfl.ch

Abstract

Reinforcement feedback can improve motor learning, but the underlying brain mechanisms remain underexplored. Especially, the causal contribution of specific patterns of oscillatory activity within the human striatum is unknown. To address this question, we exploited an innovative, non-invasive deep brain stimulation technique called transcranial Temporal Interference Stimulation (tTIS) during reinforcement motor learning with concurrent neuroimaging, in a randomised, sham-controlled, double-blind study. Striatal tTIS applied at 80Hz, but not at 20Hz, abolished the benefits of reinforcement on motor learning. This effect was related to a selective modulation of neural activity within the striatum. Moreover, 80Hz, but not 20Hz tTIS increased the neuromodulatory influence of the striatum on frontal areas involved in reinforcement motor learning. These results show for the first time that tTIS can non-invasively and selectively modulate a striatal mechanism involved in reinforcement learning, opening new horizons for the study of causal relationships between deep brain structures and human behaviour.

Keywords:

Motor learning, reward, reinforcement learning, non-invasive brain stimulation, deep brain stimulation, temporal interference stimulation, striatum, neuroimaging

1. Introduction

The ability to learn from past outcomes, often referred to as reinforcement learning, is fundamental for complex biological systems[1]. Reinforcement learning has been classically studied in the context of decision making, when agents have to decide between a discrete number of potential options[2]. Importantly, there is an increasing recognition that reinforcement learning processes are also at play in other contexts including during practice of a new motor skill[3–5]. For instance, the addition of reinforcement feedback during motor training can improve motor learning, presumably by boosting the retention of newly acquired motor memories[6,7]. Interestingly, reinforcement feedback also appears to be relevant for the rehabilitation of patients suffering from motor impairments[8–10]. Yet, despite these promising results, there is currently a limited understanding of the brain mechanisms that are critical to implement this behaviour.

A prominent hypothesis in the field is that the striatum, a structure that is particularly active both during reinforcement[11] and motor learning[12], may be causally involved in the beneficial effects of reinforcement on motor learning. As such, the striatum shares dense connexions with dopaminergic structures of the midbrain as well as with pre-frontal and motor cortical regions[13], and is therefore well positioned to mediate reinforcement motor learning[14–16]. This idea is supported by neuroimaging studies showing reward-related activation of the striatum during motor learning[17,18]. More specifically, within the striatum, oscillatory activity in specific frequency bands is suggested to be involved in aspects of reinforcement processing. Previous rodent studies have shown that striatal high gamma oscillations (~ 80 Hz) transiently increase following reward delivery[19–23], but not when reward is withheld[19]. Hence, dynamic changes of high gamma activity in the striatum[19,24,25] and in other parts of the basal ganglia[26,27] may encode the outcome of previous movements (i.e., success or failure) and support learning. Consistent with a role of such oscillatory activity in reinforcement learning, high gamma activity in the striatum shows coherence with frontal cortex oscillations and is up-regulated by dopaminergic agonists[19]. Hence, this body

of work suggests that reinforcement-related modulation of striatal oscillatory activity, especially in

the gamma range, may be crucial for reinforcement learning of motor skills. Conversely, striatal

beta oscillations (~20 Hz) have been largely associated with sensorimotor functions[28]. For

instance, beta oscillations in the striatum are exacerbated in Parkinson's disease and associated

to the severity of motor symptoms[29–31]. Consistently, excessive beta connectivity is reduced by

anti-parkinsonian treatment in proportion to the related motor improvement[32]. Taken together,

these elements suggest that striatal high gamma and beta activity may have different functional

roles preferentially associated to reinforcement and sensorimotor functions, respectively.

The studies mentioned above provide associative evidence linking the presence of

reinforcement with changes of neural activity within the striatum determined through

neuroimaging[17,18], but do not allow to draw conclusions regarding its causal role in reinforcement

motor learning in humans. The only causal evidence available to date comes from animal work

showing modulation of reinforcement-based decision-making with striatal stimulation[33,34]. A

reason for the current absence of investigations of the causal role of the striatum in human

behaviour is related to its deep localization in the brain. As such, current non-invasive brain

stimulation techniques, such as transcranial magnetic stimulation (TMS) or classical transcranial

electric stimulation (tES), do not allow to selectively target deep brain regions, because these

techniques exhibit a steep depth-focality trade-off[35,36]. Studies of patients with striatal lesions[37,38]

or invasive deep brain stimulation of connected nuclei[39,40] have provided insights into the role of

the basal ganglia in reinforcement learning. However, their conclusions are partially limited by the

fact that the studied patients also exhibit altered network properties resulting from the underlying

pathology (e.g., neurodegeneration, lesions) or from the respective compensatory mechanisms.

Here, we address these challenges by exploiting transcranial electric Temporal Interference

Stimulation (tTIS), a new non-invasive brain stimulation approach allowing to target deep brain

regions in a frequency-specific and focal manner in the physiological state[41,42].

The concept of tTIS was initially proposed and validated on the hippocampus of rodents[41]

and was then further tested through computational modelling[43–47] and in first applications on

cortical areas in humans[48,49]. tTIS requires two pairs of electrodes to be placed on the head, each

pair delivering a high frequency alternating current. One key element is that this frequency has to

be high enough (i.e., in the kHz range) to avoid direct neuronal entrainment, based on the low-

pass filtering properties of neuronal membranes[50]. The second key element is the application of a

small difference of frequency between the two alternating currents. The superposition of the

electric fields creates an envelope oscillating at this low-frequency difference, which can be

steered towards individual deep brain structures (e.g., by optimizing electrodes' placement), and

is in a range able to influence neuronal activity [41,51–53]. An interesting feature of tTIS is to stimulate

at a particular frequency of interest in order to preferentially interact with specific neuronal

processes[41,42]. Importantly, despite these exciting opportunities, current evidence for tTIS-related

neuromodulation of deep brain structures, such as the striatum, is lacking in humans.

Here, we combine tTIS with electric field modelling for target localisation, behavioural data

and functional magnetic resonance imaging (fMRI) to evaluate the causal role of specific patterns

of striatal activity in reinforcement learning of motor skills. Based on the studies mentioned above,

we hypothesised that striatal tTIS at high gamma frequency ($tTIS_{80Hz}$) would disturb the fine-tuning

of high gamma oscillatory activity in the striatum and thereby would perturb reinforcement motor

learning in contrast to beta ($tTIS_{20Hz}$) or sham ($tTIS_{Sham}$) stimulation. More specifically, we

reasoned that applying a constant high gamma rhythm in the striatum would disturb the temporally

precise and reinforcement-specific modulation of high gamma activity. Moreover, given that the

stimulation protocol was not individualised to endogenous high gamma activity and not

synchronised to ongoing activity in other hubs of the reinforcement learning network (e.g., the

frontal cortex), we anticipated disruptive rather than beneficial effects of $tTIS_{80Hz}$.

In line with our prediction, we report that $tTIS_{80Hz}$ disrupted motor learning compared to the

controls, but only in the presence of reinforcement. To evaluate the potential neural correlates of

these behavioral effects, we measured BOLD activity in the striatum and effective connectivity

between the striatum and frontal cortical areas involved in reinforcement motor learning. We found

that the disruptive effect of $tTIS_{80Hz}$ on reinforcement learning was associated to a specific

modulation of BOLD activity in the putamen and caudate, but not in the cortex, supporting the

ability of tTIS to selectively modulate striatal activity without affecting overlying cortical areas.

Moreover, $tTIS_{80Hz}$ also increased the neuromodulatory influence of the striatum on frontal cortical

areas involved in reinforcement motor learning. Overall, the present study shows for the first time

that tTIS can non-invasively and selectively modulate a striatal mechanism involved in

reinforcement learning opening new horizons for the study of causal relationships between deep

brain structures and human behaviour.

2. Results

24 healthy participants (15 women, 25.3 ± 0.1 years old; mean ± SE) performed a force tracking task in the MRI with concurrent tTIS of the striatum. The task required participants to modulate the force applied on a hand-grip force sensor in order to track a moving target with a cursor with the right, dominant hand[54,55] (**Figure 1A**). At each block, participants had to learn a new pattern of motion of the target (**Figure S1**; see Methods). In $Reinf_{ON}$ blocks, participants were provided with online reinforcement feedback during training, giving them real-time information about success or failure throughout the trial, indicated as a green or red target, respectively (please see **Video S1** for the task). The reinforcement feedback was delivered according to a closed-loop schedule[8], in which the success criterion to consider a force sample as successful was updated based on the median performance over the 4 previous trials (see Methods for more details). In $Reinf_{OFF}$ blocks, participants practiced with a visually matched random feedback (cyan/magenta). Importantly, in both types of blocks, training was performed with partial visual feedback of the cursor, a condition that has been shown to maximise reinforcement effects in various motor learning paradigms[4,56–58] and which yielded significant effects of reinforcement on motor learning as also demonstrated in an additional behavioural study testing another group of healthy participants on the same task (n = 24, Figure S2). Before and after training, participants performed Pre- and Post-training assessments with full visual feedback, no reinforcement and no tTIS, allowing us to evaluate motor learning. To assess the effect of tTIS on reinforcement-related benefits in motor learning and the associated neural changes, participants performed 6 blocks of 36 trials in the MRI, with concurrent tTIS during training, delivered with a Δf of 20 Hz ($tTIS_{20Hz}$), 80 Hz ($tTIS_{80Hz}$) or as a sham ($tTIS_{Sham}$; 3 $tTIS_{TYPE}$ x 2 $Reinf_{TYPE}$ conditions; **Figure 1B, 1C**). Notably, the order of the conditions was balanced among the 24 participants, ensuring that any potential carry-over effect would have the same impact on each experimental condition. To determine the best electrode montage to stimulate the human striatum (putamen, caudate and nucleus

161 accumbens [NAc] bilaterally), computational modelling with a realistic head model was conducted

162 with Sim4Life[59] (see Methods). The selected montage (F3-F4; TP7-TP8) generated a theoretical

163 temporal interference electric field that was ~30-40% stronger in the striatum than in the overlying

164 cortex, reaching magnitudes of 0.5 to 0.6 V/m (**Figure 1D, 1E**).

Figure 1. Striatal tTIS during reinforcement learning of motor skills in the MRI. A)
Motor learning task. Participants were required to squeeze a hand grip force sensor (depicted in
the upper right corner of the figure) in order to track a moving target (larger circle with a cross in
the center) with a cursor (black smaller circle)[54,55]. Pre- and Post-training assessments were
performed with full visual feedback of the cursor and no reinforcement. In Reinf$_{ON}$ and Reinf$_{OFF}$
trials, participants practiced the task with or without reinforcement feedback, respectively. As such,
in Reinf$_{ON}$ trials, the color of the target varied in real-time as a function of the subjects' tracking
performance. **B) Experimental procedure.** Participants performed the task in the MRI with
concomitant TI stimulation. Blocks of training were composed of 36 trials (4 Pre-, 24 Training and
8 Post-training trials) interspersed with short resting periods (represented as + on the figure). The
6 training types resulted from the combination of 3 tTIS$_{TYPES}$ and 2 Reinf$_{TYPES}$. **C) Concept of tTIS.**
On the left, two pairs of electrodes are shown on a head model and currents are applied with a

frequency f1 and f1+Δf. On the right, the interference of the two electric fields within the brain is
represented for two different locations with respectively high and low envelope modulation. $E_1(t)$
and $E_2(t)$ represent the modulation of the fields' magnitude over time. tTIS was delivered either
with a Δf of 20 or 80 Hz or as a sham (ramp-up and immediate ramp-down of high frequency
currents with flat envelope). **D) Electric field modelling with the striatal montage.** Temporal
interference exposure (electric field modulation magnitude). **E) Temporal interference exposure**
averaged in the striatum and in the overlying cortex. Magnitude of the field in the cortex was
extracted from the Brainnetome atlas (BNA[60]) regions underneath the stimulation electrodes (F3-
F4 and TP7-TP8). Error bars represent the standard deviation over the voxels in the considered
region.

tTIS$_{80Hz}$ disrupts reinforcement learning of motor skills

Task performance was evaluated by means of the Error, which was defined as the absolute

difference between the applied and target force averaged across samples for each trial, as done

previously[4,54,56] (**Figure 2A**). Across conditions, the Post-training Error was reduced compared to

the Pre-training Error (single sample t-test on the normalised Post-training data: $t_{(24)}$=-2.69;

p=0.013; Cohen's d=-0.55), indicating significant motor learning during the task (**Figure 2B**). Such

improvement was greater when participants had trained with reinforcement (Reinf$_{TYPE}$ effect in the

Linear Mixed Model (LMM): $F_{(1, 1062.2)}$=5.17; p=0.023; d=-0.14 for the post-hoc contrast Reinf$_{ON}$ –

Reinf$_{OFF}$), confirming the beneficial effect of reinforcement on motor learning[7,58]. Crucially though,

this effect depended on the type of stimulation applied during training (Reinf$_{TYPE}$ x tTIS$_{TYPE}$

interaction: $F_{(2, 1063.5)}$=2.11; p=0.034; **Figure 2C**). While reinforcement significantly improved

learning when training was performed with tTIS$_{Sham}$ (p=0.036; d=-0.22) and tTIS$_{20Hz}$ (p=0.0089;

d=-0.27), this was not the case with tTIS$_{80Hz}$ (p=0.43; d=0.083). Consistently, direct between-

condition comparisons showed that in the Reinf$_{ON}$ condition, learning was reduced with tTIS$_{80Hz}$

compared to tTIS$_{20Hz}$ (p=0.039; d=0.26) and tTIS$_{Sham}$ (p<0.001; d=0.45) but was not different

between tTIS$_{20Hz}$ and tTIS$_{Sham}$ (p=0.15; d=0.20). This disruption of motor learning with tTIS$_{80Hz}$ was

not observed in the absence of reinforcement (tTIS$_{80Hz}$ vs. tTIS$_{20Hz}$: p=0.59; d=-0.10, tTIS$_{80Hz}$ vs.

tTIS$_{Sham}$: p=0.34; d=0.15). These results strongly point to the fact that high gamma striatal tTIS

specifically disrupts the benefits of reinforcement on motor learning and not motor learning in

general.

Although training with $tTIS_{20Hz}$ did not alter the benefits of reinforcement on motor learning,

we found that learning without reinforcement was significantly impaired in this condition ($tTIS_{20Hz}$

vs. $tTIS_{Sham}$: p=0.046; d=0.25, **Figure 2C**). This suggests that $tTIS_{20Hz}$ may disrupt a qualitatively

different mechanism involved in motor learning from sensory feedback[61], in line with the role of

striatal beta oscillations in sensorimotor function[28].

Next, we evaluated the effect of tTIS on motor performance during training itself. As shown

in Figure 2A, the Error was generally higher during Training than in Test trials due to the presence

of visual uncertainty during this phase. The extent of this disruption was reduced in the presence

of reinforcement ($Reinf_{TYPE}$: $F_{(1, 3262.4)}$=339.89; p<0.001; d=-0.64 for the contrast $Reinf_{ON}$ –

$Reinf_{OFF}$), demonstrating the ability of subjects to exploit real-time reinforcement information to

improve tracking (**Figure 2D**). Notably, this effect was not modulated by $tTIS_{TYPE}$ ($Reinf_{TYPE}$ x

$tTIS_{TYPE}$: $F_{(2, 3265.8)}$=0.91; p=0.40), indicating that tTIS did not directly influence reinforcement gains

during tracking. Interestingly though, striatal stimulation did impact on general tracking

performance independently of reinforcement as indicated by a significant $tTIS_{TYPE}$ effect ($tTIS_{TYPE}$:

$F_{(2, 3262.4)}$=42.85; p<0.001). This effect was due to an increase in the Error when $tTIS_{20Hz}$ was

applied (p<0.001; d=0.28 when compared to $tTIS_{Sham}$), which was even stronger during $tTIS_{80Hz}$

(p<0.001; d=0.38 and p=0.031; d=0.11 when compared to $tTIS_{Sham}$ and $tTIS_{20Hz}$, respectively). An

additional analysis showed that the detrimental effect of tTIS on motor performance was actually

due to an impaired ability to improve performance during Training (LMM with continuous fixed

effect Trial: $tTIS_{TYPE}$ x Trial interaction: $F_{(2, 3399)}$=4.46; p=0.012, post-hoc tests: $tTIS_{Sham}$ vs. $tTIS_{20Hz}$:

p=0.013; $tTIS_{Sham}$ vs. $tTIS_{80Hz}$: p=0.068; $tTIS_{20Hz}$ vs. $tTIS_{80Hz}$: p=0.81; **Figure S3**). However, again,

this effect did not depend on the presence of reinforcement ($Reinf_{TYPE}$ x $tTIS_{TYPE}$ x Trial: $F_{(2,}$

$_{3399)}$=0.51; p=0.60). Notably, we also found that the detrimental effect of striatal tTIS did not depend

on the availability of visual information on the cursor, but rather that tTIS had a general effect on motor performance irrespective of visual and reinforcement feedback (see Supplementary materials). This analysis also confirmed that reinforcement gains in motor performance were stronger when visual information was not available (**Figure S4**), in line with the behavioural data mentioned above (Figure S2) and previous studies[57,62]. Overall, these results suggest that striatal tTIS altered motor performance in a frequency-dependent manner but did not influence the ability to rapidly adjust motor commands based on reinforcement feedback during training. Hence, $tTIS_{80Hz}$ may not disrupt real-time processing of reinforcement feedback, but may rather impair the beneficial effect of reinforcements on the retention of motor memories[6,7].

Notably, these effects could not be explained by potential differences in initial performance between conditions ($Reinf_{TYPE}$ x $tTIS_{TYPE}$: $F_{(2, 519.99)}$=1.08; p=0.34), nor by changes in the flashing properties of the reinforcement feedback (i.e., the frequency of color change during tracking; $Reinf_{TYPE}$ x $tTIS_{TYPE}$: $F_{(2, 3283)}$=0.19; p=0.82), or by differences in success rate in the $Reinf_{ON}$ blocks (i.e., the proportion of success feedback during tracking; $tTIS_{TYPE}$: $F_{(2, 1702)}$=0.17; p=0.84). The $Reinf_{TYPE}$ x $tTIS_{TYPE}$ effect on learning was also not influenced by the order of the reinforcement conditions (analysis on sub-groups based on whether participants experienced $Reinf_{ON}$ or $Reinf_{OFF}$ first; no $Reinf_{TYPE}$ x $tTIS_{TYPE}$ x $Group_{TYPE}$ interaction: $F_{(2,1105.06)}$=1.75; p=0.17; see Supplementary materials for more details on these analyses).

Finally, we confirmed that these results were not a consequence of an inefficient blinding. As such, when debriefing after the experiment, only 6/24 participants were able to successfully identify the order of the stimulation applied (e.g., real – real – placebo; chance level: 4/24; Fisher exact test on proportions: p=0.74). Consistently, the magnitude (**Figure S5A**) and type (**Figure S5B**) of tTIS-evoked sensations evaluated before the experiment were qualitatively similar across conditions and tTIS was generally well tolerated in all participants (no adverse events reported). This suggests that blinding was successful and is unlikely to explain our findings. More generally,

257 this is a first indication that tTIS evokes very limited sensations (e.g., only 2/24 and 1/24 subjects

258 rated sensations evoked at 2 mA as "strong" for tTIS$_{20Hz}$ and tTIS$_{80Hz}$, respectively; **Figure S5A**)

259 that are compatible with efficient blinding.

Figure 2. Behavioural results. A) Motor performance across training. Raw Error data (expressed in % of Maximum Voluntary Contraction [MVC]) are presented on the left panel for the different experimental conditions in bins of 4 trials. The increase in Error during Training is related to the visual uncertainty (i.e., intermittent disappearance of the cursor) that was applied to enhance reinforcement effects. On the right, the three plots represent the Pre-training normalised Error in the tTIS$_{Sham}$, tTIS$_{20Hz}$ and tTIS$_{80Hz}$ blocks. Reinforcement-related benefits represent the improvement in the Error measured in the Reinf$_{ON}$ and Reinf$_{OFF}$ blocks, during Training (reflecting benefits in motor performance) or at Post-training (reflecting benefits in learning). **B) Averaged learning across conditions.** Violin plot showing the Error distribution at Post-training (expressed in % of Pre-training) averaged across conditions, as well as individual subject data. A single-sample t-test showed that the Post-training Error was reduced compared to the Pre-training level, indicating significant learning in the task. **C) Motor learning.** Averaged Error at Post-training (normalised to Pre-training) and the corresponding individual data points in the different experimental conditions are shown on the left and right panels, respectively, for the subjects included in the analysis (i.e., after outlier detection, remaining n=23). Reduction of Error at Post-training reflects true improvement at tracking the target in Test conditions (in the absence of reinforcement, visual uncertainty or tTIS). The LMM ran on these data revealed a specific effect

of tTIS$_{80Hz}$ on reinforcement-related benefits in learning. **D) Motor performance.** Averaged Error
during Training (normalised to Pre-training) and the corresponding individual data points in the
different experimental conditions are shown on the left and right panels, respectively, for the
subjects included in the analysis (i.e., after outlier detection, n=23). Individual data points are
shown on the right panel. Error change during Training reflect the joint contribution of the
experimental manipulations (visual uncertainty, potential reinforcement and tTIS) on motor
performance. The LMM ran on these data showed a frequency-dependent effect of tTIS on motor
performance, irrespective of reinforcement. *: p<0.05. Data are represented as mean ± SE.

The effect of tTIS$_{80Hz}$ on reinforcement motor learning is related to modulation of neural

activity in the striatum

As mentioned above, task-based fMRI was acquired during Training with concomitant tTIS.

This allowed us to evaluate the neural effects of tTIS and their potential relationship to the

behavioural effects reported above. As a first qualitative evaluation of the data, we performed a

whole-brain analysis in the tTIS$_{Sham}$ condition to assess the network activated during reinforcement

motor learning (Reinf$_{ON}$ condition). Consistent with previous neuroimaging studies employing

similar tasks[63,64], we found prominent BOLD activations in a motor network including the putamen,

thalamus, cerebellum and sensorimotor cortex, particularly on the left hemisphere, contralateral

to the trained hand (**Figure S6, Table S2**). Notably though, contrasting Reinf$_{ON}$ and Reinf$_{OFF}$

conditions did not reveal any significant cluster at the whole-brain level. Hence, this first analysis

did not reveal any region specifically activated in the presence of reinforcement, but rather

confirms the involvement of a motor network engaged in this type of task irrespective of the

reinforcement feedback.

As a second step, we evaluated the effect of tTIS on striatal activity, as a function of the

type of reinforcement feedback and focusing on the very same regions of interest (ROI) that were

used to optimise tTIS exposure in the modelling. Based on this, we extracted averaged BOLD

activity within the bilateral putamen, caudate and NAc based on the Brainnetome atlas (BNA[60]),

in the different experimental conditions and considered these six striatal ROIs (ROI$_{STR}$) as fixed

effects in the LMM. This model revealed a strong enhancement of striatal activity with Reinf$_{ON}$ with

respect to $Reinf_{OFF}$ ($F_{(1, 800.01)}$=13.23; p<0.001; d=0.25 for the contrast $Reinf_{ON}$ – $Reinf_{OFF}$)

consistent with previous literature[11], but no $tTIS_{TYPE}$ effect ($F_{(2, 800.01)}$=0.46; p=0.63) and no

interaction (all p> 0.65; **Figure 3A**). Despite the absence of effects of tTIS on averaged striatal

activity, we then asked whether the behavioural effects of $tTIS_{80Hz}$ on reinforcement motor learning

(i.e., $tTIS_{80Hz}$ vs. $tTIS_{20Hz}$ and $tTIS_{Sham}$ with $Reinf_{ON}$) could be linked to modulation of activity in core

brain regions. To do so, we ran a whole-brain analysis focusing on the main behavioural effects

mentioned above. Results revealed that the effect of $tTIS_{80Hz}$ (with respect to $tTIS_{20Hz}$) on motor

learning in the $Reinf_{ON}$ condition was specifically related to modulation of activity in two clusters

encompassing the left putamen and bilateral caudate (**Figure 3B, Table S3**). Notably, the

presence of the high frequency carrier (kHz) in both stimulation conditions rules out the possibility

that the correlation was due to putative neuromodulatory effects of high frequency stimulation. No

significant clusters were found neither for the $tTIS_{80Hz}$ – $tTIS_{Sham}$ contrast, nor for the control

$tTIS_{20Hz}$ - $tTIS_{Sham}$ contrast, indicating that the reported correlation is not due to a general link

between striatal activity and reinforcement motor learning. Overall, these results provide evidence

that the detrimental effect of $tTIS_{80Hz}$ on reinforcement learning of motor skills is related to

modulation of neural activity specifically in the striatum.

Figure 3. Striatal activity. A) Striatal BOLD responses. A 3D-reconstruction of the striatal masks used in the current experiment is surrounded by plots showing averaged BOLD activity for each mask in the different experimental conditions. A LMM ran on these data showed higher striatal responses in the Reinf$_{ON}$ with respect to the Reinf$_{OFF}$ condition, but no effect of tTIS$_{TYPE}$ and no interaction. **B) Whole-brain activity associated to the behavioural effect of tTIS$_{80Hz}$ on reinforcement motor learning.** Correlation between tTIS-related modulation of striatal activity (tTIS$_{80Hz}$ – tTIS$_{20Hz}$) and learning abilities in the Reinf$_{ON}$ condition. Significant clusters of correlation were found in the left putamen and bilateral caudate (uncorrected voxel-wise FWE: p=0.001, and corrected cluster-based FDR: p=0.05). Lower panel shows individual correlations for the three significant regions highlighted in the whole-brain analysis. *: p<0.05. Data are represented as mean ± SE.

tTIS$_{80Hz}$ enhances effective connectivity between the striatum and frontal cortex.

Interactions between the striatum and frontal cortex are crucial for a variety of behaviours including motor and reinforcement learning[13]. In particular, reinforcement motor learning requires to use information about task success to guide future motor commands[4], a process for which the striatum may play an integrative role at the interface between fronto-striatal loops involved in reward processing and motor control[13,65]. In a subsequent analysis, we asked whether striatal tTIS modulates striatum to frontal cortex communication during reinforcement motor learning. More specifically, we computed effective connectivity (using the generalized psychophysiological interactions method[66]) between striatal and frontal regions classically associated with motor and reward-related functions, and thought to be involved in reinforcement motor learning[67,68]. For the motor network, we evaluated effective connectivity between motor parts of the striatum (i.e., dorso-lateral putamen (dlPu) and dorsal caudate (dCa)) and two regions strongly implicated in motor learning: the medial part of the supplementary motor area (SMA) and the part of the primary motor cortex (M1) associated to upper limb functions (**Figure 4A**). For the reward network, we assessed connectivity between parts of the striatum classically associated to limbic functions (i.e., the NAc and the ventro-medial putamen (vmPu) and two frontal areas involved in reward processing: the anterior cingulate cortex (ACC) and the ventro-medial prefrontal cortex (vmPFC; **Figure 4B**; [11]). The LMM ran with the fixed effects Reinf$_{TYPE}$, tTIS$_{TYPE}$ and Network$_{TYPE}$ showed a significant effect of tTIS$_{TYPE}$ (F$_{(2, 2264.0)}$=5.42; p=0.0045), that was due to higher connectivity in the tTIS$_{80Hz}$ condition with respect to tTIS$_{Sham}$ (p=0.0038; d=0.16) and tTIS$_{20Hz}$ (at the trend level, p=0.069; d=0.11). There was no difference in connectivity between tTIS$_{20Hz}$ and tTIS$_{Sham}$ (p=0.58; d=0.051). Hence, tTIS$_{80Hz}$, but not tTIS$_{20Hz}$, enhanced effective connectivity between the striatum and frontal cortex during motor training. This increase in effective connectivity with tTIS$_{80Hz}$ actually led to a connectivity closer to the resting state (values closer to 0, see Methods). Put differently, while the task induced

a reduction in effective connectivity between striatum and frontal cortex, $tTIS_{80Hz}$ disrupted this

modulation by bringing connectivity back to the resting state.

The LMM did not reveal any effect of $Reinf_{TYPE}$ ($F_{(1, 2264.0)}$=0.010; p=0.92), $Network_{TYPE}$ ($F_{(1,}$

$_{2264.0)}$=3.16; p=0.076) and no double interaction (note the trend for a $Reinf_{TYPE}$ x $Network_{TYPE}$ effect

though: $F_{(1, 2264.0)}$=3.52; p=0.061). Yet, we did find a significant $Reinf_{TYPE}$ x $tTIS_{TYPE}$ x $Network_{TYPE}$

interaction ($F_{(2, 2264.0)}$=4.87; p=0.0078). Such triple interaction was related to the fact that $tTIS_{80Hz}$

increased connectivity in the $Reinf_{ON}$ condition in the motor network ($Reinf_{ON}$ vs. $Reinf_{OFF}$:

p=0.0012; d=0.33; Figure 4A), while it tended to have the opposite effect in the reward network

(p=0.063; d=-0.19; Figure 4B). This increase was not present in any of the two networks when

either $tTIS_{Sham}$ or $tTIS_{20Hz}$ were applied (all p> 0.40). Moreover, in the motor network, connectivity

in the $Reinf_{ON}$ condition was higher with $tTIS_{80Hz}$ than with $tTIS_{Sham}$ (p<0.001; d=0.42) and $tTIS_{20Hz}$

(at the trend level; p=0.059; d=0.23, Figure 4A). These data suggest that $tTIS_{80Hz}$ enhanced the

neuromodulatory influence of the striatum on motor cortex during task performance, but only in

the presence of reinforcement. In the reward network, post-hocs revealed that connectivity in the

$Reinf_{OFF}$ condition was significantly higher with $tTIS_{80Hz}$ compared to $tTIS_{20Hz}$ (p=0.045; d=0.25;

Figure 4B), in line with the general effect of $tTIS_{TYPE}$ on connectivity reported above. This pattern

of results suggests that the increase of connectivity from striatum to frontal cortex observed with

$tTIS_{80Hz}$ depends on the presence of reinforcement, in particular in the motor network. Such

reinforcement-dependent increase of connectivity may reflect the preferential effect of $tTIS_{80Hz}$ on

striatal gamma oscillations[69] in a situation where these oscillations are already boosted by the

presence of reinforcement[19] (see Discussion).

In a subsequent analysis, we verified that these results did not depend on the specific

frontal ROIs considered in the analysis (ROI_{TYPE}: M1 and SMA in the motor network and ACC and

vmPFC in the reward network). Importantly, we did not find a $tTIS_{TYPE}$ x $Reinf_{TYPE}$ x ROI_{TYPE}

interaction neither in the motor ($F_{(2,1112)}$=0.83; p=0.44) nor in the reward network ($F_{(2,1112)}$=0.61;

p=0.54), suggesting that the main connectivity results were consistent within a network and were

not influenced by the specific frontal ROI included in the analysis (see Supplementary materials

for more details on this analysis). As an additional control, we verified that the effects of $tTIS_{TYPE}$

on connectivity could not be observed in a control network associated to language (as defined by

[70]), which was unlikely to be involved in the present task and did not include the striatum (see

Methods). As expected, effective connectivity within the language network was not modulated by

$Reinf_{TYPE}$ ($F_{(1, 547)}$=0.81; p=0.37), nor by $tTIS_{TYPE}$ ($F_{(2, 547)}$=0.58; p=0.56), or by $Reinf_{TYPE}$ x $tTIS_{TYPE}$

($F_{(2, 547)}$=0.45; p=0.64). Hence, tTIS and reinforcement-related changes in connectivity were

consistent within the considered fronto-striatal networks and not observed in a control network

unrelated to the task.

Notably, contrary to the BOLD results presented above, we did not find any correlations

between the effects of $tTIS_{80Hz}$ on connectivity and motor learning, neither in the motor (robust

linear regression: $tTIS_{80Hz}$ – $tTIS_{Sham}$: R^2=0.019; p=0.48; $tTIS_{80Hz}$ – $tTIS_{20Hz}$: R^2=0.034; p=0.54) nor

in the reward ($tTIS_{80Hz}$ – $tTIS_{Sham}$: R^2=0.037; p=0.46; $tTIS_{80Hz}$ – $tTIS_{20Hz}$: R^2<0.001; p=0.75)

network, suggesting some degree of independence between the effect of $tTIS_{80Hz}$ on

reinforcement motor learning and on effective connectivity.

Overall, these results highlight the ability of $tTIS_{80Hz}$, but not $tTIS_{20Hz}$, to modulate striatum

to frontal cortex connectivity, depending on the presence of reinforcement. However, the absence

of correlation with the behaviour suggests that this effect may not be directly associated to the

detrimental effect of $tTIS_{80Hz}$ on reinforcement motor learning or that $tTIS_{80Hz}$-related changes in

striato-frontal communication were linked to other aspects of reinforcement learning not captured

by our task.

Figure 4. Striatum to frontal cortex effective connectivity. A) Motor network. On the left, 3D reconstruction of the masks used for the motor network (i.e., dorso-lateral putamen, dorsal caudate, M1, SMA). On the right, plot showing effective connectivity from motor striatum to motor cortex in the different experimental conditions. Note the increase of connectivity with $tTIS_{80Hz}$ in the presence of reinforcement. B) Reward network. On the left, 3D reconstruction of the masks used for the reward network (i.e., ventro-medial putamen, NAc, vmPFC, ACC). On the right, plot showing effective connectivity from motor striatum to motor cortex in the different experimental conditions. ROIs were defined based on the BNA atlas[12] *: $p < 0.05$. Data are represented as mean ± SE.

Neural effects of tTIS$_{80Hz}$ depend on impulsivity

Determining individual factors that shape responsiveness to non-invasive brain stimulation approaches is a crucial step to better understand the mechanisms of action but also to envision stratification of patients in future clinical interventions[71]. A potential factor that could explain inter-individual differences in responsiveness to tTIS$_{80Hz}$ is the level of impulsivity. As such, impulsivity has been associated to changes of gamma oscillatory activity in the striatum of rats[72] and to the activity of fast-spiking interneurons in the striatum[73,74], a neuronal population that is strongly entrained to gamma rythms[19,21] and may therefore be particularly sensitive to tTIS$_{80Hz}$. In a subsequent exploratory analysis, we asked if the neural effects of tTIS$_{80Hz}$ were associated to impulsivity levels, as evaluated by a well-established independent delay-discounting questionnaire performed at the beginning of the experiment[75,76]. Strikingly, a whole-brain analysis revealed that impulsivity was associated to the effect of tTIS$_{80Hz}$ on BOLD activity (with respect to tTIS$_{20Hz}$) specifically in the left caudate nucleus (**Figure S7A, S7B, Table S4**). Moreover, the effect of tTIS$_{80Hz}$ on striatum to motor cortex connectivity reported above was negatively correlated to impulsivity both when contrasting tTIS$_{80Hz}$ with tTIS$_{Sham}$ (**Figure S7C, left**) and with tTIS$_{20Hz}$ (**Figure S7C, middle**). Such correlations were absent when contrasting tTIS$_{20Hz}$ with tTIS$_{Sham}$ (**Figure S7C, right**), as well as when considering the same contrasts in the reward instead of the motor network (see Supplementary materials for more details). Taken together, these results suggest that inter-individual variability in impulsivity might influence the neural responses to striatal tTIS$_{80Hz}$.

3. Discussion

In this study, we combined striatal tTIS with electric field modelling, behavioural and fMRI analyses to evaluate the causal role of the striatum in reinforcement learning of motor skills in healthy humans. $tTIS_{80Hz}$, but not $tTIS_{20Hz}$, disrupted the ability to learn from reinforcement feedback. This behavioural effect was associated to modulation of neural activity specifically in the striatum. As a second step, we show that $tTIS_{80Hz}$, but not $tTIS_{20Hz}$, increased the neuromodulatory influence of the striatum on connected frontal cortical areas involved in reinforcement motor learning. Finally, inter-individual variability in the neural effects of $tTIS_{80Hz}$ could be partially explained by impulsivity, suggesting that this trait may constitute a determinant of responsiveness to high gamma striatal tTIS. Overall, the present study shows for the first time that striatal tTIS can non-invasively modulate a striatal mechanism involved in reinforcement learning, opening new horizons for the study of causal relationships between deep brain structures and human behaviour.

We investigated the causal role of the human striatum in reinforcement learning of motor skills in healthy humans; a question that cannot be addressed with conventional non-invasive brain stimulation techniques. In particular, by stimulating at different frequencies, we aimed at dissociating striatal mechanisms involved in reinforcement and sensorimotor learning. In line with our main hypothesis, we found that striatal $tTIS_{80Hz}$ altered reinforcement learning of a motor skill. Such disruption was frequency- and reinforcement-specific: learning was not altered with striatal $tTIS_{20Hz}$ in the presence of reinforcement, or when striatal $tTIS_{80Hz}$ was delivered in the absence of reinforcement. The rationale to stimulate at high gamma frequency was based on previous work showing reinforcement-related modulation of gamma oscillations in the striatum[19–21,24,26,72,77] and in the frontal cortex[77–80]. Several neuronal mechanisms may contribute to the detrimental effect of $tTIS_{80Hz}$ on reinforcement motor learning. First, as $tTIS_{80Hz}$ consisted in a constant high gamma

oscillating field applied on the striatum, it may have perturbed the encoding of reinforcement

information into high gamma oscillations[19–21,25–27], preventing participants to learn the motor skill

based on different outcomes. Put differently, $tTIS_{80Hz}$ may specifically saturate high gamma activity

in the striatum preventing reinforcement-related modulations[81]. Moreover, because reinforcement

motor learning likely engages synchronised activity in a network of regions including fronto-striatal

loops, neuromodulation of a single node of the circuit may alter synchronisation of activity in the

network[81] and the temporal coordination with interacting rhythms[25]. Finally, because we did not

have access to electrophysiological recordings of oscillatory activity in the striatum, the applied

stimulation was not personalised as it did not take into account the individual high gamma

frequency peak associated to reward processing and the potential heterogeneity of gamma activity

within the striatum[24]. Hence, $tTIS_{80Hz}$ may have resulted in a frequency mismatch between the

endogenous high gamma activity and the externally imposed rhythm, that could paradoxically

result in a reduction of neuronal entrainment, in particular when the frequency mismatch is

relatively low[82]. Importantly, in contrast to striatal $tTIS_{80Hz}$, we found that $tTIS_{20Hz}$ reduced learning,

but only in the absence of reinforcement. This result fits well with the literature linking striatal beta

oscillations to sensorimotor functions[28,29,31,83–85]. Taken together, an interpretation of these results

is that different oscillations within the striatum support qualitatively distinct motor learning

mechanisms with beta activity contributing mostly to sensory-based learning and high gamma

activity being particularly important for reinforcement learning. This being said, it is important to

note that because we do not have concurrent electrophysiological recordings within the striatum,

we cannot be sure that the effects of $tTIS_{20Hz}$ and $tTIS_{80Hz}$ were related to frequency-specific

interactions with beta or high gamma rhythms respectively, or rather resulted from different

broadband responses when stimulating at these frequencies. Yet, these results still suggest that

sensory- and reinforcement-based motor learning rely on partially different neural mechanisms, in

line with previous literature[8,9,61,68,86,87].

Interestingly, striatal tTIS also impaired tracking performance during training, irrespective

of the presence of reinforcement. This frequency-dependent reduction of motor performance may

be due to altered neuronal processing in the sensorimotor striatum that may lead to less fine-tuned

motor control abilities[88]. Importantly though, tTIS did not modulate the ability of participants to

benefit from real-time reinforcement feedback during motor performance. This suggests that

striatal $tTIS_{80Hz}$ altered the beneficial effects of reinforcement on learning (as evaluated in Test

conditions at Post-training), but not on motor performance (as evaluated during Training). Such

dissociation between the effects of striatal $tTIS_{80Hz}$ on reinforcement-related gains in motor

performance and learning may be explained by the fact that these two phases of the protocol

probe different processes[7,54,56,89–91]. While improvement of motor performance with reinforcement

relies on rapid feedback corrections based on expected outcomes[67,92–95], reinforcement gains in

learning (i.e., probed in Test conditions without reinforcement) may rather reflect the beneficial

effect of reinforcement on the retention of motor memories[5,7,54,90]. This idea that mechanisms

underlying performance changes in training and retention phases are partially different is well

supported by previous motor learning literature[6,8,96]. For instance, in sensorimotor adaptation

paradigms, the presence of reward boosts motor memory retention but not the adaptation process

itself[7,97], and M1 transcranial direct current stimulation modulates the effect of reward on retention

but has no effect on the training phase[90]. Hence, a potential explanation for the present results is

that striatal $tTIS_{80Hz}$ did not disrupt rapid motor corrections based on recent outcomes during

training, but may rather alter the strengthening of the memory trace based on reinforcements[6,7].

Overall, these results are compatible with the view that specific patterns of oscillatory activity in

the striatum are involved in motor control and learning processes[31], and can be modulated with

electrical stimulation[69,98,99].

To better understand the neural effects and frequency-specificity of tTIS, we coupled

striatal tTIS and task performance with simultaneous fMRI acquisition. The imaging results support

the view that the effect of tTIS$_{80Hz}$ on reinforcement learning of motor skills was indeed related to

neuromodulation of the striatum. As such, when considering averaged BOLD activity, we found a

general increase of striatal activity when reinforcement was provided[11], but no effect of tTIS.

Crucially though, the detrimental effect of tTIS$_{80Hz}$ on reinforcement learning was related to a

specific modulation of activity in the caudate and putamen, providing evidence that the present

behavioural effects were indeed driven by focal neuromodulation of the striatum (Figure 3).

Interestingly, participants with stronger disruption of reinforcement learning at the behavioural

level were also the ones exhibiting stronger suppression of striatal activity with tTIS$_{80Hz}$ (compared

to tTIS$_{20Hz}$), suggesting that tTIS-induced reduction of striatal activity is detrimental for

reinforcement motor learning. Further analyses showed that tTIS$_{80Hz}$, but not tTIS$_{20Hz}$, increased

the neuromodulatory influence of the striatum on frontal areas known to be important for motor

learning and reinforcement processing[96,100]. More specifically, tTIS$_{80Hz}$ disrupted the task-related

decrease in connectivity observed with tTIS$_{Sham}$ and tTIS$_{20Hz}$, bringing connectivity closer to

resting-state values. Interestingly, this effect depended on the type of network considered (reward

vs. motor) and on the presence of reinforcement. Striatal tTIS$_{80Hz}$ coupled with reinforcement

increased connectivity between the motor striatum and the motor cortex while it tended to have

the opposite effect when considering the connectivity between limbic parts of the striatum and pre-

frontal areas involved in reward processing (Figure 4). This result may reflect the differential

influence of striatal tTIS on distinct subparts of the striatum, depending on their pattern of activity

during the task[52]. As such, a recent study in non-human primates showed that tACS can have

opposite effects on neuronal activity based on the initial entrainment of neurons to the target

frequency[82]. Hence, the present differential effects of tTIS$_{80Hz}$ on motor and reward striato-frontal

pathways may be due to different initial patterns of activity in these networks in the presence of

reinforcement. Electrophysiological recordings with higher temporal resolution than fMRI are

required to confirm or infirm this hypothesis. Overall, the present neuroimaging results support the

idea that the behavioural effects of striatal $tTIS_{80Hz}$ on reinforcement learning are associated to a selective modulation of striatal activity that influences striato-frontal communication.

The fact that we observed increased connectivity with $tTIS_{80Hz}$ and at the same time a disruption of behaviour may appear contradictory at first glance. Yet, multiple lines of evidence indicate that increases in connectivity are not necessarily beneficial for behaviour. For instance, the severity of motor symptoms in Parkinson's disease is associated with excessive connectivity in the beta band and reduction of such connectivity with treatment is associated to clinical improvement[29,32]. Moreover, there is evidence that excessive functional[101,102] as well as structural[103,104] connectivity in fronto-striatal circuits is associated to impulsivity. Hence, the increase in connectivity observed with $tTIS_{80Hz}$ appears to be compatible with the behavioural findings. This being said, contrary to the BOLD results, we did not find any correlation between the effects of $tTIS_{80Hz}$ on connectivity and on reinforcement motor learning, suggesting some degree of independence between these two effects. Future studies could aim at determining if $tTIS_{80Hz}$-related changes in striato-frontal communication are linked to other aspects of reward processing, not captured by our reinforcement motor learning task.

From a methodological point of view, the present results provide new experimental support to the idea that the effects of tTIS are related to amplitude modulation of electric fields deep in the brain and not to the high frequency fields themselves, in line with recent work[41,42,52]. As such, the different behavioural and neural effects of striatal $tTIS_{80Hz}$ and $tTIS_{20Hz}$ despite comparable carrier frequencies (centered on 2kHz) indicate that temporal interference was indeed the driving force of the present effects. Moreover, disruption of reinforcement motor learning with $tTIS_{80Hz}$ (relative to $tTIS_{20Hz}$) was specifically related to neuromodulation of the striatum, where the amplitude of the tTIS field was highest according to our simulations (see [51,53] for recent validations of comparable simulations in cadavers experiments). Hence, we believe that the frequency- and reinforcement-dependent tTIS effects reported here cannot be explained by direct modulation of neural activity

by the high frequency fields. Yet, disentangling the neural effects of the low-frequency envelope

and the high frequency carrier appears as an important next step to better characterise the

mechanisms underlying tTIS[47]. We also note that the tTIS field strengths achieved according to

our simulations (in the range of 0.5-0.6 V/m) were sufficient to induce behavioral and neural

effects, in line with recent data[52,53] (see also [48]). Determining the minimum effective dose for tTIS

is an important line of future research given recent simulation results suggesting that stimulation

via an amplitude modulation with high frequency carrier signals (such as arising during tTIS) may

require higher dosages compared to conventional electrical stimulation with low frequencies (such

as during tACS), likely due to the low-pass filtering properties of neurons[43,105].

Finally, the strength of the behavioural effects of tTIS can be considered small to medium[106]

(d=0.2-0.5). We note that these effect sizes are consistent with studies applying other types of

non-invasive brain stimulation in healthy young adults, both in the context of motor learning (see

[107] for a meta-analysis), and reward tasks (e.g., [108,109]), despite the much longer stimulation time

used in these studies (between 3 and 20 times longer). Overall, albeit moderate, we believe that

the present effect sizes are relevant and consistent with what can be expected from the non-

invasive brain stimulation literature.

579   Limitations

The present study includes some limitations that we would like to acknowledge. First, at

the imaging level, we did not find a significant effect of reinforcement at the whole-brain level. This

might be due to the short duration of the task (6x40s), combined with the fact that we did not

couple reinforcement to monetary incentives, a manipulation known to strongly boost striatal

activity in the context of motor learning[18]. Yet, when considering BOLD activity in the striatal ROIs,

we did find a significant effect of reinforcement, suggesting that our experimental manipulation did

increase striatal activity but that the strength of the effect was insufficient to survive at the whole-

brain level. Second, we did not find any effect of tTIS when considering averaged BOLD activity.

Again, the short duration of the blocks may contribute to this non-significant effect. Another

possible interpretation is that the effect of tTIS on BOLD activity is not uniform across participants

as it likely depends on individual anatomy and function of the targeted brain region, as observed

for other non-invasive brain stimulation techniques[110]. Consistently, we found a correlation

between levels of impulsivity and the neural effects of $tTIS_{80Hz}$ (both BOLD and connectivity, Figure

S7). Importantly though, when including learning as a behavioral regressor we did find significant

clusters of correlation specifically in the striatum (Figure 3), suggesting that the behavioural effects

were indeed related to modulation of activity in the target region. Notably, this result was significant

when contrasting $tTIS_{80Hz}$ to the active control ($tTIS_{20Hz}$), but not to $tTIS_{Sham}$. Overall, we believe

that the fMRI data does provide interesting support that the behavioural effects of the stimulation

were indeed related to modulation of neural activity in the striatum, also in line with the present

simulations on realistic head models (Figure 1) and the connectivity results (Figure 4). This idea

is also in agreement with another recent study investigating the effects of tTIS on motor sequence

learning[52]. Notably though, a limitation of the present dataset is the very short duration of

stimulation and imaging for each experimental condition, that may explain some inconsistencies

in the results. Hence, following this first proof-of-concept study showing robust behavioural effects

and related neural changes, future studies including longer fMRI and stimulation sessions are

required to further confirm these results.

Finally, within the present study the computational modelling was performed on a realistic,

detailed head model (i.e., the MIDA model[59], see Methods). One limitation of this approach is that

the electric field simulations do not take individual structural information into account. Such

individual modeling would require information on brain anisotropy, an aspect that is likely to

significantly influence tTIS exposure[44,111]. However, in the present study diffusion MRI to evaluate

fractional anisotropy was not acquired. Future studies including diffusion MRI data will allow for

personalised modelling, paving the way for individualised tTIS informed by brain structure[53].

Conclusion

The present findings show for the first time the ability of non-invasive striatal tTIS to

interfere with reinforcement learning in humans through a selective modulation of striatal activity

and support the causal functional role of the human striatum in reinforcement motor learning. This

deep brain stimulation was well tolerated and compatible with efficient blinding, suggesting that

tTIS provides the exciting option to circumvent the steep depth-focality trade-off of current non-

invasive brain stimulation approaches in a safe and effective way. Overall, tTIS opens new

possibilities for the study of causal brain-behaviour relationships and for the treatment of neuro-

psychiatric disorders associated to alterations of deep brain structures.

4. Methods

4.1. Participants

A total of 48 right-handed healthy volunteers participated in the study. 24 participants were enrolled for the main tTIS study (15 women, 25.3 ± 0.7 years old; mean $\pm$ SE). Another group of 24 volunteers participated in the behavioural control experiment (Figure S2, 14 women, 24.2 ± 0.5 years old). Handedness was determined via a shortened version of the Edinburgh Handedness inventory[112] (laterality index = $89.3 \pm 2.14\%$ for the main study and $86.4 \pm 2.51\%$ for the control experiment). None of the participants suffered from any neurological or psychiatric disorder, nor taking any centrally-acting medication (see Supplementary Materials for a complete list of exclusion criteria). All participants gave their written informed consent in accordance with the Declaration of Helsinki and the Cantonal Ethics Committee Vaud, Switzerland (project number 2020-00127). Finally, all participants were asked to fill out a delay-discounting monetary choice questionnaire[113], which evaluates the propensity of subjects to choose smaller sooner rewards over larger later rewards, a preference commonly associated to choice impulsivity[75,114].

4.2. Experimental procedures

The study employed a randomised, double-blind, sham-controlled design. Following screening and inclusion, participants were invited to a single experimental session including performance of a motor learning task with concurrent transcranial electric Temporal Interference stimulation (tTIS) of the striatum and functional magnetic resonance imaging (fMRI). Overall, participants practiced 6 blocks of trials, that resulted from the combination of two reinforcement feedback conditions (Reinf$_{TYPE}$: Reinf$_{ON}$ or Reinf$_{OFF}$) with three types of striatal stimulation (tTIS$_{TYPE}$: tTIS$_{Sham}$, tTIS$_{20Hz}$ or tTIS$_{80Hz}$).

4.2.1. Motor learning task

4.2.1.1. General aspects

Participants practiced an adaptation of a widely used force-tracking motor task[54,55] with a fMRI-compatible fiber optic grip force sensor (Current designs, Inc., Philadelphia, PA, USA) positioned in their right hand. The task was developed on Matlab 2018 (the Mathworks, Natick, Massachusetts, USA) exploiting the Psychophysics Toolbox extensions[115,116] and was displayed on a computer screen with a refresh rate of 60 Hz. The task required participants to squeeze the force sensor to control a cursor displayed on the screen. Increasing the exerted force resulted in the cursor moving vertically and upward in a linear way. Each trial started with a preparatory period in which a sidebar appeared at the bottom of the screen (**Figure 1A**). After a variable time interval (0.9 to 1.1 s), a cursor (black circle) popped up in the sidebar and simultaneously a target (grey larger circle with a cross in the middle) appeared, indicating the start of the movement period. Subjects were asked to modulate the force applied on the transducer to keep the cursor as close as possible to the center of the target. The target moved in a sequential way along a single vertical axis for 7 s. The maximum force required (i.e., the force required to reach the target when it was in the uppermost part of the screen; $MaxTarget_{Force}$) was set at 4% of maximum voluntary contraction (MVC) evaluated at the beginning of the experiment. This low force level was chosen based on pilot experiments to limit muscular fatigue. Finally, each trial ended with a blank screen displayed for 2 s before the beginning of the next trial.

4.2.1.2. Trial types and reinforcement manipulation

During the experiment, participants were exposed to different types of trials (**Figure 1A,**

Video S1). In Test trials, the cursor remained on the screen and the target was consistently

displayed in grey for the whole duration of the trial. These trials served to evaluate Pre- and Post-

training performance for each block, without any disturbance. In $Reinf_{ON}$ and $Reinf_{OFF}$ trials (used

during Training only), we provided only partial visual feedback to the participants in order to

increase the impact of reinforcement on learning[4,56–58]. As such, the cursor was only intermittently

displayed during the trial: it was always displayed in the first second of the trial, and then

disappeared for a total of 4.5 s randomly split on the remaining time by bits of 0.5 s. The cursor

was therefore displayed 35.7% of the time during these trials (2.5 s over the 7 s trial). Importantly,

contrary to the cursor, the target always remained on the screen for the whole trial and participants

were instructed to continue to track the target even when the cursor was away.

In addition to this visual manipulation, in $Reinf_{ON}$ trials, participants also trained with

reinforcement feedback indicating success or failure of the tracking in real time. As such,

participants were informed that, during these trials, the color of the target would vary as a function

of their performance: the target was displayed in green when tracking was considered as

successful and in red when it was considered as failure. Online success on the task was

determined based on the Error, defined as the absolute force difference between the force

required to be in the center of the target and the exerted force[4,54–56]. The Error, expressed in

percentage of MVC, was computed for each frame refresh and allowed to classify a sample as

successful or not based on a closed-loop reinforcement schedule[8]. More specifically, for each

training trial, a force sample (recorded at 60 Hz, corresponding to the refresh rate of the monitor)

was considered as successful if the computed Error was below the median Error over the 4

previous trials at this specific sample. Put differently, to be successful, participants had to

constantly beat their previous performance. This closed-loop reinforcement schedule allowed us

to deliver consistent reinforcement feedback across individuals and conditions (see control

analysis on success rates in the Supplementary materials), while maximizing uncertainty on the

presence of reinforcement, an aspect that is crucial for efficient reinforcement motor learning[117].

Notably, in addition to this closed-loop design, samples were also considered as successful if the

cursor was very close to the center of the target (i.e., within one radius around the center,

corresponding to an Error below 0.2% of MVC). This was done to prevent any conflict between

visual information (provided by the position of the cursor relative to the target) and reinforcement

feedback (provided by the color of the target), which could occur in situations of extremely good

performance (when the closed-loop Error cut-off is below 0.2% of MVC).

As a control, Reinf$_{OFF}$ trials were similar to Reinf$_{ON}$ trials with the only difference that the

displayed colors were either cyan or magenta, and were generated randomly. Participants were

explicitly told that, in this condition, colors were displayed randomly and could be ignored. The

visual properties of the target in the Reinf$_{OFF}$ condition were designed to match the Reinf$_{ON}$

condition in terms of relative luminance (cyan: RGB = [127.5 242.1 255] matched to green: [127.5

255 127.5] and magenta: [211.7 127.5 255] to red: [255 127.5 127.5]) and average frequency of

change in colors (i.e., the average number of changes in colors divided by the total duration of a

trial, see Supplementary materials).

Notably, in this task, Training trials differed from Test trials regarding not only the color of

the target (red/green or cyan/magenta in Training trials and grey in Test trials) but also the visual

feedback experienced (partial and full visual feedback in Training and Test trials, respectively).

This choice was motivated by several reasons. First, we wanted to evaluate learning in the

classical, unperturbed, version of the force-tracking task[54,55], which is compatible with clinical

translation. Second, based on additional behavioural data on another group of participants (n =

24, see Figure S2), we found that significant effects of reinforcement on learning were observed

only when training was performed with partial visual feedback (displayed on 35.7% of the trial

time, as in the present study), in line with previous results[57,62]. However, this additional study also

revealed very limited improvement of performance during training with partial visual feedback,

potentially due to ceiling effects on performance in this condition. Yet, the improvement of

performance when comparing the Pre and Post-training assessments strongly suggested that

practicing the task with partial visual feedback still induced significant learning of the skill. Finally,

the change in visual feedback between Training and Post-training was the same in all

experimental conditions; this aspect of the task is therefore unlikely to explain the reinforcement

as well as the stimulation effects reported here.

Even though our study focused on reinforcement motor learning, it is worth mentioning that

other learning mechanisms such as error-based or strategic processes are likely to be also

engaged during the force-tracking task and may have recruited other brain regions beyond the

striatum (Spampinato and Celnik, 2020). Notably though, our protocol was specifically designed

to compare in the same individuals, learning in Reinf$_{ON}$ and Reinf$_{OFF}$ conditions while keeping the

other parameters of the task constant, to specifically isolate the contribution of reinforcement

processes in motor learning.

4.2.1.3. Motor learning protocol

After receiving standardised instructions about the force-tracking task, participants

practiced 5 blocks of familiarization (total of 75 trials) without tTIS. The first block of familiarization

included 20 trials with the target moving in a regular fashion (0.5 Hz sinuoid). Then, in a second

block of familiarization, participants performed 35 trials of practice with an irregular pattern, with

the same properties as the training patterns (see below). Finally, we introduced the reinforcement

manipulation and let participants perform 2 short blocks (8 trials each) including Reinf$_{ON}$ and

Reinf$_{OFF}$ trials. These four first blocks of familiarization were performed outside the MRI

environment. A last familiarization block (4 trials) was performed after installation in the scanner,

to allow participants to get used to performing the task in the MRI. This long familiarization allowed

participants to get acquainted with the use of the force sensor, before the beginning of the

experiment.

During the main part of the experiment, participants performed 6 blocks of trials in the MRI

with concurrent striatal tTIS (**Figure 1B**). Each block was composed of 4 Pre-training trials

followed by 24 Training and 8 Post-training trials. Pre- and Post-training trials were performed in

Test conditions, without tTIS and were used to evaluate motor learning. Training trials were

performed with or without reinforcement feedback and with concomitant striatal tTIS and were

used as a proxy of motor performance. During Training, trials were interspersed with 25 s resting

periods every 4 trials (used for fMRI contrasts, see below). The order of the 6 experimental

conditions was pseudo-randomised across participants: the 6 blocks were divided into 3 pairs of

blocks with the same tTIS condition and each pair was then composed of one $Reinf_{ON}$ and one

$Reinf_{OFF}$ block. Within this structure, the order of the $tTIS_{TYPE}$ and $Reinf_{TYPE}$ conditions were

balanced among the 24 participants. Hence, this randomisation allowed us to ensure that any

order effect that may arise from the repetition of the learning blocks would have the same impact

on each experimental condition (e.g., 4 subjects experienced $tTIS_{80Hz}$ - $Reinf_{ON}$ in the first block, 4

other subjects in the second block, 4 in the third block etc.).

As mentioned above, the protocol involved multiple evaluations of motor learning within

the same experimental session. In order to limit carry-over effects from one block to the following,

each experimental block was associated to a different pattern of movement of the target (**Figure**

S1). Put differently, in each block, participants had to generate a new pattern of force to

successfully track the target. To balance the patterns' difficulty, they all consisted in the summation

of 5 sinusoids of variable frequency (range: 0.1-1.5 Hz) that presented the following properties: a)

Average force comprised between 45 and 55% of the $MaxTarget_{Force}$; b) Absolute average

derivative comprised between 54 and 66 % of the $MaxTarget_{Force}$/s; c) Number of peaks = 14

(defined as an absolute change of force of at least 1% of MaxTarget$_{Force}$). These parameters were

determined based on pilot experiments to obtain a relevant level of difficulty for young healthy

adults and consistent learning across the different patterns.

4.2.2. Transcranial Electric Temporal Interference Stimulation (tTIS) applied to the
striatum

4.2.2.1. General concept

Transcranial temporal interference stimulation (tTIS) is an innovative non-invasive brain

stimulation approach, in which two or more independent stimulation channels deliver high-

frequency currents in the kHz range (oscillating at f1 and f1 + Δf; **Figure 1C**). These high-

frequency currents are assumed to be too high to effectively modulate neuronal activity [41,50,118].

Still, by applying a small shift in frequency, they result in a modulated electric field with the

envelope oscillating at the low-frequency Δf (target frequency) where the two currents overlap.

The peak of the modulated envelope amplitude can be steered towards specific areas located

deep in the brain, by tuning the position of the electrodes and current ratio across stimulation

channels[41] (**Figure 1C, 1D**). Based on these properties, tTIS has been shown to be able to focally

target activity of deep structures in rodents, without engaging overlying tissues[41]. Here, we applied

temporal interference stimulation transcranially via surface electrodes applying a low-intensity,

sub-threshold protocol following the currently accepted cut-offs and safety guidelines for low-

intensity transcranial electric stimulation in humans[119].

4.2.2.2. Stimulators

The currents for tTIS were delivered by two independent DS5 isolated bipolar constant

current stimulators *(Digitimer Ltd, Welwyn Garden City, UK)*. The stimulation patterns were

generated using a custom-based Matlab graphical user interface and transmitted to the current

sources using a standard digital-analog converter *(DAQ USB-6216, National Instruments, Austin,*

TX, USA). Finally, an audio transformer was added between stimulators and subjects, in order to

avoid possible direct current accumulation.

4.2.2.3. Stimulation protocols

During the 6 Training blocks, we applied three different types of striatal tTIS (2 blocks

each): a stimulation with a tTIS envelope modulated at 20Hz ($tTIS_{20Hz}$), a stimulation with a tTIS

envelope modulated at 80Hz ($tTIS_{80Hz}$) and a sham stimulation ($tTIS_{Sham}$). For $tTIS_{20Hz}$, the

posterior stimulation channel (TP7-TP8, see below) delivered a 1.99 kHz stimulation while the

anterior one delivered a 2.01 kHz ($\Delta f = 20$ Hz). For $tTIS_{80Hz}$, the posterior and anterior channels

delivered 1.96 kHz and 2.04 kHz, respectively ($\Delta f = 80$ Hz). Hence in both conditions, the high

frequency component was comparable and the only difference was Δf. During each block, tTIS

was applied for 5 minutes (6 x 50 s) during Training. Each stimulation period started and ended

with currents ramping-up and -down, respectively, for 5 s. tTIS was applied only while participants

were performing the motor task and not during resting periods or Pre- and Post-training

assessments. Finally, $tTIS_{Sham}$ consisted in a ramping-up (5 s) immediately followed by a ramping-

down (5 s) of 2 kHz currents delivered without any shift in frequency. This condition allowed us to

mimic the sensations experienced during the active conditions $tTIS_{20Hz}$ and $tTIS_{80Hz}$, while

delivering minimal brain stimulation (Figure S5). A trigger was sent 5 seconds before the beginning

of each trial in order to align the beginning of the task and the beginning of the frequency shift

after the ramp-up. Other tTIS parameters were set as follows: current intensity per stimulation

channel = 2 mA (baseline-to-peak), electrode type: round, conductive rubber with conductive

cream/paste, electrode size = 3 cm^2 (see ContES checklist in Supplementary materials for more

details).

The stimulation was applied within the MRI environment (Siemens 3T MAGNETOM

Prisma; Siemens Healthcare, Erlangen, Germany) using a standard RF filter module and MRI-

compatible cables *(neuroConn GmbH, Ilmenau, Germany)*. The technological, safety and noise

tests, and methodological factors can be found in Supplementary materials (Table S1) and are

based on the ContES Checklist [120].

4.2.2.4. Modelling

Electromagnetic simulations were carried out to identify optimised electrode placement

and current steering parameters. Simulations were performed using the MIDA head model[59], a

detailed anatomical head model featuring >100 distinguished tissues and regions that was derived

from multi-modal image data of a healthy female volunteer. Importantly, for brain stimulation

modelling, the model differentiates different scalp layers, skull layers, grey and white matter,

cerebrospinal fluid, and the dura and accounts for electrical conductivity anisotropy and neural

orientation based on diffusion tensor imaging (DTI) data. Circular electrodes (radius = 0.7 cm)

were positioned on the skin according to the 10-10 system and the electromagnetic exposure was

computed using the ohmic-current-dominated electro-quasistatic solver from Sim4Life v5.0 (ZMT

Zurich MedTech AG, Switzerland), which is suitable due to the dominance of ohmic currents over

displacement currents and the long wavelength compared with the simulation domain[121]. Dielectric

properties were assigned based on the IT'IS Tissue Properties Database v4.0[122]. Rectilinear

discretization was performed, and grid convergence as well as solver convergence analyses were

used to ensure negligible numerical uncertainty, resulting in a grid that included more than 54M

voxels. Dirichlet voltage boundary conditions, and then current normalization were applied. The

electrode-head interface contact was treated as ideal. tTIS exposure was quantified according to

the maximum modulation envelope magnitude formula from Grossman et al., (2017)[41]. Then, a

sweep over 960 permutations of the four electrode positions was performed, considering

symmetric and asymmetric montages with parallel (sagittal and coronal) or crossing current paths,

while quantifying bilateral striatum (putamen [BNA regions 225, 226, 229, 230], caudate [BNA

regions 219, 220, 227, 228] and nucleus accumbens [BNA regions 223, 224]) exposure

performance according to three metrics: a) target exposure strength, b) focality ratio (the ratio of

target tissue volume above threshold compared to the whole-brain tissue volume above threshold,

a measure of stimulation selectivity), and c) activation ratio (percentage of target volume above

threshold with respect to the total target volume, a measure of target coverage). We defined the

threshold as the 98th volumetric iso-percentile level of the tTIS. From the resulting Pareto-optimal

front, two configurations stood out particularly: one that maximised focality and activation (AF3 -

AF4, P7 - P8) and a second one that accepts a reduction of these two metrics by a quarter, while

increasing the target exposure strength by more than 50% (F3-F4, TP7-TP8). This last montage

was selected, to ensure sufficient tTIS exposure in the striatum[52] (Figure 1C, 1D).

865   4.2.2.5. Electrode positioning and evaluation of stimulation-associated sensations
866

Based on the modelling approach described above, we defined the stimulation electrode

positions in the framework of the EEG 10-10 system[123]. The optimal montage leading in terms of

target (i.e. the bilateral striatum) exposure strength and selectivity, was composed of the following

electrodes: F3, F4, TP7 and TP8. Their locations were marked with a pen on the scalp and, after

skin preparation (cleaned with alcohol), round conductive rubber electrodes of 3 cm^2 were placed

adding a conductive paste (*Ten20, Weaver and Company, Aurora, CO, USA or Abralyt HiCl,*

Easycap GmbH, Woerthsee-Etterschlag, Germany) as an interface to the skin. Electrodes were

held in position with tape and cables were oriented towards the top in order to allow good

positioning inside the scanner. Impedances were checked and optimised until they were below 20

kΩ [48]. Once good contact was obtained, we tested different intensities of stimulation for each

stimulation protocol in order to familiarise the participants with the perceived sensations and to

systematically document them. $tTIS_{Sham}$, $tTIS_{20Hz}$ and $tTIS_{80Hz}$ were applied for 20 seconds with the

following increasing current amplitudes per channel: 0.5 mA, 1 mA, 1.5 mA and 2 mA. Participants

were asked to report any kind of sensation and, if a sensation was felt, they were asked to grade

the intensity from 1 to 3 (light to strong) as well as give at least one adjective to describe it (Figure

S5). Following this step, cables were removed to be replaced by MRI-compatible cables and a

bandage was added to apply pressure on the electrodes and keep them in place. An impedance

check was repeated in the MRI right before the training and then again at the end of all recordings.

4.2.3. MRI data acquisition

Structural and functional images were acquired using a 3T MAGNETOM PRISMA scanner

(*Siemens, Erlangen, Germany*). T1-weighted images were acquired via the 3D MPRAGE

sequence with the following parameters: TR = 2.3 s; TE = 2.96 ms; flip angle = 9°; slices = 192;

voxel size = 1 × 1 × 1 mm, FOV = 256 mm. Anatomical T2 images were also acquired with the

following parameters: TR = 3 s; TE = 409 ms; flip angle = 120°; slices = 208; voxel size = 0.8 ×

0.8 × 0.8 mm, FOV = 320 mm. Finally, functional images were recorded using Echo-Planar

Imaging (EPI) sequences with the following parameters: TR = 1.25 s; TE = 32 ms; flip angle =

58°; slices = 75; voxel size = 2 × 2 × 2 mm; FOV = 112 mm.

4.3.   Data and statistical analyses

Data and statistical analyses were carried out with Matlab 2018a (the Mathworks, Natick,

Massachusetts, USA) and the R software environment for statistical computing and graphics (R

Core Team 2021, Vienna, Austria). Robust linear regressions were fitted with the Matlab function

robustfit. Linear mixed models (LMM) were fitted using the lmer function of the lme4 package in R

[124]. As random effects, we added intercepts for participants and block. Normality of residuals, and

homoscedasticity of the data were systematically checked, and logarithmic transformations were

applied when necessary (i.e., when skewness of the residuals' distribution was not comprised

between - 2 and 2^{125} or when homoscedasticity was violated based on visual inspection). To

mitigate the impact of isolated influential data points on the outcome of the final model, we used

tools of the influence.ME package to detect and remove influential cases based on the following

criterion: distance > 4 * mean distance[126]. Statistical significance was determined using the anova

function with Satterthwaite's approximations of the lmerTest package[127]. For specific post-hoc

comparisons we conducted pairwise comparisons by computing estimated marginal means with

the emmeans package with Tukey adjustment of p-values[128]. Standardised effect size measures

were obtained using the eff_size function of the emmeans package[129]. The level of significance

was set at p<0.05.

4.3.1. Behavioural data

4.3.1.1. Evaluation of motor learning

The main goal of the present study was to evaluate the influence of striatal tTIS on

reinforcement motor learning. To do so, we first removed trials, in which participants did not react

within 1 s after the appearance of the cursor and target, considering that these extremely long

preparation times may reflect significant fluctuations in attention[130]. This occurred extremely rarely

(0.52 % of the whole data set). For each subject and each trial, we then quantified the tracking

Error as the absolute force difference between the applied and required force as done

previously[4,54,56]. Tracking performance during Training and Post-training trials were then

normalised according to subjects' initial level by expressing the Error data in percentage of the

average Pre-training Error for each block. In order to test our main hypothesis predicting specific

effects of striatal tTIS on reinforcement motor learning, we performed a LMM on the Post-training

data with $tTIS_{TYPE}$ and $Reinf_{TYPE}$ as fixed effects. We then also ran the same analysis on the

Training data, to evaluate if striatal tTIS also impacted on motor performance, while stimulation

was being delivered.

As a control, we checked that initial performance at Pre-training was not different between

conditions with a LMM on the Error data obtained at Pre-training. Again, $tTIS_{TYPE}$ and $Reinf_{TYPE}$

were considered as fixed effects. Finally, another LMM was fitted with the fixed effect $tTIS_{TYPE}$ to

verify that the amount of positive reinforcement (as indicated by a green target) in the $Reinf_{ON}$

blocks was similar across $tTIS_{TYPES}$.

939
940   4.3.2. fMRI data
941
942   4.3.2.1. Imaging Preprocessing
943

We analyzed functional imaging data using Statistical Parametric Mapping 12 (*SPM12;*

The Wellcome Department of Cognitive Neurology, London, UK) implemented in MATLAB

R2018a (*Mathworks, Sherborn, MA*). All functional images underwent a common preprocessing

including the following steps: slice time correction, spatial realignment to the first image,

normalization to the standard MNI space and smoothing with a 6 mm full-width half-maximal

Gaussian kernel. T1 anatomical images were then co-registered to the mean functional image and

segmented. This allowed to obtain bias-corrected gray and white matter images, by normalizing

the functional images via the forward deformation field. To select subjects with acceptable level of

head movement, framewise displacement was calculated for each run. A visual check of both non-

normalised and normalised images was performed in order to ensure good preprocessing quality.

Finally, possible tTIS-related artifacts were investigated based on signal to noise ratio maps (see

below).

957   4.3.2.2. Signal to Noise Ratio

Total signal to noise ratio (tSNR) maps were computed to check the presence of possible

artifacts induced by the electrical stimulation. The values were calculated per each voxel by

dividing the mean of the voxel time series by its standard deviation. Spherical regions of interest

were then defined both underneath the tTIS electrodes and at 4 different locations, distant from

the electrodes as a control. The center of each spherical ROI was obtained by projecting the

standard MNI coordinates of each electrode on the scalp[131] toward the center of the brain. After

visual inspection of the ROIs, average tSNR maps were extracted within each sphere. A LMM was

used to compare the average SNR underneath the electrodes versus the control regions and

between stimulation protocols. The results of this analysis are presented in Supplementary

materials (Figure S8).

970   4.3.2.3. Task-based BOLD activity analysis

A general linear model was implemented at the single-subject level in order to estimate

signal amplitude. Eight regressors were included in the model: 6 head motion parameters

(displacement and rotation) and normalised time series within the white matter and the

corticospinal fluid. Linear contrasts were then computed to estimate specific activity during the

motor task with respect to resting periods. Functional activation was also extracted within specific

ROIs individually defined based on structural images. More specifically, the Freesurfer recon-all

function   was   run   based   on   the   structural   T1w   and   T2w   images

(https://surfer.nmr.mgh.harvard.edu/). The BNA parcellation was derived on the individual subject

space and the selected ROIs were then co-registered to the functional images and normalised to

the MNI space. BOLD activity within the individual striatal masks was averaged and compared

between different striatal nuclei namely the putamen (BNA regions 225, 226, 229, 230), caudate

(BNA regions 219, 220, 227, 228) and nucleus accumbens (BNA regions 223, 224). Comparison

between conditions were presented for uncorrected voxel-wise FWE, p=0.001 and multiple

comparison corrected at the cluster level to reduce False Discovery Rate (FDR), p=0.05.

988   4.3.2.4. Effective connectivity analyses

As an additional investigation, we computed task-modulated effective functional

connectivity by means of the CONN toolbox 2021a (www.nitrc.org/projects/conn,

RRID:SCR_009550) running in Matlab R2018a (*Mathworks, Sherborn, MA*). An additional

denoising step was added by applying a band-pass filtering from 0.01 to 0.1 Hz and by regressing

potential confounders (white matter, CSF and realignment parameters). After that, generalized

Psycho-Physiological Interactions (gPPI) connectivity was extracted within specific pre-defined

customised sub-networks: a reward and a motor network. gPPI evaluates condition-specific

changes in effective connectivity, defined as the directed effect that one brain region has on

another under some model of neuronal coupling (Friston, 1994). In particular, gPPI considers a

series of equations in which activity in a ROI (pre-defined frontal areas in our case) depends on a

specific condition (the 'psychological' factor) and on activity in the seed region (striatum here, the

'physiological' factor). By solving these equations, it is possible to determine a coefficient that

represents task-modulation of effective connectivity[132]. Importantly, task-related changes in

effective connectivity are expressed relative to rest, and therefore values closer to 0 reflect a

connectivity similar to resting state.

1005

The reward network was defined as following: two regions within the striatum, namely the

NAc (BNA regions 223 and 224) and the ventro-medial putamen (BNA regions 225 and 226, left

and right respectively), and two frontal areas, namely the anterior cingulate (BNA regions 177,

179, 183 and 178, 180, 184, left and right respectively) and the orbitofrontal cortex within the

vmPFC (BNA regions 41, 45, 47, 49, 187 and 42, 46, 48, 50, 188 for left and right respectively).

The motor network included the following areas: the dorso-lateral putamen (BNA 229, 230, for left

and right respectively), the dorsal caudate (BNA regions 227, 228 for left and right respectively)

the medial part of the SMA (BNA regions 9 and 10, left and right respectively) and the part of the

M1 associated to upper limb function (BNA regions 57 and 58, left and right respectively). Notably,

we considered connectivity in the left and right motor and reward networks regardless of laterality.

These ROIs were selected based on the following rationale. First, they are consistent with previous

literature on reinforcement learning of motor skills[68,90,133,134]. Second, there is structural and

functional evidence for these fronto-striatal connections[135,136]. Third, the frontal areas included in

the analyses are well-established hubs of the motor learning (M1 and SMA, see [12] for a meta-

analysis) and reward networks (vmPFC and ACC, see [11] for a meta-analysis). Finally, gPPI was

also extracted within a control language network, defined based on the functional atlas described

by Shirer et al.(2012)[70].

Supplementary material

1024

1025   **1.** **Exclusion criteria**

1026

● Unable to consent
● Severe neuropsychiatric (e.g., major depression, severe dementia) or unstable systemic
diseases (e.g., severe progressive and unstable cancer, life threatening infectious
diseases)
● Severe sensory or cognitive impairment or musculoskeletal dysfunctions prohibiting to
understand instructions or to perform the experimental tasks
● Color blindness
● Inability to follow or non-compliance with the procedures of the study
● Contraindications for NIBS or MRI:
○ Electronic or ferromagnetic medical implants/device, non-MRI compatible metal
1037   implant
1038   ○ History of seizures
1039   ○ Medication that significantly interacts with NIBS being benzodiazepines, tricyclic
1040   antidepressant and antipsychotics
1041   ● Regular use of narcotic drugs
1042   ● Left-handedness
1043   ● Pregnancy
1044   ● Request of not being informed in case of incidental findings
1045   ● Concomitant participation in another trial involving probing of neuronal plasticity.
1046

1047   **2. <u>ContES Checklist</u>**

1048

Technological factors	
Manufacturer of Stimulator	DS5 Isolated Bipolar Constant Current Stimulator (Digitimer)
MR Conditional Electrode Details	Round, 3 cm2 conductive rubber electrodes
Electrode Positioning	F3 → F4 TP7 → TP8 A bandage is warped around the head to apply pressure and keep the electrodes in place Electrodes are oriented in order to have vertical cables entering parallel to the MRI coil Head was fixed with pillows to avoid movements
MR Conditional Skin-Electrode Interface	10-20 gel One or two drops of saline were added when impedances were too high
Amount of Contact Medium (Paste/Gel/Electrolyte)	Around 1mm of paste was manually placed on the electrodes
Electrode Placement Visualization	Pictures

RF Filter	NeuroConn DC-STIMULATOR MR RF filter module with MRI-compatible cables and electrodes
Wire Routing Pattern	10 m ethernet cables between inner and outer box pass through a conduit along the wall of the MRI room until reaching the back of the MRI. Cables are then fixed with straps on the ground and on the wall of the MRI machine in order to avoid loops until reaching the interior of the coil. Cables between the head and the inner boxes were also fixed with straps and they were oriented in order to exit the magnetic field direction as soon as possible as indicated by the red arrows of the image below.
tES-fMRI Machine Synchronization/Communication	Stimulation was triggered by the stimulus delivery PC via parallel port to BNC cable. The parallel port of the stimulus delivery PC was connected to the DAQ controlling the stimulators. Stimulus delivery PC, in turn, was also receiving the scanner trigger from the scanner via USB port.
Safety and noise tests	
MR Conditionality Specifics for tES Setting	Please refer to Section "Methods-Imaging acquisition"
tES-fMRI Setting Test - Safety Testing	Impedances were checked before and after

	the stimulation.
	No temperature tests were performed during the experiment.
	Intensity titration was performed prior to entering the MRI, testing increasing currents (0.5, 1, 1.5 and 2 mA) and asking the subject to report any type of sensation.
	A sensation questionnaire was also performed at the end of the experiment.
tES-fMRI Setting Test - Subjective Intolerance Reporting	No intolerances were reported by any subject
tES-fMRI Setting Test - Noise/Artifact	Signal to Noise Ratio (SNR) analysis was performed on the fMRI images, please refer to Section "Methods-Signal to Noise Ratio"
Impedance Testing	Impedances were checked right after electrodes positioning outside the scanner, before and after the stimulation inside.
	One or two drops of saline solution were added if impedances were higher than 20kΩ
Methodological factors	
Concurrent tES-fMRI Timing	For timings, please refer to the "Methods-Stimulation protocols" section
	To mitigate the impact of potential carry-over effects on our experimental results we used the following strategy:
	1) We stimulated for short periods in each condition (5 minutes interspersed with resting periods without stimulation; see "Methods-Stimulation protocols");
	2) We imposed breaks (~7-8 minutes) between each stimulation protocol;
	3) We randomised the order of the

	Stimulation conditions
Imaging Session Timing	All sequences were performed with TI stimulation electrodes placed on the subjects' head.
tES Experience Report	Please refer to "Results" section and to Figure S5.

1049 **Table S1. ContES checklist as recommended in Ekhtiari et al., 2022[120] for concurrent tES-fMRI**
1050 **studies.**

1051

1052 ### 3. Patterns of motion of the target used in the study

1053

1054   **Figure S1. Patterns of motion of the target.** For each block of training, participants had
1055 to learn a new pattern of motion of the target. The patterns had similar mathematical properties
1056 and their relationship to a condition was randomised (see Methods for more details).

1057

1058 ### 4. Additional behavioural experiment

1059
1060   To determine the optimal experimental parameters to study reinforcement learning of

1061 motor skills, we performed an additional behavioural experiment, in the absence of brain

1062 stimulation and imaging. In particular, we tested the relationship between the amount of visual

1063 feedback available during Training and the benefits of reinforcement in the force-tracking task.

1064 Another group of young healthy participants (n=24; 14 women, 24.2 ± 0.5 years old, independent

from the subjects tested in the main experiment) performed blocks of the task with $Reinf_{ON}$ or $Reinf_{OFF}$ and with either full visual feedback or only partial visual feedback (cursor displayed for 35.7% of the total trial duration, as in the main study). Each learning block was composed of 30 trials (vs. 36 trials in the main study) and in addition to real-time closed-loop reinforcement feedback, participants also received endpoint feedback on their overall performance after each trial during Training (i.e., indicating success or failure on the trial). The LMM ran on the Post-training data revealed a significant effect of visual feedback ($F_{(1,788.33)}$=5.90; p=0.015), reinforcement ($F_{(1,787.87)}$=11.64; p<0.001) and a significant interaction between these two factors ($F_{(1,788.03)}$=10.27; p=0.0014, **Figure S2A**). Interestingly, Tukey-corrected post-hoc tests showed that the interaction was due to the fact that while reinforcement did not improve learning when Training was performed with full visual feedback (p=0.88, d=0.014), it induced robust benefits when training with partial visual feedback (p<0.001, d=0.46, **Figure S2B**). This result is in line with previous literature showing that reinforcement feedback is particularly beneficial for motor learning when visual feedback is uncertain[57,62]. Based on the outcome of this additional study, we decided to train participants with partial visual feedback in the present experiment to evaluate the effect of tTIS in a version of the task that yielded significant reinforcement gains. Notably, this work also shows that the effect of reinforcement on motor learning observed in the $tTIS_{Sham}$ and $tTIS_{20Hz}$ conditions (Figure 2) is reproducible.

1083

Figure S2. Results of an additional behavioural experiment (n = 24). A) Motor performance across training. Raw Error data (expressed in % of Maximum Voluntary Contraction [MVC]) are presented on the left panel for the different experimental conditions in bins of 3 trials. On the right, the two plots represent the Pre-training normalised Error in the Full visual feedback and Partial visual feedback blocks (i.e., cursor displayed 35.7% of the time, as in the main experiment). Note the strong gains in motor performance, especially with partial visual feedback but also the limited improvement of performance during training in this condition. B) Motor learning. Averaged Error at Post-training (normalised to Pre-training) in the different experimental conditions are shown, for the subjects included in the analysis (i.e., after outlier detection, n=23). Reduction of Error at Post-training reflects true improvement at tracking the target in Test conditions (in the absence of reinforcement or visual uncertainty). The LMM ran on these data revealed significant a significant effect of reinforcement feedback on learning when training with partial, but not full, visual feedback. *: p<0.05. Data are represented as mean ± SE.

5. Evolution of motor performance in the different conditions

The main analysis revealed a general effect of tTIS on motor performance during Training, irrespective of the presence of reinforcement. As a subsequent analysis, we also asked whether the evolution of performance during Training depended on type of striatal stimulation applied. We ran the same LMM as in the main study (see Results) but with the addition of a continuous fixed effect Trial, allowing us to evaluate whether the slope of performance change was different according to $tTIS_{TYPE}$. (**Figure S3**). Indeed, this analysis revealed a significant $tTIS_{TYPE}$ x Trial interaction ($F_{(2, 3399)}$=4.46; p=0.012) that was due to different slopes in the $tTIS_{Sham}$ compared to the $tTIS_{20Hz}$ (p=0.013) and $tTIS_{80Hz}$ (at the trend level, p=0.068) conditions. Evolution of performance in the $tTIS_{20Hz}$ and $tTIS_{80Hz}$ conditions was not different (p=0.81). Notably, this effect

could not be explained by differences in initial performance (all p>0.21 when comparing

intercepts). Moreover, the effect of tTIS on motor improvement during Training did also not depend

on the presence of reinforcement (Reinf$_{TYPE}$, tTIS$_{TYPE}$ and Trial: $F_{(2,\,3399)}$=0.51; p=0.60). Overall,

this analysis shows that the detrimental effect of striatal tTIS on motor performance is due to an

impaired ability to improve performance with practice and further confirms that tTIS did not

modulate the ability to use reinforcement feedback during Training.

1115

1116 **Figure S3. Slopes of performance change during Training in the different stimulation**
1117 **conditions.** Modeled performance change for tTIS$_{Sham}$, tTIS$_{20Hz}$ and tTIS$_{80Hz}$ throughout Training.
1118 The tTIS$_{TYPE}$ x Trial interaction revealed that performance improved more with tTIS$_{Sham}$ compared
1119 to tTIS$_{20Hz}$ and tTIS$_{80Hz}$. Notably this effect was not modulated by the presence of reinforcement
1120 and could not be explained by differences in intercepts.

1121

1122

1123 **6.** <u>**Effect of visual and reinforcement feedback on motor performance**</u>

1124

As a control, we asked whether the tTIS and reinforcement effects reported in Figure 2

depended on the availability of visual information during Training. To do so, we computed the

normalised Error for phases with the Cursor$_{ON}$ or Cursor$_{OFF}$ (taking into account a lag of 0.25s,

corresponding to the estimated visuo-motor delay in this type of task for young healthy subjects[137])

and analysed these data in a LMM including the factors Reinf$_{TYPE}$, tTIS$_{TYPE}$ and Cursor$_{TYPE}$. As in

the main analysis, we confirmed the effect of Reinf$_{TYPE}$ ($F_{(1,\,6872)}$=344.87; p<0.001), tTIS$_{TYPE}$

($F_{(2,6872)}$=28.79; p<0.001) and the absence of interaction between these two factors ($F_{(2,6875.4)}$=0.49;

p=0.61, Figure S4). This analysis also revealed a Cursor$_{TYPE}$ effect ($F_{(2,6875.3)}$=49.66; p<0.001)

which was due to the fact that the Error was generally higher in the absence visual information on

the position of the cursor (d=0.17). Interestingly, there was also a Reinf$_{TYPE}$ x Cursor$_{TYPE}$ interaction

($F_{(2,6872)}$=29.35; p<0.001): while benefits of reinforcement were significant in both the Cursor$_{ON}$

(p<0.001, d=0.32) and Cursor$_{OFF}$ (p<0.001, d=0.58) conditions, the magnitude of the

reinforcement-related gains in performance were larger in the Cursor$_{OFF}$ condition (t-test

comparing the gains: $t_{(46)}$=2.74, p=0.0086). Moreover, post-hoc tests also revealed that the

absence of vision of the cursor was detrimental for performance in the Reinf$_{OFF}$ condition (p<0.001,

d=0.30) but not in presence of Reinf$_{ON}$ (p=0.25, d=0.039). Hence, the presence of reinforcement

was particularly beneficial when visual information was not available, in line with previous

research[57,62] and also in agreement with the results of our additional experiment (Figure S2).

Importantly, the LMM did not reveal any interaction between tTIS$_{TYPE}$ and Cursor$_{TYPE}$ ($F_{(2,6872)}$=0.49;

p=0.31) and no triple interaction ($F_{(2,6872)}$=1.53; p=0.22), confirming that striatal tTIS had a global

effect on motor performance during Training, which did not depend on the presence of visual and

reinforcement feedback.

1147

1148

1149 **Figure S4. Effect of visual and reinforcement feedback on motor performance.** Pre-
1150 training normalised Error depending on the presence of the cursor (Cursor$_{ON}$ or Cursor$_{OFF}$), and

the presence of reinforcement feedback during Training. The significant $Reinf_{TYPE}$ x $Cursor_{TYPE}$ interaction was related to the fact that the benefits of reinforcement were stronger when visual information was not available. Notably, this analysis takes into account a visuo-motor delay of 0.25s, as previously reported during a similar task (Lam and Zenon, 2021). *: p<0.05. Data are represented as mean ± SE.

7. Control analyses of behavioural data

Pre-training performance

In order to verify that our main behavioural results were not influenced by potential differences in initial performance between conditions despite randomisation, we analysed the Error at Pre-training between conditions. We did not find any $tTIS_{TYPE}$ ($F_{(2,519.15)}$=1.64; p=0.20) or $tTIS_{TYPE}$ x $Reinf_{TYPE}$ effect ($F_{(2,519.99)}$=1.08; p=0.34), suggesting that the main behavioural results could not be accounted for by differences in initial performance between conditions. However, the LMM did reveal a $Reinf_{TYPE}$ effect ($F_{(1,519.15)}$=12.47; p<0.001), that was due to the fact that Pre-training performance was generally better in $Reinf_{OFF}$ blocks. This effect, which was opposite to our learning results (generally better learning with $Reinf_{ON}$), may be related to an expectancy effect stemming from the repetitive structure of the reinforcement conditions (see Methods). However, the absence of interaction with $tTIS_{TYPE}$ is strongly suggestive that this effect did not drive any of the main findings. Put together, these data provide confidence that the differential effects of striatal tTIS on motor learning depending on the presence of reinforcement were not the result of different initial performance between conditions.

Success rate

Overall, the amount of positive reinforcement (i.e., when the target was green) averaged 52.78 +/- 0.42% and was comparable across $tTIS_{TYPES}$ ($F_{(2,1702)}$=0.17; p=0.84), suggesting that the closed-loop reinforcement schedule was successful at providing similar reinforcement feedback

despite differences in performance between conditions. Hence, different success rates during

training cannot explain the effect of the different striatal tTIS conditions on motor learning.

Frequency of flashing

Analysis of the frequency of flashing in the different conditions did not reveal any effect of

$tTIS_{TYPE}$ ($F_{(2,3283)}$=0.85; p=0.43) nor any $Reinf_{TYPE}$ x $tTIS_{TYPE}$ interaction ($F_{(2,3283)}$=0.19; p=0.82),

suggesting that the behavioural effects of tTIS could not be explained by a visual confound.

However, this analysis did reveal a $Reinf_{TYPE}$ effect ($F_{(1,3283)}$=33.62; p<0.001) which was due to the

fact that the average frequency in the $Reinf_{OFF}$ condition (4.28 ± 0.097 Hz) was slightly but

significantly higher than with $Reinf_{ON}$ (4.08 ± 0.098 Hz; $F_{(1,3283)}$=33.62; p<0.001). Notably, in

absolute terms, this difference represented only a difference of 1.4 change of color over the whole

7 s trial, which we think is unlikely to explain the improvement of performance in the $Reinf_{ON}$

condition.

Order of the reinforcement conditions

Previous exposure to reinforcement feedback may improve subsequent learning through

reinforcement[138]. Thanks to our randomisation procedure, the previous exposure to the $Reinf_{ON}$

condition was equally counterbalanced in all stimulation conditions, and should therefore not

influence our main results. Still, we performed an analysis to specifically investigate the effect of

the previous exposure to $Reinf_{ON}$. To do so, we split the participants depending on whether they

experienced $Reinf_{ON}$ or $Reinf_{OFF}$ first (12 subjects per group) and performed a new LMM on the

Post-training data with the addition of a categorical factor $Group_{TYPE}$. In particular, if the previous

exposure to the $Reinf_{ON}$ condition influenced following learning with reinforcement, we would

expect to see a $Group_{TYPE}$ x $Reinf_{TYPE}$ interaction. The analysis did not indicate any $Group_{TYPE}$

effect on learning ($F_{(1,21.96)}$=0.35; p=0.56), neither did it reveal a $Group_{TYPE}$ x $Reinf_{TYPE}$ ($F_{(1,4)}$=0.72;

p=0.44), or a triple interaction with $tTIS_{TYPE}$ ($F_{(2,1105.06)}$=1.75; p=0.17). Overall, this analysis

suggests that the order of the exposure to the reinforcement condition did not influence the present

findings.

1204

1205   **8. <u>Blinding integrity and tTIS-evoked sensations</u>**

1206

1207   **Figure S5. tTIS-related sensations. A) Magnitude of tTIS-related sensations.**
1208 Magnitude of sensations reported before the experiment for current amplitudes ranging from 0.5
1209 to 2 mA for each tTIS$_{TYPE}$. The current amplitude used in the present experiment was 2 mA. **B)**
1210 **Types of tTIS-related sensations.** Type of sensations as described by the participants, at 2 mA.
1211 Note that subjects were allowed to describe their sensations with up to two different words.
1212

1213 **9. Brain activity during reinforcement motor learning**

1214

Figure S6. Whole-brain activity during reinforcement motor learning. Activation maps
for the contrast task>rest in the tTIS$_{Sham}$, Reinf$_{ON}$ condition showing activation of key areas of the
reinforcement motor learning network including the putamen, thalamus, cerebellum and
sensorimotor network, especially on the left side. Significant clusters are shown for corrected
voxel-wise family wise error (FWE), p=0.05, and corrected cluster-based false discovery rate
(FDR), p=0.05.

1221
1222

Cluster-level				Peak-level					x	y	z	Region
$p_{FWE-corr}$	$q_{FDR-corr}$	k_E	P_{uncorr}	$p_{FWE-corr}$	$q_{FDR-corr}$	T	(Z_E)	P_{uncorr}				
<0.001	<0.001	135	<0.001	<0.001	0.005	12.63	6.84	<0.001	46	-62	4	Temporal_Mid_R
<0.001	<0.001	523	<0.001	<0.001	0.005	12.32	6.77	<0.001	-40	-8	62	Precentral_L
				<0.001	0.021	10.62	6.33	<0.001	-34	-6	52	Postcentral_L
				<0.001	0.021	10.43	6.28	<0.001	-36	-20	54	Precentral_L
<0.001	<0.001	335	<0.001	<0.001	0.018	11.08	6.46	<0.001	-8	-6	64	Supp_Motor_Area_L
				0.003	0.145	8.21	5.56	<0.001	6	6	58	Supp_Motor_Area_R
				0.003	0.145	8.20	5.55	<0.001	-4	-2	54	Supp_Motor_Area_L
<0.001	<0.001	44	<0.001	<0.001	0.021	10.65	6.34	<0.001	-10	-20	6	Thal_IL_L
<0.001	<0.001	162	<0.001	<0.001	0.021	10.36	6.26	<0.001	42	-6	56	Frontal_Mid_2_R
				<0.001	0.042	9.48	5.99	<0.001	34	-4	58	Frontal_Sup_2_R
<0.001	<0.001	175	<0.001	<0.001	0.021	10.27	6.23	<0.001	-58	10	28	Precentral_L
				<0.001	0.037	9.60	6.03	<0.001	-56	8	20	Frontal_Inf_Oper_L
				0.019	0.490	7.32	5.21	<0.001	-48	2	16	Rolandic_Oper_L
<0.001	<0.001	601	<0.001	<0.001	0.024	10.06	6.17	<0.001	2	-74	-34	Vermis_7
				<0.001	0.025	9.99	6.15	<0.001	-12	-70	-22	Cerebellum_6_L
				<0.001	0.027	9.88	6.12	<0.001	12	-70	-20	Cerebellum_6_R
<0.001	<0.001	82	<0.001	<0.001	0.070	9.14	5.88	<0.001	56	10	26	Frontal_Inf_Oper_R
				0.006	0.234	7.86	5.42	<0.001	56	10	38	Precentral_R
<0.001	<0.001	141	<0.001	0.001	0.092	8.89	5.80	<0.001	-34	-52	-24	Cerebellum_6_L
				0.002	0.117	8.47	5.65	<0.001	-28	-62	-24	Cerebellum_6_L
<0.001	<0.001	76	<0.001	0.001	0.092	8.87	5.79	<0.001	-28	-52	56	Parietal_Sup_L
				0.011	0.341	7.57	5.31	<0.001	-30	-44	48	Parietal_Inf_L
<0.001	<0.001	200	<0.001	0.001	0.092	8.77	5.76	<0.001	32	-48	-28	Cerebellum_6_R
				0.013	0.382	7.49	5.28	<0.001	34	-40	-34	Cerebellum_6_R
<0.001	<0.001	36	<0.001	0.001	0.092	8.73	5.74	<0.001	16	-54	-18	Cerebellum_4_5_R
<0.001	<0.001	28	<0.001	0.001	0.101	8.63	5.71	<0.001	26	-58	-54	Cerebellum_8_R
<0.001	<0.001	62	<0.001	0.001	0.113	8.51	5.67	<0.001	38	-62	-16	Fusiform_R
				0.002	0.117	8.45	5.64	<0.001	42	-72	-12	Occipital_Inf_R
<0.001	<0.001	21	<0.001	0.002	0.117	8.41	5.63	<0.001	-46	-68	4	Occipital_Mid_L
<0.001	<0.001	141	<0.001	0.002	0.130	8.33	5.60	<0.001	22	-56	50	Location not in atlas
				0.002	0.130	8.30	5.59	<0.001	30	-48	48	Parietal_Sup_R
				0.007	0.266	7.76	5.39	<0.001	36	-40	42	SupraMarginal_R
<0.001	<0.001	29	<0.001	0.004	0.170	8.09	5.51	<0.001	44	-50	-34	Cerebellum_Crus_1_R

<0.001	<0.001	59	<0.001	0.004	0.178	8.04	5.49	<0.001	-22	-66	-52	Cerebellum_8_L
<0.001	0.006	12	0.003	0.004	0.190	7.99	5.47	<0.001	10	-16	8	Thal_MDl_R
0.001	0.043	6	0.028	0.009	0.319	7.63	5.33	<0.001	-22	-2	6	Putamen_L
<0.001	<0.001	34	<0.001	0.009	0.319	7.63	5.33	<0.001	18	-64	-54	Cerebellum_8_R
0.001	0.300	7	0.019	0.023	0.545	7.23	5.17	<0.001	20	2	62	Frontal_Sup_2_R
0.001	0.030	7	0.019	0.024	0.560	7.21	5.16	<0.001	52	12	8	Frontal_Inf_Oper_R
0.001	0.030	7	0.019	0.025	0.568	7.19	5.16	<0.001	-44	-36	40	Parietal_Inf_L

Table S2: Significant clusters and the respective local maxima in the tTIS$_{Sham}$, Reinf$_{ON}$ condition. Related to Figure S6. Regions were identified with the Automated Anatomical Labelling atlas 3 (AAL3[139]). Significant clusters were selected for corrected voxel-wise family wise error (FWE), p=0.05, and corrected cluster-based false discovery rate (FDR), p=0.05.

10. Correlation between effect of tTIS$_{80Hz}$ on reinforcement motor learning and modulation of whole-brain activity

Cluster-level				Peak-level					x	y	z	Region
p$_{FWE-corr}$	q$_{FDR-corr}$	k$_E$	P$_{uncorr}$	p$_{FWE-corr}$	q$_{FDR-corr}$	T	(Z$_E$)	P$_{uncorr}$				
0.003	**0.005**	**157**	**<0.001**	**0.027**	**0.065**	**7.29**	**5.14**	**<0.001**	**10**	**18**	**0**	Caudate_R
				0.639	0.678	5.38	4.25	<0.001	0	0	10	Location not in atlas
				0.921	0.757	4.89	3.98	<0.001	6	6	2	Location not in atlas
0.007	**0.005**	**138**	**<0.001**	**0.693**	**0.678**	**5.30**	**4.21**	**<0.001**	**-16**	**14**	**6**	Location not in atlas
				0.923	0.757	4.88	3.98	<0.001	-22	14	-2	Putamen_L
				1.000	0.810	4.26	3.60	<0.001	-18	8	-6	Putamen_L

Table S3. Significant clusters for the correlation between the behavioural and neural effects of tTIS$_{80Hz}$ (vs. tTIS$_{20Hz}$). Related to Figure 3B. Two significant clusters were found with several local maxima. Notably, the left cluster also encompassed a portion of the left caudate (related to Figure 3). Regions were identified with the Automated Anatomical Labelling atlas 3 (AAL3[139]). Significant clusters were selected for uncorrected voxel-wise family wise error (FWE), p=0.001, and corrected cluster-based false discovery rate (FDR), p=0.05.

11. Control analysis on striatum to frontal cortex effective connectivity

The connectivity analysis showed that tTIS$_{80Hz}$, but not tTIS$_{20Hz}$, increased striatum to frontal effective connectivity and that this effect depended on the type of network considered

(reward vs. motor) and on the presence of reinforcement (Figure 4). In this analysis we considered

effective connectivity between the motor striatum and M1 and SMA for the motor network and the

limbic striatum with ACC and vmPFC for the reward network, based on a large body of

literature[11,12,135,136] (see Methods for a detailed justification of the ROIs). To verify whether our

results depended on the specific frontal ROIs included in the analysis, we performed a new

analysis. More specifically, we decomposed connectivity in each network for each frontal cortical

area (M1 and SMA in the motor network and ACC and vmPFC in the reward network) and ran two

separate LMMs on each network with $tTIS_{TYPE}$, $Reinf_{TYPE}$ as well as ROI_{TYPE} (M1 or SMA for the

LMM run on the motor network and ACC or vmPFC for the reward network) as fixed effects.

Consistent with our initial findings, we found effects of $tTIS_{TYPE}$ on both LMMs (motor network:

$F_{(2,1089.7)}=3.12$; $p=0.044$ and reward network: $F_{(2,1112)}=6.78$; $p=0.0012$). Moreover, there was a

significant $tTIS_{TYPE}$ x $Reinf_{TYPE}$ interaction in the motor network ($F_{(2,1112)}=3.36$; $p=0.035$), which was

at the trend level in the reward network ($F_{(2,1113.8)}=2.37$; $p=0.094$). Most importantly, these effects

were not modulated by ROI_{TYPE} in any network ($tTIS_{TYPE}$ x $Reinf_{TYPE}$ x ROI_{TYPE} in motor network:

$F_{(2,1112)}=0.83$; $p=0.44$, in reward network: $F_{(2,1112)}=0.61$; $p=0.54$). This analysis suggests that the

main connectivity findings were not influenced by the specific frontal ROIs considered in the

analysis.

1261

1262

1263 **12. <u>Relationship between the neural and behavioural effects of tTIS$_{80Hz}$ and impulsivity</u>**

1264

Characterising individual factors that influence responsiveness to brain stimulation is an

important line of research both for fundamental neuroscience but also to determine profiles of

responders for future clinical translation. Based on previous literature linking striatal gamma

oscillatory mechanisms and impulsivity[72], we explored the possibility that impulsivity influences

responsiveness to striatal tTIS$_{80Hz}$ (**Figure S7**).

First, we exploited the BOLD data and asked if inter-individual variability in the neural

effects of tTIS$_{80Hz}$ during reinforcement motor learning (i.e., in the Reinf$_{ON}$ condition) was related

to impulsivity at the whole-brain level. Impulsivity was evaluated by a well-established independent

delay-discounting questionnaire performed at the beginning of the experiment[75,76]. Strikingly, this

analysis revealed that impulsivity was associated to the effect of tTIS$_{80Hz}$ (with respect to tTIS$_{20Hz}$)

specifically in the left caudate nucleus (Figure S7A, Table S4). No other clusters were found. As

such, the most impulsive participants exhibited an increase of left caudate activity with tTIS$_{80Hz}$

(compared to tTIS$_{20Hz}$) while the least impulsive ones rather presented a decrease of BOLD signal,

consistent with the idea that impulsivity modulates the neuronal responsiveness to tTIS (R^2=0.47;

p<0.001; Figure S7B). No significant clusters of correlation were found for the tTIS$_{80Hz}$ − tTIS$_{Sham}$

contrast, neither for the control tTIS$_{20Hz}$ - tTIS$_{Sham}$ contrast. Hence, this analysis suggests that the

effect of tTIS$_{80Hz}$ on caudate activity depends on participants' impulsivity.

As a second step, we aimed at evaluating the association between impulsivity and the

increased striatum to motor cortex connectivity observed with tTIS$_{80Hz}$, in the presence of

reinforcement. Notably, such pattern of increased connectivity in fronto-striatal circuits has been

described as a pathophysiological mechanism in multiple neuro-psychiatric disorders involving

impulsivity[101−104]. Hence, we first asked if striatum to motor cortex connectivity was related to

impulsivity during reinforcement motor learning in the absence of stimulation (i.e., in the tTIS$_{Sham}$

condition). Indeed, we found a significant positive relationship between impulsivity and striatum to

motor cortex connectivity (robust linear regression: R^2=0.10; p=0.0038), in line with previous

results[101−104]. Then, we evaluated whether the increase of connectivity observed with tTIS$_{80Hz}$ in

the Reinf$_{ON}$ condition (Figure 4A) could be related to impulsivity. Indeed, we found that the effect

of tTIS$_{80Hz}$ on connectivity was negatively correlated to impulsivity both when contrasting tTIS$_{80Hz}$

with tTIS$_{Sham}$ (R^2=0.19; p=0.043, Figure S7C, left) and with tTIS$_{20Hz}$ (R^2=0.28; p=0.021, Figure S7C,

middle): participants with the largest increase in connectivity with tTIS$_{80Hz}$ in the Reinf$_{ON}$ condition

were also the least impulsive ones. Such correlation was absent when contrasting $tTIS_{20Hz}$ and

$tTIS_{Sham}$ (R^2=0.0031; p=0.31, Figure S7C, right), but also when considering the same contrasts in

the reward instead of the motor network (p=0.93 and p=0.86 for the $tTIS_{80Hz}$-$tTIS_{Sham}$ and $tTIS_{80Hz}$-

$tTIS_{20Hz}$ contrasts, respectively). Hence, striatum to motor cortex effective connectivity during the

task was positively correlated to impulsivity, but the change in connectivity induced by $tTIS_{80Hz}$

was rather negatively associated with impulsivity. This may be due to a ceiling effect in the most

impulsive participants: exhibiting initially high levels of connectivity may leave less room for further

modulation by $tTIS_{80Hz}$. These results suggest that inter-individual variability in impulsivity might

influence neural responses to striatal $tTIS_{80Hz}$.

Figure S7. Relationship between impulsivity and the neural effects of $tTIS_{80Hz}$. A) Whole-brain correlation between the neural effects of $tTIS_{80Hz}$ (with respect to $tTIS_{20Hz}$) and impulsivity. Correlation between tTIS-related modulation of striatal activity ($tTIS_{80Hz} - tTIS_{20Hz}$) during reinforcement motor learning ($Reinf_{ON}$) and individual impulsivity levels. A single significant cluster of correlation was found in left caudate (uncorrected voxel-wise FWE: p=0.001, and

corrected cluster-based FDR: p=0.05). **B) Correlation between left caudate activity and impulsivity.** A positive correlation was found showing that participants with higher levels of impulsivity exhibited stronger activation of the left caudate in the $tTIS_{80Hz}$ (with respect to $tTIS_{20Hz}$). **C) Correlations between impulsivity and tTIS-related modulation of effective connectivity.** Impulsivity was associated to the neural effects of $tTIS_{80Hz}$ both when contrasting to $tTIS_{Sham}$ (left) and $tTIS_{20Hz}$ (middle), but was not correlated to the effect of $tTIS_{20Hz}$ (right).

Cluster-level				Peak-level					x	y	z	Region
$p_{FWE-corr}$	$q_{FDR-corr}$	k_E	P_{uncorr}	$p_{FWE-corr}$	$q_{FDR-corr}$	T	(Z_E)	P_{uncorr}				
<0.001	**<0.001**	**254**	**<0.001**	**0.707**	**0.524**	**5.29**	**4.20**	**<0.001**	**-8**	**0**	**18**	Location not in atlas
				0.719	0.524	5.27	4.19	<0.001	-14	16	16	Caudate_L
				0.971	0.620	4.72	3.88	<0.001	-16	16	0	Location not in atlas

Table S4. Significant clusters for the correlation between impulsivity and effects of $tTIS_{80Hz}$ on BOLD activity (vs. $tTIS_{20Hz}$). Related to Figure S7A. One significant cluster encompassing the left caudate nucleus was found. Regions were identified with AAL3[139].

As a last step, we verified if impulsivity was also predictive of the behavioural effects of $tTIS_{80Hz}$ on reinforcement motor learning. We did not find any significant correlation between impulsivity and the effect of $tTIS_{80Hz}$ on motor learning ($tTIS_{80Hz} - tTIS_{Sham}$: R^2=0.098; p=0.17; $tTIS_{80Hz} - tTIS_{20Hz}$: R^2=0.11; p=0.21). Hence, impulsivity was associated to the neural, but not the behavioural effects of $tTIS_{80Hz}$.

Overall, we found that impulsivity was associated to $tTIS_{80Hz}$-related BOLD changes specifically in the left caudate and to changes of effective connectivity between the motor striatum and motor cortex during reinforcement motor learning. Hence, a possibility is that the differences in endogenous striatal gamma-related activity that have been associated to impulsive behaviour in animal models[72–74], influence the neural effects of $tTIS_{80Hz}$. If this is the case, impulsivity could constitute a behavioural factor allowing to determine responsiveness to striatal $tTIS_{80Hz}$.

1333   Conversely, an interesting avenue for future research could aim at determining whether impulsivity

1334   can be modulated by striatal $tTIS_{80Hz}$.

1335

1336

1337   **13. Imaging quality control**

1338

A threshold of 0.5 was chosen to discard subjects showing more than 40% of voxels with

framewise displacement FD higher than this threshold. In the current study cohort, no subject

exceeded the limit value, thus the whole dataset could be used. Furthermore, successful cleaning

of the data was ensured by visual checking the preprocessing results. In particular, good

registration between anatomical and functional images and normalization to standard space were

checked. Signal to noise ratio analysis showed significantly higher tSNR values underneath the

stimulating electrodes ($F_{(1,1122)}=249.25$, $p<0.001$; **Figure S5**). Moreover, an additional analysis

showed that this effect was not influenced by the $tTIS_{TYPE}$ ($Sphere_{LOCATION}$ x $tTIS_{TYPE}$:

$F_{(2,1118)}=0.0169$, $p=0.98$). This result suggests that the stimulation did not introduce additional

noise to the MR images. In summary, all controls confirmed the good quality of the imaging data.

1349

Figure S8. Total signal to noise ratio (tSNR). Total signal to noise ratio investigation. On
the top panel, the average tSNR is shown within spheres of 10mm radius underneath the 4
stimulation electrodes (F3, F4, TP7 and TP8) and underneath other 4 locations more distal from
the electrodes (C3, C4, O1 and O2). A significant higher tSNR was found underneath the
electrodes with respect to the distal locations ($F_{(1,1122)}$=249.25, p<0.001). This indicates that there
was no reduction of the tSNR due to the presence of electrical current. On the bottom panel, the
location of the spheres from where the average tSNRs were extracted: F3 and F4 in red in the
first image from the left, TP7 and TP8 in red on the second image from the left, C3 and C4 in blue
on the third image from the left, O1 and O2 in blue on the forth image from the left.

Acknowledgements

P.V. was a PhD student supported by the Fund for Research training in Industry and Agriculture (FRIA/FNRS; FC29690), and grants by the Platform for Education and Talent (Gustave Boël - Sofina Fellowships) and Wallonie-Bruxelles International. J.D. was supported by grants from the Belgian FNRS and the Fondation Médicale Reine Elisabeth (FMRE). G.D. was supported by the Belgian FNRS. The research was partially funded by the Novartis Research Foundation—FreeNovation (Basel, CH) to MJW and EN, the Bertarelli Foundation (Catalyst 'Deep-MCI-T', Gstaad, CH) to FCH, the SNSF Lead Agency (NiBS-iCog 320030L_197899) to FCH and the Defitech Foundation (Morges, CH) to FCH. We acknowledge access to and expertise of the Neuromodulation and the Neuroimaging facilities of the Human Neuroscience Platform of the Campus Biotech Geneva and of the MRI Platform of the HVS (Sion). We would also like to thank Prof. Leonardo Cohen for insightful feedback on a previous version of this manuscript.

Competing interests

E.N. is co-founder of TI Solutions AG, a company committed to producing hardware and software solutions to support tTIS research.

Data and code availability statement

The datasets generated during the current study and the code used to analyse them are available from the corresponding author on reasonable request.

References

1. Neftci, E. O. & Averbeck, B. B. Reinforcement learning in artificial and biological systems. *Nat. Mach. Intell.* **1**, 133–143 (2019).

2. Schultz, W. Neuronal Reward and Decision Signals: From Theories to Data. *Physiol. Rev.* **95**, 853–951 (2015).

3. Dhawale, A. K., Smith, M. A. & Ölveczky, B. P. The Role of Variability in Motor Learning. *Annu. Rev. Neurosci.* **40**, 479–498 (2017).

4. Vassiliadis, P. *et al.* Reward boosts reinforcement-based motor learning. *iScience* **24**, 102821 (2021).

5. Spampinato, D. & Celnik, P. Multiple Motor Learning Processes in Humans: Defining Their Neurophysiological Bases. *Neuroscientist* **27**, 246–267 (2021).

6. Huang, V. S., Haith, A., Mazzoni, P. & Krakauer, J. W. Rethinking Motor Learning and Savings in Adaptation Paradigms: Model-Free Memory for Successful Actions Combines with Internal Models. *Neuron* **70**, 787–801 (2011).

7. Galea, J. M., Mallia, E., Rothwell, J. & Diedrichsen, J. The dissociable effects of punishment and reward on motor learning. *Nat. Neurosci.* **18**, 597–602 (2015).

8. Therrien, A. S., Wolpert, D. M. & Bastian, A. J. Effective Reinforcement learning following cerebellar damage requires a balance between exploration and motor noise. *Brain* **139**, 101–114 (2016).

9. Vassiliadis, P., Derosiere, G. & Duque, J. Beyond Motor Noise : Considering Other Causes of Impaired Reinforcement Learning in Cerebellar Patients. *Eneuro* **6**, 1–4 (2019).

10.   Widmer, M. *et al.* Reward During Arm Training Improves Impairment and Activity

After Stroke: A Randomized Controlled Trial. *Neurorehabil. Neural Repair* **0**,

154596832110628 (2021).

11.   Bartra, O., McGuire, J. T. & Kable, J. W. The valuation system: A coordinate-

based meta-analysis of BOLD fMRI experiments examining neural correlates of

subjective value. *Neuroimage* **76**, 412–427 (2013).

12.   Hardwick, R. M., Rottschy, C., Miall, R. C. & Eickhoff, S. B. A quantitative meta-

analysis and review of motor learning in the human brain. *Neuroimage* **67**, 283–

297 (2013).

13.   Haber, S. N. Corticostriatal circuitry. *Dialogues Clin. Neurosci.* **18**, 7–21 (2016).

14.   Balleine, B. W., Delgado, M. R. & Hikosaka, O. The role of the dorsal striatum in

reward and decision-making. *J. Neurosci.* **27**, 8161–8165 (2007).

15.   Piray, P., den Ouden, H. E. M., van der Schaaf, M. E., Toni, I. & Cools, R.

Dopaminergic modulation of the functional ventrodorsal architecture of the human

striatum. *Cereb. Cortex* **27**, 485–495 (2017).

16.   Hori, Y. *et al.* Ventral striatum links motivational and motor networks during

operant-conditioned movement in rats. *Neuroimage* **184**, 943–953 (2019).

17.   Wachter, T., Lungu, O. V., Liu, T., Willingham, D. T. & Ashe, J. Differential Effect

of Reward and Punishment on Procedural Learning. *J. Neurosci.* **29**, 436–443

(2009).

18.   Widmer, M., Ziegler, N., Held, J., Luft, A. & Lutz, K. Rewarding feedback promotes

motor skill consolidation via striatal activity. in *Progress in Brain Research* **5**, 303–

323 (2016).

19.   Berke, J. D. Fast oscillations in cortical-striatal networks switch frequency

following rewarding events and stimulant drugs. *Eur. J. Neurosci.* **30**, 848–859

(2009).

20.   van der Meer, M. A. A. *et al.* Integrating early results on ventral striatal gamma

oscillations in the rat. *Front. Neurosci.* **4**, 1–12 (2010).

21.   van der Meer, M. A. A. & Redish, A. D. Low and high gamma oscillations in rat

ventral striatum have distinct relationships to behavior, reward, and spiking activity

on a learned spatial decision task. *Front. Integr. Neurosci.* **3**, 1–19 (2009).

22.   Dwiel, L. L., Khokhar, J. Y., Connerney, M. A., Green, A. I. & Doucette, W. T.

Finding the balance between model complexity and performance: Using ventral

striatal oscillations to classify feeding behavior in rats. *PLoS Comput. Biol.* **15**, 1–

19 (2019).

23.   Matsumoto, J. *et al.* Neuronal responses in the nucleus accumbens shell during

sexual behavior in male rats. *J. Neurosci.* **32**, 1672–1686 (2012).

24.   Kalenscher, T., Lansink, C. S., Lankelma, J. V. & Pennartz, C. M. A. Reward-

associated gamma oscillations in ventral striatum are regionally differentiated and

modulate local firing activity. *J. Neurophysiol.* **103**, 1658–1672 (2010).

25.   Cohen, M. X. *et al.* Good vibrations: Cross-frequency coupling in the human

nucleus accumbens during reward processing. *J. Cogn. Neurosci.* **21**, 875–889

(2009).

26.   Sepe-Forrest, L., Carver, F. W., Quentin, R., Holroyd, T. & Nugent, A. C. Basal

ganglia activation localized in MEG using a reward task. *Neuroimage: Reports* **1**, 100034 (2021).

27. Herrojo-Ruiz, M. *et al.* Involvement of human internal globus pallidus in the early modulation of cortical error-related activity. *Cereb. Cortex* **24**, 1502–1517 (2014).

28. Jenkinson, N. & Brown, P. New insights into the relationship between dopamine, beta oscillations and motor function. *Trends Neurosci.* **34**, 611–618 (2011).

29. Brown, P. Abnormal oscillatory synchronisation in the motor system leads to impaired movement. *Curr. Opin. Neurobiol.* **17**, 656–664 (2007).

30. McCarthy, M. M. *et al.* Striatal origin of the pathologic beta oscillations in Parkinson's disease. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 11620–11625 (2011).

31. Kondabolu, K. *et al.* Striatal cholinergic interneurons generate beta and gamma oscillations in the corticostriatal circuit and produce motor deficits. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 3159–3168 (2016).

32. Silberstein, P. *et al.* Cortico-cortical coupling in Parkinson's disease and its modulation by therapy. *Brain* **128**, 1277–1291 (2005).

33. Williams, Z. M. & Eskandar, E. N. Selective enhancement of associative learning by microstimulation of the anterior caudate. *Nat. Neurosci.* **9**, 562–568 (2006).

34. Nakamura, K. & Hikosaka, O. Facilitation of saccadic eye movements by postsaccadic electrical stimulation in the primate caudate. *J. Neurosci.* **26**, 12885–12895 (2006).

35. Deng, Z. De, Lisanby, S. H. & Peterchev, A. V. Electric field depth-focality tradeoff in transcranial magnetic stimulation: Simulation comparison of 50 coil designs.

Brain Stimul. **6**, 1–13 (2013).

36.   Wagner, T. *et al.* Transcranial direct current stimulation: A computer-based human

model study. *Neuroimage* **35**, 1113–1124 (2007).

37.   Nickchen, K. *et al.* Reversal learning reveals cognitive deficits and altered

prediction error encoding in the ventral striatum in Huntington's disease. *Brain*

Imaging Behav. **11**, 1862–1872 (2017).

38.   Schmidt, L. *et al.* Disconnecting force from money: Effects of basal ganglia

damage on incentive motivation. *Brain* **131**, 1303–1310 (2008).

39.   Seymour, B. *et al.* Deep brain stimulation of the subthalamic nucleus modulates

sensitivity to decision outcome value in Parkinson's disease. *Sci. Rep.* **6**, 1–12

(2016).

40.   Atkinson-Clement, C. *et al.* Effects of subthalamic nucleus stimulation and

levodopa on decision-making in Parkinson's disease. *Mov. Disord.* **34**, 377–385

(2019).

41.   Grossman, N. *et al.* Noninvasive Deep Brain Stimulation via Temporally Interfering

Electric Fields. *Cell* **169**, 1029-1041.e16 (2017).

42.   Song, S., Zhang, J., Tian, Y., Wang, L. & Wei, P. Temporal Interference

Stimulation Regulates Eye Movements and Neural Activity in the Mice Superior

Colliculus. *Proc. Annu. Int. Conf. IEEE Eng. Med. Biol. Soc. EMBS* 6231–6234

(2021). doi:10.1109/EMBC46164.2021.9629968

43.   Esmaeilpour, Z., Kronberg, G., Reato, D., Parra, L. C. & Bikson, M. Temporal

interference stimulation targets deep brain regions by modulating neural

oscillations. *Brain Stimul.* **14**, 55–65 (2021).

44.   Rampersad, S. *et al.* Prospects for transcranial temporal interference stimulation

in humans: A computational study. *Neuroimage* **202**, 116124 (2019).

45.   von Conta, J. *et al.* Interindividual variability of electric fields during transcranial

temporal interference stimulation (tTIS). *Sci. Rep.* **11**, 1–12 (2021).

46.   Cao, J., Doiron, B., Goswami, C. & Grover, P. The mechanics of temporal

interference stimulation. *bioRxiv* 1–6 (2020). doi:10.1101/2020.04.23.051870

47.   Mirzakhalili, E., Barra, B., Capogrosso, M. & Lempka, S. F. Biophysics of

Temporal Interference Stimulation. *Cell Syst.* **11**, 557-572.e5 (2020).

48.   von Conta, J. *et al.* Benchmarking the effects of transcranial temporal interference

stimulation (tTIS) in humans. *Cortex* **154**, 299–310 (2022).

49.   Ma, R. *et al.* High Gamma and Beta Temporal Interference Stimulation in the

Human Motor Cortex Improves Motor Functions. *Front. Neurosci.* **15**, 1–11 (2022).

50.   Hutcheon, B. & Yarom, Y. Resonance, oscillation and the intrinsic frequency

preferences of neurons. *Trends Neurosci.* **23**, 216–222 (2000).

51.   Acerbo, E. *et al.* Focal non-invasive deep-brain stimulation with temporal

interference for the suppression of epileptic biomarkers. *Front. Neurosci.* **16**, 1–12

(2022).

52.   Wessel, M. J. *et al.* LTP-like noninvasive striatal brain stimulation enhances

striatal activity and motor skill learning in humans. *bioRxiv* 1–30 (2022).

doi:10.1101/2022.10.28.514204

53.   Violante, I. R. *et al.* Non-invasive temporal interference electrical stimulation of the

human hippocampus. *biorXiv* (2022). doi:10.1101/2022.09.14.507625

54. Abe, M. *et al.* Reward improves long-term retention of a motor memory through

induction of offline memory gains. *Curr. Biol.* **21**, 557–562 (2011).

55. Steel, A., Silson, E. H., Stagg, C. J. & Baker, C. I. The impact of reward and

punishment on skill learning depends on task demands. *Sci. Rep.* **6**, 1–9 (2016).

56. Vassiliadis, P., Lete, A., Duque, J. & Derosiere, G. Reward timing matters in motor

learning. *iScience* **25**, 104290 (2022).

57. Izawa, J. & Shadmehr, R. Learning from sensory and reward prediction errors

during motor adaptation. *PLoS Comput. Biol.* **7**, 1–12 (2011).

58. Mawase, F., Uehara, S., Bastian, A. J. & Celnik, P. Motor Learning Enhances

Use-Dependent Plasticity. *J. Neurosci.* **37**, 2673–2685 (2017).

59. Iacono, M. I. *et al.* MIDA: A multimodal imaging-based detailed anatomical model

of the human head and neck. *PLoS One* **10**, (2015).

60. Jiang, T. Brainnetome: A new -ome to understand the brain and its disorders.

Neuroimage **80**, 263–272 (2013).

61. Areshenkoff, C. N., de Brouwer, A. J., Gale, D. J., Nashed, J. Y. & Gallivan, J. P.

Separate and shared low-dimensional neural architectures for error-based and

reinforcement motor learning. *biorXiv* (2022). doi:10.1101/2022.08.16.504134

62. Cashaback, J. G. A., McGregor, H. R., Mohatarem, A. & Gribble, P. L.

Dissociating error-based and reinforcement-based loss functions during

sensorimotor learning. *PLoS Comput. Biol.* **13**, 1–28 (2017).

63. Floyer-Lea, A. & Matthews, P. M. Changing brain networks for visuomotor control

with increased movement automaticity. *J. Neurophysiol.* **92**, 2405–2412 (2004).

64.   Floyer-Lea, A. & Matthews, P. M. Distinguishable brain activation networks for

short- and long-term motor skill learning. *J. Neurophysiol.* **94**, 512–518 (2005).

65.   Graybiel, A. M. & Grafton, S. T. The striatum: Where skills and habits meet. *Cold*

Spring Harb. Perspect. Biol. **7**, 1–14 (2015).

66.   McLaren, D. G., Ries, M. L., Xu, G. & Johnson, S. C. A generalized form of

context-dependent psychophysiological interactions (gPPI): A comparison to

standard approaches. *Neuroimage* **61**, 1277–1286 (2012).

67.   Codol, O., Holland, P. J., Manohar, S. G. & Galea, J. M. Reward-Based

Improvements in Motor Control Are Driven by Multiple Error-Reducing

Mechanisms. *J. Neurosci.* **40**, 3604–3620 (2020).

68.   Sidarta, A., Vahdat, S., Bernardi, N. F. & Ostry, D. J. Somatic and reinforcement-

based plasticity in the initial stages of human motor learning. *J. Neurosci.* **36**,

11682–11692 (2016).

69.   Krause, M. R., Vieira, P. G., Csorba, B. A., Pilly, P. K. & Pack, C. C. Transcranial

alternating current stimulation entrains single-neuron activity in the primate brain.

Proc. Natl. Acad. Sci. U. S. A. **116**, 5747–5755 (2019).

70.   Shirer, W. R., Ryali, S., Rykhlevskaia, E., Menon, V. & Greicius, M. D. Decoding

subject-driven cognitive states with whole-brain connectivity patterns. *Cereb.*

Cortex **22**, 158–165 (2012).

71.   Morishita, T. & Hummel, F. C. Non-invasive Brain Stimulation (NIBS) in Motor

Recovery After Stroke: Concepts to Increase Efficacy. *Curr. Behav. Neurosci.*

Reports **4**, 280–289 (2017).

72.   Donnelly, N. A. *et al.* Oscillatory activity in the medial prefrontal cortex and nucleus

accumbens correlates with impulsivity and reward outcome. *PLoS One* **9**, 14–17

(2014).

73.   Schall, T. A., Wright, W. J. & Dong, Y. Nucleus accumbens fast-spiking

interneurons in motivational and addictive behaviors. *Mol. Psychiatry* **26**, 234–246

(2021).

74.   Pisansky, M. T. *et al.* Nucleus Accumbens Fast-Spiking Interneurons Constrain

Impulsive Action. *Biol. Psychiatry* **86**, 836–847 (2019).

75.   Kirby, K. N., Petry, N. M. & Bickel, W. K. Heroin addicts have higher discount rates

for delayed rewards than non-drug-using controls. *J. Exp. Psychol. Gen.* **128**, 78–

87 (1999).

76.   Mitchell, J. M., Fields, H. L., D'Esposito, M. & Boettiger, C. A. Impulsive

responding in alcoholics. *Alcohol. Clin. Exp. Res.* **29**, 2158–2169 (2005).

77.   Catanese, J., Carmichael, J. E. & van der Meer, M. A. A. Low- and high-gamma

oscillations deviate in opposite directions from zero-phase synchrony in the limbic

corticostriatal loop. *J. Neurophysiol.* **116**, 5–17 (2016).

78.   Rothé, M., Quilodran, R., Sallet, J. & Procyk, E. Coordination of high gamma

activity in anterior cingulate and lateral prefrontal cortical areas during adaptation.

J. Neurosci. **31**, 11110–11117 (2011).

79.   Del Arco, A., Park, J., Wood, J., Kim, Y. & Moghaddam, B. Adaptive encoding of

outcome prediction by prefrontal cortex ensembles supports behavioral flexibility.

J. Neurosci. **37**, 8363–8373 (2017).

80.   Yoshimoto, A., Shibata, Y., Kudara, M., Ikegaya, Y. & Matsumoto, N.

1580   Enhancement of Motor Cortical Gamma Oscillations and Sniffing Activity by

1581   Medial Forebrain Bundle Stimulation Precedes Locomotion. *eNeuro* **9**, 1–14

1582   (2022).

1583   81.   Grover, S., Nguyen, J. A., Viswanathan, V. & Reinhart, R. M. G. High-frequency

1584   neuromodulation improves obsessive–compulsive behavior. *Nat. Med.* **27**, 232–

1585   238 (2021).

1586   82.   Krause, M. R., Vieira, P. G., Thivierge, J. P. & Pack, C. C. Brain stimulation

1587   competes with ongoing oscillations for control of spike timing in the primate brain.

1588   *PLoS Biol.* **20**, 1–27 (2022).

1589   83.   Courtemanche, R., Fujii, N. & Graybiel, A. M. Synchronous, Focally Modulated β-

1590   Band Oscillations Characterize Local Field Potential Activity in the Striatum of

1591   Awake Behaving Monkeys. *J. Neurosci.* **23**, 11741–11752 (2003).

1592   84.   Costa, R. M. *et al.* Rapid Alterations in Corticostriatal Ensemble Coordination

1593   during Acute Dopamine-Dependent Motor Dysfunction. *Neuron* **52**, 359–369

1594   (2006).

1595   85.   Engel, A. K. & Fries, P. Beta-band oscillations-signalling the status quo? *Curr.*

1596   *Opin. Neurobiol.* **20**, 156–165 (2010).

1597   86.   Uehara, S., Mawase, F. & Celnik, P. Learning similar actions by reinforcement or

1598   sensory-prediction errors rely on distinct physiological mechanisms. *Cereb. Cortex*

1599   **28**, 3478–3490 (2018).

87.   Mathis, M. W., Mathis, A. & Uchida, N. Somatosensory Cortex Plays an Essential

Role in Forelimb Motor Adaptation in Mice. *Neuron* **93**, 1493-1503.e6 (2017).

88.   Brücke, C. *et al.* Scaling of movement is related to pallidal γ oscillations in patients

with dystonia. *J. Neurosci.* **32**, 1008–1019 (2012).

89.   Soderstrom, N. C. & Bjork, R. A. Learning Versus Performance: An Integrative

Review. *Perspect. Psychol. Sci.* **10**, 176–199 (2015).

90.   Spampinato, D. A., Satar, Z. & Rothwell, J. C. Combining reward and M1

transcranial direct current stimulation enhances the retention of newly learnt

sensorimotor mappings. *Brain Stimul.* 1–8 (2019). doi:10.1016/j.brs.2019.05.015

91.   Shmuelof, L. *et al.* Overcoming Motor 'Forgetting' Through Reinforcement Of

Learned Actions. *J. Neurosci.* **32**, 14617-14621a (2012).

92.   Dhawale, A. K., Miyamoto, Y. R., Smith, M. A. & Ölveczky, B. P. Adaptive

Regulation of Motor Variability. *Curr. Biol.* **29**, 3551-3562.e7 (2019).

93.   Carroll, T. J., McNamee, D., Ingram, J. N. & Wolpert, D. M. Rapid visuomotor

responses reflect value-based decisions. *J. Neurosci.* **39**, 3906–3920 (2019).

94.   De Comite, A., Crevecoeur, F. & Lefèvre, P. Reward-Dependent Selection of

Feedback Gains Impacts Rapid Motor Decisions. *Eneuro* **9**, ENEURO.0439-

21.2022 (2022).

95.   Codol, O. *et al.* Sensorimotor feedback loops are selectively sensitive to reward.

Elife **12**, 1–25 (2023).

96.   Krakauer, J. W., Hadjiosif, A. M., Xu, J., Wong, A. L. & Haith, A. M. Motor

Learning. *Compr. Physiol.* **9**, 613–663 (2019).

97. Quattrocchi, G. *et al.* Pharmacological Dopamine Manipulation Does Not Alter

Reward-Based Improvements in Memory Retention during a Visuomotor

Adaptation Task. *Eneuro* **5**, ENEURO.0453-17.2018 (2018).

98. Johnson, L. *et al.* Dose-dependent effects of transcranial alternating current

stimulation on spike timing in awake nonhuman primates. *Sci. Adv.* **6**, 1–9 (2020).

99. Beliaeva, V., Savvateev, I., Zerbi, V. & Polania, R. Toward integrative approaches

to study the causal role of neural oscillations via transcranial electrical stimulation.

Nat. Commun. **12**, 1–12 (2021).

100. Averbeck, B. & O'Doherty, J. P. Reinforcement-learning in fronto-striatal circuits.

Neuropsychopharmacology **47**, 147–162 (2022).

101. Ma, I. *et al.* Ventral striatal hyperconnectivity during rewarded interference control

in adolescents with ADHD. *Cortex* **82**, 225–236 (2016).

102. Wang, Q. *et al.* Dissociated neural substrates underlying impulsive choice and

impulsive action. *Neuroimage* **134**, 540–549 (2016).

103. Mosley, P. E. *et al.* The structural connectivity of discrete networks underlies

impulsivity and gambling in Parkinson's disease. *Brain* **142**, 3917–3935 (2019).

104. Hampton, W. H., Alm, K. H., Venkatraman, V., Nugiel, T. & Olson, I. R.

Dissociable frontostriatal white matter connectivity underlies reward and motor

impulsivity. *Neuroimage* **150**, 336–343 (2017).

105. Negahbani, E., Kasten, F. H., Herrmann, C. S. & Fröhlich, F. Targeting alpha-band

oscillations in a cortical model with amplitude-modulated high-frequency

transcranial electric stimulation. *Neuroimage* **173**, 3–12 (2018).

106. Cohen, J. *Statistical power analysis for the behavioral sciences*. *L. Erlbaum*

Associates **13**, (1988).

107. Hashemirad, F., Zoghi, M., Fitzgerald, P. B. & Jaberzadeh, S. The effect of anodal

transcranial direct current stimulation on motor sequence learning in healthy

individuals: A systematic review and meta-analysis. *Brain Cogn.* **102**, 1–12 (2016).

108. Soutschek, A., Kang, P., Ruff, C. C., Hare, T. A. & Tobler, P. N. Brain Stimulation

Over the Frontopolar Cortex Enhances Motivation to Exert Effort for Reward. *Biol.*

Psychiatry **84**, 38–45 (2018).

109. Wischnewski, M., Zerr, P. & Schutter, D. J. L. G. Effects of Theta Transcranial

Alternating Current Stimulation Over the Frontal Cortex on Reversal Learning.

Brain Stimul. **9**, 705–711 (2016).

110. Guerra, A., López-Alonso, V., Cheeran, B. & Suppa, A. Variability in non-invasive

brain stimulation studies: Reasons and results. *Neurosci. Lett.* **719**, 133330

(2020).

111. Cassarà, A. M. *et al.* Safety Recommendations for Temporal Interference

Stimulation in the Brain. *bioRxiv* 1–69 (2022). doi:10.1101/2022.12.15.520077

112. Oldfield, R. C. The assessment and analysis of handedness: The Edinburgh

inventory. *Neuropsychologia* **9**, 97–113 (1971).

113. Kaplan, B. A. *et al.* Automating Scoring of Delay Discounting for the 21- and 27-

Item Monetary Choice Questionnaires. *Behav. Anal.* **39**, 293–304 (2016).

114. Mitchell, M. R. & Potenza, M. N. Recent Insights into the Neurobiology of

Impulsivity. *Curr. Addict. Reports* **1**, 309–319 (2014).

115.   Brainard, D. H. The Psychophysics Toolbox. *Spat. Vis.* **10**, 433–436 (1997).

116.   Pelli, D. G. The VideoToolbox software for visual psychophysics: transforming

numbers into movies. *Spat. Vis.* **10**, 437–442 (1997).

117.   Dayan, E., Averbeck, B. B., Richmond, B. J. & Cohen, L. G. Stochastic

reinforcement benefits skill acquisition. *Learn. Mem.* **21**, 140–142 (2014).

118.   Grossman, N. Modulation without surgical intervention. *Science (80-.).* **361**, 461–

462 (2018).

119.   Antal, A. *et al.* Low intensity transcranial electric stimulation: Safety, ethical, legal

regulatory and application guidelines. *Clin. Neurophysiol.* **128**, 1774–1809 (2017).

120.   Ekhtiari, H. *et al.* A checklist for assessing the methodological quality of concurrent

tES-fMRI studies (ContES checklist): a consensus study and statement. *Nat.*

Protoc. **17**, 596–617 (2022).

121.   Bossetti, C. A., Birdno, M. J. & Grill, W. M. Analysis of the quasi-static

approximation for calculating potentials generated by neural stimulation. *J. Neural*

Eng. **5**, 44–53 (2008).

122.   Hasgall, P. *et al.* "IT'IS Database for thermal and electromagnetic parameters of

biological tissues," Version 4.1. (2022). doi:10.13099/VIP21000-04-1

123.   Seeck, M. *et al.* The standardized EEG electrode array of the IFCN. *Clin.*

Neurophysiol. **128**, 2070–2077 (2017).

124.   Bates, D., Mächler, M., Bolker, B. M. & Walker, S. C. Fitting linear mixed-effects

models using lme4. *J. Stat. Softw.* **67**, (2015).

125.   Ryu, E. Effects of skewness and kurtosis on normal-theory based maximum

likelihood test statistic in multilevel structural equation modeling. *Behav. Res.*

Methods **43**, 1066–1074 (2011).

126.   Nieuwenhuis, R., te Grotenhuis, M. & Pelzer, B. Influence.ME: Tools for detecting

influential data in mixed effects models. *R J.* **4**, 38–47 (2012).

127.   Luke, S. G. Evaluating significance in linear mixed-effects models in R. *Behav.*

Res. Methods **49**, 1494–1502 (2017).

128.   Searle, S. R., Speed, F. M. & Milliken, G. A. Population marginal means in the

linear model: An alternative to least squares means. *Am. Stat.* **34**, 216–221

(1980).

129.   Ben-Shachar, M., Lüdecke, D. & Makowski, D. effectsize: Estimation of Effect Size

Indices and Standardized Parameters. *J. Open Source Softw.* **5**, 2815 (2020).

130.   Derosière, G., Billot, M., Ward, E. T. & Perrey, S. Adaptations of motor neural

structures' activity to lapses in attention. *Cereb. Cortex* **25**, 66–74 (2015).

131.   Okamoto, M. *et al.* Three-dimensional probabilistic anatomical cranio-cerebral

correlation via the international 10-20 system oriented for transcranial functional

brain mapping. *Neuroimage* **21**, 99–111 (2004).

132.   Di, X., Zhang, Z. & Biswal, B. B. Understanding psychophysiological interaction

and its relations to beta series correlation. *Brain Imaging Behav.* **15**, 958–973

(2021).

133.   Bowles, S. *et al.* Vagus nerve stimulation drives selective circuit modulation

through cholinergic reinforcement. *Neuron* **110**, 2867-2885.e7 (2022).

134.   Codol, O., Galea, J. M., Jalali, R. & Holland, P. J. Reward-driven enhancements in

motor control are robust to TMS manipulation. *bioRxiv* 2020.01.12.903419 (2020).

doi:10.1101/2020.01.12.903419

135. Draganski, B. *et al.* Evidence for segregated and integrative connectivity patterns

in the human basal ganglia. *J. Neurosci.* **28**, 7143–7152 (2008).

136. Morris, L. S. *et al.* Fronto-striatal organization: Defining functional and

microstructural substrates of behavioural flexibility. *Cortex* **74**, 118–133 (2016).

137. Lam, S. Y. & Zénon, A. Information rate in humans during visuomotor tracking.

Entropy **23**, 1–13 (2021).

138. Uehara, S., Mawase, F., Therrien, A. S., Cherry-Allen, K. M. & Celnik, P.

Interactions between motor exploration and reinforcement learning. *J.*

Neurophysiol. **122**, 797–808 (2019).

139. Rolls, E. T., Huang, C. C., Lin, C. P., Feng, J. & Joliot, M. Automated anatomical

labelling atlas 3. *Neuroimage* **206**, 116189 (2020).

1723