An Approach to Identify Expression Quantitative Super Enhancers Using Single-cell Multi-omic Data

Guoshuai Cai[1]

[1]Department of Environmental Health Science, Arnold School of Public Health, University of South Carolina, Columbia, SC, 29208, USA

Corresponding to:

Guoshuai Cai, Ph.D.

Department of Environmental Health Science, Arnold School of Public Health, University of South Carolina, Columbia, SC, 29208, USA

Discovery 449, 915 Greene St., Columbia, SC, 29208

Tel: 803-777-4120

Email: gcai@mailbox.sc.edu

Abstract

Super enhancers (SEs) drives cell identity and disease related genes. However, current methods for studying associations between SE and gene expression are time consuming, costly and have difficulties in scalability.

This study formulated an approach for screening genome-wide SE-expression

associations using single-cell multi-omic data of transcriptome and H3K27ac histone modification.

A pipeline was also constructed for an easy workflow application. Further application identified

expression quantitative SEs (EqSEs) in brain and found they mark cell types. Moreover, our

analysis provided new insights into the functional role of SEs close to *Zbtb20* and *Prkcb* in frontal

cortex near-projecting neurons and hippocampus CA1 neurons, linking to neuron development

and neurological diseases.

Collectively, this study provides a new tool for studying SE-expression associations and

identifying significant expression associated SEs, which pave the way for understanding the

regulatory role of SE in gene expression and related cellular and disease development.

Introduction

Super enhancers (SEs) are large clusters of enhancer elements, which have been found to be

cell-type specific and activate the expression of cell identity genes. Compared to the typical

enhancers, SEs bind to high levels of master transcription factors and Mediator coactivator and

drives high-level expression of their associated genes [1]. Stimulating oncogenic transcription and

other processes, SEs play key roles in the pathogenesis of various diseases, including cancers,

neurodegeneration and others [2]. Therefore, understanding the regulatory role of SEs on gene

expression will provide new insights into the cell and disease development. Genome-wide SE-

expression associations will provide key data for this aim. However, current studies mainly rely

on knock-out experiments on a single or a few SEs which are time consuming and costly, and

have difficulties in scalability. In this study, a single-cell multi-omics based approach was

developed to unblock this significant bottleneck.

The proposed approach studies SE-expression associations and identifies significant

expression associated SEs (referred to as expression quantitative SEs, EqSEs) by analyzing

single cell multi-omic data of transcriptome and H3K27ac histone modification (which is a reliable

surrogate marker for SE identification [2]). In the following application study, EqSEs were

identified and found to be activated in specific mouse brain cell types. The synchronized activity

of SE and gene expression in frontal cortex (FC) and hippocampus (HC) neuron subtypes provide

new evidence of the important role of SEs in neuron development and neurological diseases.

Methods

Datasets

Multi-omic data of H3K27ac modification and transcriptome of single cells from the frontal cortex

and hippocampus tissues of Male C57BL/6J mice were generated by Paired-Tag [3]. The

processed RNA sequencing read counts were downloaded from GEO GSE152020 and the raw

protein binding DNA sequencing reads were downloaded from the SRA project of PRJNA638110.

Single-cell transcriptomic data processing and cell type identification

Single-cell transcriptomic data were processed and analyzed using Seurat [4]. The RNA

sequencing read counts were normalized using the "LogNormalize" method and further centered

and scaled by standard deviations. 2000 highly variable feature were selected, and dimension of

their space was reduced using the top 20 principal components. Further, cell clusters were

identified using the Louvain optimization-based clustering method and were visualized in reduced

dimensions of Uniform Manifold Approximation and Projection (UMAP). Based on the expression

of markers that were used in the original study [3], 18 cell types were identified, including 7 FC

cell types (FC L2/3, FC L4, FC L5, FC PT, FC NP, FC CT, FC L6), 3 HC cell types (HC CA1, HC

CA2/3, HC DG), 3 interneurons (InNeu-CGE, InNeu-Sst, InNeu-Pvalb), Oligodendrocyte

precursor cells (OPC), Oligodendrocytes (Oligo), Astrocytes, Endothelial cells and Ependymal

cells. The cell clusters and corresponding marker expression profile were shown in Figure S1.

Single-cell histone modification data processing and SE identification

Single-cell protein binding DNA sequencing reads were processed following the method of Zhu

et al. [3] for barcode extraction and genome mapping to the mm10 mouse genome assembly.

Then, in cells of each major neuron type of FC, HC and InNeu which were identified by above

transcriptome analysis, the peaks of H3K27ac signal were identified by MACS2 [5]. From these

peaks, SEs were further identified by ROSE [1] in each single cell and H3K27ac binding DNA

sequencing reads at SE regions were counted for quantification. In a particular cell, the signals

were normalized by all reads with a consistent scale factor, 10000, and log transformation was

applied to transform data to approximately conform to normality.

EqSE identification

For each neuron type, the associations between the H3K27ac signal at each SE and the

expression of its proximal gene was assessed by Pearson correlation coefficients (r). To reduce

the bias and noise of single-cell data due to its high drop-out rate, cells with missing values were

filtered out from correlation calculation. To obtain reliable correlation assessment, only SE-gene

pairs that have non-missing multi-modal data in more than 300 cells and more than 20 in each

cell type of FC, HC and InNeu were analyzed in this study. EqSEs were considered to have a

notable correlation with $r > 0.2$ in at least one cell type.

Results

A workflow/pipeline for screening expression quantitative SEs

To enable the EqSE screening, a workflow was designed as illustrated in Figure 1. It is a multi-

step procedure for data processing, sequentially with demultiplexing, transcriptome/genome

mapping, gene expression quantification, clustering and cell type identification based on single-

cell transcriptome data, histone modification peak calling, SE identification, histone modification

quantification, multi-modal data conjugation, and EqSE identification. The details and tool

implementation were described in Methods. The outputs of this workflow are cell type-specific

super enhancer regions, single-cell gene expression and histone modification levels, SE-

expression association, and identified EqSEs. This new approach leverages single-cell multiomic

data of three key advantages: (1) the same-cell joint profiling of transcriptome and histone

modification captures cell-to-cell variation which was ignored by traditional bulk data, (2) the large

scale of single-cell measurements provides favorable statistical power in association assessment,

which was a critical challenge for traditional bulk tissue experiments, and (3) the single-cell

resolution provides an opportunity to build SE atlas in atissue which has been known to be cell

type-specific. In support of the workflow application, a pipeline was constructed and is available

in Github https://github.com/thecailab/EqSE.

EqSE activeness are cell type-specific in brain

Following the workflow, this study assessed the SE-expression association in three major neuron

types, FC, HC and interneurons (InNeu). Consistent associations (i.e., correlations) were widely

found cross these neuron types (Fig. 2). 25 EqSEs were identified ($r > 0.2$, see Methods) and the

expression of 16 associated genes statistically significantly mark brain cell types. As shown in

Figure 3A, specific expression was found on *Kcnip4* and *A830082K12Rik* in FC L2/3, L4 and L5;

Osbpl1a, *Phactr1*, *Prickle2* and *Prkcb* in FC and HC; *Sh3rf3*, *Klhl29*, *Hivep2* and *Pde4dip* in

neuron cells; *Zbtb20* in HC and non-neuron cells; *Zbtb16* in HC and InNeu-CGE; *Tiam1* and

Scn3a in HC DG; *Lrrc4* in InNeu-Pvalb; *Ctnna3* in Oligo and Ependymal cells. Despites of low

coverage, histone modification at these EqSEs also showed a corresponding pattern of

activeness (Fig. 3B).

Interestingly, the histone modification at SEs close to *Zbtb20* and *Prkcb* are specifically

correlated to gene expression in FC and HC respectively (Figs. 2 and 4). As compared to other

subtypes of FC neurons, synchronized activeness of histone modification at SE and gene

expression of *Zbtb20* were found in FC near-projecting neurons (FC NP) with relatively higher

detection rate and average level (Fig.4 left, p-value= 9.86×10^{-6} for gene expression, non-

significant for histone modification data due to the low sequencing coverage). In HC CA1 neurons,

significant activeness of *Prkcb* were observed on both histone modification at SE (p-value=

9.66×10^{-3}) and gene expression (p-value= 8.52×10^{-58}), as compared to other HC neurons (Fig.4

right).

Discussion

This study developed a novel workflow with pipeline to study SE-expression association and

identify EqSEs. It provides a new approach for interactively analyzing the same-cell transcriptome

and histone modification and investigating SE-related regulatory programs in cell populations. Its

performance was cross validated by the broad consistence of associations in three major neuron

cell types. The study also identified specific associations at *Zbtb20* in FC and at *Prkcb* in HC,

indicating SE-related regulation could be cell type-specific. Previous studies suggested an

important role of *Zbtb20* in cell fate specification and in instructing the generation of a subset of

callosal projections neurons in cortical layers II/III [6, 7]. Aligning well with these studies, active

gene expression of *Zbtb20* was found in FC L2/3 and NP cells. In addition, active expression of

Prkcb was found in HC CA1 cells, in consistent with the previous findings that CA1 pyramidal

neurons is highly vulnerable in Alzheimer's disease (AD) and was characterized by *Prkcb*, a

causative factor of AD [8, 9]. In this study, the newly identified synchronous activation of SEs in

specific cell types may play critical roles in underlying mechanisms of these related diseases.

A limitation resides in the low per cell sequencing coverage compromises the sensitivity

of the single-cell profiles, especially for histone modification detection. This limitation will be

remediated by the reducing cost of sequencing and single cell library construction. With necessary

modifications, the proposed workflow is expected to be easily extended for data from future

improved platforms, and other platforms for assessing the associations of other features, such as

methylation and gene expression.

Conflict of Interest: none declared.

References

1. Whyte, W.A., et al., *Master transcription factors and mediator establish super-enhancers at key*
cell identity genes. Cell, 2013. **153**(2): p. 307-19.
2. Hnisz, D., et al., *Super-enhancers in the control of cell identity and disease.* Cell, 2013. **155**(4): p.
934-47.
3. Zhu, C., et al., *Joint profiling of histone modifications and transcriptome in single cells from*
mouse brain. Nat Methods, 2021. **18**(3): p. 283-292.
4. Stuart, T., et al., *Comprehensive Integration of Single-Cell Data.* Cell, 2019. **177**(7): p. 1888-1902
e21.
5. Zhang, Y., et al., *Model-based analysis of ChIP-Seq (MACS).* Genome Biol, 2008. **9**(9): p. R137.
6. Medeiros de Araujo, J.A., et al., *ZBTB20 is crucial for the specification of a subset of callosal*
projection neurons and astrocytes in the mammalian neocortex. Development, 2021. **148**(16).
7. Tonchev, A.B., et al., *Zbtb20 modulates the sequential generation of neuronal layers in*
developing cortex. Mol Brain, 2016. **9**(1): p. 65.
8. Gerschutz, A., et al., *Neuron-specific alterations in signal transduction pathways associated with*
Alzheimer's disease. J Alzheimers Dis, 2014. **40**(1): p. 135-42.
9. Zhou, Z., et al., *Molecular identification of protein kinase C beta in Alzheimer's disease.* Aging
(Albany NY), 2020. **12**(21): p. 21798-21808.

Figures

Figure 1. The workflow of EqSE identification using single-cell multiomic data. It
constructs a multi-step processing of (1) sequencing data processing, including demultiplexing,
transcriptome/genome mapping and gene expression quantification, (2) cell type identification
based on single-cell transcriptomics, (3) SE identification and quantification based on cell type-
level histone modification data, (4) multi-modal data conjugation, and (5) EqSE identification.

Figure 2. Comparison of SE-expression associations in Neuron subtypes. For each gene, correlation 1 and 2 are from FC, HC or InNeu for pairwise comparisons. The significant difference of SE-expression associations in *Zbtb20* between FC and HC/InNeu and *Prkcb* between HC and InNeu were identified and labeled above corresponding dots.

Figure 3. Profiles of EqSE features in brain cell types. In a particular cell type, the level of expression of genes (upper) and histone modification at each mapped EqSE (down) were shown by dot size and color. A larger dot size indicates a higher detection rate of gene expression within each cell type, while a darker color indicates a higher average level of signal.

Figure 4. *Zbtb20* and *Prkcb* expression and histone modification at proximal SEs in neuron subtypes. The top shows the SE-expression correlation of *Zbtb20* and *Prkcb* in different neuron types. Uniform manifold approximation and projection (UMAP) plots of single-cell transcriptome with gene expression and histone modification are shown in middle and bottom respectively, for Zbtb20 in FC subtypes (left) and Prkcb in HC subtypes (right). Overlaid dot plots show average detection rates (corresponding to dot size, larger dot indicates higher detection rate) and average signal levels (corresponding to dot color, darker indicates higher signal level).

Figure S1. Brain cell clusters and marker expression profiles. For each cell type shown in
the UMAP plot (top), the detection rate (corresponding to dot size, larger dot indicates higher
detection rate) and average expression level (corresponding to dot color, darker indicates higher
expression level) of each cell-type marker are shown in the dot plot (bottom).

203
204
205
206