Normalized Semi-Covariance Co-Efficiency Analysis of Spike Proteins from SARS-

CoV-2 variant Omicron and Other Coronaviruses for their Infectivity and Virulence

Tong Xu[a], Shanyue Zhou[a,b], Jun Steed Huang[a, c], Wandong Zhang[d, e],

[a] VisionX LLC, San Jose, USA

[b] McGill University, Montreal, Canada

[c] Carleton University, Ottawa, Canada.

[d] Human Health Therapeutics, National Research Council of Canada, Ottawa, Canada

[e] University of Ottawa, Ottawa, Canada

Corresponding: Wandong.Zhang@nrc-cnrc.gc.ca

JunHuang@Cunet.Carleton.Ca

Abstract

Spectrum-based Mass-Charge modeling is increasingly used in biological analysis. To explain statistical phenomenon with positive and negative fluctuations of amino acid charges in spike protein sequences from Omicron and other coronaviruses, we propose calculation-based Mass-Charge modeling, a normalized derivation algorithm with exact Excel and MATLAB tool involving separate quadrant extension to normalized covariance, which is still compatible with Pearson covariance co-efficiency. The number of amino acids, molecular weight, isoelectric point, amino acid composition, charged residues, mass-charge ratio, hydropathicity of the proteins were taken into consideration in the analyses, and the relative peak and dip of the average with spike protein sequences based on hydrophobic mass to isoelectric charges of amino acids were also examined. The analyses with the algorithm provide more clear insights leading to revealing underline evolving trends of the viral proteins. Spike proteins from SARS-CoV-2 variants, seasonal and murine coronaviruses were taken as representative examples in this study. The analyses demonstrate that the Mass-Charge covariance co-efficiency can distinguish subtle differences between biological properties of spike proteins and correlate well with viral infectivity and virulence.

Key words: Equivalent Mass-Charge moment, normalized semi-covariance co-efficiency, coronaviruses, spike protein sequences, amino acid charge, amino acid hydrophobicity, isoelectric point, entropy.

1. Introduction

Spectrum-based Mass-Charge algorithms are used in analyzing real-world implementations. It comes as the trusted analytic solution but typically has a hardware implementation cost and time challenges, leading to a demand for more straightforward software calculation-based solutions using Mass-Charge theory. Mass-Charge algorithms have received significant attention in recent years and are increasingly used to solve real-world problems. Among those is a combination of two or more algorithms involving numerical

algorithms, analytic calculation [1], and other computational techniques, such as artificial

intelligence [2-4], gene analysis systems [5], and gene simulation [6].

Kumar and colleagues found that SARS-CoV-2 Omicron and sub-variants had a higher

positive electrostatic surface potential [7]. This could increase interactions between the

receptor-binding domain (RBD) of Omicron spike protein and the electro-negatively charged

human angiotensin-converting enzyme 2 (hACE2). We compared Omicron spike protein and

its RBD with those of Wuhan-Hu-1 (Wild type) strain. Our previous study calculated the

charges of the SARS-CoV-2 spike protein sequences using the algorithm for viral infectivity

and virulence [8]. In this study, the number of amino acids, molecular weight, theoretical pI,

amino acid composition, charged residues, mass-charge ratio, hydropathicity of the spike

proteins were all analyzed by using the improved algorithm with normalization of spike protein

length.

This study is the first to use Langland's formula to combine all the factors, including the

mass, charge, isoelectric point, hydrophilic and hydrophobic properties, and equipotential of

the amino acids to analyze the spike proteins from Omicron and other coronaviruses. The

hydrophilic and hydrophobic properties express the ability of viral spike proteins to interact

with human cells. There is a linear relationship between the isoelectric point of the amino acids

and the pH value of the amino acids. The isoelectric point and pH value of the amino acid

sequences change the equivalent charge potential. There is also a linear relationship between

hydrophilic and hydrophobic properties of the amino acids and the heat or intrinsic energy

level. The hydrophobicity and the heat/intrinsic energy change the equivalent mass. In the

following sections, we use the equivalent mass and charge value as the ratio to calculate the

overall Mass-Charge ratio. The viral basic reproduction numbers (R0 values) are also an

essential factor for us to determine the correlation of infectivity. R0 values are the expected

number of the cases directly generated by one case in a population where it assumes that all

individuals are susceptible to infection [9]. R0 value is the indicator for the level of contagious

infectious diseases. For example, the R0 value of smallpox is 3.5-6 (varies under different

medical conditions) [10]. The R0 value for the primary strain of the SARS-CoV-2 virus is

estimated to be 1.4-2.4 [11]. The detailed R0 value for each SARS-CoV-2 variant is listed in

Table 1. Another statistics used in the analysis is the death rate for each variant's virulence. A

study published by the Public Health Ontario analyzed the patients infected with the Omicron

variant (37296 cases) or Delta variant (24432 cases) and chose 9087 cases from each variant

to analyze the death rate. Their results showed that the death rate for the Omicron variant was

0.03%, while the death rate for the Delta variant was 0.3% [12].

2. Normalization of Separate Quadrant Covariance

The Mass-Charge parameter is used to measure the fluctuation-term memory of time series. It relates to the autocorrelations of the time series and the derivatives of the Poincare transform of the time series or the momentum generation function at the origin [13]. Studies involving the Mass-Charge parameter were originally developed by the Fuchsian Group in solving differential equations.

Using Langland's formula, we calculate w, h, q, and p to obtain the complete viral spike protein sequences. After the complete sequence is obtained, the N subsequences of the complete sequence are constructed with the window shift of 1, the window length of 16, and the 15 subsequences at the end to reduce the window length in turn.

In this study, we employ the Mass-Charge from Langlands program in below formula (1).

$$mc = \frac{\omega + zh}{q + zp} \qquad (1)$$

The full sequence of Mass-Charge is defined as $S = \{mc_1, mc_2, \dots, mc_n\}$, the n was the length of a full viral spike protein. In the above formula, ω is the molecular mass; h is the hydrophilic and hydrophobic index; q is the charge; p is the isoelectric point; z is equivalent to Finsler distance between viral spike protein and human cell receptor, typical range from 0.01 to 10000 (nm), and the default is 1. Excel is used to calculate the default, MATLAB is used to calculate the non-default distance variation map, and cross-check with the Excel calculation.

2.1. The Convergence and Divergence

The convergence of the viral protein are calculated by formula (2), the divergence of the viral protein are calculated by formula (3).

$$Converge = \frac{AMG(real_{cor(a,a)}) + AMG(real_{cor(b,b)})}{2} - AMG(real_{cor(a,b)}) \qquad (2)$$

$$Diverge = \frac{AMG(img_{cor(a,a)}) + AMG(img_{cor(b,b)})}{2} - AMG(img_{cor(a,b)}) \qquad (3)$$

In formula (2) and formula (3), a and b are two different viral protein sequences; AMG is average moving gate function; $real_{cor(a,b)}$ is the converging covariance of viral protein a and viral protein b, $real_{cor(a,a)}$ and $real_{cor(b,b)}$ are the special case which is that viral protein a was equal to viral protein b, they are calculated from one viral protein sequence with reference to itself instead of to other baseline one. $img_{cor(a,b)}$ is the diverging covariance of

a and b; $img_{cor(a,a)}$ and $img_{cor(b,b)}$ also the special case of $img_{cor(a,b)}$. We calculate from formula (4) and formula (5).

$$real_{cor(a,b)} = \begin{cases} (a-AMG(a) \times (b-AMG(b)), & (a-AMG(a) \times (b-AMG(b)) \geq 0 \\ 0, & (a-AMG(a) \times (b-AMG(b)) < 0 \end{cases} \quad (4)$$

$$img_{cor(a,b)} = \begin{cases} 0, & (a-AMG(a) \times (b-AMG(b)) \geq 0 \\ -(a-AMG(a) \times (b-AMG(b)), & (a-AMG(a) \times (b-AMG(b)) < 0 \end{cases} \quad (5)$$

The AMG operator is defined in here, the arithmetic mean with moving window length function.

$$Y = AMG(X) \quad (6)$$

In the operator of formula (6), $X = \{x_i | i \in [1,n]\}$, $Y = \{y_i | i \in [1,n]\}$. The calculation of AMG operator is formula (7).

$$y_i = \frac{\sum_i^{i+gap} x_i}{gap} \quad (7)$$

2.2. The Positive Correlation and Negative Correlation

Positive Correlation:

$$Cor_{positive} = \frac{\overline{real_{cor(a,b)}}}{\sqrt{\overline{(a-AMG(a))^2}} \times \sqrt{\overline{(b-AMG(b))^2}}} \quad (8)$$

Negative Correlation:

$$Cor_{negative} = \frac{\overline{img_{cor(a,b)}}}{\sqrt{\overline{(a-AMG(a))^2}} \times \sqrt{\overline{(b-AMG(b))^2}}} \quad (9)$$

Pearson Correlation:

$$Cor_{pearson} = Cor_{positive} - Cor_{negative} \quad (10)$$

2.3. The Center Gravity Positive, Center Gravity Negative and M/Q Span

Center Gravity Positive:

$$center_{Converge} = \left[\frac{\sum real_{cor(a,b)} \times I}{\sum real_{cor(a,b)}}\right] \times \delta \quad (11)$$

[] operator is Rounding operator; δ is percentage, due to different length of viral proteins;

$$\delta = \frac{N_{std}}{N_v} \quad (12)$$

N_{std} is the standard baseline viral protein sequence length, N_v is now viral protein sequence length. I is a sequence from 1 to the length of viral protein sequence.

Center Gravity Negative:

$$center_{Diverge} = \left[\frac{\sum img_{cor(a,b)} \times I}{\sum img_{cor(a,b)}}\right] \times \delta \tag{13}$$

M/Q Span:

$$MQSpan = |center_{Converge} - center_{Diverge}| + \text{ReLU}(center_{Diverge} - center_{Converge}) \tag{14}$$

Here ReLU(x) is defined as Max (0, x).

2.4. Infective Reproduction Rate and Max Position

$$RR_v = \frac{MQSpan_v \times RR_{std}}{MQSpan_{std}} \tag{15}$$

RR_{std} is the standard baseline virus sequence infective reproduction rate; RR_v is estimated viral reproduction rate; $MQSpan_{std}$ is the viral protein sequence standard M/Q Span; $MQSpan_v$ is estimated viral protein sequence M/Q Span. Reproduction rate here is used to approximate the observed R0 number.

Max Positive Position

$$MPP = mp \left|real_{cor(a,b)}\right| \tag{16}$$

The "mp| |" is the operator that get the position of max value in $real_{cor(a,b)}$.

Max Negative Position

$$MNP = mp \left|img_{cor(a,b)}\right| \tag{17}$$

2.5. M/Q Density and Virulence

$$MQDensity = \sqrt{MPP * MNP} - MQSpan_v \tag{18}$$

$$Virulence_v = \frac{MQDensity_v \times Virulence_{std}}{MQDensity_{std}} \tag{19}$$

173    $MQDensity_{std}$ is the standard baseline viral protein sequence M/Q density; $MQDensity_v$

174   is now viral protein sequence M/Q density; $Virulence_{std}$ is the viral standard virulence;

175    $Virulence_v$ is current viral virulence.

177   **3. Excel calculations of normalized semi-covariance for spike proteins from SARS-CoV-**

178   **2 and other coronaviruses**

179   To prove the usage of the simplified Mass-Charge variances, we compare the

correlation of SARS-CoV-2 viral spike protein with other coronaviral spike proteins [14].

Since Excel is capable of handling ReLU, we simplify the calculation with normalization

between two variables only, although more than two can also be done. Since each coronaviral

protein from animal or human has different electrical charge level [15], we normalize the

covariance by the variance respectively (so that the comparison is focused on the pure

difference) [16]. We calculated the whole sequences of spike proteins [17] and plotted the curve

forward starting from the low end to high end (from 1 to 1500 or 1900 depends on the length

of the spike proteins). By using the moving window of 16 neighboring amino acids [18], we

calculated the covariance and average over the same area of the sequences to make the curve

more visually smooth for easily comparisons [19]. Figures 1 to 3 show the calculation results

for the most related spike protein sequences out of past years from murine coronaviruses, from

SARS-CoV-2 and the variants Omicron, Delta, IHU, and from seasonal coronaviruses OC43,

229E, HKU1, and NL63 for human (Table 2) [20-22].

3.1. The analysis results for Wuhan strain spike protein in comparison with spike proteins

from SARS-CoV-2 variants

196   We compared the spike protein of Wuhan strain SARS-CoV-2 (NC_045512.2) with the

spike proteins from SARS-CoV2 variants including Mu (B.1.621; GISAID:

EPI_ISL_4029606), Delta (B.1.617; GISAID: EPI_ISL_1731198), Omicron (GISAID:

EPI_ISL_6951145), IHU (B.1.640; GISAID: EPI_ISL_8416940), Indonesia variant

(B.1.466.2; GenBank: QTS26735), Alpha/UK variant (B.1.1.7; EPI_ISL_744131), and BA.5

(B.1.1.529+BA; EPI_ISL_12464782). For the comparison, Wuhan strain spike protein is used

as a baseline, and Delta variant used as a reference point. We set RR_{std}=2.08 (Delta variant),

and $Virulence_{std}$=2.15% (Delta variant). The analyses results are presented in Figure 1. It

appears that most changes of the Mass-Charge ratios (the peaks and dips in Figure 1A to 1F)

occur in the N-terminal half or the middle region of the spike proteins but rarely occurs in the

C-terminal half of the spike proteins. The N-terminal half or the S1 subunit of the spike proteins
carries the receptor-binding domain (RBD) of the spike protein while the middle region of the
spike protein may have the function that directly or indirectly affects the interactions of RBD
to the ACE2 receptor. Mutations in the surrounding region of the RBD may have epistatic effect
on the binding of the spike protein to the ACE2 as we reported previously [23]. Epistasis is the
combinatory effect of two or more mutations in a genome [24].   Epistatic mutations may allow
spike protein to adopt a specific conformation for better or more efficient interaction of RBD
with ACE2 [23]. The RBD is located between 331 and 528 amino acid residues of the spike
proteins [23]. All the variants (including Figure 1A to 1F) show the changes of Mass-Charge
ratios in the RBD region of the spike proteins as compared to Wuhan strain spike protein. These
changes in the RBD region affect the infectivity and transmissibility of the viral variants. It is
also noticed that the changes of Mass-Charge ratios frequently occur closely on both sides of
the RBD region in all the variants (Fig. 1A to 1D & 1F) except the variant B.1.466.2 (Fig. 1E).
Overall, there are more changes in the RBD regions of the Omicron and its subvariant BA.5
spike proteins (Fig. 1C and 1G) as compared to others (Fig. 1A, 1B, 1D, 1E, and 1F).   As said
above, these Mass-Charge ratio changes on both sides of RBD may have direct or indirect
impacts on the interactions of RBD with ACE2 receptor. It is also noted that there are not many
changes in the C-terminal region of the variant spike proteins (Fig. 1A to 1F), indicating that
this region is relatively stable or conserved during evolution.

Figure 1. The analysis results of spike protein from Wuhan strain SARS-CoV-2 in comparison with spike proteins from SARS-CoV-2 variants. The peaks and the dips in the graphs represent variations in Mass-Charge ratio of the amino acids of the spike proteins, meaning that mutations occur in this region resulting in changes in charge/mass of the amino acids. The wider peaks or dips represent the changes involved in more amino acid changes of the spike proteins. The peaks indicate positive correlation and the dips indicate negative correlation. The dips mean opposite changes in the charge/mass of the amino acids of the spike proteins. The dips may have stronger impact on the biological function of the proteins than the peaks.   Panel A: Normalized semi-covariance for generalized mass-charge ratio of Mu variant (B.1.621) spike protein baselined on Wuhan strain spike protein. Panel B: Normalized semi-covariance for generalized mass-charge ratio of Delta variant (B.1.617) spike protein baselined on Wuhan strain spike protein. Panel C: Normalized semi-covariance for generalized mass-charge ratio of Omicron variant spike protein baselined on Wuhan strain spike protein. Panel D: Normalized semi-covariance for generalized mass-charge ratio of IHU variant (B.1.640) spike protein baselined on Wuhan strain spike protein. Panel E: Normalized semi-covariance for generalized mass-charge ratio of Indonesia variant (B.1.446.2) spike protein baselined on Wuhan strain spike protein. Panel F: Normalized semi-covariance for generalized mass-charge ratio of Alpha/UK variant (B.1.1.7) spike protein baselined on Wuhan strain spike protein. Panel G: Normalized semi-covariance for generalized mass-charge ratio of BA.5 (B.1.1.529+BA) spike protein baselined on Wuhan strain spike protein.

Further to the data presented in Figure 1, Table 1 shows the detailed analysis results for the comparison of Wuhan strain spike protein with the spike proteins from SARS-CoV-2 variants. In Table 1 (also in Tables 2 and 3 as well), the Center Convergence is the center point of positive momentum, meaning that the Mass-Charge ratio's positive momentum on both sides of the spike protein sequence is equivalent; while the Center Divergence represents the center point of negative momentum, meaning that the Mass-Charge ratio's negative momentum on both sides of the sequence is equivalent. The M/Q Span is the non-Euclidean distance between the center of convergence and the center of divergence.   The Reproduction rate in the table is the calculated result from the analyses. Max Positive Position is the maximal point of positive momentum, meaning that the Mass-Charge ratio on this point reaches the maximal positive value; while the Max Negative Position is the maximal point of negative momentum, meaning that the Mass-Charge ratio on this point reaches the maximal negative value. M/Q Density is the non-Euclidean height between the center of convergence and the center of divergence. Virulence in the Tables is the calculated results from this algorithm for the viruses. From the analysis results presented in Table 1, it appears that the calculated viral reproduction rates are very close to the R0 values reported in the literature. In particular, the calculated reproduction rate for Omicron (7.79) is almost identical to the reported R0 value for Omicron (7.0) in the

literature, suggesting that the calculated result by using the algorithm based on the Mass-Charge changes in the amino acids of the spike proteins may accurately predict the infectivity of the viral variant. More importantly, the virulence in Table 1 calculated by using the algorithm is closely within the range of death rates reported in the literature, indicating that the calculated virulence of the viral variants may predict the death rate caused by the viral variants. The calculated virulence of Omicron subvariant BA.5 is higher to that of Omicron in Table 1. A new study using multiscale investigations published in October 13, 2022 suggests that the risk of BA.4/5 to global health is greater than that of original BA.2 and that BA.4/5 is more pathogenic than BA.2 [25], which is consistent with our calculation. However, due to popular vaccination/boosters and advanced COVID therapies available, the actual death rate caused by BA.5 infection appears to be similar to that of Omicron (Table 1). BA.5 may have immune escape from current vaccination and therapies, but the symptoms and severity of the disease caused by BA.5 infection may be reduced by available vaccination and therapies [25].

Table 1: Analysis results of spike protein from SARS-CoV-2 Wuhan strain in comparison with spike proteins from SARS-CoV-2 variants.

Wuhan (baseline)/ Delta (reference)	Delta	Omicron	IHU (France)	Indonesia variant	Mu variant	UK (Alpha)	BA.5
Positive Correlation	99.44%	98.09%	98.80%	99.59%	99.28%	99.58%	98.27%
Negative Correlation	0.17%	1.00%	0.59%	0.27%	0.52%	0.32%	0.89%
Pearson Correlation	0.9927	0.9709	0.9821	0.9932	0.9876	0.9926	0.9738
Center Convergence	629	631	633	626	628	626	633
Center Divergence	557	361	419	707	478	655	417
M/Q Span	72	270	214	162	150	58	216
Reproduction Rate	2.08	7.79	6.17	4.67	4.33	1.67	6.23
R0 Value*	5.08	7.0	8.2	6.79	3.5	5.6	7.0
Max Positive Position	518	518	516	516	517	516	518
Max Negative Position	616	216	490	757	485	614	616
M/Q Density	493	64	289	463	351	505	349
Virulence	2.15%	0.28%	1.26%	2.02%	1.53%	2.20%	1.52%
Death Rate*	0.3%-3.4%	0.06%-0.3%	0.9%-2.3%	1.0%-2.3%	2.0%-3.0%	1.3-5.3%	0.06%-0.3%

* The R0 value and the death rate were obtained from Wikipedia.

3.2. Analysis results for Omicron spike protein in comparison with spike proteins from human common cold coronaviruses and murine coronaviruses

We compared the Omicron spike protein with the spike proteins from murine coronaviruses FJ64 (Murine coronavirus RA59/R13; Genbank: FJ647218.1) and AB55

(GenBank: AB551247.1) as well as from human common cold (seasonal) coronaviruses 229E

(GenBank: NC_002645.1), OC43 (GenBank: MN488635.1), NL63 (GenBank: KY554970.1),

and HKU1 (GenBank: MN488637.1). For this comparison, we use Omicron spike protein as

the baseline and OC43 spike protein as a reference. We set RR_{std} =1.56 (OC43),

$Virulence_{std}$=0.02% (OC43).   Figure 2 shows the analysis results for Omicron spike protein

in comparison with the spike proteins from murine coronaviruses and human common cold

coronaviruses. Unlike the data presented in Figure 1, there are more changes in Mass-Charge

ratios between Omicron spike protein and the spike proteins from the murine and seasonal

coronaviruses as shown in Figure 2. Phylogenetic analysis of the viral genomes shows that

Omicron and variant B.1.1.7 are closely related to Wuhan strain virus as shown in Fig. 1C and

1F; but, as compared to Wuhan strain virus and its variant B.1.1.7, Omicron is closer to seasonal

coronaviruses NL63 and 229E and then to HKU1 and OC43. The phylogenetic analysis also

shows that OC43 and HKU1 as well as NL63 and 229E are closer to murine coronaviruses in

evolution (Fig. 3). However, these differences are not apparently observed in Fig. 2. Overall,

unlike data presented in Fig. 1, there are a lot of more dissimilarities between Omicron spike

protein and the spike proteins from human common cold coronaviruses and murine

coronaviruses (Fig. 2), suggesting distant relationship between Omicron and these

coronaviruses in evolution.

Table 2: Analysis results of Omicron spike protein in comparison with spike proteins from
human seasonal common cold coronaviruses

Omicron (baseline)/OC43 (reference)	OC43	229E	NL63	HKU1
Year: 2021	1953	1965	2004	2004
Positive Correlation	59.06%	56.80%	53.97%	59.38%
Negative Correlation	20.32%	19.93%	21.87%	20.46%
Pearson Correlation	0.3874	0.3686	0.3210	0.3893
Center Convergence	695	686	681	674
Center Divergence	569	629	632	611
M/Q Span	126	57	49	63
Reproduction Rate	1.56	0.71	0.61	0.78
Max Positive Position	1180	1112	1197	1171
Max Negative Position	274	352	432	1290
M/Q Density	443	569	670	1166
Virulence	0.020%	0.026%	0.030%	0.053%
Death Rate*	0.02%	0.03%	0.045%	0.10%

307 * The data were obtained from Wikipedia.
308

Further to the data presented in Fig. 2, Table 2 shows the detailed analysis results for

the comparison of Omicron spike protein with the spike proteins from human common cold

coronaviruses. As compared to the centers of convergence presented in Table 1 (626 to 633)

for SARS-CoV-2 variants, the centers of convergence for the spike proteins from seasonal

coronaviruses is shifted to the right (674 to 695) of the sequence. The centers of divergence for

spike proteins from SARS-CoV-2 variants are scattered from 361 to 707 (Table 1); while the

centers of divergence for the spike proteins from seasonal coronaviruses are more located at

the region from 569 to 632 (Table 2). The max positive positions for SARS-CoV-2 variants are

centralized to 516 to 518 (Table 1); while the max positive positions for seasonal coronaviruses

are centralized to 1112 to 1197 (Table 2). The max negative positions for SARS-CoV-2 variants

and seasonal coronaviruses are mostly scattered on the N-terminal regions of spike proteins

(216 to 757) except HKU1 (1290) (Tables 1 and 2). More importantly, the virulence calculated

by the algorithm using the parameters for the common cold coronaviruses is very closely to the

actual death rates reported in the literature (Table 2). This further demonstrates that the

calculated virulence may predicate the virulence of the viruses, i.e., the death rate caused by

the coronavirus infection.

Figure 2. The analysis results for Omicron spike protein in comparison with spike proteins
from murine coronaviruses and human common cold coronaviruses. The peaks and the dips in
the graphs represent variations in Mass-Charge ratio of the amino acids of the spike proteins,

meaning that mutations occur in this region resulting in changes in charge/mass of the amino
acids. The wider peaks or dips represent the changes involved in more amino acid changes of
the spike proteins. The peaks indicate positive correlation and the dips indicate negative
correlation. The dips mean opposite changes in the charge/mass of the amino acids of the spike
proteins. The dips may have stronger impact on the biological function of the proteins than the
peaks.   Panel A: Normalized semi-covariance for generalized mass-charge ratio of murine
coronavirus FJ64 spike protein (Genbank: FJ647218.1) baselined on Omicron spike protein.
Panel B: Normalized semi-covariance for generalized mass-charge ratio of murine coronaviral
spike protein (GenBank: AB551247.1) baselined on Omicron spike protein. Panel C:
Normalized semi-covariance for generalized mass-charge ratio of spike protein from human
common cold coronavirus 229E baselined on Omicron spike protein. Panel D: Normalized
semi-covariance for generalized mass-charge ratio of spike protein from human common cold
coronavirus OC43 baselined on Omicron spike protein. Panel E: Normalized semi-covariance
for generalized mass-charge ratio of spike protein from human common cold coronavirus NL63
baselined on Omicron spike. Panel F: Normalized semi-covariance for generalized mass-
charge ratio of spike protein from human common cold coronavirus HKU1 baselined on
Omicron spike protein.
347

349 Figure 3. Phylogenetic tree for complete genomes of SARS-COV-2 (Wuhan strain, B.1.1.7,
350 and Omicron), human common cold coronaviruses (NL63, 229E, OC43, and HKU1), and
351 murine coronaviruses (Genbank Accession #: AB551247.1, MW620427.1, GU593319.1,
352 FJ647220.1, FJ884686.1, MF618252.1 and FJ647219.1) by using Clustal Omega.

3.3. The analysis results for OC43 spike protein in comparison with spike proteins from murine coronaviruses and seasonal coronaviruses

We compared OC43 spike protein with the spike proteins from MW62 (murine coronavirus MHV-3; GenBank: MW620427.1), FJ64 (GenBank: FJ647218.1), AB55 (GenBank: AB551247.1), GU59 (GenBank: GU593319.1), FJ88 (GenBank: FJ884686.1), and HKU1 (GenBank: MN488637.1). In this comparison, OC43 spike protein is used as the baseline and the FJ88 spike protein is used as a reference. We set RR_{std}=1.0 (FJ88), $Virulence_{std}$=60% (FJ88). The murine coronaviruses are murine hepatitis virus (MHV) and are positive single-stranded RNA coronaviruses of ~31kb. These viruses are highly infectious and fetal. MHV infection of rodents can cause ~60% death and the mortality can reach 100% in infant mice [26,27]. The standard virulence is thus set at 60% in this analysis, but the reference for virulence can be changed upon the viruses to be used in the analysis. Figure 4 shows the analysis results of OC43 spike protein in comparison with the spike proteins from murine coronaviruses and seasonal coronavirus HKU1. It can be seen from Figure 4 that the similarity of the Mass-Charge ratios between OC43 spike protein and murine/HKU1 coronaviral spike proteins is closer as compared to those between Omicron and seasonal/murine coronaviruses presented in Fig. 2, but the similarity in Fig. 4 is less than those between Wuhan strain virus and its variants presented in Fig. 1. It is also noted that the patterns of Mass-Charge ratios displayed in Fig. 4A to 4E are highly similar to each other. Another important observation is that the variations in the C-terminal part of the spike protein sequences are smaller than the variations in the N-terminal region and the middle region of the spike protein sequences (Fig. 4), further indicating that this region is relatively stable in terms of Mass-Charge ratios and protein sequence.

Table 3 presents the detailed analysis results for the comparison of OC43 spike protein with the spike proteins from murine coronaviruses, showing that Mass-Charge variances reveal more dependency and trend of each protein sequence evolution [28]. From the analyses, the virulence of MHV strains AB551247.1 and GU593319.1 is similar to the strain FJ884686.1 at ~60%; while the virulence of MHV strain FJ647218.1 (42.45%) is less than strain FJ884686.1.

Figure 4. The analysis results of OC43 spike protein in comparison with spike proteins from other coronaviruses. The peaks and the dips in the graphs represent variations in Mass-Charge ratio of the amino acids of the spike proteins, meaning that mutations occur in this region resulting in changes in charge/mass of the amino acids. The wider peaks or dips represent the changes involved in more amino acid changes of the spike proteins. The peaks indicate positive correlation and the dips indicate negative correlation. The dips mean opposite changes in the charge/mass of the amino acids of the spike proteins. The dips may have stronger impact on the biological function of the proteins than the peaks. Panel A: Normalized semi-covariance for generalized mass-charge ratio of murine coronaviral spike protein (GenBank: MW620427.1) baselined on OC43 spike protein. Panel B: Normalized semi-covariance for generalized mass-charge ratio of murine coronaviral FJ64 (GenBank: FJ647218.1) spike protein baselined on OC43 spike protein. Panel C: Normalized semi-covariance for generalized mass-charge ratio of murine coronaviral AB55 (GenBank: AB551247.1) spike protein baselined on OC43 spike protein. Panel D: Normalized semi-covariance for generalized mass-charge ratio of murine coronaviral GU59 (GenBank: GU593319.1) spike protein baselined on OC43 spike protein. Panel E: Normalized semi-covariance for generalized mass-charge ratio of murine coronaviral FJ88 (GenBank: FJ884686.1) spike protein baselined on OC43 spike protein. Panel F: Normalized semi-covariance for generalized mass-charge ratio of human seasonal coronaviral HKU1 (GenBank: MN488637.1) baselined on OC43 spike protein.

Table 3: Analysis of OC43 spike protein in comparison with spike proteins from murine coronaviruses.

OC43 (baseline)/FJ88 (reference)	FJ64	AB55	GU59	FJ88
Positive Correlation	83.28%	83.34%	85.09%	83.43%
Negative Correlation	6.37%	6.31%	5.87%	6.37%
Pearson Correlation	0.7691	0.7702	0.7922	0.7706
Center Convergence	670	751	750	755
Center Divergence	541	638	638	634
M/Q Span	129	113	112	120
Reproduction Rate	7.70	6.73	6.68	7.19
Max Positive Position	1157	1531	1531	1531
Max Negative Position	934	1293	1293	1293
M/Q Density	911	1294	1295	1287
Virulence	42.45%	60.33%	60.37%	60.00%

410
411 Fig. 5. Effects of different distance Z of the Langlands program on viral virulence (amplitude).
412 The curves highlighted in the ovary circle represent the M/Q ratio for optimal distance to
413 interact with the receptors. The arrow direction represents the increase in virulence.
414

415
416 Fig. 6. Effects of different distance Z of the Langlands program on viral infectivity (angle). The
417 arrow direction represents the increase in infectivity.
418

Fig. 5 and Fig. 6 show the effects of different Z values (binding distance) for different

strains of coronaviruses. The valley (binding depth) of the virulence for the different strains of

viruses is slightly different based on the previous analysis. For example, in Fig. 5, the Omicron

variant has a valley with a similar high magnitude to BA5 and Delta. However, a slightly bigger

horizontal Z position value than the Delta variant suggests its lower virulence. Fig. 6 shows the

effect of different Z values based on the angle (binding width). The larger the Z value and the

higher the angle is, the more contagious the virus. The SARS-CoV-1 shows the lowest angle,

while the Omicron variant BA2 and BA5 with the highest Z value is the most infectious variant

known.

Fig. 7. Comparison of different active centers of convergence for SARS-CoV-2 variant spike proteins vs. appearing month and its linear prediction of viral stabilization.

Based on our analysis of the binding coverage for each variant in the past months, we can draw the trend line to demonstrate the evolution of the coronaviruses (Fig. 7). If the binding coverage drops to zero, the viruses may not need further mutations in this region for its infectivity and virulence or the virus may be stabilized relatively. Alternatively, further mutations may not significantly increase the viral infectivity or virulence.

Conclusion

This study presents the construction of a Mass-Charge covariance for equivalent analysis of coronaviral spike protein fluctuations using an equivalent moment basis with a simple algorithm coded in Excel, and other similar tools can be used too. This novel Mass-Charge model reveals an extra performance index over the traditional model, like infective reproduction rate and virulence estimation. The study also compares the spike proteins from murine coronaviruses to the spike proteins from human common cold seasonal coronaviruses. The results of the Mass-Charge calculation show the differences between animal and human coronaviral spike proteins that traditional covariance definition and calculation may overlook. It also reveals the unique positive and negative charge (mutation) section center positions in the viral spike proteins (where the convergence and divergence sections are located). By using the normalization procedure, the new algorithm removes the self-correlation of each viral spike protein and displays only the cross-correlation between the viral spike proteins. By analyzing

various parameters of viral protein sequences using the algorithm, the infectivity and the
virulence of the viruses or viral variants may be accurately predicted, particularly when a new
virus or a new viral variant is emerging. If the viral infectivity and virulence can be estimated
or predicted, it will provide important information for preventative measures or therapeutic
preparedness for the diseases caused by the virus and its variants.

It is envisioned that the Mass-Charge model is a promising alternative for the coronaviral
spike protein analysis as well as for other human and viral protein analyses. The model will be
useful to combine inter-virus and intra-virus characterizations. The simplified Excel calculation
is very easy to use, accurate enough, and forward compatible with the traditional Pearson model
and calculations. The more complicated Matlab code is good for experienced users to do more
deep analyses for viral biology and evolution and for drug development. The example code is
available from an Excel file on the GitHub and Matlab servers:

(https://github.com/steedhuang/Poincare-Fuchs-KleinAutomorphicFunction-COVID19-
Mutations).
(https://www.mathworks.com/matlabcentral/fileexchange/106870-calculate-langlands-
automorphic-mass-charge-spectrum).
468
469

Acknowledgement: The research work conducted by Dr. W. Zhang was supported by funding

from the National Research Council of Canada and by a team grant from the Canadian Institute

of Health Research (CIHR) on the Rapid Research Response to COVID-19 Outbreak to Dr.

Marc-Andre Langlois at the University of Ottawa.

Author contributions: TX and SZ conducted data analyses, drafted the manuscript and

prepared the figures. JSH developed the algorithm, conceived the idea, wrote the manuscript

and prepared the figures. WZ conceived the idea, collected data for coronaviral spike proteins,

wrote the manuscript and prepared a figure, and finalized the manuscript.

Competing interests: The authors declare that there is no any competing interest in this study.

Code availability: The example code is available from an Excel file on the GitHub server:

(https://github.com/steedhuang/Poincare-Fuchs-KleinAutomorphicFunction-COVID19-
[Mutations](https://github.com/steedhuang/Poincare-Fuchs-KleinAutomorphicFunction-COVID19-Mutations)).

486
487

References

1. Zunino, L et al. B. A multifractal approach for Spike Protein inefficiency. Physica A. 2008; 387: 6558–6566.
2. Brooks, R.A. Elephants Don't Play Chess. Robotics and Autonomous Systems. 1990; 6: 3–15.
3. Buchanan, B. G. A (Very) Brief History of Artificial Intelligence. AI Magazine 2005; 26.4: 53-60.
4. Hernandez-Orallo, J. & Dowe, D. L. Measuring Universal Intelligence: Towards an Anytime Intelligence Test [J]. Artificial Intelligence Journal 2010; 174: 1508–1539.
5. Gerla, G. Fuzzy Logic Programming and fuzzy control. Studia Logica 2005; 79: 231-254.
6. Davidovitch, L. et al. Simulation-based Learning: The Learning-Forgetting-Relearning Process and Impact of Learning History. Computers & Education 2008; 50: 866–880.
7. Kumar, S. et al. Omicron (BA. 1) and Sub-Variants (BA. 1, BA. 2 and BA. 3) of SARS-CoV-2 Spike Infectivity and Pathogenicity: A Comparative Sequence and Structural-based Computational Assessment. J. Med. Virol. 2022; 94: 4780-4791.
8. Huang, J. S. et al. Semi-covariance co-efficiency analysis of spike proteins from SARS-CoV-2 and other coronaviruses for viral evolution and characteristics associated with fatality. Entropy 2021; 23: 512.
9. Fraser, C. et al. Pandemic potential of a strain of influenza A (H1N1): early findings. Science 2009; 324: 1557-1561.
10. Kaplan, E. H. et al. Emergency response to a smallpox attack: the case for mass vaccination. Proceedings of the National Academy of Sciences USA 2002; 99: 10935-10940.
11. Achaiah, N. C. et al. R_0 and R_e of COVID-19: can we predict when the pandemic outbreak will be contained? Indian J. Crit. Care Med. 2020; 24: 1125-1127.
12. Public Heath Ontario. Covid-19 data and surveillance. Public Health Ontario. Retrieved June 22, 2022, from https://www.publichealthontario.ca/en/data-and-analysis/infectious-disease/covid-19-data-surveillance
13. Torshin, I.Y. et al. Identification of protein folding cores using charge center model oprotein structure. The Scientific World 2002; 2(S2): 84-86.
14. Car, Z. et al. Modeling the spread of covid-19 infection using a multilayer perceptron. Comput. Math. Methods Med. 2020; 2020: 5714714.
15. Li, W. & Kaneko, K. Long-range correlation and partial $1/f^{\alpha}$ spectrum in a noncoding DNA sequence. Europhysics Letters 1992; 17: 655-660.
16. Korber B. T. et al. Covariation of mutations in the V3 loop of human immunodeficiency virus type 1 envelope protein: an information theoretic analysis. Proc. Natl. Acad. Sci. USA 1993; 90: 7176-7180.
17. Mustafa, S. et al. Peptide-Protein Interaction Studies of Antimicrobial Peptides Targeting Middle East Respiratory Syndrome Coronavirus Spike Protein: An In Silico Approach. Advances in Bioinformatics 2019; 2019: 6815105.
18. Li, W. Generating Non-trivial Long-Range Correlations and 1/f Spectra by Replication and Mutation. Int. J. Bifurcation & Chaos 1992; 2: 137-154.
19. Xu, P-R. et al. Shrinkage estimation analysis of correlated binary data with a diverging number of parameters. Science China Mathematics 2013; 56: 359-377.
20. Bianchi, M. et al. SARS-CoV-2 envelope and membrane proteins: structural differences linked to virus characteristics? Biomed. Res. Int. 2020; 2020: 4389089.
21. Gussow, A. B. et al. Genomic determinants of pathogenicity in SARS-CoV-2 and other human coronaviruses. Proc. Natl. Acad. Sci. USA 2022; 117: 15193-15199.

22. Voss, R. F. Evolution of long-range fractal correlations and 1/f noise in DNA base
sequences. Phys. Rev. Lett. 1992; 68: 3805-3808.
23. Castonguay, N et al. Meta-analysis and structural dynamics of the emergence of genetic
variants of SARS-CoV-2. Frontiers in Microbiology 2021; 12: 676314.
24. Phillips, P. C. Epistasis – the essential role of gene interactions in the structure and
evolution of genetic systems. Nat. Rev. Genet. 2008; 9: 855-867.
25. Kimura I et al. Virological characteristics of the SARS-CoV-2 Omicron BA.2 subvariants,
including BA.4 and BA.5. Cell 2022; 185(21): 3992-4007.e16.
26. Roberts, A. et al. A mouse-adapted SARS-Coronavirus causes disease and mortality in
BALB/c mice. PLoS Pathogens 2007; 3(e5): 23-37.
27. Nicklas, W. et al. Viral infections of laboratory mice. The Laboratory Mouse 2012; 2012:
427-480.
28. Tim, S. et al. COVID-19 Drug Therapy. Elsevier. Oct. 23, 2020 ([COVID-19-Drug-
Therapy_2020-8-28.pdf (elsevier.com)](#)).
551