Title: Control of *Clostridiodes difficile* virulence and physiology by the flagellin homeostasis checkpoint FliC-FliW-CsrA in the absence of motility

Short title: FliW modulates *Clostridiodes difficile* pathogenesis

Authors: Duolong Zhu[1, 2]*, Katherine J. Wozniak[1, 2], Firas Midani[1, 2], Shaohui Wang[3], Xingmin Sun[3], Robert A. Britton[1, 2]

Affiliations: [1]Department of Molecular Virology and Microbiology, Baylor College of Medicine, Houston, Texas 77030

[2]Alkek Center for Metagenomics and Microbiome Research, Baylor College of Medicine, Houston, Texas 77030

[3]Department of Molecular Medicine, Morsani College of Medicine, University of South Florida, Tampa, Florida 33612

*Corresponding author: Duolong Zhu (Email: duolong.zhu@bcm.edu)

ABSTRACT

Mutations affecting *Clostridioides difficile* flagellin (FliC) have been shown to be hypervirulent

in animal models and display increased toxin production and alterations in central metabolism.

The regulation of flagellin levels in bacteria is governed by a tripartite regulatory network

involving *fliC*, *fliW*, and *csrA*, which creates a feedback system to regulate flagella production.

Through genomic analysis of *C. difficile* clade 5 strains (non-motile), we identified they have

jettisoned many of the genes required for flagellum biosynthesis yet retain the major flagellin

gene *fliC* and regulatory gene *fliW*. We therefore investigated the roles of *fliC*, *fliW*, and *csrA* in

the clade 5 ribotype 078 strain *C. difficile* 1015, which lacks flagella and is non-motile. Analysis

of mutations in *fliC*, *fliW*, and *csrA* (and all combinations) on *C. difficile* pathogenesis indicated

that FliW plays a central role in *C. difficile* virulence as animals infected with strains carrying a

deletion of *fliW* showed decreased survival and increased disease severity. These *in vivo* findings

were supported by *in vitro* studies showing that mutations impacting the activity of FliW showed

increased toxin production. We further identified that FliW can interact with the toxin positive

regulator TcdR, indicating that modulation of toxin production via FliW occurs by sequestering

TcdR from activating toxin transcription. Furthermore, disruption of the *fliC-fliW-csrA* network

results in significant changes in carbon source utilization and sporulation. This work highlights

that key proteins involved in flagellar biosynthesis retain their regulatory roles in *C. difficile*

pathogenesis and physiology independent of their functions in motility.

IMPORTANCE

C. difficile is a leading cause of nosocomial antibiotic-associated diarrhea in developed countries

with many known virulence factors. In several pathogens, motility and virulence are intimately

linked by regulatory networks that allow coordination of these processes in pathogenesis and

physiology. Regulation of *C. difficle* toxin production by FliC has been demonstrated *in vitro* and

in vivo and has been proposed to link motility and virulence. Here we show that clinically

important, non-motile *C. difficile* strains have conserved FliC and regulatory partners FliW and

CsrA, despite lacking the rest of the machinery to produce functional flagella. Our work

highlights a novel role for flagellin outside of its role in motility and FliW in the pathogenesis

and physiology of *C. difficile*.

INTRODUCTION

Clostridioides difficile is a Gram-positive, spore-forming, toxin-producing, anaerobic bacterium that is a leading cause of nosocomial antibiotic-associated diarrhea in developed countries (1-3). Recently, *C. difficile* infection (CDI) reports from the community are rising and becoming a larger portion of total CDI cases (4). CDI can result in a spectrum of symptoms, ranging from mild diarrhea to pseudomembranous colitis and death (5). *C. difficile* possesses many virulence factors, such as toxins, binary toxins, adhesions, and flagella, among which toxin A (TcdA) and toxin B (TcdB) are the major ones (6, 7).

While flagella impact bacterial motility, biofilm formation, and host colonization, mutations in flagellin have been shown to play roles in virulence, toxin production, and fitness in several bacteria (8-10). The flagellin protein FliC has been reported to play important roles during *C. difficile* infection in motile strains (9, 11-14). Aubry et al. (11) reported that disruption of flagellar structural gene *fliC* resulted in increased toxin expression. Conversely, disruption of early-stage flagellar genes, such as *fliF*, *fliG*, or *fliM*, led to a significant reduction of toxin production in *C. difficile* 630Δ*erm*. Similarly, Baban et al. (12) reported that *tcdA* expression increased 44.4-fold in a *fliC* gene mutant, while *flgE* mutation led to a 10-fold reduction of *tcdA* expression in *C. difficile* 630Δ*erm*. Interestingly, deletion of other genes involved in flagellar assembly can abolish *C. difficile* motility completely but do not result in a significant change in toxin production that is observed in a *fliC* mutant (11). Based on previous studies, Stevenson et al. hypothesized that the regulation of the flagellar structural genes on toxin expression could be modulated by the direct change or loss of flagellar genes (such as *fliC* gene deletion) rather than the loss of the functional flagella and bacterial motility (9). Although the increase of

pathogenicity of *fliC* mutant *in vivo* has been confirmed and corroborated by several studies (12,

13, 15), how *fliC* modulates *C. difficile* toxin production and pathogenesis remains unclear.

The partner-switching mechanism Hag-FliW-CsrA, which governs flagellin homeostasis and

serves as a checkpoint of flagellar morphogenesis, was first described in *Bacillus subtilis* (16). In

this system, Hag (flagellin protein and FliC homolog), FliW (a flagella synthesis regulator), and

CsrA (a carbon storage regulator) form a sensitive regulatory feedback loop to regulate flagellum

biosynthesis in *B. subtilis*. During this feedback regulation, FliW is first released from a FliW-

Hag complex after the flagellar assembly checkpoint of hook completion and then binds to CsrA

by a noncompetitive mechanism to relieve CsrA-mediated *hag* translation repression to increase

flagellin synthesis concurrent with filament assembly (16, 17). CsrA is an ancestral protein and

has evolved to modulate various physiological processes by regulating mRNA stability and/or

translation initiation of target mRNA (18-24). CsrA typically binds to multiple specific sites that

are located near or overlapping the cognate Shine-Dalgarno (SD) sequence in target transcripts

(25, 26). In *B. subtilis*, two CsrA binding sites were identified in the 5'-UTR of *hag* transcripts

(26). Oshiro et al. (27) further quantitated the interactions of FliW, Hag, and CsrA, and

determined that Hag-FliW-CsrA$^{\text{dimer}}$ functions at nearly 1:1:1 stoichiometry in *B. subtilis*.

The regulatory effects of *csrA* on bacterial physiology and carbon metabolism in *C. difficile*

630Δ*erm* were elucidated through the overexpression of the *csrA* gene (28). However, specific

roles of FliW beyond its function as a flagellin synthesis regulator were not reported. We

previously characterized the pleiotropic roles of FliW and CsrA in *C. difficile* R20291 (29),

supporting a similar partner-switching mechanism termed FliC-FliW-CsrA. Nevertheless, more

detailed investigations into the FliC-FliW-CsrA molecular mechanism and the modulation of the

partner-switching mechanism on pathogenesis in *C. difficile* are needed.

Previous studies have shown that ribotype 078 (Clade 5) *C. difficile* strains lack flagella and

bacterial motility (9). Surprisingly, FliC and FliW are highly conserved in the non-motile clade 5

strains of *C. difficile*, despite their lack of flagella and other genes encoding regulatory and

structural genes involved in motility. Here, we investigated the roles of *fliC*, *fliW*, and *csrA* in

modulating the pathogenesis and physiology of *C. difficile*, independent of flagellum

biosynthesis. Our findings reveal that FliW plays a central role in *C. difficile* toxin production

and virulence.

RESULTS

***fliC*, *fliW*, and *csrA* are conserved and expressed in non-motile *C. difficile* strains**

We previously have shown that *fliW* and *csrA* genes are broadly found in *C. difficile* genomes

and FliW and CsrA are highly conserved in divergent *C. difficile* ribotypes (29). In *C. difficile*

R20291 (RT027 and motile), FliW and CsrA regulate flagellin (FliC) synthesis and bacterial

virulence. We found that *fliC*, *fliW*, and *csrA* are conserved in non-motile *C. difficile* strains,

although many other genes involved in flagellar synthesis have been lost (29). To evaluate the

roles of *fliC*, *fliW*, and *csrA* in non-motile clade 5 *C. difficile* strains, we compared the flagellar

regulon   genes   of   RT078   (CD1015),   RT033   (CLO_DA8024AA_AS),   RT045

(CLO_EA6022AA_AS), RT126 (CLO_EA6160AA_AS), and RT127 (CLO_BA6185AA_AS) to

the clade 2 RT027 (R20291). Clade 5 *C. difficile* strains completely lack early-stage flagellar

genes (F3) and only the late-stage flagellar genes (F1, including *fliC*, *fliW*, and *csrA*) and

glycosylation genes (F2) were found in their genomes (Fig. 1A, Fig. S1A, and Table S1). We

selected RT078 strain CD1015 for further study and confirmed that CD1015 was non-motile (Fig.

1B). To confirm that FliC was still expressed in CD1015 despite the absence of flagella, we

detected FliC in the cytoplasm of CD1015 and motile strain R20291 (Fig. 1C). We verified the

cotranscription of *fliW* and *csrA* (Fig. S1B) and measured the transcription of *fliC* and *fliW-csrA*

(Fig. S1C). These results showed that the *fliC*, *fliW*, and *csrA* genes are expressed in non-motile

C. difficile, prompting us to explore the roles of these three genes in *C. difficile* pathogenesis,

independent of their roles in flagellar biosynthesis and motility.

FliC-FliW-CsrA network negatively regulates *C. difficile* toxins production

To test whether *fliC*, *fliW* and/or *csrA* modulate toxin production and bacterial pathogenicity in

non-motile *C. difficile*, we constructed single deletions of *fliC*, *fliW*, *csrA* as well as all double

mutant combinations and the triple mutant in CD1015 using a CRISPR-Cas12a system (Fig. S2A)

(30, 31). Growth profiles of the different mutants were assessed in BHIS and TY media, and no

significant differences in growth nor motility were observed (Fig. S2B-D). Supernatants of *C.*

difficile cultures grown in TY media at 72 h post-inoculation were collected and toxin A and B

levels were measured by ELISA. All mutants displayed significantly increased toxin production

and transcription, with the notable exception of the Δ*fliC*Δ*csrA* double mutant, which produced

wild-type levels of toxins A and B (Fig. 2A-B).

Interestingly, all mutants carrying a deletion of *fliW* gene displayed increased toxin production.

To determine whether *fliW* was sufficient to restore toxin concentrations to normal levels in

several mutants, we provided *fliW* in trans to Δ*fliW*, Δ*fliC*Δ*fliW*, Δ*fliW*Δ*csrA*, and

ΔfliCΔfliWΔcsrA strains and measured toxin production. Our data indicate that *fliW*

complementation effectively reduced toxin production either to wild-type levels or below (Fig.

2C-D). We note that overexpressing *fliW* in wild-type CD1015 can also reduce toxin A and B

levels (Fig. S3), further supporting a role for FliW in negative regulation of toxin production.

These data indicate that FliW is a key node in negatively regulating *C. difficile* toxin production.

FliW is a key modulator of *C. difficile* pathogenesis in a non-motile strain

To evaluate the role of the FliC-FliW-CsrA system on CD1015 virulence *in vivo*, we utilized the

mutant strains described above in a mouse model of CDI and measured their impact on disease

severity. Antibiotic treated mice were orally challenged with CD1015, *ΔfliC*, *ΔfliW*, *ΔcsrA*,

ΔfliCΔfliW, *ΔfliCΔcsrA*, *ΔfliWΔcsrA*, or *ΔfliCΔfliWΔcsrA* vegetative cells (10^8 CFU/mouse). As

shown in Fig. 3A, mice infected with *ΔfliCΔfliW* (log-rank analysis, P □ 0.001*)*, *ΔfliW* (*P* □

0.01), *ΔfliWΔcsrA* (*P* □ 0.01), or *ΔfliCΔfliWΔcsrA* (*P* □ 0.05) showed a dramatic increase in

mortality (70-90%) compared to CD1015. The single mutant strains, *ΔfliC* and *ΔcsrA*, displayed

an intermediate phenotype in which 30-40% of the mice succumbed to infection.   The

ΔfliCΔcsrA double mutant displayed no increase in disease severity, consistent with the lack of

increase in toxin production of this strain *in vitro*. We scored the severity of disease daily using a

previously described clinical sickness scoring system (CSS) based on behavioral changes, stool

characteristics, and weight loss (32). The highest CSS score for each mouse within 7 days post-

infection was used for CSS comparison between the CD1015 and mutant strains. Our results

showed that clinical scores of all derivative mutants, except *ΔfliCΔcsrA,* were higher than that of

the parent strain, indicating more severe disease in mice colonized mutants that disrupt FliC-

FliW-CsrA regulation (Fig. 3B). These data support FliW as a key player (negative regulator) in

the regulation of toxin production in this regulatory cascade.

Characterization of protein-protein interactions of FliC-FliW-CsrA feedback network

Building on the partner-switching mechanism described in *B. subtilis* for this regulatory network,

we wanted to explore the interactions governing the partner-switching mechanism FliC-FliW-

CsrA in *C. difficile* (29). The *B. subtilis* model posits that FliC directly interacts with FliW,

FliW can bind and antagonize the function of CsrA (Fig. 4A). To test whether these interactions

occur in *C. difficile*, we performed a series of experiments to explore these interactions.

We initially examined the interactions between the proteins FliW and FliC, as well as FliW and

CsrA, using the *E. coli* two-hybrid (BACTH) system (33). We created C-terminal fusions of T25

fragment to FliW (FliW-T25) and T18 fragment to FliC (FliC-T18) and CsrA (CsrA-T18),

respectively. As shown in Fig. 4B and 4C, we detected direct interactions between FliW and FliC

(FliW-T25+FliC-T18) and between FliW and CsrA (FliW-T25+CsrA-T18) with no interaction

detected in the negative controls, as predicted by the model and consistent with previous work in

B. subtilis.

Post-transcriptional regulation of *fliC* expression by CsrA

To investigate the post-transcriptional regulation of CsrA on *fliC* expression (Fig. 4D), we

constructed a reporter plasmid (referred to as P_{lacZ}-UTR-*sfgfp*-P_{tet}-*csrA*) containing two

regulatory elements: one is an IPTG-inducible promoter (P_{lacZ}) driving the expression of the 5'

untranslated region of *fliC* (UTR) fused upstream of the reporter *sfgfp* gene (P_{lacZ}-UTR-*sfgfp*);

the other is a tetracycline-inducible promoter (P_{tet}) driving the expression of *csrA* (P_{tet}-*csrA*) (Fig.

4E). The strain containing only P_{lacZ}-UTR-*sfgfp* was used as a positive control (P_{lacZ}-UTR-*sfgfp*).

As shown in Fig. 4E, induction of *csrA* along with UTR-*sfgfp* expression (P_{lacZ}-UTR-*sfgfp*-P_{tet}-

csrA) decreased the fluorescence of the reporter strain dramatically compared to that of P_{lacZ}-

UTR-*sfgfp*, indicating negative regulation of CsrA on *fliC* expression. Meanwhile, *fliW-csrA*

coexpression (P_{lacZ}-UTR-*sfgfp*-P_{tet}-*fliW-csrA*) could restore fluorescence signal to a level similar

to the positive control strain, indicating FliW antagonizes CsrA. These results support a model in

which CsrA negatively regulates *fliC* expression by interaction with the UTR of *fliC* and FliW

antagonizes the negative regulation by CsrA. To exclude the impact of CsrA on IPTG inducible

fliC transcription, we measured *sfgfp* transcription in different reporter strains containing either

P_{lacZ}-UTR-*sfgfp*, P_{lacZ}-UTR-*sfgfp*-P_{tet}-*fliW*, P_{lacZ}-UTR-*sfgfp*-P_{tet}-*csrA*, or P_{lacZ}-UTR-*sfgfp*-P_{tet}-

fliW-csrA regulation cassette. No significant differences in *sfgfp* transcription were detected (Fig.

S4A).

In *B. subtilis*, two CsrA binding sites have been identified in the 91 bp of the 5' UTR of the *hag*

(fliC) transcript (34). We compared *C. difficile fliC* and *B. subtilis hag* 5'-UTR RNA structures

using the algorithm RNAstructure (35) and found two potential **C**srA **b**inding **s**ites (CBS1: from

-45 to -34; CBS2: from -20 to -9, overlapping *fliC* Shine-Dalgarno sequence) included in two

distinct predicted hairpin structures (Fig. 4F and Fig. S4B). To test whether the predicted binding

sites are required for CsrA post-transcriptional regulation, we deleted either CBS1, CBS2, or

both CBS1 and CBS2 (DD: **d**ouble binding sites **d**eletion) in the *fliC* 5'-UTR and assembled the

full-length UTR (UTR$_F$) and the truncated UTRs into the reporter plasmid (P_{lacZ}-UTR$_{CBS}$-*sfgfp*-

P_{tet}-*csrA*). As shown in Fig. 4G, CsrA lost its ability to regulate *fliC* post-transcriptionally when

CBS2 (P_{lacZ}-UTR_{CBS1}-P_{tet}-*csrA*) or CBS1-CBS2 (P_{lacZ}-UTR_{DD}-P_{tet}-*csrA*) were deleted but kept

similar regulation in the CBS1 deletion reporter strain (P_{lacZ}-UTR_{CBS2}-P_{tet}-*csrA*). These results

indicated that CBS2 (RBS) is required for CsrA post-transcriptional regulation on *fliC* expression.

Taken together, we showed that *fliC* is expressed in non-motile *C. difficile*, characterized the

partner-switching mechanism FliC-FliW-CsrA in *C. difficile*, and identified that CBS2 (RBS) is

required for CsrA post-transcriptional on *fliC* expression.

FliW interacts with toxin expression positive regulator TcdR

The alternative sigma factor TcdR is a key positive regulator of toxin production in *C. difficile*.

As it not only positively regulates toxin expression but also affects *C. difficile* sporulation (36),

this motivated us to investigate whether FliW interacts with TcdR. We first created C-terminal

fusion of T18 fragment to TcdR (TcdR-T18) in the pUT18 plasmid and detected the direct

interaction between FliW and TcdR using the BACTH system described above (Fig. 5A). To

quantify the interaction strength between FliW and TcdR, we conducted a β-galactosidase assay

using cell lysates from the reporter strain (BTH containing FliW-T25+TcdR-T18) by detecting

the OD_{420nm} change over one hour after adding the substrate ONPG (ortho-Nitrophenyl-β-

galactoside). As shown in Fig. 5B, although the control construct TcdR-T18+T25 showed a

weak background color change, the FliW-T25+TcdR-T18 construct exhibited a 2-fold increase in

the rate of color change compared to the TcdR-T18+T25 construct, indicating interaction

between FliW and TcdR. Overall, our data suggest that FliW interacts with TcdR, which may

antagonize TcdR's positive role in the regulation of *C. difficile* toxin expression.

Disruption of FliC-FliW-CsrA affects CD1015 strain carbon utilization

Given the central role of CsrA in bacterial carbon metabolism and that carbon utilization genes

are dramatically upregulated in *C. difficile* Δ*fliC* mutants (37-39), we measured the growth

profile of Δ*fliC*, Δ*fliW*, and Δ*csrA* mutants on 190 carbon sources using Biolog PM1 and PM2A

plates. As shown in Fig. 6A, the three single mutants showed significant growth changes in 13

carbon substrates. Mannose and trehalose displayed the largest increase in growth, therefore we

further examined *C. difficile* growth profiles in the different concentrations of these two sugars.

CD1015, Δ*fliC*, Δ*fliW* and Δ*csrA* mutants were cultured in CDMM media supplemented with 40

mM mannose or 20 mM trehalose, respectively. Our results (Fig. 6B-C) showed that mannose

poorly supports CD1015 growth in CDMM medium, but all of the mutants demonstrated robust

growth on mannose. In trehalose supplemented CDMM media, mutants displayed similar growth

rates to CD1015, but all mutants showed significantly delayed cell lysis (Fig. 6D).

To evaluate the impact of *fliC*, *fliW*, and *csrA* deletions on bacterial fitness in specific carbon

sources, *in vitro* competition assays were performed. CD1015 and each mutant were co-cultured

in CDMM media supplemented with either 40 mM mannose or 20 mM trehalose and incubated

for 12 hours. Primers 1-F/R, which can anneal to both CD1015 and the mutants' genomes, were

used for total bacterial detection and primers 42-F/R, 43-F/R, and 2-F/R which can only pair to

the wild-type genome were used for CD1015 detection (Fig. 6E). As shown in Fig. 6F-H, the

abundance of CD1015 in competition with either Δ*fliC*, Δ*fliW*, or Δ*csrA* strains significantly

decreased at 12 hours. The qPCR competition results were further verified by colony PCR test by

isolating dozens of colonies from each competition (Fig. S5). Our competition data are consistent

with the growth profile analysis, indicating that Δ*fliC*, Δ*fliW*, and Δ*csrA* mutants outcompete the

wild type strain in trehalose and mannose supplemented media. This suggests that mutants might

also outcompete the wild type *in vivo*, potentially impacting *C. difficile* pathogenesis.

DISCUSSION

Bacteria possess complex regulatory networks that govern the decision to settle in a favorable

environment or become motile to find new environments. In pathogens, the decision to make

flagella or biofilms is often intimately tied to nutrient acquisition, virulence factor expression,

and toxin production (40, 41). Clade 5 strains of *C. difficile*, which are phylogenetically distinct

from the other four clades of *C. difficile* strains, are known to colonize livestock and cause

human disease. Indeed, direct transmission between animals and humans has been described (42).

Clade 5 strains are also unique in that they are non-motile and have lost most of the genes that

encode for the machinery that builds and support the flagella, while retaining the major flagellin

protein FliC (Fig. 1A and Fig. S1A). This prompted us to explore the role of the FliC-FliW-CsrA

regulatory network on *C. difficile* physiology in the absence of motility and flagella formation.

The data presented here support that FliW is a key regulator that controls toxin production and

virulence in CD1015 (RT078). Notably, mutation of *fliW* alone or in combination with *fliC*

and/or *csrA* resulted in the greatest reduction in animal survival and increased disease severity

score compared to CD1015 and other mutations. These findings correlated well with the amount

of toxin A and B produced *in vitro*. We also found the double mutant Δ*fliC*Δ*csrA* was the only

mutant that did not increase toxin production *in vitro* and did not result in increased disease

severity *in vivo*. Finally, complementation data indicated that single *fliW* complementation can

effectively reduce toxin production in Δ*fliW*, Δ*fliC*Δ*fliW*, Δ*fliW*Δ*csrA*, and Δ*fliC*Δ*fliW*Δ*csrA*,

either to wild-type levels or below. These data support that FliW is the key node in negatively

regulating toxin production and virulence in the FliC-FliW-CsrA regulatory network. Our

previous work has also shown that *fliW* mutants cause increased toxin production in the motile

hypervirulent *C. difficile* strain R20291 (29).

Previous studies have reported that *tcdR* mutants can affect both toxin production and

sporulation in *C. difficile* (36), and that CsrA functions as a global post-transcriptional regulator

in bacteria (19), involved in stage III sporulation in *Clostridium acetobutylicum* (43). Here, we

not only verified that FliW binds to CsrA but also demonstrated that FliW can interact with TcdR,

indicating that FliW can indirectly affect *C. dificile* toxin expression and sporulation (Fig. S6A).

Because CsrA has been shown to regulate multiple aspects of central metabolism in many

different bacterial species (19, 44), we also were interested in investigating whether the

disruption of this regulatory network had an impact on carbon source metabolism. Recently, an

in vivo gene expression study of R20291Δ*fliC* strain revealed that *treA*, encoding

phosphotrehalase, increased 177-fold in the *fliC* mutant compared to the wild-type strain (13).

Our previous study showed that dietary trehalose can contribute to the virulence of epidemic *C.*

difficile (45). Strikingly, we observed that single mutations in all three genes (*fliC*, *fliW*, *csrA*)

were able to confer the ability of CD1015 to grow on the sugar mannose and altered how

trehalose is metabolized. *C. difficile* needs to balance sporulation, toxin production, and nutrient

utilization in the infected host to successfully invade and colonize the host gut. In the future, one

of the central questions moving forward with this line of research is to identify what additional

inputs regulate the FliC-FliW-CsrA to impact *C. difficile* pathogenesis and physiology.

Elegant previous work has elucidated how the Hag-FliW-CsrA regulatory loop modulates

flagellin homeostasis and bacterial motility in *B. subtilis* (26, 27). However, the export and

assembly of flagella in *B. subtilis* is a key player in the complex partner switch mechanism of

regulation, which is lost in CD1015 and other clade 5 *C. difficile* strains. Our data from two-

hybrid studies and genetic experiments agree with the proposed model of direct interactions

between FliC-FliW and CsrA-FliW with FliC being regulated post-transcriptionally by CsrA.

Notably, in two predicted CsrA binding sites, only the Shine-Dalgarno sequence (located in

CBS2) of *fliC* transcripts is required for CsrA post-transcriptional regulation, which is distinct

from the regulation in *B. subtilis*.

Based on our findings presented in this paper, we hypothesize that deleting *csrA* could enhance

FliC expression. This would increase FliC binding to FliW, reducing the amount of free FliW

(no significant difference in *fliW* transcription between 1015 and Δ*csrA*) (Fig. S6B) in the cell.

Consequently, more TcdR bound by FliW would be released, positively regulating toxin

production. In the Δ*fliC* mutant, although no FliC is present to bind to FliW, we observed lower

levels of *fliW* transcription (Fig. S6B). This is likely because no CsrA (*fliW-csrA* cotranscription)

is needed for *fliC* translation repression, suggesting a potential reduction in FliW levels, which

could also positively regulate toxin production. When both *fliC* and *csrA* were deleted, there was

no disturbance observed in *fliW* transcription or the availability of FliW within the cell, resulting

in no significant change in toxin production. We propose that FliW acts as a negative regulator

of toxin production by antagonizing TcdR whose transcription or activity is regulated by FliC

and CsrA. Disruption of the FliC-FliW-CsrA partner switch could result in less FliW available to

participate in toxin repression (Fig. 7). Since FliW functions as a binding partner for several

identified proteins, it would be worthwhile to uncover additional potential FliW binding targets

using co-immunoprecipitation to better understand its roles in *C. difficile*.

MATERIALS AND METHODS

Bacteria, plasmids, and culture conditions

Table S2 lists the strains and plasmids used in this study. *C. difficile* strains were cultured in

BHIS media (brain heart infusion broth supplemented with 0.5% yeast extract and 0.1% L-

cysteine, and 1.5% agar for agar plates) at 37 □ in an anaerobic chamber (90% N_2, 5% H_2, 5%

CO_2). For carbon substrate metabolization and Biolog analysis, *C. difficile* strains were cultured

in CDMM media supplemented with specific carbon substrates (46, 47). For spores preparation,

C. difficile strains were plated on *C. difficile* 70:30 sporulation agar plates (6.3% Bacto peptone,

0.35% proteose peptone, 0.07% $(NH_4)_2SO_4$, 0.106% Tris base, 1.11% BHI, 0.15% g yeast extract,

0.03% L- cysteine, 1.5% agar) and purified as described earlier (48). To enumerate *C. difficile*

CFU in the fecal sample, fecal solutions were plated on adapted TCCFA plates which contain 5

μg/ml lysozyme and 50 μg/ml kanamycin (49). *Escherichia coli* DH5α, *E. coli* HB101/pRK24, *E.*

coli BL21, and *E. coli* BTH101, were grown aerobically at 37 □ in LB media (1% tryptone, 0.5%

yeast extract, 1% NaCl). *E. coli* DH5α was used as a cloning host, *E. coli* HB101/pRK24 was

used as a conjugation donor host, *E. coli* BL21 was used as a protein expression host, and *E. coli*

BTH101 was used as a protein-protein interaction analysis host. Antibiotics were added when

needed: for *E. coli*, 20 μg/ml chloramphenicol, 50 μg/ml kanamycin, and 100 μg/ml ampicillin;

for *C. difficile*, 15 μg/ml thiamphenicol, 250 μg/ml D-cycloserine, 50 μg/ml kanamycin, 16

μg/ml cefoxitin, and 500 ng/ml anhydrotetracycline.

Chemicals and DNA manipulations

The DNA markers, PCR product purification kit, DNA gel extraction kit, restriction enzymes, cDNA synthesis kit, and SYBR Green RT-qPCR kit were purchased from Thermo Fisher Scientific (Waltham, USA). PCRs were performed with the high-fidelity DNA polymerase NEB Q5 Master Mix, and PCR products were assembled into target plasmids with NEBuilder HIFI DNA Assembly Master Mix (New England, UK). Primers (Table S3) were purchased from IDT (Coralville, USA). All chemicals were purchased from Sigma (St. Louis, USA) unless those stated otherwise.

DNA manipulations were carried out according to standard techniques (50). Gene *fliC*, *fliW*, and *csrA* were assembled into *EcoR*I-*BamH*I digested plasmid pMTL84153, resulting in pMTL84153-*fliC* (referred to as 153-*fliC*), pMTL84153-*fliW* (153-*fliW*), and pMTL84153-*csrA* (153-*csrA*) with primers 20-22, respectively. Gene edit plasmid pDL-1 containing Cas12a (AsCpfI) under control of tetracycline inducing promoter was used for *C. difficile* gene deletion according to the previous reports (30, 31). The target sgRNA was designed and analyzed on the Cas-OFFinder (http://www.rgenome.net/cas-offinder/) and CINDEL (http://big.hanyang.ac.kr/cindel/) websites. Plasmids were conjugated into *C. difficile* as described earlier (51). Briefly, constructed gene deletion plasmids (with primers 23-35) from *E. coli* DH5α were transformed into the donor host *E. coli* HB101/pRK24 and were conjugated into *C. difficile* strains subsequently. Transconjugants were selected on BHIS-TKC plates (15 μg/ml thiamphenicol, 50 μg/ml kanamycin, 16 μg/ml cefoxitin) and subcultured into BHIS-Tm broth (15 μg/ml thiamphenicol) to log phase, then the subsequent cultures were diluted with PBS serially and plated on induction plates (BHIS-Tm-ATc: 15 μg/ml thiamphenicol and 500 ng/ml

anhydrotetracycline). After 48 h incubation, colony PCR was examined with check primers 1-C-

F/R and 2-C-F/R to select the correct gene deletion colonies. Then the deletion plasmid was

cured by several passages in BHIS broth without antibiotics. The genome of 1015Δ*fliC* (referred

to as Δ*fliC*), 1015Δ*fliW* (Δ*fliW*), 1015Δ*csrA* (Δ*csrA*), 1015Δ*fliC*Δ*fliW* (Δ*fliC*Δ*fliW*),

1015Δ*fliC*Δ*csrA* (Δ*fliC*Δ*csrA*), 1015Δ*fliW*Δ*csrA* (Δ*fliW*Δ*csrA*), and 1015Δ*fliC*Δ*fliW*Δ*csrA*

(Δ*fliC*Δ*fliW*Δ*csrA*) were isolated and used as templates for the PCR test with check primers, and

the PCR products were sequenced to confirm the correct gene deletion.

Growth profile and motility assay

C. difficile strains were cultured to an optical density of OD_{600} of 0.8 in BHIS or TY media and

were diluted to an OD_{600} of 0.2. Then, 1% of the cultures were inoculated into fresh BHIS or TY,

followed by measuring OD_{600} for 48 h.

To analyze the difference in carbon substrate utilization between parent strain and mutants,

phenotype microarray plates (PM1 and PM2A) were used to conduct Biolog analysis. Briefly, *C.*

difficile strains were cultured to an OD_{600} of 0.6 in BHIS, then 20% of cultures were inoculated

into CDMM media with 0.5% glucose and were incubated for another 3 - 4 h (52). Following, 2

ml of cultures were centrifuged in the chamber and washed three times with CDMM base

medium, and finally were resuspended in CDMM media and adjusted into the same OD for

Biolog plates and specific carbon source (mannose and trehalose) utilization analysis. The

Biolog plates containing bacteria were incubated in a plate reader at 37°C in an anaerobic

chamber, with OD_{600} measurements taken every 10 minutes. The resulting data were

automatically analyzed using AMiGA software developed by our lab (47).

To examine the effect of *fliC* deletion on CD1015 motility, CD1015 and CD1015Δ*fliC* were

cultured to an OD_{600} of 0.8. For swarming analysis, 2 µl of cultures were dropped onto soft BHIS

agar (0.2%) plates. For swimming analysis, 2 µl of *C. difficile* cultures were penetrated into a

soft BHIS agar (0.15%) tube. The motility assay plates and tubes were incubated in the anaerobic

chamber for 48 h.

Toxin production, gene transcription, and bacteria competition assay

To evaluate toxin production in *C. difficile* strains, 10 ml of *C. difficile* cultures from different

strains cultured in TY media were collected at 72 h post incubation. The cultures were adjusted

to the same OD_{600} value, centrifuged at 4 □, 8000×*g* for 15 min, filtered with 0.22 µm filters, and

then used for ELISA test (EAGLE, CDT35-K01).

For toxin transcription analysis (with primers 37-38), 2 ml of 48 h post inoculated *C. difficile*

cultures in TY media were centrifuged at 4 □, 12000×*g* for 5 min for total RNA extraction. To

assay *manA*, *treA*, and *treA2* gene expression (with primers 39-40), 5 ml of 8 h post inoculated *C.*

difficile cultures in CDMM media supplemented with trehalose or mannose were used for RNA

isolation with TRIzol reagent. The transcription of genes were measured by RT-qPCR. All RT-

qPCRs were repeated in triplicate, independently. Data were analyzed by using the comparative

CT ($2^{-\Delta\Delta CT}$) method with 16s rRNA as a control (with primers 36).

To compare the fitness of CD1015 and mutants in specific carbon substrates, *in vitro* competition

assays were conducted. The same CFU of CD1015, Δ*fliC*, Δ*fliW*, and Δ*csrA* from the log phase

were collected, centrifuged, and washed three times with CDMM base media in the anaerobic

chamber. Following, 10^7 CFU of CD1015 was mixed with the same CFU of Δ*fliC*, Δ*fliW*, and

Δ*csrA*, respectively. The mixed seed cultures were then inoculated into CDMM media

supplemented with mannose or trehalose. After 12 h of post-inoculation, 100 μl of mixed

cultures were serially dilueted and plated on BHIS plates for colony PCR test with primers 3-C-

F/R and 4-C-F/R. Meanwhile, the total genome from mixed cultures were isolated and detected

by qPCR with primer pairs 1, 2, 42, and 43 for competition analysis. In the colony PCR test,

CFU_{1015}/CFU_{mutant} was calculated. In the qPCR test, the fold change of CD1015 accounting for

total bacteria was calculated as: $2^{(\Delta CT_{12h} - \Delta CT_{0h})}$, $\Delta CT = CT_{1015} - CT_{total}$. Among them, CT_{1015} was

acquired by primers 42-F/R in Δ*fliC*, or 43-F/R in Δ*fliW*, or 2-F/R in Δ*csrA* that only paired to

the deleted gene sequence, and CT_{total} was acquired by primers 1-F/R paired to the remaining

part of *fliC* gene after deletion. ΔCT_0 was calculated with genome from the mixed seed cultures

at 0 h.

Bacterial virulence of mutants in the mouse model of CDI

C57BL/6 female and male mice (6 weeks old) were ordered from The Jackson Laboratory,

Ellsworth, ME. All studies were approved by the Institutional Animal Care and Use Committee

of Baylor College of Medicine. The experimental design and antibiotic administration were

conducted as described earlier (53). Briefly, 80 mice were divided into 8 groups in 16 cages and

each mouse was challenged with 10^8 of *C. difficile* vegetative cells after orally administered

antibiotic cocktail (kanamycin 0.4 mg/ml, gentamicin 0.035 mg/ml, colistin 0.042 mg/ml,

metronidazole 0.215 mg/ml, and vancomycin 0.045 mg/ml) in drinking water for 6 days. After 6

days of antibiotic treatment, all mice were given autoclaved water for 2 days, followed by one

dose of clindamycin (10 mg/kg, intraperitoneal route) 24 h before *C. difficile* challenge (Day 0).

The mice were monitored daily for a week for changes in weight, diarrhea, stool characteristics,

behavior, and mortality. All survived mice were humanely euthanized on day 7 of post *C.*

difficile challenge.

Mice fecal pellets at post challenge days 1, 3, and 5 were collected from each mouse and were

transferred into the anaerobic chamber as fast as possible. Following, the feces were diluted with

PBS at a final concentration of 30 mg/ml and 100 μl of diluted feces were plated on TCCFA (5

μg/ml lysozyme and 50 μg/ml kanamycin added) plates to enumerate the total CFU of *C. difficile*.

To calculate the spores CFU in the feces, the diluted feces solution were heated at 65□ for 30

min, then plated on TCCFA plates as well. To evaluate toxin tilter in feces, 30 mg/ml of the fecal

solution was used for TcdA and TcdB ELISA analysis.

Protein-protein interaction analysis

To analyze the protein-protein interaction of FliC, FliW, and CsrA, a bacterial two-hybrid system

(BACTH system kit) was used. We got the BACTH system kit as a gift from Dr. Bouveret (Aix-

Marseille University, France) and followed the previously published protocol (33). Briefly, we

digested plasmid pUT18 and pKNT25 with *BamH*I. Then *fliW*, *fliC*, *csrA,* and *tcdR* genes were

amplified and assembled into the N-terminal of T25 and T18 tags with primers 5-10, respectively,

resulting in plasmid pKNT25-*fliW* (referred as FliW-T25), pKNT25-*fliC* (FliC-T25), pKNT25-

csrA (CsrA-T25), pUT18-*fliW* (FliW-T18), pUT18-*fliC* (FliC-T18), pUT18-*csrA* (CsrA-T18),

and pUT18-*tcdR* (TcdR-T18). Following the two recombinant plasmids encoding FliW-T25 and

FliC-T18 hybrid proteins were co-transformed into competent *E. coli* BTH101 reporter cells to

analyze the interaction between FliW and FliC. For FliW and CsrA interaction analysis, reporter

cells expressing FliW-T25 and CsrA-T18 hybrid proteins were used. Similarly, reporter cells

containing FliW-T25 and TcdR-T18 were employed to test the interaction between FliW and

TcdR. The correct cotransformants were first selected on LB plates with 100 μg/ml ampicillin

and 50 μg/ml kanamycin. Then the protein-protein interaction was analyzed by dotting the

recombinant BTH101 cultures on LB-X-Gal-IPTG indicator plates. The reporter strain

containing empty plasmids pKNT25 and pUT18 was used as a control. The blue colonies

indicate a positive result, while the white colonies indicate no interaction between two proteins.

To quantify the interaction strength between FliW and TcdR, we performed a β-galactosidase

assay according to a previous report with several modifications (33). A total of 10^{10} CFU of cell

cultures were collected and lysed by bead beating in 1 ml of Z buffer (8 g of $Na_2HPO_4 \cdot 12H_2O$,

3.125 g of $NaH_2PO_4 \cdot H_2O$, 0.375 g KCl, 0.123 g $MgSO_4 \cdot 7H_2O$ dissolved in 500 ml H_2O). The

cell lysates were then centrifuged at 4 □, 12000×g for 10 min. Subsequently, 100 µl of the

surenatants were used for the β-galactosidase assay by measuring the OD_{420nm} change after

adding the substrate ONPG. The interaction strength was calculated as $\Delta OD_{420nm} /\Delta t_{(1h)}$. Reporter

BTH101 containing TcdR-T18+T25, FliW-T25+T18, or T25+T18 were used as negative

controls, while BTH101 containing T25-Zip+ T18-Zip was used as a positive control.

Post-transcriptional regulation of CsrA on *fliC* expression

To investigate the post-transcriptional regulation of CsrA on *fliC* expression, we constructed

several recombinant reporter plasmids that containing P*lacZ*-5'UTR-*sfgfp* [5' untranslated region

(5'-UTR) of *fliC* transcript and *sfgfp*-6-His (*sfgfp*) fluorescence reporter under the control of P*lacZ*

induction promoter] and P*tet*-*csrA/fliW/fliW-csrA* (*csrA* or *fliW* or *fliW-csrA* under the control of

tetracycline induction promoter P*tet*) regulation elements with primers 11-19 in pET21b plasmid

using NEBuilder HIFI DNA Assembly Master Mix and transformed these reporter plasmids into

the expression host *E. coli* BL21. To further identify the CsrA binding sites in 5'UTR of *fliC*

transcripts, we analyzed and compared *fliC* and *hag* 5'-UTR structures on RNAstructure website

(https://rna.urmc.rochester.edu/RNAstructureWeb/Servers/Predict2/Predict2.html), following we

synthesized three truncated 5'UTR of *fliC* and assembled them into pET21b-P*lacZ*-5'UTR-*sfgfp*-

P*tet*-*csrA* to evaluate the blocking efficiency of CsrA on *sfgfp* expression.

Sporulation assay

513   *C. difficile* sporulation analysis were conducted as reported earlier (54). *C. difficile* strains were

514   cultured on *C. difficile* 70:30 sporulation agar plates for 5 days, following bacterial lawns were

515   scraped from plates and resuspended in PBS. Afterward, CFU of cultures resuspended in PBS

516   was counted on BHIS plates with 0.1% TA for sporulation ratio analysis. The sporulation ratio

517   was calculated as CFU (65 □ heated, 30 min) / CFU (not heated).

Statistical analysis

The reported experiments were conducted in independent biological triplicates, and each sample

was additionally taken in technical triplicates. Animal survivals were analyzed by Kaplan-Meier

survival analysis and compared by the Log-Rank test. Student's t-test was used for two groups

comparison. One-way analysis of variance (ANOVA) with post-hoc Tukey test was used for

more than two groups comparison. Results were expressed as mean ± standard error of the mean.

Differences were considered statistically significant if $P < 0.05$ (*).

ACKNOWLEDGMENTS

This work was supported by the National Institutes of Health (NIH) grants to R.A.B. (U01AI124290 and R01AI123278) and X.S (2R01AI132711). We thank Dr. Bouveret at Aix-Marseille University, France for gifting us the BACTH system kit.

REFERENCES

1. Lawson PA, Citron DM, Tyrrell KL, Finegold SM. 2016. Reclassification of Clostridium difficile as Clostridioides difficile (Hall and O'Toole 1935) Prevot 1938. Anaerobe 40:95-9.
2. Oren A, Garrity GM. 2018. Notification of changes in taxonomic opinion previously published outside the IJSEM. International Journal of Systematic and Evolutionary Microbiology 68:2137-2138.
3. Sebaihia M, Wren BW, Mullany P, Fairweather NF, Minton N, Stabler R, Thomson NR, Roberts AP, Cerdeno-Tarraga AM, Wang H, Holden MT, Wright A, Churcher C, Quail MA, Baker S, Bason N, Brooks K, Chillingworth T, Cronin A, Davis P, Dowd L, Fraser A, Feltwell T, Hance Z, Holroyd S, Jagels K, Moule S, Mungall K, Price C, Rabbinowitsch E, Sharp S, Simmonds M, Stevens K, Unwin L, Whithead S, Dupuy B, Dougan G, Barrell B, Parkhill J. 2006. The multidrug-resistant human pathogen Clostridium difficile has a highly mobile, mosaic genome. Nat Genet 38:779-86.
4. Fu Y, Luo Y, Grinspan AM. 2021. Epidemiology of community-acquired and recurrent Clostridioides difficile infection. Therap Adv Gastroenterol 14:17562848211016248.
5. Lessa FC, Gould CV, McDonald LC. 2012. Current Status of Clostridium difficile Infection Epidemiology. Clinical Infectious Diseases 55:S65-S70.
6. Lyras D, O'Connor JR, Howarth PM, Sambol SP, Carter GP, Phumoonna T, Poon R, Adams V, Vedantam G, Johnson S, Gerding DN, Rood JI. 2009. Toxin B is essential for virulence of Clostridium difficile. Nature 458:1176-9.
7. Kuehne SA, Cartman ST, Heap JT, Kelly ML, Cockayne A, Minton NP. 2010. The role of toxin A and toxin B in Clostridium difficile infection. Nature 467:711-3.
8. Harshey RM. 2003. Bacterial motility on a surface: many ways to a common goal. Annu Rev Microbiol 57:249-73.
9. Stevenson E, Minton NP, Kuehne SA. 2015. The role of flagella in Clostridium difficile pathogenicity. Trends Microbiol 23:275-82.
10. Duan Q, Zhou M, Zhu L, Zhu G. 2013. Flagella and bacterial pathogenicity. J Basic Microbiol 53:1-8.
11. Aubry A, Hussack G, Chen W, KuoLee R, Twine SM, Fulton KM, Foote S, Carrillo CD, Tanha J, Logan SM. 2012. Modulation of toxin production by the flagellar regulon in Clostridium difficile. Infect Immun 80:3521-32.
12. Baban ST, Kuehne SA, Barketi-Klai A, Cartman ST, Kelly ML, Hardie KR, Kansau I, Collignon A, Minton NP. 2013. The role of flagella in Clostridium difficile pathogenesis: comparison between a non-epidemic and an epidemic strain. PLoS One 8:e73026.

13.   Barketi-Klai A, Monot M, Hoys S, Lambert-Bordes S, Kuehne SA, Minton N, Collignon
A, Dupuy B, Kansau I. 2014. The flagellin FliC of Clostridium difficile is responsible for
pleiotropic gene regulation during in vivo infection. PLoS One 9:e96876.
14.   Tasteyre A, Barc MC, Collignon A, Boureau H, Karjalainen T. 2001. Role of FliC and
FliD flagellar proteins of Clostridium difficile in adherence and gut colonization. Infect
Immun 69:7937-40.
15.   Clark AE. 2016. The Clostridium difficile Flagellar System Mediates Toxin Synthesis,
Pathogenicity, and Activation of Innate Immune Responses. Doctoral. The University of
Arizona, The University of Arizona.
16.   Mukherjee S, Yakhnin H, Kysela D, Sokoloski J, Babitzke P, Kearns DB. 2011. CsrA-
FliW interaction governs flagellin homeostasis and a checkpoint on flagellar
morphogenesis in Bacillus subtilis. Mol Microbiol 82:447-61.
17.   Mukherjee S, Oshiro RT, Yakhnin H, Babitzke P, Kearns DB. 2016. FliW antagonizes
CsrA RNA binding by a noncompetitive allosteric mechanism. Proc Natl Acad Sci U S A
113:9870-5.
18.   Romeo T, Gong M, Liu MY, Brun-Zinkernagel AM. 1993. Identification and molecular
characterization of csrA, a pleiotropic gene from Escherichia coli that affects glycogen
biosynthesis, gluconeogenesis, cell size, and surface properties. J Bacteriol 175:4744-55.
19.   Timmermans J, Van Melderen L. 2010. Post-transcriptional global regulation by CsrA in
bacteria. Cell Mol Life Sci 67:2897-908.
20.   Liu MY, Yang H, Romeo T. 1995. The product of the pleiotropic Escherichia coli gene
csrA modulates glycogen biosynthesis via effects on mRNA stability. J Bacteriol
177:2663-72.
21.   Sabnis NA, Yang H, Romeo T. 1995. Pleiotropic regulation of central carbohydrate
metabolism in Escherichia coli via the gene csrA. J Biol Chem 270:29096-104.
22.   Lawhon SD, Frye JG, Suyemoto M, Porwollik S, McClelland M, Altier C. 2003. Global
regulation by CsrA in Salmonella typhimurium. Mol Microbiol 48:1633-45.
23.   Lucchetti-Miganeh C, Burrowes E, Baysse C, Ermel G. 2008. The post-transcriptional
regulator CsrA plays a central role in the adaptation of bacterial pathogens to different
stages of infection in animal hosts. Microbiology 154:16-29.
24.   Ferreiro MD, Nogales J, Farias GA, Olmedilla A, Sanjuan J, Gallegos MT. 2018.
Multiple CsrA Proteins Control Key Virulence Traits in Pseudomonas syringae pv.
tomato DC3000. Mol Plant Microbe Interact 31:525-536.
25.   Sorger-Domenigg T, Sonnleitner E, Kaberdin VR, Blasi U. 2007. Distinct and
overlapping binding sites of Pseudomonas aeruginosa Hfq and RsmA proteins on the
non-coding RNA RsmY. Biochem Biophys Res Commun 352:769-73.
26.   Yakhnin H, Pandit P, Petty TJ, Baker CS, Romeo T, Babitzke P. 2007. CsrA of Bacillus
subtilis regulates translation initiation of the gene encoding the flagellin protein (hag) by
blocking ribosome binding. Mol Microbiol 64:1605-20.
27.   Oshiro RT, Rajendren S, Hundley HA, Kearns DB. 2019. Robust Stoichiometry of FliW-
CsrA Governs Flagellin Homeostasis and Cytoplasmic Organization in Bacillus subtilis.
mBio 10.
28.   Gu H, Qi H, Chen S, Shi K, Wang H, Wang J. 2018. Carbon storage regulator CsrA plays
important roles in multiple virulence-associated processes of Clostridium difficile.
Microb Pathog 121:303-309.

29. Zhu D, Wang S, Sun X. 2021. FliW and CsrA Govern Flagellin (FliC) Synthesis and Play Pleiotropic Roles in Virulence and Physiology of Clostridioides difficile R20291. Front Microbiol 12:735616.

30. Zhu D, Patabendige H, Tomlinson BR, Wang S, Hussain S, Flores D, He Y, Shaw LN, Sun X. 2021. Cwl0971, a novel peptidoglycan hydrolase, plays pleiotropic roles in Clostridioides difficile R20291. Environ Microbiol 23:5222-5238.

31. Hong W, Zhang J, Cui G, Wang L, Wang Y. 2018. Multiplexed CRISPR-Cpf1-Mediated Genome Editing in Clostridium difficile toward the Understanding of Pathogenesis of C. difficile Infection. ACS Synth Biol 7:1588-1600.

32. Shelby RD, Tengberg N, Conces M, Olson JK, Navarro JB, Bailey MT, Goodman SD, Besner GE. 2020. Development of a Standardized Scoring System to Assess a Murine Model of Clostridium difficile Colitis. J Invest Surg 33:887-895.

33. Battesti A, Bouveret E. 2012. The bacterial two-hybrid system based on adenylate cyclase reconstitution in Escherichia coli. Methods 58:325-34.

34. Dubey AK, Baker CS, Romeo T, Babitzke P. 2005. RNA sequence and secondary structure participate in high-affinity CsrA-RNA interaction. RNA 11:1579-87.

35. Tan Z, Fu Y, Sharma G, Mathews DH. 2017. TurboFold II: RNA structural alignment and secondary structure prediction informed by multiple homologs. Nucleic Acids Res 45:11570-11581.

36. Girinathan BP, Monot M, Boyle D, McAllister KN, Sorg JA, Dupuy B, Govind R. 2017. Effect of tcdR Mutation on Sporulation in the Epidemic Clostridium difficile Strain R20291. mSphere 2.

37. Baker CS, Morozov I, Suzuki K, Romeo T, Babitzke P. 2002. CsrA regulates glycogen biosynthesis by preventing translation of glgC in Escherichia coli. Mol Microbiol 44:1599-610.

38. Hauslein I, Sahr T, Escoll P, Klausner N, Eisenreich W, Buchrieser C. 2017. Legionella pneumophila CsrA regulates a metabolic switch from amino acid to glycerolipid metabolism. Open Biol 7.

39. Sahr T, Rusniok C, Impens F, Oliva G, Sismeiro O, Coppee JY, Buchrieser C. 2017. The Legionella pneumophila genome evolved to accommodate multiple regulatory mechanisms controlled by the CsrA-system. PLoS Genet 13:e1006629.

40. Purcell EB, McKee RW, Courson DS, Garrett EM, McBride SM, Cheney RE, Tamayo R. 2017. A Nutrient-Regulated Cyclic Diguanylate Phosphodiesterase Controls Clostridium difficile Biofilm and Toxin Production during Stationary Phase. Infect Immun 85.

41. Allocati N, Masulli M, Di Ilio C, De Laurenzi V. 2015. Die for the community: an overview of programmed cell death in bacteria. Cell Death Dis 6:e1609.

42. Elliott B, Dingle KE, Didelot X, Crook DW, Riley TV. 2014. The complexity and diversity of the Pathogenicity Locus in Clostridium difficile clade 5. Genome Biol Evol 6:3159-70.

43. Tan Y, Liu Z-Y, Liu Z, Zheng H-J, Li F-L. 2015. Comparative transcriptome analysis between csrA-disruption Clostridium acetobutylicum and its parent strain. Molecular BioSystems 11:1434-1442.

44. Romeo T, Babitzke P. 2018. Global Regulation by CsrA and Its RNA Antagonists. Microbiol Spectr 6.

45. Collins J, Robinson C, Danhof H, Knetsch CW, van Leeuwen HC, Lawley TD, Auchtung JM, Britton RA. 2018. Dietary trehalose enhances virulence of epidemic Clostridium difficile. Nature 553:291-294.

46. Karasawa T, Ikoma S, Yamakawa K, Nakamura S. 1995. A defined growth medium for Clostridium difficile. Microbiology (Reading) 141 (Pt 2):371-5.

47. Midani FS, Collins J, Britton RA. 2021. AMiGA: Software for Automated Analysis of Microbial Growth Assays. mSystems 6:e0050821.

48. Perez J, Springthorpe VS, Sattar SA. 2011. Clospore: A Liquid Medium for Producing High Titers of Semi-Purified Spores of Clostridium difficile. Journal of Aoac International 94:618-626.

49. Wilson KH, Kennedy MJ, Fekety FR. 1982. Use of sodium taurocholate to enhance spore recovery on a medium selective for Clostridium difficile. J Clin Microbiol 15:443-6.

50. Chong L. 2001. Molecular cloning - A laboratory manual, 3rd edition. Science 292:446-446.

51. Heap JT, Kuehne SA, Ehsaan M, Cartman ST, Cooksley CM, Scott JC, Minton NP. 2010. The ClosTron: Mutagenesis in Clostridium refined and streamlined. Journal of Microbiological Methods 80:49-55.

52. Theriot CM, Koenigsknecht MJ, Carlson PE, Jr., Hatton GE, Nelson AM, Li B, Huffnagle GB, J ZL, Young VB. 2014. Antibiotic-induced shifts in the mouse gut microbiome and metabolome increase susceptibility to Clostridium difficile infection. Nat Commun 5:3114.

53. Sun X, Wang H, Zhang Y, Chen K, Davis B, Feng H. 2011. Mouse relapse model of Clostridium difficile infection. Infect Immun 79:2856-64.

54. Zhu D, Bullock J, He Y, Sun X. 2019. Cwp22, a novel peptidoglycan cross-linking enzyme, plays pleiotropic roles in Clostridioides difficile. Environ Microbiol 21:3076-3090.

Figures

Main figure legends

Fig. 1 Non-motile clade 5 *C. difficile* conserves and expresses *fliC*, *fliW*, and *csrA*.

(A) Schematic representation of conserved late-stage flagellar (F1) and glycosylation (F2) genes

in the non-motile clade 5 ribotype 078 *C. difficile* strain 1015. **(B)** Comparison of CD1015 and

R20291 motility. **(C)** Detection of FliC with anti-FliCD antibody in CD1015 and R20291 by

Western blotting.

Fig. 2 FliC-FliW-CsrA network negatively regulates *C. difficile* toxins production.

(A) Toxin concentration in the supernatants of CD1015 and its derivative mutants detected by

ELISA. **(B)** Transcription of *tcdA* and *tcdB* in CD1015 and its derivative mutants detected by

RT-qPCR. **(C)** TcdA production in *fliW* complementation mutants. **(D)** TcdB production in *fliW*

complementation mutants. The supernatants of different cultures were collected at 48 h of post

inoculation in TY media, following analyzed by Toxin ELISA kits at $OD_{450 \, nm}$. Bars stand for

mean ± SEM. * means the significant difference of experimental strain compared to CD1015 (A-

B) or CD1015 with empty plasmid (C-D) (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P <$

0.0001). [#] means the significant difference of *fliW* complementation strain compared to mutant

with empty plasmid (C-D) ([####]$P < 0.0001$). Differences were considered statistically significant

if $P < 0.05$. One-way ANOVA with post-hoc Tukey test was used for statistical significance.

Fig. 3 FliW is a key modulator of *C. difficile* pathogenesis in a non-motile strain.

(A) Survival curve of CD1015 and its derivative mutants in a CDI mouse model. Animal

survivals (10 mice/group) were analyzed by Kaplan-Meier survival analysis and compared by the

Log-Rank test. Differences were considered statistically significant if $P < 0.05$ (*$P < 0.05$, **$P <$

0.01, ***$P < 0.001$). **(B)** Disease severity in mice. The clinical sickness scoring system was used

to evaluate the mice disease severity. The highest CSS score for each mouse within 7 days post

infection was used for CSS comparison in different *C. difficile* strains. Animal experiments were

repeated two times, data shown here were from a representative replicate. * means the significant

difference of experimental strain compared to CD1015 (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$).

One-way ANOVA with post-hoc Tukey test was used for statistical significance.

Fig. 4 Characterization of FliC-FliW-CsrA regulatory loop in *C. difficile*.

(A) Schematic representation of direct interactions of FliC, FliW, and CsrA in *C. difficile*. In the

proposed FliC-FliW-CsrA regulatory loop, FliC binds FliW and FliW binds CsrA. **(B)** Protein-

protein interaction test of FliC-FliW. 1: FliW-T25+T18; 2: FliC-T18+T25; 3: FliW-T25+FliC-

T18; 4: T25+T18; 5: T25-Zip+T18-Zip. **(C)** Protein-protein interaction test of FliW-CsrA. 1:

FliW-T25+T18; 2: CsrA-T18+T25; 3: FliW-T25+CsrA-T18; 4: T25+T18; 5: T25-Zip+T18-Zip.

2 µl of each reporter strain were spotted onto LB plates supplemented with IPTG and X-gal for

indication. **(D)** Schematic representation of CsrA post transcriptional regulation on *fliC*

transcripts. **(E)** Verification of post transcriptional regulation of CsrA on 5'UTR of *fliC*

transcripts. Reporter plasmid pET21b-P$_{lacZ}$-UTR-*sfgfp*-P$_{tet}$-*csrA* (or *fliW* or *fliW-csrA*) was

constructed, following the fluorescence of reporter strains with different recombinant plasmids

were determined. **(F)** Prediction of potential <u>C</u>srA <u>b</u>inding <u>s</u>ites (CBS) in 5'-UTR of *fliC*. CBS1

and CBS2: CsrA potential binding sites 1 and 2 predicated by RNAstructure dynalign results

analysis. **(G)** Identification of required binding sites for CsrA post-transcriptional regulation in

5'UTR of *fliC*. Three truncated 5'UTRs of *fliC* were assembled into the reporter plasmid,

respectively, and subsequently the fluorescence of reporter strains with different truncated

5'UTR plasmids were analyzed. Statistical significance was determined at $P < 0.05$ (*), using a

paired t-test.

Fig. 5 FliW interacts with the toxin expression positive regulator TcdR.

(A) Protein-protein interaction test between FliW and TcdR *in vitro* using a bacterial two-hybrid

system. **(B)** Determination of FliW and TcdR interaction strength. To quantify the interaction

strength between FliW and TcdR, 10^{10} CFU of cell cultures were collected and lysed by bead

beating. The cell lysates were used for β-galactosidase assay by detecting the OD_{420nm} change

over one hour after adding the substrate ONPG. The interaction strength was calculated by

$\Delta OD_{420nm} / \Delta t_{(1h)}$. Reporter BTH101 containing TcdR-T18+T25, FliW-T25+T18, or T25+T18

were used as negative controls. Reporter BTH101 containing T25-Zip+ T18-Zip was used as a

positive control. 2 µl of reporter strain were dotted on LB+IPTG+X-gal indication plates when

needed. Differences were considered statistically significant if $P < 0.05$ (***$P < 0.001$). One-

way ANOVA with post-hoc Tukey test was used for statistical significance.

Fig. 6 Disruption of FliC-FliW-CsrA affects CD1015 strain carbon utilization.

(A) Carrying capacity [K, the maximum growth supported by the environment, K-log $=$

ln(OD_max)-ln(OD_initial)] of growth profile of CD1015, Δ*fliC*, Δ*fliW*, and Δ*csrA* in Biolog

PM1 and PM2A plates. **(B-D)** Growth curves of CD1015 and its derivative mutants in CDMM

media without any added carbon source (B), 40 mM mannose (C), and 20 mM trehalose (D). **(E)**

Primers used for the qPCR in the competition test. **(F-H)** Competition of CD1015 with Δ*fliC* (F),

Δ*fliW* (G), and Δ*csrA* (H). Bars stand for mean ± SEM. * means the significant difference of

CD1015 abundance in the competition mixtures compared to that of at 0 h (**$P < 0.01$, ***$P <$

0.001). One-way ANOVA with post-hoc Tukey test was used for statistical significance.

Fig. 7 Regulation model of FliC-FliW-CsrA on CD1015 pathogenesis.

Supplementary figure legends

Fig. S1 Conservation of flagellar genes in non-motile clade 5 strains and verification of

***fliW-csrA* and *fliC* transcription.**

(A) Schematic representation of flagellar genes in the non-motile clade 5 strains and motile

RT027 R20291. **(B)** Verification of *fliW* and *csrA* co-transcription by RT-PCR. M: DNA ladder;

1-3: CD1015 genomic DNA was used as PCR template; 4-6: CD1015 cDNA was used as PCR

template. 1 and 4: *fliW* PCR test; 2 and 5: *csrA* PCR test; 3 and 6: *fliW-csrA* PCR test. **(C)**

Transcription analysis of *fliC* and *fliW-csrA* in CD1015. Bars stand for mean ± SEM. Differences

were considered statistically significant if $P < 0.05$ (**$P < 0.01$, ***$P < 0.001$). One-way

ANOVA with post-hoc Tukey test was used for statistical significance.

Fig. S2 Generation of CD1015 derivative mutants, and test of bacterial growth profiles and

motility.

(A) Deletion of UTR-*fliC*, *fliW*, *csrA*, and *fliW-csrA*. 1-C-F/R were used to verify *fliC* deletion

and 2-C-F/R were used to test *fliW*, *csrA*, and *fliW-csrA* deletion. M: DNA ladder; 1 and 3:

CD1015 genome as PCR template; 2: CD1015Δ*fliC* (Δ*fliC*) genome test; 4: CD1015Δ*fliW* (Δ*fliW*)

genome test; 5: CD1015Δ*csrA* (Δ*csrA*) genome test; 6: CD1015Δ*fliW*Δ*csrA* (Δ*fliW*Δ*csrA*)

genome test. **(B)** Growth profile in BHIS media. **(C)** Growth profile in TY media. **(D)** Bacterial

motility test. Swarming and swimming analysis were tested with soft BHIS agar (0.2%) plates

and BHIS agar (0.15%) tubes, respectively.

Fig. S3. Toxin expression in the *fliW* overexpression strain.

TcdA and TcdB concentrations in the supernatants of CD1015-pMTL84153 (1015/E) and *fliW*

overexpression strain (1015/*fliW*) were detected by ELISA. Bars stand for mean ± SEM.

Differences were considered statistically significant if $P < 0.05$ (*$P < 0.05$). Statistical analysis

was performed using an unpaired two-tailed *t* test.

Fig. S4 Characterization of FliC-FliW-CsrA regulation loop in *C. difficile*.

(A) Transcription of *sfgfp* in the different recombinant reporter strains. Red column: P_{lacZ}-UTR-

sfgfp; Green column: P_{lacZ}-UTR-*sfgfp*-P_{tet}-*fliW*; Orange column: P_{lacZ}-UTR-*sfgfp*-P_{tet}-*csrA*; Blue

column: P_{lacZ}-UTR-*sfgfp*-P_{tet}-*fliW*-*csrA*; **(B)** Structure comparison of *B. subtilis* 5'-UTR of *hag*

and CD1015 5'-UTR of *fliC*. BS1 and BS2: CsrA binding sites 1 and 2 in *B. subtilis*. CBS1 and

CBS2: CsrA potential binding sites 1 and 2 in CD1015. RNAstructure dynalign results were

calculated (calculate the lowest free energy secondary structures common to two unaligned

sequences).

Fig. S5 Δ*fliC*, Δ*fliW*, and Δ*csrA* mutants outcompete CD1015 in the presence of trehalose

and mannose.

(A) Primers used for the colony PCR in the competition test. **(B-D)** CFU_{1015}/CFU_{mutant} in the

competition cultures. * means the significant difference of CD1015 CFU in the competition

mixtures compared to that of at 0 h (*$P < 0.05$, **$P < 0.01$). One-way ANOVA with post-hoc

Tukey test was used for statistical significance.

Fig. S6 Analysis of sporulation and *fliW* transcription in CD1015 and its derivative mutants

(A) Sporulation analysis. *C. difficile* strains were cultured on 70:30 sporulation agar plates for 5

days, following which the scraped cultures were 10-fold diluted and plated on BHIS plates with

0.1% TA to detect sporulation ratio. The sporulation ratio was calculated as CFU (65 □ heated,

30 min) / CFU (not heated). **(B)** Comparison of *fliW* transcription in CD1015, Δ*fliC*, Δ*csrA*, and

Δ*fliC*Δ*csrA*. Bars stand for mean ± SEM. * means the significant difference of experimental

strain compared to CD1015/E or CD1015 (*$P < 0.05$, **$P < 0.01$). One-way ANOVA with post-

hoc Tukey test was used for statistical significance.

A *C. difficile* flagellar genes

Missing F3 (30 genes) in Clade 5 strains

Late-stage flagellar genes (F1)

fliN *flgN* *fliW* *fliS1* *fliT*
flgM *flgK* *flgL* *fliD* *fliC*
csrA *fliS2*

Glycosylation genes (F2)

Early-stage flagellar genes (F3)

flgB *flgC* *fliM* *fliN*

B R20291   CD1015

C   CD1015   R20291
WT   ΔfliC   WT   ΔfliC
kDa
50
37   FliC   FliC
25

Sickness	Score
Normal	0-2
Mild	3-5
Moderate	6-8
Severe	9-12

A

FliC — activity — FliW — activity — CsrA — translation —

- FliC
- FliW
- CsrA

B

1 2 3
4 5

C

1 2 3
4 5

D

High level FliC

fliC mRNA
UTR
Translation inhibition → No FliC

Low level FliC

fliC mRNA
UTR
Translation

E

P_{lacZ} UTR RBS *sfgfp* P_{tet} *csrA* / *fliW* / *fliW csrA*

- ● Media control
- ◆ P_{lacZ}-UTR-*sfgfp*
- ◆ P_{lacZ}-UTR-*sfgfp*-P_{tet}-*fliW-csrA*
- ● P_{lacZ}-UTR-*sfgfp*-P_{tet}-*fliW*
- ■ P_{lacZ}-UTR-*sfgfp*-P_{tet}-*csrA*

RFU/OD vs Time (h), y-axis 0–20000, x-axis 0–12

F

TCTTGTCCGATTATATAAATGAAGAATTAATAAAAAAGTTAAGAG
TAGAAA**TGACAAGGATGT**CAACTATA**CTAAGGAGGG**TAAAATA
        CBS1   CBS2

P_{lacZ} RBS UTR_{CBS} *sfgfp* P_{tet} *csrA*

G

- ○ P_{lacZ}-UTR_E-*sfgfp*-P_{tet}-*csrA*
- ● P_{lacZ}-UTR_{CBS1}-*sfgfp*-P_{tet}-*csrA*
- ● P_{lacZ}-UTR_{CBS2}-*sfgfp*-P_{tet}-*csrA*
- ● P_{lacZ}-UTR_{DD}-*sfgfp*-P_{tet}-*csrA*

RFU/OD vs Time (h), y-axis 0–20000, x-axis 0–12

A

1: FliW-T25+TcdR-T18
2: TcdR-T18+T25
3: FliW-T25+T18
4: T25+T18
5: Zip (positive control)