Dynamic changes in the transcriptome of oocytes during adolescent-onset PCOS in mice

Danfeng Du [1,2#], Ke Deng [1,2#], Dengxuan Fan[1,2], Congjian Xu[1,2]*

Author information

1. Du Danfeng[1,2#]   **First author**   Attending doctor

   ORCID: 0000-0002-0753-334X

   Email: 693468525@qq.com

2. Deng Ke[1,2#]   **Co-First author**   Doctoral student

   ORCID: 0000-0002-4098-1144

   Email：20111250023@fudan.edu.cn

3. Fan Dengxuan[1,2]   **Second author**   Associate chief physician

   Email：dxfan16@fudan.edu.cn

4. Xu Congjian[1,2*]   **Corresponding author**   Professor

   ORCID: 0000-0002-7954-5892

   Email: xucongjian@fudan.edu.cn

*Xu Congjian is the **Post-Publication Corresponding author** and the **Last author**

Du Danfeng is the **First author** and will **handle correspondence** at all stages of refereeing and publication

# Deng Ke contributed equally to this work and should be listed as **Co-first author**

Affiliations

[1]Obstetrics & Gynecology Hospital, Fudan University, Shanghai, China

[2]Shanghai Key Laboratory of Female Reproductive Endocrine-Related Diseases, Shanghai, China

Address: No.128, Shenyang Road, Yangpu District, Shanghai, China

Zip code: 200000

Abstract: (1) Background: This study aimed to explore temporal changes in the transcriptome of oocytes in an adolescent-onset polycystic ovarian syndrome (PCOS) mouse model. (2) Methods: An adolescent-onset PCOS mouse model was established using DHEA. Genes with a similar expression trend over time were identified using trend analysis. KEGG pathway enrichment analysis and gene regulatory network diagrams were examined for signaling pathways to identify potential hub genes related to the pathogenesis of PCOS. (3) Results: Four main trends of gene expression were extracted, of which six combinations of Venn diagrams were generated. Differentially expressed genes were mainly enriched in oxidative phosphorylation, cell cycle, P53 signaling pathway. Cell cycle-related genes (Skp1, Ccnb1, Orc1 and 5, Wee2, Mapk3, Cdc20) were abnormally down-regulated in the DHEA group. Ptges3 was the top1 DEGs at the initial stage of PCOS modeling. (4) Conclusion: This study provides a novel insight into the altered transcriptome of oocytes from PCOS mice. mtDNA-related genes and Cell cycle-related genes play the most important role in the development of PCOS. Ptges3 was the one of the top DEGs which was up-regulated in DHEA group at the initial stage of modeling, which suggested it may play an important role in the early stage of PCOS.

Keywords: Oocyte; RNA-seq; PCOS; adolescent; mouse model

1. Introduction

Polycystic ovarian syndrome (PCOS) is a common endocrine and metabolic dysfunction. Patients with PCOS may have an increased number of retrieved oocytes during IVF treatment, but the embryo cleavage rate is low for these patients [1]. Embryo developmental competence was reduced in a PCOS model induced by dehydroepiandrosterone (DHEA) [2]. The pathogenesis of PCOS is not fully elucidated, but some evidence indicates a strong heritable component in the development of PCOS. Several genome-wide association studies have identified approximately 200 candidate genes related to the risk of PCOS [3], but not a single fully penetrant variant gene has been reported to date. All candidate genes or mutations show a low penetrance associated with many other covariant factors, such as hormones, environmental changes or lifestyles [4].

Considerable work has been done to classify PCOS in the last several decades, but the consensus regarding PCOS phenotypes remains controversial. Some researchers propose PCOS is mainly caused by ovarian-derived androgen excess, while others suggest it is a complex systemic metabolism disorder [5]. Fetal exposure to excessive androgen, nutritional excess and hyperglycemia during critical stages of prenatal development may program neurobiological circuitry [6]. Prenatally androgenized androgen receptor knockout mice and neuron-specific androgen receptor knockout mice did not develop PCOS-like symptoms [7].

PCOS can be divided into adolescent- and adult-onset types according to the time of onset. During the pubertal maturation process of the hypothalamic-pituitary-ovarian axis, girls with a predisposition to PCOS may exhibit increased gonadotropin-releasing hormone and luteinizing hormone pulse frequencies, ultimately leading to an increased luteinizing hormone/follicle stimulating hormone ratio [8]. Some characteristics of PCOS, such as acne, irregular menses and polycystic ovary morphology, may also be normal physiological symptoms of puberty [9]. Puberty may activate the susceptibility to develop PCOS, which interferes with the normal pubertal transition. Other factors, such as elevated anti-Mullerian hormone secretion, luteinizing hormone secretion and gonadotropin-releasing hormone pulse frequency, may disrupt the pubertal transition [10]. Girls who do not establish a normal hypothalamic-pituitary-ovarian axis and mature along the PCOS trajectory may eventually develop persistent PCOS [11].

At present, the origin theory of PCOS is more inclined to the 'adolescence origin or developmental origin', in which adult-onset disease originates from maternal exposure and a high androgen environment or the abnormal establishment of hypothalamic-pituitary-ovarian axis feedback mechanisms in puberty. Refractory or medicine-resistant PCOS usually arises during puberty and is marked by insulin resistance, hyperinsulinemia and hyperandrogenism [12]. Many polygenic genetic diseases may originate from

maternal-derived abnormal genes in oocytes. Mouse models have long been used to study PCOS, yet despite many reports describing the reproductive and metabolic phenotype of PCOS mouse models, research on the oocyte itself remains sparse [13]. The incidence rate of PCOS is low in women older than 35 years, and adult-onset PCOS is more likely to be self-limited with aging or relieved by medical treatment. Most of the cases with drug resistant or familial-like PCOS develop their symptoms from puberty, and may be described as having adolescent-onset PCOS. The latter may be the type of PCOS that should be given more attention in the research and clinical setting. The aim of this study was to investigate the dynamic temporal changes of the oocyte transcriptome during PCOS modeling. Transcriptome data at six consecutive time points during PCOS mouse modeling were analyzed to detect the dynamic trend of gene regulation and reveal critical genes or pathways that may be important candidate factors for PCOS development and further studies.

2. Materials and Methods

2.1 Construction of the adolescent onset PCOS mouse model

The prepubertal mouse was used to mimic the hyperandrogenic phenotype of patients with adolescent onset PCOS. Female ICR mice were purchased from Shanghai JieSiJie Laboratory Animal Co. [production license number: SCXK (Shanghai) 2018-0004, China] and housed at the laboratory animal center of Obstetrics and Gynecology Hospital of Fudan University. Mice were housed with free access to food and water, and a constant temperature, humidity and light duration. At 25 days of age, mice were subcutaneously injected with DHEA powder (MCE, HY-14650, China) (6 mg/100 g body weight/d) dissolved in corn oil (MCE, HY-1888, China) (Figure1). All procedures were performed in accordance with the guidelines of Experimental Animals Management Regulations (version 2017-03-01, China).

Figure1. The schematics of the study design.

2.2 Ovary morphology and serum testosterone measurement

Mice were executed by carbon dioxide suffocation method. Ovaries of mice from each group were removed quickly and fixed in 4% paraformaldehyde for morphological assessment. Five-μm sections of ovary tissue were mounted on glass slides and stained with hematoxylin and eosin. Ovary morphology was further examined under light microscopy. The day after the final DHEA injection, blood was collected and centrifugated at a speed of 1000×g for 10 min. Serum testosterone was measured using a mouse testosterone enzyme immunoassay kit (Westang Bio-tech Co., Ltd., F11580, China).

2.3 Oocyte collection

From 25 days of age to two weeks after the final DHEA injection, oocytes were collected every other week so that six time points were established (points 0–5, Figure 1). At each time point, a mouse from each group was selected randomly, one ovary was fixed in paraformaldehyde for morphology assessment, and the other ovary was punctured with a 1 ml syringe needle in HEPES (Sigma, 83264, USA) to collect germinal vesicle-stage oocytes. Granulosa cells were removed mechanically. Ten oocytes of the same size, translucent and in good condition, were selected from each group for RNA sequencing.

2.4 Transcriptome sequencing

The selected oocytes were processed for RNA extraction and reverse transcription within 1 h and were subjected to cDNA amplification and purification. Reverse transcription was performed using a SMARTer Ultra Low RNA Kit (Clontech, USA) directly in the cell lysate. Then, cDNA was amplified and purified using an Advantage 2 PCR Kit (Clontech). The cDNA library was sequenced using an Illumina sequencing platform (NovaSeq6000). The raw reads were cleaned by removing adaptor sequences, short sequences (length < 35 bp), low-quality bases (quality < 20) and ambiguous sequences. Hisat2 (version:2.0.4) was used to map the cleaned RNA-seq reads to the mouse mm10 genome with two mismatches, two gaps and one multi-hit allowed. The gene expression value was normalized by fragments per kilobase of exon per million mapped fragments (FPKM) and adjusted by a geometric algorithm (All the transcriptome sequencing method was provided by Shanghai Biotechnology Corporation, China, Contract number E20210017-5).

2.5 Immunofluorescence staining of oocytes

Oocytes were denuded mechanically and fixed in 1% paraformaldehyde for 30min. Then, oocytes were transferred into 0.2% Triton X-100 in phosphate buffered saline (PBS) to permeabilize for 30 min at room temperature. After blocking in QuickBlock blocking buffer (Beyotime, P0260, China) for 15 min, oocytes were incubated with Anti-Ptges3 antibody (1:250, ab92503, UK) overnight at 4°C, followed by three washes (5 min each) with PBS the next day. Oocytes were then incubated with secondary antibody (1:500 dilution of Alexa Fluor 488 goat anti-rabbit antibody, abcam, ab150077) for 1 h at room temperature protected from

light. Oocytes were then mounted in a drop of antifade mounting medium (Beyotime, P0126) on a siliconized slide. Images were taken using a laser confocal microscope (Leica, Wetzlar, Germany).

2.6 Statistical Analysis

Data analysis was performed using the OmicShare tools, an open online platform for bioinformatics analysis (https://www.omicshare.com/tools, China) (Figure 2). At each time point (Point 1–5), two group of genes were compared to identify differentially expressed genes (DEGs). To further investigate dynamic gene expression patterns during the modeling process, trend analysis was used to determine genes with the same expression trend over six consecutive time points in the control and DHEA groups. Genes with very small expression levels were considered to play a limited role in the development of the disease, so the raw data were filtered by removing genes with FPKM < 1. Data of four main trends were extracted from each group and used for further analysis. Genes with the same or different expression trends from each group were compared for overlaps and differences, and the results were visualized using Venn diagrams. Genes with different or contrary expression trends from the Venn diagrams were considered to be abnormal and were used for gene pathway enrichment analysis (Kyoto Encyclopedia of Genes and Genomes, KEGG) (https://www.kegg.jp/)[14-16] and gene regulatory network analysis to explore the function of these genes and pathways. Genes from the critical pathway of each gene regulatory network were listed as hub genes. Data of relative grey value was analyzed in SPSS 19.0 (IBM Corp, USA). A value of $P < 0.05$ was considered   statistically significant.

Figure2. RNA-seq data analysis flowchart.

3. Results

3.1 Reproductive characteristics of PCOS mice

After three weeks of DHEA injection, the PCOS mouse model was successfully established (Figure 3A). Ovarian tissue histology showed large numbers of follicles at different stages of development in the control group, while ovaries in the PCOS group exhibited significantly increased cystic follicles with degenerate granulosa cell layers. Serum testosterone levels were significantly higher in the PCOS group versus healthy controls (Figure 3B). After the final DHEA injection, ovarian morphology gradually recovered to a normal state within 3 weeks (Figure 3C).

Figure3. Ovarian histology and serum testosterone. (a) HE staining of ovarian sections from both control and PCOS model (point3); (b) Serum testosterone level of both groups at time point3; (C) HE staining of ovarian sections from DHEA group after DHEA injection stopped for 1, 2 and 3 weeks.

3.2 DEGs at different time points

Data from two groups were compared to identify genes up- or down-regulated in the DHEA group at each of the five time points examined (points 1–5). The top ten DEGs are listed below and ranked by Log2FC (Table 1-4).

Table1. Top 10 DEGs Up-regulated in DHEA injection group

	Gene ID	Gene name	Log2FC	Pvalue
	ENSMUSG00000071072	Ptges3	2.57	2.51E-28
DHEA injection for	ENSMUSG00000027367	Stard7	2.43	1.70E-24
1 week	ENSMUSG00000018363	Smurf2	2.32	6.56E-21
(Point1)	ENSMUSG00000020847	Rph3al	2.20	3.55E-20
	ENSMUSG00000051579	Tceal8	2.16	6.18E-21

	ENSMUSG00000022419	Deptor	2.12	9.36E-20
	ENSMUSG00000038679	Trps1	2.09	1.15E-16
	ENSMUSG00000072889	Nfxl1	2.04	7.62E-19
	ENSMUSG00000013973	Dedd	2.03	1.87E-19
	ENSMUSG00000054312	Mrps21	1.93	8.37E-15
	ENSMUSG00000021361	Tmem14c	7.70	8.49E-130
	ENSMUSG00000034795	Ccdc122	3.43	1.30E-40
	ENSMUSG00000035506	Slc12a8	3.22	4.47E-37
DHEA injection for	ENSMUSG00000023032	Slc4a8	2.46	1.68E-29
2 weeks	ENSMUSG00000064357	mt-Atp6	1.96	1.69E-23
(Point2)	ENSMUSG00000069867	Pabpn1l	1.96	2.08E-22
	ENSMUSG00000051579	Tceal8	1.82	3.76E-19
	ENSMUSG00000018102	Hist1h2bc	1.79	9.15E-17
	ENSMUSG00000037058	Paip2	1.66	3.70E-18
	ENSMUSG00000022012	Enox1	1.56	3.26E-16
	ENSMUSG00000018362	Kpna2	5.71	1.07E-56
	ENSMUSG00000029392	Rilpl1	5.49	7.27E-89
	ENSMUSG00000060938	Rpl26	5.16	2.48E-62
	ENSMUSG00000035295	Wdr38	4.72	7.95E-44
DHEA injection for	ENSMUSG00000020077	Srgn	4.45	1.95E-42
3 weeks	ENSMUSG00000034853	Acot11	4.33	3.39E-41
(Point3)	ENSMUSG00000023965	Fbxl17	4.27	2.37E-42
	ENSMUSG00000069867	Pabpn1l	4.11	2.10E-64
	ENSMUSG00000057147	Dph6	3.79	1.34E-49
	ENSMUSG00000041453	Rpl21	3.57	8.05E-34

Table2. Top 10 DEGs Down-regulated in DHEA injection group

	Gene ID	Gene name	Log2FC	Pvalue
DHEA injection for	ENSMUSG00000024066	Xdh	-9.12	8.27E-91
1 week	ENSMUSG00000027871	Hsd3b1	-7.35	1.46E-79
(Point1)	ENSMUSG00000021361	Tmem14c	-6.90	3.25E-109
	ENSMUSG00000060032	H2afj	-6.40	2.29E-60

	ENSMUSG00000020522	Mfap3	-3.19	5.46E-32
	ENSMUSG00000067297	Ifit1bl2	-2.41	3.05E-26
	ENSMUSG00000071226	Cecr2	-2.36	2.60E-23
	ENSMUSG00000032968	Inha	-2.12	6.05E-22
	ENSMUSG00000015837	Sqstm1	-2.03	2.14E-17
	ENSMUSG00000073705	Apitd1	-2.00	2.98E-20
	ENSMUSG00000032968	Inha	-4.62	3.08E-67
	ENSMUSG00000041536	Serpina3a	-4.36	6.44E-39
	ENSMUSG00000079033	Mef2b	-4.21	1.28E-43
DHEA injection for	ENSMUSG00000055430	Nap1l5	-4.11	4.34E-43
2 weeks	ENSMUSG00000063229	Ldha	-3.82	1.29E-39
(Point2)	ENSMUSG00000036180	Gatad2a	-3.16	2.81E-26
	ENSMUSG00000035696	Rnf38	-3.00	1.52E-29
	ENSMUSG00000028646	Rragc	-2.97	4.31E-29
	ENSMUSG00000008348	Ubc	-2.83	1.89E-30
	ENSMUSG00000059743	Fdps	-2.64	9.96E-20
	ENSMUSG00000064363	mt-Nd4	-7.00	6.11E-104
	ENSMUSG00000020282	Rhbdf1	-5.87	7.83E-60
	ENSMUSG00000061371	Zfp873	-3.27	5.50E-34
	ENSMUSG00000057135	Scimp	-3.18	5.31E-32
DHEA injection for	ENSMUSG00000076613	Ighg2b	-3.14	1.64E-31
3 weeks	ENSMUSG00000021640	Naip1	-2.82	1.70E-30
(Point3)	ENSMUSG00000107743	RP23	-2.46	5.97E-27
	ENSMUSG00000027834	Serpini1	-2.44	3.19E-26
	ENSMUSG00000032359	Ctsh	-2.23	1.35E-22
	ENSMUSG00000043289	Mei4	-2.22	6.96E-21

Table3. Top 10 DEGs Up-regulated in DHEA group

	Gene ID	Gene name	Log2FC	Pvalue
DHEA injection	ENSMUSG00000018362	Kpna2	6.11	1.22E-58
stopped for 1 week	ENSMUSG00000029392	Rilpl1	4.94	2.34E-68
(Point4)	ENSMUSG00000024066	Xdh	4.71	8.46E-50

	ENSMUSG00000069867	Pabpn1l	4.41	5.87E-65
	ENSMUSG00000057147	Dph6	3.76	1.27E-46
	ENSMUSG00000023026	Dip2b	3.61	3.62E-32
	ENSMUSG00000100618	Gm29595	3.31	1.22E-33
	ENSMUSG00000041840	Haus1	3.24	8.28E-37
	ENSMUSG00000035696	Rnf38	3.12	5.47E-35
	ENSMUSG00000042712	Wbp5	2.93	2.31E-26
	ENSMUSG00000019947	Arid5b	6.52	6.00E-58
	ENSMUSG00000034324	Tmem132c	5.98	2.49E-49
	ENSMUSG00000029392	Rilpl1	5.76	2.63E-85
	ENSMUSG00000023932	Cdc5l	5.07	7.39E-49
DHEA injection stopped for 2 weeks (Point5)	ENSMUSG00000032968	Inha	4.94	6.69E-60
	ENSMUSG00000041453	Rpl21	4.81	3.89E-45
	ENSMUSG00000034853	Acot11	4.67	6.39E-48
	ENSMUSG00000056116	H2-T22	4.43	1.42E-44
	ENSMUSG00000000184	Ccnd2	4.40	5.64E-38
	ENSMUSG00000020077	Srgn	3.85	4.57E-38

Table4. Top 10 DEGs Down-regulated in DHEA group

	Gene ID	Gene name	Log2FC	Pvalue
	ENSMUSG00000058147	Xlr3c	-2.86	1.64E-41
	ENSMUSG00000073125	Xlr3b	-1.73	2.79E-19
	ENSMUSG00000048701	Ccdc6	-1.59	7.62E-18
	ENSMUSG00000020870	Cdc34-ps	-1.54	7.08E-16
DHEA injection stopped for 1 week (Point4)	ENSMUSG00000068823	Csde1	-1.52	1.72E-16
	ENSMUSG00000020962	Gtf2a1	-1.49	4.29E-15
	ENSMUSG00000093930	Hmgcs1	-1.43	4.55E-15
	ENSMUSG00000071089	Trim75	-1.37	5.44E-13
	ENSMUSG00000021488	Nsd1	-1.34	1.38E-12
	ENSMUSG00000024077	Strn	-1.26	2.11E-12
DHEA injection stopped for 2 weeks	ENSMUSG00000076613	Ighg2b	-7.01	2.71E-93
	ENSMUSG00000076615	Ighg3	-6.44	1.15E-64

(Point5)	ENSMUSG00000026276	2-Sep	-5.03	6.39E-59
	ENSMUSG00000022661	Cd200	-4.77	1.73E-56
	ENSMUSG00000057135	Scimp	-4.44	6.97E-51
	ENSMUSG00000035506	Slc12a8	-4.07	5.97E-59
	ENSMUSG00000026205	Slc23a3	-3.84	2.49E-57
	ENSMUSG00000027367	Stard7	-3.43	9.48E-44
	ENSMUSG00000035455	Fignl1	-2.88	1.45E-34
	ENSMUSG00000107743	RP23	-2.77	1.50E-37

3.3 Trend analysis

A total of 21,928 genes were identified in mouse oocytes, and 9,382 genes had FPKM > 1. Given that each gene had a different trend over time, data in each group was first analyzed through trend analysis to summarize genes with similar trends over time. Four main trends of data were selected for further analysis. In the control group, analysis over six consecutive time points (points 0–5) showed 314 genes were continuously up-regulated, 525 genes were continuously down-regulated, 717 genes had a down-flat-up regulation trend and 859 genes had an up-flat-down regulation trend (Figure 4A). In the DHEA group over the same time points, 1491 genes were continuously up-regulated, 811 genes were continuously down-regulated, 1222 genes showed a down-flat-up regulation trend and 334 genes showed an up-flat-down regulation trend (Figure 4B).

Figure4. Trend analysis of each group. (a) In the control group, 314 genes were continuously up-regulated, 525 genes were continuously down-regulated, 717 genes had a down-flat-up r egulation trend and 859 genes had an up-flat-down regulation

228 trend. (**b**) In the DHEA group, 1491 genes were continuously up-regulated, 811 genes were continuously down-regulated,

229 1222 genes showed a down-flat-up regulation trend and 334 genes showed an up-flat-down regulation trend.

3.4 Venn diagram analysis

Four selected trends of data were plotted as Venn diagrams to identify genes with abnormal expression

trends in the DHEA group. For genes with the same trend in both groups, genes only expressed in the

DHEA group were selected. For genes with a contrary trend, overlapping genes of the two groups were

selected. Six kinds of data combinations and Venn diagrams were finally produced. Venn1 was produced

with genes expressed as 'trend1' in both the control and DHEA groups (Figure 5A). In this diagram, 1372

genes were abnormally continuously up-regulated in the DHEA group (Figure 5B). Venn2 was produced

with genes expressed as 'trend2' in both the control and DHEA groups (Figure 6A), with 743 genes

abnormally down-regulated in the DHEA group(Figure 6B). Venn3 was produced with genes expressed as

'trend1' in the control group and genes expressed as 'trend2' in the DHEA group (Figure 7A). Twenty genes

were abnormally down-regulated in the DHEA group (Figure 7B). Venn4 was produced with genes

expressed as 'trend2' in the control group and genes expressed as 'trend1' in the DHEA group (Figure 8A).

Seventy-one genes were abnormally up-regulated in the DHEA group (Figure 8B). Venn5 was produced

with genes expressed as 'trend3' in both groups (Figure 9A), in which 967 genes were abnormally

down-flat-up-regulated in the DHEA group (Figure 9B). Venn6 was produced with genes expressed as

'trend4' in both groups (Figure 10A), in which 190 genes were abnormally up-flat-down-regulated in the

PCOS group (Figure 10B). The selected genes are marked with the red dotted circle.

Figure5. Analysis of genes expressed as 'trend1' in both groups. (a) 314 and 1491 genes were continuously up-regulated in each group separately. (b) Venn1 diagram showed 1372 genes were abnormally up-regulated in DHEA group. (c) KEGG analysis of the 1372 genes abnormally up-regulated in DHEA group. (d) Gene regulatory network diagram of the 1372 g

Figure6. Analysis of genes expressed as 'trend2' in both groups. (a) 525 and 811 genes were continuously down-regulated in each group separately. (b) Venn2 diagram showed 743 genes were abnormally up-regulated in DHEA group. (c) KEGG analysis of the 743 genes abnormally down-regulated in DHEA group. (d) Gene regulatory network diagram of the 743 genes abnormally down-regulated in DHEA group.

Figure7. Analysis of genes expressed as 'trend1' in the control group and genes expressed as 'trend2' in the DHEA group. (a) 314 genes were up-regulated in control group and 811 genes were down-regulated in DHEA group. (b) Venn3 diagram showed 20 genes were abnormally down-regulated in DHEA group. (c) KEGG analysis of the 20 genes abnormally down-regulated in DHEA group. (d) Gene regulatory network diagram of the 20 genes abnormally down-regulated in DHEA group.

Figure8. Analysis of genes expressed as 'trend2' in the control group and genes expressed as 'trend1' in the DHEA group. (a) 525 genes were down-regulated in control group and 1491 genes were up-regulated in DHEA group. (b) Venn4 diagram showed 71 genes were abnormally up-regulated in DHEA group. (c) KEGG analysis of the 71 genes abnormally up-regulated in DHEA group. (d) Gene regulatory network diagram of the 71 genes abnormally up-regulated in DHEA group

Figure9. Analysis of genes expressed as 'trend3' in both groups. (a) 717 genes and 1222 genes were down-flat-up-regulated in each group. (b) 967 genes were abnormally down-flat-up-regulated in DHEA group. (c) KEGG analysis of the 967 genes abnormally down-flat-up-regulated in DHEA group. (d) Gene regulatory network diagram of the 967 genes abnormally down-flat-up-regulated in DHEA group.

Figure10. Analysis of genes expressed as 'trend4' in both groups. (a) 859 genes and 334 genes were up-flat-down-regulated in each group. (b) 190 genes were abnormally up-flat-down-regulated in DHEA group. (c) KEGG analysis of the 190 genes abnormally up-flat-down-regulated in DHEA group. (d) Gene regulatory network diagram of the 190 genes abnormally up-flat-down-regulated in DHEA group.

3.5 Genes clustering and KEGG analysis

KEGG analysis was used to detect the biological functions of the above six groups of selected genes. Genes abnormally up-regulated in the DHEA group were mainly focused on oxidative phosphorylation, thermogenesis, and the Parkinson disease pathway (Figure 5-10C). Genes abnormally down-regulated in the DHEA group were mainly focused on the cell cycle pathway. Genes abnormally up-regulated in the control group but down-regulated in the DHEA group were mainly focused on p53 signaling, PI3K-Akt, and glycolysis gluconeogenesis pathways. Genes down-regulated in the control group but up-regulated in the DHEA group were mainly focused on oxidative phosphorylation, central carbon metabolism in cancer, and the Alzheimer's disease pathway. Genes abnormally down-flat-up-regulated in the DHEA group were mainly focused on the regulation of actin cytoskeleton pathway. Genes abnormally up-flat-down-regulated in the DHEA group were mainly focused on oxidative phosphorylation, apoptosis and Huntington disease

pathways. The above results suggested that the most important pathways were oxidative phosphorylation and the cell cycle.

3.6 Gene regulatory network and hub genes

The selected genes from Venn diagrams were further investigated using gene regulatory network analysis to define the weight of different pathways. The result shows that pathways with the highest weight and strongest correlation were also oxidative phosphorylation and the cell cycle (Figure 5-10D). Some representative genes from a critical pathway were listed as potential hub genes (Table 5) and ranked by FPKM. The most significant difference was focused on mitochondrial oxidative phosphorylation. Hub genes abnormally up-regulated in the DHEA group included mtDNA coding subunits (mt-Nd1, 2, 3, 6 and mt-Co1,2), cytochrome oxidase (Cox8a, 7a2l, and 4i1), NADH-ubiquinone oxidoreductase (Ndufa1, 2, 6; Ndufb 9), and Uqcrq. Hub genes abnormally down-regulated in the DHEA group mainly focused on the cell cycle, which included Skp1, Ccnb1, origin recognition complex (ORC) subunit 1, and 5, Wee2, Mapk3, and Cdc20. All the other specific genes are listed in the supplementary documents (Table Venn1-6).

3.7 Immunofluorescence staining of Ptges3

According to the results above, Ptges3 was the top1 DEGs at time point1(Table1). The relative fluorescence intensity level of Ptges3 was significantly higher in DHEA group versus control oocytes at time point1(Figure11), which is consistent with the RNA-seq result.

Figure11. Immunofluorescence staining of Ptges3 in oocytes from control and DHEA group at time point1

4. Discussion

This study performed transcriptome sequencing throughout the course of PCOS development in a mouse model of adolescent PCOS to find the critical genes or pathways that regulate oocyte growth and quality. Identified DEGs in oocytes from the PCOS model were mainly focused on oxidative phosphorylation and the cell cycle.

Some mitochondria-related genes were abnormally and consistently up-regulated in the DHEA group, even after the DHEA injection regime was completed. Mt-Nd-x are key protein subunits of mitochondrial complex I, which is the main entrance for electrons to the respiratory chain, and its dysfunction will lead to decreased ATP production [17]. Mitochondrial complex I deficiency is associated with many clinical conditions, such as Leber hereditary optic neuropathy, mitochondrial encephalomyelopathy, and lactic acidosis [18]. Cytochrome c oxidase (COX) is the terminal enzyme of electron transport chains in mitochondria, and COX8a is required for maintenance of the structural stability of COX monomers and dimers [19]. COX7a2l is one of the three COX7a subunits, responsible for organization of the mitochondrial respiratory chain and preventing metabolic exhaustion [20]. Uqcrq is one of the ten nuclear genes encoding proteins of mitochondrial complex III, and mutations in Uqcrq are associated with a series of rare neuromuscular symptoms [21]. NDUFA-x encodes accessory subunits of mitochondrial complex I, and deficiency of this complex is the most common defect of the oxidative phosphorylation system. Mutations in NDUFA2 cause leukoencephalopathy [22], and patients with Leigh syndrome were found to harbor a homozygous mutation in NDUFA1[23] . Diseases caused by mitochondria deficiency may have some undetermined commonalities with PCOS, and the down regulation of mitochondrially encoded subunits of respiratory chain complexes may play an important role in the mitochondria dysfunction associated with PCOS [24].

Mitochondrial dysfunction has been proven to be a vital factor of many cardiovascular diseases and also plays an important role in the pathogenesis of PCOS [25]. Mitochondria convert nutrients into energy and release reactive oxygen species, but high levels of reactive oxygen species cause damage to mitochondrial components or even induce apoptosis. Growing evidence indicates that patients with PCOS have an increased level of oxidative stress and decreased levels of antioxidants [26]. The abnormal concentration of reactive oxygen species and antioxidant factors in follicular fluid from patients with PCOS may have adverse effects on oocyte quality, fertilization and embryo development [27]. mtDNA copy number is lower in women with PCOS and is negatively correlated with insulin resistance levels while positively associated with sex hormone-binding globulin levels [28]. However, some studies suggested that mtDNA number was increased in patients with PCOS and this increase was regarded as a feedback response compensating for mitochondrial dysfunction. These paradoxical results may reflect different stages of disease manifestation. mtDNA copy number increases during the early stage of PCOS as a compensatory effect, but decreases with disease development as the mitochondria fail to sustain normal function [29].

The markedly down-regulated genes in the DHEA group mainly focused on cell cycle. S-phase Kinase-Associated Protein1 (SKP1) is a core component of the Skp, Cullin, F-box containing complex, an E3 ubiquitin ligase that participates in a variety of cellular processes [30]. It was reported that reduced SKP1 expression induced chromosomal instability in both benign and malignant tumors [31]. Furthermore, SKP1 maintained synapsis in meiosis of both sexes [32], and also affected the balance of epigenetic regulation [33]. ORC proteins can bind DNA replication origins to initiate DNA synthesis, and ORC is especially important in controlling centriole and centrosome copy number [34]. Wee2 is an important oocyte-specific kinase, and its downregulation in oocytes resulted in high maturation-promoting factor activity and failure to mature to the meiosis-II stage [35]. Cyclin B1 is responsible for the composition of maturation-promoting factor, and Ccnb1-null oocytes completed meiosis I but arrested at the meiotic interphase [36]. Dominant or recessive CDC20 mutations have been linked to adverse oocyte maturation, and may act as a potential molecular markers of oocyte/embryo quality [37]. The relationship between all the hub genes mentioned above and PCOS has not been investigated until now. From these findings we propose that infertility or adverse pregnancy outcomes of patients with PCOS may result from mitochondrial dysfunction or abnormal meiotic regulation of oocytes.

At the first week of modeling, we found that Ptges3 was the top1 DEGs which was up-regulated in DHEA injection group at time point1. Ptges3 can act as an androgen receptor chaperone, which is present in many Hsp90 chaperone complexes and involved in assembling and disassembling some hormone or nuclear receptor/enhancer complexes[38]. Ptges3 and Hsp90 are of the basic minimal complexes required for the efficient folding and stabilization of steroid hormone receptors and essential for ligand responsive signaling[39]. In our research, Ptges3 was found up-regulated at the initial stage of modeling, which suggested it may play an important role in the early stage of PCOS.

5. Conclusion

This study provides a novel insight into differences in the transcriptomes of oocytes from control compared with PCOS mice. mtDNA-related genes and Cell cycle-related genes play the most important role in the development of PCOS. These findings could help focus on specific pathways or potential candidate genes that may play important roles during the development of adolescent onset PCOS. Ptges3 was the top1 DEGs which was up-regulated in DHEA group at the initial stage of modeling, which suggested it may play an important role in the early stage of PCOS.

(Figure 7B, marked by red dotted circle); Table Venn3: Genes abnormally down-regulated in the DHEA group (Figure 8B, marked by red dotted circle); Table Venn1 Venn4: Genes abnormally up-regulated in the DHEA group (Figure 9B, marked by red dotted circle); Table Venn5: Genes abnormally down-flat-up-regulated in the DHEA group (Figure 10B, marked by red dotted circle); Table Venn6: Genes were abnormally up-flat-down-regulated in the PCOS group (Figure 11B, marked by red dotted circle).

Author Contributions:

1.Du Danfeng (First Author): Methodology, Software, Investigation, Formal Analysis, Writing-Original Draft.

ORCID: 0000-0002-0753-334X

Email: 693468525@qq.com

2.Deng Ke (Co-First author): Data Curation, Investigation, Formal Analysis.

ORCID: 0000-0002-4098-1144

Email：20111250023@fudan.edu.cn

3.Fan Dengxuan: Resources, Supervision.

Email：dxfan16@fudan.edu.cn

4.Xu Congjian (Corresponding Author): Conceptualization, Funding Acquisition, Resources, Supervision.

ORCID: 0000-0002-7954-5892

Email: xucongjian@fudan.edu.cn

Funding:

This work was supported by funding from the National Natural Science Foundation of China to Xu Congjian (Grant number 82171639 and 82201807).

Ethic Statement:

All animal procedures were performed in accordance with the guidelines of Experimental Animals Management Regulations (version 2017-03-01, China).

Ethics approval was granted by the Institutional Animal Welfare and Ethics Committee Policies of Fudan University（approval number: 2020 Obstetrics and Gynecology Hospital JS-011 to Du Danfeng）.

Data Availability Statement:

Raw data underlying this article are available in Figshare (https://figshare.com/). The authors affirm that all data necessary for confirming the conclusions of the article are present within the article, figures, and tables. Data analysis was performed using the OmicShare tools (https://www.omicshare.com/tools)

Acknowledgments:

We thank Liwen Bianji (Edanz) (www.liwenbianji.cn/) for editing the English text of a draft of this manuscript.

Conflicts of Interest:

The authors declare no conflict of interest.

References

1. Qiao, J. and H.L. Feng, *Extra- and intra-ovarian factors in polycystic ovary syndrome: impact on oocyte maturation and embryo developmental competence.* Hum Reprod Update, 2011. **17**(1): p. 17-33.

2. Huang, Y., et al., *Impaired oocyte quality induced by dehydroepiandrosterone is partially rescued by metformin treatment.* PLoS One, 2015. **10**(3): p. e0122370.

3. Crespo, R.P., et al., *An update of genetic basis of PCOS pathogenesis.* Arch Endocrinol Metab, 2018. **62**(3): p. 352-361.

4. Khan, M.J., A. Ullah, and S. Basit, *Genetic Basis of Polycystic Ovary Syndrome (PCOS): Current Perspectives.* Appl Clin Genet, 2019. **12**: p. 249-260.

5. Lizneva, D., et al., *Criteria, prevalence, and phenotypes of polycystic ovary syndrome.* Fertil Steril, 2016. **106**(1): p. 6-15.

6. McCarthy, M.M., K. Herold, and S.L. Stockman, *Fast, furious and enduring: Sensitive versus critical periods in sexual differentiation of the brain.* Physiol Behav, 2018. **187**: p. 13-19.

7. Caldwell, A.S.L., et al., *Neuroendocrine androgen action is a key extraovarian mediator in the development of polycystic ovary syndrome.* Proc Natl Acad Sci U S A, 2017. **114**(16): p. E3334-E3343.

8. Rothenberg, S.S., et al., *Polycystic ovary syndrome in adolescents.* Best Pract Res Clin Obstet Gynaecol, 2018. **48**: p. 103-114.

9. Witchel, S.F., H. Roumimper, and S. Oberfield, *Polycystic Ovary Syndrome in Adolescents.* Endocrinol Metab Clin North Am, 2016. **45**(2): p. 329-44.

10. Witchel, S.F. and T.M. Plant, *Intertwined reproductive endocrinology: Puberty and polycystic ovary syndrome.* Curr Opin Endocr Metab Res, 2020. **14**: p. 127-136.

11. Pena, A.S., et al., *Adolescent polycystic ovary syndrome according to the international evidence-based guideline.* BMC Med, 2020. **18**(1): p. 72.

12. Otto-Buczkowska, E., K. Grzyb, and N. Jainta, *Polycystic ovary syndrome (PCOS) and the accompanying disorders of glucose homeostasis among girls at the time of puberty.* Pediatr Endocrinol Diabetes Metab, 2018. **24**(1): p. 40-44.

13. Chappell, N.R., et al., *Prenatal androgen induced lean PCOS impairs mitochondria and mRNA profiles in oocytes.* Endocr Connect, 2020.

14. Kanehisa, M., et al., *KEGG for taxonomy-based analysis of pathways and genomes.* Nucleic Acids Res, 2023. **51**(D1): p. D587-D592.

15. Kanehisa, M., *Toward understanding the origin and evolution of cellular organisms.* Protein Sci, 2019. **28**(11): p. 1947-1951.

16. Kanehisa, M. and S. Goto, *KEGG: kyoto encyclopedia of genes and genomes.* Nucleic Acids Res, 2000. **28**(1): p. 27-30.

17.   Formosa, L.E., et al., *Building a complex complex: Assembly of mitochondrial respiratory chain complex I.* Semin Cell Dev Biol, 2018.
76: p. 154-162.

18.   Ng, Y.S., et al., *Novel MT-ND Gene Variants Causing Adult-Onset Mitochondrial Disease and Isolated Complex I Deficiency.* Front
Genet, 2020. **11**: p. 24.

19.   Rotko, D., et al., *Molecular and Functional Effects of Loss of Cytochrome c Oxidase Subunit 8A.* Biochemistry (Mosc), 2021. **86**(1): p.
33-43.

20.   Fernandez-Vizarra, E., et al., *Two independent respiratory chains adapt OXPHOS performance to glycolytic switch.* Cell Metab, 2022.

21.   Burska, D., et al., *Homozygous missense mutation in UQCRC2 associated with severe encephalomyopathy, mitochondrial complex III*
assembly defect and activation of mitochondrial protein quality control. Biochim Biophys Acta Mol Basis Dis, 2021. **1867**(8): p.
166147.

22.   Perrier, S., et al., *Recessive mutations in NDUFA2 cause mitochondrial leukoencephalopathy.* Clin Genet, 2018. **93**(2): p. 396-400.

23.   Huttula, S., et al., *NDUFA1 p.Gly32Arg variant in early-onset dementia.* Neurobiol Aging, 2022. **114**: p. 113-116.

24.   Thompson, L.L., et al., *Reduced SKP1 Expression Induces Chromosome Instability through Aberrant Cyclin E1 Protein Turnover.*
Cancers (Basel), 2020. **12**(3).

25.   Papalou, O., V.M. Victor, and E. Diamanti-Kandarakis, *Oxidative Stress in Polycystic Ovary Syndrome.* Curr Pharm Des, 2016. **22**(18):
p. 2709-22.

26.   Sulaiman, M.A., et al., *Polycystic ovarian syndrome is linked to increased oxidative stress in Omani women.* Int J Womens Health,
2018. **10**: p. 763-771.

27.   Nunez-Calonge, R., et al., *Oxidative stress in follicular fluid of young women with low response compared with fertile oocyte donors.*
Reprod Biomed Online, 2016. **32**(4): p. 446-56.

28.   Lewis, J.G., et al., *Plasma sex hormone-binding globulin, corticosteroid-binding globulin, cortisol, and free cortisol levels in*
outpatients attending a lipid disorders clinic: a cross-sectional study of 1137 subjects. Horm Metab Res, 2010. **42**(4): p. 274-9.

29.   Zhang, J., et al., *Polycystic ovary syndrome and mitochondrial dysfunction.* Reprod Biol Endocrinol, 2019. **17**(1): p. 67.

30.   Thompson, L.L., et al., *Aberrant SKP1 Expression: Diverse Mechanisms Impacting Genome and Chromosome Stability.* Front Cell
Dev Biol, 2022. **10**: p. 859582.

31.   Lepage, C.C., et al., *Reduced SKP1 and CUL1 expression underlies increases in Cyclin E1 and chromosome instability in cellular*
precursors of high-grade serous ovarian cancer. Br J Cancer, 2021. **124**(10): p. 1699-1710.

32.   Guan, Y., et al., *SKP1 drives the prophase I to metaphase I transition during male meiosis.* Sci Adv, 2020. **6**(13): p. eaaz2129.

33.   Shen, J. and C. Spruck, *F-box proteins in epigenetic regulation of cancer.* Oncotarget, 2017. **8**(66): p. 110650-110655.

34.   Ortega, M.A., H. Nguyen, and W.S. Ward, *ORC proteins in the mammalian zygote.* Cell Tissue Res, 2016. **363**(1): p. 195-200.

35.   Sang, Q., et al., *Homozygous Mutations in WEE2 Cause Fertilization Failure and Female Infertility.* Am J Hum Genet, 2018. **102**(4):
p. 649-657.

36.   Li, J., W.P. Qian, and Q.Y. Sun, *Cyclins regulating oocyte meiotic cell cycle progressiondagger.* Biol Reprod, 2019. **101**(5): p.
878-881.

37.   Sang, Q., et al., *Genetic factors as potential molecular markers of human oocyte and embryo quality.* J Assist Reprod Genet, 2021.
38(5): p. 993-1002.

38.   Nguyen, P.M., et al., *p23 co-chaperone protects the aryl hydrocarbon receptor from degradation in mouse and human cell lines.*
Biochem Pharmacol, 2012. **84**(6): p. 838-50.

39.   Biebl, M.M. and J. Buchner, *p23 and Aha1: Distinct Functions Promote Client Maturation.* Subcell Biochem, 2023. **101**: p. 159-187.

570