The locus coeruleus broadcasts prediction errors across the cortex to promote sensorimotor plasticity

Rebecca Jordan[1,3,4] & Georg B. Keller[1,2]

[1]*Friedrich Miescher Institute for Biomedical Research, Basel, Switzerland.*
[2]*Faculty of Sciences, University of Basel, Basel, Switzerland.*
[3]*Current address: Simons Initiative for the Developing Brain, University of Edinburgh, 1 George Square,*
EH8 9JZ, Edinburgh, United Kingdom.
[4]*Correspondence to:* rjordan3@ed.ac.uk

ABSTRACT

Prediction errors are differences between expected and actual sensory input and are thought to be key computational signals that drive learning related plasticity. One way that prediction errors could drive learning is by activating neuromodulatory systems to gate plasticity. The catecholaminergic locus coeruleus (LC) is a major neuromodulatory system involved in neuronal plasticity in the cortex. Using two-photon calcium imaging in mice exploring a virtual environment, we found that the activity of LC axons in the cortex correlated with the magnitude of unsigned visuomotor prediction errors. LC response profiles were similar in both motor and visual cortical areas, indicating that LC axons broadcast prediction errors throughout the dorsal cortex. While imaging calcium activity in layer 2/3 of the primary visual cortex, we found that optogenetic stimulation of LC axons facilitated learning of a stimulus-specific suppression of visual responses during locomotion. This plasticity – induced by minutes of LC stimulation – recapitulated the effect of visuomotor learning on a scale that is normally observed during visuomotor development across days. We conclude that prediction errors drive LC activity, and that LC activity facilitates sensorimotor plasticity in the cortex, consistent with a role in modulating learning rates.

*** Dear reader, please note this manuscript is formatted in a standard submission format, and all statistical information is in Table S1. ***

INTRODUCTION

Through experience with the world, brains learn to predict the sensory feedback generated by movement. For instance, learning the precise relationship between motor commands and the resulting visual

feedback is the basis for both feedback control of movements as well as distinguishing self-generated

from externally generated sensory feedback. One way that the brain may learn to do this is via predictive

processing (Keller and Mrsic-Flogel, 2018; Rao and Ballard, 1999). In a sensorimotor version of this

framework, the brain learns an internal model which transforms motor activity into a prediction of the

sensory consequences. The framework postulates that specific neurons are responsible for computing

discrepancies – known as prediction errors – between the prediction and the actual sensory inputs

received. The neocortex has been shown to compute signals consistent with sensorimotor prediction

errors (Eliades and Wang, 2008; Keller and Mrsic-Flogel, 2018; Schneider et al., 2018). In the primary visual

cortex (V1), excitatory neurons in layer 2/3 compute mismatches between visual flow speed and

locomotion speed (Attinger et al., 2017; Jordan and Keller, 2020; Keller et al., 2012; Zmarz and Keller,

2016). Such sensorimotor prediction error responses in the cortex depend on ongoing and developmental

experience with sensorimotor coupling (Attinger et al., 2017; Schneider et al., 2018; Vasilevskaya et al.,

2022; Widmer et al., 2022), and therefore the computation underlying them is actively learned. In

predictive processing, one main function of prediction errors is to drive corrective plasticity in the internal

model that generates predictions (Keller and Mrsic-Flogel, 2018).

During the first sensorimotor experience in life, prediction errors should strongly drive learning of an

internal model of the world, while later in life, prediction errors should update internal models primarily

when overall prediction errors are abundant (for instance, in novel or volatile situations). One way to

implement such a developmental or contextual shift in the amount of plasticity induced by a particular

prediction error would be a gating signal. The characteristics of a gating signal are that it is activated by

prediction errors and that it permits increases in plasticity. In addition, because it is not driving plasticity,

but simply gating it, it can be low dimensional (i.e., the same for all neurons). Prime candidates for a gating

signal are neuromodulatory systems. Dopaminergic neurons in the midbrain signal reward prediction

errors (Kim et al., 2020; Schultz, 2016) and action prediction errors (Greenstreet et al., 2022), which are

thought to gate plasticity in downstream targets to support learning of stimulus-reward associations and

goal directed behavior (Cox and Witten, 2019; Flagel et al., 2011; Reynolds et al., 2001; Steinberg et al.,

2013). Indeed, the catecholamines – dopamine and noradrenaline – regulate and gate plasticity in the

cortex (Choi et al., 2005; He et al., 2015; Seol et al., 2007). However, the midbrain dopaminergic system

projects only sparsely to most of the sensory neocortex of the mouse, including V1 (Nomura et al., 2014),

where visuomotor predictions are thought to be learned (Attinger et al., 2017; Widmer et al., 2022).

Instead, the major catecholaminergic input to the cortex is the locus coeruleus (LC). LC activity has

primarily been investigated in the context of classical stimulus presentations and reinforcement tasks.

Such studies have shown that the LC responds to many different events, including unexpected task

outcomes (Bouret and Sara, 2004; Breton-Provencher et al., 2022), novelty (Takeuchi et al., 2016; Vankov

et al., 1995), and unexpected sensory stimuli in general (Aston-Jones and Bloom, 1981; Deitcher et al.,

2019; Foote et al., 1980; Hervé-Minvielle and Sara, 1995). In zebrafish, noradrenergic neurons increase

their activity during the prolonged visuomotor mismatches of failed swim attempts to drive 'giving up'

behavior (Mu et al., 2019). We hypothesized that if the mouse LC is activated during similar visuomotor

mismatches, this could represent visuomotor prediction error signaling that would function to gate

prediction error driven plasticity in output circuits like the cortex.

To test fundamental aspects of this idea, we imaged calcium activity in LC axons in the cortex during

visuomotor mismatches, and optogenetically manipulated LC activity levels during visuomotor experience

to probe for experience dependent changes in cortical activity. We found that LC axons in both sensory

and motor cortical regions respond to unsigned visuomotor prediction errors, i.e., unpredicted visual

motion or unpredicted visual halts during locomotion. We then combined optogenetic LC axon stimulation

with two-photon calcium imaging of layer 2/3 neurons in V1, to show that the LC greatly facilitates a form

of plasticity consistent with predictive visuomotor learning: learned suppression of nasotemporal visual

flow responses during forward locomotion. These results support the idea that a key function of the LC is

to facilitate prediction error driven cortical plasticity.

RESULTS

We first assessed whether locus coeruleus (LC) axons convey sensorimotor prediction errors to the cortex.

To record LC axonal activity, we expressed axon-targeted, Cre-dependent GCaMP6s in noradrenergic

neurons of the LC via stereotactic viral vector injections in 13 NET-Cre mice (NET = norepinephrine

transporter) (Wagatsuma et al., 2018). We confirmed that these injections labelled only cells in the LC,

and that GCaMP6s positive cells were also immunoreactive for tyrosine hydroxylase, a marker for

catecholaminergic neurons (**Figure S1A**). For two-photon imaging, we implanted a 4 mm imaging window

either over the right V1 and surrounding cortical regions (posterior cortex, 9 mice), or over the right

primary and secondary motor cortices (anterior cortex, 4 mice). Prior to imaging, mice were habituated

to a virtual reality (VR) system consisting of a virtual tunnel, with walls patterned with vertical sinusoidal

gratings (**Figure 1A**). In the closed loop condition, locomotion was coupled to visual movement of the

tunnel, and in open loop conditions, the walls moved independent of mouse locomotion. After

habituation to head-fixation, we began imaging across up to five non-overlapping fields of view (FoV) in

each imaging window, yielding a total of 31 posterior cortex FoVs and 9 anterior cortex FoVs. Axonal

fluorescence showed strong correlations with pupil diameter (**Figure S1B**) typical for LC axons (Reimer et

al., 2016), rapid responses to air puffs (**Figure S1C**) as previously reported (Breton-Provencher et al., 2022;

Deitcher et al., 2019), and increases in fluorescence at locomotion onset that decayed during the

locomotion bout (**Figure S2**) consistent with earlier work (Reimer et al., 2016). We found that within a

FoV, individual axon segments exhibited highly correlated activity (**Figures 1B and S1D-F**). Given that the

algorithms that identify axons (i.e., determine whether two axon segments belong to the same axon) rely

on temporal correlations of activity (Leinweber et al., 2017; Mukamel et al., 2009), axon segmentation is

difficult in these data. Thus, to prevent oversampling of the data, we pooled data from all axon segments

within a FoV for further analysis.

Locus coeruleus cortical axons signal unsigned visuomotor errors

To determine whether LC axons respond to visuomotor errors, we first measured their responses to

visuomotor mismatch stimuli (unexpected 1 s halts of visual flow during locomotion in the closed loop

condition). Indeed, LC axons showed significant increases in fluorescence following mismatch

presentations (**Figure 1C-D**). Assuming these responses reflect visuomotor prediction errors, we would

expect to find two things:

First, the responses should depend on precise coupling between visual flow speed and locomotion speed

(i.e., when visual flow is predictable from locomotion). Responses to visuomotor mismatch in closed loop

conditions should be higher than responses to a replay of the same visual flow halt (termed 'playback

halt') in open loop conditions, even when the mouse is locomoting. Playback halts are visually identical to

mismatches but occur independent of whether the mouse is stationary or locomoting (**Figure 1A**).

Responses of LC axons to playback halts when the mouse was stationary were negative and substantially

different from mismatch responses (**Figure 1C-D**), demonstrating that the response of LC axons to

visuomotor mismatch cannot be explained by the visual stimulus alone. The mismatch responses also

could not be explained by a locomotion driven increase in gain of the visual halt response: a) The average

playback halt response when the mouse was stationary was negative, b) there was no correlation between

playback halt responses and mismatch responses across different FoVs ($R^2 = 0.04$, p = 0.31, 25 FoVs, linear

regression), and c) mismatch responses were significantly larger than playback halt responses that

occurred during locomotion in the open loop condition (**Figure 1D**). This indicated that responses in LC

axons were acutely dependent on visuomotor coupling, since the only difference between a mismatch

stimulus and a playback halt during locomotion is that the former occurs during visuomotor coupling and

the latter does not.

Second, the mismatch response should scale with the size of the error between visual flow speed and

locomotion speed, since the latter is a proxy for the prediction of visual flow speed. Since visual flow speed

is zero during mismatch, and visual flow is perfectly coupled to locomotion prior to mismatch, the

visuomotor error is proportional to the locomotion speed during mismatch (**Figure 1A**). Indeed, we found

that LC axonal mismatch responses increased monotonically with the locomotion speed of the mouse

(**Figure 1E**). Thus, LC axonal responses to visuomotor mismatch are consistent with a visuomotor

prediction error signal.

Visuomotor mismatches evoke negative prediction errors because they are a condition where there is *less*

visual flow than expected. However, these are not the only type of prediction error in our paradigm. In

the open loop condition, visual flow onsets result in positive prediction errors (i.e., a condition with *more*

visual flow than expected), especially when the mouse is stationary. We thus assessed the responses of

LC axons to visual flow onsets in the open loop condition (i.e., uncoupled from locomotion). We found

that LC axons showed significant responses to visual flow onsets, but only when the mouse was stationary

(**Figure 1F-G**). Visual flow onsets during locomotion did not evoke significant responses on average.

Altogether, LC axonal responsiveness to visual and visuomotor stimuli was best explained by the absolute

(i.e., unsigned) error between locomotion and visual flow speeds ($|Error| = |Speed_{locomotion} - Speed_{visual}$

$_{flow}|$). Changes in the absolute error resulted in corresponding changes in the activity in LC axons (**Figures**

1C and 1F), and responses scaled with the size of $\Delta|Error|$ (i.e., the change in unsigned error magnitude,

see Methods) across the different stimulus conditions (**Figure 1H**).

Noradrenaline broadcasts visuomotor prediction errors across the dorsal cortex

Individual LC neurons can preferentially target different cortical areas (Chandler et al., 2019; Kebschull et

al., 2016), but it is unclear whether these projection-specific neurons have differing response profiles. To

assess the heterogeneity of LC signaling across the cortex, we first compared the calcium responses of LC

axons between two imaging window locations: the anterior cortex (the primary and secondary motor

cortex), and the posterior cortex (V1 and surrounding regions) (**Figure 2A**). We found very similar

responses to visuomotor mismatches, playback halts (stationary) and visual flow onsets (stationary)

(**Figure 2B**), as well as air puffs and locomotion onsets at these two locations (**Figure 2C**), and there were

no significant differences in the responses between the two locations. These results are consistent with

the idea that the LC sends the same visuomotor and locomotion related signals to distinct regions of the

cortex, in line with recent findings showing that LC axons in different cortical regions show responses to

the same sensory stimuli (Deitcher et al., 2019).

We find no evidence of an effect of locus coeruleus axon stimulation on average responses in layer 2/3 of V1

What could be the function of prediction errors signaled by the LC to the cortex? One possibility is that

cortical prediction errors, like mismatch responses in V1 (Keller et al., 2012), are enhanced by LC output

in order to augment error driven learning. Indeed, a prevalent idea is that noradrenaline is a mediator of

gain modulation in the cortex (Ferguson and Cardin, 2020), including the gain increase in V1 during

locomotion (Polack et al., 2013). To test whether the LC may enhance prediction error responses in cortical

layer 2/3, we assessed the impact of transient LC axon stimulation on cortical responses using an

optogenetic strategy (**Figure 3A**). First, we injected an AAV vector encoding Cre-dependent excitatory red-

light activated opsin, ChrimsonR, into the right LC of 8 NET-Cre mice, while injecting a vector encoding

jGCaMP8m into the right V1 to drive non-specific neuronal expression. An imaging window was then

implanted over the right V1. Mice underwent habituation to the virtual reality system before calcium

imaging of V1 neurons during 1) a closed loop condition with mismatches and 2) an open loop condition

with 1 s fixed-speed visual flow stimuli. During imaging, we stimulated ChrimsonR by directing a 637 nm

laser through the objective into the imaging window (27 mW/mm^2, in 15 ms pulses presented at 20 Hz).

This stimulation was presented a) alone in closed loop conditions, b) coupled to a random 50% of

mismatches (**Figure 3B**), and c) coupled to a random 50% of open loop visual flow stimuli. To assess the

visual effects of optical stimulation of ChrimsonR (Danskin et al., 2015) in absence of the effects of

optogenetically stimulating LC axons, we also imaged from a second group of 6 mice treated in an identical

manner, but without ChrimsonR expression in the LC (control mice). Two mice were excluded from the

LC-injected group due to low density of axonal labelling with tdTomato in V1, leaving 6 ChrimsonR-

expressing mice in the dataset (see Methods).

Optogenetic stimulation, when presented in isolation, resulted in dilations of the right pupil in mice

expressing ChrimsonR (**Figure 3C**), but not in control mice. This potentially resulted from antidromic

activation of the LC, direct stimulation of which is known to produce strong ipsilateral pupil dilation (Liu

et al., 2017), and increased our confidence that we were indeed driving LC axons with the parameters
selected. Onset of the optogenetic stimulation laser caused a small calcium response across the
population in ChrimsonR-expressing mice, which was indistinguishable from that caused in control mice
(**Figure 3D**). This was likely due to a visual effect of the stimulation laser. The lack of a difference in this
stimulation response between ChrimsonR-expressing mice and control mice indicated that optogenetic
stimulation of LC axons did not have a direct measurable effect on calcium activity of L2/3 neurons in V1.

We then analyzed the effect of optogenetic activation of LC axons on stimulus responses of the layer 2/3
population. Stimulation of ChrimsonR in LC axons had no significant effect on average visuomotor
mismatch responses of layer 2/3 neurons, and the difference in response size compared between trials
with and without laser stimulation was indistinguishable from that seen in control mice (**Figure 3E**). The
same was true for visual flow responses (**Figure 3F**). The only observable difference in responses during
optogenetic stimulation was that responses to visual flow onsets (i.e., at the beginning of the visual flow
stimulus, or at the offset of mismatch) appeared slightly more pronounced (**Figure 3E-F**).

In V1, there is electrophysiological and molecular evidence for at least three different layer 2/3 pyramidal
neuron types, including negative prediction error (NPE) neurons which respond to visuomotor
mismatches, positive prediction error neurons (PPE) which respond to unexpected visual flow (Jordan and
Keller, 2020; O'Toole et al., 2022), and an intermediate group. It is therefore conceivable that LC output
could have differential effects on these groups that are masked when analyzed at the population level. To
quantify potential cell-type specific effects of LC axon stimulation, we thus split the population of neurons
into three groups based on locomotion onset responses (see **Figure S3** and Methods): the group with the
strongest visuomotor mismatch responses (NPE), the group with the strongest visual flow responses
(PPE), and an intermediate population (Other) (**Figure S3**). We found no evidence for an effect of
optogenetic stimulation of LC axons on average mismatch and visual flow responses across any of these
cell groups (**Figure 3G-H**). We thus concluded that transient LC axon stimulation in the awake state has
only a minor direct effect on the responses of neurons in layer 2/3 of V1.

Phasic LC output enhances sensorimotor plasticity in layer 2/3 of V1
If transient LC activity in the awake state does not overtly affect average layer 2/3 responses directly, what
is the function of visuomotor prediction errors signaled by the LC in the cortex? The LC is thought to be
involved in cortical plasticity (Kasamatsu et al., 1985; Kasamatsu and Pettigrew, 1976; Martins and
Froemke, 2015; Shepard et al., 2015). Catecholamine receptors in V1 are known to modulate synaptic

plasticity by activating intracellular signaling cascades and interacting with eligibility traces (Choi et al.,

2005; He et al., 2015; Nomura et al., 2014; Seol et al., 2007). One plausible function of LC sensorimotor

prediction errors is to gate the plasticity that underlies learning of sensorimotor predictions, e.g., the

prediction of visual feedback during locomotion. Primary sensory cortices can indeed learn to suppress

the sensory consequences of locomotion, driving selective reductions of responses to locomotion-coupled

inputs (Schneider et al., 2018; Widmer et al., 2022).

While our optogenetic stimulation of LC axons had no measurable effects on mismatch or visual flow

response size (**Figure 3**), we did notice one striking difference between ChrimsonR-expressing and control

mice. In V1, there is a known increase in the size of visual responses during locomotion compared to

stationary periods (Bennett et al., 2013; Erisken et al., 2014; Niell and Stryker, 2010; Pakan et al., 2016;

Polack et al., 2013; Zmarz and Keller, 2016). Analyzing only trials where the optogenetic stimulation laser

was not on, we found that this canonical gain increase was present for control mice but reversed in sign

in ChrimsonR-expressing mice: Visual responses during locomotion were suppressed on average relative

to stationary periods (**Figure 3I**). This resulted in a significantly lower locomotion modulation index (see

Methods) for ChrimsonR-expressing relative to control mice (**Figure 3I**). Since episodic LC axon stimulation

occurred throughout the whole imaging session (i.e., across 10 minutes in the closed loop condition, prior

to the presentation of open loop visual flow stimuli), it is possible that the reversal of locomotion

modulation index reflects a result of plasticity induced by earlier LC axon stimulation.

We wanted to directly test the idea that transient LC activation enhances plasticity that leads to a

reduction in locomotion modulation index over the course of visuomotor coupling experience. To do this,

we designed an experiment to assess the impact of LC axon stimulation on neuronal learning about

visuomotor coupling (**Figure 4A**). We expressed jGCaMP8m in neurons in V1 of 19 mice. 12 mice also had

AAV vector injections into the LC to express ChrimsonR in NET-positive neurons, while the remaining 7

underwent no injection into the LC to serve as controls for the non-optogenetic effects of the protocol.

Mice were habituated to the VR as before and we then proceeded to two-photon imaging in layer 2/3 of

V1. During the imaging session, mice would first be exposed to 5 minutes of 1 s fixed-speed open loop

visual flow presentations, to quantify the gain-change of visual flow responses during locomotion (**Figure

4A**). Mice were then exposed for 5-10 minutes to either a) a closed loop condition where visual flow and

locomotion are coupled, or b) an open loop replay of visual flow generated from a previous closed loop

session. During this session, LC axons would be periodically optogenetically stimulated as before (15 ms

pulses, presented at 20 Hz for 1 s every 7 seconds on average) to simulate phasic LC prediction error

responses (**Figure 4A**). Afterwards, the open loop visual flow presentations would be repeated to quantify

how the previous LC axon stimulation affected visual responses. To assess the stimulus specificity with

which visual responses changed, we presented both nasotemporal and temporonasal visual flow (mice

see the former during forward locomotion in the closed loop condition) (**Figure 4A**). If LC axon stimulation

results in enhanced plasticity of locomotion based predictions of visual flow, we would expect the

following observations: 1) Suppression of visual flow responses by locomotion is enhanced after

stimulation of LC axons in closed loop conditions, 2) this suppression is specific to the type of visual flow

seen during forward locomotion (nasotemporal), 3) the effect depends on ChrimsonR expression in the

LC, and 4) the effect is absent when LC axons are instead stimulated during open loop conditions (i.e., the

effect depends on visuomotor coupling).

Optogenetic stimulation caused a degree of pupil dilation that was strongly predictable from axonal

ChrimsonR expression levels in V1 quantified postmortem (R^2 = 0.89, p < 10^{-5}, 12 mice, linear regression)

(**Figure 4B**). We split the LC-injected dataset into high ChrimsonR expression (six mice) and low ChrimsonR

expression (six mice) according to V1 axonal expression level (**Figures 4B and S4A**). Low ChrimsonR-

expressing mice had optogenetically evoked pupil dilations that were not statistically distinguishable from

controls, while high ChrimsonR-expressing mice had overt pupil dilations (**Figures 4B and S4B**). For the

following analyses, we excluded 8 (5) of 49 (25) sessions from the ChrimsonR (control) dataset due to lack

of locomotion during those sessions (see Methods). At the beginning of the imaging session, prior to

optogenetic stimulation, neurons in layer 2/3 showed a pronounced positive locomotion modulation

index for both the nasotemporal and temporonasal visual flow responses, both in high and low

ChrimsonR-expressing mice and in control mice (**Figures 4C-E and S4D**). Quantifying visual flow responses

in high ChrimsonR-expressing mice after optogenetic stimulation during closed loop visuomotor

experience revealed a reduction in the locomotion modulation index for nasotemporal visual flow

responses (**Figure 4C**). This originated from a reduction in the response to visual flow during locomotion,

while responses during stationary periods remained stable (**Figure 4G**). This effect was specific to the

direction of visual flow seen during forward locomotion, since locomotion modulation index was

preserved for temporonasal visual flow responses (**Figure 4D**), with negligible changes in response size

both during locomotion and stationary periods (**Figure 4H**). The effect was also specific to high ChrimsonR-

expressing mice, as locomotion modulation index was largely preserved in both control and low

ChrimsonR-expressing mice (**Figures 4E, 4I, and S4D**).

The selective reduction in locomotion modulation index of nasotemporal visual flow responses in high
ChrimsonR-expressing mice could be due to enhanced learning to suppress reafferent visual feedback
seen during forward locomotion, or it could simply be due to general exposure to nasotemporal visual
flow during optogenetic stimulation (e.g., enhanced adaptation). In the latter case, response size
reduction should be the same if optogenetic stimulation takes place during open loop replay of the same
visual flow seen during closed loop conditions. To assess this, we repeated the same experiment, but
instead of stimulating LC axons during the closed loop condition, we stimulated LC axons during open loop
replays of visual flow from previous closed loop sessions of the same mice (**Figure 4A**). Instead of a
stimulus selective reduction in locomotion gain, this paradigm drove a nonsignificant increase in
locomotion modulation index for both nasotemporal (**Figures 4F and S4F**) and temporonasal visual flow
(**Figure S4E**) responses. This confirmed that the reduction in locomotion modulation index after
stimulation of LC axons in the closed loop condition (**Figure 4C**) depends on visuomotor coupling
experience and is not simply due to exposure to nasotemporal visual flow.

We conclude that LC axon stimulation over the course of minutes increases locomotion driven
suppression of visual flow responses in layer 2/3 in a manner that is specific to reafferent stimulus
properties and dependent on both ChrimsonR expression and visuomotor coupling (**Figure 4J**).
Importantly, the effect could not be explained by differences in locomotion behavior between any of the
groups analyzed (**Figure S5**).

LC axon stimulation over minutes recapitulates visuomotor plasticity seen over days
In the virtual reality environment, mouse movement and visual flow are constrained to the forward-
backward dimension, and the visuomotor gain is constant: changes in visuomotor gain that would usually
occur during changes in environment geometry are absent. Mice head-fixed in the VR therefore
experience a vastly reduced diversity of visuomotor coupling compared to freely moving conditions. The
reduction in locomotion modulation index resulting from LC axon stimulation (**Figure 4C**) may therefore
result from overlearning of this simplified visuomotor coupling.

If this interpretation is correct, then extensive training within the VR should also lead to visual flow
response suppression during locomotion even in mice without artificial LC axon stimulation. To assess this
idea, we reanalyzed two-photon calcium imaging data from V1 layer 2/3 in coupled trained mice reared
with visual experience entirely constrained to that of the visuomotor coupling in the VR across days (and
otherwise dark reared) (Attinger et al., 2017; Widmer et al., 2022) (**Figure 5A**). We compared this data to

similar data from normally reared mice that also have experience of the virtual reality system, but that had experience of normal visuomotor coupling in freely moving conditions (Vasilevskaya et al., 2022; Zmarz and Keller, 2016) (**Figure 5A**). Analyzing responses to nasotemporal visual flow onsets during open loop replay, we found that visual flow responses were amplified during locomotion relative to stationary periods in normally reared mice but were suppressed during locomotion in coupled trained mice (**Figure 5B**). This resulted in a significantly lower locomotion modulation index in coupled trained mice compared to normally reared mice (**Figure 5C**). The effect was not due to dark rearing, as non-coupled trained mice (also dark reared, but with only open loop virtual reality experience (**Figure 5A**) (Attinger et al., 2017)) showed a positive locomotion modulation index on average that did not significantly differ from that of normally reared mice (**Figure 5C**).

Qualitatively, the comparison of locomotion modulation index between normally reared and coupled trained mice (**Figure 5B-C**) was similar to the change in locomotion modulation index in ChrimsonR-expressing mice for nasotemporal visual flow after LC axon stimulation (**Figure 4C**). This suggests that the increase in locomotion driven suppression after LC axon stimulation is caused by a similar form of plasticity to that seen when visuomotor experience is constrained to the virtual reality across days. If this is the case, we should see similarities in the properties of the locomotion driven suppression. First, we looked at the locomotion speed dependence of the suppression. If the negative locomotion modulation index results from learning to suppress visual responses according to a locomotion driven prediction of visual flow, the degree of suppression should depend on locomotion speed. As locomotion speed increased, visual flow responses were increasingly suppressed in coupled trained mice (**Figure 5D**), while this was not evident in normally reared mice (**Figure 5E**). For nasotemporal visual flow responses in ChrimsonR-expressing mice, locomotion modulation index was positive and did not change with locomotion speed at the start of the imaging session, but after stimulation of LC axons in closed loop conditions, locomotion modulation index became increasingly negative as locomotion speed increased (**Figure 5F-G**). Thus, the locomotion driven suppression of visual flow emerging after stimulating LC axons in closed loop conditions was locomotion speed dependent in a similar manner to the locomotion driven suppression seen in coupled trained mice. Note that in high ChrimsonR-expressing mice, suppression of visual flow was seen at high locomotion speeds only (**Figure 5G**), but in coupled trained mice, some level of suppression was evident across all speeds (**Figure 5E**). This could be due to differences in visual flow speed: for coupled trained mice, the stimuli analyzed are visual flow onsets during open loop replay; thus, the visual flow speed will cover a similar range as the locomotion speed distribution of the mice. For high ChrimsonR-

expressing mice, the stimuli are presented at a fixed visual flow speed that often exceeds the locomotion

speed of the mouse. Thus, the fact that suppression only appears at the highest locomotion speed in these

mice is consistent with a suppression of visual flow input based on a locomotion speed dependent

expectation. In addition to the similarities in locomotion speed dependence of suppression in the two

groups, the pattern of change in locomotion modulation index across different functional layer 2/3 cell

groups also appeared similar between high ChrimsonR-expressing and coupled trained mice (**Figure S6**).

Overall, the pattern of differences in locomotion modulation was remarkably similar for ChrimsonR-

expressing (**Figures 5G, and S6D**) and coupled trained mice (**Figures 5E and S6B**). This indicates that the

functional consequence of plasticity evoked by minutes of LC axon stimulation during visuomotor coupling

is similar to that seen after experience is developmentally constrained to visuomotor coupling for days.

DISCUSSION

The locus coeruleus (LC) has long been hypothesized by computational models to report quantities such

as unexpected uncertainty (Yu and Dayan, 2005) or high level state-action prediction errors (Sales et al.,

2019). What these quantities have in common is that they essentially are metrics for the inaccuracy of the

brain's internal models. One way to implement such a measure would be to integrate prediction errors

from across the brain and then broadcast this signal back across the brain. Indeed, we found that the LC

sends unsigned visuomotor prediction errors non-specifically across the dorsal cortex (**Figures 1 and 2**).

Models hypothesizing that the LC reports the global inaccuracy of the brain's internal models suggest that

the purpose of this is to increase learning rates in output circuits like the cortex, enabling internal models

to be modified more rapidly. In line with this, we found that stimulating cortical LC axons during closed

loop visuomotor coupling over minutes reduces and even reverses the canonical locomotion related gain

of visual flow responses in layer 2/3 of V1, in a manner specific to the direction of the movement-coupled

stimulus (**Figure 4**). Surprisingly, this enhanced plasticity seems to happen even with no overt direct effect

of LC axon stimulation on average calcium responses (**Figure 3**). This plasticity, on the timescale of

minutes, recapitulates the effect of restricting visuomotor experience to the closed loop coupling in virtual

reality across days (**Figure 5**), suggesting that it reflects an acceleration of a slower form of visuomotor

plasticity (i.e., an increased learning rate).

The LC responds to unexpected stimuli of various modality, including reward prediction errors, and novel

or intense unexpected sensory stimuli (Aston-Jones and Bloom, 1981; Bouret and Sara, 2004; Breton-

Provencher et al., 2022; Deitcher et al., 2019; Foote et al., 1980; Hervé-Minvielle and Sara, 1995; Takeuchi
et al., 2016; Vankov et al., 1995). The responses we found in LC axons to the various visual flow stimuli
were proportional to the change in the degree of error between visual flow and locomotion speeds (**Figure**
1). This supports a model in which the LC signals prediction errors, and combined with previous findings,
suggests that prediction errors of all modalities are a general feature of LC signaling. Based on the
multimodality and unsigned nature of LC responses, and the integrate and broadcast nature of its input-
output anatomy (Schwarz et al., 2015), our working model of LC function is as follows: The LC integrates
prediction errors of all modalities to compute a measure of how inaccurately internal models are currently
predicting the global sensory inputs of the animal. The LC does not signal what *type* of prediction error
occurred, but what the global rate of prediction errors currently is (i.e., a measure of surprise). The
resulting LC output does not directly *drive* model updating in cortical output areas, but it modulates the
rate at which local cortical prediction errors drive plasticity in internal models. Thus, the specificity of how
internal models are changed depends on ongoing prediction errors in the cortex, and the LC input acts to
control the learning rate, modulating the extent to which these cortical prediction errors drive plasticity.
This would allow the LC to be a controller of plasticity in internal models: when prediction error rates are
high (i.e., during developmental, novel, or volatile situations), updating of internal models will be more
rapid as a result of increased LC output.

While the idea of a brain wide average over sensory prediction errors can explain many of the observed
LC responses, other responses are less easily explained by prediction error. LC calcium responses to
locomotion onsets, which have been shown previously in LC axons (Deitcher et al., 2019; Reimer et al.,
2016), were generally larger than visual flow or visuomotor mismatch responses (**Figures 2 and S2**). This
could be the result of a direct movement related input unrelated to prediction errors, or the result of
prediction errors that are not driven by differences between forward locomotion and backward visual
flow. Increases in activity occurring after locomotion onset could reflect vestibulomotor prediction errors
that result from the fact that the mouse is head-fixed, or visuomotor prediction errors for motor attempts
we do not record, like turns or head movements. We do find that visuomotor prediction errors contribute
to locomotion onset responses, as locomotion onset responses in the open loop condition are smaller
during visual flow, when locomotion evoked visuomotor error is lower (**Figure S2B**). However, LC calcium
activity also rises prior to locomotion onset, creating the additional possibility of an internally generated
movement onset response. Other kinds of movement are also correlated with LC firing, including pup
retrievals (Dvorkin and Shea, 2022) and lever pressing (Breton-Provencher et al., 2022). These movement

onset responses could also reflect other forms of prediction error, including action prediction errors
(Greenstreet et al., 2022) or temporal difference reward prediction errors (in cases where the movement
is associated with reward expectation) similar to those documented in the dopaminergic system (Kim et
al., 2020). Future work will need to investigate these possibilities to build a complete general model of LC
computation.

Depleting noradrenaline has been shown to prevent forms of sensory-guided developmental plasticity in
the cortex, including ocular dominance plasticity in V1 (Kasamatsu and Pettigrew, 1976) and changes in
the tonotopic map in the auditory cortex (Shepard et al., 2015). Loss of function experiments can suffer
from the potential confound that catecholamines may simply be necessary for normal neuronal activity,
and the loss of plasticity may be a secondary effect of a disruption in neuronal function. Indeed, acute
noradrenergic receptor block can silence neurons in layer 2/3, and massively reduce the amplitude of
visual inputs (Polack et al., 2013). In gain of function experiments, electrical stimulation of the LC has been
shown to partially restore ocular dominance plasticity in adult cats (Kasamatsu et al., 1985). Here we also
performed a gain of function experiment: we show that a few minutes of optogenetic LC axon stimulation
in V1 of adult mice during visuomotor coupling can recapitulate the developmental effects of constrained
exposure to visuomotor coupling in the virtual reality (**Figure 4 and 5**). While our optogenetic stimulation
is likely far more constrained to LC neurons than previous electrical stimulations, we cannot rule out a
systemic effect of the LC axon stimulation. We know that we are likely antidromically activating the LC
with our optogenetic stimulation owing to the resulting pupil dilation (**Figures 3C and 4B**). Given that LC
neurons are electrically coupled (McKinney et al., 2022) and have widely divergent axons, our stimulation
likely has effects on other LC output targets, such as other neuromodulatory systems, and other areas of
the cortex. Thus, the plasticity effect that we measure may not entirely result from LC output in V1 alone.
However, since the LC will never be acting in isolation and broadcasts its output to different cortical areas,
the functional impact of LC axon stimulation in this experiment is strongly suggestive of a role in facilitating
plasticity. Future work will be needed to understand the circuit locus of plasticity generated by this
paradigm, and its dependence on particular neuromodulators (e.g., dopamine and noradrenaline).

Prediction error driven catecholamine release is thought to gate cortical synaptic plasticity during
reinforcement learning (He et al., 2015; Roelfsema and Holtmaat, 2018). The results presented here
expand this idea from reward associations to sensorimotor associations. We propose that a general
function of catecholamines is to gate the plasticity underlying predictive learning across all modalities.

AUTHOR CONTRIBUTIONS

RJ designed and performed the experiments and analyzed the data. Both authors wrote the manuscript.

ACKNOWLEDGEMENTS

We thank all members of the Keller lab for discussion and support, Tingjia Lu for the production of viral vectors, Susumu Tonegawa for sharing NET-Cre mice, and Andreas Lüthi for reagents and equipment. This project has received funding from Human Frontier Science Program (LT000077/2019-L, long term fellowship to RJ), the Swiss National Science Foundation, the Novartis Research Foundation, and the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement no. 865617).

DATA AND CODE AVAILABILITY.

All data and analysis code are available at https://data.fmi.ch/PublicationSupplementRepo/.

METHODS

Viral vector injections and surgery. All animal procedures were approved by and carried out in accordance with guidelines of the Veterinary Department of the Canton Basel-Stadt, Switzerland. For all surgical procedures, mice were anesthetized with an intraperitoneal injection of a mixture of fentanyl (0.05 mg/kg), medetomidine (0.5 mg/kg), and midazolam (5 mg/kg) and provided with both peri- and postoperative general and local analgesics: Metacam (5mg/kg, s.c.) and buprenorphine (0.1 mg/kg, s.c.). A lidocaine and ropivacaine mixture was injected subcutaneously into the scalp prior to incision. Metacam analgesia was provided for a further 48 hours post-surgery. Mice were allowed to recover for 1-3 weeks prior to first head-fixation.

Surgeries for two-photon imaging of LC axons. 13 male and female NET-Cre (Wagatsuma et al., 2018) mice (4-6 weeks old) were unilaterally injected at stereotactic coordinates (relative to bregma and brain surface): 5.45 mm posterior, 1.10 mm lateral, 3.65 mm deep. 250 to 500 nl of AAV2/5-hSyn-DIO-axon-GCaMP6s (2.7 x 10^{13} genome copies/ml) was injected into each site. A 4 mm diameter circular imaging window was then implanted. A circular craniotomy was made overlying visual cortex (8 mice) or motor cortex (4 mice) and a durectomy was performed. A glass window was then placed onto the craniotomy

and fixed in place with cyanoacrylate. A custom titanium headplate was attached to the skull using dental

cement (Paladur, Kulzer).

Surgeries for two-photon imaging in layer 2/3 V1 during optogenetic stimulation of LC axons. Male and

female NET-Cre mice (4-6 weeks old) were unilaterally injected with 250 to 500 nl of AAV2/1-hSyn-DIO-

ChrimsonR-tdTomato (1.2 x 10^{13} to 3.4 x 10^{13} genome copies/ml) into the right LC at stereotactic

coordinates (relative to bregma and brain surface): 5.45 mm posterior, 1.10 mm lateral, 3.65 mm deep. A

maximum of 250 nl of AAV2/1-hSyn-soma-jGCaMP8m (titer range from 10^{13} to 10^{15} genome copies/ml)

was then injected across 3-5 locations spanning right V1, and a window and headplate were implanted as

above, but with the additional measure of black pigmenting of the dental cement for the reduction of

optogenetic stimulation light reaching the eyes of the mouse. For control mice, the injection into the LC

was omitted. For the dataset in **Figure 3**, eight mice were injected in the LC, but two were excluded due

to lack of expression (see section Axon density). Six mice were not injected in the LC to serve as controls.

For the dataset in **Figure 4**, 12 mice were injected in the LC, with 6 mice classified as high ChrimsonR

expression and 6 classified as low ChrimsonR expression (see section Axon Density). 7 mice were not

injected in the LC to serve as controls.

Immunohistochemistry. For all experiments with vector injections in LC to express either GCaMP or

ChrimsonR-tdTomato, we histologically verified that expression was confined to the neurons within the

anatomical location and spatial pattern expected of the LC. In a subset of mice used for imaging LC axons,

we also confirmed that the transgene expressing neurons were catecholaminergic by immunostaining for

tyrosine hydroxylase and GFP (**Figure S1A**). After transcardial perfusion, brains were extracted and placed

in 4% paraformaldehyde overnight. Brains were then embedded in 4% agar and sliced at 40 μm on a

vibratome. Free floating slices were then washed in PBS-triton, blocked for 2 hours with 10% goat serum,

and incubated overnight with primary antibodies (1:1000 dilution rabbit anti-tyrosine hydroxylase (Abcam

ab13970), and 1:500 dilution chicken anti-GFP (Abcam ab13970). The following day, slices were washed

again in PBS-triton and incubated for 2 hours with the secondary antibodies at 1:1000 dilution (Alexa Fluor

488 anti-chicken (Jackson ImmunoResearch 703-545-155) and Alexa Fluor 568 anti-rabbit (ThermoFisher

Scientific, A10042)). Slices were mounted in Fluoroshield with DAPI (F6057, Sigma Aldrich).

Virtual reality and visual stimuli. A virtual tunnel, with walls patterned with vertical sinusoidal gratings

was projected onto a toroidal screen spanning 240 degrees horizontally and 100 degrees vertically of the

visual field. The projector output was gated by a 24 kHz TTL such that it only turned on at the turnaround

points of the two-photon resonance scanner, minimizing light artifacts during imaging. Mice were head-

fixed on an air-floated polystyrene ball of 20 cm diameter. Movement of the virtual tunnel walls could

occur only in one dimension (forward and backwards), with the ball restricted to rotation around the pitch

axis using a pin. In closed loop conditions, visual flow speed in the tunnel was coupled to the rotation of

the ball, with the exception of brief 1 s mismatch stimuli. These were triggered every 12 ± 2 s (mean ± SD)

regardless of locomotion behavior and would clamp the visual flow speed to zero. In open loop conditions,

the visual flow speed was controlled irrespective of mouse locomotion. Three different types of open loop

session were used: 1) for LC axon imaging (**Figures 1 and 2**), as well as optogenetic stimulation of LC axons

during open loop replays (**Figure 4**), the visual flow from a previous closed loop session would be replayed.

For LC axon imaging, the visual flow onset responses and playback halt responses were analyzed during

this session. 2) For stimulation of LC axons to assess the effect on cortical responses (**Figure 3**), 1 s fixed

speed nasotemporal visual flow stimuli would be presented every 9 ± 3 s (mean ± SD). We presented 3

different speeds of visual flow in a pseudorandom order. 3) For assessing the effects of LC axon stimulation

on visuomotor plasticity (**Figure 4**), we presented 1 s duration fixed speed visual flow stimuli every 7 ± 1 s

(mean ± SD), but this time only one speed was presented (equivalent to the speed seen during 24 cm/s

locomotion in closed loop conditions), however, the direction was either nasotemporal or temporonasal,

in a pseudorandom order. All mice imaged were habituated to head-fixation and the virtual reality tunnel

for at least four daily one-hour sessions prior to imaging, until they showed regular, comfortable

locomotion.

Two-photon imaging. Imaging was performed on a modified Thorlabs Bergamo II two-photon microscope

system. Excitation illumination was provided by a Ti-Sapphire laser with a wavelength at 930 nm and a

laser power between 20 and 30 mW under the objective. When imaging neurons in V1, a piezo scanner

was used to move the objective to image 4 separate layers sequentially (15 Hz effective frame rate), but

for LC axons, imaging was fixed in a single layer (60 Hz frame rate). Field of view size was 750 × 400 pixels

(approximately 300 x 300 µm). Custom software was used to acquire the imaging data. For mice

undergoing imaging of layer 2/3 V1 neurons (**Figures 3 and 4**), two-photon imaging sites were confirmed

as being in V1 by mapping V1 boundaries using intrinsic signal optical imaging.

Optogenetic stimulation of LC axons. The beam from a 637 nm OBIS laser (Coherent) was focused onto

the imaging site via the two-photon microscope objective. The laser was gated to turn on only during the

turnaround times of the two-photon resonance scanner, to minimize stimulation induced light artifacts.

During stimulation, the laser was presented in 15 ms pulses at 20 Hz (30% duty cycle) for a total duration

of 1 s. For stimulation during visual stimulus presentations (**Figure 3**), the power was 27 mW/mm^2. Since

this evoked a small visual response in the V1 calcium activity of both ChrimsonR-expressing and control

mice (**Figure 3C**), the power was reduced for subsequent experiments to 20 mW/mm^2 (**Figure 4**), at which

point the positive visual response disappeared (**Figure S4C**). For the optogenetic stimulation during visual

stimulus presentations (**Figure 3**), optogenetic stimulation occurred simultaneous with a random 50% of

mismatch stimuli and visual flow stimuli, or in isolation, every 12 s on average, during closed loop

conditions. For optogenetic stimulation in the plasticity experiments (**Figure 4**), stimulation occurred

every 7 s on average. In 17/61 included imaging sites, this was gated by mouse locomotion such that

stimulation only occurred when locomotion speed exceeded 4 cm/s. Since the results looked very similar

for both locomotion gated and ungated stimulations, the data were pooled for analysis.

ΔF/F calculation. Raw images were full-frame registered to correct for lateral brain motion. For V1 layer

2/3 neurons, neuronal somata were manually selected based on mean and maximum fluorescence

images. Average fluorescence per region of interest (ROI) was corrected for slow fluorescence drift over

time using an 8th percentile filter and a 1000 frame window, and divided by the median value over the

entire trace to calculate ΔF/F (Dombeck et al., 2007). For LC axonal data (**Figures 1 and 2**), there were a

number of alternative/additional steps. First, all ROIs within an image above a manually set fluorescence

threshold were selected. Non-axonal ROIs from this set were then removed in two ways: first, a 'circularity

index' (circularity index = 4π x area/perimeter^2) and area threshold were used. ROIs with circularity index

above 0.14 or a total area below 150 pixels were excluded – this served to remove non-axonal structures.

Second, any ROI without evidence of calcium activity was removed: a fast Fourier transform was used to

obtain the power spectrum of the ΔF/F trace. From this, the signal-to-noise ratio of the calcium signal was

calculated, where noise was calculated as the average power between 3 and 8 Hz. ROIs with power below

a signal-to-noise ratio of 10 in the frequency band from 0.05 to 1 Hz (classified as signal) were excluded.

This served to remove uncommon elongated non-axonal ROIs, such as blood vessel walls. Next, due to the

lower signal to noise of the axonal recordings, the ΔF/F recorded at 60 Hz was low pass filtered at 10 Hz.

Finally, due to the high correlation of ΔF/F between LC axonal ROIs within a field of view (**Figure S1D-F**),

ΔF/F was averaged across all axonal ROIs within a field of view.

Calculation of visuomotor error (Figure 1). Visuomotor error is calculated as the difference between

locomotion speed and visual flow speed, assuming a constant gain between the two that the mouse

predominantly experiences in the virtual reality. The change in absolute error (Δ|Error|) is calculated for

each trial as the change in absolute (i.e., unsigned) visuomotor error during the stimulus window, minus

the absolute visuomotor error in the 1 s preceding the stimulus. For each type of stimulus and each FoV,

Δ|Error| was then averaged across trials. Note that for visuomotor mismatch, Δ|Error| is equivalent to

the locomotion speed during mismatch (since visual flow speed during mismatch is zero, and the

preceding 1 s is closed loop, where the error is zero). In all stimulus conditions (**Figure 1H**), the Δ|Error|

is calculated in a window 0.66 s (40 frames) preceding the window used to calculate ΔF/F responses.

Locomotion onsets and visual flow onsets. Locomotion onsets were defined as the locomotion speed

crossing a threshold of 0.4 cm/s, where the average speed in the previous two seconds was < 0.4 cm/s,

and where the average speed in the subsequent 1 s exceeded 2 cm/s. The same criteria were used to

determine visual flow onsets during open loop replays, using the visual flow speed.

LC axonal responses. For each visual stimulus trigger, the average ΔF/F in the 1 s window prior to stimulus

onset was subtracted from the response, before averaging across trials to get the average response of

each axon segment. Mismatch and playback halt responses were calculated for each FoV in the 1 s window

beginning 0.66 s (40 frames) after stimulus onset. For visual flow onsets during open loop replays, the

window was delayed by 400 ms (25 frames) to take into account the slower onset of the stimulus

compared to mismatches. Trials were classified as occurring during locomotion if the locomotion speed

in the 1 s after the onset of visual flow or mismatch exceeded 1 cm/s, while they were classified as

occurring during stationary periods if the locomotion speed was below 1 cm/s. For **Figure 1H**, fast

locomotion during mismatch was classified as locomotion speed exceeding 10 cm/s, while slow

locomotion was classified as locomotion speed between 1 cm/s and 10 cm/s. For locomotion onset

responses (**Figure S2**), the average ΔF/F in the 500 ms window 2 s prior to locomotion onset was

subtracted from the response, before averaging across trials. Locomotion onsets were considered to take

place during the absence of visual flow if average visual flow speed in the 2 s window centered on

locomotion onset was below that corresponding to a 1 cm/s locomotion speed in closed loop conditions.

If visual flow speed exceeded this value, the trial was considered to take place during visual flow.

Responses for all stimuli were averaged for each FoV across trials only if there were at least three valid

trials. For all stimulus conditions that were selected based on locomotion speed (e.g., mismatches), 120

sham triggers were generated and sham responses calculated from triggers that were sub-selected based

on the same locomotion selection criteria. The average sham response was subtracted from the stimulus
response.

V1 layer 2/3 neuron stimulus responses. For each stimulus trigger, the average ΔF/F in the 1 s window
prior to stimulus onset was subtracted from the response, before averaging across trials to get the average
response of the neuron. Visual flow and mismatch responses were calculated for each neuron in the
window 0.33 s to 2.33 s (5 to 30 frames) after stimulus onset. Trials were classified as occurring during
locomotion if the locomotion speed in the 1 s during visual flow or mismatch exceeded 4 cm/s, while they
were classified as during stationary periods if the locomotion speed was below 1 cm/s. For mismatches,
only locomotion trials were included. Note that for **Figure 3**, visual responses were not segregated into
locomotion and stationary trials: all trials were included. For analysis of locomotion modulation index in
different locomotion speed conditions (**Figure 5D-G**), slow, medium, and fast locomotion speed trials
were categorized where average locomotion speed during visual flow was between 1 and 5 cm/s, 5 and
10 cm/s, and above 10 cm/s respectively. For the previously published datasets (**Figure 5**, CT, NR, and NT)
(Attinger et al., 2017; Vasilevskaya et al., 2022; Widmer et al., 2022), differences in the measurement of
locomotion speed across the datasets made it difficult to use a common threshold value. Thus, to make
the locomotion modulation indices comparable to those calculated from the newly acquired dataset
(**Figure 5D-G**), we used the 95[th] percentile of the locomotion speed to normalize thresholds across
datasets. In addition, since the visual responses used to measure locomotion modulation index were
visual flow onsets in open loop conditions rather than rapid onset fixed speed stimuli, the analysis window
relative to stimulus onset was shifted by 400 ms to account for the slower onset. Responses were
averaged for each neuron across trials only if there were at least three valid trials. For all stimulus
conditions that were selected based on locomotion speed (e.g., mismatches), the average response to
1000 sham triggers sub-selected based on the same locomotion conditions was calculated. This sham
response was subtracted from the stimulus response.

Locomotion modulation index. To quantify the change in mean visual flow response during locomotion
for each neuron normalized by the overall size of response, we calculated a locomotion modulation index
(LMI) for each neuron as follows:

594
$$LMI = \frac{R_{VFloco} - R_{VFstat}}{\sqrt{R_{VFloco}{}^2 + R_{VFstat}{}^2}}$$

Where R_{VFloco} is the average visual flow response during locomotion, and R_{VFstat} is the average visual flow

response during stationary periods.

Exclusion criteria in analysis of visual plasticity (Figure 4). Imaging sessions during which the mouse spent

less than 15% of the time locomoting at a speed exceeding 4 cm/s were excluded (13 of 71 sessions), as

this would limit both the amount of visuomotor coupling experienced during the plasticity assay, and our

ability to analyze visual responses during locomotion.

Layer 2/3 functional cell type classification (Figure S3). Neurons were grouped into visually suppressed

and mismatch activated NPE neurons, visually activated PPE neurons, and an intermediate group ('other').

Since we were analyzing the visual and mismatch responses of these groups, we used locomotion onset

responses to group neurons to avoid a circular analysis. To compute each neuron's locomotion onset

response, for each onset the average ΔF/F in the 1 s window beginning 2 s prior to locomotion onset was

subtracted from the response, before averaging across trials. Average responses to locomotion onsets

(quantified in the window 0.66 to 4 s (10 to 60 frames) after onset) in open loop conditions were

subtracted from locomotion onset responses in closed loop conditions (where visual flow is concurrent

with locomotion) for each neuron. Note that for analyzing the dataset in **Figure 4**, only the open loop

locomotion onsets before LC axon stimulation were used. It was previously shown that the difference

between locomotion onset responses in open and closed loop conditions correlates with these cell groups

(Jordan and Keller, 2020). For each dataset, the 33rd and 66th percentile of these values were used as

thresholds. Neurons with response differences exceeding the 66th percentile were classified as PPE

neurons, and those with response differences below the 33rd percentile were classified as NPE neurons,

with the remaining neurons were classified as 'other'. Responses to mismatch and visual flow stimuli were

consistent with these groupings in all three datasets (**Figure S3**).

Axon density. PFA fixed brains of mice injected with a vector to express ChrimsonR-tdTomato in NET-

positive neurons of the LC were sliced at 40 μm and after mounting, examined under a confocal

microscope at 20x magnification. For all mice, images were taken from four sites in V1, each one from a

different cortical slice, with sites chosen based on somatic labelling in the green channel (indicating

GCaMP expression and a likely two-photon imaging site). To quantify LC axon density, the red channel

images were then analyzed. tdTomato labelled axonal segments were manually traced in ImageJ using the

NeuronJ plugin (Meijering et al., 2004). The total combined length of the axon segments was then divided

by the total area of the cortex in the image to calculate axon density in each image. The average density

across four images was taken as the axon density for each mouse. This value was used to distinguish high

expression and low expression mice based on a threshold of 0.002 µm per µm^2 (total axon length per unit

area of the cortex) (**Figures 4B and S4A**). Since only two mice were categorized to have low expression in

the dataset used in **Figure 3**, low expression mice were not included as a separate group in these analyses,

and instead excluded.

Pupillometry. Images of the right eye, ipsilateral to the side of LC stimulation, were recorded with a CMOS

infrared camera at 30 Hz. Pupil diameter was measured offline by fitting a circle to the pupil, which was

backlit by the 930 nm laser light of the two-photon microscope. Pupil diameter traces were z-scored by

subtracting the mean of the entire trace and dividing by the standard deviation of the entire trace. To

calculate optogenetically induced pupil dilations, the average baseline pupil diameter (in the 1 s prior to

stimulus onset) was subtracted from the response for each trial, then the pupil diameter response was

averaged across trials. Trials including blinks were excluded.

Statistical tests. All statistical information for the tests performed in this manuscript are provided in **Table**

S1. For data where the experimental unit was neurons, we used hierarchical bootstrapping (Saravanan et

al., 2020) for statistical testing due to the nested structure (neurons and mice) of the data. To do this, we

first resampled the data (with replacement) at the level of imaging sites. From the selected set of imaging

sites, we then resampled the data (with replacement) at the level of neurons. We then computed the

mean of this bootstrap sample. This would be repeated 10 000 times to generate a distribution of mean

values. For paired tests, the p-value was calculated as the proportion of this distribution that was higher

or lower than zero, depending on the null hypothesis. For unpaired tests, the p-value was calculated as

the proportion of the distribution higher or lower than the values from the distribution of the compared

dataset, depending on the hypothesis. For data where the experimental unit was FoV (**Figures 1 and 2**) or

mice (pupil dilation, **Figure S4B**), t-tests were used to compare datasets that conformed with normality

and did not show statistically distinguishable variances. Datasets that did not conform with normality or

had unequal variances were compared with Rank-sum tests. Paired t-tests were used to test the mean of

a population against zero when the data was normally distributed, otherwise signed-rank tests were used

to test the median against zero (**Figure 1**).

 MAIN FIGURES

Figure 1. LC axonal calcium activity reflects unsigned visuomotor prediction errors.

(**A**) Left: Schematic of the two-photon microscope and virtual reality system. Right: Schematics of the two virtual reality conditions. Top: closed loop condition in which visual flow speed (green) is yoked to locomotion speed (purple). Mismatches consist of 1 s halts in visual flow during locomotion (orange shading). Bottom: open loop condition in which visual flow from the closed loop session is replayed to the mouse uncoupled from locomotion. Playback halts are 1 s halts in visual flow in the open loop condition (gray shading).

(**B**) Two-photon calcium imaging of LC axons in the dorsal cortex. Left: NET-Cre mice underwent stereotactic viral vector injections to express axon-targeted GCaMP6s in the LC. 4 weeks later, axons were imaged in two regions of the cortex. Inset shows example two-photon field of view (FoV). Right: Example

ΔF/F traces (black) from three different axons segments in the same FoV, recorded during the closed loop
condition. Purple and green traces show locomotion and visual flow speed respectively. Orange shading
indicates mismatch events. Gray trace shows pupil diameter.

(**C**) Responses of LC axons to visual flow halt stimuli (left: mismatches, and right: playback halts while the
mouse is stationary). Top: Locomotion speed (purple), visual flow speed (green) and absolute error (red;
absolute difference between visual flow speed and locomotion speed) each averaged across trials. Middle:
Heat map of the average responses of different fields of view (FoVs) sorted according to response
magnitude. Bottom: Response averaged across FoVs. Shading indicates SEM.

(**D**) Average responses per FoV to mismatches and playback halts, while the mouse was either stationary
or locomoting. Error bars indicate SEM across FoVs. Here and in other panels, n.s.: not significant, *: p <
0.05, **: p < 0.01, ***: p < 0.001. For complete statistical information see **Table S1**.

(**E**) Mean mismatch response of LC axons averaged across FoVs as a function of locomotion speed. Top:
Heat map of the average mismatch response, sorted according to locomotion speed (speeds correspond
to below plot). Bottom: Mean and SEM (error bars) of the mismatch responses averaged over FoVs as a
function of locomotion speed. Only datapoints with at least 10 FoVs in each speed bin are included.

(**F**) Responses of LC axons to visual flow onsets in the open loop condition during locomotion (left) or
stationary periods (right), plotted as in panel **C**.

(**G**) Average responses of LC axons to visual flow onsets while the mouse was either locomoting or
stationary, plotted as in panel **D**.

(**H**) Responses of LC axons to the different visual flow stimuli in different conditions averaged across FoVs,
color coded according to the legend, and plotted as a function of the change in absolute error between
locomotion speed and visual flow speed during the stimulus (relative to 1 s prior to stimulus onset). Error
bars indicate SEM.

Figure 2. LC axonal responses are indistinguishable in sensory (posterior) and motor (anterior) cortical areas.

(**A**) Schematic of the two different locations of the two-photon imaging windows used. Posterior windows were centered on V1 and included surrounding structures, while anterior windows included the primary and secondary motor cortex.

(**B**) Responses of LC axons to mismatch, playback halt, and visual flow onset averaged across FoVs, compared between posterior and anterior imaging locations. Error bars indicate SEM. Here and in other panels, n.s.: not significant. For complete statistical information see **Table S1**.

(**C**) As in **B**, but for air puff responses (which were infrequently used to evoke locomotion) and locomotion onset responses in various conditions, each averaged in the 3 s window after onset. Note that locomotion_Visual flow and locomotion_No visual flow occur in open loop conditions (see also **Figure S2**).

Figure 3. LC axon stimulation has no measurable effect on stimulus responses in layer 2/3 of V1.

(**A**) AAV vector injections were used to express ChrimsonR-tdTomato in LC NET-positive neurons, and jGCaMP8m in layer 2/3 neurons of V1. Simultaneous two-photon calcium imaging and optogenetic stimulation with a 637 nm laser would take place in layer 2/3. Injections into the LC were omitted in control mice. Example histology images indicate expression of jGCaMP8m (green) and tdTomato (magenta); note that blood vessels and pia mater (see arrows labelled 'BV' and 'pia' respectively) are also visible in the magenta channel. Arrow labelled 'Ax' indicates an axon.

(**B**) Example ΔF/F traces for three somatic ROIs in layer 2/3 of V1 (black) and the corresponding visual flow speed (green) and locomotion speed (purple) traces. Orange shading indicates visuomotor mismatch events. Red marks indicate concurrent laser stimulation to activate ChrimsonR on a random subset of mismatch trials.

(**C**) Average pupil diameter response to stimulation with the optogenetic laser presented in isolation for
ChrimsonR-expressing mice (pink), and control mice (gray). Shading represents SEM.

(**D**) Left: Average population ΔF/F response of layer 2/3 neurons in V1 to the onset of the optogenetic
stimulation laser presented in isolation. Right: Boxplots to compare average response (quantified in the
window 0.3 to 1.6 s after optogenetic stimulation onset) for the two groups of mice. Here and in other
panels, n.s.: not significant. For complete statistical information see **Table S1**.

(**E**) Analysis of mismatch responses of all layer 2/3 neurons. Left: Average population ΔF/F response of
layer 2/3 neurons in V1 of ChrimsonR-expressing mice to visuomotor mismatch either with (red) or
without (black) concurrent optogenetic laser stimulation. Right: Boxplots to compare the difference in
average mismatch response between trials with and without optogenetic laser stimulation. Pink points
indicate data for each neuron from ChrimsonR-expressing mice, and gray points indicate data from control
mice. Note, statistical tests against zero for the effect in ChrimsonR-expressing mice were also insignificant
(see **Table S1**).

(**F**) As for **E**, but for responses to 1 s fixed speed visual flow stimuli in open loop conditions.

(**G**) As for boxplots in panel **E**, but for mismatch responses of three different functionally defined neuronal
groups: NPE = negative prediction error neurons, with large responses to mismatches, PPE = positive
prediction error neurons, with large responses to visual flow, and an intermediate group 'other'. See
Methods and **Figure S3** for information on neuronal types. Note, statistical tests against zero for the effect
in ChrimsonR-expressing mice were also insignificant (see **Table S1**).

(**H**) As for **G**, but for responses to 1 s fixed speed visual flow stimuli in open loop conditions.

(**I**) Left: Population average responses to visual flow stimuli during stationary periods (black) and during
locomotion (purple), for control mice (bottom), and ChrimsonR-expressing mice (top). Shading represents
SEM. Note that only trials *without* concurrent optogenetic stimulation are included here. Right:
Histograms and boxplots to show distribution of locomotion modulation index for the visual responses
recorded in control (gray) and ChrimsonR-expressing mice (pink). Here, **: $p < 0.01$ (see **Table S1**).

Figure 4. Phasic LC output enhances sensorimotor plasticity in layer 2/3 of V1.

(**A**) Diagram of the experiment used to determine whether LC axon stimulation during different visuomotor coupling conditions can modulate plasticity. Visual responses are compared before and after 5-10 minutes of either closed or open loop conditions, during which LC axons were stimulated every 7 s on average.

(**B**) Left: Scatter plot to show the relationship between density of ChrimsonR-tdTomato labelled axons in V1 (total axon length per unit area of the cortex, analyzed postmortem) and the average evoked pupil dilation during optogenetic stimulation. Green dashed line is a linear regression fit to the data, and red dotted line indicates axon density threshold used to categorize low (blue) and high (pink) ChrimsonR-expressing mice. Right: Average pupil diameter response to stimulation with the optogenetic laser for 6 mice with high ChrimsonR expression in LC axons (pink), 6 mice with low ChrimsonR expression (blue), and 7 control mice that did not receive a vector injection into the LC (black). Shading represents SEM over sessions.

(**C**) Analysis of plasticity in nasotemporal visual flow responses in ChrimsonR-expressing mice undergoing
optogenetic laser stimulation during closed loop visuomotor experience. Histograms and boxplots show
distribution of locomotion modulation indices for the visual responses of layer 2/3 V1 neurons recorded
before (dark gray) and after the stimulation period (red). Here and in all other panels: n.s.: not significant,
*: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$. See **Table S1** for all statistical information.

(**D**) As for panel **C**, but for temporonasal visual flow responses.

(**E**) As for panel **C**, but for low ChrimsonR-expressing mice.

(**F**) As for panel **C**, but for the condition in which optogenetic stimulation occurred during open loop
replays of visual flow (i.e., no visuomotor coupling).

(**G**) Analysis of plasticity in nasotemporal visual flow responses in high ChrimsonR-expressing mice
undergoing optogenetic laser stimulation during the closed loop condition. Left: Boxplots of the change
in response to visual flow after optogenetic stimulation for responses recorded during locomotion
(purple) and those recorded during stationary periods (black). Right: Average population response to
visual flow during locomotion (purple) and during stationary periods (black), before (top) and after
(bottom) optogenetic stimulation. Shading indicates SEM.

(**H**) As for panel **G**, but for temporonasal visual flow responses.

(**I**) As for panel **G**, but for low ChrimsonR-expressing mice.

(**J**) Change in locomotion modulation index after optogenetic stimulation, for the four sets of
experiments from panels **C** (nasotemporal), **D** (temporonasal), **Figure S4C** (Control), **E** (low ChrimsonR
expression), and **F** (open loop) (presented in the respective sequence).

Figure 5. LC axon stimulation over minutes recapitulates the visuomotor plasticity seen over days.

(**A**) Diagram of the three groups of mice reared with different visuomotor experience. Normally reared mice (NR, gray) were raised with a normal light/dark cycle in their cages, with the full diversity of visuomotor coupling in freely moving conditions. Coupled trained mice (CT, blue) were reared in the dark, and their only visuomotor experience prior to imaging was during closed loop conditions in the virtual reality system. Non-coupled trained mice (NT, orange) were reared similar to coupled trained mice, except their only visual experience was during open loop conditions in the virtual reality system.

(**B**) Average population responses of layer 2/3 neurons in V1 to the onset of visual flow during open loop replay sessions. Green shows the average speed profile of the visual flow stimulus, purple shows the average population responses during locomotion, and black shows the average population responses during stationary periods for normally reared (top) and coupled trained (bottom) mice. Shading indicates SEM.

(**C**) Distribution of the locomotion modulation indices of all V1 layer 2/3 neurons for the visual flow onset responses recorded in normally reared (gray), coupled trained (blue), and non-coupled trained (orange) mice. Here and in other panels, n.s.: not significant, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$. For complete statistical information see **Table S1**.

(**D**) Average population responses of layer 2/3 neurons in V1 in coupled trained mice to the onset of visual flow during stationary periods (black), and during slow (top) and fast (bottom) locomotion (purple). Shading indicates SEM.

(**E**) Comparison of locomotion modulation indices averaged across neurons for visual flow onset responses
recorded during slow, medium, and fast locomotion speeds, in normally reared (left) and coupled trained
(right) mice. Error bars indicate SEM. Green dashed line indicates linear regression fit to the data, and
shading indicates the 95% confidence interval of the regression. Green asterisks and 'n.s.' indicate
significance of the fit.

(**F**) As for panel **D**, but for nasotemporal visual flow responses in high ChrimsonR-expressing mice after LC
axon stimulation in closed loop conditions.

(**G**) As for panel **E**, but for nasotemporal visual flow responses in high ChrimsonR-expressing mice before
(left) and after (right) LC axon stimulation in closed loop visuomotor coupling.

801 SUPPLEMENTARY FIGURES

Figure S1. Additional information and analyses on two-photon imaging of LC axons. Related to Figure 1.

(**A**) Confocal micrographs showing immunolabelling for tyrosine hydroxylase (magenta) and GCaMP6s (green) in the two LC. GCaMP6s was expressed via stereotactic viral vector injection into only the right LC of NET-Cre mice.

(**B**) Cross-correlations between LC axonal ΔF/F and pupil diameter (black) or the time derivative of pupil diameter (gray) averaged across 40 FoVs. Shading indicates SEM. Positive axonal lag values indicate time delays where axonal fluorescence lags relative to pupil diameter or its derivative.

(**C**) Responses to air puff stimuli applied to the flank which were infrequently used to evoke locomotion (here during the closed loop condition). Top: Average locomotion speed (purple) and visual flow speed (green) for trials that evoked locomotion. Middle: Heat maps of the average air puff responses of different FoVs sorted according to response magnitude, for trials that evoked locomotion (top heat map) and rare trials that did not evoke locomotion (bottom heat map). Bottom: Response averaged across FoVs for air puff trials that evoked locomotion (black) and those that did not (gray). Shading indicates SEM.

(**D**) Example two-photon field of view of LC axons, with 18 regions of interest (ROI) superimposed in different colors.

(**E**) Heat map of ΔF/F of the 18 ROIs in panel **D** from a segment of closed loop experience.

(**F**) Correlation matrix of the activity of the 18 different ROIs in panels **D** and **E**.

Figure S2. Locomotion onset responses of LC axons. Related to Figure 2.

(**A**) Responses to the onset of locomotion occurring during the closed loop condition (left), during visual flow in the open loop condition (middle), and during no visual flow in the open loop condition (right). Top: Average visual flow speed (green) and locomotion speed (purple) averaged across trials. Note that the window for defining the occurrence of visual flow is in the 2 s window centered on the locomotion onset – therefore the incidence of visual flow increases away from this onset. Red trace shows the absolute error between locomotion speed and visual flow speed averaged across trials (note that, since absolute error is averaged across trials independently, it is *not* equivalent to the absolute difference in averaged locomotion speed and visual flow speed traces plotted above. In addition, note that the minor error occurring in the closed loop condition is due to mismatches). Middle: Heat map of the average responses of different fields of view (FoV) sorted according to response magnitude. Bottom: Response averaged over FoVs. Shading indicates SEM.

(**B**) Comparison of average sustained response of LC axons to locomotion onsets in the open loop condition (quantified in the window 1 to 10 s after locomotion onset), during either visual flow or no visual flow, across FoVs. Each point indicates the average response of an individual FoV. Error bars indicate SEM. *: $p < 0.05$. For statistical information see **Table S1**.

Figure S3. Classification of functional neuronal types in V1 layer 2/3. Related to Figures 3, 4, and 5.

(**A**) Percentage of neurons as a function of the difference in response to locomotion onsets in closed loop conditions and open loop conditions. Neurons with a responses difference below the 33rd percentile (open loop response > closed loop response), were classified as visually suppressed NPE neurons (red), while those with a response difference exceeding the 66th percentile (closed loop response > open loop response), were classified as visually driven PPE neurons (turquoise), as shown. Data for this panel comes from **Figure 3** dataset.

(**B**) Top: Diagram of visual flow and locomotion speed traces to indicate closed loop and open loop locomotion onsets (black dashed lines). Bottom: Responses to locomotion onset during closed loop and open loop, averaged across neurons classified as NPE, PPE, and the intermediate group (Other). Shading indicates SEM. Data for this panel comes from **Figure 3** dataset.

(**C**) Bar graph to show visual flow and visuomotor mismatch responses, averaged across neurons, compared for the three functional neuron types: NPE, Other, and PPE neurons. Error bars indicate SEM. Data for this panel comes from **Figure 3** dataset.

(**D**) Bar graph to show nasotemporal and temporonasal visual flow responses (split further into occurring during locomotion or stationary periods), averaged across neurons, compared for the three functional neuron types: NPE, Other, and PPE neurons. Error bars indicate SEM. Data for this panel comes from **Figure 4** dataset (High ChrimsonR only). Mismatch data was not acquired for this dataset.

(**E**) Bar graph to show visual flow onset and visuomotor mismatch responses, averaged across neurons, compared between NPE, Other, and PPE cell classes. Error bars indicate SEM. Data for this panel comes from **Figure 5** datasets (coupled trained and normally reared mice).

Figure S4. Additional information and analyses on optogenetic stimulation of LC axons to drive plasticity. Related to Figure 4.

(**A**) Two example confocal microscopy images from V1 showing jGCaMP8m expression (green) and tdTomato expression (magenta), indicating labelled LC axons. The right image is from an animal classified as high ChrimsonR-expressing, based on labelled axon density, and left image is from an animal classified as low ChrimsonR-expressing.

(**B**) Comparison of average pupil diameter response (quantified in the window 1.3 to 4.7 s after optogenetic stimulation onset) for control, low ChrimsonR-expressing and high ChrimsonR-expressing mice. Here and in other panels, n.s.: p > 0.05, *: p < 0.05, **: p < 0.01, ***: p < 0.001. For complete statistical information see **Table S1**.

(**C**) Average population ΔF/F response of layer 2/3 neurons in V1 to the onset of the optogenetic stimulation laser. Shading indicates SEM. Note that optogenetic laser power was lowered compared to the value used in **Figure 3**, and the positive response seen in **Figure 3D** is no longer evident.

(**D**) Analysis of plasticity in nasotemporal visual flow responses in control mice which underwent optogenetic laser stimulation during the closed loop condition. Left: Distribution of locomotion modulation index for the visual responses recorded before (dark gray) and after the stimulation period (red). Right: Change in ΔF/F response to visual flow after optogenetic stimulation for responses recorded during locomotion (purple) and during stationary periods (black).

(**E**) As for panel **D**, but for temporonasal visual flow responses recorded in high ChrimsonR-expressing
mice, where optogenetic stimulation occurred during open loop replays of visual flow (i.e., uncoupled
from locomotion).

(**F**) Analysis of plasticity in nasotemporal visual flow responses for high ChrimsonR-expressing mice which
underwent optogenetic laser stimulation during open loop replays of visual flow (i.e., uncoupled from
locomotion). Boxplot shows the change in ΔF/F response to visual flow after optogenetic stimulation for
responses recorded during locomotion (purple) and during stationary periods (black). Traces show
average population response to visual flow during locomotion (purple) and during stationary periods
(black), before (top) and after (below) optogenetic stimulation. Shading indicates SEM.

Figure S5. Differences in locomotion during stimulation cannot explain differences in the change of locomotion modulation index between groups. Related to Figure 4.

(**A**) Average locomotion speed (excluding stationary periods) for: High ChrimsonR-expressing mice during closed loop stimulation, Control mice during closed loop stimulation, low ChrimsonR-expressing mice during closed loop stimulation, and high ChrimsonR-expressing mice during open loop stimulation. Points show data for each imaging session.

(**B**) As for panel **A**, but for the time spent engaged in locomotion during the stimulation period.

(**C**) Change in locomotion modulation index after optogenetic stimulation plotted against the time spent engaged in locomotion during the laser stimulation period of the imaging session. Each point shows the averages for a single imaging session.

Figure S6. Comparison of locomotion modulation index across different functional neuronal types in layer 2/3 V1. Related to Figure 5.

(**A**) Average population responses of layer 2/3 neurons in V1 in coupled trained mice to the onset of visual flow during stationary periods (black), and during locomotion (purple) for visually driven positive prediction error neurons (PPE, top) and visually suppressed negative prediction error neurons (NPE, bottom). Shading indicates SEM.

(**B**) Comparison of locomotion modulation index for visual flow onset responses of positive (turquoise), negative (orange) prediction error neurons, and an intermediate population of neurons (gray), recorded in normally reared (left) and coupled trained (right) mice. For information on functional neuron types, see **Figure S3**. Here and in other panels, n.s.: p > 0.05, *: p < 0.05, **: p < 0.01, ***: p < 0.001. For complete statistical information see **Table S1**.

(**C**) As for panel **A**, but for nasotemporal visual flow responses in high ChrimsonR-expressing mice after LC axon stimulation in closed loop conditions.

(**D**) As for panel **B**, but for nasotemporal visual flow responses in high ChrimsonR-expressing mice before (left) and after (right) LC axon stimulation in closed loop conditions.

TABLES

918 **Table S1.** All statistical information.

Figure panel	Comparison	Value compared	Mean 1 *(delta shown for paired tests)*	SD 1	Mean 2 *(n/a for paired tests)*	SD 2	Test type	P-value	N (ROIs)	N (Sites)	N (Mice)
1D	Mean response *vs* zero	Playback halt response (stationary) [% ΔF/F]	-1.07	1.71	n/a	n/a	Paired t-test	**0.0045**	n/a	25	10
	Mismatch *vs* Playback halt (locomotion)	Mean response [% ΔF/F]	2.24	2.30	0.64	3.51	Rank-sum	**0.0497**	n/a	40,24	10,13
	Playback halt locomotion *vs* stationary	Mean response [% ΔF/F]	0.64	3.51	-1.07	1.71	Rank-sum	**0.0080**	n/a	24,25	10,10
	Mean response *vs* zero	Playback halt response (locomotion) [% ΔF/F]	0.64	3.51	n/a	n/a	Paired t-test	0.3841	n/a	24	10
	Mean response *vs* zero	Mismatch response [% ΔF/F]	2.24	2.30	n/a	n/a	Paired t-test	**< 10⁻⁵**	n/a	40	13
	Mismatch *vs* Playback halt (stationary)	Mean response [% ΔF/F]	2.24	2.30	-1.07	1.71	Unpaired t-test	**< 10⁻⁵**	n/a	40,25	10,13
1G	Mean response *vs* zero	Visual flow response (stationary) [% ΔF/F]	3.19	5.31	n/a	n/a	Signed-rank test	**0.0031**	n/a	28	11
	Locomotion *vs* stationary	Mean visual response [% ΔF/F]	-0.55	4.24	3.19	5.31	Rank-sum	**0.0133**	n/a	21,28	11,11
	Mean response *vs* zero	Visual flow response (locomoting) [% ΔF/F]	-0.55	4.24	n/a	n/a	Paired t-test	0.5598	n/a	21	11
2B	Posterior *vs* Anterior	Mismatch response [% ΔF/F]	2.56	2.03	1.18	2.94	Unpaired t-test	0.1127	n/a	31, 9	9,4
	Posterior *vs* Anterior	Playback halt response (stationary) [% ΔF/F]	-1.11	1.56	-1.00	2.05	Unpaired t-test	0.8801	n/a	16, 9	6, 4
	Posterior *vs* Anterior	Visual flow response (stationary) [% ΔF/F]	2.99	5.49	3.60	5.20	Unpaired t-test	0.7855	n/a	19, 9	7, 4
2C	Posterior *vs* Anterior	Locomotion response (visual flow) [% ΔF/F]	10.85	5.89	8.12	8.46	Rank-sum	0.3025	n/a	14, 6	6, 3
	Posterior *vs* Anterior	Locomotion response (closed loop) [% ΔF/F]	14.99	7.07	13.33	7.86	Rank-sum	0.4257	n/a	28, 9	9,4
	Posterior *vs* Anterior	Locomotion response (no visual flow) [% ΔF/F]	14.42	8.05	14.96	11.93	Unpaired t-test	0.9202	n/a	8, 6	6, 3
	Posterior *vs* Anterior	Air puff response [% ΔF/F]	24.73	10.03	24.48	11.23	Unpaired t-test	0.9581	n/a	13, 8	6,4
3D	Mean response *vs* zero	ChrimsonR, laser response [% ΔF/F]	0.40	2.69	n/a	n/a	Bootstrap	0.0775	1985	15	6
	Mean response *vs* zero	Control, laser response [% ΔF/F]	0.30	2.24	n/a	n/a	Bootstrap	0.1785	1026	10	6
	Control *vs* ChrimsonR	Laser response [% ΔF/F]	0.30	2.24	0.40	2.69	Bootstrap	0.3929	1026, 1985	10,15	6,6
3E	Control *vs* ChrimsonR	Mismatch response difference [% ΔF/F]	-0.26	3.39	0.05	2.86	Bootstrap	0.2244	1053, 1911	10,14	6,6
	Optogenetic stimulation *vs* no stimulation	ChrimsonR, mismatch response [% ΔF/F]	0.05	2.86	n/a	n/a	Bootstrap	0.4189	1911	14	6
3F	Control *vs* ChrimsonR	Visual flow response difference [% ΔF/F]	-0.04	2.43	0.06	2.43	Bootstrap	0.3852	1053, 2233	10,16	6,6
	Optogenetic stimulation *vs* no stimulation	ChrimsonR, visual flow response [% ΔF/F]	0.06	2.34	n/a	n/a	Bootstrap	0.3869	2233	16	6
3G	Control *vs* ChrimsonR	PPE, mismatch response difference [% ΔF/F]	-0.20	2.97	0.25	2.93	Bootstrap	0.1432	369, 541	10,12	6,6
	Control *vs* ChrimsonR	NPE, mismatch response difference [% ΔF/F]	-0.47	4.41	0.21	3.81	Bootstrap	0.1771	382, 525	10,12	6,6
	Optogenetic stimulation *vs* no stimulation	ChrimsonR PPE, mismatch response [% ΔF/F]	0.25	2.93	n/a	n/a	Bootstrap	0.2177	541	12	6
	Optogenetic stimulation *vs* no stimulation	ChrimsonR NPE, mismatch response [% ΔF/F]	0.21	3.81	n/a	n/a	Bootstrap	0.3433	525	12	6
	Control *vs* ChrimsonR	Other, mismatch response difference [% ΔF/F]	-0.02	2.22	-0.08	1.92	Bootstrap	0.5335	275, 526	10,12	6,6
	Optogenetic stimulation *vs* no stimulation	ChrimsonR Other, mismatch response [% ΔF/F]	-0.08	1.92	n/a	n/a	Bootstrap	0.6313	526	12	6

Panel	Comparison	Measure					Test	p-value			
3H	Control vs ChrimsonR	PPE, visual flow response difference [% ΔF/F]	0.33	2.86	0.09	3.19	Bootstrap	0.7156	369, 541	10,12	6,6
	Control vs ChrimsonR	NPE, visual flow response difference [% ΔF/F]	-0.20	2.18	-0.19	2.31	Bootstrap	0.4995	382, 525	10,12	6,6
	Optogenetic stimulation vs no stimulation	ChrimsonR PPE, visual flow response [% ΔF/F]	0.09	3.19	n/a	n/a	Bootstrap	0.3983	541	12	6
	Control vs ChrimsonR	Other, visual flow response difference [% ΔF/F]	-0.13	1.79	-0.08	1.49	Bootstrap	0.4584	275, 526	10,12	6,6
	Optogenetic stimulation vs no stimulation	ChrimsonR Other, visual flow response [% ΔF/F]	-0.08	1.49	n/a	n/a	Bootstrap	0.6316	526	12	6
	Optogenetic stimulation vs no stimulation	ChrimsonR NPE, visual flow response [% ΔF/F]	-0.19	2.31	n/a	n/a	Bootstrap	0.7564	525	12	6
3I	Control vs ChrimsonR	Locomotion modulation index	0.22	0.84	-0.15	0.72	Bootstrap	**0.0012**	816, 1687	8,12	5,6
4C	Before vs After	Locomotion modulation index	-0.42	1.00	n/a	n/a	Bootstrap	**< 10⁻⁵**	1066	11	5
4D	Before vs After	Locomotion modulation index	0.04	1.08	n/a	n/a	Bootstrap	0.6271	1111	11	5
4E	Before vs After	Locomotion modulation index	0.06	1.25	n/a	n/a	Bootstrap	0.5922	1140	10	5
4F	Before vs After	Locomotion modulation index	0.24	0.98	n/a	n/a	Bootstrap	0.9278	832	8	4
4G	Before vs After	Visual response (locomotion) [% ΔF/F]	-0.88	3.54	n/a	n/a	Bootstrap	**0.0013**	1066	11	5
	Before vs After	Visual response (stationary) [% ΔF/F]	0.17	2.36	n/a	n/a	Bootstrap	0.6929	1276	13	5
4H	Before vs After	Visual response (stationary) [% ΔF/F]	-0.17	2.61	n/a	n/a	Bootstrap	0.2579	1276	13	5
	Before vs After	Visual response (locomotion) [% ΔF/F]	-0.06	3.34	n/a	n/a	Bootstrap	0.4043	1111	11	5
4I	Before vs After	Visual response (stationary) [% ΔF/F]	-0.28	2.97	n/a	n/a	Bootstrap	0.2404	1140	10	5
	Before vs After	Visual response (locomotion) [% ΔF/F]	-0.08	2.61	n/a	n/a	Bootstrap	0.3878	1344	12	5
4J	Open loop vs Nasotemporal	Δ locomotion modulation index	0.24	0.98	-0.42	1.00	Bootstrap	**0.0007**	832, 1066	8, 11	4, 5
	Low expression vs Nasotemporal	Δ locomotion modulation index	0.06	1.25	-0.42	1.00	Bootstrap	**0.0264**	1140, 1066	10, 11	5, 5
	Control vs Nasotemporal	Δ locomotion modulation index	0.03	1.01	-0.42	1.00	Bootstrap	**0.0002**	646, 1066	7, 11	3, 5
	Temporonasal vs Nasotemporal	Δ locomotion modulation index	0.04	1.08	-0.42	1.00	Bootstrap	**0.0014**	1111, 1066	11, 11	5, 5
5C	CT vs NR	Locomotion modulation index	-0.14	0.92	0.08	0.87	Bootstrap	**0.0164**	4247, 4383	24,26	19,23
	NT vs CT	Locomotion modulation index	0.08	0.92	-0.14	0.92	Bootstrap	**0.0308**	1969, 4247	12, 24	8,19
	NR vs NT	Locomotion modulation index	0.08	0.87	0.08	0.92	Bootstrap	0.5093	4383, 1969	26,12	23,8
5E	NR vs CT	Locomotion modulation index, fast locomotion	0.09	0.88	-0.27	0.90	Bootstrap	**0.0008**	3042, 2298	16,15	15,12
	NR vs CT	Locomotion modulation index, slow locomotion	0.16	0.87	-0.07	0.88	Bootstrap	**0.0118**	4669, 2907	31,18	29,16
	NR vs CT	Locomotion modulation index, medium locomotion	0.10	0.89	-0.14	0.94	Bootstrap	**0.0178**	4098, 3481	25,21	22,17
5G	Before vs After	Locomotion modulation index, fast locomotion	0.27	0.97	-0.25	0.92	Bootstrap	**0.0024**	389, 755	4,7	3,4
	Before vs After	Locomotion modulation index, medium locomotion	0.35	0.90	-0.03	0.93	Bootstrap	**0.0429**	872, 723	9,8	5,5
	Before vs After	Locomotion modulation index, slow locomotion	0.36	0.85	0.28	0.91	Bootstrap	0.3475	1123, 1043	12,11	5,5
S2B	With vs without visual flow	Locomotion onset response (open loop) [% ΔF/F]	3.03	7.82	7.17	6.45	Rank-sum	**0.0442**	n/a	20,14	9,9
S4B	Control vs High expression	Pupil diameter response [z-score]	-0.09	0.12	0.14	0.09	Unpaired t-test	**0.0024**	n/a	n/a	7,6
	High expression vs Low expression	Pupil diameter response [z-score]	0.14	0.09	0.00	0.05	Unpaired t-test	**0.0071**	n/a	n/a	6,6
	Low expression vs Control	Pupil response [z-score]	0.00	0.05	-0.09	0.12	Unpaired T-test	0.1242	n/a	n/a	6,7
S4D	Before vs After	Visual response (stationary) [% ΔF/F]	-0.70	3.12	n/a	n/a	Bootstrap	**0.003**	793	8	4
	Before vs After	Visual response (locomotion) [% ΔF/F]	-0.57	3.83	n/a	n/a	Bootstrap	0.0534	1163	12	5
	Before vs After	Locomotion modulation index	0.03	1.01	n/a	n/a	Bootstrap	0.5909	646	7	3

S4E	Before *vs* After	Locomotion modulation index	0.17	1.02	n/a	n/a	Bootstrap	0.8755	718	7	4
S4F	Before *vs* After	Visual response (stationary) [% ΔF/F]	-0.44	2.31	n/a	n/a	Bootstrap	0.1088	832	8	4
	Before *vs* After	Visual response (locomotion) [% ΔF/F]	-0.07	3.47	n/a	n/a	Bootstrap	0.4273	832	8	4
S6B	NR *vs* CT	Locomotion modulation index, other	0.07	0.89	-0.18	0.94	Bootstrap	**0.0341**	1276, 1384	23,24	22,19
	NR *vs* CT	Locomotion modulation index, NPE	-0.10	0.90	-0.22	0.91	Bootstrap	0.1196	1510, 1108	23,24	22,19
	NR *vs* CT	Locomotion modulation index, PPE	0.11	0.81	-0.07	0.90	Bootstrap	**0.0436**	986, 1755	23,24	22,19
S6D	Before *vs* After	Locomotion modulation index, other	0.33	0.90	-0.13	0.93	Bootstrap	**0.0239**	378, 408	13,13	5,5
	Before *vs* After	Locomotion modulation index, NPE	0.12	0.88	-0.18	0.87	Bootstrap	0.0706	401, 418	13,13	5,5
	Before *vs* After	Locomotion modulation index, PPE	0.51	0.80	0.25	0.92	Bootstrap	0.1006	372, 365	13,13	5,5

REFERENCES

Aston-Jones, G., Bloom, F.E., 1981. Activity of norepinephrine-containing locus coeruleus neurons in behaving rats anticipates fluctuations in the sleep-waking cycle. J. Neurosci. 1, 876–886. https://doi.org/10.1523/JNEUROSCI.01-08-00876.1981

Attinger, A., Wang, B., Keller, G.B., 2017. Visuomotor Coupling Shapes the Functional Development of Mouse Visual Cortex. Cell 169, 1291-1302.e14. https://doi.org/10.1016/j.cell.2017.05.023

Bennett, C., Arroyo, S., Hestrin, S., 2013. Subthreshold mechanisms underlying state-dependent modulation of visual responses. Neuron 80, 350–357. https://doi.org/10.1016/j.neuron.2013.08.007

Bouret, S., Sara, S.J., 2004. Reward expectation, orientation of attention and locus coeruleus-medial frontal cortex interplay during learning. Eur. J. Neurosci. 20, 791–802. https://doi.org/10.1111/j.1460-9568.2004.03526.x

Breton-Provencher, V., Drummond, G.T., Feng, J., Li, Y., Sur, M., 2022. Spatiotemporal dynamics of noradrenaline during learned behaviour. Nature 1–7. https://doi.org/10.1038/s41586-022-04782-2

Chandler, D.J., Jensen, P., McCall, J.G., Pickering, A.E., Schwarz, L.A., Totah, N.K., 2019. Redefining Noradrenergic Neuromodulation of Behavior: Impacts of a Modular Locus Coeruleus Architecture. J. Neurosci. 39, 8239–8249. https://doi.org/10.1523/JNEUROSCI.1164-19.2019

Choi, S.-Y., Chang, J., Jiang, B., Seol, G.-H., Min, S.-S., Han, J.-S., Shin, H.-S., Gallagher, M., Kirkwood, A., 2005. Multiple receptors coupled to phospholipase C gate long-term depression in visual cortex. J. Neurosci. Off. J. Soc. Neurosci. 25, 11433–11443. https://doi.org/10.1523/JNEUROSCI.4084-05.2005

Cox, J., Witten, I.B., 2019. Striatal circuits for reward learning and decision-making. Nat. Rev. Neurosci. 20, 482–494. https://doi.org/10.1038/s41583-019-0189-2

Danskin, B., Denman, D., Valley, M., Ollerenshaw, D., Williams, D., Groblewski, P., Reid, C., Olsen, S., Waters, J., 2015. Optogenetics in Mice Performing a Visual Discrimination Task: Measurement and Suppression of Retinal Activation and the Resulting Behavioral Artifact. PLOS ONE 10, e0144760. https://doi.org/10.1371/journal.pone.0144760

Deitcher, Y., Leibner, Y., Kutzkel, S., Zylbermann, N., London, M., 2019. Nonlinear relationship between multimodal adrenergic responses and local dendritic activity in primary sensory cortices. https://doi.org/10.1101/814657

Dombeck, D.A., Khabbaz, A.N., Collman, F., Adelman, T.L., Tank, D.W., 2007. Imaging Large-Scale Neural Activity with Cellular Resolution in Awake, Mobile Mice. Neuron 56, 43–57. https://doi.org/10.1016/j.neuron.2007.08.003

Dvorkin, R., Shea, S.D., 2022. Precise and Pervasive Phasic Bursting in Locus Coeruleus during Maternal Behavior in Mice. J. Neurosci. Off. J. Soc. Neurosci. 42, 2986–2999. https://doi.org/10.1523/JNEUROSCI.0938-21.2022

Eliades, S.J., Wang, X., 2008. Neural substrates of vocalization feedback monitoring in primate auditory cortex. Nature 453, 1102–1106. https://doi.org/10.1038/nature06910

Erisken, S., Vaiceliunaite, A., Jurjut, O., Fiorini, M., Katzner, S., Busse, L., 2014. Effects of Locomotion Extend throughout the Mouse Early Visual System. Curr. Biol. 24, 2899–2907. https://doi.org/10.1016/j.cub.2014.10.045

Ferguson, K.A., Cardin, J.A., 2020. Mechanisms underlying gain modulation in the cortex. Nat. Rev. Neurosci. 21, 80–92. https://doi.org/10.1038/s41583-019-0253-y

Flagel, S.B., Clark, J.J., Robinson, T.E., Mayo, L., Czuj, A., Willuhn, I., Akers, C.A., Clinton, S.M., Phillips, P.E.M., Akil, H., 2011. A selective role for dopamine in stimulus–reward learning. Nature 469, 53–57. https://doi.org/10.1038/nature09588

Foote, S.L., Aston-Jones, G., Bloom, F.E., 1980. Impulse activity of locus coeruleus neurons in awake rats and monkeys is a function of sensory stimulation and arousal. Proc. Natl. Acad. Sci. U. S. A. 77, 3033–3037.

Greenstreet, F., Vergara, H.M., Pati, S., Schwarz, L., Wisdom, M., Marbach, F., Johansson, Y., Rollik, L., Moskovitz, T., Clopath, C.M., Stephenson-Jones, M., 2022. Action prediction error: a value-free dopaminergic teaching signal that drives stable learning. https://doi.org/10.1101/2022.09.12.507572

He, K., Huertas, M., Hong, S.Z., Tie, X., Hell, J.W., Shouval, H., Kirkwood, A., 2015. Distinct Eligibility Traces for LTP and LTD in Cortical Synapses. Neuron 88, 528–538. https://doi.org/10.1016/j.neuron.2015.09.037

Hervé-Minvielle, A., Sara, S.J., 1995. Rapid habituation of auditory responses of locus coeruleus cells in anaesthetized and awake rats. Neuroreport 6, 1363–1368. https://doi.org/10.1097/00001756-199507100-00001

Jordan, R., Keller, G.B., 2020. Opposing Influence of Top-down and Bottom-up Input on Excitatory Layer 2/3 Neurons in Mouse Primary Visual Cortex. Neuron. https://doi.org/10.1016/j.neuron.2020.09.024

Kasamatsu, T., Pettigrew, J.D., 1976. Depletion of brain catecholamines: failure of ocular dominance shift after monocular occlusion in kittens. Science 194, 206–209. https://doi.org/10.1126/science.959850

Kasamatsu, T., Watabe, K., Heggelund, P., Schöller, E., 1985. Plasticity in cat visual cortex restored by electrical stimulation of the locus coeruleus. Neurosci. Res. 2, 365–386. https://doi.org/10.1016/0168-0102(85)90047-1

Kebschull, J.M., Garcia da Silva, P., Reid, A.P., Peikon, I.D., Albeanu, D.F., Zador, A.M., 2016. High-Throughput Mapping of Single-Neuron Projections by Sequencing of Barcoded RNA. Neuron 91, 975–987. https://doi.org/10.1016/j.neuron.2016.07.036

Keller, G.B., Bonhoeffer, T., Hübener, M., 2012. Sensorimotor mismatch signals in primary visual cortex of the behaving mouse. Neuron 74, 809–815. https://doi.org/10.1016/j.neuron.2012.03.040

Keller, G.B., Mrsic-Flogel, T.D., 2018. Predictive Processing: A Canonical Cortical Computation. Neuron 100, 424–435. https://doi.org/10.1016/j.neuron.2018.10.003

Kim, H.R., Malik, A.N., Mikhael, J.G., Bech, P., Tsutsui-Kimura, I., Sun, F., Zhang, Y., Li, Y., Watabe-Uchida, M., Gershman, S.J., Uchida, N., 2020. A Unified Framework for Dopamine Signals across Timescales. Cell 183, 1600-1616.e25. https://doi.org/10.1016/j.cell.2020.11.013

Leinweber, M., Ward, D.R., Sobczak, J.M., Attinger, A., Keller, G.B., 2017. A Sensorimotor Circuit in Mouse Cortex for Visual Flow Predictions. Neuron 95, 1420-1432.e5. https://doi.org/10.1016/j.neuron.2017.08.036

Liu, Y., Rodenkirch, C., Moskowitz, N., Schriver, B., Wang, Q., 2017. Dynamic Lateralization of Pupil Dilation Evoked by Locus Coeruleus Activation Results from Sympathetic not Parasympathetic Contributions. Cell Rep. 20, 3099–3112. https://doi.org/10.1016/j.celrep.2017.08.094

Martins, A.R.O., Froemke, R.C., 2015. Coordinated forms of noradrenergic plasticity in the locus coeruleus and primary auditory cortex. Nat. Neurosci. 18, 1483–1492. https://doi.org/10.1038/nn.4090

McKinney, A., Hu, M., Hoskins, A., Mohammadyar, A., Naeem, N., Jing, J., Patel, S.S., Sheth, B.R., Jiang,
X., 2022. Cellular and Circuit Organization of the Locus Coeruleus of Adult Mice.
https://doi.org/10.1101/2022.03.01.481656
Meijering, E., Jacob, M., Sarria, J.-C.F., Steiner, P., Hirling, H., Unser, M., 2004. Design and validation of a
tool for neurite tracing and analysis in fluorescence microscopy images. Cytom. Part J. Int. Soc.
Anal. Cytol. 58, 167–176. https://doi.org/10.1002/cyto.a.20022
Mu, Y., Bennett, D.V., Rubinov, M., Narayan, S., Yang, C.-T., Tanimoto, M., Mensh, B.D., Looger, L.L.,
Ahrens, M.B., 2019. Glia Accumulate Evidence that Actions Are Futile and Suppress Unsuccessful
Behavior. Cell 178, 27-43.e19. https://doi.org/10.1016/j.cell.2019.05.050
Mukamel, E.A., Nimmerjahn, A., Schnitzer, M.J., 2009. Automated Analysis of Cellular Signals from Large-
Scale Calcium Imaging Data. Neuron 63, 747–760. https://doi.org/10.1016/j.neuron.2009.08.009
Niell, C.M., Stryker, M.P., 2010. Modulation of visual responses by behavioral state in mouse visual
cortex. Neuron 65, 472–479. https://doi.org/10.1016/j.neuron.2010.01.033
Nomura, S., Bouhadana, M., Morel, C., Faure, P., Cauli, B., Lambolez, B., Hepp, R., 2014. Noradrenalin
and dopamine receptors both control cAMP-PKA signaling throughout the cerebral cortex.
Front. Cell. Neurosci. 8. https://doi.org/10.3389/fncel.2014.00247
O'Toole, S.M., Oyibo, H.K., Keller, G.B., 2022. Prediction error neurons in mouse cortex are molecularly
targetable cell types. https://doi.org/10.1101/2022.07.20.500837
Pakan, J.M., Lowe, S.C., Dylda, E., Keemink, S.W., Currie, S.P., Coutts, C.A., Rochefort, N.L., 2016.
Behavioral-state modulation of inhibition is context-dependent and cell type specific in mouse
visual cortex. eLife 5, e14985. https://doi.org/10.7554/eLife.14985
Polack, P.-O., Friedman, J., Golshani, P., 2013. Cellular mechanisms of brain-state-dependent gain
modulation in visual cortex. Nat. Neurosci. 16, 1331–1339. https://doi.org/10.1038/nn.3464
Rao, R.P., Ballard, D.H., 1999. Predictive coding in the visual cortex: a functional interpretation of some
extra-classical receptive-field effects. Nat. Neurosci. 2, 79–87. https://doi.org/10.1038/4580
Reimer, J., McGinley, M.J., Liu, Y., Rodenkirch, C., Wang, Q., McCormick, D.A., Tolias, A.S., 2016. Pupil
fluctuations track rapid changes in adrenergic and cholinergic activity in cortex. Nat. Commun. 7,
13289. https://doi.org/10.1038/ncomms13289
Reynolds, J.N.J., Hyland, B.I., Wickens, J.R., 2001. A cellular mechanism of reward-related learning.
Nature 413, 67–70. https://doi.org/10.1038/35092560
Roelfsema, P.R., Holtmaat, A., 2018. Control of synaptic plasticity in deep cortical networks. Nat. Rev.
Neurosci. 19, 166–180. https://doi.org/10.1038/nrn.2018.6
Sales, A.C., Friston, K.J., Jones, M.W., Pickering, A.E., Moran, R.J., 2019. Locus Coeruleus tracking of
prediction errors optimises cognitive flexibility: An Active Inference model. PLOS Comput. Biol.
15, e1006267. https://doi.org/10.1371/journal.pcbi.1006267
Saravanan, V., Berman, G.J., Sober, S.J., 2020. Application of the hierarchical bootstrap to multi-level
data in neuroscience. Neurons Behav. Data Anal. Theory 3,
https://nbdt.scholasticahq.com/article/13927-application-of-the-hierarchical-bootstrap-to-
multi-level-data-in-neuroscience.
Schneider, D.M., Sundararajan, J., Mooney, R., 2018. A cortical filter that learns to suppress the acoustic
consequences of movement. Nature 561, 391–395. https://doi.org/10.1038/s41586-018-0520-5
Schultz, W., 2016. Dopamine reward prediction error coding. Dialogues Clin. Neurosci. 18, 23–32.
Schwarz, L.A., Miyamichi, K., Gao, X.J., Beier, K.T., Weissbourd, B., DeLoach, K.E., Ren, J., Ibanes, S.,
Malenka, R.C., Kremer, E.J., Luo, L., 2015. Viral-genetic tracing of the input–output organization
of a central noradrenaline circuit. Nature 524, 88–92. https://doi.org/10.1038/nature14600

1053 Seol, G.H., Ziburkus, J., Huang, S., Song, L., Kim, I.T., Takamiya, K., Huganir, R.L., Lee, H.-K., Kirkwood, A.,
1054    2007. Neuromodulators control the polarity of spike-timing-dependent synaptic plasticity.
1055    Neuron 55, 919–929. https://doi.org/10.1016/j.neuron.2007.08.013
1056 Shepard, K.N., Liles, L.C., Weinshenker, D., Liu, R.C., 2015. Norepinephrine is necessary for experience-
1057    dependent plasticity in the developing mouse auditory cortex. J. Neurosci. Off. J. Soc. Neurosci.
1058    35, 2432–2437. https://doi.org/10.1523/JNEUROSCI.0532-14.2015
1059 Steinberg, E.E., Keiflin, R., Boivin, J.R., Witten, I.B., Deisseroth, K., Janak, P.H., 2013. A causal link
1060    between prediction errors, dopamine neurons and learning. Nat. Neurosci. 16, 966–973.
1061    https://doi.org/10.1038/nn.3413
1062 Takeuchi, T., Duszkiewicz, A.J., Sonneborn, A., Spooner, P.A., Yamasaki, M., Watanabe, M., Smith, C.C.,
1063    Fernández, G., Deisseroth, K., Greene, R.W., Morris, R.G.M., 2016. Locus coeruleus and
1064    dopaminergic consolidation of everyday memory. Nature 537, 357–362.
1065    https://doi.org/10.1038/nature19325
1066 Vankov, A., Hervé-Minvielle, A., Sara, S.J., 1995. Response to novelty and its rapid habituation in locus
1067    coeruleus neurons of the freely exploring rat. Eur. J. Neurosci. 7, 1180–1187.
1068    https://doi.org/10.1111/j.1460-9568.1995.tb01108.x
1069 Vasilevskaya, A., Widmer, F.C., Keller, G.B., Jordan, R., 2022. Locomotion-induced gain of visual
1070    responses cannot explain visuomotor mismatch responses in layer 2/3 of primary visual cortex.
1071    https://doi.org/10.1101/2022.02.11.479795
1072 Wagatsuma, A., Okuyama, T., Sun, C., Smith, L.M., Abe, K., Tonegawa, S., 2018. Locus coeruleus input to
1073    hippocampal CA3 drives single-trial learning of a novel context. Proc. Natl. Acad. Sci. 115, E310–
1074    E316. https://doi.org/10.1073/pnas.1714082115
1075 Widmer, F.C., O'Toole, S.M., Keller, G.B., 2022. NMDA receptors in visual cortex are necessary for
1076    normal visuomotor integration and skill learning. eLife 11, e71476.
1077    https://doi.org/10.7554/eLife.71476
1078 Yu, A.J., Dayan, P., 2005. Uncertainty, neuromodulation, and attention. Neuron 46, 681–692.
1079    https://doi.org/10.1016/j.neuron.2005.04.026
1080 Zmarz, P., Keller, G.B., 2016. Mismatch Receptive Fields in Mouse Visual Cortex. Neuron 92, 766–772.
1081    https://doi.org/10.1016/j.neuron.2016.09.057
1082

1083