The genome of the bee louse fly reveals deep convergences in the evolution of social

inquilinism

Hélène Legout[1], David Ogereau[1], Julie Carcaud[1], Jonathan Filée[1], Lionel Garnery[1], Clément

Gilbert[1], Fabrice Requier[1], Jean-Christophe Sandoz[1], Amir Yassin[1], Héloïse Bastide[1]

[1]Laboratoire Évolution, Génomes, Comportement et Écologie, CNRS, IRD, Université Paris-

Saclay – Institut Diversité, Écologie et Évolution (IDEEV), 12 route 128, 91190 Gif-sur-

Yvette, France.

Corresponding author: Héloïse Bastide (heloise.bastide@universite-paris-saclay.fr)

Abstract

The nests of social insects often harbor a rich fauna of intruders, known as inquilines. Close relatedness between the host and the inquiline prevails due to potential genetic predispositions but how phylogenetically distant inquilines adapt to their hosts remains unclear. Here, we analyzed the genome of the wingless and blind bee louse fly *Braula coeca*, an inquiline of the Western honey bee *Apis mellifera*. We found that unlike many parasites, such as the human louse, the bee louse genome did not show significant erosion or strict reliance on an endosymbiont, likely due to a relatively recent age of inquilinism. However, a strikingly parallel evolution in a set of gene families was observed between the honey bee and the bee louse. Convergences included genes potentially involved in metabolism and immunity, and the loss of nearly all bitter-tasting gustatory receptors in agreement with life in a protective hive and a major diet of honey, pollens, and beeswax. Vision-related and odorant receptor genes also exhibited rapid losses. Only genes whose orthologs in the closely related *Drosophila melanogaster* respond to components of the honey bee alarm pheromones or floral aroma were retained, whereas the losses included orthologous receptors responsive to the anti-ovarian honey bee queen pheromone. These results establish a new model for the study of major morphological and neuroethological transitions and indicate that deep genetic convergences between phylogenetically distant organisms can underlie the evolution of social inquilinism.

Keywords: parasitism; morphological evolution; behavioral adaptations; gene family evolution; Drosophila.

Introduction

Inquilinism is a form of interspecific interactions wherein an organism, the inquiline, inhabits the "home" of another organism (Hegner 1926). Examples abound in nature (Moser 1964; Luczkovich *et al.* 1991; Sanver and Hawkins 2000; Kneitel and Miller 2002), and ancient cases are even present in the fossil record (Landman *et al.* 2014). The relationship between the inquiline and its host can range from commensalism to parasitism depending on the overlap in and availability of resources and space. Factors favoring the evolution of inquilinism greatly depend on the inquiline's capacity to hide its presence from the host. This is particularly true in the case of social inquilines, which constitute the most frequent case of inquilinism. wherein the efficient nest cleaning and care for the offspring performed by social organization are often fatal for the intruders. Therefore, phylogenetic relatedness with the host was suggested to play a major role in facilitating social inquilinism, since the inquiline would share some common genetic factors promoting camouflage, known as Emery's rule (Cini *et al.* 2019). Although recent phylogenetic studies have revised and sometimes rejected Emery's rule at the specific level (Huang and Dornhaus 2008; Lopez-Osorio *et al.* 2015; Romiguier *et al.* 2018; but see Savolainen and Vepsäläinen 2003; Degueldre *et al.* 2021), most cases of social inquilines remain related to the same genus (Jansen *et al.* 2010; Cardinal *et al.* 2010), tribe (Schrader *et al.* 2021), family (Ronquist 1994; Cardinal *et al.* 2010), or even order (Gilbert *et al.* 2012). The genetic basis underlying inquilines belonging to distinct orders or across wide phylogenetic distances is still less understood.

The hive of the Western honey bee *Apis mellifera* comprises several parasites and inquilines that are attracted by the hive's rich resources (honey, pollen, beeswax, and the bees themselves) and its clean and protective shelter (Winston 1987). These include several insects belonging to distinct orders such as wax moths, hive beetles, and endoparasitoid flies. None of these has endured as profound morphological changes that even its affiliation at the order-level was confounded, as the apterous and quasi-blind bee louse fly, *Braula coeca* (Figure 1A-C). The female lays eggs in honey (not brood) cells, and the hatched larvae eat pollen and wax, where they borrow tunnels in which they pupate without forming true puparia (Skaife 1922; Imms 1942). Following emergence, the adults attach to the body of worker bees, migrating from one individual to another until reaching the queen. There, they move to the head, stimulate regurgitation and imbibe from her mouth the honey and nectar (Skaife 1922; Imms 1942). The bee louse is considered an

inquiline kleptoparasite with potential negative effects on bee colony health due to

galleries in bee combs and the facilitation of transmitting serious pathogenic viruses to

the bees (Avalos *et al.* 2019).

The phylogenetic positioning within the Diptera of the family Braulidae, which

contains the bee louse, has long been puzzling due to its modified morphology (Grimaldi

and Underwood 1986). Interestingly, recent phylotranscriptomic and phylogenomic

analyses show the bee louse to constitute a basal lineage within the Drosophilidae

(Bayless *et al.* 2021; Winkler *et al.* 2022). This proximity to *Drosophila melanogaster*, the

most investigated insect at the genetic, developmental, and neurobiological levels, makes

the bee louse a unique model for the study of the genomic changes underlying major

morphological and ecological shifts. We present here an annotated assembly of the bee

louse *B. coeca* genome and compare the evolution of its genomic architecture and gene

content with those of *D. melanogaster* (Adams *et al.* 2000), the honey bee *A. mellifera*

(Weinstock *et al.* 2006), as well as to its homonym and morphologically-similar the

ectoparasitic human louse *Pediculus humanus* (Kirkness *et al.* 2010). The comparisons

revealed striking evidence of cross-order genomic parallelism and shed new light on the

evolution of social inquilinism between phylogenetically distant organisms.

Results

Genome architecture

We sequenced whole genome from a pooled sample of 15 unsexed *Braula coeca*

individuals, all collected from the same geographical location, the Island of Ouessant in

France. We used a hybrid approach to assemble a draft genome using both long-read

Oxford Nanopore Technology (ONT) and short-read Illumina sequencing (see Methods).

Size. The final assembly of the bee louse showed a size of 309 Mb, an N50 of 347227 bp

and a GC content of 34.95%. BUSCO gave a score of 95.8% of the Dipteran conserved

single-copy orthologs with 1.3% of duplicated genes. Analysis of heterozygous k-mers

pairs distribution indicated that the genome is diploid (Supplementary Figure 1A) and

genome size prediction using k-mers distribution spectra predict a genome size of 308

Mb, concordant with the assembly size (Supplementary Figure 1B). The bee louse genome

is therefore larger than the genomes of the honey bee (227 Mb), *Drosophila melanogaster*

(143 Mb), and the ectoparasitic human louse (108 Mb).

Endosymbionts. The taxonomic assignment of each contigs show that most of them (96%) match with arthropods indicating no (or very few) DNA contamination (Supplementary Figure 1C,D). Unlike in the human louse, no evidence for an obligate endosymbiont was detected in the bee louse. Imms (1942) discussed the possibility that the bee louse larvae have gut microbes that facilitate the digestion of the beeswax. We cannot rule out this hypothesis, but our results show that such microbiota, if present, likely do not persist in the adult stage.

Gene content. The annotation of the bee louse genome yielded 11,221 protein-coding genes. This number is higher than in *A. mellifera* and *P. humanus* where 9,935 and 10,773 protein-coding genes are found, respectively, but lower than that of *D. melanogaster* (13,968 protein-coding genes). The Annotation Edit Distance (AED), which measures the congruence between gene annotation and its supporting evidence was ≤ 0.5 for 96.7% of our gene models, indicating the near completeness of our annotation. Similarly, 79.79% of the corresponding proteins had a Pfam domain, which is another indication of annotation completeness since it varies between 57% and 75% in most eukaryotes.

Synteny. Orthologous genes were used to map *B. coeca* scaffolds to their corresponding Muller's elements in *D. melanogaster*. The alignment showed strong consistency indicating persistent synteny between the two lineages (Figure 1D). However, given the short length of the scaffolds it was difficult to assess how much collinearity and rearrangement events took place since their divergence. There is no karyotypic map of *B. coeca* so it remains difficult at this stage to infer the chromosomal number from the assembly alone.

Mitochondrial genome. The mitochondrial genome consisted of a single scaffold of 19 kb (Figure 1E) unlike in the human louse, which has multiple fragments inside the mitochondria. Compared to conserved gene content and collinearity of the *D. melanogaster* and *A. mellifera* mitogenomes (Crozier and Crozier 1993), the mitochondrial gene content of the bee louse was incomplete, lacking the *ND1* gene, included several duplications of the *ND2* and *ND4L* genes that are truncated and/or display frameshifts and had a rearrangement between the *COX1* and *COX2* genes. The nuclear genomes contained 165 mitochondrial DNA insertions (NUMTs) distributed on 88 contigs and totalizing 278 kb (~0.09% of the genome). The proportion of NUMTs in the bee louse approaches that of the honey bee (0.08%, (Behura 2007)) but exceeds that of the genus *Drosophila* (0.03%, (Rogers and Griffiths-Jones 2012).

Transposable elements. The large genome size of the bee louse despite its low gene content compared to *D. melanogaster* suggest an increase in repetitive sequences. RepeatModeler and RepeatMasker analyses indicated that nearly 41.41% of the *B. coeca* genome consist of such sequences, compared to 22.15% and 11.14% in *D. melanogaster* and *A. mellifera*, respectively (Supplementary Table 1). There is a far larger proportion of long interspersed nuclear elements (LINEs) retrotransposons in *B. coeca* (15.18%) compared to only 2.38% and none in *D. melanogaster* and *A. mellifera*, respectively. Long terminal repeat (LTR) elements on the other hand were fewer in *B. coeca* (0.57%) and *A. mellifera* (0.17%) than in *D. melanogaster* (6.99%).

Because host-parasite relationships have repeatedly be invoked as a factor that may favor horizontal transfer of transposable elements (TE) (Gilbert *et al.* 2010; Ortiz *et al.* 2015; Venner *et al.* 2017), we searched for evidence of such transfers between *B. coeca* and *A. mellifera* (Supplementary Text 1). We found one TE, *Famar1-like* element, previously described in the earwig *Forficula auricularia* (Barry *et al.* 2004a) that shows high similarity between *B. coeca* and *A. mellifera* but was absent in *D. melanogaster*, highly suggestive of acquisition through horizontal transfer (Supplementary Figure 2). However, phylogenetic analysis of multiple copies of this TE extracted from 37 widely divergent animal species (Supplementary Figure 3) shows that it was most likely acquired through independent transfer events in *B. coeca* and *A. mellifera*, and that the tight ecological connection between the two species are unlikely to have played a direct role in these transfers (Supplementary Text 1).

Gene family evolution

Families with excess losses. We identified gene families that underwent reduction or extension as compared to *D. melanogaster* using DAVID (Sherman *et al.* 2022). Underrepresented families (27 families with False Discovery Rate (FDR) < 0.05) showed a striking parallelism with the honey bee, with three of the most underrepresented families being similar (given in Honey bee Genome Sequencing Consortium 2006; Supplementary Table 2). These included InterPro-defined families such as Peptidase S1 (IPR001254), Ecdysteroid kinase-like (IPR004119) and Zinc finger AD-type (IPR012934).

Peptidases play a major role in detoxification and digestion of protein-rich diets. Their reduction in both the honey bee and the bee louse could be due to the overprotectiveness of their mutual habitat, the hive, and/or the low protein content of

their food, nectar and honey. The Ecdysteroid kinase-like (EcKL) family is also suspected
to include proteins involved in detoxification (Scanlan *et al.* 2020). We also noted a
functional affinity between two underrepresented families involved in the formation of
the exoskeleton: the Insect cuticle protein (IPR000618) in the honey bee and Chitin-
binding domain (IPR002557) in the bee louse. Cuticles could act as barriers against
environmental toxins, which may not be highly encountered in the hive. Remarkably, *B.*
coeca is unique among Cyclorrhaphan Diptera as its pupa, similarly to the honey bee's
(Winston 1987), is contained in the unmodified cuticle of the third instar larva, and no
sclerotized puparium is formed (Skaife 1922; Imms 1942).

Significant biological processes terms (FDR < 0.05) were mostly related to
proteolysis and lipid metabolism (e.g., fatty-acyl-co-A biosynthesis), whereas cellular
components terms were associated with the membranes and extracellular space in
agreement with a biased loss of genes potentially involved with metabolism,
detoxification and/or immunity (Supplementary Table 2).

Families with excess gains. We did not find a similar important overlap in
overrepresented gene families between the bee louse and the honey bee (5 families with
FDR < 0.05; Supplementary Table 3). Expanding gene families unique to the honey bee
included 7TM odorant receptor (IPR004117), Ankyrin (IPR002110), Yellow/royal jelly
protein (IPR003534) and LysR substrate-binding (IPR005119) (Weinstock *et al.* 2006).
We found one family, Pleckstrin homology domain (IPR001849), that was
overrepresented in both species. This domain is a part of several lipases that are known
to be involved in wax ingestion. Indeed, larvae of the bee louse mine the beeswax (Imms
1942), whereas honey bee workers use their salivary lipases while chewing the wax to
form the hive combs (Kurstjens *et al.* 1985).

Gene Ontology (GO) biological and cellular terms with significant enrichment at
FDR < 0.05 associated with morphological (e.g., imaginal disc-derived leg morphogenesis,
autophagy, and dorsal closure) and/or neurological (e.g., dendrite morphogenesis, axon
guidance, sensory perception of sound, and neuromuscular junction) developments
(Supplementary Table S3). The bee louse duplicated genes hence may play a role in the
evolution of the particular morphologies that helped adaptation to the phoretic lifestyle.

Vision. The species Latin name refers to the early assumption that the bee louse was blind
due to the reduction of the eye size and the loss of the ocelli. However, thin optic nerves
connect the brain to the rudimentary eyes, which lack ommatidia and pigments

(Müggenburg 1892), and the bee louse demonstrates negative phototaxis, indicating a certain degree of light perception (Kaschef 1959). In agreement with reduced vision in the bee louse, we found three out of the seven rhodopsin genes, which are responsible for colored vision and the positive phototaxis of *D. melanogaster*. Two of the three genes, *Rh1* and *Rh6*, are expressed in the ommatidia and are sensitive to light with long wavelengths (Senthilan and Helfrich-Förster 2016), whereas the third one, *Rh7*, is expressed in the brain and regulates light-dependent circadian entrainment (Ni *et al.* 2017). The role of these opsins in light detection despite the absence of ommatidia is unclear. Remarkably, *Rh1*, *Rh6* and *Rh7* are structurally required in mechanosensory bristles to control larval locomotion (Zanini *et al.* 2018), and *Rh1* and *Rh6* also detect temperature (Leung and Montell 2017), whereas *Rh1* and *Rh7* detect low concentrations of a bitter plant component (Leung *et al.* 2020). Therefore, the retention of these rhodopsins in the bee louse could mainly be due to their unconventional functions. On the other hand, the red-sensitive rhodopsin *Rh2*, which is exclusively expressed in the ocelli and used for horizon detection in *D. melanogaster* (Mishra *et al.* 2021), is among those which were lost in the bee louse, in agreement with the loss of the ocelli in the bee louse.

Olfaction. Odorant receptors (ORs) are essential to detect volatile chemical signals from the environment. In most *Drosophila* species, there are nearly 61 ORs, whereas in the honey bee this family has expanded to reach 160 (Robertson and Wanner 2006). We found 29 ORs in the bee louse in addition to the Orco ortholog. Of these four had no direct orthologs in *D. melanogaster*, but the remaining 25 genes were orthologous to 21 genes in *D. melanogaster* and 12 were duplicates specific to the bee louse lineage (Figure 2A; Supplementary Figure 3). Judging from the response of those ORs to different volatiles in *D. melanogaster* as curated in the DOOR database (Münch and Galizia 2016) and assuming potential conservation of function, we can divide the bee louse ORs into three categories. First, 11 ORs respond to different components of the honey bee workers alarm and mandibular gland pheromones, e.g., 2-heptanol, propyl acetate, 2-heptanone, 1-hexanol, butyl acetate, isopentyl acetate, etc. These included *Or13a* (x2 paralogues), *Or42a*, *Or42b/Or59b*, *Or43a*, *Or47b*, *Or49a/Or85f*, *Or67b* (x4), *Or74a*, *Or85b/Or85c*, and *Or85e* (x2). Second, 12 ORs respond to different floral, pollen and nectar aromas, such as acetophenones, ethylguaiacol, geranyl acetate or fenchone, but most importantly to benzaldehyde, a major volatile of honey (Machado *et al.* 2020; Starowicz *et al.* 2021). These included *Or30a*, *Or45b* (x2), *Or59a*, *Or67b* (x4), *Or82a*, *Or85e* (x2) and *Or94a*. Note

that the multiple *B. coeca* paralogs orthologous to *D. melanogaster Or67b* and *Or85e* genes belong to both categories. Third, one OR, *Or67d*, which is involved in sexual pheromone 11-cis-vaccenyl acetate perception (Ha and Smith 2006), is present in the bee louse. Three among the ORs that were lost in the bee louse compared to other drosophilids, *Or49b*, *Or56a* and *Or98a*, are responsible to *D. melanogaster* response to the anti-ovarian honey bee queen mandibular pheromones 9-oxo-2-decenoic acid (9ODA) and 10-hydroxy-2-decenoic acid (10HDA) (Galang *et al.* 2019). Their loss most likely protected the capacity of the bee louse to reproduce in the hive.

Taste. Gustatory receptors (GRs) allow detecting soluble chemical signals. There are nearly 68 GRs in *D. melanogaster*, that respond mostly to sweet, bitter and carbon dioxide (CO_2) tastes (Weiss *et al.* 2011). Unlike their expanded OR family, the honey bee has only 10 GRs, of which 7 are orthologous to sweet *Drosophila* GRs (Robertson and Wanner 2006). This is likely due to the bees' food reliance on sweet floral nectars and honey. We found 13 GRs in the bee louse, with no duplications (Figure 2B; Supplementary Figure 4). These GRs could be classified according to their *D. melanogaster* orthology into three categories. First, 6 GRs belong to the sweet class, namely *Gr43a*, *Gr61a* and *Gr64a,b,e,f* which usually respond to sucrose, maltose and fructose and other major honey sugars. Second, 5 GRs belong to the bitter class, namely *Gr33a*, *Gr57a*, *Gr66a*, *Gr93a* and *Gr94a*, suggesting that nearly 50 mostly bitter tasting drosophilid GRs were lost in the bee louse. Note that in *D. melanogaster*, *Gr33a* and *Gr66a*, are potentially involved in sexual pheromones detection (Lacaille *et al.* 2007; Moon *et al.* 2009). Third, 2 GRs, *Gr21a* and *Gr63a*, which detects CO_2 odor in *D. melanogaster* (Jones *et al.* 2007) are present in the bee louse. Those GRs are absent in the honey bee despite its ability to perceive CO_2 (Stange and Diesendorf 1973). High CO_2 concentrations are probably characteristic of largely populated hives and induce fanning response in bees (Seeley 1974). The quasi-blind bee louse may therefore use CO_2 concentrations to detect the bees in the dark hive environment.

Discussion

That the enigmatic bee louse is indeed a drosophilid (Winkler *et al.* 2022), a lineage within the most investigated insect family with more than 100 fully sequenced genomes (Kim *et al.* 2021), is undoubtedly one of the most exciting discoveries in Dipteran phylogeny. How could a fly with a typical drosophilid genome become partly ecologically

like a bee and partly morphologically like a louse? Our cross-order comparisons of the bee louse to its homonyms shed significant light on the genomic basis of these spectacular convergences.

The genome of the human louse is among the smallest sequenced insect genomes (Kelley *et al.* 2014). Loss of significant portions of genomic and gene contents is a characteristic of obligate parasites specializing on specific hosts or inhabiting extreme environments. Indeed, the gene content of the bee louse genome approached that of the human louse, but the ~309 Mb-long genome of the bee louse is longer than that of most drosophilid species, even being slightly longer from the largest genome in this family (~304 Mb-long; Kim *et al.* 2021). Besides, the human louse has an obligatory bacterial endosymbiont that may compensate the loss of its genes (Kirkness *et al.* 2010) but no evidence for such associations is present in the bee louse. These differences may mainly indicate the relative recency of the shift to inquilinism in the bee louse compared to the 230 million years (myr) of specialization in true lice (Misof *et al.* 2014). *Megabraula*, the closest relative to the genus *Braula*, is an inquiline of the giant honey bee *Apis laboriosa*, whose divergence from the Western honey bee *A. mellifera* is around 23 myr ago (Grimaldi and Underwood 1986; Cardinal *et al.* 2010). The crown age of the Drosophilidae and the divergence time between *Apis* and its closest pollen-basket (corbiculate) bees are estimated at 66-70 myr ago (Cardinal *et al.* 2010; Suvorov *et al.* 2021). This indicates that association between braulids and *Apis* has arisen between 70 to 20 myr ago.

The origin of social inquilinism requires the evolution of multiple convergences that can locate the host, deceive it to enter the colony, adapt to its social organization, and remain undetected or not easily removed from the colony. These phenotypes, which are often host-specific, evolve more easily when the inquiline and the host are phylogenetically close (Cini *et al.* 2019). Despite the distance of the bee louse from honey bees, gene family analysis provided strong evidence for convergent evolution mostly for genes potentially involved in immunity, detoxification, metabolism and chemical perception. Although *Braula* has lost nearly half of the typical drosophilids odorant receptor repertoire, in contrast to the major expansion of this family in the honey bee, it predominantly retained genes whose orthologs in *D. melanogaster* detect compounds of the honey bee pheromones and/or floral aroma and honey odors. Low concentrations of isopentyl acetate (IPA), the major component of the alarm pheromone, released by unstressed workers at hive entries attract the parasitic hive beetle *Aethina tumida* (Torto

et al. 2007), suggesting that the detection of the host odors could be a common strategy
among phylogenetically distant social inquilines and parasites. We also noted that the bee
louse has multiple copies related to *Or67b* which responds to several components of bee
alarm pheromones, such as 1-hexanol, 2-heptanone, 1-butanol and 3-methyl-1-butanol,
as well as to benzaldehyde, the major honey volatile, in *D. melanogaster* (Münch and
Galizia 2016), suggesting a possible dual ancestral function that might have facilitated
association with the honey bees. The multiple copies related to *Or67b* are remarkable
since the copy number expansion of this gene was associated with the evolution of
herbivory and strong plant-association in the drosophilid genus *Scaptomyza* (Goldman-
Huertas *et al.* 2015; Matsunaga *et al.* 2022). Therefore, the bee louse might have evolved
from a flower-breeding or plant-associated lineage within the Drosophilidae, a lifestyle
that has recurrently evolved in this family (Yassin 2013).

Whereas major molecular convergences could exist between the inquiline and its
social host, divergent strategies to adapt to the eusocial lifestyle requirements are still
needed. The loss of the three odorant receptors, *Or56a*, *Or49b* and *Or98a*, that respond in
D. melanogaster to the honey bee queen's pheromones which "sterilize" the bee workers,
mainly 9-oxo-2-decenoic acid (9ODA) and possibly 9-hydroxy-2-decenoic acid (9HDA)
(Galang *et al.* 2019), is a notable example. The queen's pheromones elicit anti-ovarian
response in other insects including *D. melanogaster* mostly through the activation of the
neurons bearing these receptors. Therefore, a *sine qua non* condition for reproducing in a
beehive is to protect against the effects of those pheromones, with the loss of the
responding receptors being a preliminary and effective strategy. However, this raises the
question of how the bee lice recognize the queen, which they preferentially infest (Imms
1942). It is therefore possible that higher sensitivity to worker pheromones, as suggested
by the repertoire of retained odorant receptors (see above), may help to mostly
discriminate the workers hence facilitating the recognition of the queen, who has a
rudimentary sting gland. Another possibility is that strong CO_2 emission by the court
surrounding the queen may be an indicator of her location, as could be suggested from
the retention of the two CO_2-smelling gustatory receptors, *Gr21a* and *Gr63a*, whose
orthologs are absent in the honey bee. It also remains unclear how the bee lice sexually
communicate in the hive given the predominance of chemical camouflage; the
hydrocarbon profile of bee lice from different colonies mimicked that of the colony of
origin (Martin and Bayfield 2014). The retention of some chemosensory genes whose

orthologs are involved in sexual pheromone detection in *D. melanogaster*, such as *Or67d*, *Gr33a* and *Gr66a* (Kurtovic *et al.* 2007; Lacaille *et al.* 2007; Moon *et al.* 2009) suggest that pheromonal communications may still be present in the bee louse but perhaps at low undetectable levels.

Small size, loss of wings and the evolution of strongly clinging legs are all morphological adaptations that could prevent the honey bees getting rid of the bee lice. All these adaptations are convergent with ectoparasitic true lice, and for some, such as apterism, represent major recurrent changes that have responded to distinct pressures throughout the history of insects (Roff 1990). We found intact most of the main wing development genes whose mutations severely reduce the wing in *D. melanogaster*, such as *wingless*, *apterous* or *vestigial*. This means that the major morphological changes more likely resulted from regulatory changes of these core genes or modifications of other genes. The regression of visual systems and their underlying genes in the bee louse spending most of its life cycle in the bee hives is a common phenomenon in animals inhabiting dark environments, such as fossorial mammals (Partha *et al.* 2017) and cavefishes (Policarpo *et al.* 2021). With its genetic relatedness to *Drosophila* and ecological association to *Apis*, two major laboratory models, and with new genomic tools presented here, the bee louse *Braula coeca* is a promising model to address questions related to deep convergences that are still difficult to approach in multiple highly specializing animals.

Materials and Methods

Sample collection and genomic library preparation

Samples of *Braula coeca* were collected from honey bee colonies on the Island of Ouessant in France and kindly provided to us by the *Association Conservatoire de l'Abeille Noire Bretonne* (A.C.A.N.B.). Genomic DNA was extracted from 15 unsexed individuals conserved in alcohol, using the Nucleobond AXG20 kit and buffer set IV from Macherey-Nagel (ref. 740544 and 740604, https://www.mn-net.com, Düren, Germany).

Genome sequencing and assembly

We used a hybrid approach to assemble a draft genome of *B. coeca* using both long-read Oxford Nanopore Technology (ONT) and short-read Illumina sequencing, as in Miller et al. (2019). Before nanopore sequencing, a size selection was conducted on the DNA

using the SRE XS from Circulomics (https://www.circulomics.com/, Baltimore, Maryland,

USA). The SQK-LSK110 kit from Oxford Nanopore Technology (Lu *et al.* 2016;

https://nanoporetech.com/) was then used to prepare the samples for nanopore

sequencing following manufacturer's protocol. The library was loaded and sequenced on

a R9.4.1 flow cell (ref FLO-Min106) for sequencing. Raw data were basecalled using

Guppy v5.0.11 and the "sup" algorithm. Illumina paired-end sequencing was performed

by Novogene Company Limited (https://en.novogene.com, Cambridge, UK) on the same

DNA library.

We used MaSuRCA v4.0.3 (Zimin *et al.* 2017) to produce the hybrid assembly of

our genome, using the Cabog assembler. We obtained a final assembly size of 309,35Mb

in 2477 contigs, with a N50 of 347211 pb. The completeness of the assembly was

estimated to 95,8% with Busco v5.0 on the *diptera_odb10* dataset

(C:95.8%[S:94.6%,D:1.2%],F:0.7%,M:3.5%,n:3285).

Estimation of genome size, endosymbionts detection and mitogenome assembly

K-mers frequencies within short-read data were obtained with KMC 3 (Kokot *et al.*

2017). Genome size and ploidy were inferred using GenomeScope v2.0 with k-mer size =

21 and Smudgeplot (Ranallo-Benavidez *et al.* 2020). Contig taxonomy was performed

using Blobtools (Laetsch and Blaxter 2017) with Diamond as search engine (Buchfink *et*

al. 2015) against the UniProt database using a local copy of the NCBI TaxID file for the

taxonomic assignation of the best hit. Minimap2 (Li 2018) was used for read mapping.

Genome annotation

The *B. coeca* genome was annotated using Maker v2.31.10 (Cantarel *et al.* 2008),

following the protocol given in Muller *et al.* (2021), wherein multiple rounds of Maker

supported by the training of the SNAP v.2006-07-28 (Korf 2004) and Augustus v.3.3.3

(König *et al.* 2016) gene finding and prediction tools, were conducted. Transcriptome of

B. coeca (NCBI accession no. SRR2046564; 1KITE Consortium; (Bayless *et al.* 2021)) and

proteomes of five *Drosophila* species, namely *D. innubila* (Hill *et al.* 2019), *D. albomicans*

(Mai *et al.* 2020), *D. bipectinata* (Kim *et al.* 2021), *D. melanogaster* (Adams *et al.* 2000) and

D. virilis (Clark *et al.* 2007) were used to guide the annotation. Protein-Protein BLAST

2.9.0+ (Altschul *et al.* 1997) (-evalue 1e-6 -max_hsps 1 -max_target_seqs 1) was then used

to assess putative protein functions in *B. coeca* by comparing the protein sequences given

by Maker to the protein sequences from the annotated genome of *D. melanogaster*. The

colocalization of *B. coeca* and their *D. melanogaster* orthologs on their respective contigs

and chromosomal arms was analyzed to test for synteny.

Transposable elements annotation and transfer

Transposable elements were identified in each species following a two-step

protocol. First, we used RepeatModeler v 2.0.1 (Flynn *et al.* 2020) with default parameters

to generate a *de novo* library of repetitive regions. RepeatMasker v 4.0.9 (Flynn *et al.*

2020) was then run with the newly generated library and the options -a (create a .align

output file) and -s (slow search; more sensitive) to create a summary of the families of

transposable elements found in each reference genome along with the percentage of the

genome they represent. Horizontal transfer analyses protocols of the *Famar1*-like

element are given in Supplementary Text 1.

Gene family evolution

The Database for Annotation, Visualization and Integrated Discovery DAVID 2021

(Sherman *et al.* 2022) was used to test for gene ontology (GO) terms enrichments among

lists of *D. melanogaster* orthologs annotated on the *B. coeca* genome (see above). The

program was either fed with the list of *D. melanogaster* genes that were absent or

duplicated in the *B. coeca* genome to test for under- and overrepresented gene families,

respectively, compared to *D. melanogaster* genome, as denoted by the False Discovery

Rate (FDR) correction for multiple tests. Each analysis was conducted for biological

processes, molecular function, and cytological components.

Chemosensory superfamilies evolution

Protein sequences of the odorant (ORs) and gustatory (GRs) receptors of *A.*

mellifera and *D. melanogaster* were obtained from Robertson and Wanner (2006) to

which we added the annotated ORs and GRs protein sequences of *B. coeca*. Sequences

were aligned using Molecular Evolutionary Genetic Analysis (MEGA X) software package

(Kumar *et al.* 2018), which was also used to infer a maximum-likelihood phylogenetic tree

for each family. We used iToL v4 (Letunic and Bork 2019) to visualize the trees. *B. coeca*

protein sequences are given in Supplementary Datasets 2 and 3 for ORs and GRs,

respectively.

Acknowledgments

We are very grateful to Dr. Frédéric Marion-Poll for insightful comments on an early draft of the manuscript and to the *Association Conservatoire de l'Abeille Noire Bretonne* (A.C.A.N.B.) for help collecting *B. coeca* flies. *Braula* genome sequencing was funded by a grant from Université Paris Saclay (ADAPAR) to HB.

Conflicts of interest

The authors declare no conflicts of interest.

References

Adams M. D., S. E. Celniker, R. A. Holt, C. A. Evans, J. D. Gocayne, *et al.*, 2000 The Genome Sequence of Drosophila melanogaster. Science 287: 2185–2195. https://doi.org/10.1126/science.287.5461.2185

Altschul S. F., T. L. Madden, A. A. Schäffer, J. Zhang, Z. Zhang, *et al.*, 1997 Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucl Acids Res 25: 3389–3402. https://doi.org/10.1093/nar/25.17.3389

Avalos J., H. Rosero, G. Maldonado, and F. J. Reynaldi, 2019 Honey bee louse (Braula schmitzi) as a honey bee virus vector? Journal of Apicultural Research 58: 427–429. https://doi.org/10.1080/00218839.2019.1565726

Barry E. G., D. J. Witherspoon, and D. J. Lampe, 2004a A Bacterial Genetic Screen Identifies Functional Coding Sequences of the Insect *mariner* Transposable Element *Famar1* Amplified From the Genome of the Earwig, *Forficula auricularia*. Genetics 166: 823–833. https://doi.org/10.1534/genetics.166.2.823

Barry E. G., D. J. Witherspoon, and D. J. Lampe, 2004b A Bacterial Genetic Screen Identifies Functional Coding Sequences of the Insect mariner Transposable Element Famar1 Amplified From the Genome of the Earwig, Forficula auricularia. Genetics 166: 823–833. https://doi.org/10.1093/genetics/166.2.823

Bayless K. M., M. D. Trautwein, K. Meusemann, S. Shin, M. Petersen, *et al.*, 2021 Beyond Drosophila: resolving the rapid radiation of schizophoran flies with phylotranscriptomics. BMC Biol 19: 23. https://doi.org/10.1186/s12915-020-00944-8

Behura S. K., 2007 Analysis of Nuclear Copies of Mitochondrial Sequences in Honeybee (Apis mellifera) Genome. Molecular Biology and Evolution 24: 1492–1505.

https://doi.org/10.1093/molbev/msm068

Buchfink B., C. Xie, and D. H. Huson, 2015 Fast and sensitive protein alignment using

DIAMOND. Nat Methods 12: 59–60. https://doi.org/10.1038/nmeth.3176

Cantarel B. L., I. Korf, S. M. C. Robb, G. Parra, E. Ross, *et al.*, 2008 MAKER: An easy-to-use

annotation pipeline designed for emerging model organism genomes. Genome Res. 18: 188–

196. https://doi.org/10.1101/gr.6743907

Cardinal S., J. Straka, and B. N. Danforth, 2010 Comprehensive phylogeny of apid bees reveals

the evolutionary origins and antiquity of cleptoparasitism. Proceedings of the National

Academy of Sciences 107: 16207–16211. https://doi.org/10.1073/pnas.1006299107

Cini A., S. Sumner, and R. Cervo, 2019 Inquiline social parasites as tools to unlock the secrets

of insect sociality. Philosophical Transactions of the Royal Society B: Biological Sciences 374:

20180193. https://doi.org/10.1098/rstb.2018.0193

Clark A. G., M. B. Eisen, D. R. Smith, C. M. Bergman, B. Oliver, *et al.*, 2007 Evolution of

genes and genomes on the Drosophila phylogeny. Nature 450: 203–218.

https://doi.org/10.1038/nature06341

Crozier R. H., and Y. C. Crozier, 1993 The mitochondrial genome of the honeybee Apis

mellifera: complete sequence and genome organization. Genetics 133: 97–117.

https://doi.org/10.1093/genetics/133.1.97

Degueldre F., P. Mardulyn, A. Kuhn, A. Pinel, C. Karaman, *et al.*, 2021 Evolutionary history

of inquiline social parasitism in Plagiolepis ants. Molecular Phylogenetics and Evolution 155:

107016. https://doi.org/10.1016/j.ympev.2020.107016

Emery C., 1909 Über den Ursprung der dulotischen, parasitischen und myrmekophilen

Ameisen. Biologisches Centralblatt 29: 352–362.

Flynn J. M., R. Hubley, C. Goubert, J. Rosen, A. G. Clark, *et al.*, 2020 RepeatModeler2 for

automated genomic discovery of transposable element families. Proceedings of the National

Academy of Sciences 117: 9451–9457. https://doi.org/10.1073/pnas.1921046117

Galang K. C., J. R. Croft, G. J. Thompson, and A. Percival-Smith, 2019 Analysis of the

Drosophila melanogaster anti-ovarian response to honey bee queen mandibular pheromone.

Insect Molecular Biology 28: 99–111. https://doi.org/10.1111/imb.12531

Gilbert C., S. Schaack, J. K. Pace, P. J. Brindley, and C. Feschotte, 2010 A role for host-parasite

interactions in the horizontal transfer of transposons across phyla. Nature 464: 1347–50.

https://doi.org/10.1038/nature08939

Gilbert J. D. J., L. A. Mound, and S. J. Simpson, 2012 Biology of a new species of socially

parasitic thrips (Thysanoptera: Phlaeothripidae) inside Dunatothrips nests, with evolutionary

implications for inquilinism in thrips. Biological Journal of the Linnean Society 107: 112–122.

https://doi.org/10.1111/j.1095-8312.2012.01928.x

Goldman-Huertas B., R. F. Mitchell, R. T. Lapoint, C. P. Faucher, J. G. Hildebrand, *et al.*, 2015

Evolution of herbivory in Drosophilidae linked to loss of behaviors, antennal responses, odorant

receptors, and ancestral diet. PNAS 112: 3026–3031. https://doi.org/10.1073/pnas.1424656112

Grimaldi D., and B. A. Underwood, 1986 Megabraula, a new genus for two new species of

Braulidae (Diptera), and a discussion of braulid evolution. Systematic Entomology 11: 427–

438. https://doi.org/10.1111/j.1365-3113.1986.tb00534.x

Ha T. S., and D. P. Smith, 2006 A Pheromone Receptor Mediates 11-cis-Vaccenyl Acetate-

Induced Responses in Drosophila. Journal of Neuroscience 26: 8727–8733.

https://doi.org/10.1523/JNEUROSCI.0876-06.2006

Hegner R. W., 1926 The Biology of Host-Parasite Relationships Among Protozoa Living in

Man. The Quarterly Review of Biology 1: 393–418.

Hill T., B. S. Koseva, and R. L. Unckless, 2019 The Genome of Drosophila innubila Reveals

Lineage-Specific Patterns of Selection in Immune Genes. Molecular Biology and Evolution 36:

1405–1417. https://doi.org/10.1093/molbev/msz059

Huang M. H., and A. Dornhaus, 2008 A meta-analysis of ant social parasitism: host

characteristics of different parasitism types and a test of Emery's rule. Ecological Entomology

33: 589–596. https://doi.org/10.1111/j.1365-2311.2008.01005.x

Imms A. D., 1942 On Braula coeca Nitsch and its affinities. Parasitology 34: 88–100.

https://doi.org/10.1017/S0031182000016012

Jansen G., R. Savolainen, and K. Vepsäläinen, 2010 Phylogeny, divergence-time estimation,

biogeography and social parasite–host relationships of the Holarctic ant genus Myrmica

(Hymenoptera: Formicidae). Molecular Phylogenetics and Evolution 56: 294–304.

https://doi.org/10.1016/j.ympev.2010.01.029

Jones W. D., P. Cayirlioglu, I. Grunwald Kadow, and L. B. Vosshall, 2007 Two chemosensory

receptors together mediate carbon dioxide detection in Drosophila. Nature 445: 86–90.

https://doi.org/10.1038/nature05466

Kaschef A.-H., 1959 The sensory physiology and behaviour of the Honeybee LouseBraula

coeca Nitzcsh (Diptera, Braulidae). Ins. Soc 6: 313–342. https://doi.org/10.1007/BF02225778

Kelley J. L., J. T. Peyton, A.-S. Fiston-Lavier, N. M. Teets, M.-C. Yee, *et al.*, 2014 Compact

genome of the Antarctic midge is likely an adaptation to an extreme environment. Nat Commun

5: 4611. https://doi.org/10.1038/ncomms5611

Kim B. Y., J. R. Wang, D. E. Miller, O. Barmina, E. Delaney, *et al.*, 2021 Highly contiguous

assemblies of 101 drosophilid genomes, (G. Coop, P. J. Wittkopp, and T. B. Sackton, Eds.). eLife 10: e66405. https://doi.org/10.7554/eLife.66405

Kirkness E. F., B. J. Haas, W. Sun, H. R. Braig, M. A. Perotti, *et al.*, 2010 Genome sequences of the human body louse and its primary endosymbiont provide insights into the permanent parasitic lifestyle. Proceedings of the National Academy of Sciences 107: 12168–12173. https://doi.org/10.1073/pnas.1003379107

Kneitel J. M., and T. E. Miller, 2002 Resource and Top-Predator Regulation in the Pitcher Plant (sarracenia Purpurea) Inquiline Community. Ecology 83: 680–688. https://doi.org/10.1890/0012-9658(2002)083[0680:RATPRI]2.0.CO;2

Kokot M., M. Długosz, and S. Deorowicz, 2017 KMC 3: counting and manipulating k-mer statistics. Bioinformatics 33: 2759–2761. https://doi.org/10.1093/bioinformatics/btx304

König S., L. W. Romoth, L. Gerischer, and M. Stanke, 2016 Simultaneous gene finding in multiple genomes. Bioinformatics 32: 3388–3395. https://doi.org/10.1093/bioinformatics/btw494

Korf I., 2004 Gene finding in novel genomes. BMC Bioinformatics 5: 59. https://doi.org/10.1186/1471-2105-5-59

Kumar S., G. Stecher, M. Li, C. Knyaz, and K. Tamura, 2018 MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. Molecular Biology and Evolution 35: 1547–1549. https://doi.org/10.1093/molbev/msy096

Kurstjens S. P., H. R. Hepburn, F. R. L. Schoening, and B. C. Davidson, 1985 The conversion of wax scales into comb wax by African honeybees. J Comp Physiol B 156: 95–102. https://doi.org/10.1007/BF00692930

Kurtovic A., A. Widmer, and B. J. Dickson, 2007 A single class of olfactory neurons mediates behavioural responses to a Drosophila sex pheromone. Nature 446: 542–546. https://doi.org/10.1038/nature05672

Lacaille F., M. Hiroi, R. Twele, T. Inoshita, D. Umemoto, *et al.*, 2007 An Inhibitory Sex Pheromone Tastes Bitter for Drosophila Males. PLOS ONE 2: e661. https://doi.org/10.1371/journal.pone.0000661

Laetsch D. R., and M. L. Blaxter, 2017 BlobTools: Interrogation of genome assemblies

Landman N. H., R. H. B. Lraaije, S. M. Klofak, N. L. Larson, G. A. Bishop, *et al.*, 2014 Inquilinism of a Baculite by a Dynomenid Crab from the Upper Cretaceous of South Dakota. novi 2014: 1–16. https://doi.org/10.1206/3818.1

Letunic I., and P. Bork, 2019 Interactive Tree Of Life (iTOL) v4: recent updates and new developments. Nucleic Acids Research 47: W256–W259. https://doi.org/10.1093/nar/gkz239

Leung N. Y., and C. Montell, 2017 Unconventional Roles of Opsins. Annu. Rev. Cell Dev.

Biol. 33: 241–264. https://doi.org/10.1146/annurev-cellbio-100616-060432

Leung N. Y., D. P. Thakur, A. S. Gurav, S. H. Kim, A. Di Pizio, *et al.*, 2020 Functions of

Opsins   in   Drosophila   Taste.   Current   Biology   30:   1367-1379.e6.

https://doi.org/10.1016/j.cub.2020.01.068

Li H., 2018 Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34: 3094–

3100. https://doi.org/10.1093/bioinformatics/bty191

Lopez-Osorio F., A. Perrard, K. M. Pickett, J. M. Carpenter, and I. Agnarsson, 2015

Phylogenetic tests reject Emery's rule in the evolution of social parasitism in yellowjackets and

hornets (Hymenoptera: Vespidae, Vespinae). Royal Society Open Science 2: 150159.

https://doi.org/10.1098/rsos.150159

Lu H., F. Giordano, and Z. Ning, 2016 Oxford Nanopore MinION Sequencing and Genome

Assembly.   Genomics,   Proteomics   &   Bioinformatics   14:   265–279.

https://doi.org/10.1016/j.gpb.2016.05.004

Luczkovich J. J., G. M. Watters, and B. L. Olla, 1991 Seasonal Variation in Usage of a Common

Shelter Resource by Juvenile Inquiline Snailfish (Liparis inquilinus) and Red Hake (Urophycis

chuss). Copeia 1991: 1104–1109. https://doi.org/10.2307/1446107

Machado A. M., M. G. Miguel, M. Vilas-Boas, and A. C. Figueiredo, 2020 Honey Volatiles as

a Fingerprint for Botanical Origin—A Review on their Occurrence on Monofloral Honeys.

Molecules 25: 374. https://doi.org/10.3390/molecules25020374

Mai D., M. J. Nalley, and D. Bachtrog, 2020 Patterns of Genomic Differentiation in the

Drosophila nasuta Species Complex. Molecular Biology and Evolution 37: 208–220.

https://doi.org/10.1093/molbev/msz215

Martin S. J., and J. Bayfield, 2014 Is the bee louse Braula coeca (Diptera) using chemical

camouflage   to   survive   within   honeybee   colonies?   Chemoecology   24:   165–169.

https://doi.org/10.1007/s00049-014-0158-1

Matsunaga T., C. E. Reisenman, B. Goldman-Huertas, P. Brand, K. Miao, *et al.*, 2022 Evolution

of Olfactory Receptors Tuned to Mustard Oils in Herbivorous Drosophilidae. Molecular

Biology and Evolution 39: msab362. https://doi.org/10.1093/molbev/msab362

Mishra A. K., C. Fritsch, R. Voutev, R. S. Mann, and S. G. Sprecher, 2021 Homothorax controls

a binary Rhodopsin switch in Drosophila ocelli. PLOS Genetics 17: e1009460.

https://doi.org/10.1371/journal.pgen.1009460

Misof B., S. Liu, K. Meusemann, R. S. Peters, A. Donath, *et al.*, 2014 Phylogenomics resolves

the   timing   and   pattern   of   insect   evolution.   Science   346:   763–767.

603 https://doi.org/10.1126/science.1257570

604 Moon S. J., Y. Lee, Y. Jiao, and C. Montell, 2009 A Drosophila Gustatory Receptor Essential

605 for Aversive Taste and Inhibiting Male-to-Male Courtship. Current Biology 19: 1623–1627.

606 https://doi.org/10.1016/j.cub.2009.07.061

607 Moser J. C., 1964 Inquiline Roach Responds to Trail-Marking Substance of Leaf-Cutting Ants.

608 Science 143: 1048–1049. https://doi.org/10.1126/science.143.3610.1048

609 Müggenburg F. H., 1892 *Der Rüssel der Diptera pupipara*. Nicolai.

610 Muller H., D. Ogereau, J.-L. Da Lage, C. Capdevielle, N. Pollet, *et al.*, 2021 Draft nuclear

611 genome and complete mitogenome of the Mediterranean corn borer, Sesamia nonagrioides, a

612 major pest of maize. G3 Genes|Genomes|Genetics 11: jkab155.

613 https://doi.org/10.1093/g3journal/jkab155

614 Münch D., and C. G. Galizia, 2016 DoOR 2.0 - Comprehensive Mapping of *Drosophila*

615 *melanogaster* Odorant Responses. Scientific Reports 6: 21841.

616 https://doi.org/10.1038/srep21841

617 Ni J. D., L. S. Baik, T. C. Holmes, and C. Montell, 2017 A rhodopsin in the brain functions in

618 circadian photoentrainment in Drosophila. Nature 545: 340–344.

619 https://doi.org/10.1038/nature22325

620 Ortiz M. F., G. L. Wallau, D. A. Graichen, and E. L. Loreto, 2015 An evaluation of the

621 ecological relationship between Drosophila species and their parasitoid wasps as an opportunity

622 for horizontal transposon transfer. Molecular Genetics and Genomics 290: 67–78.

623 https://doi.org/10.1007/s00438-014-0900-y

624 Partha R., B. K. Chauhan, Z. Ferreira, J. D. Robinson, K. Lathrop, *et al.*, 2017 Subterranean

625 mammals show convergent regression in ocular genes and enhancers, along with adaptation to

626 tunneling, (D. T. Odom, Ed.). eLife 6: e25884. https://doi.org/10.7554/eLife.25884

627 Policarpo M., J. Fumey, P. Lafargeas, D. Naquin, C. Thermes, *et al.*, 2021 Contrasting Gene

628 Decay in Subterranean Vertebrates: Insights from Cavefishes and Fossorial Mammals.

629 Molecular Biology and Evolution 38: 589–605. https://doi.org/10.1093/molbev/msaa249

630 Ranallo-Benavidez T. R., K. S. Jaron, and M. C. Schatz, 2020 GenomeScope 2.0 and

631 Smudgeplot for reference-free profiling of polyploid genomes. Nat Commun 11: 1432.

632 https://doi.org/10.1038/s41467-020-14998-3

633 Robertson H. M., and K. W. Wanner, 2006 The chemoreceptor superfamily in the honey bee,

634 Apis mellifera: Expansion of the odorant, but not gustatory, receptor family. Genome Res. 16:

635 1395–1403. https://doi.org/10.1101/gr.5057506

636 Roff D. A., 1990 The Evolution of Flightlessness in Insects. Ecological Monographs 60: 389–

421. https://doi.org/10.2307/1943013

Rogers H. H., and S. Griffiths-Jones, 2012 Mitochondrial Pseudogenes in the Nuclear Genomes

of Drosophila. PLOS ONE 7: e32593. https://doi.org/10.1371/journal.pone.0032593

Romiguier J., J. Rolland, C. Morandin, and L. Keller, 2018 Phylogenomics of palearctic

Formica species suggests a single origin of temporary parasitism and gives insights to the

evolutionary pathway toward slave-making behaviour. BMC Evolutionary Biology 18: 40.

https://doi.org/10.1186/s12862-018-1159-4

Ronquist F., 1994 Evolution of Parasitism Among Closely Related Species: Phylogenetic

Relationships and the Origin of Inquilinism in Gall Wasps (hymenoptera, Cynipidae).

Evolution 48: 241–266. https://doi.org/10.1111/j.1558-5646.1994.tb01310.x

Sanver D., and B. A. Hawkins, 2000 Galls as habitats: the inquiline communities of insect galls.

Basic and Applied Ecology 1: 3–11. https://doi.org/10.1078/1439-1791-00001

Savolainen R., and K. Vepsäläinen, 2003 Sympatric speciation through intraspecific social

parasitism. Proceedings of the National Academy of Sciences 100: 7169–7174.

https://doi.org/10.1073/pnas.1036825100

Scanlan J. L., R. S. Gledhill-Smith, P. Battlay, and C. Robin, 2020 Genomic and transcriptomic

analyses in Drosophila suggest that the ecdysteroid kinase-like (EcKL) gene family encodes

the 'detoxification-by-phosphorylation' enzymes of insects. Insect Biochemistry and Molecular

Biology 123: 103429. https://doi.org/10.1016/j.ibmb.2020.103429

Schrader L., H. Pan, M. Bollazzi, M. Schiøtt, F. J. Larabee, *et al.*, 2021 Relaxed selection

underlies genome erosion in socially parasitic ant species. Nat Commun 12: 2918.

https://doi.org/10.1038/s41467-021-23178-w

Seeley T. D., 1974 Atmospheric carbon dioxide regulation in honey-bee (Apis mellifera)

colonies. Journal of Insect Physiology 20: 2301–2305. https://doi.org/10.1016/0022-

1910(74)90052-3

Senthilan P. R., and C. Helfrich-Förster, 2016 Rhodopsin 7–The unusual Rhodopsin in

Drosophila. PeerJ 4: e2427. https://doi.org/10.7717/peerj.2427

Sherman B. T., M. Hao, J. Qiu, X. Jiao, M. W. Baseler, *et al.*, 2022 DAVID: a web server for

functional enrichment analysis and functional annotation of gene lists (2021 update). Nucleic

Acids Research 50: W216–W221. https://doi.org/10.1093/nar/gkac194

Skaife S. H., 1922 On *Braula Coeca* , Nitzsch, a Dipterous parasite of the honey bee.

Transactions of the Royal Society of South Africa 10: 41–48.

https://doi.org/10.1080/00359192209519263

Stange G., and M. Diesendorf, 1973 The response of the honeybee antennal CO2-receptors to

N2O and Xe. J. Comp. Physiol. 86: 139–158. https://doi.org/10.1007/BF00702534

Starowicz M., P. Hanus, G. Lamparski, and T. Sawicki, 2021 Characterizing the Volatile and

Sensory Profiles, and Sugar Content of Beeswax, Beebread, Bee Pollen, and Honey. Molecules

26: 3410. https://doi.org/10.3390/molecules26113410

Suvorov A., B. Y. Kim, J. Wang, E. E. Armstrong, D. Peede, *et al.*, 2021 Widespread

introgression across a phylogeny of 155 Drosophila genomes. Current Biology 0.

https://doi.org/10.1016/j.cub.2021.10.052

Torto B., D. G. Boucias, R. T. Arbogast, J. H. Tumlinson, and P. E. A. Teal, 2007 Multitrophic

interaction facilitates parasite–host relationship between an invasive beetle and the honey bee.

Proceedings of the National Academy of Sciences 104: 8374–8378.

https://doi.org/10.1073/pnas.0702813104

Venner S., V. Miele, C. Terzian, C. Biémont, V. Daubin, *et al.*, 2017 Ecological networks to

unravel the routes to horizontal transposon transfers. PLOS Biology 15: e2001536.

https://doi.org/10.1371/journal.pbio.2001536

Weinstock G. M., G. E. Robinson, R. A. Gibbs, G. M. Weinstock, G. M. Weinstock, *et al.*,

2006 Insights into social insects from the genome of the honeybee Apis mellifera. Nature 443:

931–949. https://doi.org/10.1038/nature05260

Weiss L. A., A. Dahanukar, J. Y. Kwon, D. Banerjee, and J. R. Carlson, 2011 The Molecular

and Cellular Basis of Bitter Taste in Drosophila. Neuron 69: 258–272.

https://doi.org/10.1016/j.neuron.2011.01.001

Winkler I. S., A. H. Kirk-Spriggs, K. M. Bayless, J. Soghigian, R. Meier, *et al.*, 2022

Phylogenetic resolution of the fly superfamily Ephydroidea–Molecular systematics of the

enigmatic and diverse relatives of Drosophilidae. PLOS ONE 17: e0274292.

https://doi.org/10.1371/journal.pone.0274292

Winston M. L., 1987 *The Biology of the Honey Bee*. Harvard University Press.

Yassin A., 2013 Phylogenetic classification of the Drosophilidae Rondani (Diptera): the role of

morphology in the postgenomic era. Systematic Entomology 38: 349–364.

Zanini D., D. Giraldo, B. Warren, R. Katana, M. Andrés, *et al.*, 2018 Proprioceptive Opsin

Functions in Drosophila Larval Locomotion. Neuron 98: 67-74.e4.

https://doi.org/10.1016/j.neuron.2018.02.028

Zhang H.-H., J. Peccoud, M.-R.-X. Xu, X.-G. Zhang, and C. Gilbert, 2020 Horizontal transfer

and evolution of transposable elements in vertebrates. Nat Commun 11: 1362.

https://doi.org/10.1038/s41467-020-15149-4

Zimin A. V., D. Puiu, M.-C. Luo, T. Zhu, S. Koren, *et al.*, 2017 Hybrid assembly of the large

and highly repetitive genome of Aegilops tauschii, a progenitor of bread wheat, with the

MaSuRCA   mega-reads   algorithm.   Genome   Res.   27:   787–792.

https://doi.org/10.1101/gr.213405.116

Legends of figures

Figure 1 – The bee louse fly (*Braula coeca*) morphology and genome. A) Two adults attached to a honey bee worker preserved in alcohol. B) Dorsal view of an adult showing the loss of the wings, halters and scutum, the reduction of the mesonotum and the robustness of the legs. C) Frontal view of an adult showing the reduction of the eyes and the loss of the ocelli. Scale bars = 1 mm in A and 0.5 mm in B and C. D) Orthologous genes colocalization in *D. melanogaster* Muller's elements and *B. coeca* contigs demonstrating the predominant conservation of synteny. E) The mitochondrial genome of *B. coeca*. Blue and violet ribbons indicate the protein-encoding genes in sens and anti-sens respectively, red and yellow ribbons indicate the rRNA and tRNA genes in sens and anti-sens respectively. Names with asterix indicate the presence of inactivated paraloguous genes.

Figure 2 – Maximum-likelihood phylogenetic tree for A) odorant receptors (ORs) and B) gustatory receptors (GRs) of *A. mellifera* (green), *B. coeca* (red) and *D. melanogaster* (blue). Genes commented in the text are labelled. For full labels see Supplementary Figures 4 and 5. Sequences for *B. coeca* are given in Supplementary Datasets 2 and 3 for ORs and GRs, respectively. Sequences for *A. mellifera* and *D. melanogaster* are from Robertson and Wanner (2006). a.a. = amino acids.

0.12) was calculated over the transposase open reading from of one copy of the element

extracted from the *A. mellifera* genome and another copy extracted from the *B. coeca*

genome.

Supplementary Figure 3 – Phylogeny of *Famar1*-like copies from 38 animal species. The

ten *Famar1*-like copies showing the highest nucleotide identity to the *Famar* element

initially described in the earwig (Barry *et al.* 2004b) were retrieved using online blastn

(see Supplementary Text 1) and extracted from the genome of 37 animal species. Filled

circles indicate bootstrap value higher than 70%, with the diameter of the circle

proportional to each individual value. Name of the copies are composed of the name of

the species from which they were extracted and a unique number that allows making a

correspondence with sequences provided in Supplementary Dataset 1.

Supplementary Figure 4 – Labelled maximum-likelihood phylogeny of odorant

receptors (ORs) of *A. mellifera* (green), *B. coeca* (blue), and *D. melanogaster* (red) show in

Figure 2A.

Supplementary Figure 5 – Labelled maximum-likelihood phylogeny of gustatory

receptors (GRs) of *A. mellifera* (green), *B. coeca* (blue), and *D. melanogaster* (red) show in

Figure 2B.

Supplementary Table 1 – Proportions of transposable elements in the genomes of *B.*

coeca, *D. melanogaster* and *A. mellifera*.

Supplementary Table 2 – Gene Ontology (GO) enrichment for *D. melanogaster* genes that

are absent from *B. coeca* genome as inferred using DAVID 2021.

Supplementary Table 3 – Gene Ontology (GO) enrichment for *D. melanogaster* genes that

have additional copies in *B. coeca* genome as inferred using DAVID 2021.

Supplementary Text 1 – Analysis of horizontal transfer of transposable elements

between *B. coeca* and *A. mellifera*.

Figure 1

D

B. coeca contigs

Muller's element
(*D. melanogaster*)

F B C D E A

Figure 2

A

DmOr67b

DmOrco

AmOr11

DmOr56a

DmOr49b

DmOr98b

B

DmGr61a, DmGr64a-e (sweet)

DmGr21a, DmGr63a
(CO₂)

DmGr43a (sweet)

DmGr66a (bitter)

AmGr10 (a.a.)

DmGr33a (bitter)

DmGr57a (bitter)

DmGr93a, DmGr94a (bitter)