Single-molecule long-read methylation profiling reveals regional DNA methylation

regulated by Elongator Complex Subunit 2 in Arabidopsis roots experiencing

spaceflight

Mingqi Zhou[1], Alberto Riva[2], Marie-Pierre L. Gauthier[3], Michael P. Kladde[3], Robert J.

Ferl[1,4,*], Anna-Lisa Paul[1,2,*]

[1]Department of Horticultural Sciences, University of Florida, 2550 Hull Road, Fifield Hall,

Gainesville FL, 32611, USA

[2]Interdisciplinary Center for Biotechnology Research, University of Florida, 2033 Mowry

Road, Gainesville FL, 32610, USA

[3]Department of Biochemistry and Molecular Biology, University of Florida, 2033 Mowry

Rd, Gainesville, FL 32608, USA

[4]Office of Research, University of Florida, 1523 Union Rd, Grinter Hall, Gainesville, FL,

32611, USA

*Corresponding authors: robferl@ufl.edu (RJF), alp@ufl.edu (ALP)

Abstract

The Advanced Plant Experiment-04 - Epigenetic Expression (APEX04-EpEx)

experiment onboard the International Space Station examined the spaceflight-altered

cytosine methylation in two genetic lines of *Arabidopsis thaliana*, wild-type Col-0 and the

mutant *elp2-5,* which is deficient in an epigenetic regulator Elongator Complex Subunit

2 (ELP2). Whole-genome bisulfite sequencing (WGBS) revealed distinct spaceflight

associated methylation differences, presenting the need to examine specific space-

altered methylation at single molecule resolution to associate specific changes over

large regions of spaceflight related genes. Therefore flap-enabled next-generation

capture (FENGC), a novel targeted multiplexed DNA capture and enrichment technique

allowing cleavage at any specified sites, was applied to survey spaceflight-altered DNA

methylation at the single-molecule level. The FENGC capture panel contained 108

targets ranging from 509 to 704 nt within the promoter or gene body regions of gene

targets derived from spaceflight whole-genome data sets. In addition to genes with

significant changes in expression and average methylation levels between spaceflight

and ground control, targets without differentially methylated WGBS but with space-

altered distributions of the proportion of methylated cytosines per molecule were further

identified. Moreover, trends of co-methylation of different cytosine contexts were

exhibited in the same DNA molecules. We further identified significant DNA methylation

changes in three previously biological process-unknown genes, and two of them

showed enhanced root growth rate in loss-of-function mutants (named as *ELP2-*

regulated Methylation in Orbit 1 and *2*). FENGC makes multiplexed targeted single-

molecule profiling of methylation simple and cost-effective, which provides an additional

power to population-based data for methylation analysis and identification of

epigenetically modified substrates in plants. This case study has revealed regional

spaceflight-altered methylation in selected targets, which were not identified by short-

read methods in spaceflight plants. The newly identified *EMO1* and *EMO2* are root

growth regulators, which can be epigenetically involved in plant adaptation to

spaceflight.

Introduction

DNA methylation epigenetically modulates gene expression and impacts chromatin

structure to function in physiological adaptation to environmental changes in eukaryotes

(Zhang *et al.*, 2018). Plants will play an important role in the human life support systems

envisioned for spaceflight and planetary exploration habitats (Wheeler, 2010; Zabel *et*

al., 2016). Spaceflight presents a unique environmental challenge for plants, an

environment that is outside the evolutionary experience of any terrestrial organism.

Epigenetic change plays an important role in environmental adaptation, and it is crucial

to understand the role of epigenetics in plant adaptation to this novel environment (Paul

et al., 2021; Zhou *et al.*, 2019). The Advanced Plant Experiment-04 - Epigenetic

Expression (APEX-04) experiment onboard the International Space Station (ISS)

examined the spaceflight-altered cytosine methylation in *Arabidopsis thaliana* (Arabidopsis) using *elp2-5*, a mutant of an epigenetic regulator, Elongator Complex Subunit 2 (ELP2), along with Col-0, a wild-type control (Paul *et al.*, 2021). RNA sequencing (RNAseq) and Whole genome bisulfite sequencing (WGBS) revealed remodeled transcriptomes and DNA methylomes during spaceflight, remodeling that was regulated by ELP2. The spaceflight *elp2-5* phenotype was distinct from the space-grown wild-type plants in that root growth did not present a typical spaceflight directionality, which orients root growth away from the germinating seed, but rather displayed a distinct clumping of roots clustered around the seed. The molecular phenotype exhibited distinct differences in the transcriptome and methylome compare to the ground controls, with a majority of differentially expressed genes in *elp2-5* roots being also differentially methylated between spaceflight samples and ground controls (Paul *et al.*, 2021).

However, the short-read sequencing used in WGBS precludes the ability to phase regional epigenetic signatures along individual molecules such as consecutive methylated cytosines relative to transcription factor binding, nucleosome positioning, histone incorporation and modification (Edwards *et al.*, 2017; Zhang *et al.*, 2018). Moreover, the WGBS detection of differentially methylated cytosines (DmCs) can be affected by sequencing depth and false positives due to overestimation of 5mC level (Feng *et al.*, 2020; NEB), and further downstream analyses including differentially methylated region analysis rely on DmCs (Chen *et al.*, 2016; Kishore and Pelizzola, 2018).

Flap-enabled next-generation capture (FENGC) is a novel, strand-specific DNA capture method that allows DNA methylation analysis with long reads of user-defined target regions (Zhou *et al.*, 2022). FENGC enables sensitive characterization of contiguous single molecules over larger regions, thereby providing a deeper insight into the methylation state than that afforded by the averaged state of methylation obtained by multiple short reads. In addition, compared with WGBS, FENGC increases coverage depth in regions of interest, which improves data quality and minimizes cost of reads in unwanted areas. Furthermore, FENGC has advantages of liberty in target region

selection and cost-effectiveness that are superior to other targeted-sequencing methods.

First, the formation of flap structure on DNA strands of target regions using user-

designed oligos and cleavage by flap endonuclease avoid the limits due to sequence

dependence of enzymes for target selection. For DNA methylation survey, the target

capture of FENGC occurs prior to deamination of 5mC, also avoiding the primer design

limits of regular methyl-PCR due to potentially methylated sites. Second, the utilization

of standard oligos for targeting without modified bases as well as the single-tube

experimental protocol significantly lower the cost (Kladde *et al.*, 2022; Zhou *et al.*, 2022).

Hence, this tool is ideal for targeting customer-designed specific panels especially for

DNA methylation analysis of plants possessing 5-methylation in three types of cytosine

context, CG/CHG/CHH (Grimanelli and Ingouff, 2020). The accuracy and efficiency of

enzymatic methyl-seq, which is part of the FENGC procedure, has been verified in

plants (Feng *et al.*, 2020); however, FENGC has yet to be applied to methylation

analysis in plants.

Therefore, we set out to demonstrate the effective application of FENGC in plant

methylation contexts. The observed phenotypical and molecular alteration in roots of

Col-0 and *elp2-5* grown on the ISS in APEX-04 present a short-read methylation

dataset for a comparison with methylation profiles assayed by FENGC single-molecule

long reads. WGBS-unidentified differential modification of DNA methylation substrates

in functionally-annotated genes as well as unknown genes of plants grown in orbit have

been detected.

Results and discussion

Quality of FENGC assay

In the FENGC assay, strand cleavage was directed by precise targeting of FEN1 on flap

formation, which was user-defined through oligo design so that sequence-based limits,

such as reliance on restriction sites or specific motif sequences, could be avoided. The

target panel for capture and assay contained 108 genic regions ranging from 509 to 704

nt (Table S1), with mean of 580 nt and median of 572 nt, within the promoter or gene

body regions of 87 epigenetically regulated genes derived from our APEX-04

spaceflight and related experiments (Paul *et al.*, 2021, 2013; Paul, Zhou, *et al.*, 2017;

Zhou *et al.*, 2019). This design therefore provides strand specific methylation data over

the entire range of the targeted sequences.

The capture was performed as a single-tube multiplex reaction followed by enzymatic

methyl-PCR (oligos are listed in Table S2). The FENGC products were subjected to

PacBio long-read HiFi sequencing. For APEX04 samples, including Ground control Col-

0 Roots (GCR), Flight Col-0 Roots (FCR), Ground control elp2-5 Roots (GER) and

Flight elp2-5 Roots (FER), capture products consistently yielded around 90% reads on-

target ratio and, on average, 82% targets were detected by at least 10 reads after

filtering and combining the four biological replicates from each sample (Figure 1A and

1B; Table S3). The GC content of the 108 targets, ranging from 19% to 52%, was

positively correlated with the aligned reads number (Figure 1C; Figure S1), whereas

there was a modest negative correlation in FENGC targets with an average GC content

above 50% (Zhou *et al.*, 2022). This indicates that both high and low GC content of

targets influenced the reads yield of FENGC products.

Targeted DNA capture and sequencing analyses have been widely deployed in plant

genomic studies (Ostezan *et al.*, 2021). For invitro targeted assay in plant genomes,

FENGC possesses multiple advantages compared with other methods, including high

liberty for customer design, high performance in sequencing depth, target number, on-

target ratio, and low cost (Table 1). Especially for targeted DNA methylation detection in

plants, to date, only singleplexed or multiplexed bisulfite-PCR has been regularly used,

which suffers from the limits in sequencing read length, primer region selection, and

assay capability. To our knowledge, FENGC provides the unique opportunity for plant

research community for multiplexed, targeted sequencing analysis on DNA methylation.

Regional methylation detection by FENGC in comparison with WGBS data

One major advantage of single-molecule profiling of DNA methylation is that the

contiguous long reads allow assessment of every cytosine within same DNA molecule,

thereby presenting a coherent regional methylation status of that entire gene element

within a cell. In addition, the average methylation levels in cytosines across all

molecules can be derived from the data for comparison to WGBS data. For the present

study, long-read, single-molecule sequencing identified gene targets showing both

significant differential expression and average methylation level changes between

spaceflight and ground control. Moreover, space-altered distributions of the proportion

of methylated cytosines per molecule were further revealed.

Combining the four biological replicates, GCR, FCR, GER and FER yielded 79, 93, 88

and 96 target regions with coverage of at least 10 reads (Table S3; Table S4). The

distributions of the proportion of methylated cytosines per molecule in each target

region were calculated. Among them, 12 targets in 12 genes showed differential

methylation proportion per molecule (DmPPM) for the comparison between spaceflight

and ground control in Col-0 or *elp2-5* (Table S1). Meanwhile, the APEX04 WGBS-

identified DmCs (methylation difference > 0.2, FDR < 0.01) between spaceflight and

ground control were examined within 108 target regions of FENGC. Col-0 and *elp2-5*

showed 8/2/7 and 5/2/2 targets possessing DmPPM for CG/CHG/CHH methylation,

respectively. Among these regions, only one target with DmCHH for Col-0 and two

targets with DmCHG for *elp2-5* were identified by WGBS (Table S1). The comparison of

detection in DmPPM and DmC demonstrated that the changes of contiguous

methylation status observable in a specific subpopulation of molecules in FENGC may

be indiscernible in a population-based analysis.

Next, we present four examples associated with defense responses to stimuli and three

biological process-unknown genes regulated by ELP2 during plant growth in spaceflight

to exhibit details of space-altered methylation status that are newly uncovered by

FENGC (Table S1). Considering these seven targets, the methylation levels of

cytosines exhibited correlation coefficients above 0.93 between any two biological

replicates and above 0.95 between FENGC and WGBS results (Figure 1D). In each

region, 93%-100% of the mapped reads are full-length, containing all CG/CHG/CHH

sites of each target region (Figure S2). These results verify the high data quality for

captured targets in the FENGC assay of space plants.

Identification of previously unidentified space-altered DNA methylation in defense-related gene targets

Among 12 genes exhibiting DmPPM of spaceflight vs ground control, four defense-related genes showed space-triggered DNA methylation modification that was not discernible in WGBS. The first three examples were examined with CG methylation, for which DmPPM was shown in CG sites but no DmCG (methylation difference > 0.2, FDR < 0.01) between spaceflight and ground control was detected by WGBS in these target regions. *CML46* (AT5G39670), a gene encoding a calmodulin-like protein showing significant transcriptional down-regulation during spaceflight in Col-0 but not *elp2-5* plants, was assayed in a promoter region containing a MYB-like DNA binding element (Figure 2A and 2B). Flight and ground control Col-0 roots (FCR and GCR), but not roots from *elp2-5* (FER and GER), showed a significant difference in the CG methylation level in an area flanking the MYB-like element, which contained five CG sites (Figure 2C, in dashed frame). Consistent with these data, CG methylation proportion per molecule (mPPM) in this target region was significantly higher in FCR than GCR (Figure 2D). A significant difference between GCR and GER was also observed. FENGC showed more molecules containing consecutively methylated CG sites in FCR (12%) compared to GCR (6.9%) in the area flanking this *cis*-regulatory element (Figure 2E). Consistency between the levels of gene transcription, WGBS-derived population-based methylation status, and FENGC-derived mPPM, support the biological conclusion that ELP2 played a negative role in expression and a positive role in promoter methylation of *CML46* in spaceflight.

ABCB17 (AT3G28380), an ATP-binding cassette transporter gene that also showed a significantly lower transcription level in FCR than GCR, was assayed in a gene body region (Figure S3A and S3B). The overall patterns of averaged regional CG methylation levels were very similar between the four samples, which did not show significant changes between any pair of samples (Figure S3C); however, the CG mPPM in this target region was significantly higher in FCR than in GCR (Figure S3D). The FENGC single-molecule data showed that 3.7% of the reads in FCR possessed at least 6

consecutive methylated CG sites, whereas in GCR the percentage was 0.8% (Figure
S3E). Biologically, *ABCB17* was repressed by spaceflight and ELP2 contributed to its
transcription and gene body methylation.

Similarly, in the promoter region of *DEFL* (AT4G22217), a defensin-like gene (Figure
S4A and S4B), the patterns of averaged CG methylation levels were highly similar and
no significant alterations were detected between any two samples (Figure S4C);
however, CG mPPM was significantly higher in FER than GER (Figure S4D and S4E).
This higher methylation proportion correlates with the lower transcription level of this
gene in FER compared to GER (Figure S4B). ELP2 enhanced expression and
repressed promoter methylation of *DEFL* in spaceflight. These examples suggest that
long-read FENGC single-molecule level methylation profiling can be more revealing of
consequential methylation changes compared with population-based methylation
methods such as WGBS, especially when alterations potentially occur in a
subpopulation of cells within a sample.

For the fourth example, a promoter fragment containing a MeJA-responsive element
was selected in *PDF1.1* (AT1G75830), a gene coding a pathogenesis-related protein
belonging to the plant defensin family and showing a significant spaceflight induction in
elp2-5 (Figure 3A and 3B). The average methylation level per site over all sequenced
molecules showed significant differences at CHG and CHH sites between GER and
GCR, as well as in CHH sites between FCR and GCR (Figure 3C). For mPPM,
significant differences were detected in all pairwise comparisons between flight
conditions and genotypes at cytosines in all three methylation contexts (Figure 3D).
DmCs of WGBS between GER and GCR correlated with methylation changes detected
by FENGC (Figure 3D; Table S1), while between spaceflight and ground control only
three DmCGs in *elp2-5* were identified by WGBS, and the FENGC determined
methylation level changes were smaller than 0.2 at these three sites (Figure 3C), which
was in line with the WGBS overestimation of methylation level compared with EM-seq,
especially in CHG sites (Feng *et al.*, 2020). The patterns of methylation proportion in
DNA molecules demonstrated that three cytosine contexts showed a trend of co-
methylation in this region among cells in the population (Figure S5). Within each

molecule, methylation proportion of CG is higher than CHG in 97.9% reads, and that of

CHG is higher than CHH in 87.5% reads (Figure 4A). Although three types of cytosine

methylation exhibited a trend of coordination in the whole cell population, they could

spread independently within subpopulations of cells. The CG methylation proportions

were above 0.57 in all reads of GCR and above 0.28 in GER, while FCR and FER had

reads with CG methylation proportion as low as 0. These data indicate that more

molecules were hypomethylated in this region in *elp2-5* compared with Col-0 and in

spaceflight compared with ground control.

The profiling of methylation data in this example also showcases another advantage of

long-read, single-molecule FENGC assay of DNA methylation in plants, which is that all

three methylation sequence contexts (CG/CHG/CHH) can be identified and plotted for

the same molecule, thereby examining the entire methylation status of a gene region.

To determine the spatial distribution of these methylation differences, three separate but

correlated panels were plotted for CG, CHG and CHH methylation in the same DNA

molecule per row (Figure 3E). In these plots, black areas are unmethylated regions, red

areas are methylated regions, and gray areas are transition zones between methylated

and unmethylated sites. It is visually apparent that spaceflight results in more

unmethylated regions in Col-0, with the deepest loss of methylation concentrated near

the MeJA-responsive element. It is also visually apparent that *elp2-5* shows a deeper

loss of methylation in that region. For the CG methylation context, a footprint showing

consecutive non-methylation of the first 2 CG sites in flanking area of the MeJA-

responsive element was identified in 8.4%, 18.0%, 39.4% and 44.0% reads in GCR,

FCR, GER and FER, respectively, quantifying the loss of CG methylation in this area.

Interestingly, 60%, 86.5%, 97.5% and 97.6% reads with consecutive non-methylation in

the second and the third CG sites in GCR, FCR, GER and FER, respectively, were also

included in this footprint and showed consecutive non-methylation of the first 3 CG sites.

In addition, CHG and CHH methylation proportions were significantly lower in the

footprint area in the reads containing this footprint (Figure 4B). These results are

consistent with the previously observed high frequency of co-methylation/demethylation

in nearby cytosines in Arabidopsis (Tran *et al.*, 2018), and demonstrate that FENGC

provides further opportunities for studying all three types of co-methylation methylation

along each single molecule .

Four targets shown here are associated with plant defense signaling, which is largely

utilized in spaceflight adaptation of plants and the related genes can be involved in

regulation of root growth phenotypes during spaceflight (Califar *et al.*, 2020; Paul, Sng,

et al., 2017). It has also been reported that ELP2 modulates the epigenetic control of

plant immune responses and defense processes (Wang *et al.*, 2013). The newly

identified space-altered methylation in these targets is involved in ELP2-regulated

defense/immunity-related pathways that may function in adaptation of plant roots grown

in orbit.

Identification of novel spaceflight-related epigenetic modification in biological
process-unknown genes

The 12 genes with DmPPM between spaceflight and ground control were annotated

using the Database for Annotation, Visualization and Integrated Discovery (DAVID,

https://david.ncifcrf.gov/). Three genes were returned without gene ontology (GO) or

functional annotation for biological process. Further investigation verified that the three

target regions within these three genes possessed significant alteration between

spaceflight and ground control in both average cytosine methylation level as well as

mPPM in Col-0 or *elp2-5*. Moreover, loss-of-function mutants of two of these genes

showed a root growth phenotype. We dubbed the two genes that could be

epigenetically involved in plant adaptation to spaceflight under regulation of ELP2,

AT1G27565 and AT5G52710, as *ELP2-regulated Methylation in Orbit* (*EMO*) *1* and *2*,

respectively.

A promoter region was captured by FENGC for *EMO1* (AT1G27565), containing an

anaerobic-responsive element (ARE) beside the transcription start site (TSS) (Figure

5A). The presence of the ARE is consistent with the observed induction of hypoxia-

responsive genes in spaceflight transcriptome data sets (Choi *et al.*, 2019; Zhou *et al.*,

2019). *EMO1* was significantly up-regulated by spaceflight in *elp2-5*, but not in Col-0

(Figure 5B). Regional average methylation level analysis detected significant

differences between genotypes and between spaceflight and ground control in Col-0

only for CG sites (Figure 5C). The calculation of mPPM showed significant alteration

between spaceflight and ground control in all three types of cytosine context for Col-0

and in CG sites for *elp2-5* (Figure 5D). Visualization of three types of methylated

cytosines in each single molecule revealed footprints of methylation-free areas in areas

in subpopulations of epialleles (Figure 5E). Significantly, more molecules in *elp2-5*

lacked cytosine methylation within this region than Col-0, and more molecules with

methylation-free areas beside TSS were identified in spaceflight samples than ground

control. These results were consistent with the observation of lower correlation of

mPPM between GG and CHH in GCR than other samples (Figure S6 and S7). After

demethylation initiated by spaceflight or mutation of ELP2 in a small proportion of cells,

the co-methylation for different types of cytosine context in those molecules had been

changed accordingly. Overall, ELP2 played a negative role in *EMO1* expression and

maintained cytosine methylation of the *EMO1* promoter in a subpopulation of cells in

roots. Spaceflight promoted demethylation in flanking areas of *EMO1* TSS, while the

induction of *EMO1* transcription in space was repressed by ELP2.

The *EMO2* (AT5G52710) gene belongs to copper transport protein family (de Abreu-

Neto *et al.*, 2013). The captured promoter region contains two MeJA-responsive

elements (Figure 6A). The transcription level of *EMO2* was lower in *elp2-5* than Col-0,

and lower in spaceflight than ground control (Figure 6B). Accordingly, the regional

average methylation levels were significantly higher in *elp2-5* than Col-0 for CG and

CHG sites, which was consistent with DmC analysis of WGBS (Figure 6C). In contrast,

CHH differential methylation levels detected by FENGC were smaller than that of

WGBS. The mPPMs were higher in spaceflight than ground control in all three cytosine

types for *elp2-5*, which explained the space-repressed expression of *EMO2* when ELP2

was deficient (Figure 6B and 6D). Intriguingly, the areas of two MeJA-responsive

elements both harbored a footprint of low CG methylation, whereas CHG sites localized

within these elements showed methylation peaks, especially in *elp2-5* (Figure 6C and

6E). When ELP2 was inactivated, the two CG methylation footprints were largely lost.

Moreover, the correlations of mPPM between different types of cytosine context were

lower in *elp2-5* than Col-0 (Figure S8 and S9). These results indicate the role of ELP2 in

the maintenance of hypomethylated motifs and coordination of methylation for different

cytosine types in this chromatin region, which could contribute to activation of *EMO2*

transcription.

The AT4G04990 gene encodes a serine/arginine repetitive matrix-like protein and the

transcripts can be transported between roots and shoots (Thieme *et al.*, 2015). The

captured promoter region contains a TC-rich repeat that is related to defense and stress

responses (Figure S10A). The transcription level of AT4G04990 was down-regulated by

both spaceflight and mutation of ELP2 (Figure S10B). No DmC in WGBS or significant

difference of average methylation level in FENGC was identified for this region, while

DmPPMs of CG and CHH were detected between spaceflight and ground control in Col-

0 (Figure S10C and S10D). In this region, a stronger correlation was observed between

mPPMs of CHG and CHH than that of CG and CHH (Figure S10E; Figure S10 and S11).

Correlation between mPPMs of CG and CHG increased by spaceflight in Col-0 (Figure

S11), which was partially due to space-induced demethylation of CG sites in a

subpopulation of cells (Figure S10D and S10E; Figure S12). The coordination of CG

and non-CG methylation might be related to the transcription repression of AT4G04990

in spaceflight and *elp2-5*.

The features in alteration of DNA methylation at the single molecule level, such as

DmPPM and methylation footprints, would be more informative for detection of

epigenetic changes and better connect the regulation of gene expression to epigenetic

modification, compared with population-based DmC levels that would lose epigenetic

information in the single molecules. Therefore, FENGC data provide additional clues for

further selection of target genes of interest for downstream analysis, especially when

candidates include unknown genes.

EMO1 and EMO2 are negative regulators of root growth

T-DNA insertion lines of the three biological process-unknown genes identified by

FENGC were characterized. Two lines of each gene tested showed disrupted gene

expression levels compared with Col-0 (Figure S13). Thus, SALK_092504 and SAIL_344_F07 were named as *emo1-1* and *emo1-2*, while SALK_140748C and SALK_091636C were named as *emo2-1* and *emo2-2*, respectively. In a constant LED light condition that was similar with Veggie facility onboard ISS, *emo1* and *emo2* mutants showed significantly enhanced root growth rates compared with Col-0 (Figure 7A-C). The primary root length of *emo1-1*, *emo1-2*, *emo2-1*, and *emo2-2* were significantly longer than that of Col-0 in 4-day and 7-day old seedlings. The two mutants of AT4G04990 exhibited similar growth phenotype with wild type. Meanwhile, treatments using 2-D clinorotation at 4 rpm or reorientation to 90° did not show altered gravitropic responses in roots of any line tested, although roots of *EMO1* and *EMO2* mutants still grew faster than Col-0 on the clinostat (Figure S14 and S15). These results demonstrated that *EMO1* and *EMO2* are two negative regulators of root growth, but not regulators of root gravitropism.

Remodeled plant root growth has been reported in orbital spaceflight (Paul *et al.*, 2012). Both direction of root tips during growth and the growth rate of roots were impacted in Arabidopsis plants grown on the ISS (Kwon *et al.*, 2015; Paul *et al.*, 2012; Paul, Sng, *et al.*, 2017). The primary root length of 8.5-day old seedlings for both Col-0 and Ws were significantly shorter in flight samples than ground control. This influence on root growth from spaceflight involves epigenetic and transcriptional modulation (Paul *et al.*, 2021, 2013). As a epigenetic modulator of root growth and development, ELP2 modifies DNA methylation or histone acetylation of a range of transcription factors and other developmental regulators in roots (Jia *et al.*, 2015). On the ISS, the *elp2-5* mutant exhibited severe growth retardation in roots compared with Col-0 (Paul *et al.*, 2021). Here, *EMO1* and *EMO2* have been identified as two downstream targets of DNA methylation modification led by ELP2 and/or caused by extraterritorial environment. Together with the root growth rate phenotype in their mutants, *EMO1* and *EMO2* can play a role in root growth regulation during space adaptation of plants, which may involve the epigenetic signaling pathways controlled by ELP2. Moreover, although the mutants of AT4G04990 did not show root growth alteration on the ground, further investigation will be needed to examine its function in space.

Conclusion

Whereas short-read WGBS only detects alteration of average methylation level in
cytosines, targeted long-read, single-molecule profiling reveals the detailed epigenetic
modifications occurring in cell subpopulations and identifies epialleles present at low
frequencies. On one hand, FENGC and WGBS show consistency in the detection of
cytosine methylation levels. On the other hand, FENGC enables single-molecule
determination of methylation alteration, which greatly extends interpretation beyond that
of WGBS. Analysis of DmPPM serves as critical addition to DmC data in methylation
survey. The capacity of target cleavage at any specified sites enables wide application
of FENGC, such as studies on plant development, environmental responses, and
generation of biomarkers. In the current work, FENGC allowed us to survey genic
regions of interest from primary transcriptomic and WGBS screening, successfully
identifying two novel root growth regulators through analysis of DNA methylation
alteration. In summary, FENGC provides the plant and space biology communities with
a powerful tool for the practical application of multiplexed targeted sequencing and
coordinated methylation analysis at single-molecule level as routine experiments. This
FENGC screen of spaceflight-associated genes revealed candidate genes with
regulatory regions that harbored significant alterations in consecutive spans of
methylation. Further genetic analysis identified *EMO1* and *EMO2*, which are novel
candidates for further investigation of root growth regulatory signaling elicited by ELP2
in plant adaptation to spaceflight.

Materials and methods

Plant materials and operations of spaceflight and ground controls

The *Arabidopsis thaliana* lines Col-0 and *elp2-5*, as well as spaceflight experimental
operations, have been previously documented (Paul *et al.*, 2021; Wang *et al.*, 2013).
The Advanced Plant Experiment 04 – Epigenetic Expression (APEX-04-EPEX)
spaceflight experiment was launched onto International Space Station (ISS) through the

SpaceX mission CRS-10. In brief, seeds were sterilized and sown on growth medium

containing 0.5% phytagel, 0.5× Murashige–Skoog salts, 0.5% (w/v) sucrose, and 1 ×

Gamborg's Vitamin Mixture, in Petri dishes (100 mm × 15 mm; Fisher Scientific,

Pittsburgh, USA), which were then sealed with breathable tape (3M Micropore,

Maplewood, USA). Plants were grown in the Vegetable Production System (Veggie)

facility on the ISS and the Veggie hardware in the ISS environmental simulator (ISSES)

chamber at Kennedy Space Center under constant LED light of 100–135 µmol/m^{2}s

Photosynthetically Active Radiation (PAR) until 11 days after germination, prior to

harvest using RNAlater (Ambion, Grand Island, USA) in Kennedy Space Center fixation

tubes (KFTs). Plants from each plate were collected into individual KFTs and stored

at -80°C until delivery to the lab for analysis.

Genomic DNA Isolation and flap-enabled next-generation capture (FENGC) assay

DNA extraction was conducted using the protocol designed for RNAlater preserved

plant samples (LeFrois *et al.*, 2016). The same set of DNA samples from Col-0 and

elp2-5 root tissues used for whole-genome bisulfite sequencing (WGBS) (Paul *et al.*,

2021) were subjected to flap-enabled next-generation capture (FENGC) assay. The

samples included four biological replicates of each condition: Ground control Col-0

Roots (GCR), Flight Col-0 Roots (FCR), Ground control *elp2-5* Roots (GER) and Flight

elp2-5 Roots (FER).

FENGC oligo-1, -2 and -3 were designed using FENGC Oligonucleotide Designer

(FOLD) program (https://github.com/uf-icbr-bioinformatics/FOLD), to bind target DNA to

form 5′ flap structures that were cleaved by flap endonuclease (FEN), which enabled

ligation between target fragments and universal adaptors (Zhou *et al.*, 2022). There

were 108 targets selected within promoter and gene body regions of 87 genes of

interest derived from previous spaceflight-associated transcriptome and DNA

methylome studies (Paul *et al.*, 2021, 2013; Paul, Zhou, *et al.*, 2017; Zhou *et al.*, 2019).

The detailed information of the 108 targets is in Table S1. All oligos used in this study

are listed in Table S2. All captured sequences are from the coding strand of each gene.

FENGC experiments were performed as described by Kladde and colleagues (Zhou *et*
al., 2022), with minor modifications. In brief, approximately 200 ng of Arabidopsis
genomic DNA was fragmented by sonication using a Bioruptor Pico (Diagenode,
Denville, USA) for 6 cycles of 5 sec on and 30 sec off in 20 µl ddH$_2$O. The concentration
of oligo-1, -2, and -3 was 18 nM each in the master tube as recommended (Zhou *et al.*,
2022). Flap cleavage was performed in the presence of FENGC universal oligo 1-T (U1-
T oligo), flap oligos-1 and -2 specific for 108 targets, and thermostable FEN1 (New
England Biolabs, Ipswich, USA). Ligation was initiated by adding a stock mixture of
oligos-3 for all 108 targets, U2 oligo, and Ampligase (Lucigen, Middleton, USA) to the
same tube. DNA not ligated to U2 oligo and therefore not protected by 3' C3 Spacer
was removed by addition of exonucleases I and III (NEB, Ipswich, USA). The enriched
products were purified and then subjected to enzymatic methylation conversion
according to the manual of NEBNext Enzymatic Methyl-seq Conversion Module (NEB,
Ipswich, USA). Converted DNA was amplified for 23 cycles using U1 and U2 primers
and KAPA HiFi HotStart Uracil+ DNA polymerase (Roche, Pleasanton, USA), according
to the manufacturer's instructions. FENGC amplification products were purified and
product quality and quantity were determined using Agilent TapeStation D1000 system
(Agilent, Santa Clara, USA) and Qubit 3 Fluorometer (Thermo Fisher, Waltham, USA),
respectively.

Pacific Biosciences Sequel IIe SMRT sequencing and data analysis

The library construction and sequencing of FENGC products on the Pacific Biosciences
(PacBio) Sequel IIe SMRT platform was performed by Interdisciplinary Center for
Biotechnology Research (ICBR) at the University of Florida. The HiFi reads were
generated using 30-h movie of circular consensus (CCS) mode with filtering for ≥3
passes per single polymerase read, with default settings for other parameters. HiFi
reads mapping and visualization of DNA methylation states of single molecules were
performed using MethylMapper as previously described (Nabilsi *et al.*, 2014), with
modifications to accommodate plant genomes. Reads with more than three consecutive
methylated CHH sites were removed (Liu *et al.*, 2021).

Demultiplexed CCS reads data files of PacBio Sequel IIe SMRT sequencing have been

deposited with the accession number of PRJNA857037 to BioProject database

(https://www.ncbi.nlm.nih.gov/bioproject/) in National Center for Biotechnology

Information (NCBI), and with the OSD identifier of OSD-625 (DOI: 10.26030/rp5n-0d70)

in NASA GeneLab.

Roots characterization assay

The *Arabidopsis thaliana* T-DNA insertion lines SALK_092504 (*emo1-1*),

SAIL_344_F07 (*emo1-2*), SALK_140748C (*emo2-1*), SALK_091636C (*emo2-2*),

SAIL_205_A08, and WiscDsLox441G3 were obtained from Arabidopsis Biological

Resource Center (https://abrc.osu.edu/). Seeds were sterilized, incubated in ddH_2O at

4°C for 3 days, and then sown on 0.5x Murashige–Skoog medium in Petri dishes as

described above. Plates were placed on vertical racks at 21°C in 24 h LED light

condition at approximately 100 µmol/m^{2}s PAR. The 4-day old were photographed. Then

plants were subjected to three treatments. The first set of plates were kept quiescent

and seedlings were grown to 7-day old and photographed. The second set of plates

were mounted to a 2-dimentional clinostat to be rotated at 4 rounds per min until plants

were 7-day old, and then photographed. The third set of plates were reoriented 90° to

the horizontal, and photographed at 0 h, 6 h and 24 h after reorientation. Roots were

traced and measured using JFilament plugin of ImageJ (Li *et al.*, 2009; Smith *et al.*,

2010). Root growth measurements were processed using the R script

RootMeasurement (https://github.com/eschultzphd/RootMeasurement/tree/master)

(Schultz *et al.*, 2017). The quantitative real-time PCR (qPCR) assay to verify gene

knock-out or knock-down was performed using roots of 7-day old seedlings as

previously described (Zhou *et al.*, 2017). UBQ11 (AT4G05050) was used as the internal

control and primers are listed in Table S2.

FENGC materials availability

501 Requests for a detailed FENGC protocol and updated versions of FOLD should be

502 directed to Michael Kladde (kladde@ufl.edu).

Funding

This work was supported by NASA grants NNX14AT24G and 80NSSC19K0130 to A-LP and RJF. Development of FENGC was supported by Defense Threat Reduction Agency grant HDTRA1-16-1-0048 awarded to MPK as a co-PI and NIH grant R01 CA155390 awarded to MPK.

Acknowledgments

We thank members of UF Space Plants Lab for helpful discussion and assistance in experiments, as well as members of UF ICBR, NextGen DNA sequencing (RRID:SCR_019152), Bioinformatics (RRID:SCR_019120) and Gene Expression and Genotyping (RRID:SCR_019145) cores, for their support and services. We also thank Jason O. Brant for his assistance in developing the FOLD program.

Conflict of interest

A United States patent application (#18/551,004) was filed on September 18, 2023, and is pending. M.Z. and M.P.K. are two of the co-inventors.

Author contributions

MZ, RJF and A-LP designed this study. MZ conducted the experiments and analyzed the data. MZ and MPK developed the FENGC methodology. AR, M-PLG and MPK contributed to the FENGC oligo design program. MZ, MPK, RJF and A-LP wrote the manuscript.

Data availability

528 Demultiplexed CCS reads data files of PacBio Sequel IIe SMRT sequencing have been

529 deposited with the accession number of PRJNA857037 to BioProject database

530 (https://www.ncbi.nlm.nih.gov/bioproject/) in National Center for Biotechnology

531 Information (NCBI), and with the OSD identifier of OSD-625 (DOI: 10.26030/rp5n-0d70)

532 in NASA GeneLab.

References

de Abreu-Neto, J.B., Turchetto-Zolet, A.C., de Oliveira, L.F.V., Bodanese Zanettini,
M.H., and Margis-Pinheiro, M. (2013) *Heavy metal-associated isoprenylated*
plant protein (HIPP): characterization of a family of proteins exclusive to plants.
The FEBS Journal, **280**, 1604–1616.
Califar, B., Sng, N.J., Zupanska, A., Paul, A.-L., and Ferl, R.J. (2020) *Root Skewing-*
Associated Genes Impact the Spaceflight Response of Arabidopsis thaliana.
Frontiers in Plant Science, **11**.
Chen, D.-P., Lin, Y.-C., and Fann, C.S.J. (2016) *Methods for identifying differentially*
methylated regions for sequence- and array-based data. Briefings in Functional
Genomics, **15**, 485–490.
Choi, W.-G., Barker, R.J., Kim, S.-H., Swanson, S.J., and Gilroy, S. (2019) *Variation in*
the transcriptome of different ecotypes of Arabidopsis thaliana reveals signatures
of oxidative stress in plant responses to spaceflight. American Journal of Botany,
106, 123–136.
Edwards, J.R., Yarychkivska, O., Boulard, M., and Bestor, T.H. (2017) *DNA methylation*
and DNA methyltransferases. Epigenetics Chromatin, **10**, 23.
Feng, S., Zhong, Z., Wang, M., and Jacobsen, S.E. (2020) *Efficient and accurate*
determination of genome-wide DNA methylation patterns in Arabidopsis thaliana
with enzymatic methyl sequencing. Epigenetics & Chromatin, **13**, 42.
Giolai, M., Paajanen, P., Verweij, W., Witek, K., Jones, J.D.G., and Clark, M.D. (2017)
Comparative analysis of targeted long read sequencing approaches for
characterization of a plant's immune receptor repertoire. BMC Genomics, **18**, 564.
Grimanelli, D. and Ingouff, M. (2020) *DNA Methylation Readers in Plants. Journal of*
Molecular Biology, **432**, 1706–1717.
Hill, C.B., Wong, D., Tibbits, J., Forrest, K., Hayden, M., Zhang, X.-Q., et al. (2019)
Targeted enrichment by solution-based hybrid capture to identify genetic
sequence variants in barley. Sci Data, **6**, 12.
Jia, Y., Tian, H., Li, H., Yu, Q., Wang, L., Friml, J., and Ding, Z. (2015) *The Arabidopsis*
thaliana elongator complex subunit 2 epigenetically affects root development.
Journal of Experimental Botany, **66**, 4631–4642.
Kishore, K. and Pelizzola, M. (2018) *Identification of Differentially Methylated Regions in*
the Genome of Arabidopsis thaliana. In: *Plant Chromatin Dynamics: Methods*
and Protocols Methods in Molecular Biology (Bemer,M. and Baroux,C., eds) , pp.
61–69. New York, NY: Springer.

Kladde, M.P., ZHOU, M., and NABILSI, N.H. (2022) *Methods and kits for targeted*
cleavage and enrichment of nucleic acids for high-throughput analyses of user-
defined genomic regions.
Kwon, T., Sparks, J.A., Nakashima, J., Allen, S.N., Tang, Y., and Blancaflor, E.B. (2015)
Transcriptional response of Arabidopsis seedlings during spaceflight reveals
peroxidase and cell wall remodeling genes associated with root hair development.
American Journal of Botany, **102**, 21–35.
LeFrois, C.E., Zhou, M., Amador, D.M., Sng, N., Paul, A.-L., and Ferl, R.J. (2016)
Enabling the Spaceflight Methylome: DNA Isolated from Plant Tissues Preserved
in RNAlater Is Suitable for Bisulfite PCR Assay of Genome Methylation.
Gravitational and Space Research, **4**, 28–37.
Li, H., Shen, T., Smith, M.B., Fujiwara, I., Vavylonis, D., and Huang, X. (2009)
Automated actin filament segmentation, tracking and tip elongation
measurements based on open active contour models. In: *2009 IEEE*
International Symposium on Biomedical Imaging: From Nano to Macro , pp.
1302–1305.
Liu, W., Gallego-Bartolomé, J., Zhou, Y., Zhong, Z., Wang, M., Wongpalee, S.P., et al.
(2021) *Ectopic targeting of CG DNA methylation in Arabidopsis with the bacterial*
SssI methyltransferase. Nat Commun, **12**, 3130.
López-Girona, E., Davy, M.W., Albert, N.W., Hilario, E., Smart, M.E.M., Kirk, C., et al.
(2020) *CRISPR-Cas9 enrichment and long read sequencing for fine mapping in*
plants. Plant Methods, **16**, 121.
Nabilsi, N.H., Deleyrolle, L.P., Darst, R.P., Riva, A., Reynolds, B.A., and Kladde, M.P.
(2014) *Multiplex mapping of chromatin accessibility and DNA methylation within*
targeted single molecules identifies epigenetic heterogeneity in neural stem cells
and glioblastoma. Genome Res, **24**, 329–339.
NEB *NEBNext® Enzymatic Methyl-seq (EM-seq^{TM}).*
Onda, Y., Takahagi, K., Shimizu, M., Inoue, K., and Mochida, K. (2018) *Multiplex PCR*
Targeted Amplicon Sequencing (MTA-Seq): Simple, Flexible, and Versatile SNP
Genotyping by Highly Multiplexed PCR Amplicon Sequencing. Frontiers in Plant
Science, **9**.
Ostezan, A., McDonald, S.C., Tran, D.T., Souza, R.S.E., and Li, Z. (2021) *Target region*
sequencing and applications in plants. J. Crop Sci. Biotechnol., **24**, 13–26.
Paul, A.-L., Amalfitano, C.E., and Ferl, R.J. (2012) *Plant growth strategies are*
remodeled by spaceflight. BMC Plant Biology, **12**, 232.
Paul, A.-L., Haveman, N., Califar, B., and Ferl, R.J. (2021) *Epigenomic Regulators*
Elongator Complex Subunit 2 and Methyltransferase 1 Differentially Condition the
Spaceflight Response in Arabidopsis. Frontiers in Plant Science, **12**.
Paul, A.-L., Sng, N.J., Zupanska, A.K., Krishnamurthy, A., Schultz, E.R., and Ferl, R.J.
(2017) *Genetic dissection of the Arabidopsis spaceflight transcriptome: Are some*
responses dispensable for the physiological adaptation of plants to spaceflight?
PLOS ONE, **12**, e0180186.
Paul, A.-L., Zhou, M., Callaham, J.B., Reyes, M., Stasiak, M., Riva, A., et al. (2017)
Patterns of Arabidopsis gene expression in the face of hypobaric stress. AoB
PLANTS, **9**, plx030.

Paul, A.-L., Zupanska, A.K., Schultz, E.R., and Ferl, R.J. (2013) *Organ-specific remodeling of the Arabidopsis transcriptome in response to spaceflight. BMC Plant Biology*, **13**, 112.

Schultz, E.R., Zupanska, A.K., Sng, N.J., Paul, A.-L., and Ferl, R.J. (2017) *Skewing in Arabidopsis roots involves disparate environmental signaling pathways. BMC Plant Biology*, **17**, 31.

Smith, M.B., Li, H., Shen, T., Huang, X., Yusuf, E., and Vavylonis, D. (2010) *Segmentation and tracking of cytoskeletal filaments using open active contours. Cytoskeleton*, **67**, 693–705.

Thieme, C.J., Rojas-Triana, M., Stecyk, E., Schudoma, C., Zhang, W., Yang, L., et al. (2015) *Endogenous Arabidopsis messenger RNAs transported to distant tissues. Nature Plants*, **1**, 1–9.

Tran, H., Zhu, H., Wu, X., Kim, G., Clarke, C.R., Larose, H., et al. (2018) *Identification of Differentially Methylated Sites with Weak Methylation Effects. Genes*, **9**, 75.

Wang, Y., An, C., Zhang, X., Yao, J., Zhang, Y., Sun, Y., et al. (2013) *The Arabidopsis elongator complex subunit2 epigenetically regulates plant immune responses. Plant Cell*, **25**, 762–776.

Wheeler, R.M. (2010) *Plants for human life support in space: from Myers to Mars. Gravitational and Space Biology*, **23**, 25–36.

Zabel, P., Bamsey, M., Schubert, D., and Tajmar, M. (2016) *Review and analysis of over 40 years of space plant growth systems. Life Sciences in Space Research*, **10**, 1–16.

Zhang, H., Lang, Z., and Zhu, J.-K. (2018) *Dynamics and function of DNA methylation in plants. Nat Rev Mol Cell Biol*, **19**, 489–506.

Zhou, M., Nabilsi, N.H., Wang, A., Gauthier, M.-P.L., Murray, K.O., Azari, H., et al. (2022) *Flap-enabled next-generation capture (FENGC): precision targeted single-molecule profiling of epigenetic heterogeneity, chromatin dynamics, and genetic variation.* 2022.11.08.515732.

Zhou, M., Paul, A.-L., and Ferl, R.J. (2017) *Data for characterization of SALK_084889, a T-DNA insertion line of Arabidopsis thaliana. Data in Brief*, **13**, 253–258.

Zhou, M., Sng, N.J., LeFrois, C.E., Paul, A.-L., and Ferl, R.J. (2019) *Epigenomics in an extraterrestrial environment: organ-specific alteration of DNA methylation and gene expression elicited by spaceflight in Arabidopsis thaliana. BMC Genomics*, **20**, 205.

Supporting Information

Table S1. Information of 108 capture targets.

Table S2. Oligos used in this study.

Table S3. Filtered HiFi reads on target and targets detected.

Table S4. Number of filtered HiFi reads aligned to 108 targets.

Figure S15. Root gravitropism in mutants of *EMO1*, *EMO2* or AT4G04990.

Figure Legends

Figure 1. Application of flap-enabled next-generation capture (FENGC) in root
samples of *Arabidopsis* experiencing spaceflight.

(A) Percentage of filtered HiFi reads on target for all 108 targets. Reads with more than
three consecutive methylated CHH sites were removed. GCR, Ground Col Roots; FCR,
Flight Col Roots; GER, Ground control *elp2-5* Roots; FER, Flight *elp2-5* Roots. (B)
Percentage of targets detected by at least 1 filtered HiFi read or at least 10 filtered HiFi
reads. (C) Correlation between numbers of total filtered HiFi reads of all samples
aligned to targets and the GC content of target regions. Log transformation of reads
number was done by $\log_2$(read number +1). R-squared value and *p* value of logarithmic
regression analysis are shown. (D) Correlation coefficients between biological replicates
of FENGC and between FENGC and WGBS for average cytosine methylation levels in
seven example targets.

Figure 2. FENGC results for *CML46.*

(A) Genic region of *CML46*. The target region is indicated by brown rectangle (target ID:
AT5G39670-1 in S1 Table). A MYB-like binding element is indicated by green brick. The
transcription start site is indicated by arrow. (B) Expression levels of *CML46* in Ground
control Col-0 Roots (GCR), Flight Col-0 Roots (FCR), Ground control *elp2-5* Roots
(GER) and Flight *elp2-5* Roots (FER). Gene expression is indicated by average count
per million mapped reads (CPM) in RNA-seq reported by Paul et al. 2021. Lowercase
letters indicate significant differences between samples (FDR < 0.05 in DESeq2). (C)
Average CG methylation level within the target region. The relative position of MYB-like
binding element shown in (A) is indicated by green brick. The region indicated by
dashed frame shows significant differences of CG methylation level between FCR and
GCR (One-way ANOVA with post-hoc Tukey HSD test; *, *p* < 0.05). (D) Distribution of
proportion of methylated CG sites per molecule in filtered, full-length HiFi reads aligned

to the target region. The box indicates the upper and lower quartiles of the data set. The

whiskers extend to the largest or the smallest data point that is within 1.5 times the

interquartile range. The data points of mPPM are represented by dots. The line and "x"

in the box indicate median and mean, respectively. The significant differences of

methylation proportion are shown (Kruskal-Wallis test; *, $p < 0.05$; **, $p < 0.01$). (E)

Heatmaps showing CG methylation of single molecules. Red indicates two

consecutively methylated cytosines, black indicates two consecutively unmethylated

cytosines, and gray indicates borders between methylated and unmethylated cytosines.

The reads from the four biological replicates (Table S4) were combined to generate

heatmaps for each sample. The relative position of MYB-like binding element shown in

(A) and (C) is indicated by green brick. The region used for the CG methylation level

comparison in (C) is indicated by dashed frame. Around 6.9% reads in GCR and 12%

reads in FCR containing at least 2 consecutively methylated CG sites in both areas

flanking the MYB-like binding element are labeled by blue bricks. One-way ANOVA with

post-hoc Tukey HSD test was performed for pairwise comparisons of percentage of

reads with this signature. FER vs FCR, $p < 0.01$.

Figure 3. FENGC results for *PDF1.1.*

(A) Genic region of *PDF1.1*. The target region is indicated by brown rectangle (target ID:

AT1G75830-1 in Table S1). A MeJA-responsive element is indicated by green brick.

The transcription start site is indicated by the arrow. (B) Expression levels of *PDF1.1*

that are presented in the same way of Figure 2B. (C) Average CG, CHG or CHH

methylation levels over all sequenced molecules of the target region. The relative

position of MeJA-responsive element shown in (A) is indicated by green brick. The

significant differences of methylation levels between samples for CHG and CHH sites in

the target region are shown. One-way ANOVA with post-hoc Tukey HSD test; *, $p <$

0.05; **, $p < 0.01$; ***, $p < 0.001$. Black arrows indicate DmCs detected in WGBS using

the threshold of methylation difference > 0.2 and FDR < 0.01. Arrows of solid line and

dot line indicate DmCs of GER vs GCR and FER vs GER, respectively. (D) Distribution

of proportion of methylated CG, CHG or CHH sites per molecule in filtered full-length

HiFi reads aligned to the target region. The significant differences of methylation proportion are shown (Kruskal-Wallis test; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$). (E) Heatmaps showing the cytosine methylation in single molecules. Reads from the four biological replicates (Table S4) were combined to generate heatmaps for each sample. If the number of combined reads exceeded 1,000, 1,000 reads were randomly selected and plotted. The three heatmap panels show the patterns of methylation at CG, CHG, and CHH sites (vertical white lines) as indicated at bottom along the same DNA molecule in each row. The color legend is the same as that in Figure 2E. Reads were hierarchically clustered using Kendall's Tau based on CG methylation. The relative position of MeJA-responsive element shown in (A) is indicated by green brick. A footprint (blue ovals) with consecutive protection against methylation of the first 2 CG sites in the area flanking the MeJA-responsive element is present in 8.4%, 18.0%, 39.4% and 44.0% molecules in GCR, FCR, GER and FER, respectively. One-way ANOVA with post-hoc Tukey HSD test was performed for pairwise comparisons of percentage of reads with this footprint. FCR vs GCR, $p < 0.05$; GER vs GCR, $p < 0.001$; FER vs FCR, $p < 0.001$.

Figure 4. DNA methylation within same molecules in the target region of *PDF1.1.*

(A) Proportion of methylated CG, CHG and CHH sites within the same molecule in the captured region of AT1G75830. There are 120, 1009, 345 and 1310 full-length reads covering all CG, CHG and CHH sites in this region (target ID: AT1G75830-1 in Table S1), which are mapped for GCR, FCR, GER and FER samples, respectively. Ochre, gold and dark green dots indicate CG, CHG and CHH methylation proportions, respectively, in the same column, each representing one read. (B) Comparisons of methylation proportions per molecule in reads with a footprint and in all reads in the captured region of AT1G75830. CHG and CHH methylation in GCR, FCR, GER and FER samples are plotted, respectively. The footprint is defined as the first 2 CG sites in this target region that are both unmethylated. The significant differences of methylation proportion are shown (Kruskal-Wallis test; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$).

Figure 5. FENGC results for *EMO1*.

(A) Genic region of *EMO1*. The target region is indicated by brown rectangle (target ID: AT1G27565-1 in Table S1). An anaerobic-responsive element (ARE) is indicated by green brick. The transcription start site (TSS) is indicated by arrow. (B) Expression levels of *EMO1* that are presented in the same way of Figure 2B. (C) Average CG, CHG or CHH methylation levels in cytosines within the target region. The relative position of ARE and TSS shown in (A) are indicated by green brick and arrow, respectively. The significant differences of CG methylation level between samples in the target region are shown, in which the first CG site is omitted. One-way ANOVA with post-hoc Tukey HSD test; *, $p < 0.05$; ***, $p < 0.001$. Black arrows indicate DmCs of FER vs GER detected in WGBS using the threshold of methylation difference > 0.2 and FDR < 0.01. (D) Distribution of proportion of methylated CG, CHG or CHH sites per molecule in filtered full-length HiFi reads aligned to the target region. The significant differences of methylation proportion are shown (Kruskal-Wallis test; *, $p < 0.05$). (E) Heatmaps showing the cytosine methylation in single molecules. The heatmap panels and color legend are the same as that in Figure 3E. Reads were hierarchically clustered using Euclidean distance based on CG methylation. The percentages of reads without methylated CG sites and reads with at least 9 consecutive unmethylated CG sites at the 3' end are shown on the left and right side of the CG panel, respectively. One-way ANOVA with post-hoc Tukey HSD test was performed for pairwise comparisons. For percentages of reads without methylated CG sites, FER vs GER, $p < 0.05$; GER vs GCR, $p < 0.001$; FER vs FCR, $p < 0.01$. For percentages of reads with at least 9 consecutive unmethylated CG sites at the 3' end, FCR vs GCR, $p < 0.01$; GER vs GCR, $p < 0.01$.

Figure 6. FENGC results for *EMO2*.

(A) Genic region of *EMO2*. The target region is indicated by brown rectangle (target ID: AT5G52710 in Table S1). Two MeJA-responsive elements are indicated by green bricks. The transcription start site (TSS) is indicated by arrow. (B) Expression levels of *EMO2* that are presented in the same way of Figure 2B. (C) Average CG, CHG or CHH

methylation levels in cytosines within the target region. The relative position of MeJA-

responsive elements shown in (A) are indicated by green bricks. The significant

differences of CG and CHG methylation levels between samples in the target region are

shown. One-way ANOVA with post-hoc Tukey HSD test; **, $p < 0.01$; ***, $p < 0.001$.

Black arrows indicate the DmCs detected in WGBS using the threshold of methylation

difference > 0.2 and FDR < 0.01. Arrows with solid line, triangle, round end, and dot line

indicate DmCs of GER vs GCR, FER vs FCR, FCR vs GCR, and FER vs GER,

respectively. (D) Distribution of proportion of methylated CG, CHG or CHH sites per

molecule in filtered full-length HiFi reads aligned to the target region. The significant

differences of methylation proportion are shown (Kruskal-Wallis test; ***, $p < 0.001$). (E)

Heatmaps showing the cytosine methylation in single molecules. The reads clustering,

heatmap panels and color legend are the same as that in Figure 5E.

Figure 7. Enhanced root growth in mutants of *EMO1* or *EMO2*.

(A) 7-day old seedlings of Col-0, *emo1-1*, *emo1-2*, *emo2-1*, and *emo2-2*. Scale bars

indicate 10 mm. (B) Primary root length of 4-day old and 7-day old seedlings for each

line. (C) Average root growth rate (mm per day) from day-0 to day-4, and from day-4 to

day-7. Student's t test, *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. Data ± SD, n = 33 to 55

for each condition.

Table 1. Comparison of multiplexed, targeted DNA sequencing methods in plants.

Light gray indicates advantages and dark gray indicates disadvantages.

Method	Custom design liberty	Read length	Example of sequencing depth	Example of on-target read % in plant genome	Example of target number	Cost	Adaptation in DNA methylation detection
Multiplexed PCR	limited by low complexity of sequences	Short (e.g., < 200 bp) (Onda *et al.*, 2018)	1,000x (Onda *et al.*, 2018)	50.9 % (Onda *et al.*, 2018)	443 markers (Onda *et al.*, 2018)	low	no

Probe hybridization-based	**limited by commercial panels and costly design**	**short/long (e.g., 3.5 kb)** (Giolai *et al.*, 2017)	**55x** (Giolai *et al.*, 2017)	**52.9%** (Hill *et al.*, 2019)	**190 genes** (Hill *et al.*, 2019)	**high due to modified probes**	**no**
CRISPR-based	**limited by PAM sites and cleavage efficiency**	**short/long (e.g. 8 kb)** (López-Girona *et al.*, 2020)	**~200x** (López-Girona *et al.*, 2020)	**3.04%** (López-Girona *et al.*, 2020)	**1 gene** (López-Girona *et al.*, 2020)	**high due to RNA oligos and specific proteins**	**no**
FENGC	**cleavage for any site**	**short/long (e.g., > 500 bp)**	**10-5,000x**	**~90%**	**102 genic regions**	**low**	**long-read EM-seq**

A

-2132   -1490

MeJA responsiveness: TGACG

AT1G75830
PDF1.1
plant defensin 1.1

B

PDF1.1

C

GER vs GCR **

FCR vs GCR ***
GER vs GCR *

D

E

Col   8.4%   18%

elp2   39.4%   44.0%

CG   CHG   CHH   CG   CHG   CHH

Ground control   Spaceflight

Methylated   Unmethylated
CG/CHG/CHH   CG/CHG/CHH

Border

Full-length reads mapped to captured region AT1G75830-1

A

Col-0   *emo1-1*   *emo1-2*   *emo2-1*   *emo2-2*

B

Root length (mm)

- Col-0
- *emo1-1*
- *emo1-2*
- *emo2-1*
- *emo2-2*

4-day old   7-day old

*** *

***

C

Average growth rate (mm/d)

day-0 to day-4   day-4 to day-7

*** *

*** **