Single-molecule long-read methylation profiling reveals regional DNA

methylation regulated by Elongator Complex Subunit 2 in *Arabidopsis*

roots experiencing spaceflight

Mingqi Zhou[1], Marie-Pierre L. Gauthier[2], Michael P. Kladde[2], Robert J. Ferl[1,3,], Anna-*

Lisa Paul[1,4,]*

[1]Department of Horticultural Sciences, University of Florida, 2550 Hull Road, Fifield Hall,

Gainesville FL, 32611, USA

[2]Department of Biochemistry and Molecular Biology, University of Florida, 2033 Mowry

Rd, Gainesville, FL 32608, USA

[3]Office of Research, University of Florida, 1523 Union Rd, Grinter Hall, Gainesville, FL,

32611, USA

[4]Interdisciplinary Center for Biotechnology Research, University of Florida, 2033 Mowry

Road, Gainesville FL, 32610, USA

*Correspondance: Robert J. Ferl (robferl@ufl.edu), **Anna-Lisa Paul** (alp@ufl.edu)

Abstract

Flap-Enabled Next-Generation Capture (FENGC), a novel targeted DNA capture

technique, has been applied to *Arabidopsis* root samples from the Advanced Plant

Experiment-04 - Epigenetic Expression (APEX-04) experiment onboard the International

Space Station. Single-molecule long-read profiling reveals regional DNA methylation

alteration that was not previously uncovered by population-based methods during

spaceflight.

DNA methylation epigenetically modulates gene expression and impacts chromatin

structure to function in physiological adaptation to environmental changes in eukaryotes

(Zhang et al., 2018). Plants will play an important role in the human life support systems

envisioned for spaceflight and planetary exploration habitats (Wheeler, 2010; Zabel et

al., 2016). Spaceflight presents a unique environmental challenge for plants, an

environment that is outside the evolutionary experience of any terrestrial organism.

Epigenetic change plays an important role in environmental adaptation, and it is crucial

to understand the role of epigenetics in plant adaptation to this novel environment (Paul

et al., 2021; Zhou et al., 2019). The Advanced Plant Experiment-04 - Epigenetic

Expression (APEX-04) experiment onboard the International Space Station (ISS)

examined the spaceflight-altered cytosine methylation in *Arabidopsis thaliana* using

elp2-5, a mutant of an epigenetic regulator, Elongator Complex Subunit 2 (ELP2), along

with Col-0, a wild-type control (Paul et al., 2021). Genome-wide analysis revealed

remodeled transcriptomes and DNA methylomes during spaceflight, remodeling that

was regulated by ELP2. Phenotypically, root growth in *elp2-5* did not set up a typical

spaceflight directionality, and a majority of differentially expressed genes in *elp2-5* roots

were also differentially methylated between spaceflight samples and ground controls

(Paul et al., 2021).

Flap-Enabled Next-Generation Capture (FENGC) is a novel, strand-specific DNA

capture method that allows DNA methylation analysis with long reads of up to ~1 kb in

user defined target regions (Zhou et al. 2022). FENGC allows the characterization of a

single molecule over long ranges, thereby providing a deeper insight into the

methylation state afforded by averaged state of methylation obtained by multiple short

reads. In addition, compared with whole genome bisulfite sequencing (WGBS), FENGC

increases coverage depth in regions of interest and avoids cost of reads in unwanted

areas. The accuracy and efficiency of enzymatic methyl-seq, which is part of the

FENGC procedure, is verified in plants (Feng et al., 2020); however, FENGC has yet to

be applied to methylation analysis in plants.

Therefore, we set out to demonstrate the effective application of FENGC in plant

methylation contexts. The observed phenotypical and molecular alteration in roots of

Col-0 and *elp2-5* grown on ISS in APEX-04 present a well-characterized, short-read

methylation dataset for comparison of methylation profiles assayed by FENGC single-

molecule long reads.

The target panel for capture and assay contained 108 genic regions ranging from 509 to 704 nt (Supplemental Table 1), with mean of 580 nt and median of 572 nt, within the promoter or gene body regions of epigenetically regulated genes derived from our APEX-04 spaceflight and related experiments (Paul et al., 2017a; Paul et al., 2021; Paul et al., 2013; Zhou et al., 2019). The capture was performed in a single-tube multiplex reaction followed by enzymatic methyl-PCR (Oligos are listed in Supplemental Table 2). The FENGC products were subjected to PacBio long-read HiFi sequencing. The samples consistently yielded ~90% on- : off-target ratio and, on average, 82% targets were detected by at least 10 reads after filtering and combining the four biological replicates from each sample (Figure 1A, 1B, and Supplemental Table 3). The GC content of the 108 targets ranged from 19% to 52%, which positively correlated with the aligned reads number (Figure 1C, Supplemental Figure 1), which was opposite to FENGC results in DNA materials with GC content above 50% in most of targets (Zhou et al. 2022), indicating that both high and low GC content of targets influenced the reads yield of FENGC products.

One major advantage of single-molecule profiling of DNA methylation is that the contiguous long reads allow assessment of every cytosine within same DNA molecule, thereby presenting a coherent regional methylation status of that entire gene element within a cell. In addition, the average methylation levels in cytosines across all molecules can be derived from the data for comparison to WGBS data. Among the test targets for this study, we identified genes that differentially responded to spaceflight in *elp2-5* or Col-0 and showed significant changes in average methylation level in cytosines as well as in distribution of the proportion of methylated sites per molecule using FENGC.

We present four examples associated with defense responses involved in spaceflight (Supplemental Figure 2), the methylation levels of which exhibited correlation coefficients above 0.96 between any two of biological replicates and above 0.95 between FENGC and WGBS results (Figure 1D). The first three examples examine FENGC data for CG methylation sites as no significant difference was observed for CHG and CHH (data not shown). *CML46* (AT5G39670), a gene encoding a calmodulin-

like protein showing significant transcriptional down-regulation during spaceflight in Col-
0 but not *elp2-5* plants, was assayed in a promoter region containing a MYB-like DNA
binding element (Supplemental Figure 3A). More than 99% of the mapped reads are
full-length, containing all CG sites in this region of *CML46* (Supplemental Figure 3B).
Flight Col-0 Roots (FCR) and Ground control Col-0 Roots (GCR), but not roots from
elp2-5 (FER and GER), showed a significant difference in the CG methylation level in
an area flanking the MYB-like element, which contained five CG sites (Supplemental
Figure 3C, in dotted frame). Consistent with these data, CG methylation proportion per
molecule in this target region was significantly higher in FCR than GCR (Supplemental
Figure 3D). A significant difference between GCR and GER was also observed. FENGC
showed more molecules containing consecutive methylated CG sites in FCR (12%)
compared to GCR (6.9%) in the area flanking this *cis*-regulatory element (Supplemental
Figure 3E). Consistency between transcriptional gene expression, population-based
methylation level derived from WGBS, and FENGC-derived methylation proportion per
molecule supported the biological conclusion that ELP2 played a negative role in
expression and a positive role in promoter methylation of *CML46* in spaceflight.
ABCB17 (AT3G28380), an ATP-binding cassette transporter gene that also showed a
significantly lower transcriptional expression level in FCR than GCR (Supplemental
Figure 2), was assayed in a gene body region (Supplemental Figure 4A). More than 98%
of the mapped reads contained all CG sites in this region (Supplemental Figure 4B).
The averaged regional CG methylation levels were in a same trend, which did not show
significant changes between any pair of samples (Supplemental Figure 4C); however,
the CG methylation proportion per molecule in this target region was significantly higher
in FCR than in GCR (Supplemental Figure 4D). The FENGC single-molecule data
showed that 3.8% of the reads in FCR possessed at least 6 consecutive methylated CG
sites while in GCR the percentage was 0.8% (Supplemental Figure 4E). Biologically,
ABCB17 was repressed by spaceflight and ELP2 contributed to its transcription and
gene body methylation. Similarly, in the promoter region of *DEFL* (AT4G22217), a
defensin-like gene (Supplemental Figure 5A and 5B), the patterns of averaged CG
methylation levels were highly similar and no significant alterations were detected
between any two samples (Supplemental Figure 5C); however, CG methylation

proportion per molecule was significantly higher in FER than GER (Supplemental Figure
5D and 5E). This higher methylation proportion correlates with the lower transcription
level of this gene in FER compared to GER (Supplemental Figure 2). ELP2 enhanced
expression and repressed promoter methylation of *DEFL* in spaceflight. These
examples suggest that FENGC single-molecule level methylation profiling can be more
revealing of consequential methylation changes compared with population-based
methylation methods such as WGBS, especially when alterations potentially occur in a
subpopulation of cells within the sample.

The fourth example examines another advantage of long-read single-molecule FENGC
assay of DNA methylation in plants, which is that all three methylation sequence
contexts (CG/CHG/CHH) can be identified and plotted for the same molecule, thereby
aggregating the entire methylation status of a gene region. For this example, we chose
a promoter fragment containing a MeJA-responsive element in *PDF1.1* (AT1G75830), a
gene coding a pathogenesis-related protein belonging to plant defensin family (Figure
1E). This target was captured with more than 92.6% full-length, aligned reads
containing three types of cytosines (Figure 1F). The average methylation level in this
region showed significant differences at CHG and CHH sites between GER and GCR,
as well as in CHH sites between FCR and GCR (Figure 1G). These data correlated with
the previous WGBS data (Paul et al., 2021). For methylation proportion per molecule,
significant differences were detected in all pairwise comparisons between flight
conditions and genotypes at cytosines in all three methylation contexts (Figure 1H). The
patterns of methylation proportion in DNA molecules demonstrated that three types of
cytosines showed a trend of co-methylation in this region among cells in the population
(Supplementary Figure 6). Within the same molecule, methylation proportion of CG is
higher than CHG in 97.9% reads, and that of CHG is higher than CHH in 87.5% reads
(Supplemental Figure 7). The CG methylation proportions were above 0.57 in all reads
of GCR and above 0.28 in GER, while FCR and FER had reads with CG methylation
proportion as low as 0. These data indicate that more molecules were hypomethylated
in this region in *elp2-5* compared with Col-0, and in spaceflight compared with ground
control.

To examine the spatial distribution of these methylation differences, three separate but

correlated panels were plotted for CG, CHG and CHH methylation in the same DNA

molecules per row (Figure 1I). In these plots, black areas are unmethylated regions, red

areas are methylated regions and gray areas are transition zones between methylated

and unmethylated regions. It is visually apparent that spaceflight results in more

unmethylated regions in Col-0, with the deepest loss of methylation concentrated near

the MeJA-responsive element. It is also visually apparent that *elp2-5* shows a deeper

loss of methylation in that region. For the CG methylation context, a footprint showing

consecutive non-methylation of the first 2 CG sites in flanking area of the MeJA-

responsive element was identified in 8.4%, 18.0%, 39.4% and 44.0% reads in GCR,

FCR, GER and FER, respectively, quantifying the loss of CG methylation in this area.

Interestingly, 60%, 86.5%, 97.5% and 97.6% reads with consecutive non-methylation in

the second and the third CG sites in GCR, FCR, GER and FER, respectively, were also

included in this footprint and showed consecutive non-methylation of the first 3 CG sites.

In addition, CHG and CHH methylation proportions were significantly lower in the

footprint area in the reads containing this footprint (Supplementary Figure 8). These

were consistent with previously observed high frequency of co-

methylation/demethylation in nearby cytosines in *Arabidopsis* (Tran et al., 2018), and

demonstrated that FENGC provides further opportunities for studying all three types of

co-methylation in same molecules.

These changes in average methylation level of cytosines or methylation proportion per

molecule could be due to modulation by ELP2 in a diverse manner during spaceflight.

Four targets shown here are associated with plant defense signaling, which is largely

utilized in spaceflight adaptation of plants and may be involved in regulation of root

growth phenotypes during spaceflight (Califar et al., 2020; Paul et al., 2017b). The

targets possessing changes that cannot be observed in a population-based average

level but shown by consecutive methylation status in each same molecule would be

novel candidates for further investigation of downstream pathways elicited by ELP2.

While short-read WGBS only detects average methylation level alteration in cytosines,

targeted single-molecule profiling further reveals the detailed epigenetic modification

occurring in cell subpopulations, identifies epialleles with low frequency, and therefore

provides an additional power to population-based data for methylation analysis and
identification of epigenetic modification substrates.

Supplemental Information

Supplemental materials and methods

Supplemental Figure 1. Correlation between numbers of filtered HiFi reads aligned to
targets and the GC content of target regions in each sample.

Supplemental Figure 2. Expression level of AT5G39670, AT3G28380, AT4G22217, and
AT1G75830 in roots of Col-0 and elp2-5 in APEX04 spaceflight experiment.

Supplemental Figure 3. FENGC results for AT5G39670.

Supplemental Figure 4. FENGC results for AT3G28380.

Supplemental Figure 5. FENGC results for AT4G22217.

Supplemental Figure 6. Correlation between proportions of different types of methylated
cytosines within the same molecule in the captured region of AT1G75830.

Supplemental Figure 7. Proportion of methylated CG, CHG and CHH sites within the
same molecule in the captured region of AT1G75830.

Supplemental Figure 8. Comparisons of methylation proportions per molecule in reads
with a footprint and in all reads.

Supplemental Table 1. Information of 108 capture targets.

Supplemental Table 2. Oligos used in this study.

Supplemental Table 3. HiFi reads on target and targets detected after removing reads
with more than three consecutive methylated CHH sites.

Supplemental Table 4. Number of filtered HiFi reads aligned to 108 targets.

Funding

This work was supported by NASA grants NNX14AT24G and 80NSSC19K0130 to A.-L.P. and R.J.F. Development of FENGC was supported by Defense Threat Reduction Agency grant HDTRA1-16-1-0048 awarded to M.P.K. as a co-PI and NIH grant R01 CA155390 awarded to M.P.K.

Author contributions

M.Z., R.J.F. and A.-L.P. designed this study. M.Z. conducted the experiments and analyzed the data. M.-P.L.G. and M.P.K. contributed to the FENGC oligo design program. M.Z., R.J.F., A.-L.P. and M.P.K. wrote the manuscript.

Accession numbers

Demultiplexed CCS reads data files of PacBio Sequel IIe SMRT sequencing have been deposited to BioProject database (https://www.ncbi.nlm.nih.gov/bioproject/) under accession number PRJNA857037 in National Center for Biotechnology Information (NCBI).

Acknowledgments

We thank members of UF Space Plants Lab for helpful discussion and members of UF ICBR, NextGen DNA sequencing (RRID:SCR_019152), Bioinformatics (RRID:SCR_019120) and Gene Expression and Genotyping (RRID:SCR_019145) cores, for their support and services. A patent has been submitted as an international application (No. PCT/US2022/020624). M.Z. and M.P.K. are two of co-inventors of this patent. The priority date is March 16, 2021, and international filing date is March 16, 2022. The patent release date is Sept. 22, 2022.

 Legends

Figure 1. Application of flap-enabled next-generation capture (FENGC) in root

samples of *Arabidopsis* experiencing spaceflight.

(A) Percentage of filtered HiFi reads on target for all 108 targets. Reads with more than

three consecutive methylated CHH sites were removed. GCR, Ground Col Roots; FCR,

Flight Col Roots; GER, Ground control *elp2* Roots; FER, Flight *elp2* Roots.

(B) Percentage of targets detected by at least 1 filtered HiFi read or at least 10 filtered

HiFi reads.

(C) Correlation between numbers of total filtered HiFi reads of all samples aligned to

targets and the GC content of target regions. Log transformation of reads number was

done by Log_2(read number +1). R-squared value and *p* value for logarithmic regression

analysis are shown.

(D) Correlation coefficients between biological replicates of FENGC and between

FENGC and whole-genome bisulfite sequencing (WGBS) for average methylation levels

in cytosines.

(E) Genic region of AT1G75830 (PDF1.1). The target region is indicated by brown

rectangle (target ID: AT1G75830-1 in Supplemental Table 1). A MeJA-responsive

element is indicated by green brick. The transcription start site is indicated by the arrow.

(F) Percentage of full-length reads containing all CG, CHG or CHH sites in the target

region, and truncated reads with 1 or more cytosines missing. For CHH sites, 1 site

missing is allowed in the full-length reads.

(G) Average CG, CHG or CHH methylation levels in cytosines within the target region.

The relative position of MeJA-responsive element shown in **(E)** is indicated by green

brick. The significant differences of methylation levels between samples for CHG and

CHH sites in the target region are shown. One-way ANOVA with post-hoc Tukey HSD

test; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. Black arrows indicate the differentially

methylated cytosines (DmCs) detected in WGBS using the threshold of methylation

difference > 0.2 and $p < 0.01$. All six sites were significantly different between GER and
GCR.

(H) Distribution of proportion of methylated CG, CHG or CHH sites per molecule in
filtered full-length HiFi reads aligned to the target region. The box indicates the upper
and lower quartiles of the data set. The whiskers extend to the largest or the smallest
data point that is within 1.5 times the interquartile range. The data points of methylation
proportion per molecule are represented by dots. The line and "x" in the box indicate
median and mean, respectively. The significant differences of methylation proportion
are shown (Kruskal-Wallis test; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$).

(I) Heatmaps showing the cytosine methylation in single molecules. Reads from the four
biological replicates (Supplemental Table 4) were combined to generate heatmaps for
each sample. When the number of combined reads exceeded 1000, 1000 reads were
randomly selected and plotted. The three heatmap panels show the patterns of
methylation at CG, CHG, and CHH sites (vertical white lines) as indicated at bottom
along the same DNA molecule in each row. Red indicates two consecutively methylated
cytosines, black indicates two consecutively unmethylated cytosines, and gray indicates
borders between methylated and unmethylated cytosines. Reads were hierarchically
clustered using Kendall's Tau based on CG methylation. The relative positions of MeJA-
responsive element shown in **(E)** and differentially methylated cytosines detected by
WGBS shown in **(G)** in the target region are indicated by green brick and black arrows,
respectively. A footprint (blue ovals) with consecutive protection against methylation of
the first 2 CG sites in the area flanking the MeJA-responsive element is present in 8.4%,
18.0%, 39.4% and 44.0% molecules in GCR, FCR, GER and FER, respectively. One-
way ANOVA with post-hoc Tukey HSD test was performed for pairwise comparisons of
percentage of reads with this footprint. FCR vs GCR, $p < 0.05$; GER vs GCR, $p < 0.001$;
FER vs FCR, $p < 0.001$.

References

Califar, B., Sng, N.J., Zupanska, A., Paul, A.-L., and Ferl, R.J. (2020). Root Skewing-Associated
Genes Impact the Spaceflight Response of. Front Plant Sci **11**:239. 10.3389/fpls.2020.00239.

Feng, S., Zhong, Z., Wang, M., and Jacobsen, S.E. (2020). Efficient and accurate determination of
genome-wide DNA methylation patterns in Arabidopsis thaliana with enzymatic methyl sequencing.
Epigenetics Chromatin **13**:42. 10.1186/s13072-020-00361-9.

Paul, A.-L., Zhou, M., Callaham, J., Reyes, M., Stasiak, M., Riva, A., Zupanska, A., Dixon, M., and
Ferl, R. (2017a). Patterns of Arabidopsis gene expression in the face of hypobaric stress. Aob Plants
910.1093/aobpla/plx030.

Paul, A.-L., Zupanska, A.K., Schultz, E.R., and Ferl, R.J. (2013). Organ-specific remodeling of the
Arabidopsis transcriptome in response to spaceflight. BMC Plant Biol **13**:112. 10.1186/1471-2229-13-112.

Paul, A.-L., Haveman, N., Califar, B., and Ferl, R.J. (2021). Epigenomic Regulators *Elongator Complex*
Subunit 2 and *Methyltransferase 1* Differentially Condition the Spaceflight Response in Arabidopsis. Front
Plant Sci **12**:691790. 10.3389/fpls.2021.691790.

Paul, A.-L., Sng, N.J., Zupanska, A.K., Krishnamurthy, A., Schultz, E.R., and Ferl, R.J. (2017b).
Genetic dissection of the Arabidopsis spaceflight transcriptome: Are some responses dispensable for the
physiological adaptation of plants to spaceflight? PLoS One **12**:e0180186. 10.1371/journal.pone.0180186.

Tran, H., Zhu, H., Wu, X., Kim, G., Clarke, C.R., Larose, H., Haak, D.C., Askew, S.D., Barney, J.N.,
Westwood, J.H., et al. (2018). Identification of Differentially Methylated Sites with Weak Methylation
Effects. Genes (Basel) **9**10.3390/genes9020075.

Wheeler, R. (2010). Plants for human life support in space: From Myers to Mars. Gravitational and Space
Biology **23**:25-35.

Zabel, P., Bamsey, M., Schubert, D., and Tajmar, M. (2016). Review and analysis of over 40 years of
space plant growth systems. Life Sci Space Res (Amst) **10**:1-16. 10.1016/j.lssr.2016.06.004.

Zhang, H., Lang, Z., and Zhu, J.K. (2018). Dynamics and function of DNA methylation in plants. Nat Rev
Mol Cell Biol **19**:489-506. 10.1038/s41580-018-0016-z.

Zhou, M., Sng, N., LeFrois, C., Paul, A.-L., and Ferl, R. (2019). Epigenomics in an extraterrestrial
environment: organ-specific alteration of DNA methylation and gene expression elicited by spaceflight in
Arabidopsis thaliana. Bmc Genomics **20**10.1186/s12864-019-5554-z.

Zhou, M., Nabilsi, N.H., Wang, A., Gauthier, M.-P.L., Murray, K.O., Azari, H., Owens, W.S., Newman,
J.R.B., Pardo-Palacios, F.J., Conesa, A., Riva, A., Clanton, T.L., Reynolds, B.A., Concannon P.,
Brant, J.O., Bacher, R., Kladde, M.P. (2022). Flap-enabled next-generation capture (FENGC): precision
targeted single-molecule profiling of epigenetic heterogeneity, chromatin dynamics, and genetic variation.
BioRxiv. doi: https://doi.org/10.1101/2022.11.08.515732.