Single-molecule long-read methylation profiling reveals regional DNA methylation regulated by Elongator Complex Subunit 2 in Arabidopsis roots experiencing spaceflight

Mingqi Zhou[1], Alberto Riva[2], Marie-Pierre L. Gauthier[3], Michael P. Kladde[3], Robert J. Ferl[1,4,*], Anna-Lisa Paul[1,2,*]

[1]Department of Horticultural Sciences, University of Florida, 2550 Hull Road, Fifield Hall, Gainesville FL, 32611, USA

[2]Interdisciplinary Center for Biotechnology Research, University of Florida, 2033 Mowry Road, Gainesville FL, 32610, USA

[3]Department of Biochemistry and Molecular Biology, University of Florida, 2033 Mowry Rd, Gainesville, FL 32608, USA

[4]Office of Research, University of Florida, 1523 Union Rd, Grinter Hall, Gainesville, FL, 32611, USA

*Corresponding authors: robferl@ufl.edu (RJF), alp@ufl.edu (ALP)

Abstract

Background

The Advanced Plant Experiment-04 - Epigenetic Expression (APEX04-EpEx)

experiment onboard the International Space Station examined the spaceflight-altered

cytosine methylation in two genetic lines of *Arabidopsis thaliana*, wild-type Col-0 and the

mutant *elp2-5,* which is deficient in an epigenetic regulator Elongator Complex Subunit

2 (ELP2). Whole-genome bisulfite sequencing (WGBS) revealed distinct spaceflight

associated methylation differences, presenting the need to explore specific space-

altered methylation at single-molecule resolution to associate specific changes over

large regions of spaceflight related genes. To date, tools of multiplexed targeted DNA

methylation sequencing remain limited for plant genomes.

Results

To provide methylation data at single-molecule resolution, Flap-enabled next-generation

capture (FENGC), a novel targeted multiplexed DNA capture and enrichment technique

allowing cleavage at any specified sites, was applied to survey spaceflight-altered DNA

methylation in genic regions of interest. The FENGC capture panel contained 108

targets ranging from 509 to 704 nt within the promoter or gene body regions of gene

targets derived from spaceflight whole-genome data sets. In addition to genes with

significant changes in expression and average methylation levels between spaceflight

and ground control, targets with space-altered distributions of the proportion of

methylated cytosines per molecule were identified. Moreover, trends of co-methylation

of different cytosine contexts were exhibited in the same DNA molecules. We further

identified significant DNA methylation changes in three previously biological process-

unknown genes, and loss-of-function mutants of two of these genes (named as *EMO1*

and *EMO2* for *ELP2-regulated Methylation in Orbit 1* and *2*) showed enhanced root

growth rate.

Conclusions

FENGC simplifies and reduces the cost of multiplexed, targeted, single-molecule

profiling of methylation in plants, providing additional resolution along each DNA

molecule that is not seen in population-based short-read data such as WGBS. This

case study has revealed spaceflight-altered regional modification of cytosine

methylation occurring within single DNA molecules of cell subpopulations, which were

not identified by WGBS. The single-molecule survey by FENGC can lead to

identification of novel functional genes. The newly identified *EMO1* and *EMO2* are root

growth regulators which may be epigenetically involved in plant adaptation to

spaceflight.

Keywords

Arabidopsis, DNA capture, DNA methylation, FENGC, Single-molecule level,

Spaceflight

Background

DNA methylation can impact chromatin structure to epigenetically modulate gene

expression in eukaryotes. In plants, DNA methylation plays a substantial role in the

physiological adaptation to environmental changes and stresses [1]. Spaceflight

presents a unique physiological challenge for plants, constituting an environment that is

outside the evolutionary experience of any terrestrial organism. Plants will play a key

role in the human life support systems envisioned for spaceflight and planetary

exploration habitats [2,3], thus it is important to understand the nature of the responses

plants mount in their physiological adaptation to this novel environment. Epigenetic

change plays a crucial role in environmental adaptation, including adaptation to

spaceflight [4, 5]. The Advanced Plant Experiment-04 - Epigenetic Expression (APEX-

04) experiment onboard the International Space Station (ISS) examined the spaceflight-

altered cytosine methylation in *Arabidopsis thaliana* (Arabidopsis) using *elp2-5*, a

mutant of an epigenetic regulator, Elongator Complex Subunit 2 (ELP2), along with Col-

0, a wild-type control [4]. RNA sequencing (RNA-seq) and Whole genome bisulfite

sequencing (WGBS) revealed remodeled transcriptomes and DNA methylomes during

spaceflight, remodeling that was regulated by ELP2. The spaceflight *elp2-5* phenotype

was distinct from the space-grown wild-type plants in that root growth did not present a

typical spaceflight directionality, which orients root growth away from the germinating

seed, but rather displayed a distinct clumping of roots clustered around the seed. The

molecular phenotype exhibited distinct differences in the transcriptome and methylome

compare to the ground controls, with a majority of differentially expressed genes in *elp2-*

5 roots being also differentially methylated between spaceflight samples and ground

controls [4].

However, the short-read sequencing used in WGBS precludes the ability to phase

regional epigenetic signatures along individual molecules such as consecutive

methylated cytosines relative to transcription factor binding, nucleosome positioning,

histone incorporation and modification [1, 6]. Furthermore, detection of differentially

methylated cytosines (DmCs) by WGBS is affected by sequencing depth and false

positives due to overestimation of 5mC levels [7], and many downstream analyses are

restricted to DmCs [8, 9].

Flap-enabled next-generation capture (FENGC) is a novel method for sensitive, strand-

specific capture of long DNA sequences from user-defined target regions for

subsequent genotyping and detection of DNA modification [10]. In the single-tube

FENGC protocol, standard oligonucleotides are designed to form 5' flaps at any desired

base pair position for cost-effective, specific cleavage of target sequences by flap

endonuclease. In contrast to WGBS, targeted sequencing increases coverage depth,

thereby improving data quality and further maximizing cost savings by limiting off-target

sequencing. FENGC also reports 5mC over long, contiguous single molecules, and

therefore readily identifies distinct epigenetic configurations of extended regions that

cannot be distilled from short WGBS reads. Furthermore, because 5mC does not affect

flap endonuclease activity and target sequence capture by FENGC occurs prior to

deamination of C to U, target sequence primers are designed irrespective of DNA

methylation [10, 11]. Hence, FENGC is not subject to the primer design constraints

incumbent to bisulfite sequencing, namely, avoidance of potentially methylated sites or

insertion of degenerate bases. Therefore, FENGC offers unprecedented latitude in

targeted sequencing, especially for surveys of plant genomes that possess 5mC in

three cytosine contexts, CG, CHG, and CHH [12]. The accuracy and efficiency of

enzymatic methyl-seq, which is deployed in the methylation detection procedure of

FENGC, has been verified in plants [7]; however, FENGC has yet to be applied to

methylation analysis in plants.

Therefore, we set out to demonstrate the effective application of FENGC in plant

methylation contexts and its power in identification of functional loci. The observed

phenotypical and molecular alteration in roots of Col-0 and *elp2-5* grown on the ISS in

APEX-04 present a short-read methylation dataset for a comparison with methylation

profiles assayed by FENGC single-molecule long reads. WGBS-unidentified differential

modification of DNA methylation substrates in functionally-annotated genes as well as

unknown genes of plants grown in orbit have been detected, which led to identification

of novel regulators of root growth.

Results and Discussion

Quality of FENGC assay

In the FENGC assay, strand cleavage was directed by precise targeting of flap

endonuclease 1 (FEN1) to 5' flaps formed by sequence-specific oligonucleotides,

avoiding reliance on restriction sites or other specific motifs. The oligo panel targeted

108 regions in total, with 107 regions within the promoter or gene bodies of 86 genes of

interest derived from our APEX-04 spaceflight and related experiments. These genes

were differentially expressed in one or more spaceflight-associated environment(s) [4,

5, 13, 14]. A region of AT2G46830 with nearly no detectable 5mC in the APEX-04

WGBS experiment was included as an internal control to gauge efficient conversion and

to rule out 5mC false positives. The captured set of DNA strands ranged from 509 to

704 nt (Additional file 1: Table S1), with mean of 580 nt and median of 572 nt. The

sequence capture was performed as a single-tube multiplex reaction followed by

enzymatic methyl-PCR (oligos listed in Additional file 2: Table S2). The FENGC

products amplified from each reaction were subjected to PacBio long-read HiFi

sequencing. For APEX-04 samples, including Ground control Col-0 Roots (GCR), Flight

Col-0 Roots (FCR), Ground control *elp2-5* Roots (GER) and Flight *elp2-5* Roots (FER),

the reads from the sequenced capture libraries were consistently ~90% on-target (Figs.

1A and 1B; Additional file 3: Table S3). After filtering and combining the four biological

replicates from each sample, on average, 82% of the 108 targeted regions were

detected by at least 10 reads, with median coverage of 144, 612, 211 and 791x in four

samples, respectively (Fig. 1C; Additional file 4: Table S4). The GC content of the 108

targets, ranging from 19% to 52%, was positively correlated with the aligned reads

number (Fig. 1D; Additional file 5: Fig. S1), whereas there was a modest negative

correlation in FENGC targets with an average GC content above 50% [10]. This

indicates that both high and low GC content of targets influenced the reads yield of

FENGC products.

The efficiency of C to U conversion by EM-seq and hence quality of 5mC detection was

calculated, using all cytosines from 558 bp of the internal control region of AT2G46830

promoter (Additional file 1: Table S1; Fig. 1E). The EM-seq conversion rates of the four

samples pooled from the four biological replicates were all above 99%, on par with that

of WGBS from 97% to 99% (Fig. 1E). In this study, we analyzed single molecules from

seven target genes of interest, containing both hyper- and hypomethylated cytosines

(described below). Collectively, levels of 5mC across CG, CHG, and CHH motifs

exhibited correlation coefficients above 0.93 between any two biological replicates and

above 0.95 between FENGC and WGBS results of APEX04 samples (Fig. 1F). The high

reproducibility of FENGC targeted sequencing and strong agreement with WGBS

results at diverse regions validate the FENGC application.

This case demonstrates multiple advantages for *in vitro* targeted measurement of 5mC

in plant genomes by FENGC, including flexible target design, high sequencing depth,

scalability, outstanding on-target ratio, and low cost (Table 1). To date, targeted DNA

methylation detection in plants has predominantly relied on bisulfite sequencing of PCR

amplicons. Primer design for bisulfite sequencing requires inclusion of degenerate

bases to cover all possible methylation configurations of potential methylation sites that

cannot be avoided, which can lead to assay bias and low multiplexing capacity.

Recently, direct calling of DNA base modifications for plant sequences has extended

the methods for profiling DNA methylation in plants [15, 16]; however, these

applications require micrograms of input DNA and/or are limited to high-copy number

sequences. The increasing need to examine large panels of specific single-copy loci,

(as for differentially expressed genes of interest in an RNA-seq experiment) for

epigenetic regulation by DNA methylation can pose substantial challenges, particularly

when only nanograms of DNA are available and budgetary considerations are factors.

FENGC overcomes these limitations, and therefore provides a unique approach that

facilitates multiplexed, targeted DNA methylation analysis for plant research.

Regional methylation detection by FENGC in comparison with WGBS data

One major advantage of long-read, single-molecule profiling of DNA methylation is assessment of 5mC at every cytosine along each contiguous DNA molecule, thereby presenting a coherent regional methylation status of an entire region from a single cell. In addition, the average 5mC levels across all molecules can be derived for comparison to WGBS data. For the present study, long-read, single-molecule sequencing identified gene targets showing not only average methylation level changes between spaceflight and ground control, but also space-altered distributions of the proportion of methylated cytosines per molecule.

Combining the four biological replicates, GCR, FCR, GER and FER samples yielded 79, 93, 88 and 96 target regions, respectively, with coverage of at least 10 reads (Additional file 3: Table S3 and Additional file 4: Table S4). The distributions of the proportion of methylated cytosines per molecule in each target region were calculated. Among them, 12 targets in 12 genes showed differential methylation proportion per molecule (DmPPM) for the comparison between spaceflight and ground control in Col-0 or *elp2-5* (Additional file 1: Table S1). Meanwhile, the APEX-04 WGBS-identified DmCs (methylation difference > 0.2, FDR < 0.01) between spaceflight and ground control were examined within 108 target regions of FENGC. Col-0 and *elp2-5* showed 8/2/7 and 5/2/2 targets possessing DmPPM for CG/CHG/CHH methylation, respectively. Among these regions, only one target with DmCHH for Col-0 and two targets with DmCHG for *elp2-5* were identified by WGBS (Additional file 1: Table S1). The comparison of detection in DmPPM and DmC demonstrated that the changes of contiguous

methylation status observable in a specific subpopulation of molecules in FENGC may

be indiscernible in a population-based analysis.

Next, we present four examples associated with defense responses to stimuli and three

biological process-unknown genes regulated by ELP2 during plant growth in spaceflight

to exhibit details of space-altered methylation status that are newly uncovered by

FENGC (Additional file 1: Table S1). In each region, 93%-100% of the mapped reads

are full-length, containing all CG/CHG/CHH sites of each target region (Additional file 5:

Fig. S2). These results verify the high data quality for captured targets in the FENGC

assay of space plants.

Identification of previously unidentified space-altered DNA methylation in

defense-related gene targets

Among 12 genes exhibiting DmPPM of spaceflight vs ground control, four defense-

related genes showed space-triggered DNA methylation modification that was not

discernible in WGBS. The first three examples were examined with CG methylation, for

which DmPPM was shown in CG sites but no DmCG (methylation difference > 0.2, FDR

< 0.01) between spaceflight and ground control was detected by WGBS in these target

regions. *CML46* (AT5G39670), a gene encoding a calmodulin-like protein showing

significant transcriptional down-regulation during spaceflight in Col-0 but not *elp2-5*

plants, was assayed in a promoter region containing a MYB-like DNA binding element

(Figs. 2A and 2B). Flight and ground control Col-0 roots (FCR and GCR), but not roots

from *elp2-5* (FER and GER), showed a significant difference in the CG methylation level

in an area flanking the MYB-like element, which contained five CG sites (Fig. 2C, in

dashed frame). Consistent with these data, CG methylation proportion per molecule

(mPPM) in this target region was significantly higher in FCR than GCR (Fig. 2D). A

significant difference between GCR and GER was also observed. FENGC showed more

molecules containing consecutively methylated CG sites in FCR (12%) compared to

GCR (6.9%) in the area flanking this *cis*-regulatory element (Fig. 2E). Consistency

between the levels of gene transcription, WGBS-derived population-based methylation

status, and FENGC-derived mPPM, support the biological conclusion that ELP2 played

a negative role in expression and a positive role in promoter methylation of *CML46* in

spaceflight.

ABCB17 (AT3G28380), an ATP-binding cassette transporter gene that also showed a

significantly lower transcription level in FCR than GCR, was assayed in a gene body

region (Additional file 5: Figs. S3A and S3B). The overall patterns of averaged regional

CG methylation levels were very similar between the four samples, which did not show

significant changes between any pair of samples (Additional file 5: Fig. S3C); however,

the CG mPPM in this target region was significantly higher in FCR than in GCR

(Additional file 5: Fig. S3D). The FENGC single-molecule data showed that 3.7% of

FCR reads possessed at least 6 consecutively methylated CG sites, whereas in GCR

the percentage was 0.8% (Additional file 5: Fig. S3E). Biologically, *ABCB17* was

repressed by spaceflight and ELP2 contributed to its transcription and gene body

methylation.

Similarly, in the promoter region of *DEFL* (AT4G22217), a defensin-like gene (Additional

file 5: Figs. S4A and S4B), the patterns of averaged CG methylation levels were highly

similar and no significant alterations were detected between any two samples

(Additional file 5: Fig. S4C); however, CG mPPM was significantly higher in FER than

GER (Additional file 5: Figs. S4D and S4E). This higher methylation proportion

correlates with the lower transcription level of this gene in FER compared to GER

(Additional file 5: Fig. S4B). ELP2 enhanced expression and repressed promoter

methylation of *DEFL* in spaceflight. These examples suggest that long-read FENGC

single-molecule level methylation profiling can be more revealing of consequential

methylation changes compared with population-based methylation methods such as

WGBS, especially when alterations occur in a subpopulation of cells within a sample.

For the fourth example, a promoter fragment containing a MeJA-responsive element

was selected in *PDF1.1* (AT1G75830), a gene coding a pathogenesis-related protein

belonging to the plant defensin family and showing significant spaceflight induction in

elp2-5 (Figs. 3A and 3B). The average methylation level per site over all sequenced

molecules showed significant differences at CHG and CHH sites between GER and

GCR, as well as in CHH sites between FCR and GCR (Fig. 3C). For mPPM, significant

differences were detected in all pairwise comparisons between flight conditions and

genotypes at cytosines in all three methylation contexts (Fig. 3D). DmCs of WGBS

between GER and GCR correlated with methylation changes detected by FENGC (Fig.

3D; Additional file 1: Table S1), while between spaceflight and ground control only three

DmCGs in *elp2-5* were identified by WGBS, and the FENGC determined methylation

level changes were smaller than 0.2 at these three sites (Fig. 3C), which was in line with

the WGBS overestimation of methylation level compared with EM-seq, especially at

CHG sites [7]. The patterns of methylation proportion in DNA molecules demonstrated

that three cytosine contexts showed a trend of co-methylation in this region among cells

in the population (Additional file 5: Fig. S5). Within each molecule, methylation

proportion of CG is higher than CHG in 97.9% reads, and that of CHG is higher than

CHH in 87.5% reads (Fig. 4A). Although three types of cytosine methylation exhibited a

trend of coordination in the whole cell population, they could spread independently

within subpopulations of cells. The CG methylation proportions were above 0.57 in all

reads of GCR and above 0.28 in GER, while FCR and FER had reads with CG

methylation proportion as low as 0. These data indicate that more molecules were

hypomethylated in this region in *elp2-5* compared with Col-0 and in spaceflight

compared with ground control.

The profiling of methylation data in this example also showcases another advantage of

long-read, single-molecule FENGC assay of DNA methylation in plants, which is that all

three methylation sequence contexts (CG/CHG/CHH) can be identified and plotted for

the same molecule, thereby examining the entire methylation status of a gene region.

To determine the spatial distribution of these methylation differences, three separate but

correlated panels were plotted for CG, CHG and CHH methylation in the same DNA

molecule per row (Fig. 3E). In these plots, black areas are unmethylated regions, red

areas are methylated regions, and gray areas are transition zones between methylated

and unmethylated sites. It is visually apparent that spaceflight results in more

unmethylated regions in Col-0, with the deepest loss of methylation concentrated near

the MeJA-responsive element. It is also visually apparent that *elp2-5* shows a deeper

loss of methylation in that region. For the CG methylation context, a footprint showing

consecutive non-methylation of the first two CG sites in flanking area of the MeJA-

responsive element was identified in 8.4%, 18.0%, 39.4% and 44.0% reads in GCR,

FCR, GER and FER, respectively, quantifying the loss of CG methylation in this area.

Interestingly, 60%, 86.5%, 97.5% and 97.6% reads with consecutive non-methylation in

the second and the third CG sites in GCR, FCR, GER and FER, respectively, were also

included in this footprint and showed consecutive non-methylation of the first three CG

sites. In addition, CHG and CHH methylation proportions were significantly lower in the

footprint area in the reads containing this footprint (Fig. 4B). These results are

consistent with the previously observed high frequency of co-methylation/demethylation

in nearby cytosines in Arabidopsis [17], and demonstrate that FENGC provides further

opportunities for studying co-methylation of all three types of methylation along each

single molecule.

Four targets shown here are associated with plant defense signaling, which is largely

utilized in spaceflight adaptation of plants and the related genes can be involved in

regulation of root growth phenotypes during spaceflight [18, 19]. It has also been

reported that ELP2 modulates the epigenetic control of plant immune responses and

defense processes [20]. The newly identified space-altered methylation in these targets

is involved in ELP2-regulated defense/immunity-related pathways that may function in

adaptation of plant roots grown in orbit.

Identification of novel spaceflight-related epigenetic modification in biological

process-unknown genes

The 12 genes with DmPPM between spaceflight and ground control were annotated

using the Database for Annotation, Visualization and Integrated Discovery (DAVID,

https://david.ncifcrf.gov/). Three genes were returned without gene ontology (GO) or

functional annotation for biological process. Further investigation verified that the three

target regions within these three genes possessed significant alteration between

spaceflight and ground control in both average cytosine methylation level as well as

mPPM in Col-0 or *elp2-5*. Moreover, loss-of-function mutants of two of these genes

showed a root growth phenotype. We dubbed the two genes that could be

epigenetically involved in plant adaptation to spaceflight under regulation of ELP2,

AT1G27565 and AT5G52710, as *ELP2-regulated Methylation in Orbit* (*EMO*) *1* and *2*,

respectively.

A promoter region was captured by FENGC for *EMO1* (AT1G27565), containing an

anaerobic-responsive element (ARE) beside the transcription start site (TSS) (Fig. 5A).

The presence of the ARE is consistent with the observed induction of hypoxia-

responsive genes in spaceflight transcriptome data sets [5, 21]. *EMO1* was significantly

up-regulated by spaceflight in *elp2-5*, but not in Col-0 (Fig. 5B). Regional average

methylation level analysis detected significant differences between genotypes and

between spaceflight and ground control in Col-0 only for CG sites (Fig. 5C). The

calculation of mPPM showed significant alteration between spaceflight and ground

control in all three types of cytosine context for Col-0 and in CG sites for *elp2-5* (Fig.

5D). Visualization of three types of methylated cytosines in each single molecule

revealed footprints of methylation-free areas in areas in subpopulations of epialleles

(Fig. 5E). Significantly, more molecules in *elp2-5* lacked cytosine methylation within this

region than Col-0, and more molecules with methylation-free areas beside TSS were

identified in spaceflight samples than ground control. These results were consistent with

the observation of lower correlation of mPPM between GG and CHH in GCR than other

samples (Additional file 5: Figs. S6 and S7). After demethylation initiated by spaceflight

or mutation of ELP2 in a small proportion of cells, the co-methylation for different types

of cytosine context in those molecules had been changed accordingly. Overall, ELP2

played a negative role in *EMO1* expression and maintained cytosine methylation of the

EMO1 promoter in a subpopulation of cells in roots. Spaceflight promoted demethylation

in flanking areas of *EMO1* TSS, while the induction of *EMO1* transcription in space was

repressed by ELP2.

The *EMO2* (AT5G52710) gene belongs to copper transport protein family [22]. The

captured promoter region contains two MeJA-responsive elements (Fig. 6A). The

transcription level of *EMO2* was lower in *elp2-5* than Col-0, and lower in spaceflight than

ground control (Fig. 6B). Accordingly, the regional average methylation levels were

significantly higher in *elp2-5* than Col-0 for CG and CHG sites, which was consistent

with DmC analysis of WGBS (Fig. 6C). In contrast, CHH differential methylation levels

detected by FENGC were smaller than that of WGBS. The mPPMs were higher in

spaceflight than ground control in all three cytosine types for *elp2-5*, which explained

the space-repressed expression of *EMO2* when ELP2 was deficient (Figs. 6B and 6D).

Intriguingly, the areas of two MeJA-responsive elements both harbored a footprint of low

CG methylation, whereas CHG sites localized within these elements showed

methylation peaks, especially in *elp2-5* (Figs. 6C and 6E). When ELP2 was inactivated,

the two CG methylation footprints were largely lost. Moreover, the correlations of mPPM

between different types of cytosine context were lower in *elp2-5* than Col-0 (Additional

file 5: Figs. S8 and S9). These results indicate the role of ELP2 in the maintenance of

hypomethylated motifs and coordination of methylation for different cytosine types in

this chromatin region, which could contribute to activation of *EMO2* transcription.

The AT4G04990 gene encodes a serine/arginine repetitive matrix-like protein and the

transcripts can be transported between roots and shoots [23]. The captured promoter

region contains a TC-rich repeat that is related to defense and stress responses

(Additional file 5: Fig. S10A). The transcription level of AT4G04990 was down-regulated

by both spaceflight and mutation of ELP2 (Additional file 5: Fig. S10B). No DmC in

WGBS or significant difference of average methylation level in FENGC was identified for

this region, while DmPPMs of CG and CHH were detected between spaceflight and

ground control in Col-0 (Additional file 5: Figs. S10C and S10D). In this region, a

stronger correlation was observed between mPPMs of CHG and CHH than that of CG

and CHH (Additional file 5: Figs. S10E and S11). Correlation between mPPMs of CG

and CHG increased by spaceflight in Col-0 (Additional file 5: Fig. S11), which was

partially due to space-induced demethylation of CG sites in a subpopulation of cells

(Additional file 5: Figs. S10D, S10E and S12). The coordination of CG and non-CG

methylation might be related to the transcription repression of AT4G04990 in spaceflight

and *elp2-5*.

The features in alteration of DNA methylation at the single molecule level, such as

DmPPM and methylation footprints, would be more informative for detection of

epigenetic changes and better connect the regulation of gene expression to epigenetic

modification, compared with population-based DmC levels that lose contextual

epigenetic information of the single molecules. Therefore, FENGC data provide

additional clues for further selection of target genes of interest for downstream analysis,

especially when candidates include unknown genes.

EMO1 and *EMO2* are negative regulators of root growth

T-DNA insertion lines of the three biological process-unknown genes identified by

FENGC were characterized. Two lines of each gene tested showed disrupted gene

expression levels compared with Col-0 (Additional file 5: Fig. S13). Thus, SALK_092504

and SAIL_344_F07 were named as *emo1-1* and *emo1-2*, while SALK_140748C and

SALK_091636C were named as *emo2-1* and *emo2-2*, respectively. In a constant LED

light condition that was similar with Veggie facility onboard ISS, *emo1* and *emo2*

mutants showed significantly enhanced root growth rates compared with Col-0 (Figs.

7A-7C). The primary root length of *emo1-1*, *emo1-2*, *emo2-1*, and *emo2-2* were

significantly longer than that of Col-0 in 4-day and 7-day old seedlings. The two mutants

of AT4G04990 exhibited similar growth phenotype with wild type. Meanwhile,

treatments using 2-D clinorotation at 4 rpm or reorientation to 90˚ did not show altered

gravitropic responses in roots of any line tested, although roots of *EMO1* and *EMO2*

mutants still grew faster than Col-0 on the clinostat (Additional file 5: Figs. S14 and

S15). These results demonstrated that *EMO1* and *EMO2* are two negative regulators of

root growth, but not regulators of root gravitropism.

Remodeled plant root growth has been reported in orbital spaceflight [24]. Both

direction of root tip growth and root growth rate were impacted in Arabidopsis plants

grown on the ISS [18, 24, 25]. The primary root length of 8.5-day old seedlings for both

Col-0 and Ws were significantly shorter in flight samples than ground control. This

influence on root growth from spaceflight involves epigenetic and transcriptional

modulation [4, 13]. As a epigenetic modulator of root growth and development, ELP2

modifies DNA methylation or histone acetylation of a range of transcription factors and

other developmental regulators in roots [26]. On the ISS, the *elp2-5* mutant exhibited

severe growth retardation in roots compared with Col-0 [4]. Here, *EMO1* and *EMO2*

have been identified as two downstream targets of DNA methylation modification led by

ELP2 and/or caused by extraterritorial environment. Together with the root growth rate

phenotype in their mutants, *EMO1* and *EMO2* can play a role in root growth regulation

during space adaptation of plants, which may involve the epigenetic signaling pathways

controlled by ELP2. Moreover, although the mutants of AT4G04990 did not show root

growth alteration on the ground, further investigation will be needed to examine its

function in space.

Conclusion

The targeted long-read, single-molecule profiling enabled by FENGC reveals detailed

epigenetic modifications occurring in cell subpopulations and identifies epialleles

present at low frequencies, whereas short-read WGBS only detects alteration of the

average methylation level in cytosines. On one hand, FENGC and WGBS show

consistency in the detection of cytosine methylation levels. On the other hand, FENGC

enables single-molecule determination of methylation alteration in genes of interest

defined by users, which greatly extends interpretation beyond that of WGBS. Analysis of

DmPPM serves as critical addition to DmC data in the methylation survey. The capacity

of target cleavage at any specified sites enables wide application of FENGC, such as

studies on plant development, environmental responses, and generation of biomarkers.

In the current work, FENGC enabled a richer contextual survey of regional methylation

in genes of interest identified from primary transcriptomic and WGBS screening,

successfully identifying two novel root growth regulators through analysis of DNA

methylation alteration. In summary, FENGC provides the plant and space biology

communities with a powerful tool for the practical, routine application of multiplexed

targeted sequencing and coordinated methylation analysis at single-molecule

resolution. This FENGC screen of spaceflight-associated genes revealed candidate

genes with regulatory regions that harbored significant alterations in consecutive spans

of methylation. Further genetic analysis identified *EMO1* and *EMO2*, which could be

epigenetically modulated by ELP2 for both transcription and DNA methylation during

spaceflight. The DmPPM of *EMO1* and *EMO2* between *elp2-5* and Col-0 indicated that

ELP2-mediated cytosine modification in these two genic regions occurred in a specific

subpopulation of cells in space. These two genes are novel candidates for further

investigation of root growth regulatory signaling elicited by ELP2 in plant adaptation to

spaceflight.

Methods

Plant materials and operations of spaceflight and ground controls

The *Arabidopsis thaliana* lines Col-0 and *elp2-5*, as well as spaceflight experimental

operations, have been previously documented [4, 20]. The Advanced Plant Experiment

04 – Epigenetic Expression (APEX-04-EPEX) spaceflight experiment was launched

onto International Space Station (ISS) through the SpaceX mission CRS-10. In brief,

seeds were sterilized and sown on growth medium containing 0.5% phytagel, 0.5×

Murashige–Skoog salts, 0.5% (w/v) sucrose, and 1 × Gamborg's Vitamin Mixture, in

Petri dishes (100 mm × 15 mm; Fisher Scientific, Pittsburgh, USA), which were then

sealed with breathable tape (3M Micropore, Maplewood, USA). Plants were grown in

the Vegetable Production System (Veggie) facility on the ISS and the Veggie hardware

in the ISS environmental simulator (ISSES) chamber at Kennedy Space Center under

constant LED light of 100–135 µmol/m^{2}s Photosynthetically Active Radiation (PAR) until

11 days after germination, prior to harvest using RNAlater (Ambion, Grand Island, USA)

in Kennedy Space Center fixation tubes (KFTs). Plants from each plate were collected

into individual KFTs and stored at -80°C until delivery to the lab for analysis.

Genomic DNA Isolation and flap-enabled next-generation capture (FENGC) assay

DNA extraction was conducted using the protocol designed for RNAlater preserved

plant samples [27]. The same set of DNA samples from Col-0 and *elp2-5* root tissues

used for whole-genome bisulfite sequencing (WGBS) [4] were subjected to flap-enabled

next-generation capture (FENGC) assay. The samples included four biological

replicates of each condition: Ground control Col-0 Roots (GCR), Flight Col-0 Roots

(FCR), Ground control *elp2-5* Roots (GER) and Flight *elp2-5* Roots (FER).

FENGC oligo-1, -2 and -3 were designed using FENGC Oligonucleotide Designer

(FOLD) program (https://github.com/uf-icbr-bioinformatics/FOLD), to bind target DNA to

form 5′ flap structures that were cleaved by flap endonuclease (FEN), which enabled

ligation between target fragments and universal adaptors [10]. There were 108 targets

selected within promoter and gene body regions of 87 genes of interest derived from

previous spaceflight-associated transcriptome and DNA methylome studies [4, 5, 13,

14]. The detailed information of the 108 targets is in Additional file 1: Table S1. All oligos

used in this study are listed in Additional file 2: Table S2. All captured sequences are

from the coding strand of each gene.

FENGC experiments were performed as described by Zhou et al. [10], with

modifications to adapt to plant genome. In brief, approximately 200 ng of Arabidopsis

genomic DNA was fragmented by sonication using a Bioruptor Pico (Diagenode,

Denville, USA) for 6 cycles of 5 sec on and 30 sec off in 20 µl ddH$_2$O. The concentration

of oligo-1, -2, and -3 was 18 nM each in the master tube as recommended [10]. Flap

cleavage was performed in the presence of FENGC universal oligo 1-T (U1-T oligo),

flap oligos-1 and -2 specific for 108 targets, and thermostable FEN1 (New England

Biolabs, Ipswich, USA). Ligation was initiated by adding a stock mixture of oligos-3 for

all 108 targets, U2 oligo, and Ampligase (Lucigen, Middleton, USA) to the same tube.

DNA not ligated to U2 oligo and therefore not protected by 3' C3 Spacer was removed

by addition of exonucleases I and III (NEB, Ipswich, USA). The enriched products were

purified and then subjected to enzymatic methylation conversion according to the

manual of NEBNext Enzymatic Methyl-seq Conversion Module (NEB, Ipswich, USA).

Converted DNA was amplified for 23 cycles using U1 and U2 primers and KAPA HiFi

HotStart Uracil+ DNA polymerase (Roche, Pleasanton, USA), according to the

manufacturer's instructions. FENGC amplification products were purified and product

quality and quantity were determined using Agilent TapeStation D1000 system (Agilent,

Santa Clara, USA) and Qubit 3 Fluorometer (Thermo Fisher, Waltham, USA),

respectively.

Pacific Biosciences Sequel IIe SMRT sequencing and data analysis

The library construction and sequencing of FENGC products on the Pacific Biosciences

(PacBio) Sequel IIe SMRT platform was performed by Interdisciplinary Center for

Biotechnology Research (ICBR) at the University of Florida. The HiFi reads were

generated using 30-h movie of circular consensus (CCS) mode with filtering for ≥3

passes per single polymerase read, with default settings for other parameters. HiFi

reads mapping and visualization of DNA methylation states of single molecules were

performed using MethylMapper as previously described [28], with modifications to

accommodate plant genomes. Reads with more than three consecutive methylated

CHH sites were removed [29].

Demultiplexed CCS reads data files of PacBio Sequel IIe SMRT sequencing have been

deposited with the accession number of PRJNA857037 to BioProject database

(https://www.ncbi.nlm.nih.gov/bioproject/) in National Center for Biotechnology

Information (NCBI), and with the OSD identifier of OSD-625 (DOI: 10.26030/rp5n-0d70)

in NASA GeneLab.

Roots characterization assay

The *Arabidopsis thaliana* T-DNA insertion lines SALK_092504 (*emo1-1*),

SAIL_344_F07 (*emo1-2*), SALK_140748C (*emo2-1*), SALK_091636C (*emo2-2*),

SAIL_205_A08, and WiscDsLox441G3 were obtained from Arabidopsis Biological

Resource Center (https://abrc.osu.edu/). Seeds were sterilized, incubated in ddH$_2$O at

4°C for 3 days, and then sown on 0.5x Murashige–Skoog medium in Petri dishes as

described above. Plates were placed on vertical racks at 21°C in 24 h LED light

condition at approximately 100 µmol/m^{2}s PAR. The 4-day old were photographed. Then

plants were subjected to three treatments. The first set of plates were kept quiescent

and seedlings were grown to 7-day old and photographed. The second set of plates

were mounted to a 2-dimentional clinostat to be rotated at 4 rounds per min until plants

were 7-day old, and then photographed. The third set of plates were reoriented 90˚ to

the horizontal, and photographed at 0 h, 6 h and 24 h after reorientation. Roots were

traced and measured using JFilament plugin of ImageJ [30, 31]. Root growth

measurements were processed using the R script RootMeasurement

(https://github.com/eschultzphd/RootMeasurement/tree/master) [32]. The quantitative

real-time PCR (qPCR) assay to verify gene knock-out or knock-down was performed

using roots of 7-day old seedlings as previously described [33]. UBQ11 (AT4G05050)

was used as the internal control and primers are listed in Additional file 2: Table S2.

Availability of data and materials

Demultiplexed CCS reads data files of PacBio Sequel IIe SMRT sequencing have been

deposited with the accession number of PRJNA857037 to BioProject database

(https://www.ncbi.nlm.nih.gov/bioproject/) in National Center for Biotechnology

Information (NCBI), and with the OSD identifier of OSD-625 (DOI: 10.26030/rp5n-0d70)

in NASA GeneLab. Requests for updated versions of FENGC protocol, FOLD and

MethylMapper scripts should be directed to Michael Kladde (kladde@ufl.edu).

Abbreviations

ABCB17: ATP-binding cassette B17

ANOVA: Analysis of Variance

APEX-04-EPEX: Advanced Plant Experiment 04 – Epigenetic Expression

ARE: Anaerobic-responsive element

CCS: Circular consensus sequencing

CML46: Calmodulin like 46

CPM: Count per million mapped reads

Col-0: Columbia-0

DEFL: Defensin-like

DmC: Differentially methylated cytosine

DmPPM: Differential methylation proportion per molecule

ELP2: Elongator complex subunit 2

EMO: ELP2-regulated methylation in orbit

EM-seq: Enzymatic methylation sequencing

FCR: Flight Col-0 roots

FDR: False discovery rate

FEN1: Flap endonuclease 1

FENGC: Flap-enabled next-generation capture

FER: Flight elp2-5 roots

FOLD: FENGC oligonucleotide designer

GCR: Ground control Col-0 roots

GER: Ground control elp2-5 roots

GO: Gene ontology

HSD: Honestly significant difference

ICBR: Interdisciplinary Center for Biotechnology Research

ISS: International Space Station

ISSES: ISS environmental simulator

KFT: Kennedy Space Center fixation tube

Methyl-seq: Methylation sequencing

MYB: Myeloblastosis

mPPM: Methylation proportion per molecule

PacBio: Pacific Biosciences

PAR: Photosynthetically Active Radiation

PDF1.1: Plant defensin 1.1

qPCR: Quantitative real-time polymerase chain reaction

RNA-seq: RNA sequencing

TSS: Transcriptional start site

T-DNA: Transfer DNA

UBQ11: Ubiquitin 11

Veggie: Vegetable Production System

WGBS: Whole-genome bisulfite sequencing

5mC: 5-Methylcytosine

References

1. Zhang H, Lang Z, Zhu J-K. Dynamics and function of DNA methylation in plants. Nat
Rev Mol Cell Biol. 2018;19:489–506.

2. Wheeler RM. Plants for human life support in space: from Myers to Mars.
Gravitational and Space Biology. 2010;23:25–36.

3. Zabel P, Bamsey M, Schubert D, Tajmar M. Review and analysis of over 40 years of
space plant growth systems. Life Sciences in Space Research. 2016;10:1–16.

4. Paul A-L, Haveman N, Califar B, Ferl RJ. Epigenomic Regulators Elongator Complex
Subunit 2 and Methyltransferase 1 Differentially Condition the Spaceflight Response in
Arabidopsis. Frontiers in Plant Science. 2021;12.

5. Zhou M, Sng NJ, LeFrois CE, Paul A-L, Ferl RJ. Epigenomics in an extraterrestrial
environment: organ-specific alteration of DNA methylation and gene expression elicited
by spaceflight in Arabidopsis thaliana. BMC Genomics. 2019;20:205.

6. Edwards JR, Yarychkivska O, Boulard M, Bestor TH. DNA methylation and DNA
methyltransferases. Epigenetics Chromatin. 2017;10:23.

7. Feng S, Zhong Z, Wang M, Jacobsen SE. Efficient and accurate determination of
genome-wide DNA methylation patterns in Arabidopsis thaliana with enzymatic methyl
sequencing. Epigenetics & Chromatin. 2020;13:42.

8. Chen D-P, Lin Y-C, Fann CSJ. Methods for identifying differentially methylated
regions for sequence- and array-based data. Briefings in Functional Genomics.
2016;15:485–90.

9. Kishore K, Pelizzola M. Identification of Differentially Methylated Regions in the
Genome of Arabidopsis thaliana. In: Bemer M, Baroux C, editors. Plant Chromatin
Dynamics: Methods and Protocols. New York, NY: Springer; 2018. p. 61–9.

10. Zhou M, Nabilsi NH, Wang A, Gauthier M-PL, Murray KO, Azari H, et al. Flap-
enabled next-generation capture (FENGC): precision targeted single-molecule profiling
of epigenetic heterogeneity, chromatin dynamics, and genetic variation.
2022;:2022.11.08.515732.

11. Kladde MP, Zhou M, Nabilsi NH. Methods and kits for targeted cleavage and
enrichment of nucleic acids for high-throughput analyses of user-defined genomic
regions. 2022. WO Patent WO2022197852A3.

12. Grimanelli D, Ingouff M. DNA Methylation Readers in Plants. Journal of Molecular
Biology. 2020;432:1706–17.

13. Paul A-L, Zupanska AK, Schultz ER, Ferl RJ. Organ-specific remodeling of the
Arabidopsis transcriptome in response to spaceflight. BMC Plant Biology. 2013;13:112.

14. Paul A-L, Zhou M, Callaham JB, Reyes M, Stasiak M, Riva A, et al. Patterns of
Arabidopsis gene expression in the face of hypobaric stress. AoB PLANTS.
2017;9:plx030.

15. Ni P, Huang N, Nie F, Zhang J, Zhang Z, Wu B, et al. Genome-wide detection of
cytosine methylations in plant from Nanopore data using deep learning. Nat Commun.
2021;12:5976.

16. Mo W, Shu Y, Liu B, Long Y, Li T, Cao X, et al. Single-molecule targeted
accessibility and methylation sequencing of centromeres, telomeres and rDNAs in
Arabidopsis. Nat Plants. 2023;9:1439–50.

17. Tran H, Zhu H, Wu X, Kim G, Clarke CR, Larose H, et al. Identification of
Differentially Methylated Sites with Weak Methylation Effects. Genes. 2018;9:75.

18. Paul A-L, Sng NJ, Zupanska AK, Krishnamurthy A, Schultz ER, Ferl RJ. Genetic
dissection of the Arabidopsis spaceflight transcriptome: Are some responses
dispensable for the physiological adaptation of plants to spaceflight? PLOS ONE.
2017;12:e0180186.

19. Califar B, Sng NJ, Zupanska A, Paul A-L, Ferl RJ. Root Skewing-Associated Genes
Impact the Spaceflight Response of Arabidopsis thaliana. Frontiers in Plant Science.
2020;11.

20. Wang Y, An C, Zhang X, Yao J, Zhang Y, Sun Y, et al. The Arabidopsis elongator
complex subunit2 epigenetically regulates plant immune responses. Plant Cell.
2013;25:762–76.

21. Choi W-G, Barker RJ, Kim S-H, Swanson SJ, Gilroy S. Variation in the
transcriptome of different ecotypes of Arabidopsis thaliana reveals signatures of
oxidative stress in plant responses to spaceflight. American Journal of Botany.
2019;106:123–36.

22. de Abreu-Neto JB, Turchetto-Zolet AC, de Oliveira LFV, Bodanese Zanettini MH,
Margis-Pinheiro M. Heavy metal-associated isoprenylated plant protein (HIPP):
characterization of a family of proteins exclusive to plants. The FEBS Journal.
2013;280:1604–16.

23. Thieme CJ, Rojas-Triana M, Stecyk E, Schudoma C, Zhang W, Yang L, et al.
Endogenous Arabidopsis messenger RNAs transported to distant tissues. Nature
Plants. 2015;1:1–9.

24. Paul A-L, Amalfitano CE, Ferl RJ. Plant growth strategies are remodeled by
spaceflight. BMC Plant Biology. 2012;12:232.

25. Kwon T, Sparks JA, Nakashima J, Allen SN, Tang Y, Blancaflor EB. Transcriptional
response of Arabidopsis seedlings during spaceflight reveals peroxidase and cell wall
remodeling genes associated with root hair development. American Journal of Botany.
2015;102:21–35.

26. Jia Y, Tian H, Li H, Yu Q, Wang L, Friml J, et al. The Arabidopsis thaliana elongator
complex subunit 2 epigenetically affects root development. Journal of Experimental
Botany. 2015;66:4631–42.

27. LeFrois CE, Zhou M, Amador DM, Sng N, Paul A-L, Ferl RJ. Enabling the
Spaceflight Methylome: DNA Isolated from Plant Tissues Preserved in RNAlater Is
Suitable for Bisulfite PCR Assay of Genome Methylation. Gravitational and Space
Research. 2016;4:28–37.

28. Nabilsi NH, Deleyrolle LP, Darst RP, Riva A, Reynolds BA, Kladde MP. Multiplex
mapping of chromatin accessibility and DNA methylation within targeted single
molecules identifies epigenetic heterogeneity in neural stem cells and glioblastoma.
Genome Res. 2014;24:329–39.

29. Liu W, Gallego-Bartolomé J, Zhou Y, Zhong Z, Wang M, Wongpalee SP, et al.
Ectopic targeting of CG DNA methylation in Arabidopsis with the bacterial SssI
methyltransferase. Nat Commun. 2021;12:3130.

30. Li H, Shen T, Smith MB, Fujiwara I, Vavylonis D, Huang X. Automated actin filament
segmentation, tracking and tip elongation measurements based on open active contour
models. In: 2009 IEEE International Symposium on Biomedical Imaging: From Nano to
Macro. 2009. p. 1302–5.

31. Smith MB, Li H, Shen T, Huang X, Yusuf E, Vavylonis D. Segmentation and tracking
of cytoskeletal filaments using open active contours. Cytoskeleton. 2010;67:693–705.

32. Schultz ER, Zupanska AK, Sng NJ, Paul A-L, Ferl RJ. Skewing in Arabidopsis roots
involves disparate environmental signaling pathways. BMC Plant Biology. 2017;17:31.

33. Zhou M, Paul A-L, Ferl RJ. Data for characterization of SALK_084889, a T-DNA
insertion line of Arabidopsis thaliana. Data in Brief. 2017;13:253–8.

34. Onda Y, Takahagi K, Shimizu M, Inoue K, Mochida K. Multiplex PCR Targeted
Amplicon Sequencing (MTA-Seq): Simple, Flexible, and Versatile SNP Genotyping by
Highly Multiplexed PCR Amplicon Sequencing. Frontiers in Plant Science. 2018;9.

35. Giolai M, Paajanen P, Verweij W, Witek K, Jones JDG, Clark MD. Comparative
analysis of targeted long read sequencing approaches for characterization of a plant's
immune receptor repertoire. BMC Genomics. 2017;18:564.

36. Hill CB, Wong D, Tibbits J, Forrest K, Hayden M, Zhang X-Q, et al. Targeted
enrichment by solution-based hybrid capture to identify genetic sequence variants in
barley. Sci Data. 2019;6:12.

37. López-Girona E, Davy MW, Albert NW, Hilario E, Smart MEM, Kirk C, et al.
CRISPR-Cas9 enrichment and long read sequencing for fine mapping in plants. Plant
Methods. 2020;16:121.

Acknowledgments

696 We thank members of UF Space Plants Lab for helpful discussion and assistance in

697 experiments, as well as members of UF ICBR, NextGen DNA sequencing

698 (RRID:SCR_019152), Bioinformatics (RRID:SCR_019120) and Gene Expression and

699 Genotyping (RRID:SCR_019145) cores, for their support and services. We also thank

700 Jason O. Brant for his assistance in developing the FOLD program.

Funding

703 This work was supported by NASA grants NNX14AT24G and 80NSSC19K0130 to A-LP

704 and RJF. Development of FENGC was supported by Defense Threat Reduction Agency

705 grant HDTRA1-16-1-0048 awarded to MPK as a co-PI and NIH grant R01 CA155390

706 awarded to MPK.

Author contributions

MZ, RJF and A-LP designed this study. MZ conducted the experiments and analyzed

the data. MZ and MPK developed the FENGC methodology. AR, M-PLG and MPK

contributed to the FENGC oligo design program. MZ, MPK, RJF and A-LP wrote the

manuscript.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

A United States patent application (#18/551,004) for FENGC was filed on September

18, 2023, and is pending. MZ and MPK are two of the co-inventors.

Supplementary Information

Additional file 1: Table S1. Information of 108 capture targets.

Additional file 2: Table S2. Oligos used in this study.

Additional file 3: Table S3. Filtered HiFi reads on target and targets detected.

Additional file 4: Table S4. Number of filtered HiFi reads aligned to 108 targets.

Additional file 5: Figs. S1-S15.

Fig. S1. Correlation between numbers of filtered HiFi reads aligned to targets and the

GC content of target regions in each sample. **Fig. S2.** Percentage of full-length reads

and truncated reads mapped in seven selected target regions. **Fig. S3.** FENGC results

for *ABCB17*. **Fig. S4.** FENGC results for AT4G22217. **Fig. S5.** Correlation between

proportions of methylated cytosines in different sequence contexts within the same

molecule in the captured region of *PDF1.1*. **Fig. S6.** Correlation between proportions of

methylated cytosines in different sequence contexts within the same molecule in the

captured region of *EMO1* (AT1G27565). **Fig. S7.** Proportion of methylated CG, CHG

and CHH sites within the same molecule in the captured region of *EMO1* (AT1G27565).

Fig. S8. Correlation between proportions of methylated cytosines in different sequence

contexts within the same molecule in the captured region of *EMO2* (AT5G52710). **Fig.**

S9. Proportion of methylated CG, CHG and CHH sites within the same molecule in the

captured region of *EMO2* (AT5G52710). **Fig. S10.** FENGC results for AT4G04990. **Fig.**

S11. Correlation between proportions of methylated cytosines in different sequence

contexts within the same molecule in the captured region of AT4G04990. **Fig. S12.**

Proportion of methylated CG, CHG and CHH sites within the same molecule in the

captured region of AT4G04990. **Fig. S13.** Relative transcription levels of *EMO1*, *EMO2*

or AT4G04990 in their mutants, respectively. **Fig. S14.** Characterization of root growth

in mutants of *EMO1*, *EMO2* or AT4G04990. **Fig. S15.** Root gravitropism in mutants of

EMO1, *EMO2* or AT4G04990.

Figures

Fig. 1. Application of flap-enabled next-generation capture (FENGC) in root

samples of *Arabidopsis* experiencing spaceflight.

(A) Percentage of filtered HiFi reads on target for all 108 targets. Reads with more than

three consecutive methylated CHH sites were removed. GCR, Ground Col Roots; FCR,

Flight Col Roots; GER, Ground control *elp2-5* Roots; FER, Flight *elp2-5* Roots. (B)

Percentage of targets detected by at least 1 filtered HiFi read or at least 10 filtered HiFi

reads. (C) Filtered HiFi read coverage of 108 targets. Log transformation of reads

number was done by $\log_2$(read number +1). In boxplots, central solid line indicates

median; central dot line indicates mean; box limits indicate upper and lower quartiles;

whiskers indicate upper/lower adjacent values. Data dots are shown beside the

boxplots. (D) Correlation between numbers of total filtered HiFi reads of all samples

aligned to targets and the GC content of target regions. R-squared value and *p* value of

logarithmic regression analysis are shown. (E) Conversion rates of C to T in sequencing

results within the internal control region of 558 bp in AT2G46830 promoter. The region

coordinates are in Additional file 4: Table S4. (F) Correlation coefficients between

biological replicates of FENGC and between FENGC and WGBS for average cytosine

methylation levels in seven example targets.

Fig. 2. FENGC results for *CML46.*

(A) Genic region of *CML46*. The target region is indicated by brown rectangle (target ID: AT5G39670-1 in S1 Table). A MYB-like binding element is indicated by green brick. The transcription start site is indicated by arrow. (B) Expression levels of *CML46* in each condition. Gene expression is indicated by average count per million mapped reads (CPM) in RNA-seq reported by Paul et al. 2021. Lowercase letters indicate significant differences between samples (FDR < 0.05 in DESeq2). (C) Average CG methylation level within the target region. The relative position of MYB-like binding element shown in (A) is indicated by green brick. The region indicated by dashed frame shows significant differences of CG methylation level between FCR and GCR (one-way ANOVA with post-hoc Tukey HSD test; *, $p < 0.05$). (D) Distribution of proportion of methylated CG sites per molecule in filtered, full-length HiFi reads aligned to the target region. The box indicates upper/lower quartiles and whisker indicates upper/lower adjacent values. The data points of mPPM are represented by dots. The line and "x" in the box indicate median and mean, respectively. The significant differences of methylation proportion are shown (Kruskal-Wallis test; *, $p < 0.05$; **, $p < 0.01$). (E) Heatmaps showing CG methylation of single molecules. Red indicates two or more consecutively methylated cytosines, black indicates two or more consecutively unmethylated cytosines, and gray indicates transitions between methylated and unmethylated cytosines. The reads from the four biological replicates (Additional file 4: Table S4) were combined to generate heatmaps for each sample. The relative position of MYB-like binding element shown in (A) and (C) is indicated by green brick. The region used for the CG methylation level comparison in (C) is indicated by dashed frame. Approximately 6.9% of reads in GCR

793   and 12% of reads in FCR containing at least two consecutively methylated CG sites in

794   both areas flanking the MYB-like binding element are labeled by blue bricks. One-way

795   ANOVA with post-hoc Tukey HSD test was performed for pairwise comparisons of

796   percentage of reads with this signature. FER vs FCR, $p < 0.01$.

Fig. 3. FENGC results for *PDF1.1.*

(A) Genic region of *PDF1.1*. The target region is indicated by brown rectangle (target ID: AT1G75830-1 in Additional file 1: Table S1). A MeJA-responsive element is indicated by green brick. The transcription start site is indicated by the arrow. (B) Expression levels of *PDF1.1* in each condition. (C) Average CG, CHG or CHH methylation levels over all sequenced molecules of the target region. The relative position of MeJA-responsive element shown in (A) is indicated by green brick. The significant differences of methylation levels between samples for CHG and CHH sites in the target region are shown. One-way ANOVA with post-hoc Tukey HSD test; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. Black arrows indicate DmCs detected in WGBS using the threshold of methylation difference > 0.2 and FDR < 0.01. Arrows of solid line and dot line indicate DmCs of GER vs GCR and FER vs GER, respectively. (D) Distribution of proportion of methylated CG, CHG or CHH sites per molecule in filtered full-length HiFi reads aligned to the target region. The significant differences of methylation proportion are shown (Kruskal-Wallis test; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$). (E) Heatmaps showing the cytosine methylation in single molecules. Reads from the four biological replicates (Additional file 4: Table S4) were combined to generate heatmaps for each sample. If the number of combined reads exceeded 1,000, 1,000 reads were randomly selected and plotted. The three heatmap panels show the patterns of methylation at CG, CHG, and CHH sites (vertical white lines) as indicated at bottom along the same DNA molecule in each row. The color legend is the same as that in Fig. 2E. Reads were hierarchically clustered using Kendall's Tau based on CG methylation. The relative position of MeJA-responsive element shown in (A) is indicated by green brick. A

footprint (blue ovals) with consecutive protection against methylation of the first 2 CG

sites in the area flanking the MeJA-responsive element is present in 8.4%, 18.0%,

39.4% and 44.0% molecules in GCR, FCR, GER and FER, respectively. One-way

ANOVA with post-hoc Tukey HSD test was performed for pairwise comparisons of

percentage of reads with this footprint. FCR vs GCR, $p < 0.05$; GER vs GCR, $p < 0.001$;

FER vs FCR, $p < 0.001$.

Fig. 4. DNA methylation within same molecules in the target region of *PDF1.1.*

(A) Proportion of methylated CG, CHG and CHH sites within the same molecule in the

captured region of AT1G75830. There are 120, 1009, 345 and 1310 full-length reads

covering all CG, CHG and CHH sites in this region (target ID: AT1G75830-1 in

Additional file 1: Table S1), which are mapped for GCR, FCR, GER and FER samples,

respectively. Ochre, gold and dark green dots indicate CG, CHG and CHH methylation

proportions, respectively, in the same column, each representing one read. (B)

Comparisons of methylation proportions per molecule in reads with a footprint and in all

reads in the captured region of AT1G75830. CHG and CHH methylation in GCR, FCR,

GER and FER samples are plotted, respectively. The footprint is defined as the first 2

CG sites in this target region that are both unmethylated. The significant differences of

methylation proportion are shown (Kruskal-Wallis test; *, $p < 0.05$; **, $p < 0.01$; ***, $p <$

0.001).

Full-length reads mapped to captured region AT1G75830-1

Fig. 5. FENGC results for *EMO1*.

(A) Genic region of *EMO1*. The target region is indicated by brown rectangle (target ID: AT1G27565-1 in Additional file 1: Table S1). An anaerobic-responsive element (ARE) is indicated by green brick. The transcription start site (TSS) is indicated by arrow. (B) Expression levels of *EMO1* in each condition. (C) Average CG, CHG or CHH methylation levels in cytosines within the target region. The relative position of ARE and TSS shown in (A) are indicated by green brick and arrow, respectively. The significant differences of CG methylation level between samples in the target region are shown, in which the first CG site is omitted. One-way ANOVA with post-hoc Tukey HSD test; *, $p <$ 0.05; ***, $p < 0.001$. Black arrows indicate DmCs of FER vs GER detected in WGBS using the threshold of methylation difference > 0.2 and FDR < 0.01. (D) Distribution of proportion of methylated CG, CHG or CHH sites per molecule in filtered full-length HiFi reads aligned to the target region. The significant differences of methylation proportion are shown (Kruskal-Wallis test; *, $p < 0.05$). (E) Heatmaps showing the cytosine methylation in single molecules. The heatmap panels and color legend are the same as that in Fig. 3E. Reads were hierarchically clustered using Euclidean distance based on CG methylation. The percentages of reads without methylated CG sites and reads with at least 9 consecutive unmethylated CG sites at the 3' end are shown on the left and right side of the CG panel, respectively. One-way ANOVA with post-hoc Tukey HSD test was performed for pairwise comparisons. For percentages of reads without methylated CG sites, FER vs GER, $p < 0.05$; GER vs GCR, $p < 0.001$; FER vs FCR, $p < 0.01$. For percentages of reads with at least 9 consecutive unmethylated CG sites at the 3' end, FCR vs GCR, $p < 0.01$; GER vs GCR, $p < 0.01$.

Fig. 6. FENGC results for *EMO2*.

(A) Genic region of *EMO2*. The target region is indicated by brown rectangle (target ID:

AT5G52710 in Additional file 1: Table S1). Two MeJA-responsive elements are

indicated by green bricks. The transcription start site (TSS) is indicated by arrow. (B)

Expression levels of *EMO2* in each condition. (C) Average CG, CHG or CHH

methylation levels in cytosines within the target region. The relative position of MeJA-

responsive elements shown in (A) are indicated by green bricks. The significant

differences of CG and CHG methylation levels between samples in the target region are

shown. One-way ANOVA with post-hoc Tukey HSD test; **, $p < 0.01$; ***, $p < 0.001$.

Black arrows indicate the DmCs detected in WGBS using the threshold of methylation

difference > 0.2 and FDR < 0.01. Arrows with solid line, triangle, round end, and dot line

indicate DmCs of GER vs GCR, FER vs FCR, FCR vs GCR, and FER vs GER,

respectively. (D) Distribution of proportion of methylated CG, CHG or CHH sites per

molecule in filtered full-length HiFi reads aligned to the target region. The significant

differences of methylation proportion are shown (Kruskal-Wallis test; ***, $p < 0.001$). (E)

Heatmaps showing the cytosine methylation in single molecules. The reads clustering,

heatmap panels and color legend are the same as that in Fig. 5E.

Fig. 7. Enhanced root growth in mutants of *EMO1* or *EMO2*.

(A) 7-day old seedlings of Col-0, *emo1-1*, *emo1-2*, *emo2-1*, and *emo2-2*. Scale bars indicate 10 mm. (B) Primary root length of 4-day old and 7-day old seedlings for each line. In violin/boxplots, central solid line indicates median; central dot line indicates mean; box limits indicate upper and lower quartiles; whiskers indicate upper/lower adjacent values. Kruskal-Wallis test; *, $p < 0.05$; ***, $p < 0.001$. (C) Average root growth rate (mm per day) from day-0 to day-4, and from day-4 to day-7. Student's t test, *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. Data ± SD, n = 55, 38, 33, 36, 34, 41, 38, 33, 36, 34 from left to right.

894 **Table 1. Comparison of multiplexed, targeted DNA sequencing methods in plants.**

895 Light gray indicates advantages and dark gray indicates disadvantages.

Method	Custom design flexibility	Read length	Example of sequencing depth	Example of on-target read % in plant genome	Example of target number	Cost	Adaptation in DNA methylation detection
Multiplexed PCR	limited by low complexity of sequences and potentially methylated sites	Short (e.g., < 200 bp) [34]	1,000x [34]	50.9 % [34]	443 markers [34]	low	no
Probe hybridization-based	limited by commercial panels and costly design	short/long (e.g., 3.5 kb) [35]	55x [35]	52.9% [36]	190 genes [36]	high due to modified probes	no
CRISPR-based	limited by PAM sites and cleavage efficiency	short/long (e.g. 8 kb) [37]	~200x [37]	3.04% [37]	1 gene [37]	high due to RNA oligos and specific proteins	no
FENGC	cleavage for any specified sites	short/long (e.g., > 500 bp)	10-5,000x	~90%	102 genic regions	low	long-read EM-seq