# Energy conservation by collective movement in schooling fish

Yangfan Zhang[1]*, George V. Lauder[1],

[1]Department of Organismic and Evolutionary Biology, Harvard University, 26 Oxford St, Cambridge, Massachusetts, 02138, USA

*Corresponding author. Email: yangfan_zhang@fas.harvard.edu

**Short title:** Energy conservation by schooling fish

**Abstract**

Many animals moving through fluids exhibit highly coordinated group movement that is thought to reduce the cost of locomotion. However, direct energetic measurements demonstrating the energy-saving benefits of fluid-mediated collective movements remain elusive. By characterizing both aerobic and anaerobic metabolic energy contributions in schools of giant danio (*Devario aequipinnatus*), we discovered that fish schools have a concave upward shaped metabolism–speed curve, with a minimum metabolic cost at ~1 body length s$^{-1}$. We demonstrate that fish schools reduce total energy expenditure (TEE) per tail beat by up to 56% compared to solitary fish. When reaching their maximum sustained swimming speed, fish swimming in schools had a 44% higher maximum aerobic performance and used 65% less non-aerobic energy compared to solitary individuals, which lowered the TEE and total cost of transport by up to 53%, near the lowest recorded for any aquatic organism. Fish in schools also recovered from exercise 43% faster than solitary fish. The non-aerobic energetic savings that occur when fish in schools actively swim at high speed can considerably improve both peak and repeated performance which is likely to be beneficial for evading predators. These energetic savings may underlie the prevalence of coordinated group locomotion in fishes.

**One-Sentence Summary:** Fish schools showed a *U*-shaped metabolism-speed curve and reduced the energy use per tail beat up to 56% at high swimming speeds compared to solitary fish.

## Introduction

Newton's laws of motion underpin animal locomotion, from fine-scale maneuvers to long-distance migrations (*1*). As speed increases or during migration involving sustained movement over long distances, animals often move in coordinated groups: *e.g.*, migratory birds in V-formation (*2*), ducklings swimming in formation (*3*), cyclists in a peloton (*4*), elite marathon runners (*5*), and fish schools (*6*). To overcome gravitational or fluid dynamic resistance to forward motion, animals use both aerobic and glycolytic metabolism (oxidative and substrate-level phosphorylation at the cellular level) to generate energy and sustain movement in the air, on land, and in the water. In water, a fluid that is 50 times more viscous than air and contains much less $O_2$ per Kg than air, the need for aquatic animals to reduce fluid dynamic drag for energy conservation is even greater than for aerial or terrestrial locomotion. Hence, we use fish schooling behaviour as a model system to explore whether the fluid dynamics of collective movement (active and directional movement of animal groups along a common trajectory (*7*)) can enable energetic savings compared to locomotion by a solitary individual. Fish schooling behaviours are some of the most prominent social and group activities exhibited by vertebrates.

Fish may school for many reasons, including foraging, reproduction, and migration, and fish schools often exhibit high-speed movement during evasion from predators. Hence, natural selection can put pressure on the formation of schools and individual interactions within a school. Besides 'safety in numbers' where individuals in a large school have a lower probability of becoming prey, can there be more tangible energetic benefits of high-speed swimming as a school? Since fluid drag scales as velocity squared (*8*), movement at higher speeds places a premium on mechanisms to conserve energy. Fast and unsustainable swimming is fueled by aerobic metabolism with a major contribution from anaerobic glycolysis (*9*) (*10*). During high-speed swimming aerobic capacity is maximized (*10*), and energy use is more likely to compete

with other activities than in low-speed swimming which only occupies a small portion of aerobic

capacity. The direct consequences of unsustainably engaging anaerobic glycolysis include both

fatigue and metabolic perturbation (such as changing blood acidity) that need time to recover

after bouts of intense locomotion. Animals become vulnerable during recovery because of their

hindered ability to repeat peak locomotor performance in the presence of predators.

Another factor of importance in the experimental evaluation of the energetic cost of

collective locomotion in fishes is that endurance and efficiency are often key to achieving

lifetime fitness, especially in fish species that undergo significant migrations. Fish groups usually

undergo long-distance migrations at slow speeds ranging from 0.25–1.0 BL s$^{-1}$ as recorded by

tags on migrating fish (Fig. S1). The common migratory speed of ~1 BL s$^{-1}$ for marine and

anadromous fish could be related to the minimum aerobic cost of transport (*11*). However, there

is currently no direct measurement of the absolute cost of teleost fish locomotion demonstrating

a minimum group metabolic rate ($\dot{M}O_2$) at an optimal speed ($U_{opt}$). Fish schools in nature swim

over a wide range of speeds (Fig. S1) but no study has directly characterized the total energy

expenditure (TEE), and accounted for the potentially substantial anaerobic metabolic costs

involved in higher speed locomotion. As a result, the swimming TEE performance curve of fish

schools from the minimum swimming speed to the critical swimming speed ($U_{crit}$) has not yet

been measured.

More broadly, despite the widespread notion that collective motion saves energy (*6*), very

few studies have directly measured the energetic cost of collective movement compared to the

cost of movement by solitary individuals. In the canonical case of V-formation flight in birds,

energetic saving is exclusively inferred from indirect measurements, *e.g.,* heart rate (*2*), and

flapping frequency and phase (*12*) (Table 1).  There are no direct measurements of metabolic

energy consumption of collective movement in birds, and indeed this would be extremely

challenging to accomplish. Although some energetic aspects of schooling fishes have been

studied (*13*) (*14*) (*15*) (*16*) (*17*), these studies have analyzed a limited range of speeds and

focused on aerobic metabolism (*see* Table 1). In fact, several field and laboratory kinematic

studies suggest that both bird flocks and fish schools do not always conserve aerobic energy (*17*)

(*18*) (*19*) and indicate that moving in a group can actually involve *increased* costs. Without

directly measuring the energetic cost of movement, inferences solely based on kinematics do not

include complex group interactions, and the interaction between collective behaviour and fluid

dynamics. Consequently, we do not yet have a holistic and mechanistic view of where and how

the collective movement by fish might conserve energy.

For swimming fish, fluid dynamic experiments have shown how collective movement

can improve swimming efficiency due to interactions among deforming bodies and through

interactions between moving animals and the local fluid environment (these energy-saving

mechanisms are summarized in Fig. 1). Our current understanding of energy saving mechanisms

during collective locomotion in fishes is largely based on computational fluid dynamic models

with a few analyses using robotic systems (*6*) (*20*) (*21*) (*22*) (*23*) at low speed, but the link

between such models and fish metabolic energy saving over a wide speed range is not yet

established. We expect that the need to conserve energy in animals moving against air or water

resistance should be greater at higher speeds. We hypothesize that fish in schools can reduce the

total cost of high-speed locomotion relative to solitary movement, leading to reduced non-

aerobic energetic costs and time of recovery for schooling fish swimming at speeds occupying

the majority (> 50%) of their aerobic capacity. Our secondary objective is to determine the

differences (if any) between the swimming performance curves (metabolism versus speed) of

solitary fish and fish schools.

To evaluate these hypotheses, we directly measured both the aerobic and non-aerobic

energy used by schooling fishes over a wide range of water velocities (0.3–8.0 body lengths s$^{-1}$;

BL s$^{-1}$), and then compared the swimming performance curve to that of solitary fish. We

equipped a high-resolution (*see* Methods for our approach to enhancing the signal-to-noise ratio

in our respirometer) swim-tunnel respirometer with two orthogonal high-speed cameras to

quantify three-dimensional (3-D) fish kinematics. This enabled us to simultaneously measure

energetics, 3-D dynamics of fish schools (n=5 replicate schools), and the kinematics of solitary

fish (n=5) of a model species, giant danio (*Devario aequipinnatus*), that exhibits an active

directional group swimming from near still water to maximum sustained speeds (equivalent to a

Reynolds number range of $6.4 \cdot 10^3$ to $1.8 \cdot 10^5$; Fig. S2), using a $U_{crit}$ test.

**Results**

We discovered that both solitary fish and fish schools have a concave upward shaped

aerobic metabolic rate ($\dot{M}O_2$)-speed curve over the lower portion of their speed range (0.3–3 BL

s$^{-1}$, Fig. 2C, D). The aerobic locomotor cost at 0.3–3 BL s$^{-1}$ showed no statistical difference

between solitary fish and fish schools ($F_{1,80} \leq 2.40$, $p \geq 0.13$). Fish schools swimming at ~1.0 BL

s$^{-1}$ consume less energy than at slower speeds ($F_{9,40} = 24.7$, $p \leq 0.0007$), while swimming at 3 BL

s$^{-1}$ consumes a similar amount of energy to maintain position at a water velocity of 0.3 BL s$^{-1}$ ($p$

$= 0.85$; Fig. 2D). Danio schools have a minimum aerobic cost ($\dot{M}O_{2min} = 212.9$ mg O$_2$ kg$^{-1}$ h$^{-1}$) of

1.25 BL s$^{-1}$. $\dot{M}O_{2min}$ at this optimal speed ($U_{opt}$) was *lower* than both the $\dot{M}O_2$ of aggregating

behaviours (328.5 mg O$_2$ kg$^{-1}$ h$^{-1}$) exhibited before $U_{crit}$ test (Table S1, Fig. S3) and for the group

at rest in still water $\dot{M}O_2$ (267.9 mg O$_2$ kg$^{-1}$ h$^{-1}$) (by 35% and 21% respectively; $F_{2,12} = 16.7$, $p \leq$

0.02, Fig. 2A, B). Swimming at $U_{opt}$ reduces the energetic cost below that of swimming at either

slower or higher speeds. Indeed, the Strouhal number of fish schools decreased from 2.1 at the

lowest speed to 0.3 at the energetic minimum $U_{opt}$, and then increased to 0.4 at the higher $U_{crit}$.

We also characterized the kinematic features that result in elevated $\dot{M}O_2$ at low water

velocities. Danio maintaining body position in low-velocity aggregations had a higher tail beat

frequency ($f_{TB}$) than solitary fish ($F_{1,80} \geq 9.8$, $p \leq 0.002$). However, at $U_{opt}$ and the highest speed

of active directional schooling, fish in schools had a lower $f_{TB}$ ($F_{1,80} \geq 4.6$, $p \leq 0.035$) than

solitary fish (Fig. 4B). The 3-D angular heading of schooling fish transitioned from

omnidirectional to pointing against water flow when water velocity is above 0.75 BL s$^{-1}$, and

individual fish orient into the flow starting at 0.3 BL s$^{-1}$ (Fig. 4C). Although the body angle and

turning frequency of solitary fish and schools decreased with water velocity, schooling fish had a

higher turning frequency ($F_{1,80} \geq 15.6$, $p < 0.002$) below 0.75 BL s$^{-1}$ than solitary fish (Fig.

4D,E).

We discovered that, across the entire 0.3–8 BL s$^{-1}$ range, $\dot{M}O_2$-speed curves of fish

schools are concave upward shaped (reached 1053.5 mg $O_2$ kg$^{-1}$ h$^{-1}$ at 8 BL s$^{-1}$) whereas solitary

fish showed an upward concave curve that reaching a plateau of 760.2 mg $O_2$ kg$^{-1}$ h$^{-1}$ (~10% CV

in 6–8 BL s$^{-1}$, Fig. 2C), demonstrating that schooling dynamics results in a 44% higher

maximum aerobic performance ($F_{1,80} = 30.0$, $p < 0.001$). This increased maximum aerobic

performance translated to a lower use of aerobic capacity compared to solitary fish over 0.3–8

BL s$^{-1}$ (*see* the discussion for how fish swimming in school can improve aerobic performance).

Collectively, fish schools used a 36% lower proportion of their aerobic scope than solitary fish

(Wilcoxon test: $p = 0.0002$, Fig. 2F). Fish schools used a 38% lower proportion of aerobic scope

(34 vs. 55%) at 4 BL s$^{-1}$ (50% $U_{crit}$ & > 50% aerobic scope onward), and consistently used ~25%

lower proportion of their aerobic scope at 6 and 7 BL s$^{-1}$ than solitary fish ($F_{1,103} \geq 4.8$, $p \leq 0.03$,

Fig. 2E).

Given that fish schools had a higher maximum aerobic performance, generally used a lower proportion of their aerobic scope, and individuals within the schools have a 14% lower $f_{TB}$ than the solitary fish at 8 BL s$^{-1}$ (11.5 vs. 13.3 Hz, $F_{1,80} = 15.1$, $p < 0.001$), we predicted that fish schools, compared with solitary fish, use less anaerobic energy to supplement aerobic energy for the high-speed movement approaching their aerobic limit. Hence, we measured post-exercise O$_2$ utilization, a majority of which is used to restore high-energy phosphate storage and glycolytically induced metabolic perturbations (EPOC: Excess Post-exercise O$_2$ Consumption). We discovered that fish schools have a 65% lower EPOC (0.69 vs. 1.95 mg O$_2$; $t_8 = 4.5$, $p = 0.0021$), and recover 43% faster than solitary fish (8 vs. 14 h; $t_8 = 2.8$, $p = 0.025$; Fig. 2G, H).

To estimate the relative proportions of aerobic and anaerobic energy contributions for locomotion at each swimming speed, we modeled EPOC in addition to $\dot{M}O_2$, the aerobic cost (> 50% $U_{crit}$ & aerobic scope, Fig. 3A) (Table S2). We also estimated the total O$_2$ cost during the entire swimming process for each school and individual and calculated total energy expenditure (TEE) (10). The TEE of fish schools was 38–53% lower than that of the solitary fish between 5–8 BL s$^{-1}$ ($F_{1,103} \geq 7.4$, $p \leq 0.008$; Fig. 3D). TEE of fish schools was only 42–143 % higher than the aerobic metabolic rate at 5–8 BL s$^{-1}$ ($F_{1,96} \geq 3.5$, $p \leq 0.001$) and anaerobic metabolic energy only accounted for 29–58 % of the TEE depending on speed (Fig. 3B). In contrast, TEE of solitary fish was 131–465 % higher than the aerobic metabolic rate between 5–8 BL s$^{-1}$ ($F_{1,112} \geq 10.7$, $p \leq 0.001$), where anaerobic energy accounted for 62–81% of the TEE depending on speed (Fig. 3C).

Schooling dynamics reduced the total cost (aerobic plus anaerobic) of transport (TCOT) by an average of 43% compared to swimming alone ($F_{1,103} = 6.9$, $p = 0.01$), and most of this energy conservation happens at higher speeds when fish approach their aerobic limit. Schooling dynamics in danio enables an extremely shallow rate of increase in TCOT with speed compared

to that of an individual (Fig. 3E). The TCOT of solitary fish increased by 490% (6.5 kJ km$^{-1}$ kg$^{-1}$) at 8 BL s$^{-1}$, whereas the schooling TCOT increased by only 200% (3.6 kJ km$^{-1}$ kg$^{-1}$) at 8 BL s$^{-1}$. Therefore, individual fish form a more energy-efficient biological entity when they collectively move as a school.

To answer the question of how schooling dynamics reduces TEE, we combine video analysis of fish tail beat kinematics with simultaneous aerobic and anaerobic measurements to compute energy expended *per tail beat* (TEE•beat$^{-1}$), and compared values for fish in schools to those for solitary fish. Schooling fish reduced TEE•beat$^{-1}$ by 30–56% at higher speeds compared to solitary fish ($F_{1,81} \geq 7.3$, $p \leq 0.008$), a substantial reduction in TEE•beat$^{-1}$ consumed both by the school, and by individual fish within a school (Fig. 4A). Notably, the energetic benefits during active directional schooling occur when fish schools become more streamlined, as the length of fish school increased with speed and plateaued beyond 2 BL s$^{-1}$ (Fig. 4F), while the 3-D distance among individuals stayed relatively constant at ~1.2 BL (Fig 4G). These results directly demonstrate schooling dynamics benefits the swimming kinematics of individual fish within the school and results in a net outcome of up to 53% TEE reduction in fish schools compared to solitary fish.

## Discussion

Simultaneous characterization of energetics and kinematics enables an integrated understanding of both the physiology and physics of fish schooling behaviour. Hydrodynamic models, kinematic measurements, and robotic analyses of fish schools indicate that the cost of swimming can be reduced when fish swim beside neighbouring fish (*20*), behind or in front of another fish (*24*), or behind leading individuals (*6*) (Fig. 1). We have demonstrated here that fish schools downshifted an overall concave upward swimming energetics performance curve at

higher speeds by ~43%. Energy use as swimming speed increases changes in a non-linear

manner, with higher energy use at the lowest speeds (Fig. 2D), and the performance curve below

3.0 BL s$^{-1}$ is *U*-shaped. Energy saving occurs through a substantial reduction in the non-aerobic

energy contribution when fish approach aerobic limits. One of the key ecological benefits of the

reduced use of glycolysis is a faster recovery time from fatigue (Fig. 2H). This would enable fish

schools in nature to repeat high-performance movements. Considering that high-speed

maneuvers are extremely common in predatory evasion (*25*) food-searching group motion (*26*),

and in fish schools in the ocean (Fig. S1), energy savings by fish schooling at high speed and the

faster recovery that followed could have a considerable impact on lifetime fitness (*27*) (*28*).

Moreover, considerable energy savings occur even though we observed that individual danio

regularly change positions within the school and do not maintain stable inter-individual

locations. We expect that this is because individuals within the school can use multiple

hydrodynamic mechanisms to save energy depending on their location (Fig. 1). We propose that

energetic savings by collective movement may not require either fixed positional arrangements

among individuals or specific kinematic features (such as tail beat synchronization) compared to

solitary locomotion. Based on these results, we regard fish schooling as a highly robust

behaviour that provides considerable energetic benefits that are not necessarily sensitive to

specific fish locations or movements. Future studies are necessary to decipher each of the

specific hydrodynamic mechanisms for energy saving.

**Schooling dynamics enhances aerobic performance and reduces non-aerobic energy use**

We discovered that a significant amount of energy conservation for active directional

swimming occurs at speeds above 3 BL s$^{-1}$. In nature, fish schools routinely exhibit active

directional collective locomotion above ~6 BL s$^{-1}$ (Fig. S1, Table S3) (*29*), a speed that engages

anaerobic glycolysis. Yet, there are no previous measurements of the anaerobic cost of schooling in fish. Hence, important and previously unrecognized benefits of active directional collective locomotion are (1) increased aerobic performance, (2) a reduced use of aerobic capacity and anaerobic energy, and (3) a resultant faster recovery from the associated metabolic perturbations and costs of swimming at higher speeds. When animals are approaching their maximum metabolic rate, the highest attainable rate of $O_2$ uptake limits aerobic performance during the unsteady state of high-speed locomotion and anaerobic glycolysis is engaged to support the maximum metabolic demand.

Fish swimming at higher speeds (> 50% $U_{crit}$) routinely exhibit a burst-&-glide gait mode of locomotion (*30*, *31*). Repeated bursting is substantially fueled by anaerobic glycolysis and the gliding phase enables peak $\dot{M}O_2$ (*32*) to replenish the venous $O_2$ content (*33*). Fish within a school are known to increase gliding time and decrease burst time by 19% when they are trailing the leading fish (*34*), *e.g,* a lower $f_{TB}$ of fish schools at 8 BL s$^{-1}$ is related to the extended gliding time (Fig. 4B). We reason that the higher proportion of time spent gliding likely enables more bouts of peak $\dot{M}O_2$ and enhances the maximum aerobic performance (*32*) (*33*). This enhanced maximum aerobic performance increases the metabolic ceiling and enables fish to use a lower proportion of their aerobic scope for locomotion (Fig. 2E). Physiological studies of exercise metabolism and locomotion suggest that a lower proportional use of aerobic scope during movement relates to a reduced accumulation of anaerobic end-products (*9*). Collectively, our results showed that the increased aerobic performance (Fig. 2C) and the reduced use of the aerobic capacity (Fig. 2E) for swimming above 50% $U_{crit}$ in fish schools likely plays a key role in reducing the use of glycolysis and the accumulation of lactate (*e.g.* EPOC, Fig. 2G) for high-speed swimming (*10*, *31*).

We present direct evidence of substantial non-aerobic energy saving by demonstrating the 65% lower EPOC (Fig. 2G) used by fish that swam in schools. Anaerobic glycolysis is crucial in permitting continued movement when aerobic limits are reached at high swimming speeds (35). Often vertebrates at higher speeds use more fast twitch fibres that generate high-frequency contractile force in part through anaerobic glycolysis (36) (37). The higher EPOC for single fish is unlikely to have been confounded by any possible stress effect of swimming as a solitary fish. Stressed solitary fish would have elevated aerobic metabolic rates in low water velocity, which we show is not the case for our experiments (*see* Fig. 2D). Indeed, to mitigate the stress response, we acclimated fish to the solitary conditions and habituated fish to the respirometer for a quiescent day before the experiment (*see* more details in experimental animal and protocols). When vertebrates move at high speed, the typical 'stress' neurotransmitters (*e.g.* adrenaline, catecholamines & cortisol) usually increase (38) (39), regardless of whether vertebrates move as an individual or in a group. These endocrinological responses underpin the physiological mechanisms that enable higher aerobic performance (*e.g.* rising heart rate) (40) (41). Moreover, after the highest swimming speeds immediately followed by an active recovery process, fish likely prioritize the recovery of the essential functions (as the levels of the neurotransmitters decline fairly rapidly) for the potential of repeated locomotion performance (speedy recovery allows fish to restore their swimming performance) (31) (42) (43).

To present a complete energetic profile showing the total energy expenditure (TEE) of locomotion for fish schooling, we model the non-aerobic cost (EPOC) on top of the aerobic swimming performance curve. The model is rooted in the use of aerobic and glycolytic energy when fish swim above 50% $U_{crit}$ (30) and a commonly observed inflection point of faster anaerobic end-product accumulation is at ~50% aerobic scope (9) (*see* Methods for detailed physiological bases and criteria for EPOC modeling). As a result, we observed a TEE being ~2.5

fold higher than the aerobic metabolic energy expenditure which agrees with the theoretical

estimates based on individual yearling sockeye salmon (*44*). This model elucidates how aerobic

and anaerobic metabolic energy constitute TEE, fuel muscles, generate locomotor thrust and

overcome hydrodynamic drag during swimming.

We also demonstrated that the need for energy saving in fish schools at lower speeds (< 3

BL s$^{-1}$ or 38% $U_{crit}$ Fig. 2E, 3D, E) is not as crucial as at higher speeds where we demonstrated

substantial energy savings in the schools compared to solitary individuals. Fluid drag is

exponentially less at lower speeds compared to higher speeds. Fish predominately use aerobic

metabolism to support low-speed steady swimming for prolonged periods (*10*) which does not

require lengthy recovery as swimming at higher speeds where glycolysis contributes to the

energetic demands. The costs of low-speed swimming are often less than < 20% of aerobic

capacity (Fig. 2E), which leaves the majority of aerobic capacity (>50%) for other activities.

Therefore, the benefits of reducing the energetic cost of locomotion are likely not major factors

underlying behaviours such as low-speed milling (*45*) and aggregation (*46*) in fishes, where other

ecological drivers such as feeding, predator avoidance, and reproduction likely dominate the

fitness landscape.

**Schooling dynamics and energy conservation**

Our 3D kinematic analyses shed light on the complex interactions between schooling and

hydrodynamics that enable energy saving by fish collective movement. One of the key possible

mechanisms is local interactions among individuals, where the kinematics of individual fish

respond to the wakes shed by neighbours (*47*). Kinematic and simulation studies indicate that 2–

3 coordinated fish can save energy (*48*) (*20*) through local interactions. We directly demonstrate

here that eight coordinated fish can save energy, and recent simulations show that schooling

benefits can extend to at least 23 fish (*21*). Although these studies appear to suggest that the

energetic benefits of schooling are scalable, we are still some ways from proving the exact

mechanisms of how the large school size in nature is coordinated and whether or not local

interactions are still one of the mechanisms for energy saving when scaling up school size.

Future hydrodynamic and long-term 3-D tracking studies are needed to investigate the specific

schooling formations and hydrodynamic mechanisms used by live fish for energy saving (as

summarized in Fig.1) (*49*).

We show that danio in a school keep a relatively consistent 3-D distance from neighbours

and have small variation in mean position even as individuals routinely change position within

the school (Fig. 4G). Computational fluid dynamic analyses of fish schools show that a small 3D

neighbouring distance (*e.g.* < 0.4 BL s$^{-1}$) increases drag and can increase swimming costs,

whereas a large 3-D neighbouring distance (*e.g.* > 2 BL s$^{-1}$) reduces the hydrodynamic benefits

(*48*). There may thus be an optimal mean distance among individuals within a school to

maximize energetic savings. Whether inter-individual distance should change as swimming

speed increases remains an open question and may be a function of school size. While we cannot

completely preclude the possibility that elongation of danio fish school might be due to the

weaker individuals falling behind, the fact that the school maintained a stable 3-D distance as

speed increased and that fish continued to change position within the school suggests that the

elongated school volume at higher speeds may reflect changing hydrodynamic interactions

among individuals (Fig. 1).

To encapsulate the complex interaction among animal physiology, kinematics and fluid

dynamics, more comprehensive quantification beyond simple kinematic metrics are necessary.

For example, the wing flapping frequency of birds in a V-formation can be higher than for

solitary flight despite strong indirect evidence of overall energy saving by V-formation flying (*2*)

(*12*) (*19*), and flapping phase of adjacent birds may be informative for understanding the fluid

dynamic advantages of V-formation flying (*12*). Likewise, phase matching of body motion

between neighbouring fish can help individual fish within the school to boost thrust or reduce

drag (*20*). Fish can also adjust their body stiffness and maintain $f_{TB}$ while reducing the amount of

muscle activity needed to generate movement (*50*) (*51*) (*52*). Thus, we did not observe consistent

and substantial changes in the $f_{TB}$ of fish within the schools compared to solitary swimmers

despite demonstrating substantial energetic savings by the group at high speed. Although

reduced $f_{TB}$ has been used in past studies as an indirect indicator of energy saving in fishes (*53*)

(*54*), we caution that $f_{TB}$ is not necessarily an indicator of energy use in group dynamics where

fish constantly and dynamically interact with complex vortices and flow generated by other

individuals. By simultaneously measuring kinematics and energetics, we discovered a

downshifted performance curve of TEE per tail beat in fish schools at higher speeds (Fig. 4A),

even with limited alteration in $f_{TB}$ at lower swimming speeds within the school. This suggests

that the locomotor muscle fibres in the body musculature of fish within the school need to

generate less force as indicated by the lower measured TEE. Fish can possibly fine-tune

undulatory motion to harness kinetic energy from nearby vortices (*20*) (*48*), reducing biological

energy contribution for thrust.

**Aerobic metabolic rate–speed curve of fish schools**

We discovered that the aerobic metabolic rate ($\dot{M}O_2$)–speed curve at speeds less than 3

BL s$^{-1}$ is *U*-shaped, with swimming at 0.3 BL s$^{-1}$ utilizing the same aerobic energy as moving at 3

BL s$^{-1}$ (Fig. 2D).  The entire aerobic energy-speed performance curve is concave upward in an

extended *J*-shape (Fig. 2C), and non-linear, in contrast to several previous analyses of energy use

with speed in fish which have shown linear relationships (*55*) (*56*) (*57*). By quantifying energy

use over a wide speed range and by measuring aerobic energy use at 14 distinct speeds, we

showed that locomotion at the lowest speeds involves increased energy use relative to swimming

at speeds up to 3 BL s$^{-1}$, similar to previous results from skate locomotion (*58*). This suggests

that analyses of fish energy use as speed increases would benefit from increasing the number of

tested speeds, and that efforts to extend speed measurements to both the lowest and highest

speeds would enable a greater level of precision in measuring fish locomotor performance.

One key finding in this regard is that fish schools show a minimal absolute energetic cost

($\dot{M}O_2$) at a mean group swimming speed of ~1.0 BL s$^{-1}$ that is higher than the minimum

swimming speeds tested. Thus, fish schools can swim at a speed with the least amount of energy

use and potentially extend the distance travelled with the same amount of metabolic substrate

onboard. This swimming speed of minimum cost closely matches the migratory speeds of

carangiform and subcarangiform migratory fishes (often as schools) which are in the range of 0.5

to 1.5 BL s$^{-1}$ as recorded by tags on migrating fish (*59*, *60*) (Fig. S1; Tables S1, S3). Although *D.*

*aequipinnatus* is not a migratory species, our results suggest that the migratory speed of fish

schools likely occurs at the speed showing the minimum aerobic metabolic rate for long-distance

locomotion. Indeed, the Strouhal number when fish schools swim at $U_{opt}$, showing a minimum

aerobic energy cost of locomotion, is 0.3, a hallmark of efficient swimming for many fish species

(*61*). Fish of different size swimming at 1 BL s$^{-1}$ will necessarily move at different Reynolds

numbers, and hence the scaling of body size to swimming speed needs to be considered in future

analyses of other species that differ in size.

The exact energetic mechanisms underpinning the chosen migratory speeds of fish

schools would benefit from more in-depth studies. Since both fish schools and solitary

individuals have $U_{opt}$ of ~1 BL s$^{-1}$, the $\dot{M}O_2$–speed curve of a fish school might be the average of

individual curves from fish within the school. However, pushing and pulling forces from

hydrodynamic interactions of neighbouring fish could help swimmers settle into stable

arrangements, a phenomenon known as the Lighthill conjecture (*62*). Further studies will need to

explore whether individuals within the school are more frequently located at points

recommended by the Lighthill conjecture which could potentially result in minimum metabolic

costs of locomotion at $U_{opt}$ in fish schools (*49*). Research on fish swimming in large aquaculture

circular tanks has discovered that holding a large school of salmonids at a water velocity of ~1

BL s$^{-1}$ resulted in healthy growth and good conditions (*63*). This phenomenon seems to suggest

that the minimum metabolic costs of locomotion yield the largest available aerobic capacity for

growth and other functions.

Higher locomotor costs at the lowest speeds are likely caused by higher postural costs for

active stability adjustments in the near-still fluid. The direct link between higher energetic costs

caused by a higher 3-D body angle and higher turning frequency at the lowest speeds in solitary

danio (Fig. 4D, E) supports previous results in skates (*58*) where energetic costs are high at the

very low swimming speeds due to the increased energetic cost involved in maintaining body

stability and generating lift in negatively buoyant fishes. We expect more species that move in

the fluid to show a higher postural cost in near-still fluid but careful measurements of both body

kinematics and position along with high-resolution respirometry and accurate low-speed flow

control will be required to demonstrate this phenomenon in a diversity of fish species. Elevated

costs should be indicated at the lowest speeds as a direct increase in $\dot{M}O_2$ and not just as COT.

When the denominator (*i.e.* speed) for deriving COT is less than 1, COT tends to skew towards

the higher value at the lowest speeds. Mechanistically, fish at low fluid speeds have lower

muscle and propeller efficiencies (*44*) with reduced fluid-assisted stability. Increased active

stability adjustments such as fin movements (*58*) and tilting behaviours that enable the fish's

body to act as a hydrofoil (*64*) are used for stability control (see supplementary video).

Moreover, danio do not show intermittent swimming gaits at low speed as observed in some labriform swimmers (65) (66), and hence gait-specific kinematics most likely do not play a role in the elevated locomotion cost that we observe at the lowest speeds. Also, the higher turning frequency at < 0.75 BL s$^{-1}$ in fish schools can relate to the need for coordination with neighbours. Nevertheless, as fish move more rapidly over a speed range, we note that the total cost of transport of ~1.2 kJ km$^{-1}$ kg$^{-1}$ at 3 BL s$^{-1}$ is among the lowest recorded for aquatic organisms, and is less than the TCOT (1.5 kJ km$^{-1}$ kg$^{-1}$) for jellyfish (the most energetically efficient low-speed swimmers) (4) swimming at < 2 BL s$^{-1}$ (67).

In summary, our experiments on giant danio have demonstrated substantial energy conservation resulting from schooling dynamics across a wide range of speeds in fish. Direct measurement of both aerobic and non-aerobic energy use is critical for understanding the rapid collective movement of animals. Fish schooling in the high-drag viscous aquatic medium serves as a model for understanding how group movement by animals can be a more energy-efficient biological collective than movement by isolated individuals. By increasing maximum aerobic performance, fish schools save anaerobic energy and reduce the recovery time after peak swimming performance. Furthermore, by decreasing the proportion of metabolic capacity and recovery time devoted to locomotion, animals can apportion more energy to other fitness-related activities, such as digestion, growth and reproduction. More broadly, comprehending how the collective dynamics of animal movements in the water, land, and air can modify the energy use profiles of individuals provide a better understanding of the ecological and evolutionary implications of group locomotion.

**Materials and Methods**

**Experimental animals**

The experiments were performed on giant danio (*Devario aequipinnatus*) that were acquired from a local commercial supplier near Boston, Massachusetts USA (Table S4). Five schooling groups are randomly distributed and housed separately in five 37.9 l aquaria (n=8 per tank). The five solitary individuals are housed separately in five 9.5 l aquaria (n=1 per tank). The individual housing condition acclimated the single *D. aequipinnatus* to the solitary environment and helped to reduce any isolation stress that might elevate whole-organism metabolic rate. In fact, the aerobic locomotion cost of solitary individuals showed no statistical difference from (in fact, being numerically lower) that of fish schools at a very low testing speed. The flow speed is similar to some areas of the aerated home aquarium for each individual fish. This suggests that the stress of solitary fish likely does not meaningfully contribute to the higher locomotor costs (*see* experimental protocol for more details on mitigating the stress). The condition factor showed no difference between solitary fish and fish schools (0.81 vs. 0.99; $t_8 = 2.14$, $p = 0.065$). All aquaria have self-contained thermal control (28 °C), an aeration system (>95 % air saturation, % sat.) and a filtration system. Water changes (up to 50% exchange ratio) were carried out weekly. Fish were fed *ad libitum* daily (TetraMin, Germany). Animal holding and experimental procedures were approved by the Harvard Animal Care IACUC Committee (protocol number 20-03-3).

**Integrated Biomechanics & Bioenergetic Assessment System (IBAS)**

The core of our Integrated Biomechanics & Bioenergetic Assessment System (IBAS) (Fig. S4) is a 9.35-l (respirometry volume plus tubing) customized Loligo® swim-tunnel respirometer (Tjele, Denmark). The respirometer has an electric motor, and a sealed shaft attached to a

452 propeller located inside the respirometer. Regulating the revolutions per minute (RPM) of the

453 motor controls water velocity of the tunnel. The linear regression equation between RPM and

454 water velocity (V) is established ($V = 0.06169 \cdot RPM - 5.128$, $R^2 = 0.9988$, $p < 0.0001$) by

455 velocity field measured by particle image velocimetry (PIV). Hence, the aerobic costs during

456 locomotion are measured through the regulation of the water velocity.

457   The swim-tunnel respirometer is oval-shaped. The central hollow space of the respirometer

increases the turning radius of the water current. As a result, the water velocity passing the cross-

section of the swimming section ($80 \times 80 \times 225$ mm) is more homogenous (validated by direct

particle image velocimetry (PIV) of flow in the working section of the respirometer following

the procedures in (68)). Moreover, a honeycomb flow straightener ($80 \times 80 \times 145$ mm) is

installed in the upstream section of the swimming section that was specifically designed to create

relatively uniform flow across the working section (also confirmed by direct flow imaging).

Based on our flow visualization analyses, the effective distance of slow-moving fluid due to

boundary layer was <2.5mm at speeds above 2 BL s$^{-1}$. The boundary layer played an ever-

diminishing role at higher speeds (> 4 BL s$^{-1}$) when energy saving of fish schools becomes more

predominant. Danio (~10 mm wide) cannot effectively hide in the narrow boundary layer created

by our flow baffle system. In addition, the convex hull volume of the fish school did not change

as speed increased, suggesting that the fish school was not flattening against the wall of the swim

tunnel, a typical feature when fish schools are benefiting from wall effects. In nature, fish in the

centre of the school effectively swim against a 'wall' of surrounding fish where they can benefit

from hydrodynamic interactions with neighbours.

473   To standardize the experimental apparatus and avoid instrument variation, solitary fish and

fish schools are measured in the same swim-tunnel respirometer, and the same overall

experimental design is used as in previous studies (14) (13) (15). The ratios of

respirometer:individual volume ($r_{RI}$) in our experiments was 2200 for individual fish (we used

larger solitary *D. aequipinnatus* to increase the signal-to-noise ratio), and 693 for fish schools.

The $r_{RI}$ is essentially a balance between giving enough space for fish to exhibit natural

locomotion and reducing wall effects and generating a reliable signal-to-noise ratio to measure

the decline of dissolved $O_2$ (DO) in water (*69*). The increase in the signal-to-noise ratio can come

from the better technology of the $O_2$ probe (we used an optical $O_2$ probe which has a higher

measurement sensitivity than the Winkler method or Electrode & Galvanic probes used in the

previous studies, *see* (*70*) for review) and modifications to the swim-tunnel respirometer. We

used a water loop is installed 95 cm downstream of the propeller with water returned to the

respirometer 240 cm upstream of the swimming section. Flow in the water loop moves

(produced by an in-line circulation pump, Universal 600, EHEIM GmbH & Co KG, Deizisau,

Germany) in the same direction as the water flow in the swimming tunnel. A high-resolution

fibre optic $O_2$ probe (Robust oxygen probe OXROB2, PyroScience GmbH, Aachen, Germany) is

sealed in the loop in the downstream of the circulation pump where there is better mixing to

continuously measure the DO level in the water (recording frequency ~1 Hz, response time <

15s). The designated water sampling loop together with the water mixing by the propeller and

water pumps effectively reduces the noise-to-signal ratio. As a result, the respirometry system

reaches a stable signal-to-noise ratio once the sampling window is longer than 1.67 mins (*see*

Fig. S5), well within the duration of the velocity step to obtain a stable signal-to-noise ratio for

calculating $\dot{M}O_2$ (*32*).

The oxygen probe was calibrated to anoxic (0 % sat., a solution created by super-saturated

sodium sulphite and bubbling nitrogen gas) and fully aerated water (100 % sat.). After fish

removal, the background $\dot{M}O_2$ in the swim-tunnel respirometer was measured for a 20-min

sealed period before and after each trial to calculate the average background $\dot{M}O_2$ (< 6% of fish

$\dot{M}O_2$), which was used to correct for the $\dot{M}O_2$ of fish (*69*). The pre-filtered water (laboratory

grade filtration system) is constantly disinfected by UV light (JUP-01, SunSun, China) located in

an external water reservoir to suppress the growth of microbial elements. Water changes of 60%

total volume occurred every other day and a complete disinfection by sodium hypochlorite is

conducted weekly (Performance bleach, Clorox & 1000 ppm).

To simultaneously measure schooling dynamics and swimming kinematics, the customized

oval-shaped swim-tunnel respirometer is located on a platform with an open window beneath the

swimming section. The platform is elevated 243 mm above the base to allow a front surface

mirror to be installed at a 45° angle. This mirror allows a high-speed camera (FASTCAM Mini

AX50 type 170K-M-16GB, Photron Inc., United States, lens: Nikon 50mm F1.2, Japan) to

record the ventral view. The second camera (FASTCAM Mini AX50 type 170K-M-16GB,

Photron Inc., United States, lens: Nikon 50mm F1.2, Japan) is positioned 515 mm to the side of

the swimming section to record a lateral view. Synchronized lateral and ventral video recordings

were made at 125 fps, and each frame was 1024 by 1024 pixels. To avoid light refraction passing

through the water and distorting the video recordings, the swim-tunnel respirometry is not

submerged in a water bath. Temperature regulation of the respirometer is achieved by regulating

room temperature, installing thermal insulation layers on the respirometer and replenishing the

water inside the respirometer from a thermally regulated (28 °C, heater: ETH 300, Hydor, United

States & chiller: AL-160, Baoshishan, China) water reservoir (insulated 37.9-l aquarium) located

externally.

The aerated (100% sat., air pump: whisper AP 300, Tetra, China) reservoir water is flushed

(pump: Universal 2400, EHEIM GmbH & Co KG, Deizisau, Germany) to the respirometer

through an in-line computer-controlled motorized ball valve (U.S. Solid) installed at the in-flow

tube. The other in-line one-way valve is installed at the out-flow tube. The out-flow tube is also

equipped with a one-way valve. The valve is shut during the measurement period, a

precautionary practice to eliminate the exchange of water between the respirometer and the

external reservoir when the water moves at a high velocity inside the respirometer. This flushing

was manually controlled to maintain DO above 80 % sat. Every time the respirometer was closed

to measure $\dot{M}O_2$, the water temperature fluctuates no more than 0.2 °C. The water temperature

inside the respirometer is measured by a needle temperature probe (Shielded dipping probe,

PyroScience GmbH, Aachen, Germany) sealed through a tight rubber port of the respirometer.

To allow fish to reach the undisturbed quiescent state during the trial (another stress

mitigation practice), the entire IBAS is covered by laser blackout sheets (Nylon Fabric with

Polyurethane Coating; Thorlabs Inc, New Jersey, United States). The room lights are shut off

and foot traffic around IBAS is restrained to the absolute minimum. Fish are orientated by dual

small anterior spots of white light (lowest light intensity, Model 1177, Cambridge Instruments

Inc, New York, United States) for orientation (one to the top and the other to the side) of the

swimming section. The test section is illuminated by infrared light arrays to allow high-speed

video recording with minimal disturbance of fish behaviour.

**Experimental Protocol**

The energy use of vertebrates at lower speeds is primarily aerobic, while for high-speed

movement anaerobic metabolic pathways are activated to supply the additional (largely shorter-

term) energy needs (*9*). While whole-animal aerobic metabolism is measured by oxygen ($O_2$)

uptake rate ($\dot{M}O_2$), the non-aerobic $O_2$ cost (mostly through high-energy phosphate and

anaerobic glycolysis) is measured as excess post-exercise $O_2$ consumption (EPOC)(*10*). Both

metabolic energy sources contribute to the total energy expenditure (TEE) required for

movement. For a given workload, the higher the maximum aerobic performance of the animal,

the less the need for the anaerobic energy contribution (*9*, *35*). The experimental protocol

captures both metabolic energy contributions by measuring $\dot{M}O_2$ during locomotion and EPOC

afterwards.

We studied five replicate schools and five replicate individuals. Swimming performance test

trials were conducted with *Devario aequipinnatus* fasted for 24 hours, a sufficient period for a

small size species at 28 °C (*i.e.* high resting $\dot{M}O_2$) to reach an absorptive state. In fact, we

observed no specific dynamic action, the amount of $O_2$ consumed for digestion during the first

diurnal cycle (Fig. S3). Prior to the swimming performance test, testing fish were gently weighed

and placed in the swim-tunnel respirometer. The fish swam at 35% $U_{crit}$ for 30 mins to help

oxidize the inevitable but minor lactate accumulation during the prior handling and help fish

become accustomed to the flow conditions in the swim-tunnel respirometer (*43*). After this time,

the fish to be tested were habituated (>20 hours) to the respirometer environment under quiescent

and undisturbed conditions. The long habituation period also helps to reduce the stress and

further reduce the likelihood that the solitary fish might be more stressed than the fish schools

and showed an elevated $\dot{M}O_2$ (*13*). During this time, we used an automatic system to measure the

resting $\dot{M}O_2$ for at least 19 hours. Relays (Cleware GmbH, Schleswig, Germany) and software

(AquaResp v.3, Denmark) were used to control the intermittent flushing of the respirometer with

fresh water throughout the trial to ensure $O_2$ saturation of the respirometer water. $\dot{M}O_2$ was

calculated from the continuously recorded DO level (at 1 Hz) inside the respirometer chamber.

The intermittent flow of water into the respirometer occurred over 930 s cycles with 30 s where

water was flushed into the respirometer and 900 s where the pumps were off and the

respirometer was a closed system. The first 240 s after each time the flushing pump was turned

off were not used to measure $\dot{M}O_2$ to allow $O_2$ levels inside the respirometer to stabilize. The

remaining 660 s when the pumps were off during the cycle were used to measure $\dot{M}O_2$. The in-

line circulation pump for water in the $O_2$ measurement loop stayed on throughout the trial.

We characterize the aerobic costs for the swimming performance of fish using an

established incremental step-wise critical swimming speed ($U_{crit}$) test (*10*). The first preliminary

trial determined the $U_{crit}$ of this population of *Devario aequipinnatus* as 8 BL s$^{-1}$. Characterizing

the swimming performance curve required a second preliminary trial to strategically select 9

water velocities (0.3, 0.5, 0.8, 1.0, 1.3, 1.5, 1.8, 2.3, 2.8 BL s$^{-1}$) to bracket the hypothesized

upward concave shaped aerobic metabolism-speed curve at the lower speed (< 40% $U_{crit}$).

Additional five water velocities (3.8, 4.9, 5.9, 6.9, 8.0 BL s$^{-1}$) are used to characterize the

exponentially increasing curve to the maximum and sustained swimming speed, $U_{crit}$ (*see* Fig.

S6). Altogether, 14 points provide a reliable resolution to characterize the swimming

performance curve. The cross-section of the danio school is ~10% of the cross-sectional area of

the swim tunnel, hence the blocking effect is negligible (*71*).  At each water velocity, fish swam

for 10 mins (*58*) to reach a steady state in $\dot{M}O_2$ at low speed (*see* Fig. S7). Above 40% $U_{crit}$, $\dot{M}O_2$

can become more variable (*32*). Hence, in this protocol, we focus on measuring the sustained

aerobic energy expenditure by calculating the average $\dot{M}O_2$ for each 10-min velocity step using

Eqn 1. At the 5$^{th}$ min of each velocity step, both ventral and lateral-view cameras are triggered

simultaneously to record 10-sec footage at 125 frames per second, at 1/1000 shutter speed and

1024 ×1024 pixel resolution. Thus, both data streams of $\dot{M}O_2$ and high-speed videos are recorded

simultaneously. The $U_{crit}$ test is terminated when 12.5% of fish in the school or a solitary

individual touches the back grid of the swimming section for more than 20 secs (*43*). The $U_{crit}$

test lasted ~140 mins and estimates the aerobic portion of TEE over the entire range of

swimming performance.

To measure the contribution of non-aerobic $O_2$ cost, where the majority of the cost is related

to substrate-level phosphorylation, and to calculate the TEE for swimming over the entire speed

range, we measured EPOC after the $U_{crit}$ test for the ensuing 19 hours, recorded by an automatic

system. Most previous measurements of EPOC have used a duration of ~5 hours (*see* review

(*42*)), but our extended measurement period ensured that longer duration recovery $O_2$

consumption during EPOC was measured completely as fish were exercised to $U_{crit}$ (*see*

summary table in *16*). The intermittent flow of water into the respirometer occurred over 30 s to

replenish the DO level to ~95% sat. For the following 900 s the flushing pump remained closed,

and the respirometer becomes a closed system, with the first 240 s to allow $O_2$ saturation inside

the respirometer to stabilize. The remaining 660 s when the flushing pump was off during the

cycle were used to measure $\dot{M}O_2$ (*see* Eqn. 1). The cycle is automated by computer software

(AquaResp v.3) and provided 74 measurements of $\dot{M}O_2$ to compute EPOC. Upon the completion

of the three-day protocol, the school or individual fish are returned to the home aquarium for

recovery. The fish condition was closely monitored during the first 48 hours after the

experiment, during which no mortality was observed.

**Bioenergetic measurement and modeling**

To estimate the steady-rate whole-animal aerobic metabolic rate, $\dot{M}O_2$ values were

calculated from the sequential interval regression algorithm (Eqn. 1) using the DO points

continuously sampled (~1 Hz) from the respirometer.

$$\dot{M}O_2 = \left[ \frac{d_{DO[i,(i+a)]}}{d_{t[i,(i+a)]}} \bullet \left( V_r - V_f \right) \bullet S_O \right] \Big/ \left( t \bullet M_f \right) \qquad \text{(Eqn. 1)}$$

Where $\frac{d_{DO}}{d_t}$ is the change in $O_2$ saturation with time, $V_r$ is the respirometer volume, $V_f$ is

the fish volume (1 g body mass = 1 ml water), $S_o$ is the water solubility of $O_2$ (calculated by

AquaResp v.3 software) at the experimental temperature, salinity and atmospheric pressure, $t$ is a

time constant of 3600 s h$^{-1}$, $M_f$ is fish mass, and $a$ is the sampling window duration, $i$ is the next

$PO_2$ sample after the preceding sampling window.

To account for allometric scaling, the $\dot{M}O_2$ values of solitary fish were transformed to match

the size of the individual fish in the school (*see* Table S4) using an allometric scaling exponent

(b = 0.7546). The calculation of the scaling relationship [$Log_{10}(\dot{M}O_2)$ = b•Log10($M$) + Log10($a$),

where $M$ is the body mass & $a$ is a constant] was performed by least squares linear regression

analysis (y = 0.7546 • x + 0.2046; $R^2$ = 0.6727, $p < 0.0001$) on the 180 data points of metabolic

rate and body mass from a closely related species (the best available dataset to our knowledge) .

(The mass scaling for tail beat frequency was not conducted because of the lack of data for *D.*

*aequipinnatus* and its related species. Using the scaling exponent of distant species for mass

scaling of tail beat frequency will introduce errors of unknown magnitude.) The allometrically

scaled $\dot{M}O_2$ values were used to derive other energetic metrics (listed below & such as aerobic

scope) for the solitary fish. The energetic metrics of fish schools are calculated from the mass-

specific $\dot{M}O_2$.

The resting oxygen uptake ($\dot{M}O_{2rest}$), the minimum resting metabolic demands of a group of

fish or a solitary individual, is calculated from a quantile 20% algorithm (*74*) using the $\dot{M}O_2$

estimated between the 10$^{th}$–18$^{th}$ hour and beyond the 32$^{nd}$ –51$^{st}$ hour of the trial. These are the

periods of quiescent state when fish completed the EPOC from handling and swimming test.

The $\dot{M}O_2$ for the aggregation (Fig. 2B, $\dot{M}O_{2aggregate}$) is calculated as the average value using

the $\dot{M}O_2$ estimated between the 10$^{th}$–18$^{th}$ hour without the effects of any tests.

Minimum oxygen uptake ($\dot{M}O_{2min}$) is the lowest $\dot{M}O_2$ value recorded in the entire trial, which always occurred at the optimal speed when a school of fish collectively reached the lowest $\dot{M}O_2$ value.

Active oxygen uptake ($\dot{M}O_{2active}$) is the highest average $\dot{M}O_2$ when fish are actively swimming (*10*).

The aerobic scope (the metric for aerobic capacity) is the numerical difference between $\dot{M}O_{2active}$ and $\dot{M}O_{2min}$ (*i.e.* $\dot{M}O_{2active} - \dot{M}O_{2min}$) (*10*).

The percentage of aerobic scope (% aerobic scope) is calculated by normalizing the $\dot{M}O_2$ value at a water velocity as a % aerobic scope: $[(\dot{M}O_2 - \dot{M}O_{2min}) / (\dot{M}O_{2active} - \dot{M}O_{2min})]*100\%$. The apportioning of aerobic scope to swimming performance is computed across the entire range of swim speeds.

The excess post-exercise oxygen consumption (EPOC) is an integral area of $\dot{M}O_2$ measured during post-exercise recovery, from the end of $U_{crit}$ until reached $\dot{M}O_{2rest}$ plus 10% or within 24 hours post-exercise, whichever endpoint occurred first (*72*). This approach reduces the likelihood of overestimating EPOC due to any potential spontaneous activities (*72*). To account for the allometric scaling effect, we used the total amount of $O_2$ consumed (mg $O_2$) by the standardized body mass of fish (1.66 g) for fish schools and solitary fish.

We model EPOC (*i.e.* non-aerobic $O_2$ cost) and $\dot{M}O_2$ during $U_{crit}$ to estimate a *total* $O_2$ cost over the duration of the swimming performance test. Our conceptual approach was pioneered by Brett (*10*) in fish (*75*) and is also used in sports science (*9*). Mathematical modeling of EPOC and $\dot{M}O_2$ during swimming was applied to study the effects of temperature on the total cost of swimming for migratory salmon (*43*). We improved the mathematical modeling by applying the following physiological and physics criteria. The first criterion is that significant accumulation of glycolytic end-product occurred when fish swimming above 50% $U_{crit}$ (*31*) which corresponds to

> 40% $\dot{M}O_{2max}$ (or ~ 50% aerobic scope) (*9*). This is also when fish start unsteady-state burst-&-

glide swimming gait (*31*). The second criterion is that the integral area for the non-aerobic $O_2$

cost during swimming can only differ by ≤ 0.09% when compared to EPOC. The non-aerobic $O_2$

cost during swimming is the area bounded by modeled $\dot{M}O_2$ and measured $\dot{M}O_2$ as a function of

time when fish swim > 50% $U_{crit}$ (*see* Fig. 2A & Table S2). The third criterion is that total energy

expenditure is expected to increase exponentially with swimming speed (Fig. S8). Specifically,

these curves were fitted by power series or polynomial models, the same models that describe the

relationship between water velocity and total power and energy cost of transport (Fig. S8).

Following these criteria, the non-aerobic $O_2$ cost at each swimming speed is computed by a

percentage (%) modifier based on the aerobic $O_2$ cost (Table S2). The exponential curve of total

$O_2$ cost as swimming speed of each fish school or solitary individual was derived by an iterative

process until the difference between non-aerobic $O_2$ cost and EPOC met the 2nd criterion. The

sum of non-aerobic $O_2$ cost and aerobic cost gives the total $O_2$ cost.

Total energy expenditure (TEE) is calculated by converting total $O_2$ cost to kJ × kg$^{-1}$ using

an oxy-calorific equivalent of 3.25 cal per 1 mg $O_2$ (*76*).

Cost of transport (COT), in kJ × km$^{-1}$ × kg$^{-1}$ is calculated by dividing TEE by speed (in km

× h$^{-1}$) (*58*).

**Three-dimensional kinematic data extraction from high-speed videography**

We used two synchronized 10-sec high-speed videos (lateral and ventral views, at each

speed) for kinematic analyses. We calibrated the field of view of the high-speed cameras using a

direct linear transformation for three-dimensional (3-D) kinematic reconstruction (DLTdv8)(*77*)

by applying a stereo calibration to the swimming section of the respirometer (*see* Fig. S9). We

digitized the anatomical landmarks of fish (*see* Fig. S10) to obtain the X, Y, and Z coordinates

for each marker at the 1st sec, 5th sec and 10th sec for videos recorded at each speed. These

coordinates are used to calculate the following kinematic parameters. All the calculations are

validated on the known length and angle of test objects inserted into the tank working section.

The 3-D distance between the tip of the nose of each fish in the school per frame is

calculated in vector Eqn. 2

$$d = \sqrt{(Xa - Xb)^2 + (Ya - Yb)^2 + (Za - Zb)^2} \qquad \text{(Eqn. 2)}$$

Where spatial coordinates of two neighbouring fish are $(X_a, Y_a, Z_a)$ and $(X_b, Y_b, Z_b)$

respectively.

The 3-D angle of each fish in the school to the frontal plane per frame is calculated by Eqn.

3

$$\theta = \arccos \frac{(X2-X1)\cdot(X3-X1)+(Y2-Y1)\cdot(Y3-Y1)+(Z2-Z1)\cdot(Z3-Z1)}{\sqrt{(X2-X1)^2\cdot(Y2-Y1)^2\cdot(Z2-Z1)^2}\cdot\sqrt{(X3-X1)^2\cdot(Y3-Y1)^2\cdot(Z3-Z1)^2}} \qquad \text{(Eqn. 3)}$$

where the spatial coordinates of the caudal peduncle, nose of fish and right-angle crosshair

between the peduncle and the nose are $(X_1, Y_1, Z_1)$, $(X_3, Y_3, Z_3)$ and $(X_2, Y_2, Z_2)$ respectively (*see*

Fig. S9).

The fish's angle to water flow per frame is calculated using the arctangent function in Excel

(Microsoft, United States) using the spatial coordinates of the caudal peduncle and nose of the

fish.

The school length (in *X*-axis) is calculated as a 3-D distance (Eqn. 2) between the nose of

the first fish of the school to the caudal peduncle of the last fish in the school.

The values (3-D distance of the individuals, 3-D body angle, 3-D school length) calculated

above were averaged among the three frames at each speed as a representative kinematic feature

of the schools (or solitary fish) for each speed. The standard deviation of the angle to water flow

in fish schools is calculated from the angular values of the individuals.

Tail beat frequency ($f_{TB}$), in Hz, is calculated as the number of tail beats observed within

one second. We sampled 10 tail beats from fish exhibiting steady-state swimming to obtain an

average and representative $f_{TB}$ for each solitary fish at each speed. We sampled 10 tail beats of

steady-state swimming for three individuals in the center of the fish school for an average and

representative $f_{TB}$ for each fish school.

Turning frequency is the total number of events when fish made a 90° turn in the 10-sec

video. To make the total number of turns per individual and fish schools comparable (multiple

individuals inevitably have a larger absolute number of turns), we derived the average turning

frequency per schooling fish by dividing the total number of turns of fish school by the number

of fish in the school.

Total energy expenditure per tail beat (kJ kg$^{-1}$ beat$^{-1}$) is calculated by TEE • ($f_{TB}$ • 60 )$^{-1}$ to

estimate the total metabolic energy spent to achieve one stride length.

Additional calculations of fluid dynamic metrics were:

Strouhal number = ($f_{TB}$ • tail beat amplitude) • $U^{-1}$ (8). Tail beat amplitude is measured as

the average distance of five peak-to-peak oscillation amplitudes of the tip of the fish's tail. The

measurement is conducted on the calibrated high-speed video in video analysis software

(Phontron FASTCAM Viewer 4, Photron USA, Inc.).

Reynolds number = (water density • $U$ • fish fork length) • water dynamic viscosity $^{-1}$ (8).

Water density and dynamic viscosity are given at 28 °C.

**Statistical analyses**

Measurement points are presented as mean ± s.e.m. For the metrics that failed normality

tests, logarithm transformations were applied to meet the assumptions of normality of residuals,

homoscedasticity of the residuals, and no trend in the explanatory variables. Since fish schools

exhibit the features of a coherent functional unit, the statistical model treated one school as one

sample size, and the experiment measured five replicates of fish schools. The majority of

statistical comparisons used the General Linear Model (Fixed factors: solitary fish vs. fish

schools & swimming speeds, the label of solitary fish & fish schools as a random effect) with

Holm–Šídák *post-hoc* tests. The few comparisons that used other statistical models are listed

below. The comparison of $\dot{M}O_{2min}$ at $U_{opt}$, $\dot{M}O_2$ of aggregating behaviours exhibited at the lowest

speed ($\dot{M}O_{2aggregate}$) and resting $\dot{M}O_2$ ($\dot{M}O_{2rest}$) in fish schools used one-way ANOVA with Holm–

Šídák *post-hoc* tests. The comparison of EPOC between solitary fish and fish schools used a two-

tailed Student's t-test. The comparison of the duration of EPOC between solitary fish and fish

schools used a two-tailed Student's t-test. The overall difference in percentage aerobic scope (%

aerobic scope) between fish schools and solitary individuals is compared by the Wilcoxon

singed-rank test over the entire range of 14 swimming speeds. The statistical analyses were

conducted in SPSS v.28 (SPSS Inc. Chicago, IL, USA). The best-fitting regression analyses were

conducted using Prism v.9.4.1 (GraphPad Software, San Diego, CA, USA). 95% C.I. values

were presented for all regression models as shaded areas around the regression or data points.

Statistical significance is denoted by *, **, ***, **** for *p*-values of $\leq 0.05$, $\leq 0.01$, $\leq 0.001$, $\leq$

0.0001 respectively.

**References and Notes**

1.  A. Biewener, S. Patek, *Animal Locomotion* (Oxford University Press, 2018).

2.  H. Weimerskirch, J. Martin, Y. Clerquin, P. Alexandre, S. Jiraskova, Energy saving in flight formation. *Nature* **413**, 697–698 (2001).

3.  F. E. Fish, Kinematics of ducklings swimming in formation: Consequences of position. *Journal of Experimental Zoology* **273**, 1–11 (1995).

4.  B. Blocken, T. van Druenen, Y. Toparlar, F. Malizia, P. Mannion, T. Andrianne, T. Marchal, G.-J. Maas, J. Diepens, Aerodynamic drag in cycling pelotons: New insights by CFD simulation and wind tunnel testing. *Journal of Wind Engineering and Industrial Aerodynamics* **179**, 319–337 (2018).

5.   I. Shinichiro, Aerodynamic effects by marathon pacemakers on a main runner. *Transactions of the Japan*
*Society of Mechanical Engineers Part B* **73**, 1975–1980 (2007).

6.   D. Weihs, Hydromechanics of fish schooling. *Nature* **241**, 290–291 (1973).

7.   Y. Zhang, G. V. Lauder, Energetics of collective movement in vertebrates. *Journal of Experimental Biology*
**226**, jeb245617 (2023).

8.   S. Vogel, *Life in Moving Fluids: The Physical Biology of Flow - Revised and Expanded Second Edition*
(Princeton University Press, 1981).

9.   J. Laforgia, R. T. Withers, C. J. Gore, Effects of exercise intensity and duration on the excess post-exercise
oxygen consumption. *Journal of Sports Sciences* **24**, 1247–1264 (2006).

10.  J. R. Brett, The respiratory metabolism and swimming performance of young sockeye salmon. *J. Fish. Res. Bd.*
*Can.* **21**, 1183–1226 (1964).

11.  F. W. H. Beamish, "Swimming capacity" in *Fish Physiology* (London: Academic Press, W.S. Hoar and D. J.
Randall., 1978; https://cir.nii.ac.jp/crid/1571135650623593984)vol. VII, pp. 101–187.

12.  S. J. Portugal, T. Y. Hubel, J. Fritz, S. Heese, D. Trobe, B. Voelkl, S. Hailes, A. M. Wilson, J. R. Usherwood,
Upwash exploitation and downwash avoidance by flap phasing in ibis formation flight. *Nature* **505**, 399–402
(2014).

13.  F. R. Parker Jr., Reduced metabolic rates in fishes as a result of induced schooling. *Transactions of the*
*American Fisheries Society* **102**, 125–131 (1973).

14.  M. V. Abrahams, P. W. Colgan, Risk of predation, hydrodynamic efficiency and their influence on school
structure. *Environ Biol Fish* **13**, 195–202 (1985).

15.  E. Burgerhout, C. Tudorache, S. A. Brittijn, A. P. Palstra, R. P. Dirks, G. E. E. J. M. van den Thillart, Schooling
reduces energy consumption in swimming male European eels, Anguilla anguilla L. *Journal of Experimental*
*Marine Biology and Ecology* **448**, 66–71 (2013).

16.  M. Currier, J. Rouse, D. J. Coughlin, Group swimming behaviour and energetics in bluegill Lepomis
macrochirus and rainbow trout Oncorhynchus mykiss. *Journal of Fish Biology* **98**, 1105–1111 (2021).

17.  M. Hvas, F. Oppedal, Influence of experimental set-up and methodology for measurements of metabolic rates
and critical swimming speed in Atlantic salmon Salmo salar. *Journal of Fish Biology* **95**, 893–902 (2019).

18.  B. L. Partridge, T. J. Pitcher, Evidence against a hydrodynamic function for fish schools. *Nature* **279**, 418–419
(1979).

19.  J. R. Usherwood, M. Stavrou, J. C. Lowe, K. Roskilly, A. M. Wilson, Flying in a flock comes at a cost in
pigeons. *Nature* **474**, 494–497 (2011).

20.  L. Li, M. Nagy, J. M. Graving, J. Bak-Coleman, G. Xie, I. D. Couzin, Vortex phase matching as a strategy for
schooling in robots and in fish. *Nat Commun* **11**, 5408 (2020).

21.  J. Kelly, Y. (潘宇) Pan, A. Menzer, H. (董海波) Dong, Hydrodynamics of body–body interactions in dense
synchronous elongated fish schools. *Physics of Fluids* **35**, 041906 (2023).

22.  F. E. Fish, C. M. Schreiber, K. W. Moored, G. Liu, H. Dong, H. Bart-Smith, Hydrodynamic performance of
aquatic flapping: Efficiency of underwater flight in the Manta. *Aerospace* **3**, 20 (2016).

23. M. Kurt, K. W. Moored, Flow interactions of two- and three-dimensional networked bio-inspired control
elements in an in-line arrangement. *Bioinspir. Biomim.* **13**, 045002 (2018).

24. K. D. Jones, C. M. Dohring, M. F. Platzer, Experimental and computational investigation of the Knoller-Betz
effect. *AIAA Journal* **36**, 1240–1246 (1998).

25. J. D. Stewart, A. Barroso, R. H. Butler, R. J. Munns, Caught at the surface: myctophids make easy prey for
dolphins and devil rays. *Ecology* **99**, 1894–1896 (2018).

26. T. J. Pitcher, A. E. Magurran, I. J. Winfield, Fish in larger shoals find food faster. *Behav Ecol Sociobiol* **10**,
149–151 (1982).

27. K. E. Jain, A. P. Farrell, Influence of seasonal temperature on the repeat swimming performance of rainbow
trout Oncorhynchus mykiss. *Journal of Experimental Biology* **206**, 3569–3579 (2003).

28. P. E. Holder, C. M. Wood, M. J. Lawrence, T. D. Clark, C. D. Suski, J.-M. Weber, A. J. Danylchuk, S. J.
Cooke, Are we any closer to understanding why fish can die after severe exercise? *Fish and Fisheries* **23**,
1400–1417 (2022).

29. O. A. Misund, A. Aglen, Swimming behaviour of fish schools in the North Sea during acoustic surveying and
pelagic trawl sampling. *ICES Journal of Marine Science* **49**, 325–334 (1992).

30. S. J. Peake, A. P. Farrell, Fatigue is a behavioural response in respirometer-confined smallmouth bass. *Journal
of Fish Biology* **68**, 1742–1755 (2006).

31. S. J. Peake, A. P. Farrell, Postexercise Physiology and Repeat Performance Behaviour of Free-Swimming
Smallmouth Bass in an Experimental Raceway. *Physiological and Biochemical Zoology* **78**, 801–807 (2005).

32. Y. Zhang, M. J. H. Gilbert, A. P. Farrell, Finding the peak of dynamic oxygen uptake during fatiguing exercise
in fish. *Journal of Experimental Biology* **222** (2019).

33. A. P. Farrell, S. M. Clutterham, On-line venous oxygen tensions in rainbow trout during graded exercise at two
acclimation temperatures. *Journal of Experimental Biology* **206**, 487–496 (2003).

34. F. E. Fish, J. F. Fegely, C. J. Xanthopoulos, Burst-and-coast swimming in schooling fish (Notemigonus
crysoleucas) with implications for energy economy. *Comparative Biochemistry and Physiology Part A:
Physiology* **100**, 633–637 (1991).

35. G. A. Gaesser, G. A. Brooks, Metabolic bases of excess post-exercise oxygen consumption: a review. *Med Sci
Sports Exerc* **16**, 29–43 (1984).

36. B. C. Jayne, G. V. Lauder, New data on axial locomotion in fishes: How speed affects diversity of kinematics
and motor patterns. *American Zoologist* **36**, 642–655 (1996).

37. L. C. Rome, P. T. Loughna, G. Goldspink, Temperature Acclimation: Improved Sustained Swimming
Performance in Carp at Low Temperatures. *Science* **228**, 194–196 (1985).

38. C. L. Milligan, Metabolic recovery from exhaustive exercise in rainbow trout. *Comparative Biochemistry and
Physiology Part A: Physiology* **113**, 51–60 (1996).

39. M. Kjaer, J. Bangsbo, G. Lortie, H. Galbo, Hormonal response to exercise in humans: influence of hypoxia and
physical training. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* **254**,
R197–R203 (1988).

40. H. F. Brown, D. Difrancesco, S. J. Noble, How does adrenaline accelerate the heart? *Nature* **280**, 235–236
(1979).

841. C. A. Ross, D. A. Ruggiero, D. H. Park, T. H. Joh, A. F. Sved, J. Fernandez-Pardal, J. M. Saavedra, D. J. Reis,
838      Tonic vasomotor control by the rostral ventrolateral medulla: effect of electrical or chemical stimulation of the
839      area containing C1 adrenaline neurons on arterial pressure, heart rate, and plasma catecholamines and
840      vasopressin. *J. Neurosci.* **4**, 474–494 (1984).

42. Y. Zhang, G. Claireaux, H. Takle, S. M. Jørgensen, A. P. Farrell, A three-phase excess post-exercise oxygen
     consumption in Atlantic salmon Salmo salar and its response to exercise training. *Journal of Fish Biology* **92**,
     1385–1403 (2018).

43. C. G. Lee, A. P. Farrell, A. Lotto, S. G. Hinch, M. C. Healey, Excess post-exercise oxygen consumption in
     adult sockeye (Oncorhynchus nerka) and coho (O. kisutch) salmon following critical speed swimming.
     *Journal of Experimental Biology* **206**, 3253–3260 (2003).

44. P. Webb PW;, Hydrodynamics and Energetics of Fish Propulsion. *Bulletin of the Fisheries Research Board of*
     *Canada* **Bulletin 190** (1975).

45. A. Costanzo, C. K. Hemelrijk, Spontaneous emergence of milling (vortex state) in a Vicsek-like model. *J. Phys.*
     *D: Appl. Phys.* **51**, 134004 (2018).

46. R. A. Rountree, Association of Fishes with Fish Aggregation Devices: Effects of Structure Size on Fish
     Abundance. *Bulletin of Marine Science* **44**, 960–972 (1989).

47. S. Camazine, J.-L. Deneubourg, N. R. Franks, J. Sneyd, G. Theraula, E. Bonabeau, *Self-Organization in*
     *Biological Systems* (Princeton University Press, 2020;
     https://www.degruyter.com/document/doi/10.1515/9780691212920/html).

48. S. Verma, G. Novati, P. Koumoutsakos, Efficient collective swimming by harnessing vortices through deep
     reinforcement learning. *Proceedings of the National Academy of Sciences* **115**, 5849–5854 (2018).

49. H. Ko, G. Lauder, R. Nagpal, The role of hydrodynamics in collective motions of fish schools and bioinspired
     underwater robots. *Journal of The Royal Society Interface* **20**, 20230357 (2023).

50. D. N. Beal, F. S. Hover, M. S. Triantafyllou, J. C. Liao, G. V. Lauder, Passive propulsion in vortex wakes.
     *Journal of Fluid Mechanics* **549**, 385–402 (2006).

51. J. C. Liao, D. N. Beal, G. V. Lauder, M. S. Triantafyllou, The Kármán gait: novel body kinematics of rainbow
     trout swimming in a vortex street. *J Exp Biol* **206**, 1059–1073 (2003).

52. D. Quinn, G. Lauder, Tunable stiffness in fish robotics: mechanisms and advantages. *Bioinspir. Biomim.* **17**,
     011002 (2021).

53. G. V. Zuyev, V. V. Belyayev, An experimental study of the swimming of fish in groups as exemplified by the
     horsemackerel [Trachurus mediterraneus ponticus Aleev]. *Journal of ichthyology* **10**, 545–549 (1970).

54. S. Marras, S. S. Killen, J. Lindström, D. J. McKenzie, J. F. Steffensen, P. Domenici, Fish swimming in schools
     save energy regardless of their spatial position. *Behav Ecol Sociobiol* **69**, 219–226 (2015).

55. J. Herskin, J. F. Steffensen, Energy savings in sea bass swimming in a school: measurements of tail beat
     frequency and oxygen consumption at different swimming speeds. *Journal of Fish Biology* **53**, 366–376
     (1998).

56. P. G. Bushnell, J. F. Steffensen, K. Johansen, Oxygen Consumption and Swimming Performance in Hypoxia-
     Acclimated Rainbow Trout Salmo Gairdneri. *Journal of Experimental Biology* **113**, 225–235 (1984).

57. C. Sepulveda, K. A. Dickson, Maximum Sustainable Speeds and Cost of Swimming in Juvenile Kawakawa
Tuna (Euthynnus Affinis) and Chub Mackerel (Scomber Japonicus). *Journal of Experimental Biology* **203**,
3089–3101 (2000).

58. V. Di Santo, C. P. Kenaley, G. V. Lauder, High postural costs and anaerobic metabolism during swimming
support the hypothesis of a U-shaped metabolism–speed curve in fishes. *Proceedings of the National*
*Academy of Sciences* **114**, 13048–13053 (2017).

59. B. A. Block, D. Booth, F. G. Carey, Direct Measurement of Swimming Speeds and Depth of Blue Marlin.
*Journal of Experimental Biology* **166**, 267–284 (1992).

60. B. A. Block, S. L. H. Teo, A. Walli, A. Boustany, M. J. W. Stokesbury, C. J. Farwell, K. C. Weng, H. Dewar,
T. D. Williams, Electronic tagging and population structure of Atlantic bluefin tuna. *Nature* **434**, 1121–1127
(2005).

61. M. S. Triantafyllou, G. S. Triantafyllou, An Efficient Swimming Machine. *Scientific American* **272**, 64–70
(1995).

62. M. Kurt, J. Cochran-Carney, Q. Zhong, A. Mivehchi, D. B. Quinn, K. W. Moored, Swimming freely near the
ground leads to flow-mediated equilibrium altitudes. *Journal of Fluid Mechanics* **875**, R1 (2019).

63. G. Timmerhaus, C. C. Lazado, N. A. R. Cabillon, B. K. M. Reiten, L.-H. Johansen, The optimum velocity for
Atlantic salmon post-smolts in RAS is a compromise between muscle growth and fish welfare. *Aquaculture*
**532**, 736076 (2021).

64. P. He, C. S. Wardle, Tilting behaviour of the Atlantic mackerel, Scomber scombrus, at low swimming speeds.
*Journal of Fish Biology* **29**, 223–232 (1986).

65. K. Cathcart, S. Y. Shin, J. Milton, D. Ellerby, Field swimming performance of bluegill sunfish, Lepomis
macrochirus: implications for field activity cost estimates and laboratory measures of swimming performance.
*Ecology and Evolution* **7**, 8657–8666 (2017).

66. E. D. Gellman, T. R. Tandler, D. J. Ellerby, Swimming from coast to coast: a novel fixed-gear swimming gait
in fish. *Biology Letters* **15**, 20190270 (2019).

67. N. W. Xu, J. O. Dabiri, Low-power microelectronics embedded in live jellyfish enhance propulsion. *Science*
*Advances* **6**, eaaz3194 (2020).

68. K. Y. Lee, S.-J. Park, D. G. Matthews, S. L. Kim, C. A. Marquez, J. F. Zimmerman, H. A. M. Ardoña, A. G.
Kleber, G. V. Lauder, K. K. Parker, An autonomously swimming biohybrid fish designed with human cardiac
biophysics. *Science* **375**, 639–647 (2022).

69. M. B. S. Svendsen, P. G. Bushnell, J. F. Steffensen, Design and setup of intermittent-flow respirometry system
for aquatic organisms. *Journal of Fish Biology* **88**, 26–50 (2016).

70. Y. Zhang, "Interpreting species, intraspecific and intra-individual variability by comprehensively characterizing
a fish's respiratory phenotype with valid measures of oxygen uptake," thesis, University of British Columbia
(2021).

71. R. J. Kline, D. C. Parkyn, D. J. Murie, Empirical Modelling of Solid-blocking Effect in a Blazka Respirometer
for Gag, a Large Demersal Reef Fish. *Advances in Zoology and Botany* **3**, 193–202 (2015).

72. Y. Zhang, G. Claireaux, H. Takle, S. M. Jørgensen, A. P. Farrell, A three-phase excess post-exercise oxygen
consumption in Atlantic salmon Salmo salar and its response to exercise training. *J Fish Biol* **92**, 1385–1403
(2018).

73. H. F. Wootton, J. R. Morrongiello, T. Schmitt, A. Audzijonyte, Smaller adult fish size in warmer water is not
explained by elevated metabolism. *Ecology Letters* **25**, 1177–1188 (2022).

74. D. Chabot, J. F. Steffensen, A. P. Farrell, The determination of standard metabolic rate in fishes. *Journal of*
*Fish Biology* **88**, 81–121 (2016).

75. J. C. Svendsen, C. Tudorache, A. D. Jordan, J. F. Steffensen, K. Aarestrup, P. Domenici, Partition of aerobic
and anaerobic swimming costs related to gait transitions in a labriform swimmer. *Journal of Experimental*
*Biology* **213**, 2177–2183 (2010).

76. A. E. Brafield, D. J. Solomon, Oxy-calorific coefficients for animals respiring nitrogenous substrates.
*Comparative Biochemistry and Physiology Part A: Physiology* **43**, 837–841 (1972).

77. T. L. Hedrick, Software techniques for two- and three-dimensional kinematic measurements of biological and
biomimetic systems. *Bioinspir. Biomim.* **3**, 034001 (2008).

78. M. Saadat, F. Berlinger, A. Sheshmani, R. Nagpal, G. V. Lauder, H. Haj-Hariri, Hydrodynamic advantages of
in-line schooling. *Bioinspir Biomim* **16** (2021).

79. F. E. Fish, C. A. Hui, Dolphin swimming–a review. *Mammal Review* **21**, 181–195 (1991).

80. M. Daghooghi, I. Borazjani, The hydrodynamic advantages of synchronized swimming in a rectangular pattern.
*Bioinspir. Biomim.* **10**, 056018 (2015).

**Acknowledgments:** Many thanks to members of Lauder Laboratory for numerous discussions
about fish schooling behaviour, for comments on the manuscript, and to Cory Hahn for fish care.

**Funding:** Funding provided by the National Science Foundation grant 1830881 (GVL), the
Office of Naval Research grants N00014-21-1-2661 (GVL), N00014-16-1-2515 (GVL), 00014-
22-1-2616 (GVL), and a Postdoctoral Fellowship of Natural Sciences and Engineering Research
Council of Canada PDF - 557785 – 2021 (YZ).

**Author contributions:**

Y.Z. and G.L. conceptualized the study. Y.Z. performed experiments and data analyses and wrote
the manuscript. Y.Z. and G.L. provided manuscript edits and comments and approved the final
version.

**Competing interests:** Authors declare that they have no competing interests.

**Data and materials availability:** All data generated or analysed during this study are included
in this published article (and its supplementary information files).

**Supplementary Materials:**
Introductory text
Figs. S1 to S9
Tables S1 to S4
951
952

TABLE  HEADING

**Table 1. A summary of experimental studies that directly estimate the energetic saving of group movement in aerial and aquatic vertebrates that move through a fluid environment.** No energetic measurements have made for freely-moving bird flocks. Three studies measured the energetics of fish schools over a range of narrow speeds; two other studies measured the energetics of fish schooling at one speed.  But no studies have quantified *both* the aerobic and anaerobic energetic cost of swimming as a group compared to solitary locomotion.

| Animal group | Species | Testing speeds | Group size | Measurements | Results | Reference |
|---|---|---|---|---|---|---|
| Bird | *Pelecanus onocrotalus* | Voluntary flights | < 8 | Heart rate | Group movement saves energy | Weimerskirch et al.,2001 |
| Bird | *Columba livia* | Voluntary flights | 6 | Flap frequency & Accelerations | Group movement costs more energy | Usherwood et al.,2011 |
| Bird | *Geronticus eremita* | Voluntary flights | 14 | Wing beat phase, spatial distribution | V-formation birds use phasing strategy to cope with wakes | Portugal et al., 2014 |
| Ray-finned fish | 15 species | 1 speed, unknown speed | 6–12 individuals | Aerobic energy | Grouping reduces metabolic rate | Parker, 1973 |
| Ray-finned fish | *Notropis heterodon* | 1 speed ~1.2 BL s$^{-1}$ | 3 individuals | Aerobic energy | Only schooling of 6 cm fish saves energy | Abrahams & Colgan, 1985* |
| Ray-finned fish | *Anguilla anguilla* L | 6 speeds 1-2.2 BL s$^{-1}$ | 7 individuals | Aerobic energy | Schooling save energy at low speeds | Burgerhout et al.,2013* |
| Ray-finned fish | *Lepomis macrochirus* *Oncorhynchus mykiss* | 4 speeds, 1.5-3 BL s$^{-1}$ | 1-8 individuals | Aerobic energy | Schooling saves energy | Currier et al., 2020* |
| Ray-finned fish | *Salmo salar* | 6 speeds 1-3 BL s$^{-1}$ | 10 individuals | Aerobic energy | Schooling does not save energy | Hvas & Oppedal, 2019 |
| Ray-finned fish | *Devario aequipinnatus* | 14 speeds, 0.3–8 BL s$^{-1}$ | 8 individuals | Aerobic & anaerobic energy | Schooling saves energy at high speed | Present study* |

961    *Studies used group and individual fish in the same size swim-tunnel respirometer

962    †Other uncited studies of fish schoolings use kinematic calculations and computer simulation to indirectly show that fish schooling conserve energy

**Fig. 1. A summary of biomechanical principles underlying proposed hydrodynamic advantages of schooling behaviour in fish.** When fish swim into free-stream flow ($U_{fs}$), experimental data show that fish schools are dynamic with fish changing position frequently. Regardless of fish position within a school and changes in relative position, theoretical and robotic analyses have demonstrated that at least four mechanisms (indicated by numbers) provide an advantage in the form of reduced power consumption. 1. Reduced oncoming velocity ($U_2$) requires less thrust for a fish swimming in the wake between two leading fish (*6*); 2. The Knoller-Betz effect of leading edge suction reduces costs for a trailing fish due to accelerated oscillating flow at the head ($U_1$) (*24*) (*78*); 3. Added mass "push" from follower to leader can reduce costs for the leader in front of another fish (*79*) (*23*) (*78*); 4. Wall effects benefit neighbouring fish where swimming next to another fish reduces swimming costs (*80*) (*20*). These principles suggest that regardless of the relative positions of the individuals within the fish school, the fish school as a collective unit should be able to save metabolic energy compared to a solitary fish swimming in $U_{fs}$.

**Fig. 2. Measurements of aerobic and anaerobic locomotor cost of fish schools and solitary fish.** (**A**) Average traces of metabolic rate ($\dot{M}O_2$) of fish schools over a 40-hour experiment. Following the first 18-h quiescent state, a critical swimming speed ($U_{crit}$) test quantifies the aerobic cost of active swimming. The ensuing 18-h measurement of excess post-exercise oxygen consumption (EPOC) quantifies the anaerobic cost. (**B**) Comparison of $\dot{M}O_2$ for conditions of aggregating behaviour, minimum demand speed, and resting condition with minimal flow ($\dot{M}O_{2aggregate}$, $\dot{M}O_{2min}$, $\dot{M}O_{2rest}$) (**C**) Comparisons of concave upward shaped $\dot{M}O_2$-speed curve over the entire range (0.3–8 body length s$^{-1}$, BL s$^{-1}$) and (**D**) the concave shaped $\dot{M}O_2$-speed curve at the lower speeds (0.3–3 BL s$^{-1}$). (**E, F**) Percentage (%) aerobic scope used by fish schools and solitary fish during the $U_{crit}$ test. (**G, H**) Comparisons of EPOC and EPOC durations between fish schools and solitary fish. Statistical significance is denoted by asterisk(s). Green colour = school data (n=5 schools); blue colour = solitary fish data (n=5 individual fish); shading indicates the 95% confidence interval. Statistical details are available in the statistical analyses section.

| Non-aerobic contribution: | 18 | 29 | 39 | 51 | 58% |
| Aerobic contribution: | 82 | 71 | 61 | 49 | 42% |

| Non-aerobic contribution: | 45 | 62 | 71 | 77 | 81% |
| Aerobic contribution: | 55 | 38 | 29 | 23 | 19% |

**Fig. 3. Modeling of simultaneous aerobic and non-aerobic costs of fish schools and solitary**

**fish for a critical swimming speed ($U_{crit}$) test. (A)** Modeling the $O_2$ cost of the metabolic rate

($\dot{M}O_2$)-speed curve and the ensuing recovery cost (excess post-exercise oxygen consumption,

EPOC) as a function of speed. After $U_{crit}$, fish returned to the same resting $\dot{M}O_2$ ($\dot{M}O_{2rest}$) as a

pre-test. **(B, C)** In addition to $\dot{M}O_2$ (solid line & filled symbols), we modeled the total $O_2$ cost

(dash line & half-filled symbols) for fish schools and solitary fish and when performing the $U_{crit}$

swimming test. The estimated partitioning of aerobic and non-aerobic contributions to swimming

are denoted (red-&-bold) with respect to speed for 4–8 BL s$^{-1}$ is shown below each graph. **(D, E)**

Using total $O_2$ cost, we computed total energy expenditure (10-min period per point) and the

total cost of transport (including both aerobic metabolism, high-energy phosphate, and anaerobic

glycolysis) for both fish schools and solitary fish. Statistical significance is denoted by

asterisk(s). Green colour = school data (n=5); blue colour = solitary fish data (n=5); shading

indicates the 95% confidence interval. See methods for modeling and statistical details.

1012

**Fig. 4. Three-dimensional characterization of swimming kinematics and fish schooling dynamics as a function of speed. (A)** Total energy expenditure (TEE) per tail beat, **(B)** Tail beat frequency, **(C)** The angle of fish to free-stream water flow, measured as the mean and the S.D. of the angles of the individuals within the school. **(D)** Three-dimensional angle of fish to the frontal plane. **(E)** Turning frequency, **(F)** Three-dimensional school length. **(G)** Three-dimensional distances among all individuals in the school and the S.D. of the distance. The upper and lower boundaries of the metrics are indicated. Statistical significance is denoted by asterisk(s). Green colour = school data (n=3-4); blue colour = solitary fish data (n=3-4); shading indicates the 95% confidence interval.  See methods for details of three-dimensional reconstruction and statistics.