1  **Longitudinal map of transcriptome changes in the Lyme pathogen *Borrelia burgdorferi***
2  **during tick-borne transmission**
3

4  Anne L. Sapiro[1*], Beth M. Hayes[1], Regan F. Volk[2], Jenny Y. Zhang[1], Diane M. Brooks[3], Calla
5  Martyn[1], Atanas Radkov[1], Ziyi Zhao[1], Margie Kinnersley[3], Patrick R. Secor[3], Balyn W. Zaro[2],
6  Seemay Chou[1*]
7
8  1 Department of Biochemistry & Biophysics, University of California, San Francisco, California,
9  USA
10  2 Department of Pharmaceutical Chemistry and Cardiovascular Research Institute, University of
11  California, San Francisco, California, USA
12  3 Division of Biological Sciences, University of Montana, Missoula, Montana, USA
13
14  *Corresponding authors:
15  Anne L. Sapiro: annesapiro@gmail.com & Seemay Chou: seemaychou@gmail.com

16
**ABSTRACT**

*Borrelia burgdorferi* (*Bb*), the causative agent of Lyme disease, must adapt to vastly different
environments as the bacterium cycles between the tick vector and a vertebrate host. During a
bloodmeal, *Bb* migrates from the tick midgut to the salivary glands and changes its gene
expression, priming *Bb* for vertebrate infection. These tick-specific transmission events are
dependent on key transcriptional regulators; however, the full range of transcriptional changes
that occur over several days inside of the tick are technically challenging to capture. We
developed an experimental approach to enrich *Bb* cells from *Ixodes* ticks during a transmitting
bloodmeal to longitudinally define their global transcriptomic landscape. We identified 192 genes
that substantially change expression over the course of the tick bloodmeal, most of which were
located on plasmids of the *Bb* genome. The majority of upregulated genes encode proteins found
at the cell envelope or proteins of unknown function, including 45 outer surface lipoproteins
embedded in the unusual protein-rich coat of *Bb*. As these proteins may facilitate *Bb* interactions
with host tissues or immune systems, we also utilized mass spectrometry to identify candidate
tick proteins that physically associate with *Bb*. The *ex vivo Bb* transcriptomes and candidate tick
interacting proteins presented here provide an important roadmap for investigating key
determinants of *Bb* priming and transmission during the tick stage of its unique transmission cycle.

**INTRODUCTION**

Vector-borne microbial pathogens are transmitted by the bite of arthropods and have evolved sophisticated ways to adapt to vastly different environments as they move between vector and host. Uncovering their adaptive mechanisms can not only open avenues for disrupting pathogen transmission but also provide fundamental insights into vector physiology and microbial symbioses (Shaw and Catteruccia, 2018). Lyme disease, the most reported vector-borne disease in North America, is caused by the bacterial pathogen *Borrelia burgdorferi* (*Bb*) (Rosenberg et al., 2018; Steere et al., 2016)*.* Its primary vector, the blacklegged tick *Ixodes scapularis,* acquires and transmits *Bb* through two separate multi-day bloodmeals, one in which *Bb* is acquired from an infected vertebrate host and a second, during the subsequent life stage, in which *Bb* is transmitted to a new host (Tilly et al., 2008). During the transmission bloodmeal, *Bb* proliferate in the tick midgut before a subset of these cells disseminate to the salivary glands (Dunham-Ems et al., 2009). *Bb* is deposited into the new host via the tick saliva extruded into the bite site (Spielman et al., 1987). Given the prolonged nature of *I. scapularis* feeding, the high specificity of vector-pathogen relationships, and the complicated array of events needed for successful *Bb* transmission, the tick bloodmeal provides an opportune intervention point for preventing pathogen spread. However, we do not currently have a clear understanding of the molecular mechanisms involved in this process.

*Bb* must adapt to dramatically different environments as it cycles from tick to vertebrate host, and understanding the genes involved in this process will enable the identification of key interactions to target to prevent transmission. When an infected tick feeds, *Bb* responds to bloodmeal-induced environmental changes and undergoes cellular modifications driven by key transcriptional circuits, including the RpoN-RpoS sigma factor cascade and the Hk1/Rrp1 two-component system (Radolf et al., 2012). *In vitro* analyses of *Bb* cells cultured in tick- or mammalian-like growth conditions have pointed to additional genetic determinants of tick-borne transmission by revealing widespread transcriptome remodeling during host switching (reviewed in Samuels et al., 2021). However, it is not fully clear how these *in vitro* expression changes correspond to complex *in vivo* changes over the course of a transmission bloodmeal, which comprises several physiologically distinct stages for the tick vector. Capturing comprehensive, longitudinal data on *Bb* expression from inside its vector has been hampered by technical challenges given the dynamic nature of the bloodmeal and the general low abundance of bacterial cells relative to the tick (Samuels et al., 2021). Although microarray approaches have been used to identify large scale changes in *Bb* gene expression between the first and second tick bloodmeals (Iyer et al., 2015), we have limited

temporal resolution into the molecular transitions that happen across key steps of tick feeding.

This problem necessitates novel approaches to capture the transcriptomic changes of *Bb* within

the natural tick environment.

While the full landscape of *Bb* transmission determinants is not yet known, we do have a growing

knowledge of functional processes that are critical during the tick stage, such as motility,

metabolism, and immune evasion (Kurokawa et al., 2020; Phelan et al., 2019). These functions

often rely on the unique protein-rich *Bb* outer surface. Notably, several specific tick–*Bb* protein–

protein interactions are important for *Bb* survival, migration, or transmission to the next host. *Bb*

encodes an extensive <u>o</u>uter <u>s</u>urface <u>p</u>rotein (Osp) family with members that are differentially

expressed during host switching. One of these proteins, OspA, binds a tick cell surface protein,

<u>t</u>ick <u>r</u>eceptor for <u>OspA</u> (TROSPA), which is required for successful *Bb* colonization of the tick

midgut during the first acquisition bloodmeal (Pal et al., 2004). Several other proteins have also

been linked to *Bb* migration within the tick (Pal et al., 2022). For example, BBE31 binds a tick

protein TRE31, and disruption of this interaction decreases the number of *Bb* cells that

successfully migrate from the tick gut to salivary glands (Zhang et al., 2011). However, these

interactions alone are not sufficient to block *Bb* growth or migration in ticks, suggesting there are

likely additional molecular factors from *Bb* and ticks at play during tick-borne transmission.

To provide a more comprehensive set of *Bb* determinants driving tick-borne transmission of this

important human pathogen, we longitudinally mapped *Bb* genome-wide expression changes for

bacterial cells isolated from inside ticks during the transmission bloodmeal. We developed a novel

sequencing-based strategy for *ex vivo* transcriptomic profiling of *Bb* populations within feeding

ticks. We identified 192 highly differentially expressed genes, including both genes previously

implicated in *Bb* transmission as well as many others. Genes upregulated during tick transmission

were enriched for outer surface lipoproteins, suggesting *Bb* dramatically remodels its cell

envelope as it migrates through the tick. Mass spectrometry analyses revealed dramatic changes

in the tick environment over feeding, identifying new potential determinants of a more extensive

and diverse set of tick–microbe molecular interactions than previously appreciated.

**RESULTS**

**A two-step enrichment process facilitates robust transcriptional profiling of *Bb* during the**

**tick bloodmeal**

To gain a more comprehensive understanding of *Bb* gene expression throughout the tick phase

of the transmission cycle, we developed an experimental approach to characterize the *Bb*

transcriptome of spirochetes isolated from ticks during a bloodmeal. We aimed to establish a

longitudinal transcriptional profile encompassing key pathogen transmission events each day of

feeding after ticks attached to their mouse bloodmeal hosts (**Figure 1A**). The major bottleneck for

such an RNA sequencing (RNA-seq) approach is capturing sufficient quantities of *Bb* transcripts

from complex multi-organism samples in which pathogen transcripts represent a very small

minority. Our initial attempts to uncover *Bb* mRNA by simply removing tick mRNA with polyA-

depletion and removing tick rRNA sequences using Depletion of Abundant Sequences by

Hybridization (DASH) (Dynerman et al., 2020; Gu et al., 2016) were unsuccessful. This approach

resulted in an average of only 0.09% of RNA-seq reads mapping to *Bb* mRNA – approximately

10-fold less than we estimated would be needed to feasibly obtain robust transcriptome-wide

gene expression analysis (Haas et al., 2012).

To dramatically increase *Bb* transcript representation in our libraries, we physically enriched *Bb*

cells in tick lysates prior to library preparation by adding an initial step of immunomagnetic

separation (**Figure 1B**). We took advantage of a commercial antibody previously generated

against whole *Bb* cells (α*Bb*, RRID: AB_1016668). By western blot analysis, we confirmed that

α*Bb* specifically recognized several *Bb* proteins, including surface protein OspA, which is highly

prevalent on the *Bb* surface in ticks (**Figure S1A**). In addition, immunofluorescence microscopy

with α*Bb* showed clear recognition of *Bb* cells from within the tick at each day of feeding (**Figure**

**S1B**). After collecting infected

ticks from mice one, two, three,

and four days post-attachment,

we used α*Bb* and magnetic

beads to enrich *Bb* cells away

from the majority of the tick

material in the lysates. We

tracked relative *Bb* enrichment

through RT-qPCR of *Bb flaB*

RNA and tick *gapdh* RNA in the

separated samples. Measuring

*Bb flaB* RNA from both *Bb*-

enriched samples and their

**Figure S1. Anti-*Bb* antibody recognizes ospA and binds *Bb* in the tick throughout the bloodmeal.** (**A**) Western blot with anti-*Bb* on lysate from cultured *Bb*: wildtype (A3, left), a mutant lacking ospA (ospA1), and the mutant with ospA restored (ospA+B1). Anti-*Bb* recognizes ospA among other proteins. (**B**) Immunofluorescence microscopy with anti-*Bb* (green) and propidium iodide (PI) (DNA, red) on each day of feeding (yellow is merge). Anti-*Bb* antibody recognizes *Bb* in the tick across the bloodmeal.

137 matched *Bb*-depleted fractions, we found over 95% of total *Bb flaB* RNA was present in enriched

138 fractions (**Figure 1C**), suggesting our approach captured the vast majority of *Bb* transcripts from

139 the tick.

**Figure 1. A two-step enrichment process facilitates robust transcriptional profiling of *Bb* during the tick bloodmeal.** (**A**) Schematic of *Bb* during tick feeding. *Bb* in the nymphal tick midgut respond to the bloodmeal by multiplying and changing their transcriptional state. At the same time, the tick gut undergoes numerous changes to digest the bloodmeal. After two to three days of feeding, a small number of *Bb* leave the midgut and enter the salivary glands (blue), while the majority are left behind in the gut after engorgement. (**B**) Schematic of *Bb* enrichment process from feeding ticks. Whole ticks are dissociated, anti-*Bb* antibodies are added to lysates, and antibodies and *Bb* are captured magnetically. RNA is extracted and RNA-seq libraries are prepared. DASH is then used to remove rRNA before sequencing. This process increases *Bb* reads in the resulting sequencing. (**C**) RT-qPCR results showing the percentage of *Bb flaB* and tick *gapdh* RNA in the enriched versus depleted fractions after the enrichment process. Data come from 4 replicates each from day 2, day 3, and day 4, mean +/- SE. Nearly all *Bb flaB* RNA was found in the enriched fraction, while the majority of tick *gapdh* was found in the depleted fraction. (**D**) The percentage of reads mapping to rRNA before and after DASH. Data are shown as mean +/- SD. rRNA reads are drastically reduced after DASH. (**E**) The percentage of reads in RNA-seq libraries mapping to *Bb*. *Bb* mRNA reads make up a small but sufficient proportion of libraries for transcriptomics. Data are shown as mean +/- SD. (**F**) The number of reads in millions (M) mapped to *Bb* for each day. Data are shown as mean +/- SD. An average of 4.3 million reads per sample mapped to Bb genes, covering 92% of genes with at least 10 reads.

Total RNA recovered from the enrichment process was used to create RNA-seq libraries that subsequently underwent depletion of rRNA sequences from ticks, mouse, and *Bb* using DASH, which targets unwanted sequences for degradation by Cas9 (Gu et al., 2016; Ring et al., 2022). DASH reduced unwanted sequences from 94% to 9% of our total libraries, greatly increasing the relative abundance of *Bb* transcripts (**Figure 1D**). For resulting libraries generated across feeding (**Table S1**), between 0.6% and 3.4% of reads mapped to *Bb* coding regions (**Figure 1E**), which translated to an average of 4.3 million *Bb* reads per sample (**Figure 1F**). Read coverage was sufficient for downstream differential expression analyses for the vast majority of *Bb* genes, with at least 10 reads mapping to 92% of *Bb* genes. To evaluate whether our approach introduced any major artifacts, we sequenced and compared RNA-seq libraries from *in vitro* cultured *Bb* cells before and after immunomagnetic enrichment. We found minimal expression differences (29 genes with p<0.05, fold changes between 0.83-1.12) (**Figure S2, Table S2**), suggesting experimental enrichment did not significantly alter global transcriptome profiles for *Bb*. Thus, we developed an approach to enable genome-wide analysis of *Bb* population-level expression changes that occur within the feeding tick during transmission.

**Figure S2. Enrichment process does not induce large scale gene expression changes in *in vitro* cultured *Bb*.** Log$_2$ fold changes vs. mean normalized number of counts comparing cultured *Bb* input and samples after enrichment with anti-*Bb*. n=3. Red dots, p-value < 0.05, Wald tests. The gene expression changes induced during processing are much smaller than those observed between days of feeding.

**Global *ex vivo* profiling of *Bb* reveals extent and kinetics of transcriptional changes**

To provide a broad overview of *Bb* expression changes during feeding, we performed principal component analysis (PCA) on the *Bb* transcriptome data from one, two, three, and four days post-attachment (n=4). We reasoned that if many longitudinal expression changes were occurring across *Bb* populations, we would observe greater data variability between time points than between biological replicates. Indeed, we found replicates from each day grouped together, whereas distinct time points were largely non-overlapping. The first principal component, which explained 64% of the variance in our data, correlated well with day of feeding (**Figure 2A**). The global pattern suggested that *Bb* gene expression changes generally trended in the same

direction over the course of feeding with the most dramatic differences between flanking

timepoints on day 1 and day 4.

Using day 1 (early attachment) as a baseline, we performed differential expression analysis for

all *Bb* genes at subsequent time points and examined changes above a twofold threshold (**Figure**

**2B, Table S3**). These analyses mirrored the global longitudinal expression pattern predicted by

our PCA. The total number of differentially expressed (DE) genes increased steadily over time

and peaked at day 4, which when compared to day 1 had 186 DE genes, including 153

upregulated and 33 downregulated (**Figure 2C-E**). Across time points, DE genes were highly

overlapping and largely changed in the same directions. In all three comparisons, we found 192

DE genes in total (**Table S4**). However, there were some notable differences between genes,

such as the overall timing and kinetics of expression changes. Transcript levels for some DE

**Figure 2. Global *ex vivo* profiling of *Bb* reveals extent and kinetics of transcriptional changes.** (**A**) Principal component analysis of samples from across feeding. PC1 correlates strongly with day of feeding. (**B**) Schematic depicting how data was analyzed, as pairwise comparisons between the first day after attachment and all other days. (**C-E**) Volcano plots of differentially expressed genes of day 2 versus day 1 after attachment (**C**), day 3 versus day 1 (**D**), and day 4 versus day 1 (**E**). The total number of upregulated genes is shown in the top right and downregulated genes is shown in the top left. Yellow dots are genes that change expression between day 1 and day 2, red dots are genes that change expression between day 1 and day 3, and purple dots are genes that change expression between day 1 and day 4. Two genes with $\log_2$ fold changes >4 are shown at 4, and five genes with $-\log_{10}$(padj) >60 are shown at 60. Only genes with at least a 2-fold change are highlighted. p-value < 0.05, Wald tests. By day 4 of feeding, 153 genes are upregulated and 33 genes are downregulated from day 1 baseline levels.

genes changed suddenly over the course of feeding, while others were more gradual. To our knowledge, this is the first comprehensive report of global *Bb* expression changes within feeding ticks.

To help assess the integrity of our dataset, we examined expression profiles for previously characterized targets of major transcriptional programs activated at the onset of the bloodmeal, the Hk1/Rrp1 two-component system and the RpoN/RpoS sigma factor cascade (Caimano et al., 2019, 2015). Between day 1 and day 4, the majority of genes previously reported to be activated or repressed by Rrp1 *in vitro* (Caimano et al., 2015) trended significantly in the expected direction *ex vivo* (111/148 upregulated, 37/57 downregulated, $p<0.05$, Wald tests, **Figure S3A**). The majority of the genes previously reported to be activated by RpoS in a mammalian context, including canonical targets *ospC* and *dbpA*, were also significantly upregulated during tick feeding (33/55, $p<0.05$, Wald tests) (**Figure S3B**), along with *rpoS* itself (**Figure S3C**), as expected (Hübner et al., 2001). However, very few genes repressed by RpoS in mammals were similarly repressed in ticks (2/33, $p<0.05$, Wald tests). These data support the hypothesis that while RpoS-dependent gene activation occurs across both vertebrate and vector stages of *Bb* transmission, RpoS-dependent repression is specific to spirochetes infecting a mammalian host (Caimano et

**Figure S3.** *Ex vivo* **RNA-seq corroborates key transcriptional programs in the tick.** (**A**) Volcano plot of DE genes comparing day 4 to day 1, with Rrp1-upregulated (blue) and downregulated (pink) genes. Rrp1-regulated genes correlate well with up and downregulated genes during feeding. (**B**) Volcano plot of DE genes comparing day 4 to day 1, with RpoS-upregulated (blue) and downregulated (pink) genes. Genes upregulated by RpoS in mammals increase during feeding, but genes repressed by RpoS in mammals are not repressed in the tick. (**C**) Tukey style boxplot of Transcripts Per Million (TPM) on each day for *rpoS*. Black dots represent replicates. ****$p$-value $< 0.00001$, Wald test. *rpoS* expression increases over the course of feeding. (**D**) Volcano plot of DE genes comparing day 4 to day 1, with Rel$_{Bbu}$-upregulated (blue) and downregulated (pink) genes. About half of Rel$_{Bbu}$ genes change in the expected direction over feeding.

al., 2019, 2007). We also compared our DE genes to those regulated by Rel$_{Bbu}$ as part of the

stringent response, another major transcriptional program active in *Bb* in the tick during nutrient

starvation (Drecktrah et al., 2015). About half of Rel$_{Bbu}$-regulated genes changed in the direction

expected if the stringent response was active during this time (**Figure S3D**); however, more of

the genes that were downregulated over feeding have been reported to be regulated by Rel$_{Bbu}$

than either RpoS or Rrp1, suggesting it may play some role during feeding.

By looking at these previously reported transcriptional regulons in our data, we were able to verify

that our data captured expected transcriptional programs active during tick feeding. Nevertheless,

25% of the twofold DE genes were not previously identified through these RNA-seq studies as

dependent upon RpoS, Rrp1, or Rel$_{Bbu}$ (Caimano et al., 2019, 2015; Drecktrah et al., 2015), which

are the three main *Bb* regulatory programs active in the tick (Samuels et al., 2021). These

additional genes highlight the necessity of measuring transcription in the tick environment and

suggest we uncovered gene expression changes specific to the tick-stage of the *Bb* enzootic

cycle. The nature and dynamics of these changes provide insights into potential genetic

determinants of *Bb* survival, proliferation, and dissemination in the tick during transmission.

***Bb* genes upregulated during feeding are found predominantly on plasmids**

*Bb* has a complex, highly fragmented genome (Barbour, 1988) (**Figure 3A**), including numerous

plasmids that are necessary during specific stages of the enzootic cycle (reviewed in Schwartz et

al., 2021) suggesting they contain genes that are crucial for pathogen transmission and survival.

In fact, many genes found on the plasmids have been previously shown to alter expression upon

environmental changes or in different host environments (Iyer et al., 2015; Ojaimi et al., 2005;

Revel et al., 2002; Tokarz et al., 2004). Thus, we reasoned that many of the 192 *Bb* genes that

change expression over the course of the tick bloodmeal would reside on the plasmids, and we

examined their distribution throughout the genome. Consistent with these previous reports, we

found that most of the upregulated genes were located on the plasmids (143/158; 90%), while

fewer were found on the chromosome (15/158; 10%) (**Figure 3B**), which is home to the majority

of metabolic and other housekeeping genes. In contrast, the majority of the downregulated genes

were found on the chromosome (27/34, 79%) (**Fig 3C**).

Several plasmid-encoded genes that were longitudinally upregulated in our datasets have known

roles during the tick bloodmeal or in mammalian infection. For example, linear plasmid 54 (lp54),

**Figure 3. *Bb* genes upregulated during feeding are found predominantly on plasmids.** (**A**) Schematic of the chromosome and plasmids in the *Bb* B31-S9 genome. (**B-C**) The number of genes from each chromosome or plasmid that increased (**B**) or decreased expression (**C**) twofold during feeding. Upregulated genes are distributed across plasmids, while downregulated genes are found on the chromosome and lp54.

which is an essential plasmid present in all *Bb* isolates (Casjens et al., 2012), contained the largest

number of upregulated genes. Many of the genes on lp54 are known to be regulated by RpoS

during feeding, including those encoding adhesins DbpA and DbpB, which are important for

infectivity in the host (Blevins et al., 2008). This set also included five members of a paralogous

family of outer surface lipoproteins BBA64, BBA65, BBA66, BBA71, and BBA73. BBA64 and

BBA66 are necessary for optimal transmission via the tick bite (Gilmore et al., 2010; Patton et al.,

2013). These findings indicate our dataset captures key *Bb* transcriptional responses known to

be important for survival inside the tick during a bloodmeal.

Many upregulated genes were encoded by cp32 plasmid prophages. *Bb* strain B31-S9 harbors

seven cp32 isoforms that are highly similar to each other (Casjens et al., 2012). When cp32

prophages are induced, phage virions called φBB1 are produced (Eggers and Samuels, 1999). In

addition to phage structural genes, cp32 contain loci that encode various families of paralogous

outer surface proteins (Stevenson et al., 2000). Amongst the cp32 genes that increased over

feeding were members of the RevA, Erp, and Mlp families, which are known to increase

expression during the bloodmeal (Gilmore et al., 2001). We also found several phage genes that

were upregulated, including those encoding proteins annotated as phage terminases on cp32-3,

cp32-4, and cp32-7 (BBS45, BBR45, and BBO44). Some cp32 genes have been shown to

change expression in response to temperature changes (Tokarz et al., 2004) and as a part of the

stringent response regulated by Rel$_{Bbu}$ (Drecktrah et al., 2015). Our data suggest that some

prophage genes are upregulated over the course of tick feeding, raising the possibility that cp32
prophage are induced towards the end of feeding. Overall, our data support the long-held idea
that the *Bb* plasmids, which house many genes encoding cell envelope proteins, proteins of
unknown function, and prophage genes, play a critical role in the enzootic cycle during the key
transition period of tick feeding.

**_Bb_ genes encoding outer surface proteins are enriched among upregulated genes**

To gain a better overall sense of the types of genes that changed over feeding and the timing of
those changes, we grouped DE genes into functional categories. Since a high proportion of
plasmid genes encode lipoproteins within the unique protein-rich outer surface of *Bb*, genes of
unknown function, and predicted prophage genes (Casjens et al., 2000; Fraser et al., 1997), we
expected that many of the DE genes would fall into these categories. We classified the genes as
related to either: cell envelope, bacteriophage, cell division, DNA replication and repair,
chemotaxis and motility, metabolism, transporter proteins, transcription, translation, stress
response, protein degradation, or unknown (as in Drecktrah et al., 2015)(see **Table S4**).

Of the genes that increased at least twofold over feeding, the clear majority each day of feeding
fell into two broad categories, cell envelope (55 of 158 total genes) and proteins of unknown
function (69 of 158 total genes), with fewer genes related to metabolism, chemotaxis and motility,
transporters, bacteriophage, cell division, and transcription (**Figure 4A**). In contrast to the
upregulated genes, genes downregulated during feeding were more evenly distributed among the
functional categories – including translation, protein degradation, transcription, and metabolism –
consistent with many of them being located on the chromosome.

When looking at changes across these functional categories, the overrepresentation of cell
envelope proteins was striking, while not unexpected. The *Bb* outer surface is covered with
lipoproteins (**Figure 4B**), and these proteins are critical determinants in *Bb* interactions with the
various environments encountered during the enzootic cycle (Kurokawa et al., 2020). We found
that more than half of these outer surface lipoproteins changed expression during this time, with
46 of 83 total outer surface lipoproteins (Dowdell et al., 2017) differentially expressed twofold over
feeding (**Figure 4C**). These data suggested widespread changes may be occurring on the *Bb*
outer surface during feeding.

**Figure 4.** *Bb* **genes encoding outer surface proteins are enriched among upregulated genes.** (**A**) The number of *Bb* genes that change over the course of tick feeding sorted into functional categories. Genes that first change 2 days after attachment are shown in yellow, 3 days after attachment in red, and 4 days after attachment in purple. A majority of upregulated genes fall into cell envelope and unknown categories. (**B**) Schematic of the outer membrane of *Bb* showing the location of outer surface lipoproteins. (**C**) Heat map of the average Transcripts Per Million (TPM) of all genes encoding outer surface lipoproteins across the 4 days of tick feeding. Genes in blue were twofold upregulated and genes in pink twofold downregulated over feeding. A majority of genes encoding outer surface proteins increased in expression throughout feeding, while having different magnitudes of expression.

To understand the functional implications of expression changes in a majority of outer surface

lipoproteins, we also compared their relative expression. The magnitude of expression varied

greatly, with *ospA*, *ospB*, *ospC*, and *bba59* being the most highly expressed outer surface protein

transcripts. Many of the outer surface protein genes that we found increased expression over

feeding were much less abundantly expressed (**Table S5**). However, even the genes that

appeared to have low expression in these population level measurements could play important

roles in transmission if they are highly expressed in a small number of crucial cells, such as those

that ultimately escape the midgut. While bulk RNA-seq cannot distinguish what is happening at

the single cell level, our data suggest that during the bloodmeal, *Bb* are undergoing a complex

outer surface transformation driven by increases in a majority of the genes encoding these crucial

lipoproteins.

**Identification of candidate tick interaction partners of *Bb* cells *ex vivo***

Our RNA-seq data suggested that the outer surface of *Bb* transforms over the course of feeding

as *Bb* are primed for transmission to a vertebrate host. At the same time, the tick midgut

environment is changing as the tick begins to digest its bloodmeal. Tick-*Bb* interactions are likely

crucial at the beginning of the tick bloodmeal, as *Bb* adhere to the tick gut epithelium before

becoming motile and migrating out of the midgut and into the salivary glands (Dunham-Ems et

al., 2009). Some *Bb* outer surface proteins, such as BBE31 and BBA52 play key roles in pathogen

migration through interactions with the tick environment (Kurokawa et al., 2020). We wanted to

explore the changing tick environment to identify tick proteins with which *Bb* could interact

throughout the tick bloodmeal. Since our *Bb* enrichment process retained some tick material, we

reasoned that tick proteins that interact with *Bb* would be present in these samples.

To outline the changes occurring in the tick during feeding and to identify candidate *Bb*-interacting

tick proteins, we used mass spectrometry to survey the content of the tick material that was

enriched along with the *Bb* cells we sequenced at early and late stages during feeding. We purified

proteins from the α*Bb*-enriched fraction of crushed infected ticks on day 1 after attachment and

day 4 after attachment in triplicate. As controls for each day, we also performed the *Bb* enrichment

process on lysate from uninfected ticks mixed with *in vitro* cultured *Bb* to help rule out proteins

**Figure 5. Identification of candidate tick interaction partners of *Bb* cells *ex vivo*.** (**A**) Schematic of experiment to determine candidate tick proteins interacting with *Bb* over the course of feeding. Ticks were collected 1 day after attachment and 4 days after attachment. Uninfected ticks at the same time points were mixed with cultured *Bb* as controls. *Bb* was enriched with anti-*Bb* antibody as in RNA-seq experiments and then subjected to mass spectrometry to identify tick proteins present in the samples. Venn diagram depicts the proteins enriched in day 1 and day 4 samples over controls. Tick proteins that are enriched with *Bb* vary greatly over the course of feeding. (**B**) Tick proteins uniquely identified one day after attachment that are annotated as extracellular matrix (ECM) proteins. (**C**) Tick proteins uniquely identified four days after attachment that are annotated as low-density lipoprotein receptors. ECM and membrane proteins may be good candidates for *Bb*-interacting proteins.

that were not pulled down through *in vivo* tick-*Bb* interactions (**Figure 5A**). We identified between
414 and 2240 protein groups per sample replicate. To identify proteins of interest, we looked for
those that were detected in at least two of three replicates with infected ticks and had a mean
average coverage twice that of uninfected ticks mixed with cultured *Bb*. We found 256 proteins
that were enriched with *Bb* from infected ticks one day after attachment and 226 proteins that
were enriched with *Bb* from infected ticks four days after attachment (**Table S6**). Of these proteins,
only 27 were detectable on both days, suggesting the tick proteins present upon *Bb* enrichment
change dramatically over the course of feeding. The vast majority of all detected proteins were
from ticks (98%). Amongst the small number of *Bb* proteins, we identified OspC in the samples
from day 4 after attachment but not day 1 after attachment, confirming the expression change we
saw in our RNA-seq. The distinct sets of tick proteins we identified at each timepoint suggest
dramatic changes occur in the *Bb*-infected tick midgut environment during feeding that may alter
the landscape of tick-*Bb* interactions.

Some of the proteins enriched with *Bb* from the tick may be good candidates for key *Bb*-interacting
partners during feeding, especially if they are localized to the surface of tick cells where they may
encounter *Bb*. The scarcity of both predicted and experimentally validated functions and
localizations for tick proteins makes it difficult to fully assess the potential for tick protein
interactions with extracellular *Bb*. Nevertheless, of the proteins found exclusively one day after
attachment, 10 were categorized as extracellular matrix proteins using the PANTHER gene
database (Thomas et al., 2022), and this category was statistically enriched (Fisher's exact test,
FDR=0.000093) (**Figure 5B, Table S7**). 30 additional proteins were annotated with a cellular
component as plasma membrane. Four days after attachment, we did not detect any annotated
extracellular matrix proteins; however, we identified 31 proteins that are likely to be found at the
membrane, including two proteins annotated as putative low-density lipoprotein receptors (**Figure
5C, Table S8**). These extracellular matrix and membrane proteins may be the most likely to
directly interact with *Bb* during this timeframe and are candidates for tick proteins important in the
*Bb* dissemination process. The proteins present in the changing tick environment may be key
determinants of pathogen transmission as *Bb* remodels its outer surface while preparing to
migrate through the tick to a new host.

**DISCUSSION**
Vector-borne pathogens must adapt to distinct environments as they are transmitted by
arthropods to colonize new bloodmeal hosts. The tick-borne Lyme disease pathogen *Bb*

undergoes transcriptional changes inside its vector over the course of a single bloodmeal that are
important for a number of key transmission events. For example, expression changes enable
survival in the feeding tick (He et al., 2011), facilitate transmission across several internal
compartments into the bloodmeal host (Kurokawa et al., 2020), and prime bacteria for successful
infection of the next host (Kasumba et al., 2016). Capturing these changes as they occur *in vivo*
has been challenging due to the low relative abundance of *Bb* material inside of the rapidly
growing, blood-filled tick. To date, many advances in our understanding of *Bb* expression during
transmission have come from tracking changes in small subsets of genes during tick feeding
(Gilmore et al., 2001), leveraging *Bb* culture conditions that approximate environmental changes
across its lifecycle (Ojaimi et al., 2005; Revel et al., 2002; Tokarz et al., 2004), and defining
transcriptional regulons outside of the tick (Caimano et al., 2019, 2015; Drecktrah et al., 2015).
Here, we developed an experimental RNA-seq-based strategy to more directly and longitudinally
profile gene expression for *Bb* populations isolated from ticks over the course of a transmitting
bloodmeal.

Our longitudinal collection of transcriptomes for *Bb* cells isolated from ticks serves as a resource
and starting point for delineating the functional determinants of *Bb* adaptation and transmission.
Our transcriptome profiles reveal more extensive expression changes for *Bb* outer surface
proteins than previously appreciated. These results raise the possibility that *Bb* is actively
remodeling its outer coat to navigate the dynamic tick environment during feeding, akin to
wardrobe changes across seasons. These outer surface changes may play roles in the
persistence of *Bb* in the tick, cell adhesion, or in immune evasion, either inside of the tick or later
in the vertebrate host (Kenedy et al., 2012). It has long been observed that molecular interactions
between pathogens and the midgut of their vectors are key determinants of transmission (Barillas-
Mury et al., 2022). Modifications to the *Bb* surface could facilitate different tick–pathogen
interactions that are critical for its physical movement through tick compartments. Notably, we
also identified numerous genes of unknown function that change expression during feeding.
Although challenging to address, efforts focused on uncovering the contribution of these genes
to the *Bb* life cycle could be groundbreaking for understanding the unique aspects of tick-borne
transmission.

More comprehensive knowledge about *Bb* outer surface proteins and their functional
consequences could lead to new avenues for curtailing the spread of Lyme disease. Protective
vaccines against vector-borne pathogens have often targeted surface proteins encoded by

pathogens (Kovacs-Simon et al., 2010). While previous Lyme disease vaccine efforts effectively
targeted the highly expressed *Bb* outer surface protein OspA (Steere et al., 1998), more targets
could increase the likelihood of a successful vaccine. Our study provides a catalog of new *Bb*
candidates that could be explored. In addition, our biochemical pull-downs using tick-isolated *Bb*
cells as bait unearthed a preliminary list of potential tick proteins that could be involved in tick–*Bb*
molecular interactions. Blocking molecular interactions that are functionally critical for
transmission could also be explored as a therapeutic strategy (Barillas-Mury et al., 2022; Manning
and Cantaert, 2019).

Our transcriptomic resource provides critical insights into *Bb* population-level changes during the
vector stage of its lifecycle, an important starting point for understanding the primary drivers of
tick-borne transmission. Strikingly few *Bb* cells out of the total pathogen population in ticks are
ultimately transmitted to the next host during feeding. There are several major bottlenecks to
infection, including dissemination to the tick salivary glands and survival upon inoculation of the
host (Dunham-Ems et al., 2009; Rego et al., 2014). There may be important physiological or
molecular variations across *Bb* cells within the population residing in the tick that contribute to
these differential outcomes. An important clue is the fact that *Bb* proteins are indeed
heterogeneously expressed across cells within the feeding tick midgut (Ohnishi et al., 2001).
Conducting transcriptomic analyses at the single-cell level will be key. Given that only a small
number of *Bb* escape the tick midgut, genes that appear unaltered or show low expression levels
across the bulk population could still play an outsized role in infection for a minority of the cells.
Our work provides a foundational methodology that can be leveraged to greatly improve the
resolution of tick–microbe studies. Technological advances stemming from this work will provide
molecular details that potentiate new questions and may unearth surprising mechanistic insights
into the unique lifestyles of tick-borne pathogens.

**METHODS**
***B. burgdorferi* culture**
*Bb* strain B31-S9 (Rego et al., 2011) was provided by Dr. Patricia Rosa (NIAID, NIH, RML) and
cultured in BSK II media at 35°C, 2.5% $CO_2$.

**Tick feeding experiments**
*I. scapularis* larvae were purchased from the Tick Lab at Oklahoma State University (OSU) (for
RNA-seq experiments) or provided by BEI Resources, a division of the Center for Disease Control

(for mass spectrometry experiments). Before and after feeding, ticks were maintained in glass

jars with a relative humidity of 95% (saturated solution of potassium nitrate) in a sealed incubator

at 22°C with a light cycle of 16h/8h (light/dark). Animal experiments were conducted in

accordance with the approval of the Institutional Animal Care and Use Committee (IACUC) at

UCSF. Ticks were fed on young (4–6-week-old) female C3H/HeJ mice acquired from Jackson

Laboratories. Mice were anesthetized with ketamine/xylazine before placement of ≤ 100 larval or

≤ 30 nymphal ticks. Ticks were either pulled off isoflurane anesthetized mice at various times

during feeding (1-3 days after placement) or allowed to feed to repletion and collected from mouse

cages. Replete larval ticks were placed in the incubator to molt before being used as nymphs in

experiments.

**Western blot with anti-*Bb* antibody**

To determine whether the anti-*Bb* antibody targeted ospA, we cultured wildtype *Bb* (B31-A3),

ospA1-mutants (ospA1) and ospA-restored *Bb* (ospA$^+$B1) (Battisti et al., 2008) to approximately

$5x10^7$ *Bb*/mL and centrifuged 3 mLs of culture for 7 minutes at 8000 x g, washing twice with PBS.

Pelleted cells were lysed in 50 µL of water, and 25 ng per sample was mixed with 5X loading dye

(0.25% Bromophenol Blue, 50% Glycerol, 10% Sodium Dodecyl Sulfate, 0.25M Tris-Cl pH 6.8,

10% B-Mercaptoethanol), run on a Mini-PROTEAN TGX 4-15% gel (Biorad), and transferred

using the Trans-Blot Turbo Transfer System (Biorad). After transfer, the blot was blocked for 30

minutes at 4°C in TBST (Tris buffered saline with .1% tween) with 5% milk, then treated with anti-

*Bb* antibody (Invitrogen: PA1-73004; RRID: AB_1016668) diluted 1:10,000 for 1 hour at room

temperature, followed by anti-rabbit HRP secondary antibody (Advansta) diluted 1:5,000 for 45

minutes at room temperature with short PBST washes between each step. Blots were exposed

using Clarity Western ECL Substrate (Biorad) and imaged using the Azure C400 imaging system

(Azure Biosystems). This experiment was repeated three times.

**Enrichment of *Bb* from feeding ticks**

To sequence RNA from *Bb* from feeding ticks, we enriched *Bb* to increase the ratio of *Bb* to tick

material. Larval ticks were fed to repletion on three mice that were infected with *Bb* through

intraperitoneal and subcutaneous injection with $10^4$ total *Bb*. Approximately five months later, the

molted nymphal ticks were fed on eight mice. We estimated that 83% of the ticks were infected

with *Bb* by crushing 12 unfed nymphs in BSK II media and checking for viable *Bb* days later. Ticks

were pulled from all mice and pooled into four replicates 1 day after placement (14 ticks per

replicate), 2 days after placement (12 ticks per replicate), and 3 days after placement (6 ticks per

replicate) and collected from cages 4 days after placement (7 ticks per replicate). Shortly after

collection, ticks were washed with water and placed in a 2 mL glass dounce grinder (Kimble) in

500 µL of phosphate-buffered saline (PBS). Ticks were homogenized first with the large clearance

pestle and then the small clearance pestle. The homogenate was transferred to a 1.5 mL

Eppendorf tube and 500 µL of PBS was added to total 1 mL. At this stage, 50 µL of homogenate

was removed as an input sample and mixed with 500 µL of TRIzol (Invitrogen) for RNA extraction.

2 µL of anti-*Bb* antibody (Invitrogen: PA1-73004; RRID: AB_1016668) was added to the

homogenate, which was then placed on a nutator at 4°C for 30 minutes. During incubation, 50 µL

of Dynabeads Protein G (Invitrogen) per sample were washed twice in PBS. After incubation with

the antibody, the homogenate and antibody mixture were added to the beads. This mixture was

placed on a nutator at 4°C for 30 minutes. Tubes were then placed on a magnet to secure beads,

and the homogenate was removed and saved to create depleted samples. The depleted

homogenate was centrifuged at 8,000 x g for 7 minutes, 900 µL of supernatant was removed, and

500 µL of TRIzol was added to the pellet to create depleted samples. The beads were washed

twice with 1 mL of PBS, resuspending the beads each time. The second wash was removed and

500 µL of TRIzol was added to the beads to create enriched samples. RNA was extracted from

all input, enriched, and depleted samples using the Zymo Direct-zol RNA Microprep Kit with on-

column DNase treatment (Zymo Research). The step-by-step *Bb* enrichment protocol is available

at: https://www.protocols.io/view/b-burgdorferi-enrichment-from-feeding-ticks-36wgqjrbovk5/v1

**Enrichment of *Bb* from culture**

To test whether the *Bb* enrichment process altered gene expression levels, we performed the

enrichment protocol on cultured *Bb*. We grew three tubes of *Bb* in BSK II media to 9x10$^4$ *Bb*/mL

at 35°C. 1 mL of culture was spun down at 8,000 x g for 7 minutes, media was removed, *Bb* were

washed in 1 mL of PBS and spun again. Pelleted *Bb* were resuspended in 1 mL of fresh PBS.

These samples were used as starting homogenate for the *Bb* enrichment protocol and input,

enriched, and depleted fractions were collected as above. RNA-seq libraries from these samples

were prepared and sequenced as below.

**RNA-seq library preparation and sequencing**

To make RNA-seq libraries from enriched *Bb* RNA, we used 50 ng of total RNA as input into the

NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (New England BioLabs). We

prepared libraries following the manufacturer's protocol for use of the kit with purified mRNA or

rRNA-depleted RNA, despite starting with total RNA. Libraries were barcoded using NEBNext
Multiplex Oligos for Illumina Dual Index (New England BioLabs).

To deplete reads from the libraries that were from tick, *Bb*, or mouse rRNA, we used Depletion of
Abundant Sequences by Hybridization (DASH) (Gu et al., 2016), which targets Cas9 to unwanted
reads in RNA-seq libraries using custom dual-guide RNAs (dgRNAs). Our dgRNAs targeted short
sequences within tick rRNA, mouse rRNA, and *Bb* rRNA that were designed using DASHit
software (Dynerman et al., 2020). We ordered crRNAs that targeted our sequences of interest
(Ring et al., 2022). To transcribe the dgRNAs, we followed our protocol for In Vitro Transcription
for dgRNA V2 (Lyden et al., 2019b). Both tracrRNA and pooled crRNA DNA templates were
annealed to equimolar amounts of T7 primer by heating to 95°C for 2 minutes and slowly cooling
to room temperature. Annealed templates were used in 1 mL in vitro transcription reactions with:
120 µL10X T7 buffer (400 mM Tris pH 7.9, 200 mM MgCl2, 50 mM DTT, 20 mM spermidine
(Sigma)), 100 µL of T7 enzyme (custom prepped enzyme from E. Crawford, diluted 1:100 in T7
buffer, final concentration: 100 µg/mL), 300 µl NTPs (25 mM each, Thermo Fisher Scientific), 4
µg of annealed crRNA template or 8 µg of annealed tracrRNA template, and water to 1 mL. In
vitro transcription was performed for 2 hours at 37°C. Reactions were purified twice with the Zymo
RNA Clean & Concentrator-5 Kit (Zymo Research). To form the dgRNA complex for DASH, crRNA
and tracrRNAs were diluted to 80 µM, mixed in equimolar amounts and annealed by heating to
95ºC for 30 seconds and cooling slowly to room temperature.

After transcription of dgRNAs, we performed DASH Protocol Version 4 (Lyden et al., 2019a) on
each library individually. Cas9 and transcribed dgRNAs were prepped by mixing: 2.5 µL 10X Cas9
buffer, 5 µL Cas9 (New England BioLabs), and 5 µL of 40 µM transcribed dgRNAs. The mixture
was incubated at 37°C for 5 minutes before 7.5 µL of RNA-seq library (2.8 nM) was added. The
mixture was incubated at 37°C for 1 hour and then purified with Zymo DNA Clean & Concentrator-
5 (Zymo Research) following the PCR product protocol and eluting DNA into 10.5 µL of water.
During cleanup, Cas9 was again mixed with buffer and dgRNAs and incubated at 37°C for 5
minutes. Following cleanup, eluted DNA was added to the second Cas9-dgRNA mixture and
incubated at 37°C for 1 hour for a second time. Then, 1 µL of proteinase K (New England Biolabs)
was added, and the mixture was incubated at 50°C for 15 min. The libraries were then purified
with 0.9x volume of sparQ PureMag Beads (QuantaBio) following the standard protocol, eluting
in 24 µL of water. rRNA-depleted RNA-seq libraries were then amplified in a BioRad CFX96 using
the Kapa HiFi Real-Time Amplification Kit in a 50 µL reaction with 25 µL master mix, 23 µL of the

DASHed library pool, and 2 µL of 25 µM mix of Illumina P5 (5'-

AATGATACGGCGACCACCGAGATCT) and P7 (5'- CAAGCAGAAGACGGCATACGAGAT)

primers. The qPCR program for amplification was as follows: 98°C for 45 sec (1 cycle), (98°C for

15 sec, 63°C for 30 sec, 72°C for 45 sec, plate read, 72°C for 20 sec) for 10 cycles (day 3 and

day 4 samples) or 11 cycles (day 1 and day 2 samples). The libraries were removed from cycling

conditions before leaving the exponential phase of amplification and then purified with 0.9X

volume of sparQ PureMag Beads according to the standard protocol.

Following DASH, RNA-seq libraries were sequenced on an Illumina NovaSeq S2 (2 lanes) with

paired-end 100 base pair reads. Libraries from *in vitro* cultured control experiment were

sequenced on an Illumina NextSeq with paired-end 75 base pair reads. FASTQ files and raw *Bb*

read counts for *in vitro* control experiment (GSE217146) and *ex vivo* experiment (GSE216261)

have been deposited in NCBI's Gene Expression Omnibus (Edgar et al., 2002) under SuperSeries

accession number GSE217236.

**RNA-seq data analysis**

To measure the success of our rRNA depletion through DASH, we used DASHit software

(Dynerman et al., 2020) to determine the percentage of reads that would be DASHable by our

guide RNAs (Ring et al., 2022). For pre-DASH data, we sequenced the input of each of our RNA-

seq libraries before performing DASH on a MiSeq V2 Micro (Illumina). We tested DASHability on

a random subset of 200,000 paired-end reads chosen by seqtk (https://github.com/lh3/seqtk) from

each pre- and post-DASH library.

To map our RNA-seq data to *Bb*, we wanted to optimize for mapping reads that came from the

many paralogous gene families found across the plasmids of the genome. We used the

pseudoalignment tool Salmon v1.2.1 (Patro et al., 2017), which is used to accurately map reads

coming from different isoforms of the same gene, for this reason. While using Salmon to map to

CDS sequences may improve mapping to paralogous genes, it may also have a tradeoff of

reduced mapping of reads that fall on the ends of genes that reside in operons. Nevertheless, all

samples should be similarly affected, and any undercounting should not change differential

expression results. Reads were first trimmed of bases with quality scores less than 20 using Trim

Galore v0.6.5 (https://github.com/FelixKrueger/TrimGalore). We mapped our reads to *Bb* CDS

sequences as a reference transcriptome: NCBI Genbank GCA_000008685.2 ASM868v2 (with

plasmids lp5, cp9, and lp56 removed as they are not present in B31-S9) using Salmon with the

following parameters: --validateMappings --seqBias --gcBias. Before mapping, the transcriptome

was first indexed using the Salmon index command with the whole genome as decoys and the

parameter --keepDuplicates to keep all duplicate genes.

Read counts from Salmon were used as input into DESeq2 v1.24.0 (Love et al., 2014) for

differential expression analysis in R version 3.6.1. DESeq2 function PlotPCA() was used to create

a PCA plot from read counts after running the varianceStabilizingTransformation() function. For

differential expression analysis between days, a DESeq object was created from count data using

the DESeq() function. The lfcShrink() function with the apeglm method (Zhu et al., 2018) was

used to calculate fold changes between days. Code used for differential expression analysis is

available at https://github.com/annesapiro/Bb-tick-feeding.

**Gene classification**

To classify genes into functional groups, functional categories were sourced from Drecktrah et

al., 2015 where available. Other gene functions were sourced from Fraser et al., 1997. Genes

found within the co-transcribed "late" bacteriophage operon (Zhang and Marconi, 2005) were

considered "bacteriophage" even if their function is unknown. Outer surface lipoproteins were

those found in Dowdell et al., 2017 plus additional lipoproteins listed in Iyer et al., 2015 that were

found in Dowdell et al. 2017 categories SpII and SpI and evidence of outer surface localization.

Outer surface and periplasmic lipoproteins were classified as "cell envelope" in the absence of

other classifications. Gene family information from Casjens et al., 2000 was considered to aid in

classification.

**RT-qPCR measuring *Bb* enrichment**

To test the efficacy of our enrichment protocol, we used RT-qPCR to quantify *Bb flaB* and *I.*

*scapularis gapdh* transcript levels in enriched and depleted fractions. We synthesized cDNA from

8 µL of RNA extracted from *Bb* enrichment samples and their matched depleted samples from

feeding day 2, day 3, and day 4 using the qScript cDNA Synthesis Kit (Quantabio). cDNA was

diluted 2X before use in qPCR. To measure *flaB* copies, we created standards of known

concentration from purified PCR products. These standards were made from PCR with primers

with the following sequences: 5'-CACATATTCAGATGCAGACAGAGGTTCTA and 5'-

GAAGGTGCTGTAGCAGGTGCTGGCTGT. We ran a dilution series with ten-fold dilutions

between $10^6$ copies and $10^1$ copies of this PCR template alongside our enriched and depleted

samples. We performed qPCR using Taqman Universal PCR Master Mix (Applied Biosystems).

The primers used to amplify *flaB* were: 5'- TCTTTTCTCTGGTGAGGGAGCT and 5'-

TCCTTCCTGTTGAACACCCTCT (used at 900 nM) and the probe was /56-
FAM/AAACTGCTCAGGCTGCACCGGTTC/36-TAMSp (used at 250 nM). For tick *gapdh* RT-
qPCR, the cDNA samples were diluted an additional 2X. Standards of known concentration were
created using the qPCR primer sequences. Primer sequences were: 5'-
TTCATTGGAGACACCCACAG and 5'-CGTTGTCGTACCACGAGATAA (used at 900 nM). qPCR
was performed using PowerUp SYBR Green Master Mix (Applied Biosystems). For both *flaB* and
*gapdh*, the number of copies in each sample was calculated based on the standards of known
concentration. Three technical replicates were averaged from each of four biological replicates at
each time point tested. We totaled the number of copies in each matched enriched and depleted
fraction to calculate the fraction of *flaB* or *gapdh* that was found in either sample. All qPCR was
performed on the QuantStudio3 Real-Time PCR System (Applied Biosystems).

**Immunofluorescence microscopy**
To test whether the anti-*Bb* antibody recognized *Bb* inside of the tick, ticks at each day of feeding
were crushed in 50 µL of PBS. 10 µL of lysate was spotted onto slides and allowed to air dry
before slides were heated briefly three times over a flame. Heat fixed slides were then treated
with acetone for one hour. Slides were incubated with anti-*Bb* primary antibody (1:100 diluted in
PBS + 0.75% BSA) for 30 minutes at 37°C in a humid chamber. A control without primary antibody
was also used for each day. Slides were washed once in PBS for 15 minutes at room temperature,
then rinsed in distilled water and air dried. Anti-rabbit IgG Alexa 488 (Invitrogen) diluted 1:100 in
PBS + 0.75% BSA was added for 30 minutes at 37°C in a humid chamber. Slides were washed
in PBS for 15 minutes at room temperature three times, adding 1:100 Propidium Iodide
(Invitrogen) during the second wash. Slides were then rinsed with distilled water and air dried
before the addition of mounting media (Fluoromount-G, SouthernBiotech) and cover slips.
Fluorescence imaging was performed on a Nikon Ti2 inverted microscope for widefield
epifluorescence using a 100x/1.40 objective. Images were captured with NIS-Elements AR View
5.20 and then processed with ImageJ software (Schneider et al., 2012). No strong florescence
signal was observed on the control slides without primary antibody.

**Mass spectrometry of *Bb*-enriched samples**
To identify which tick proteins were found in samples after *Bb* enrichment across feeding, we fed
both uninfected and infected ticks on mice. We collected three replicates of uninfected ticks one
day after attachment (11 ticks per replicate), infected ticks one day after attachment (27 ticks per
replicate), uninfected ticks four days after attachment (8 ticks per replicate), and infected ticks

four days after attachment (16 ticks per replicate). Before anti-*Bb* enrichment, the uninfected tick

samples were mixed with *Bb* grown in culture that was washed with PBS (3 x$10^4$ *Bb* one day after

attachment and 3x$10^6$ *Bb* four days after attachment) and mixed lysates were rotated at room

temperature for 30 minutes. Infected tick samples underwent the *Bb* enrichment process

immediately. The enrichment process followed the same protocol used for RNA-seq. Sample

volumes were increased to 1 mL as needed, and then 2 µL of anti-*Bb* antibody (Invitrogen: PA1-

73004; RRID: AB_1016668) was added, and samples were rotated at 4℃ for 30 minutes. 50 µL

of Dynabeads Protein G per sample were washed in PBS during this incubation and added to the

lysates, which were rotated at 4°C for 30 minutes. The beads were washed twice with 1 mL of

PBS, and then placed into 50 µL of lysis buffer (iST LYSE, PreOmics). Samples were boiled at

95°C for 5 minutes, and lysates were removed from beads and frozen for mass spectrometry

preparation.

For mass spectrometry, a nanoElute was attached in line to a timsTOF Pro equipped with a

CaptiveSpray Source (Bruker). Chromatography was conducted at 40°C through a 25 cm

reversed-phase C18 column (PepSep) at a constant flowrate of 0.5 µL min−1. Mobile phase A

was 98/2/0.1% water/MeCN/formic acid (v/v/v) and phase B was MeCN with 0.1% formic acid

(v/v). During a 108 min method, peptides were separated by a 3-step linear gradient (5% to 30%

B over 90 min, 30% to 35% B over 10 min, 35% to 95% B over 4 min) followed by a 4 min isocratic

flush at 95% for 4 min before washing and a return to low organic conditions. Experiments were

run as data-dependent acquisitions with ion mobility activated in PASEF mode. MS and MS/MS

spectra were collected with m/z 100 to 1700 and ions with z = +1 were excluded.

Raw data files were searched using PEAKS Online Xpro 1.6 (Bioinformatics Solutions Inc.). The

precursor mass error tolerance and fragment mass error tolerance were set to 20 ppm and 0.03

respectively. The trypsin digest mode was set to semi-specific and missed cleavages was set to

2. The *I. scapularis* reference proteome (Proteome ID UP000001555, taxon 6945) and *Bb*

reference proteome (Proteome ID UP000001807, strain ATCC 35210/B31) was downloaded from

Uniprot, totaling 21,774 entries. The *I. scapularis* proteome was the primary search reference and

the *Bb* was used as a secondary to identify any bacterial proteins present. Carbamidomethylation

was selected as a fixed modification. Oxidation (M) and Deamidation (NQ) was selected as a

variable modification.

Experiments were performed in biological triplicate, with samples being a single run on the

instrument. Proteins present in a database search (−10 log(p-value) ≥ 20, 1% peptide and protein

FDR) were subjected to the following filtration process. We first filtered to include proteins found

in 2 out of 3 biological replicates, within each respective day (one or four days after attachment).

We calculated the mean area of proteins found in uninfected and infected samples. Proteins with

missing values (i.e. not identified in a sample) were set to 1. We calculated the ratio of mean area

for each protein as infected/uninfected, and enriched proteins were identified by having a

infected:uninfected ratio greater than 2 within their respective feeding day (one or four days after

attachment).

To classify the proteins we identified into functional groups, we used a PANTHER

Overrepresentation Test (released 07/12/2022) (Mi et al., 2019) with PANTHER version 17.0

(released 02/22/2022) (Thomas et al., 2022). We used all *Ixodes scapularis* genes in the database

as our reference list and analyzed the lists of proteins enriched on each day for their PANTHER

Protein Class. The test type used was Fisher's Exact, calculating a false discovery rate as the

correction.

Raw data files and searched datasets are available on the Mass Spectrometry Interactive Virtual

Environment (MassIVE), a full member of the Proteome Xchange consortium under the identifier:

MSV000090560.

**SUPPLEMENTAL TABLES**

**Table S1.** Overview of mapping statistics from 16 *Bb* sequencing samples.

**Table S2.** Differential expression analysis results between *in vitro* cultured *Bb* before and after

*Bb* enrichment.

**Table S3.** Transcriptome-wide differential expression analysis results from *Bb* across tick

feeding timepoints.

**Table S4.** Twofold differentially expressed *Bb* genes from across tick feeding timepoints.

**Table S5.** Transcripts per million (TPM) for all *Bb* genes across feeding timepoints.

**Table S6.** Mass spectrometry analysis for *Bb*-enriched samples.

**Table S7.** Annotation and GO term enrichment for tick proteins enriched on feeding day 1.

**Table S8.** Annotation and GO term enrichment for tick proteins enriched on feeding day 4.

**ACKNOWLEDGMENTS**

We are grateful to all members of the Chou lab for their feedback throughout the project and

specifically to Fauna Yarza and Patrick Rockefeller Grimes for assistance with tick feeding and

Ethel Enoex-Godonoo for administrative assistance. We thank Amy Lyden and Emily Crawford

for assistance with and reagents for DASH along with Olga Botvinnik and The Chan Zuckerberg

Biohub sequencing team for their input and sequencing assistance. We thank Patricia Rosa,

Jenny Wachter, Scott Samuels, and Meghan Lybecker for their feedback on the project. We also

thank William Hatleberg for assistance with figure schematics. This work was funded by: a Life

Sciences Research Foundation fellowship from the SVCF-Wave Fund to Anne Sapiro, a Beckman

Young Investigator award from the Arnold and Mabel Beckman Foundation to Balyn Zaro, grants

from CZ Biohub, the Pew Biomedical Research Foundation, and NIH funding 1R01AI132851 to

Seemay Chou, and NIH INBRE funding P20GM103474 to Patrick Secor and Margie Kinnersley.

**COMPETING INTERESTS**

Seemay Chou is president and CEO of Arcadia Biosciences.

**REFERENCES**

Barbour AG. 1988. Plasmid analysis of Borrelia burgdorferi, the Lyme disease agent. *J Clin*
*Microbiol* 26:475–478. doi:10.1128/jcm.26.3.475-478.1988

Barillas-Mury C, Ribeiro JMC, Valenzuela JG. 2022. Understanding pathogen survival and
transmission by arthropod vectors to prevent human disease. *Science* 377:eabc2757.
doi:10.1126/science.abc2757

Battisti JM, Bono JL, Rosa PA, Schrumpf ME, Schwan TG, Policastro PF. 2008. Outer Surface
Protein A Protects Lyme Disease Spirochetes from Acquired Host Immunity in the Tick
Vector. *Infect Immun* 76:5228–5237. doi:10.1128/iai.00410-08

Blevins JS, Hagman KE, Norgard MV. 2008. Assessment of decorin-binding protein A to the
infectivity of Borrelia burgdorferi in the murine models of needle and tick infection. *Bmc*
*Microbiol* 8:82. doi:10.1186/1471-2180-8-82

Caimano MJ, Dunham-Ems S, Allard AM, Cassera MB, Kenedy M, Radolf JD. 2015. Cyclic di-
GMP Modulates Gene Expression in Lyme Disease Spirochetes at the Tick-Mammal
Interface To Promote Spirochete Survival during the Blood Meal and Tick-to-Mammal
Transmission. *Infect Immun* 83:3043–3060. doi:10.1128/iai.00315-15

Caimano MJ, Groshong AM, Belperron A, Mao J, Hawley KL, Luthra A, Graham DE, Earnhart
CG, Marconi RT, Bockenstedt LK, Blevins JS, Radolf JD. 2019. The RpoS Gatekeeper in
Borrelia burgdorferi: An Invariant Regulatory Scheme That Promotes Spirochete Persistence
in Reservoir Hosts and Niche Diversity. *Front Microbiol* 10:1923.
doi:10.3389/fmicb.2019.01923

Caimano MJ, Iyer R, Eggers CH, Gonzalez C, Morton EA, Gilbert MA, Schwartz I, Radolf JD.
2007. Analysis of the RpoS regulon in Borrelia burgdorferi in response to mammalian host
signals provides insight into RpoS function during the enzootic cycle. *Mol Microbiol* 65:1193–
1217. doi:10.1111/j.1365-2958.2007.05860.x

Casjens S, Palmer N, Vugt RV, Huang WM, Stevenson B, Rosa P, Lathigra R, Sutton G,
Peterson J, Dodson RJ, Haft D, Hickey E, Gwinn M, White O, Fraser CM. 2000. A bacterial
genome in flux: the twelve linear and nine circular extrachromosomal DNAs in an infectious
isolate of the Lyme disease spirochete Borrelia burgdorferi. *Mol Microbiol* 35:490–516.
doi:10.1046/j.1365-2958.2000.01698.x

Casjens SR, Mongodin EF, Qiu W-G, Luft BJ, Schutzer SE, Gilcrease EB, Huang WM,
Vujadinovic M, Aron JK, Vargas LC, Freeman S, Radune D, Weidman JF, Dimitrov GI,
Khouri HM, Sosa JE, Halpin RA, Dunn JJ, Fraser CM. 2012. Genome Stability of Lyme
Disease Spirochetes: Comparative Genomics of Borrelia burgdorferi Plasmids. *Plos One*
7:e33280. doi:10.1371/journal.pone.0033280

Dowdell AS, Murphy MD, Azodi C, Swanson SK, Florens L, Chen S, Zückert WR. 2017.
Comprehensive Spatial Analysis of the Borrelia burgdorferi Lipoproteome Reveals a
Compartmentalization Bias toward the Bacterial Surface. *J Bacteriol* 199:e00658-16.
doi:10.1128/jb.00658-16

Drecktrah D, Lybecker M, Popitsch N, Rescheneder P, Hall LS, Samuels DS. 2015. The
Borrelia burgdorferi RelA/SpoT Homolog and Stringent Response Regulate Survival in the
Tick Vector and Global Gene Expression during Starvation. *Plos Pathog* 11:e1005160.
doi:10.1371/journal.ppat.1005160

Dunham-Ems SM, Caimano MJ, Pal U, Wolgemuth CW, Eggers CH, Balic A, Radolf JD. 2009.
Live imaging reveals a biphasic mode of dissemination of Borrelia burgdorferi within ticks. *J*
*Clin Invest* 119:3652–3665. doi:10.1172/jci39401

Dynerman D, Lyden A, Quan J, Caldera S, McGeever A, Dimitrov B, King R, Cirolia G, Tan M,
Sit R, Berge M van den, Kerstjens HAM, Faiz A, Christenson S, Langelier C, DeRisi J,
Crawford E. 2020. Designing and implementing programmable depletion in sequencing
libraries with DASHit. *Biorxiv* 2020.01.12.891176. doi:10.1101/2020.01.12.891176

Edgar R, Domrachev M, Lash AE. 2002. Gene Expression Omnibus: NCBI gene expression
and hybridization array data repository. *Nucleic Acids Res* 30:207–210.
doi:10.1093/nar/30.1.207

Eggers CH, Samuels DS. 1999. Molecular evidence for a new bacteriophage of Borrelia
burgdorferi. *J Bacteriol* 181:7308–13.

Fraser CM, Casjens S, Huang WM, Sutton GG, Clayton R, Lathigra R, White O, Ketchum KA,
Dodson R, Hickey EK, Gwinn M, Dougherty B, Tomb J-F, Fleischmann RD, Richardson D,
Peterson J, Kerlavage AR, Quackenbush J, Salzberg S, Hanson M, Vugt R van, Palmer N,
Adams MD, Gocayne J, Weidman J, Utterback T, Watthey L, McDonald L, Artiach P,
Bowman C, Garland S, Fujii C, Cotton MD, Horst K, Roberts K, Hatch B, Smith HO, Venter
JC. 1997. Genomic sequence of a Lyme disease spirochaete, Borrelia burgdorferi. *Nature*
390:580–586. doi:10.1038/37551

Gilmore RD, Howison RR, Dietrich G, Patton TG, Clifton DR, Carroll JA. 2010. The bba64 gene
of Borrelia burgdorferi, the Lyme disease agent, is critical for mammalian infection via tick
bite transmission. *P Natl Acad Sci Usa* 107:7515–20. doi:10.1073/pnas.1000268107

Gilmore RD, Mbow ML, Stevenson B. 2001. Analysis of Borrelia burgdorferi gene expression
during life cycle phases of the tick vector Ixodes scapularis. *Microbes Infect* 3:799–808.
doi:10.1016/s1286-4579(01)01435-6

Gu W, Crawford ED, O'Donovan BD, Wilson MR, Chow ED, Retallack H, DeRisi JL. 2016.
Depletion of Abundant Sequences by Hybridization (DASH): using Cas9 to remove
unwanted high-abundance species in sequencing libraries and molecular counting
applications. *Genome Biol* 17:41. doi:10.1186/s13059-016-0904-5

Haas BJ, Chin M, Nusbaum C, Birren BW, Livny J. 2012. How deep is deep enough for RNA-
Seq profiling of bacterial transcriptomes? *Bmc Genomics* 13:734. doi:10.1186/1471-2164-13-
734

He M, Ouyang Z, Troxell B, Xu H, Moh A, Piesman J, Norgard MV, Gomelsky M, Yang XF.
2011. Cyclic di-GMP is Essential for the Survival of the Lyme Disease Spirochete in Ticks.
*Plos Pathog* 7:e1002133. doi:10.1371/journal.ppat.1002133

Hübner A, Yang X, Nolen DM, Popova TG, Cabello FC, Norgard MV. 2001. Expression of
Borrelia burgdorferi OspC and DbpA is controlled by a RpoN–RpoS regulatory pathway. *Proc
National Acad Sci* 98:12724–12729. doi:10.1073/pnas.231442498

Iyer R, Caimano MJ, Luthra A, Axline D, Corona A, Iacobas DA, Radolf JD, Schwartz I. 2015.
Stage-specific global alterations in the transcriptomes of Lyme disease spirochetes during
tick feeding and following mammalian host adaptation. *Mol Microbiol* 95:509–538.
doi:10.1111/mmi.12882

Kasumba IN, Bestor A, Tilly K, Rosa PA. 2016. Virulence of the Lyme disease spirochete before
and after the tick bloodmeal: a quantitative assessment. *Parasite Vector* 9:129.
doi:10.1186/s13071-016-1380-1

Kenedy MR, Lenhart TR, Akins DR. 2012. The role of Borrelia burgdorferi outer surface
proteins. *Fems Immunol Medical Microbiol* 66:1–19. doi:10.1111/j.1574-695x.2012.00980.x

Kovacs-Simon A, Titball RW, Michell SL. 2010. Lipoproteins of Bacterial Pathogens. *Infect
Immun* 79:548–561. doi:10.1128/iai.00682-10

Kurokawa C, Lynn GE, Pedra JHF, Pal U, Narasimhan S, Fikrig E. 2020. Interactions between
Borrelia burgdorferi and ticks. *Nat Rev Microbiol* 1–14. doi:10.1038/s41579-020-0400-5

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* 15:550. doi:10.1186/s13059-014-0550-8

Lyden A, Crawford E, Quan J, Caldera S, Dynerman D. 2019a. DASH Protocol v4.
doi:10.17504/protocols.io.6rjhd4n

Lyden A, Crawford E, Quan J, Caldera S, Pesce-Ares L. 2019b. In Vitro Transcription for
dgRNA v2. doi:10.17504/protocols.io.3bpgimn

Manning JE, Cantaert T. 2019. Time to Micromanage the Pathogen-Host-Vector Interface:
Considerations for Vaccine Development. *Nato Adv Sci Inst Se* 7:10.
doi:10.3390/vaccines7010010

Mi H, Muruganujan A, Huang X, Ebert D, Mills C, Guo X, Thomas PD. 2019. Protocol Update for
large-scale genome and gene function analysis with the PANTHER classification system
(v.14.0). *Nat Protoc* 14:703–721. doi:10.1038/s41596-019-0128-8

Ohnishi J, Piesman J, Silva AM de. 2001. Antigenic and genetic heterogeneity of Borrelia
burgdorferi populations transmitted by ticks. *Proceedings of the National Academy of
Sciences* 98:670–5. doi:10.1073/pnas.98.2.670

Ojaimi C, Mulay V, Liveris D, Iyer R, Schwartz I. 2005. Comparative Transcriptional Profiling of
Borrelia burgdorferi Clinical Isolates Differing in Capacities for Hematogenous Dissemination.
*Infect Immun* 73:6791–6802. doi:10.1128/iai.73.10.6791-6802.2005

Pal U, Kitsou C, Drecktrah D, Yaş OB, Fikrig E. 2022. Interactions Between Ticks and Lyme
Disease Spirochetes. *Curr Issues Mol Biol* 42:113–144. doi:10.21775/cimb.042.113

Pal U, Li X, Wang T, Montgomery RR, Ramamoorthi N, deSilva AM, Bao F, Yang X, Pypaert M,
Pradhan D, Kantor FS, Telford S, Anderson JF, Fikrig E. 2004. TROSPA, an Ixodes
scapularis Receptor for Borrelia burgdorferi. *Cell* 119:457–468.
doi:10.1016/j.cell.2004.10.027

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017. Salmon provides fast and bias-
aware quantification of transcript expression. *Nat Methods* 14:417 419.
doi:10.1038/nmeth.4197

Patton TG, Brandt KS, Nolder C, Clifton DR, Carroll JA, Gilmore RD. 2013. Borrelia burgdorferi
bba66 Gene Inactivation Results in Attenuated Mouse Infection by Tick Transmission. *Infect
Immun* 81:2488–2498. doi:10.1128/iai.00140-13

Phelan JP, Kern A, Ramsey ME, Lundt ME, Sharma B, Lin T, Gao L, Norris SJ, Hyde JA, Skare
JT, Hu LT. 2019. Genome-wide screen identifies novel genes required for Borrelia
burgdorferi survival in its Ixodes tick vector. *Plos Pathog* 15:e1007644.
doi:10.1371/journal.ppat.1007644

Radolf JD, Caimano MJ, Stevenson B, Hu LT. 2012. Of ticks, mice and men: understanding the
dual-host lifestyle of Lyme disease spirochaetes. *Nat Rev Microbiol* 10:87 99.
doi:10.1038/nrmicro2714

Rego ROM, Bestor A, Rosa PA. 2011. Defining the Plasmid-Borne Restriction-Modification
Systems of the Lyme Disease Spirochete Borrelia burgdorferi. *J Bacteriol* 193:1161–1171.
doi:10.1128/jb.01176-10

Rego ROM, Bestor A, Štefka J, Rosa PA. 2014. Population Bottlenecks during the Infectious
Cycle of the Lyme Disease Spirochete Borrelia burgdorferi. *Plos One* 9:e101009.
doi:10.1371/journal.pone.0101009

Revel AT, Talaat AM, Norgard MV. 2002. DNA microarray analysis of differential gene
expression in Borrelia burgdorferi, the Lyme disease spirochete. *Proc National Acad Sci*
99:1562–1567. doi:10.1073/pnas.032667699

Ring K, Couper LI, Sapiro AL, Yarza F, Yang XF, Clay K, Mateusiak C, Chou S, Swei A. 2022.
Host blood meal identity modifies vector gene expression and competency. *Mol Ecol*
31:2698–2711. doi:10.1111/mec.16413

Rosenberg R, Lindsey NP, Fischer M, Gregory CJ, Hinckley AF, Mead PS, Paz-Bailey G,
Waterman SH, Drexler NA, Kersh GJ, Hooks H, Partridge SK, Visser SN, Beard CB,
Petersen LR. 2018. Vital Signs: Trends in Reported Vectorborne Disease Cases — United
States and Territories, 2004–2016. *Morbidity Mortal Wkly Rep* 67:496–501.
doi:10.15585/mmwr.mm6717e1

Samuels DS, Lybecker MC, Yang XF, Ouyang Z, Bourret TJ, Boyle WK, Stevenson B,
Drecktrah D, Caimano MJ. 2021. Gene Regulation and Transcriptomics. *Curr Issues Mol Biol*
42:223–266. doi:10.21775/cimb.042.223

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods* 9:671–675. doi:10.1038/nmeth.2089

Schwartz I, Margos G, Casjens SR, Qiu W-G, Eggers CH. 2021. Multipartite Genome of Lyme
Disease Borrelia : Structure, Variation and Prophages. *Curr Issues Mol Biol* 42:409–454.
doi:10.21775/cimb.042.409

Shaw WR, Catteruccia F. 2018. Vector biology meets disease control: using basic research to
fight vector-borne diseases. *Nat Microbiol* 4:20–34. doi:10.1038/s41564-018-0214-7

Spielman A, Ribeiro JMC, Mather TN, Piesman J. 1987. Dissemination and Salivary Delivery of
Lyme Disease Spirochetes in Vector Ticks (Acari: Ixodidae). *J Med Entomol* 24:201–205.
doi:10.1093/jmedent/24.2.201

Steere AC, Sikand VK, Meurice F, Parenti DL, Fikrig E, Schoen RT, Nowakowski J, Schmid CH,
Laukamp S, Buscarino C, Krause DS. 1998. Vaccination against Lyme Disease with
Recombinant Borrelia burgdorferi Outer-Surface Lipoprotein A with Adjuvant. *New Engl J*
*Medicine* 339:209–215. doi:10.1056/nejm199807233390401

Steere AC, Strle F, Wormser GP, Hu LT, Branda JA, Hovius JWR, Li X, Mead PS. 2016. Lyme
borreliosis. *Nat Rev Dis Primers* 2:nrdp201690. doi:10.1038/nrdp.2016.90

Stevenson B, Zückert WR, Akins DR. 2000. Repetition, conservation, and variation: the multiple
cp32 plasmids of Borrelia species. *J Mol Microb Biotech* 2:411–22.

Thomas PD, Ebert D, Muruganujan A, Mushayahama T, Albou L, Mi H. 2022. PANTHER:
Making genome-scale phylogenetics accessible to all. *Protein Sci* 31:8–22.
doi:10.1002/pro.4218

Tilly K, Rosa PA, Stewart PE. 2008. Biology of Infection with Borrelia burgdorferi. *Infect Dis Clin*
*N Am* 22:217 234. doi:10.1016/j.idc.2007.12.013

Tokarz R, Anderton JM, Katona LI, Benach JL. 2004. Combined Effects of Blood and
Temperature Shift on Borrelia burgdorferi Gene Expression as Determined by Whole
Genome DNA Array. *Infect Immun* 72:5419–5432. doi:10.1128/iai.72.9.5419-5432.2004

Zhang H, Marconi RT. 2005. Demonstration of Cotranscription and 1-Methyl-3-Nitroso-
Nitroguanidine Induction of a 30-Gene Operon of Borrelia burgdorferi: Evidence that the 32-
Kilobase Circular Plasmids Are Prophages. *J Bacteriol* 187:7985–7995.
doi:10.1128/jb.187.23.7985-7995.2005

Zhang L, Zhang Y, Adusumilli S, Liu L, Narasimhan S, Dai J, Zhao YO, Fikrig E. 2011.
Molecular Interactions that Enable Movement of the Lyme Disease Agent from the Tick Gut
into the Hemolymph. *Plos Pathog* 7:e1002079. doi:10.1371/journal.ppat.1002079

Zhu A, Ibrahim JG, Love MI. 2018. Heavy-tailed prior distributions for sequence count data:
removing the noise and preserving large differences. *Bioinformatics* 35:2084–2092.
doi:10.1093/bioinformatics/bty895