Longitudinal map of transcriptome changes in the Lyme pathogen _Borrelia burgdorferi_

during tick-borne transmission

Anne L. Sapiro[1*], Beth M. Hayes[1], Regan F. Volk[2], Jenny Y. Zhang[1], Diane M. Brooks[3], Calla

Martyn[1], Atanas Radkov[1], Ziyi Zhao[1], Margie Kinnersley[3], Patrick R. Secor[3], Balyn W. Zaro[2],

Seemay Chou[1*]

1 Department of Biochemistry & Biophysics, University of California, San Francisco, California,

USA

2 Department of Pharmaceutical Chemistry and Cardiovascular Research Institute, University of

California, San Francisco, California, USA

3 Division of Biological Sciences, University of Montana, Missoula, Montana, USA

*Corresponding authors:

Anne L. Sapiro: annesapiro@gmail.com & Seemay Chou: seemaychou@gmail.com

ABSTRACT

Borrelia burgdorferi (_Bb_), the causative agent of Lyme disease, adapts to vastly different

environments as it cycles between tick vector and vertebrate host. During a tick bloodmeal, _Bb_

alters its gene expression to prepare for vertebrate infection; however, the full range of

transcriptional changes that occur over several days inside of the tick are technically challenging

to capture. We developed an experimental approach to enrich _Bb_ cells to longitudinally define

their global transcriptomic landscape inside nymphal _Ixodes scapularis_ ticks during a transmitting

bloodmeal. We identified 192 _Bb_ genes that substantially change expression over the course of

the bloodmeal from one to four days after host attachment. The majority of upregulated genes

encode proteins found at the cell envelope or proteins of unknown function, including 45 outer

surface lipoproteins embedded in the unusual protein-rich coat of _Bb_. As these proteins may

facilitate _Bb_ interactions with the host, we utilized mass spectrometry to identify candidate tick

proteins that physically associate with _Bb_. The _Bb_ enrichment methodology along with the _ex vivo_

_Bb_ transcriptomes and candidate tick interacting proteins presented here provide a resource to

facilitate investigations into key determinants of _Bb_ priming and transmission during the tick stage

of its unique transmission cycle.

INTRODUCTION

Vector-borne microbial pathogens are transmitted by the bite of arthropods and have evolved sophisticated ways to adapt to vastly different environments as they move between vector and host. Uncovering their adaptive mechanisms can not only open avenues for disrupting pathogen transmission but also provide fundamental insights into vector physiology and microbial symbioses (Shaw and Catteruccia, 2018). Lyme disease, the most reported vector-borne disease in North America, is caused by the bacterial pathogen *Borrelia burgdorferi* (*Bb*) (Rosenberg et al., 2018; Steere et al., 2016)*.* Its primary vector, the blacklegged tick *Ixodes scapularis,* acquires and transmits *Bb* through two separate multi-day bloodmeals, one in which *Bb* is acquired from an infected vertebrate host and a second, during the subsequent life stage, in which *Bb* is transmitted to a new host (Tilly et al., 2008). During the transmission bloodmeal, *Bb* proliferate in the tick midgut before a subset of these cells disseminate to the salivary glands (Dunham-Ems et al., 2009). *Bb* is deposited into the new host via the tick saliva extruded into the bite site (Spielman et al., 1987). Given the prolonged nature of *I. scapularis* feeding, the high specificity of vector-pathogen relationships, and the complicated array of events needed for successful *Bb* transmission, the tick bloodmeal provides an opportune intervention point for preventing pathogen spread. However, we do not currently have a clear understanding of the molecular mechanisms involved in this process.

Bb must adapt to dramatically different environments as it cycles from tick to vertebrate host, and understanding the genes involved in this process will enable the identification of key interactions to target to prevent transmission. When an infected tick feeds, *Bb* responds to bloodmeal-induced environmental changes and undergoes cellular modifications driven by key transcriptional circuits, including the RpoN-RpoS sigma factor cascade and the Hk1/Rrp1 two-component system (Radolf et al., 2012). *In vitro* analyses of *Bb* cells cultured in tick- or mammal-like growth conditions have pointed to additional genetic determinants of tick-borne transmission by revealing widespread transcriptome remodeling during host switching (reviewed in Samuels et al., 2021). However, it is not fully clear how these *in vitro* expression changes correspond to complex *in vivo* changes over the course of a transmission bloodmeal. Capturing comprehensive, longitudinal data on *Bb* gene expression from inside its vector has been hampered by technical challenges due to the dynamic nature of the bloodmeal and the general low abundance of bacterial cells relative to the tick (Samuels et al., 2021). Some progress has been made in making transcriptome-wide measurements from the tick. *Bb* sequence enrichment coupled with microarrays identified large scale changes in *Bb* gene expression between the first and second tick bloodmeals (Iyer et al.,

2015). More recently, enriching *Bb* sequences from infected tick RNA-seq libraries through

TDBCapSeq has provided clearer resolution into differences in *Bb* gene expression as it cycles

between the fed tick and mammalian host (Grassmann et al., 2023). Still, we have limited temporal

resolution into the molecular transitions that happen across key steps of tick feeding. This problem

necessitates novel approaches to capture the transcriptomic changes of *Bb* within the natural tick

environment.

While the full landscape of *Bb* transmission determinants is not yet known, we do have a growing

knowledge of functional processes that are critical during the tick stage, such as motility,

metabolism, and immune evasion (Kurokawa et al., 2020; Phelan et al., 2019). These functions

often rely on the unique protein-rich *Bb* outer surface. Notably, several specific tick–*Bb* protein–

protein interactions are important for *Bb* survival, migration, or transmission to the next host. *Bb*

encodes an extensive <u>o</u>uter <u>s</u>urface <u>p</u>rotein (Osp) family with members that are differentially

expressed during host switching. One of these proteins, OspA, binds a tick cell surface protein,

<u>t</u>ick <u>r</u>eceptor for <u>OspA</u> (TROSPA), which is required for successful *Bb* colonization of the tick

midgut during the first acquisition bloodmeal (Pal et al., 2004). Several other proteins have also

been linked to *Bb* migration within the tick (Pal et al., 2021). For example, BBE31 binds a tick

protein TRE31, and disruption of this interaction decreases the number of *Bb* cells that

successfully migrate from the tick gut to salivary glands (Zhang et al., 2011). However, these

interactions alone are not sufficient to block *Bb* growth or migration in ticks, suggesting there are

likely additional molecular factors from *Bb* and ticks at play during tick-borne transmission.

To provide a more comprehensive set of *Bb* determinants driving tick-borne transmission of this

important human pathogen, we developed a novel sequencing-based strategy for *ex vivo*

transcriptomic profiling of *Bb* populations within infected nymphal *I. scapularis* ticks as they

transmit *Bb* to a mouse host. We used this method to longitudinally map genome-wide *Bb*

expression changes for bacterial cells isolated from inside ticks during the transmission bloodmeal

from one to four days after attachment. We identified 192 highly differentially expressed genes,

including genes previously implicated in *Bb* transmission as well as many others. Genes

upregulated during tick transmission were enriched for outer surface lipoproteins, suggesting *Bb*

dramatically remodels its cell envelope as it migrates through the tick. Mass spectrometry

analyses revealed dramatic changes in the tick environment over feeding, identifying new

potential determinants of a more extensive and diverse set of tick–microbe molecular interactions

than previously appreciated. The *Bb* enrichment method and resulting datasets provide an

important community resource to facilitate further investigations into the key determinants of *Bb*

transmission.

RESULTS

A two-step enrichment process facilitates robust transcriptional profiling of *Bb* during the

tick bloodmeal

To gain a more comprehensive understanding of *Bb* gene expression throughout the tick phase

of the transmission cycle, we developed an experimental approach to characterize the *Bb*

transcriptome of spirochetes isolated from nymphal *I. scapularis* ticks during a days-long

bloodmeal in which *Bb* is transmitted to a vertebrate host. We aimed to establish a longitudinal

transcriptional profile encompassing key pathogen transmission events each day of feeding after

ticks attached to their mouse bloodmeal hosts (**Figure 1A**). We fed *Bb*-infected nymphal ticks on

naïve mice and collected the feeding ticks at daily intervals after the start of feeding until four days

after attachment, at which time the ticks had fully engorged and detached from the mice. The

major bottleneck for such an RNA sequencing (RNA-seq) approach is capturing sufficient

quantities of *Bb* transcripts from complex multi-organism samples in which pathogen transcripts

represent a very small minority. Our initial attempts to uncover *Bb* mRNA by simply removing tick

mRNA with polyA-depletion and removing tick rRNA sequences using Depletion of Abundant

Sequences by Hybridization (DASH) (Dynerman et al., 2020; Gu et al., 2016) were unsuccessful.

This approach resulted in an average of only 0.09% of RNA-seq reads mapping to *Bb* mRNA –

approximately 10-fold less than we estimated would be needed to feasibly obtain robust

transcriptome-wide differential

gene expression analysis (Haas

et al., 2012).

To dramatically increase *Bb*

transcript representation in our

libraries, we physically enriched

Bb cells in tick lysates prior to

library preparation by adding an

initial step of immunomagnetic

separation (**Figure 1B**). We took

advantage of a commercial

antibody previously generated

Figure S1. α*Bb* antibody recognizes OspA and binds *Bb* in the tick throughout the bloodmeal. (**A**) Western blot with α*Bb* on lysate from cultured *Bb*: wildtype (A3, left), a mutant lacking *ospA* (ospA1), and the mutant with *ospA* restored (ospA+B1)(Battisti et al., 2008). α*Bb* recognizes OspA among other proteins. (**B**) Immunofluorescence microscopy with α*Bb* (green, left) and propidium iodide (PI) (DNA, red, center) on each day of feeding (merge is yellow, right). α*Bb* antibody recognizes *Bb* in the tick across the bloodmeal.

against whole *Bb* cells (α*Bb*, RRID: AB_1016668). By western blot analysis, we confirmed that

α*Bb* specifically recognized several *Bb* proteins, including surface protein OspA (**Figure S1A**),

Figure 1. A two-step enrichment process facilitates robust transcriptional profiling of *Bb* during the tick bloodmeal. (**A**) Schematic of *Bb* during nymphal *I. scapularis* feeding. *Bb* in the nymphal tick midgut respond to the nutrient-rich bloodmeal by multiplying and changing their transcriptional state (Ouyang et al., 2012; de Silva and Fikrig, 1995). At the same time, the tick gut undergoes numerous changes to digest the bloodmeal (Caimano et al., 2015; Sonenshine and Anderson, 2014). After two to three days of feeding, a small number of *Bb* leave the midgut and enter the salivary glands (blue), while the majority are left behind in the gut after engorgement (Dunham-Ems et al., 2009). (**B**) Schematic of *Bb* enrichment process from feeding ticks. Whole ticks are dissociated, α*Bb* antibodies are added to lysates, and antibodies and *Bb* are captured magnetically. RNA is extracted and RNA-seq libraries are prepared. DASH is then used to remove rRNA before sequencing. This process increases *Bb* reads in the resulting sequencing data. (**C**) RT-qPCR results showing the percentage of *Bb flaB* and *I. scapularis gapdh* RNA in the enriched versus depleted fractions after the enrichment process. Data come from 4 replicates each from day 2, day 3, and day 4, mean +/- SE. ****p-value<0.0001, paired t test. Nearly all *Bb flaB* RNA was found in the enriched fraction. (**D**) The percentage of reads mapping to rRNA before and after DASH. n=4. Data are shown as mean +/- SD. ****p-value<0.0001, paired t test. rRNA reads are drastically reduced after DASH. (**E**) The percentage of reads in RNA-seq libraries mapping to *Bb*. *Bb* mRNA reads make up a larger proportion of libraries than without enrichment. Data are shown as mean +/- SD. (**F**) The number of reads in millions (M) mapped to *Bb* for each day. n=4. Data are shown as mean +/- SD. An average of 4.3 million reads per sample mapped to *Bb* genes, covering 92% of genes with at least 10 reads.

which is highly prevalent on the *Bb* surface in the tick (Radolf et al., 2012). In addition,
immunofluorescence microscopy with α*Bb* showed clear recognition of *Bb* cells from within the
tick at each day of feeding (**Figure S1B**). After collecting infected nymphal ticks from mice one,
two, three, and four days post-attachment, we used α*Bb* and magnetic beads to enrich *Bb* cells
from the tick material in the lysates. We tracked relative *Bb* enrichment through RT-qPCR of *Bb*
flaB RNA and tick *gapdh* RNA in the separated samples. Measuring *Bb flaB* RNA from both *Bb*-
enriched samples and their matched *Bb*-depleted fractions, we found over 95% of total *Bb flaB*
RNA was present in enriched fractions (**Figure 1C**), suggesting our approach captured the vast
majority of *Bb* transcripts from the tick.

Total RNA recovered from the enrichment process was used to create RNA-seq libraries that
subsequently underwent depletion of rRNA sequences from the tick, mouse, and *Bb* using DASH,
which targets unwanted sequences for degradation by Cas9 (Gu et al., 2016; Ring et al., 2022).
DASH reduced unwanted sequences from 94% to 9% of our total libraries, greatly increasing the
relative abundance of *Bb* transcripts (**Figure 1D**). For resulting libraries generated across feeding,
between 0.6% and 3.4% of reads mapped to *Bb* coding regions (**Figure 1E**), which translated to
an average of 4.3 million *Bb* genic reads per sample (**Figure 1F** and **Table S1**).

As expected, for samples pulled from the mice one, two, and three days after attachment, the
majority of the remaining sequencing reads mapped to the *I. scapularis* genome (72-83%) (**Table
S1**). On day four, when ticks were fully engorged and were recovered from mouse cages rather
than pulled from the mice, we found that fewer reads mapped to the *I. scapularis* genome (24-
44%). Only a small percentage of reads from all samples mapped to the host *Mus musculus*
genome (1-3%). To identify the source of the remaining reads in the day 4 samples, we ran the
data through a publicly available computational pipeline that identifies microbes in sequencing
datasets, CZ ID (Kalantar et al., 2020). This analysis led us to discover that a large percentage of
day 4 reads mapped to bacterial species *Pseudomonas fulva* (41-64%) (**Table S1**), which may
have been present in our mouse cages. While these samples had a lower percentage of reads
that mapped to *Bb*, broad transcriptome coverage was still obtained by increasing total
sequencing depth. Across all samples, at least 10 reads mapped to 92% of *Bb* genes. The median
number of reads per gene in each sample varied from 338 to 1167 reads (**Table S1**). This
coverage was sufficient for statistically significant downstream differential expression analyses
for the vast majority of *Bb* genes.

To evaluate whether our approach introduced any major
artifacts in *Bb* expression, we sequenced and compared
RNA-seq libraries from *in vitro* cultured *Bb* cells before and
after immunomagnetic enrichment. We found minimal
expression differences (29 genes with p<0.05, fold changes
between 0.83-1.12) (**Figure S2** and **Table S2**), suggesting
experimental enrichment did not significantly alter global
transcriptome profiles for *Bb*. Thus, our enrichment
approach enabled genome-wide analysis of *Bb* population-
level expression changes that occur within the feeding
nymph as *Bb* is transmitted to the host.

Figure S2. Enrichment process does not induce large scale gene expression changes in *in vitro* cultured *Bb*. Log$_2$ fold changes vs. mean normalized number of counts comparing cultured *Bb* input and samples after enrichment with α*Bb*. n=3. Red dots, p-value < 0.05, Wald tests. The gene expression changes induced during processing are much smaller than those observed between days of feeding.

Global *ex vivo* profiling of *Bb* reveals extent and
kinetics of transcriptional changes

To provide a broad overview of *Bb* expression changes in
the tick during the nymphal *I. scapularis* transmission
bloodmeal, we performed principal component analysis (PCA) on the *Bb* transcriptome data from
one, two, three, and four days after attachment (n=4). We reasoned that if many longitudinal
expression changes were occurring across *Bb* populations, we would observe greater data
variability between time points than between biological replicates. Indeed, we found replicates
from each day grouped together, whereas distinct time points were largely non-overlapping. The
first principal component, which explained 64% of the variance in our data, correlated well with
day of feeding (**Figure 2A**). The global pattern suggested that *Bb* gene expression changes
generally trended in the same direction over the course of feeding with the most dramatic
differences between flanking timepoints on day 1 and day 4.

Using day 1 (early attachment) as a baseline, we performed differential expression analysis for
all *Bb* genes at subsequent time points (day 2, day 3, day 4) (**Figure 2B** and **Table S3**). We
examined changes with p-values < 0.05 when adjusted for multiple hypothesis testing and fold
changes above a two-fold threshold (listed in **Table S4**). These analyses mirrored the global
longitudinal expression pattern predicted by the PCA. The total number of differentially expressed
(DE) genes when compared to day 1 increased with each subsequent timepoint to day 4. By day
4 there were 186 DE genes, including 153 upregulated and 33 downregulated (**Figure 2C-E**).
Across all later time point comparisons to day 1, DE genes were highly overlapping and largely

changed in the same directions. For example, of the DE genes that increased on day 2, 29 of 30

were still increased on day 3, and 29 of 30 were still increased on day 4. In the day 2, day 3, and

day 4 comparisons to the day 1 baseline, we found 192 DE genes in total (**Table S4**). There were

some notable differences between genes, such as the overall timing and kinetics of expression

changes. Transcript levels for some DE genes changed suddenly over the course of feeding,

while others were more gradual. To our knowledge, this is the first comprehensive report of global

Bb expression changes over multiple stages of a tick feeding.

Figure 2. Global *ex vivo* profiling of *Bb* reveals extent and kinetics of transcriptional changes. (**A**) Principal component analysis of samples from across feeding. PC1 correlates strongly with day of feeding. (**B**) Schematic depicting how data was analyzed, as pairwise comparisons between the first day after attachment and all other days. (**C-E**) Volcano plots of differentially expressed genes comparing day 2 versus day 1(**C**), day 3 versus day 1 (**D**), and day 4 versus day 1 (**E**). The total number of upregulated genes is shown in the top right and the number of downregulated genes is shown in the top left. Yellow dots are genes that first change expression between day 1 and day 2, red dots are genes that first change expression between day 1 and day 3, and purple dots are genes that first change expression between day 1 and day 4. Two genes with log$_2$ fold changes >4 are shown at x=4, and five genes with -log$_{10}$(padj) >60 are shown at y=60. Only genes with p-value < 0.05 from Wald tests and at least a two-fold change are highlighted. n=4. By day 4 of feeding, 153 genes are upregulated and 33 genes are downregulated from day 1 baseline levels.

To assess the integrity of our dataset, we first examined expression profiles of previously

characterized targets of major transcriptional programs activated at the onset of the bloodmeal:

the RpoN/RpoS sigma factor cascade and the Hk1/Rrp1 two-component system (Caimano et al.,

2015; Grassmann et al., 2023). Between day 1 and day 4, the expression of *rpoS* increased

(**Figure S3A**), as expected (Hübner et al., 2001), and the majority of genes activated by RpoS in

the feeding tick (Grassmann et al., 2023), including canonical targets *ospC* and *dbpA*, were also

significantly upregulated (79/89, $p<0.05$, Wald tests) (**Figure S3B**). The majority of genes

activated or repressed by Rrp1 *in vitro* (Caimano et al., 2015) also trended significantly in the

expected direction *ex vivo* between day 1 and day 4 (111/148 upregulated, 37/57 downregulated,

$p<0.05$, Wald tests, **Figure S3C**). We also examined the expression trends of genes regulated by

Rel$_{Bbu}$ as part of the stringent response, another major transcriptional program active in *Bb* in the

tick during nutrient starvation (Drecktrah et al., 2015). About half of Rel$_{Bbu}$-regulated genes

changed in the direction expected if the stringent response was active during this time (129/251

upregulated, 111/226 downregulated, $p<0.05$, Wald tests) (**Figure S3D**); however, more of the

genes that were two-fold downregulated over feeding are regulated by Rel$_{Bbu}$ than either RpoS or

Rrp1, suggesting it may play a role during this time frame.

Figure S3. *Ex vivo* **RNA-seq corroborates transcriptional programs in the tick. (A)** Tukey style boxplot of Transcripts Per Million (TPM) on each day for *rpo*S. Black dots represent replicates. ****p-value < 0.00001, Wald test. *rpoS* expression increases over the course of feeding. (**B**) Volcano plot of DE genes comparing day 4 to day 1, with RpoS-upregulated genes (blue). Genes upregulated by RpoS in ticks increase during feeding. (**C**) Volcano plot of DE genes comparing day 4 to day 1, with Rrp1-upregulated (blue) and downregulated (pink) genes. Rrp1-regulated genes correlate well with genes up and downregulated during feeding. (**D**) Volcano plot of DE genes comparing day 4 to day 1, with Rel$_{Bbu}$-upregulated (blue) and downregulated (pink) genes. About half of Rel$_{Bbu}$ genes change in the expected direction over feeding.

We then compared the 192 two-fold DE genes in our longitudinal time course to those identified

in two previous studies that measured *Bb* gene expression changes from culture conditions

approximating the unfed tick and fed tick through modulation of temperature and/or pH (Ojaimi et

al., 2003; Revel et al., 2002). 31% of the DE genes upregulated from day 1 (49/158) were more

highly expressed in "fed tick" conditions compared to "unfed tick" conditions in one or both studies,

while 24% of the DE genes downregulated from day 1 (8/24) were more highly expressed in

"unfed tick" conditions in one or both studies (**Figure S4A** and **Table S4**). The studies become

more concordant when focusing on the DE genes that were upregulated on day 2, which were

generally the genes that changed the most dramatically in the time course. 70% of DE genes

upregulated on day 2 (21/30) were more highly expressed in "fed tick" conditions in these previous

studies, suggesting that the majority of the most dramatic gene expression changes we saw

across feeding agree with what was observed in these previous studies.

Figure S4. Genes changing over tick feeding overlap with genes that change expression in previously probed tick feeding contexts. (**A**) The overlap of two-fold changed genes with genes that changed expression in *Bb* cultures grown in conditions mimicking feeding ticks. Genes are grouped based on the first day that they changed two-fold from day 1. "Fed tick" culture conditions were 37ºC, pH 6.8 in Revel et al. 2002 and 35ºC, pH 7.4 in Ojaimi et al. 2003, and genes elevated in these conditions in one or both studies are highlighted in red. "Unfed tick" culture conditions were 23ºC, pH 7.5 in Revel et al. 2002 and 23ºC, pH 7.4 in Ojaimi et al. 2003, and genes elevated in these conditions in one or both studies are highlighted in teal. Genes that were not elevated in either condition in those studies are in gray. Particularly for genes that increase on day 2, there is a large overlap with genes elevated in "fed tick" culture conditions in previous studies. (**B**) The overlap of two-fold changed genes with genes that changed expression between fed nymphs and dialysis membrane chambers (DMCs) mimicking mammalian conditions. Genes are grouped based on the first day that they changed two-fold from day 1. *Bb* expression in fed nymphs versus in DMCs was compared by Iyer et al. 2015 using bacterial RNA amplification and microarray, while Grassmann et al. 2023 used TBDCapSeq. Genes elevated in fed nymphs in one or both studies are highlighted in red, while genes elevated in DMCs in one or both studies are highlighted in purple. Genes that were elevated in conflicting conditions between the two studies are in dark gray, while genes not elevated in either condition are in light gray. For genes that increase on day 2, there is a large overlap with genes elevated in fed nymphs, while genes that increase first on days 3 and 4 have a larger overlap with genes elevated in DMCs.

We also compared the DE genes to two studies that assessed *Bb* gene expression differences in

fed nymphs versus dialysis membrane chambers (DMCs), which mimic *Bb* conditions in the

mammal (Grassmann et al., 2023; Iyer et al., 2015). 63% of all upregulated DE genes (100/158)

were differentially expressed between fed nymphs and DMCs in one or both studies (**Figure S4B**

and **Table S4**). The genes that were more highly expressed in nymphs were most concentrated

amongst the day 2 DE genes (17/30, 57%), while the genes more highly expressed in DMCs were

concentrated amongst the day 3 and day 4 DE genes (55/128, 43%). These comparisons

suggested that the timing and magnitude of gene expression changes during feeding may indicate

whether gene expression will peak in the tick or continue rising once *Bb* is transmitted to the host.

Through comparisons to these previous studies, we were able to verify that our data captured

many expected transcriptional trends occurring during tick feeding. Nevertheless, 14% of the two-

fold DE genes were not previously found to change expression in these different tick-feeding

contexts (Grassmann et al., 2023; Iyer et al., 2015; Ojaimi et al., 2003; Revel et al., 2002) or

identified in these RNA-seq studies as dependent upon RpoS, Rrp1, or Rel_{Bbu} (Caimano et al.,

2015; Drecktrah et al., 2015; Grassmann et al., 2023), which are the three main *Bb* regulatory

programs active in the tick (Samuels et al., 2021). These additional genes highlight the necessity

of measuring transcription in the tick environment and suggest we uncovered gene expression

changes specific to the tick-stage of the *Bb* enzootic cycle. The nature and dynamics of these

changes provide insights into potential genetic determinants of *Bb* survival, proliferation, and

dissemination in the tick during transmission.

***Bb* genes upregulated during feeding are found predominantly on plasmids**

Bb has a complex, highly fragmented genome (Barbour, 1988) (**Figure 3A**), including numerous

plasmids that are necessary during specific stages of the enzootic cycle (Schwartz et al., 2021)

suggesting they contain genes that are crucial for pathogen transmission and survival. In fact,

many genes found on the plasmids have been previously shown to alter expression upon

environmental changes or in different host environments (Iyer et al., 2015; Ojaimi et al., 2005;

Revel et al., 2002; Tokarz et al., 2004). Thus, we reasoned that many of the 192 DE *Bb* genes

that change expression from day 1 to any later feeding time point (**Table S4**) would reside on the

plasmids, and we examined their distribution throughout the genome. Consistent with these

previous reports, we found that most of the upregulated genes were located on the plasmids

(143/158; 90%), while fewer were found on the chromosome (15/158; 10%) (**Figure 3B**), which

is home to the majority of metabolic and other housekeeping genes. In contrast, the majority of

the downregulated genes were found on the chromosome (27/34, 79%) (**Figure 3C**).

Figure 3. *Bb* genes upregulated during feeding are found predominantly on plasmids. (**A**) Schematic of the chromosome and plasmids in the *Bb* B31-S9 genome. Plasmid names denote whether the plasmid is linear (lp) or circular (cp) and the length of plasmids in kilobases (kb). For example, lp17 is a 17 kb linear plasmid. Genome is shown approximately to scale. (**B-C**) The number of genes from each chromosome or plasmid that increased (**B**) or decreased expression (**C**) twofold during feeding. Upregulated genes are distributed across plasmids, while downregulated genes are found on the chromosome and lp54.

Several plasmid-encoded genes that were longitudinally upregulated in our dataset have known

roles during the tick bloodmeal or in mammalian infection. Linear plasmid 54 (lp54),

which is an essential plasmid present in all *Bb* isolates (Casjens et al., 2012), contained the largest

number of upregulated genes. Many of the genes on lp54 are known to be regulated by RpoS

during feeding, including those encoding adhesins DbpA and DbpB, which are important for

infectivity in the host (Blevins et al., 2008). This set also included five members of a paralogous

family of outer surface lipoproteins BBA64, BBA65, BBA66, BBA71, and BBA73. BBA64 and

BBA66 are necessary for optimal transmission via the tick bite (Gilmore et al., 2010; Patton et al.,

2013). These findings indicate our dataset captures key *Bb* transcriptional responses known to

be important for survival inside the tick during a bloodmeal.

Many upregulated genes were also encoded by cp32 plasmid prophages. *Bb* strain B31-S9

harbors seven cp32 isoforms that are highly similar to each other (Casjens et al., 2012). When

cp32 prophages are induced, phage virions called φBB1 are produced (Eggers and Samuels,

1999). In addition to phage structural genes, cp32 contain loci that encode various families of

paralogous outer surface proteins (Stevenson et al., 2000). Amongst the cp32 genes that

increased over feeding were members of the RevA, Erp, and Mlp families, which are known to

increase expression during the bloodmeal (Gilmore et al., 2001). We also found several phage

genes that were upregulated, including those encoding proteins annotated as phage terminases
on cp32-3, cp32-4, and cp32-7 (BBS45, BBR45, and BBO44). Some cp32 genes have been
shown to change expression in response to the presence of blood (Tokarz et al., 2004), and as a
part of the stringent response regulated by Rel_{Bbu} (Drecktrah et al., 2015), while BBD18 and RpoS
regulate prophage production in the tick midgut after feeding (Wachter et al., 2023). Our data
suggest that some prophage genes are upregulated over the course of tick feeding, raising the
possibility that cp32 prophage are induced towards the end of feeding. Overall, our data support
the long-held idea that the *Bb* plasmids, which house many genes encoding cell envelope
proteins, proteins of unknown function, and prophage genes, play a critical role in the enzootic
cycle during the key transition period of tick feeding.

Bb genes encoding outer surface proteins are highly prevalent among upregulated genes
To gain a better overall sense of the types of genes that changed over feeding and the timing of
those changes, we grouped DE genes into functional categories. Since a high proportion of
plasmid genes encode lipoproteins within the unique protein-rich outer surface of *Bb*, genes of
unknown function, and predicted prophage genes (Casjens et al., 2000; Fraser et al., 1997), we
expected that many of the DE genes would fall into these categories. We classified the genes as
related to either: cell envelope, bacteriophage, cell division, DNA replication and repair,
chemotaxis and motility, metabolism, transporter proteins, transcription, translation, stress
response, protein degradation, or unknown (Drecktrah et al., 2015) (see **Table S4**).

Of the genes that increased two-fold over feeding, the clear majority each day of feeding fell into
two broad categories, cell envelope (55 of 158 total genes) and proteins of unknown function (69
of 158 total genes), with fewer genes related to metabolism, chemotaxis and motility, transporters,
bacteriophage, cell division, and transcription (**Figure 4A**). In contrast to the upregulated genes,
genes downregulated during feeding were more evenly distributed among the functional
categories – including translation, protein degradation, transcription, and metabolism – consistent
with many of them being located on the chromosome.

When looking at changes across these functional categories, the overrepresentation of cell
envelope proteins was striking, while not unexpected. The *Bb* outer surface is covered with
lipoproteins (**Figure 4B**), and these proteins are critical determinants in *Bb* interactions with the
various environments encountered during the enzootic cycle (Kurokawa et al., 2020). We found

Figure 4. *Bb* **genes encoding outer surface proteins are highly prevalent among upregulated genes. (A)** The number of *Bb* genes that change over the course of tick feeding sorted into functional categories. Genes that first change 2 days after attachment are shown in yellow, 3 days after attachment in red, and 4 days after attachment in purple. A majority of upregulated genes fall into cell envelope and unknown categories. **(B)** Schematic of the outer membrane of *Bb* showing outer surface lipoproteins. Lipoproteins can also reside in the periplasmic space. **(C)** Heat map of expression levels of all genes encoding outer surface lipoproteins as average Transcripts Per Million (TPM) across the 4 days of tick feeding. Gene names highlighted in blue were two-fold upregulated and genes in pink two-fold downregulated over feeding (see **Figure 2**). A majority of genes encoding outer surface proteins increased in expression throughout feeding, while having different magnitudes of expression.

that more than half (46 of 83) of annotated outer surface lipoproteins (Dowdell et al., 2017; Iyer

et al., 2015) changed expression two-fold over the time course (**Figure 4C**). These data

suggested widespread changes may be occurring on the *Bb* outer surface during feeding.

To understand the functional implications of expression changes in a majority of outer surface

lipoproteins, we also compared their relative expression. The magnitude of expression varied

greatly, with *ospA*, *ospB*, *ospC*, and *bba59* being the most highly expressed outer surface protein

transcripts. Many of the outer surface protein genes that we found increased expression over

feeding were much less abundantly expressed (**Figure 4C** and **Table S5**). However, even the

genes that appeared to have low expression in these population level measurements could play

important roles in transmission if they are highly expressed in a small number of crucial cells,

such as those that ultimately escape the midgut. While bulk RNA-seq cannot distinguish what is

happening at the single cell level, our data suggest that during the bloodmeal, *Bb* are undergoing

a complex outer surface transformation driven by increases in transcription of a majority of the
genes encoding these lipoproteins.

Identification of candidate tick interaction partners of *Bb* cells *ex vivo*

Our RNA-seq data suggested that the outer surface of *Bb* transforms over the course of feeding
as *Bb* are primed for transmission to a vertebrate host. At the same time, the tick midgut
environment is changing as the tick begins to digest its bloodmeal (Sonenshine and Anderson,
2014). Tick-*Bb* interactions are likely crucial at the beginning of the tick bloodmeal, as *Bb* adhere
to the tick gut epithelium before becoming motile and migrating out of the midgut and into the
salivary glands (Dunham-Ems et al., 2009). Some *Bb* outer surface proteins, such as BBE31 and
BBA52 play key roles in pathogen migration through interactions with the tick environment
(reviewed in Kurokawa et al., 2020). We wanted to explore the changing tick environment to
identify tick proteins with which *Bb* could interact throughout the tick bloodmeal. Since our *Bb*
enrichment process retained some tick material, we reasoned that tick proteins that interact with
Bb would be present in these samples.

To outline the changes occurring in the tick during feeding and to identify candidate *Bb*-interacting
tick proteins, we used mass spectrometry to survey the content of the tick material that was
enriched along with the *Bb* cells we sequenced at early and late stages during feeding. We purified
proteins from the α*Bb*-enriched fraction of crushed infected ticks one day after attachment and
four days after attachment in triplicate. As controls for each day, we also performed the *Bb*
enrichment process on lysate from uninfected ticks mixed with *in vitro* cultured *Bb* to help rule out
proteins that were not pulled down through *in vivo* tick-*Bb* interactions (**Figure 5A**). When
querying against the *Bb* and *I. scapularis* proteomes, we identified between 414 and 2240 protein
groups per sample replicate. The vast majority of all detected proteins were from ticks (2801/2858,
98%). To identify proteins of interest, we looked for those that were detected in at least two of
three replicates within infected ticks and had a mean average coverage twice that of uninfected
ticks mixed with cultured *Bb*. We found 256 proteins (251 from *I. scapularis* and 5 from *Bb*) that
were enriched with *Bb* from infected ticks one day after attachment and 226 proteins (220 from *I.
scapularis* and 6 from *Bb*) that were enriched with *Bb* from infected ticks four days after
attachment (**Table S6**). Of these proteins, only 27 (24 from *I. scapularis* and 3 from *Bb*) were
detectable on both days, suggesting the tick proteins present upon *Bb* enrichment change
dramatically over the course of feeding (**Figure 5A**). Amongst the small number of *Bb* proteins,
we identified OspC in the samples from day 4 after attachment but not day 1 after attachment,

confirming the expression change we saw in our RNA-seq. The distinct sets of tick proteins we
identified at each timepoint suggest dramatic changes occur in the *Bb*-infected tick midgut
environment during feeding that may alter the landscape of tick-*Bb* interactions.

Figure 5. Identification of candidate tick interaction partners of *Bb* cells *ex vivo*. (**A**) Schematic of experiment to determine candidate tick proteins interacting with *Bb* over the course of feeding. Ticks were collected 1 day after attachment and 4 days after attachment. Uninfected ticks at the same time points were mixed with cultured *Bb* as controls. *Bb* was enriched with α*Bb* antibody as in RNA-seq experiments and then subjected to mass spectrometry to identify tick proteins present in the samples. Venn diagram depicts the proteins enriched in day 1 and day 4 samples over controls in at least two of three replicates. Tick proteins that are enriched with *Bb* vary greatly over the course of feeding. (**B**) Tick proteins uniquely identified one day after attachment that are annotated as extracellular matrix (ECM) proteins. (**C**) Tick proteins uniquely identified four days after attachment that are annotated as low-density lipoprotein receptors. ECM and membrane proteins may be good candidates for *Bb*-interacting proteins.

Some of the proteins enriched with *Bb* from the tick may be good candidates for key *Bb*-interacting
partners during feeding, especially if they are localized to the surface of tick cells where they may
encounter *Bb*. The scarcity of both predicted and experimentally validated functions and
localizations for tick proteins makes it difficult to fully assess the potential for tick protein
interactions with extracellular *Bb*. Nevertheless, of the proteins found exclusively one day after
attachment, 10 were categorized as extracellular matrix proteins using the PANTHER gene
database (Thomas et al., 2022), and this category was statistically enriched (Fisher's exact test,
FDR=0.000093) (**Figure 5B** and **Table S7**). 30 additional proteins were annotated with a cellular
component as plasma membrane. Four days after attachment, we did not detect any annotated
extracellular matrix proteins; however, we identified 31 proteins that are likely to be found at the
membrane, including two proteins annotated as putative low-density lipoprotein receptors (**Figure**
5C and **Table S8**). These extracellular matrix and membrane proteins may be the most likely to
directly interact with *Bb* during this timeframe and are candidates for tick proteins important in the

Bb dissemination process. The proteins present in the changing tick environment may be key

determinants of pathogen transmission as *Bb* remodels its outer surface while preparing to

migrate through the tick to a new host.

DISCUSSION

Vector-borne pathogens must adapt to distinct environments as they are transmitted by

arthropods to colonize new bloodmeal hosts. The tick-borne Lyme disease pathogen *Bb*

undergoes transcriptional changes inside its vector over the course of a single bloodmeal that are

important for a number of key transmission events. For example, expression changes enable

survival in the feeding tick (He et al., 2011), facilitate transmission across several internal

compartments into the bloodmeal host (Kurokawa et al., 2020), and prime bacteria for successful

infection of the next host (Kasumba et al., 2016). Capturing these changes as they occur *in vivo*

has been challenging due to the low relative abundance of *Bb* material inside of the rapidly

growing, blood-filled tick. To date, many advances in our understanding of *Bb* expression during

transmission from tick to host have come from tracking changes in small subsets of genes during

tick feeding (Bykowski et al., 2007; Gilmore et al., 2001; Narasimhan et al., 2002), leveraging *Bb*

culture conditions that approximate environmental changes across its lifecycle (Ojaimi et al.,

2005; Revel et al., 2002; Tokarz et al., 2004), and defining transcriptional regulons outside of the

tick (Caimano et al., 2019, 2015; Drecktrah et al., 2015). Here, we developed an experimental

RNA-seq-based strategy to more directly and longitudinally profile gene expression for *Bb*

populations isolated from ticks over the course of a transmitting bloodmeal.

Our longitudinal collection of transcriptomes for *Bb* cells isolated from ticks serves as a resource

and starting point for delineating the functional determinants of *Bb* adaptation and transmission.

We focused our analysis on 192 genes that changed two-fold between day 1 and later days in

feeding. While these genes were highly concordant with those found in previous studies probing

transcriptional changes throughout the *Bb* enzootic cycle (Caimano et al., 2019, 2015; Drecktrah

et al., 2015; Grassmann et al., 2023; Iyer et al., 2015; Ojaimi et al., 2003; Revel et al., 2002), our

transcriptome profiles revealed changes in expression of 26 genes not observed in these previous

studies. These novel gene expression changes, like all of the 192 DE genes, were concentrated

in genes of unknown function and those encoding outer surface proteins. Although challenging to

address, efforts focused on uncovering the contribution of the genes of unknown function to the

Bb life cycle could be groundbreaking for understanding the unique aspects of tick-borne

transmission. Our data reveal more extensive expression changes for *Bb* outer surface proteins

than previously appreciated, with nine additional genes encoding cell envelope proteins
upregulated over feeding that were not found to change expression in the previous studies
examining unfed versus fed tick conditions or nymphs versus DMCs. The extensive changes in
genes encoding outer surface proteins suggest *Bb* is actively remodeling its outer coat to navigate
the dynamic tick environment during feeding, akin to wardrobe changes across seasons. These
outer surface changes may play roles in the persistence of *Bb* in the tick, cell adhesion, or in
immune evasion, either inside of the tick or later in the vertebrate host (Kenedy et al., 2012). It
has long been observed that molecular interactions between pathogens and the midgut of their
vectors are key determinants of transmission (Barillas-Mury et al., 2022). Modifications to the *Bb*
surface could facilitate different tick–pathogen interactions that are critical for its physical
movement through tick compartments.

More comprehensive knowledge about *Bb* outer surface proteins and their functional
consequences could lead to new avenues for curtailing the spread of Lyme disease. Protective
vaccines against vector-borne pathogens have often targeted surface proteins encoded by
pathogens (Kovacs-Simon et al., 2010). While previous Lyme disease vaccine efforts effectively
targeted the highly expressed *Bb* outer surface protein OspA (Steere et al., 1998), more targets
could increase the likelihood of a successful vaccine. Our study provides a catalog of new *Bb*
candidates that could be explored. In addition, our biochemical pull-downs using tick-isolated *Bb*
cells as bait unearthed a preliminary list of potential tick proteins that could be involved in tick–*Bb*
interactions. Blocking molecular interactions that are functionally critical for transmission could
also be explored as a therapeutic strategy (Barillas-Mury et al., 2022; Manning and Cantaert,
2019).

To identify the genes changing expression during tick feeding, we used an antibody targeting *Bb*
to overcome the low abundance of *Bb* in the tick. This method produced broad coverage of the
transcriptome over our time course, but there are caveats. While we confirmed that the hour-long
enrichment procedure does not cause significant gene expression changes in cultured *Bb*, this
may still influence gene expression in our *ex vivo* samples. The α*Bb* antibody, which targets OspA
and a number of other proteins, captured the vast majority, but not all, *Bb* from inside of the tick.
This loss may introduce some degree of bias. Further, a large number of day 4 reads mapping to
P. fulva suggests that this particular antibody may enrich other bacterial species. It is also
important to note that our method was unable to produce consistent expression data from *Bb* from
unfed ticks, limiting the full longitudinal scope of the study and potentially overlooking key changes

occurring in the first 24 hours of feeding. We predict this difficulty may be due to low numbers of

Bb in flat nymphs (de Silva and Fikrig, 1995) or the stress of the prolonged nutrient-deprived

period between tick feedings (Kung et al., 2013). The transcriptional landscape of unfed ticks may

be better probed through enrichment after RNA-seq library preparation such as TBDCapSeq

(Grassmann et al., 2023), or perhaps a combination of the two enrichment strategies. Despite

these caveats, this type of antibody-based enrichment strategy is a simple and flexible technique

that can probe gene expression from *Bb* from multiple timepoints in feeding ticks. With small

modifications, this protocol may be able to assist in facilitating sequencing of *Bb* or other *Borrelia*

species from other milieu or enrich other tick-borne bacterial species.

Our method has produced a transcriptomic resource providing critical insights into *Bb* population-

level changes during the vector stage of its lifecycle, which serves as a starting point for

understanding the primary drivers of tick-borne transmission. Strikingly few *Bb* cells out of the

total pathogen population in ticks are ultimately transmitted to the next host during feeding

(Dunham-Ems et al., 2009; Rego et al., 2014). There may be important molecular variations

across *Bb* cells within the population residing in the tick that contribute to these differential

outcomes including heterogeneously expressed proteins across cells within the feeding tick

midgut (Ohnishi et al., 2001). Genes that appear unaltered or show low expression levels across

the bulk population could still play an outsized role in infection for a minority of the cells, and

conducting transcriptomic analyses at the single-cell level will be key. Our work provides a

foundational methodology that can be leveraged to greatly improve the resolution of tick–microbe

studies. Technological advances stemming from this work will provide molecular may unearth

surprising mechanistic insights into the unique lifestyles of tick-borne pathogens.

METHODS

***B. burgdorferi* culture**

Bb strain B31-S9 (Rego et al., 2011) was provided by Dr. Patricia Rosa (NIAID, NIH, RML) and

cultured in BSK II media at 35°C, 2.5% CO_2. B31-S9 was used for all RNA-seq and *Bb* enrichment

experiments. Wildtype *Bb* strain B31-A3, ospA1-mutants (ospA1) and ospA-restored *Bb*

(ospA$^+$B1) (Battisti et al., 2008) used in α*Bb* western blot were also provided by Dr. Rosa.

Tick feeding experiments

I. scapularis larvae were purchased from the Tick Lab at Oklahoma State University (OSU) for

RNA-seq experiments or provided by BEI Resources, a division of the Center for Disease Control,

for mass spectrometry experiments. Before and after feeding, ticks were maintained in glass jars

with a relative humidity of 95% (saturated solution of potassium nitrate) in a sealed incubator at

22°C with a light cycle of 16h/8h (light/dark). Animal experiments were conducted in accordance

with the approval of the Institutional Animal Care and Use Committee (IACUC) at UCSF, Project

Number AN183452. Ticks were fed on young (4–6-week-old) female C3H/HeJ mice acquired from

Jackson Laboratories. Mice were anesthetized with ketamine/xylazine before placement of ≤ 100

larval or ≤ 30 nymphal ticks. Replete larval ticks were placed in the incubator to molt before being

used as nymphs in experiments. Nymphal ticks were either pulled off isoflurane anesthetized mice

at various times during feeding (1-3 days after placement) or allowed to feed to repletion and

collected from mouse cages (4 days after placement).

Western blot with α*Bb* antibody

To determine whether the α*Bb* antibody targeted ospA, wildtype *Bb* (B31-A3), ospA1-mutants

(ospA1) and ospA-restored *Bb* (ospA$^+$B1) (Battisti et al., 2008) were cultured to approximately

5x10^7 *Bb*/mL. 3 mLs of culture were centrifuged for 7 minutes at 8000 x g, washing twice with

PBS. Pelleted cells were lysed in 50 μL of water, and 25 μg of protein per sample were mixed

with 5X loading dye (0.25% Bromophenol Blue, 50% Glycerol, 10% Sodium Dodecyl Sulfate,

0.25M Tris-Cl pH 6.8, 10% B-Mercaptoethanol), run on a Mini-PROTEAN TGX 4-15% gel

(Biorad), and transferred using the Trans-Blot Turbo Transfer System (Biorad). After transfer, the

blot was blocked for 30 minutes at 4°C in TBST (Tris buffered saline with 0.1% tween) with 5%

milk, then treated with α*Bb* antibody (Invitrogen: PA1-73004; RRID: AB_1016668) diluted

1:10,000 for 1 hour at room temperature, followed by anti-rabbit HRP secondary antibody

(Advansta: R-05072-500; RRID: AB_10719218) diluted 1:5,000 for 45 minutes at room

temperature with 3 short PBST washes between each step. Blots were exposed using Clarity

Western ECL Substrate (Biorad) and imaged using the Azure C400 imaging system (Azure

Biosystems). This experiment was repeated three times.

Enrichment of *Bb* from feeding ticks

To sequence RNA from *Bb* from feeding ticks, *Bb* were enriched to increase the ratio of *Bb* to tick

material. Larval ticks were fed to repletion on three mice that were infected with *Bb* through

intraperitoneal and subcutaneous injection with 10^4 total *Bb*. Approximately five months later, the

molted nymphal ticks were fed on eight mice, which were housed individually during the feeding.

We estimated that 83% of the ticks were infected with *Bb* by crushing 12 unfed nymphs in BSK II

media and checking for viable *Bb* days later. Ticks were pulled from all mice and pooled into four

biological replicates 1 day after placement (14 ticks per replicate), 2 days after placement (12

ticks per replicate), and 3 days after placement (6 ticks per replicate) and collected from cages 4

days after placement (7 ticks per replicate). Shortly after collection, ticks were washed with water

and placed in a 2 mL glass dounce grinder (Kimble) in 500 μL of phosphate-buffered saline (PBS).

Ticks were homogenized first with the large clearance pestle and then the small clearance pestle.

The homogenate was transferred to a 1.5 mL Eppendorf tube and 500 μL of PBS was added to

total 1 mL. At this stage, 50 μL of homogenate was removed as an input sample and mixed with

500 μL of TRIzol (Invitrogen) for RNA extraction. 2 μL of α*Bb* antibody (Invitrogen: PA1-73004;

RRID: AB_1016668) was added to the homogenate, which was then placed on a nutator at 4°C

for 30 minutes. During incubation, 50 μL of Dynabeads Protein G (Invitrogen) per sample were

washed twice in PBS. After incubation with the antibody, the homogenate and antibody mixture

were added to the beads. This mixture was placed on a nutator at 4°C for 30 minutes. Tubes were

then placed on a magnet to secure beads, and the homogenate was removed and saved to create

depleted samples. The depleted homogenate was centrifuged at 8,000 x g for 7 minutes, 900 μL

of supernatant was removed, and 500 μL of TRIzol was added to the pellet to create depleted

samples. The beads were washed twice with 1 mL of PBS, resuspending the beads each time.

The second wash was removed and 500 μL of TRIzol was added to the beads to create enriched

samples. RNA was extracted from all input, enriched, and depleted samples using the Zymo

Direct-zol RNA Microprep Kit with on-column DNase treatment (Zymo Research). The step-by-

step *Bb* enrichment protocol is available at: dx.doi.org/10.17504/protocols.io.36wgqjrbovk5/v1.

Enrichment of *Bb* from culture

To test whether the *Bb* enrichment process altered gene expression levels, we performed the

enrichment protocol on cultured *Bb*. Tubes of *Bb* in BSK II media were grown to 9x10^4 *Bb*/mL at

35°C. 1 mL of culture was spun down at 8,000 x g for 7 minutes, media was removed, *Bb* were

washed in 1mL of PBS and spun again. Pelleted *Bb* were resuspended in 1 mL of fresh PBS.

These samples were used as starting homogenate for the *Bb* enrichment protocol and input,

enriched, and depleted fractions were collected as above. RNA-seq libraries from these samples

were prepared and sequenced as below.

RNA-seq library preparation and sequencing

To make RNA-seq libraries from enriched *Bb* RNA, 50 ng of total RNA was used as input into the

NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (New England BioLabs). Libraries

were prepared following the manufacturer's protocol for use of the kit with purified mRNA or rRNA-

depleted RNA, despite starting with total RNA. Libraries were barcoded using NEBNext Multiplex

Oligos for Illumina Dual Index (New England BioLabs).

To deplete reads from the libraries that were from tick, *Bb*, or mouse rRNA, we used Depletion of

Abundant Sequences by Hybridization (DASH) (Gu et al., 2016), which targets Cas9 to unwanted

reads in RNA-seq libraries using custom dual-guide RNAs (dgRNAs). The dgRNAs targeted short

sequences within tick rRNA, mouse rRNA, and *Bb* rRNA that were designed using DASHit

software (Dynerman et al., 2020). We ordered crRNA oligos (**Table S9**) that targeted these

sequences (Ring et al., 2022). To transcribe the dgRNAs, we followed the protocol for *In Vitro*

Transcription for dgRNA V2 (Lyden et al., 2019b) as follows. Both tracrRNA and pooled crRNA

DNA templates were annealed to equimolar amounts of T7 primer by heating to 95°C for 2

minutes and slowly cooling to room temperature. Annealed templates were used in 1 mL *in vitro*

transcription reactions with: 120 µL10X T7 buffer (400 mM Tris pH 7.9, 200 mM MgCl2, 50 mM

DTT, 20 mM spermidine (Sigma)), 100 µL of T7 enzyme (custom prepped enzyme from E.

Crawford, diluted 1:100 in T7 buffer, final concentration: 100 µg/mL), 300 µl NTPs (25 mM each,

Thermo Fisher Scientific), 4 µg of annealed crRNA template or 8 µg of annealed tracrRNA

template, and water to 1 mL. *In vitro* transcription was performed for 2 hours at 37°C. Reactions

were purified twice with the Zymo RNA Clean & Concentrator-5 Kit (Zymo Research). To form the

dgRNA complex for DASH, crRNA and tracrRNAs were diluted to 80 µM, mixed in equimolar

amounts and annealed by heating to 95ºC for 30 seconds and cooling slowly to room temperature.

After transcription of dgRNAs, we performed DASH Protocol Version 4 (Lyden et al., 2019a) on

each library individually as follows. Cas9 and transcribed dgRNAs were prepped by mixing: 2.5

µL 10X Cas9 buffer, 5 µL 20 µM Cas9 (New England BioLabs), and 5 µL of 40 µM transcribed

dgRNAs. The mixture was incubated at 37°C for 5 minutes before 7.5 µL of RNA-seq library (2.8

nM) was added. The mixture was incubated at 37°C for 1 hour and then purified with Zymo DNA

Clean & Concentrator-5 (Zymo Research) following the PCR product protocol and eluting DNA

into 10.5 µL of water. During cleanup, Cas9 was again mixed with buffer and dgRNAs and

incubated at 37°C for 5 minutes. Following cleanup, eluted DNA was added to the second Cas9-

dgRNA mixture and incubated at 37°C for 1 hour for a second time. Then, 1 µL of proteinase K

(New England Biolabs) was added, and the mixture was incubated at 50°C for 15 min. The

libraries were then purified with 0.9X volume of sparQ PureMag Beads (QuantaBio) following the

standard protocol, eluting in 24 µL of water. rRNA-depleted RNA-seq libraries were then amplified

in a BioRad CFX96 using the Kapa HiFi Real-Time Amplification Kit (Roche) in a 50 µL reaction

with 25 µL master mix, 23 µL of the DASHed library pool, and 2 µL of a 25 µM mix of Illumina P5

(5'-AATGATACGGCGACCACCGAGATCT) and P7 (5'- CAAGCAGAAGACGGCATACGAGAT)

primers. The qPCR program for amplification was as follows: 98°C for 45 sec (1 cycle), (98°C for

15 sec, 63°C for 30 sec, 72°C for 45 sec, plate read, 72°C for 20 sec) for 10 cycles (day 3 and

day 4 samples) or 11 cycles (day 1 and day 2 samples). The libraries were removed from cycling

conditions before leaving the exponential phase of amplification and then purified with 0.9X

volume of sparQ PureMag Beads according to the standard protocol.

Following DASH, RNA-seq libraries were sequenced on an Illumina NovaSeq S2 (2 lanes) with

paired-end 100 base pair reads. Libraries from *in vitro* cultured control experiment were

sequenced on an Illumina NextSeq with paired-end 75 base pair reads. FASTQ files and raw *Bb*

read counts for *in vitro* control experiment (GSE217146) and *ex vivo* experiment (GSE216261)

have been deposited in NCBI's Gene Expression Omnibus (Edgar et al., 2002) under SuperSeries

accession number GSE217236.

RNA-seq data analysis

DASH

To measure the success of our rRNA depletion through DASH, we used DASHit software

(Dynerman et al., 2020) to determine the percentage of reads that would be DASHable by our

guide RNAs (Ring et al., 2022). For pre-DASH data, we sequenced the input of each of our RNA-

seq libraries before performing DASH on a MiSeq V2 Micro (Illumina). We tested DASHability on

a random subset of 200,000 paired-end reads chosen by seqtk (https://github.com/lh3/seqtk) from

each pre- and post-DASH library. Paired t test comparing DASHable reads before and after DASH

was performed using GraphPad Prism v9.5.1.

Differential expression analysis

To map our RNA-seq data to *Bb*, we wanted to optimize for mapping reads that came from the

many paralogous gene families found across the plasmids of the genome. We used the

pseudoalignment tool Salmon v1.2.1 (Patro et al., 2017), which is used to accurately map reads

coming from different isoforms of the same gene, for this reason. While using Salmon to map to

CDS sequences may improve mapping to paralogous genes, it may also have a tradeoff of

reduced mapping of reads that fall on the ends of genes that reside in operons. Nevertheless, all

samples should be similarly affected, and any undercounting should not change differential

expression results. Reads were first trimmed of bases with quality scores less than 20 using

Cutadapt (Martin, 2011) via Trim Galore v0.6.5 (https://github.com/FelixKrueger/TrimGalore).

Reads were mapped to *Bb* CDS sequences as a reference transcriptome: NCBI Genbank
GCA_000008685.2 ASM868v2 (with plasmids lp5, cp9, and lp56 removed as they are not present
in B31-S9) using Salmon with the following parameters: --validateMappings --seqBias --
gcBias. Before mapping, the transcriptome was indexed using the Salmon index command with
the whole genome as decoys and the parameter --keepDuplicates to keep all duplicate genes.
636
Read counts from Salmon were used as input into DESeq2 v1.24.0 (Love et al., 2014) for
differential expression analysis in R version 3.6.1. DESeq2 function PlotPCA() was used to create
a PCA plot from read counts after running the varianceStabilizingTransformation() function. For
differential expression analysis between days, a DESeq object was created from count data using
the DESeq() function. The lfcShrink() function with the apeglm method (Zhu et al., 2018) was
used to calculate fold changes between days. DESeq2 uses a Benjamini-Hochberg multiple
testing correction, and we focused the majority of our analysis on genes that had an adjusted p-
value < 0.05 and used an additional cutoff requiring genes to change two-fold between conditions.
Code used for differential expression analysis is available at: https://github.com/annesapiro/Bb-
tick-feeding.
647

Mapping to other species
To determine the source of non-*Bb* reads in our RNA-seq libraries, trimmed reads were mapped
to tick and mouse genomes using STAR v2.7.3a (Dobin et al., 2012). The *I. scapularis* ISE6
genome (RefSeq assembly GCF_002892825.2, ISE6_asm2.2_dedeplicated) (Miller et al., 2018)
was indexed using STAR run mode genomeGenerate with option --genomeChrBinNbits 18. The
Mus musculus genome GRCm39 (RefSeq assembly GCF_000001635.27) was indexed using
STAR run mode genomeGenerate with basic options. Reads were mapped using STAR to each
genome using basic options. The percentage of reads that mapped to these genomes was
determined by adding the percentage of uniquely mapped reads, reads mapped to multiple loci,
and reads mapped to too many loci. To identify the potential source of reads that did not map to
tick, mouse, or *Bb* in day 4 samples, one million reads from day 1 and day 4 libraries were used
as input into CZ ID (Kalantar et al., 2020), which determined that a large number of reads mapped
to bacterial species *Pseudomonas fulva*. Full RNA-seq libraries were then mapped to the *P. fulva*
genome (NCBI GenBank GCF_001186195.1 ASM118619v1), using the standard options of
Bowtie2 (Langmead and Salzberg, 2012) to calculate the overall alignment rate.
663

Comparisons to other studies

Genes identified in previous studies were compared to time course expression changes. Here we
considered RpoS-regulated genes as those found in Grassmann et al. 2023 that were upregulated
by RpoS in both fed nymphs and DMCs and those upregulated by RpoS only during tick
transmission (Grassmann et al. Supplemental Tables 5 and 6). RpoS did not suppress the
expression of any genes in fed nymphs in the study. Genes up- and down-regulated by RpoS in
DMCs only (Grassmann et al. Supplemental Tables 7 and 8) are noted in **Table S3** and **Table S4**
for reference along with genes found to be regulated by RpoS in DMCs in Caimano et al. 2019
(Tables 2 and 3), which were used for RpoS comparisons in previous versions of this study. Rrp1
up- and down-regulated genes were those identified *in vitro* in Caimano et al. 2015, Table S2.
Rel$_{Bbu}$ up- and down-regulated genes were examined by Drecktrah et al. 2015 in three different
in vitro conditions: starvation (Tables S6 and S9), recovery (Tables S7 and S10), and stationary
phase (Tables S5 and S8). For simplicity, we considered genes as Rel$_{Bbu}$-regulated if they were
up- or down-regulated in one or more of these conditions (**Tables S3** and **S4**). One gene was
regulated in opposing directions across conditions and is noted in our tables as "both" and was
excluded from the comparison analysis. Genes changing between "unfed tick" and "fed tick"
culture conditions in Revel et al. 2002 were those in Table 3. Genes from Ojaimi et al. 2003 Table
4 with increased expression *in vitro* at 35°C relative to 25°C were considered higher in "fed tick"
while those in Ojaimi et al. 2003 Table 5 with increased expression at 25°C relative to 35°C were
considered higher in "unfed tick." Genes more highly expressed in nymphs than DMCs from Iyer
et al. 2015 were found in Table S4, and genes more highly expressed in DMCs than nymphs were
found in Table S8. Genes differentially expressed between nymphs and DMCs in Grassmann et
al. 2023 were determined from the DESeq2 comparison between WT DMC vs Fed Nymphs found
in Supplemental Table 3, in accordance with their cutoffs of at least a three-fold difference and q-
value < 0.05. As many of these studies used different strains of *Bb* and different genome
annotations, some genes were not examined here as they were not present in the B31-S9 strain
used here.

Gene classification
To classify genes into functional groups, functional categories were sourced from (Drecktrah et
al., 2015) where available. Other gene functions were sourced from (Fraser et al., 1997). Genes
found within the co-transcribed "late" bacteriophage operon (Zhang and Marconi, 2005) were
considered "bacteriophage" even if their function is unknown. Outer surface proteins were those
found in Dowdell et al., 2017 plus additional outer surface proteins listed in Iyer et al., 2015 that
were also found in Dowdell et al. 2017 Supporting Table S2 categories SpII and SpI as evidence

of outer surface localization. Outer surface and periplasmic lipoproteins were classified as "cell

envelope" in the absence of other classifications. Gene family information from (Casjens et al.,

2000) was considered to aid in classification. **Table S4** contains the classification source for each

gene.

RT-qPCR measuring *Bb* enrichment

To test the efficacy of the *Bb* enrichment protocol, RT-qPCR was used to quantify *Bb flaB* and *I.*

scapularis gapdh transcript levels in enriched and depleted fractions. cDNA was synthesized from

8 µL of RNA extracted from *Bb* enrichment samples and their matched depleted samples from

day 2, day 3, and day 4 post-attachment using the qScript cDNA Synthesis Kit (Quantabio). cDNA

was diluted 2X before use in qPCR. To measure *flaB* copies, standards of known concentration

were created from purified PCR products. These standards were made from PCR with primers

with the following sequences: 5'-CACATATTCAGATGCAGACAGAGGTTCTA and 5'-

GAAGGTGCTGTAGCAGGTGCTGGCTGT. A dilution series with ten-fold dilutions between 10^6

copies and 10^1 copies of this PCR template was run alongside enriched and depleted samples.

qPCR was performed using Taqman Universal PCR Master Mix (Applied Biosystems). The

primers used to amplify *flaB* were: 5'- TCTTTTCTCTGGTGAGGGAGCT and 5'-

TCCTTCCTGTTGAACACCCTCT (used at 900 nM) and the probe was /56-

FAM/AAACTGCTCAGGCTGCACCGGTTC/36-TAMSp (used at 250 nM). For tick *gapdh* RT-

qPCR, the cDNA samples were diluted an additional 2X. Standards of known concentration were

created using the qPCR primer sequences: 5'-TTCATTGGAGACACCCACAG and 5'-

CGTTGTCGTACCACGAGATAA (used at 900 nM). qPCR was performed using PowerUp SYBR

Green Master Mix (Applied Biosystems). For both *flaB* and *gapdh*, the number of copies in each

sample was calculated based on the standards of known concentration. Three technical replicates

were averaged from each of four biological replicates at each time point tested. We totaled the

number of copies in each matched enriched and depleted fraction to calculate the percentage of

flaB or *gapdh* that was found in either sample. All qPCR was performed on the QuantStudio3

Real-Time PCR System (Applied Biosystems). Paired t tests were performed using GraphPad

Prism v9.5.1.

Immunofluorescence microscopy

To test whether the α*Bb* antibody recognized *Bb* inside of the tick, ticks at each day of feeding

were crushed in 50 µL of PBS. 10 µL of lysate was spotted onto slides and allowed to air dry

before slides were heated briefly three times over a flame. Heat fixed slides were then treated

with acetone for one hour. Slides were incubated with αBb primary antibody (1:100 diluted in PBS

+ 0.75% BSA) for 30 minutes at 37°C in a humid chamber. A control without primary antibody

was also used for each day. Slides were washed once in PBS for 15 minutes at room temperature,

then rinsed in distilled water and air dried. Anti-rabbit IgG Alexa 488 (Invitrogen: A-11008; RRID:

AB_143165) diluted 1:100 in PBS + 0.75% BSA was added for 30 minutes at 37°C in a humid

slide chamber. Slides were washed in PBS for 15 minutes at room temperature three times,

adding 1:100 Propidium Iodide (Invitrogen) during the second wash. Slides were then rinsed with

distilled water and air dried before the addition of mounting media (Fluoromount-G,

SouthernBiotech) and cover slips. Fluorescence imaging was performed on a Nikon Ti2 inverted

microscope for widefield epifluorescence using a 100X/1.40 objective. Images were captured with

NIS-Elements AR View 5.20 and then processed with ImageJ software (Schneider et al., 2012).

No strong florescence signal was observed on the control slides without primary antibody.

Mass spectrometry of *Bb*-enriched samples

To identify which tick proteins were found in samples after *Bb* enrichment across feeding, both

uninfected and infected ticks were fed on mice. Three biological replicates of uninfected ticks one

day after attachment (11 ticks per replicate), infected ticks one day after attachment (27 ticks per

replicate), uninfected ticks four days after attachment (8 ticks per replicate), and infected ticks

four days after attachment (16 ticks per replicate) were collected. Before αBb enrichment, the

uninfected tick samples were mixed with *Bb* grown in culture that was washed with PBS (3×10^4

Bb one day after attachment and 3×10^6 *Bb* four days after attachment) and mixed lysates were

rotated at room temperature for 30 minutes. Infected tick samples underwent the *Bb* enrichment

process immediately. The enrichment process followed the same protocol used for RNA-seq.

Sample volumes were increased to 1 mL as needed, and then 2 µL of αBb antibody (Invitrogen:

PA1-73004; RRID: AB_1016668) was added, and samples were rotated at 4°C for 30 minutes.

50 µL of Dynabeads Protein G per sample were washed in PBS during this incubation and added

to the lysates, which were rotated at 4°C for 30 minutes. The beads were washed twice with 1 mL

of PBS, and then placed into 50 µL of lysis buffer (iST LYSE, PreOmics). Samples were boiled at

95°C for 5 minutes, and lysates were removed from beads and frozen for mass spectrometry

preparation.

For mass spectrometry, a nanoElute was attached in line to a timsTOF Pro equipped with a

CaptiveSpray Source (Bruker). Chromatography was conducted at 40°C through a 25 cm

reversed-phase C18 column (PepSep) at a constant flowrate of 0.5 µL min−1. Mobile phase A

was 98/2/0.1% water/MeCN/formic acid (v/v/v) and phase B was MeCN with 0.1% formic acid

(v/v). During a 108 min method, peptides were separated by a 3-step linear gradient (5% to 30%

B over 90 min, 30% to 35% B over 10 min, 35% to 95% B over 4 min) followed by a 4 min isocratic

flush at 95% for 4 min before washing and a return to low organic conditions. Experiments were

run as data-dependent acquisitions with ion mobility activated in PASEF mode. MS and MS/MS

spectra were collected with m/z 100 to 1700 and ions with z = +1 were excluded.

Raw data files were searched using PEAKS Online Xpro 1.6 (Bioinformatics Solutions Inc.). The

precursor mass error tolerance and fragment mass error tolerance were set to 20 ppm and 0.03

respectively. The trypsin digest mode was set to semi-specific and missed cleavages was set to

2. The *I. scapularis* reference proteome (Proteome ID UP000001555, taxon 6945) and *Bb*

reference proteome (Proteome ID UP000001807, strain ATCC 35210/B31) was downloaded from

Uniprot, totaling 21,774 entries. The *I. scapularis* proteome was the primary search reference and

the *Bb* was used as a secondary to identify any bacterial proteins present. Carbamidomethylation

was selected as a fixed modification. Oxidation (M) and Deamidation (NQ) was selected as a

variable modification.

Experiments were performed in biological triplicate, with samples being a single run on the

instrument. Proteins present in a database search ($-10\log$(p-value) $\geq$ 20, 1% peptide and protein

FDR) were subjected to the following filtration process. Proteins were filtered to include only those

found in 2 out of 3 biological replicates, within each respective day (one or four days after

attachment). The mean area of proteins found in uninfected and infected samples was calculated.

Proteins with missing values (i.e. not identified in a sample) were set to 1. The ratio of mean area

for each protein was calculated as infected/uninfected, and enriched proteins were identified by

having an infected to uninfected ratio greater than 2 within their respective feeding day (one or

four days after attachment).

To classify the identified proteins into functional groups, a PANTHER Overrepresentation Test

(released 07/12/2022) was used (Mi et al., 2019) with PANTHER version 17.0 (released

02/22/2022) (Thomas et al., 2022). All *I. scapularis* genes in the database were used as the

reference list to and all proteins enriched on each day were analyzed for their PANTHER Protein

Class. The test type used was Fisher's Exact, calculating a false discovery rate as the correction.

Raw data files and searched datasets are available on the Mass Spectrometry Interactive Virtual
Environment (MassIVE), a full member of the Proteome Xchange consortium under the identifier:
MSV000090560.
803

DATA AVAILABILITY
For sequencing data, FASTQ files and raw *Bb* read counts for *in vitro* control experiment
(GSE217146) and *ex vivo* experiment (GSE216261) have been deposited in NCBI's GEO
database under SuperSeries accession number GSE217236. For mass spectrometry data, raw
data files and searched datasets are available on the Mass Spectrometry Interactive Virtual
Environment (MassIVE) under the identifier: MSV000090560. Code used for data analysis is
available at: https://github.com/annesapiro/Bb-tick-feeding.
811

SUPPLEMENTAL TABLES
Table S1. Overview of mapping statistics from 16 *Bb* sequencing samples.
Table S2. Differential expression analysis results between *in vitro* cultured *Bb* before and after
Bb enrichment.
Table S3. Transcriptome-wide differential expression analysis results from *Bb* across tick
feeding timepoints.
Table S4. Two-fold differentially expressed *Bb* genes from across tick feeding timepoints.
Table S5. Transcripts per million (TPM) for all *Bb* genes across feeding timepoints.
Table S6. Mass spectrometry analysis for *Bb*-enriched samples.
Table S7. Annotation and GO term enrichment for tick proteins enriched on feeding day 1.
Table S8. Annotation and GO term enrichment for tick proteins enriched on feeding day 4.
Table S9. crRNAs targeting tick, mouse, and *Bb* rRNA sequences used in DASH.
824

ACKNOWLEDGMENTS
We are grateful to all members of the Chou lab for their feedback throughout the project,
specifically to Fauna Yarza and Patrick Rockefeller Grimes for assistance with tick feeding and
Ethel Ethel Enoex-Godonoo for administrative assistance. We thank Amy Lyden and Emily
Crawford for assistance with and reagents for DASH along with Olga Botvinnik and The Chan
Zuckerberg Biohub sequencing team for their input and sequencing assistance. We thank Patricia
Rosa, Jenny Wachter, Scott Samuels, and Meghan Lybecker for their feedback on the project.
We also thank William Hatleberg for assistance with figure schematics. This work was funded by:
a Life Sciences Research Foundation fellowship from the SVCF-Wave Fund to Anne Sapiro, a

Beckman Young Investigator award from the Arnold and Mabel Beckman Foundation to Balyn

Zaro, grants from CZ Biohub, the Pew Biomedical Research Foundation, and NIH funding

1R01AI132851 to Seemay Chou, and NIH INBRE funding P20GM103474 to Patrick Secor and

Margie Kinnersley.

COMPETING INTERESTS

Seemay Chou is president and CEO of Arcadia Biosciences.

REFERENCES

Barbour AG. 1988. Plasmid analysis of *Borrelia burgdorferi*, the Lyme disease agent. *J Clin*
Microbiol 26:475–478. doi:10.1128/jcm.26.3.475-478.1988

Barillas-Mury C, Ribeiro JMC, Valenzuela JG. 2022. Understanding pathogen survival and
transmission by arthropod vectors to prevent human disease. *Science* 377:eabc2757.
doi:10.1126/science.abc2757

Battisti JM, Bono JL, Rosa PA, Schrumpf ME, Schwan TG, Policastro PF. 2008. Outer Surface
Protein A Protects Lyme Disease Spirochetes from Acquired Host Immunity in the Tick
Vector. *Infect Immun* 76:5228–5237. doi:10.1128/iai.00410-08

Blevins JS, Hagman KE, Norgard MV. 2008. Assessment of decorin-binding protein A to the
infectivity of *Borrelia burgdorferi* in the murine models of needle and tick infection. *Bmc*
Microbiol 8:82. doi:10.1186/1471-2180-8-82

Bykowski T, Woodman ME, Cooley AE, Brissette CA, Brade V, Wallich R, Kraiczy P, Stevenson
B. 2007. Coordinated Expression of *Borrelia burgdorferi* Complement Regulator-Acquiring
Surface Proteins during the Lyme Disease Spirochete's Mammal-Tick Infection Cycle. *Infect*
Immun 75:4227–4236. doi:10.1128/iai.00604-07

Caimano MJ, Dunham-Ems S, Allard AM, Cassera MB, Kenedy M, Radolf JD. 2015. Cyclic di-
GMP Modulates Gene Expression in Lyme Disease Spirochetes at the Tick-Mammal
Interface To Promote Spirochete Survival during the Blood Meal and Tick-to-Mammal
Transmission. *Infect Immun* 83:3043–3060. doi:10.1128/iai.00315-15

Caimano MJ, Groshong AM, Belperron A, Mao J, Hawley KL, Luthra A, Graham DE, Earnhart
CG, Marconi RT, Bockenstedt LK, Blevins JS, Radolf JD. 2019. The RpoS Gatekeeper in
Borrelia burgdorferi: An Invariant Regulatory Scheme That Promotes Spirochete Persistence
in Reservoir Hosts and Niche Diversity. *Front Microbiol* 10:1923.
doi:10.3389/fmicb.2019.01923

Casjens S, Palmer N, Van Vugt R, Huang WM, Stevenson B, Rosa P, Lathigra R, Sutton G,
Peterson J, Dodson RJ, Haft D, Hickey E, Gwinn M, White O, Fraser CM. 2000. A bacterial
genome in flux: the twelve linear and nine circular extrachromosomal DNAs in an infectious

isolate of the Lyme disease spirochete *Borrelia burgdorferi*. *Mol Microbiol* 35:490–516. doi:10.1046/j.1365-2958.2000.01698.x

Casjens SR, Mongodin EF, Qiu W-G, Luft BJ, Schutzer SE, Gilcrease EB, Huang WM, Vujadinovic M, Aron JK, Vargas LC, Freeman S, Radune D, Weidman JF, Dimitrov GI, Khouri HM, Sosa JE, Halpin RA, Dunn JJ, Fraser CM. 2012. Genome Stability of Lyme Disease Spirochetes: Comparative Genomics of *Borrelia burgdorferi* Plasmids. *Plos One* 7:e33280. doi:10.1371/journal.pone.0033280

de Silva AM, Fikrig E. 1995. Growth and Migration of *Borrelia burgdorferi* in *Ixodes* Ticks during Blood Feeding. *Am J Tropical Medicine Hyg* 53:397–404. doi:10.4269/ajtmh.1995.53.397

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2012. STAR: ultrafast universal RNA-seq aligner. *Bioinform Oxf Engl* 29:15–21. doi:10.1093/bioinformatics/bts635

Dowdell AS, Murphy MD, Azodi C, Swanson SK, Florens L, Chen S, Zückert WR. 2017. Comprehensive Spatial Analysis of the *Borrelia burgdorferi* Lipoproteome Reveals a Compartmentalization Bias toward the Bacterial Surface. *J Bacteriol* 199:e00658-16. doi:10.1128/jb.00658-16

Drecktrah D, Lybecker M, Popitsch N, Rescheneder P, Hall LS, Samuels DS. 2015. The *Borrelia burgdorferi* RelA/SpoT Homolog and Stringent Response Regulate Survival in the Tick Vector and Global Gene Expression during Starvation. *Plos Pathog* 11:e1005160. doi:10.1371/journal.ppat.1005160

Dunham-Ems SM, Caimano MJ, Pal U, Wolgemuth CW, Eggers CH, Balic A, Radolf JD. 2009. Live imaging reveals a biphasic mode of dissemination of *Borrelia burgdorferi* within ticks. *J Clin Invest* 119:3652–3665. doi:10.1172/jci39401

Dynerman D, Lyden A, Quan J, Caldera S, McGeever A, Dimitrov B, King R, Cirolia G, Tan M, Sit R, van den Berge M, Kerstjens HAM, Faiz A, Christenson S, Langelier C, DeRisi J, Crawford E. 2020. Designing and implementing programmable depletion in sequencing libraries with DASHit. *Biorxiv* 2020.01.12.891176. doi:10.1101/2020.01.12.891176

Edgar R, Domrachev M, Lash AE. 2002. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res* 30:207–210. doi:10.1093/nar/30.1.207

Eggers CH, Samuels DS. 1999. Molecular evidence for a new bacteriophage of *Borrelia burgdorferi*. *J Bacteriol* 181:7308–7313. doi:10.1128/jb.181.23.7308-7313.1999

Fraser CM, Casjens S, Huang WM, Sutton GG, Clayton R, Lathigra R, White O, Ketchum KA, Dodson R, Hickey EK, Gwinn M, Dougherty B, Tomb J-F, Fleischmann RD, Richardson D, Peterson J, Kerlavage AR, Quackenbush J, Salzberg S, Hanson M, van Vugt R, Palmer N, Adams MD, Gocayne J, Weidman J, Utterback T, Watthey L, McDonald L, Artiach P, Bowman C, Garland S, Fujii C, Cotton MD, Horst K, Roberts K, Hatch B, Smith HO, Venter JC. 1997. Genomic sequence of a Lyme disease spirochaete, *Borrelia burgdorferi*. *Nature* 390:580–586. doi:10.1038/37551

Gilmore RD, Howison RR, Dietrich G, Patton TG, Clifton DR, Carroll JA. 2010. The *bba64* gene
of *Borrelia burgdorferi*, the Lyme disease agent, is critical for mammalian infection via tick
bite transmission. *P Natl Acad Sci Usa* 107:7515–7520. doi:10.1073/pnas.1000268107

Gilmore RD, Mbow ML, Stevenson B. 2001. Analysis of *Borrelia burgdorferi* gene expression
during life cycle phases of the tick vector *Ixodes scapularis*. *Microbes Infect* 3:799–808.
doi:10.1016/s1286-4579(01)01435-6

Grassmann AA, Tokarz R, Golino C, McLain MA, Groshong AM, Radolf JD, Caimano MJ. 2023.
BosR and PlzA reciprocally regulate RpoS function to sustain *Borrelia burgdorferi* in ticks
and mammals. *J Clin Invest* 133. doi:10.1172/jci166710

Gu W, Crawford ED, O'Donovan BD, Wilson MR, Chow ED, Retallack H, DeRisi JL. 2016.
Depletion of Abundant Sequences by Hybridization (DASH): using Cas9 to remove
unwanted high-abundance species in sequencing libraries and molecular counting
applications. *Genome Biol* 17:41. doi:10.1186/s13059-016-0904-5

Haas BJ, Chin M, Nusbaum C, Birren BW, Livny J. 2012. How deep is deep enough for RNA-
Seq profiling of bacterial transcriptomes? *Bmc Genomics* 13:734. doi:10.1186/1471-2164-13-
734

He M, Ouyang Z, Troxell B, Xu H, Moh A, Piesman J, Norgard MV, Gomelsky M, Yang XF.
2011. Cyclic di-GMP is Essential for the Survival of the Lyme Disease Spirochete in Ticks.
Plos Pathog 7:e1002133. doi:10.1371/journal.ppat.1002133

Hübner A, Yang X, Nolen DM, Popova TG, Cabello FC, Norgard MV. 2001. Expression of
Borrelia burgdorferi OspC and DbpA is controlled by a RpoN–RpoS regulatory pathway. *Proc
National Acad Sci* 98:12724–12729. doi:10.1073/pnas.231442498

Iyer R, Caimano MJ, Luthra A, Axline D Jr, Corona A, Iacobas DA, Radolf JD, Schwartz I. 2015.
Stage-specific global alterations in the transcriptomes of Lyme disease spirochetes during
tick feeding and following mammalian host adaptation. *Mol Microbiol* 95:509–538.
doi:10.1111/mmi.12882

Kalantar KL, Carvalho T, de Bourcy CFA, Dimitrov B, Dingle G, Egger R, Han J, Holmes OB,
Juan Y-F, King R, Kislyuk A, Lin MF, Mariano M, Morse T, Reynoso LV, Cruz DR, Sheu J,
Tang J, Wang J, Zhang MA, Zhong E, Ahyong V, Lay S, Chea S, Bohl JA, Manning JE, Tato
CM, DeRisi JL. 2020. IDseq—An open source cloud-based pipeline and analysis service for
metagenomic pathogen detection and monitoring. *Gigascience* 9:giaa111.
doi:10.1093/gigascience/giaa111

Kasumba IN, Bestor A, Tilly K, Rosa PA. 2016. Virulence of the Lyme disease spirochete before
and after the tick bloodmeal: a quantitative assessment. *Parasite Vector* 9:129.
doi:10.1186/s13071-016-1380-1

Kenedy MR, Lenhart TR, Akins DR. 2012. The role of *Borrelia burgdorferi* outer surface
proteins. *Fems Immunol Medical Microbiol* 66:1–19. doi:10.1111/j.1574-695x.2012.00980.x

Kovacs-Simon A, Titball RW, Michell SL. 2010. Lipoproteins of Bacterial Pathogens. *Infect
Immun* 79:548–561. doi:10.1128/iai.00682-10

948 Kung F, Anguita J, Pal U. 2013. *Borrelia burgdorferi* and tick proteins supporting pathogen
949  persistence in the vector. *Future Microbiol* 8:41–56. doi:10.2217/fmb.12.121

950 Kurokawa C, Lynn GE, Pedra JHF, Pal U, Narasimhan S, Fikrig E. 2020. Interactions between
951  *Borrelia burgdorferi* and ticks. *Nat Rev Microbiol* 18:587–600. doi:10.1038/s41579-020-0400-
952  5

953 Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods*
954  9:357–359. doi:10.1038/nmeth.1923

955 Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
956  RNA-seq data with DESeq2. *Genome Biol* 15:550. doi:10.1186/s13059-014-0550-8

957 Lyden A, Crawford E, Quan J, Caldera S, Dynerman D. 2019a. DASH Protocol v4.
958  doi:10.17504/protocols.io.6rjhd4n

959 Lyden A, Crawford E, Quan J, Caldera S, Pesce-Ares L. 2019b. In Vitro Transcription for
960  dgRNA v2. doi:10.17504/protocols.io.3bpgimn

961 Manning JE, Cantaert T. 2019. Time to Micromanage the Pathogen-Host-Vector Interface:
962  Considerations for Vaccine Development. *Nato Adv Sci Inst Se* 7:10.
963  doi:10.3390/vaccines7010010

964 Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
965  *Embnet J* 17:10–12. doi:10.14806/ej.17.1.200

966 Mi H, Muruganujan A, Huang X, Ebert D, Mills C, Guo X, Thomas PD. 2019. Protocol Update for
967  large-scale genome and gene function analysis with the PANTHER classification system
968  (v.14.0). *Nat Protoc* 14:703–721. doi:10.1038/s41596-019-0128-8

969 Miller JR, Koren S, Dilley KA, Harkins DA, Stockwell TB, Shabman RS, Sutton GG. 2018. A
970  draft genome sequence for the *Ixodes scapularis* cell line, ISE6. [version 1; peer review: 2
971  approved]. *F1000research* 7:297. doi:10.12688/f1000research.13635.1

972 Narasimhan S, Santiago F, Koski RA, Brei B, Anderson JF, Fish D, Fikrig E. 2002. Examination
973  of the *Borrelia burgdorferi* Transcriptome in *Ixodes scapularis* during Feeding. *J Bacteriol*
974  184:3122–3125. doi:10.1128/jb.184.11.3122-3125.2002

975 Ohnishi J, Piesman J, de Silva AM. 2001. Antigenic and genetic heterogeneity of *Borrelia*
976  *burgdorferi* populations transmitted by ticks. *Proceedings of the National Academy of*
977  *Sciences* 98:670–675. doi:10.1073/pnas.98.2.670

978 Ojaimi C, Brooks C, Casjens S, Rosa P, Elias A, Barbour A, Jasinskas A, Benach J, Katona L,
979  Radolf J, Caimano M, Skare J, Swingle K, Akins D, Schwartz I. 2003. Profiling of
980  Temperature-Induced Changes in *Borrelia burgdorferi* Gene Expression by Using Whole
981  Genome Arrays. *Infect Immun* 71:1689–1705. doi:10.1128/iai.71.4.1689-1705.2003

Ojaimi C, Mulay V, Liveris D, Iyer R, Schwartz I. 2005. Comparative Transcriptional Profiling of
Borrelia burgdorferi Clinical Isolates Differing in Capacities for Hematogenous Dissemination.
Infect Immun 73:6791–6802. doi:10.1128/iai.73.10.6791-6802.2005

Ouyang Z, Narasimhan S, Neelakanta G, Kumar M, Pal U, Fikrig E, Norgard MV. 2012.
Activation of the RpoN-RpoS regulatory pathway during the enzootic life cycle of *Borrelia*
burgdorferi. *Bmc Microbiol* 12:44. doi:10.1186/1471-2180-12-44

Pal U, Kitsou C, Drecktrah D, Yaş OB, Fikrig E. 2021. Interactions Between Ticks and Lyme
Disease Spirochetes. *Curr Issues Mol Biol* 42:113–144. doi:10.21775/cimb.042.113

Pal U, Li X, Wang T, Montgomery RR, Ramamoorthi N, de Silva AM, Bao F, Yang X, Pypaert M,
Pradhan D, Kantor FS, Telford S, Anderson JF, Fikrig E. 2004. TROSPA, an *Ixodes*
scapularis Receptor for *Borrelia burgdorferi*. *Cell* 119:457–468.
doi:10.1016/j.cell.2004.10.027

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017. Salmon provides fast and bias-
aware quantification of transcript expression. *Nat Methods* 14:417–419.
doi:10.1038/nmeth.4197

Patton TG, Brandt KS, Nolder C, Clifton DR, Carroll JA, Gilmore RD. 2013. *Borrelia burgdorferi*
bba66 Gene Inactivation Results in Attenuated Mouse Infection by Tick Transmission. *Infect*
Immun 81:2488–2498. doi:10.1128/iai.00140-13

Phelan JP, Kern A, Ramsey ME, Lundt ME, Sharma B, Lin T, Gao L, Norris SJ, Hyde JA, Skare
JT, Hu LT. 2019. Genome-wide screen identifies novel genes required for *Borrelia*
burgdorferi survival in its Ixodes tick vector. *Plos Pathog* 15:e1007644.
doi:10.1371/journal.ppat.1007644

Radolf JD, Caimano MJ, Stevenson B, Hu LT. 2012. Of ticks, mice and men: understanding the
dual-host lifestyle of Lyme disease spirochaetes. *Nat Rev Microbiol* 10:87–99.
doi:10.1038/nrmicro2714

Rego ROM, Bestor A, Rosa PA. 2011. Defining the Plasmid-Borne Restriction-Modification
Systems of the Lyme Disease Spirochete *Borrelia burgdorferi*. *J Bacteriol* 193:1161–1171.
doi:10.1128/jb.01176-10

Rego ROM, Bestor A, Štefka J, Rosa PA. 2014. Population Bottlenecks during the Infectious
Cycle of the Lyme Disease Spirochete *Borrelia burgdorferi*. *Plos One* 9:e101009.
doi:10.1371/journal.pone.0101009

Revel AT, Talaat AM, Norgard MV. 2002. DNA microarray analysis of differential gene
expression in *Borrelia burgdorferi*, the Lyme disease spirochete. *Proc National Acad Sci*
99:1562–1567. doi:10.1073/pnas.032667699

Ring K, Couper LI, Sapiro AL, Yarza F, Yang XF, Clay K, Mateusiak C, Chou S, Swei A. 2022.
Host blood meal identity modifies vector gene expression and competency. *Mol Ecol*
31:2698–2711. doi:10.1111/mec.16413

Rosenberg R, Lindsey NP, Fischer M, Gregory CJ, Hinckley AF, Mead PS, Paz-Bailey G,
Waterman SH, Drexler NA, Kersh GJ, Hooks H, Partridge SK, Visser SN, Beard CB,
Petersen LR. 2018. Vital Signs: Trends in Reported Vectorborne Disease Cases — United
States and Territories, 2004–2016. *Morbidity Mortal Wkly Rep* 67:496–501.
doi:10.15585/mmwr.mm6717e1

Samuels DS, Lybecker MC, Yang XF, Ouyang Z, Bourret TJ, Boyle WK, Stevenson B,
Drecktrah D, Caimano MJ. 2021. Gene Regulation and Transcriptomics. *Curr Issues Mol Biol*
42:223–266. doi:10.21775/cimb.042.223

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods* 9:671–675. doi:10.1038/nmeth.2089

Schwartz I, Margos G, Casjens SR, Qiu W-G, Eggers CH. 2021. Multipartite Genome of Lyme
Disease *Borrelia* : Structure, Variation and Prophages. *Curr Issues Mol Biol* 42:409–454.
doi:10.21775/cimb.042.409

Shaw WR, Catteruccia F. 2018. Vector biology meets disease control: using basic research to
fight vector-borne diseases. *Nat Microbiol* 4:20–34. doi:10.1038/s41564-018-0214-7

Sonenshine DE, Anderson JM. 2014. Mouthparts And Digestive System In: Sonenshine DE,
Roe RM, editors. Biology of Ticks. Oxford University Press. pp. 122–162.

Spielman A, Ribeiro JMC, Mather TN, Piesman J. 1987. Dissemination and Salivary Delivery of
Lyme Disease Spirochetes in Vector Ticks (Acari: Ixodidae). *J Med Entomol* 24:201–205.
doi:10.1093/jmedent/24.2.201

Steere AC, Sikand VK, Meurice F, Parenti DL, Fikrig E, Schoen RT, Nowakowski J, Schmid CH,
Laukamp S, Buscarino C, Krause DS. 1998. Vaccination against Lyme Disease with
Recombinant *Borrelia burgdorferi* Outer-Surface Lipoprotein A with Adjuvant. *New Engl J*
Medicine 339:209–215. doi:10.1056/nejm199807233390401

Steere AC, Strle F, Wormser GP, Hu LT, Branda JA, Hovius JWR, Li X, Mead PS. 2016. Lyme
borreliosis. *Nat Rev Dis Primers* 2:16090. doi:10.1038/nrdp.2016.90

Stevenson B, Zückert WR, Akins DR. 2000. Repetition, conservation, and variation: the multiple
cp32 plasmids of *Borrelia* species. *J Mol Microb Biotech* 2:411–422.

Thomas PD, Ebert D, Muruganujan A, Mushayahama T, Albou L-P, Mi H. 2022. PANTHER:
Making genome-scale phylogenetics accessible to all. *Protein Sci* 31:8–22.
doi:10.1002/pro.4218

Tilly K, Rosa PA, Stewart PE. 2008. Biology of Infection with *Borrelia burgdorferi*. *Infect Dis Clin*
N Am 22:217–234. doi:10.1016/j.idc.2007.12.013

Tokarz R, Anderton JM, Katona LI, Benach JL. 2004. Combined Effects of Blood and
Temperature Shift on *Borrelia burgdorferi* Gene Expression as Determined by Whole
Genome DNA Array. *Infect Immun* 72:5419–5432. doi:10.1128/iai.72.9.5419-5432.2004

Wachter J, Cheff B, Hillman C, Carracoi V, Dorward DW, Martens C, Barbian K, Nardone G,
Olano LR, Kinnersley M, Secor PR, Rosa PA. 2023. Coupled induction of prophage and
virulence factors during tick transmission of the Lyme disease spirochete. *Nat Commun*
14:198. doi:10.1038/s41467-023-35897-3

Zhang H, Marconi RT. 2005. Demonstration of Cotranscription and 1-Methyl-3-Nitroso-
Nitroguanidine Induction of a 30-Gene Operon of *Borrelia burgdorferi*: Evidence that the 32-
Kilobase Circular Plasmids Are Prophages. *J Bacteriol* 187:7985–7995.
doi:10.1128/jb.187.23.7985-7995.2005

Zhang L, Zhang Y, Adusumilli S, Liu L, Narasimhan S, Dai J, Zhao YO, Fikrig E. 2011.
Molecular Interactions that Enable Movement of the Lyme Disease Agent from the Tick Gut
into the Hemolymph. *Plos Pathog* 7:e1002079. doi:10.1371/journal.ppat.1002079

Zhu A, Ibrahim JG, Love MI. 2018. Heavy-tailed prior distributions for sequence count data:
removing the noise and preserving large differences. *Bioinformatics* 35:2084–2092.
doi:10.1093/bioinformatics/bty895

1069