Gac is a transcriptional repressor of the Lyme disease spirochete's

OspC virulence-associated surface protein

Tatiana N. Castro-Padovani [1, §], Timothy C. Saylor[1, §], Olivia T. Husted[1, § ,†],

Andrew C. Krusenstjerna[1, §], and Brian Stevenson [1,2, *]

1. Department of Microbiology, Immunology, and Molecular Genetics, University of

Kentucky College of Medicine, Lexington, Kentucky

2. Department of Entomology, University of Kentucky College of Agriculture, Food,

and Ecology, Lexington, Kentucky

§ T.N. Castro-Padovani, T.C. Saylor, O.T. Husted, and A.C. Krusenstjerna contributed

equally to this work, and are to be considered equal co-first authors

* For correspondence:

Brian Stevenson, MS421 Medical Center, Department of Microbiology, Immunology,

and Molecular Genetics, University of Kentucky College of Medicine, Lexington,

Kentucky, USA, 40536-0298

† Current address for O.T. Husted: Venture Laboratories, Lexington, KY

Abstract

The OspC outer-surface lipoprotein is essential for the Lyme disease spirochete's initial

phase of vertebrate infection. Bacteria within the midguts of unfed ticks do not produce

OspC, but produce high levels when ticks begin to ingest blood. Lyme disease

spirochetes cease production of OspC within 1-2 weeks of vertebrate infection, and

bacteria that fail to downregulate OspC are cleared by host antibodies. Thus, tight

regulation of OspC levels is critical for survival of Lyme borreliae, and therefore an

attractive target for development of novel treatment strategies. Previous studies

determined that a DNA region 5′ of the *ospC* promoter, the *ospC* operator, is required

for control of OspC production, but the mechanism was not determined. Hypothesizing

that the *ospC* operator may bind a regulatory factor, DNA affinity pulldown was

performed, and identified binding by the Gac protein. Gac is encoded by the C-terminal

half of the *gyrA* open reading frame, from an internal promoter, ribosome-binding site,

and initiation codon. Our analyses determined that Gac exhibits a substantially greater

affinity for *ospC* operator and promoter DNAs than for other tested borrelial sequences.

In vitro and in vivo analyses demonstrated that Gac is a transcriptional repressor of

ospC. These results constitute a substantial advance to our understanding the

mechanisms by which the Lyme disease spirochete controls production of OspC.

Importance

Borrelia burgdorferi (sensu lato) requires its surface-exposed OspC protein in order to

establish infection of humans and other vertebrate hosts. Lyme disease spirochetes

initiate production of OspC when the vector tick begins to feed on the blood of a

vertebrate host, and the spirochetes ceases production of OspC soon after establishment

of infection. Bacteria that either do not produce OspC during transmission, or fail to

repress OspC after infection is established, are rapidly cleared by the host. The

molecular mechanisms controlling OspC production are largely unstudied. Herein, we

identified a borrelial protein, Gac, that binds with high affinity to the *ospC* promoter

and 5′ adjacent DNA. A combination of biochemical analyses and investigations of

genetically-manipulated bacteria demonstrated that Gac is a transcriptional repressor of

ospC. This is a substantial advance toward understanding how the Lyme disease

spirochete controls production of the essential OspC virulence factor, and identifies a

novel target for preventative and curative therapies.

Introduction

The Lyme disease spirochete, *Borrelia burgdorferi* sensu lato (*B. burgdorferi* sl),

requires its OspC protein during the initial stages of vertebrate infection, and mutant

bacteria that cannot produce this outer surface-exposed lipoprotein are unable to

initiate infection of vertebrates (1-5). Production of OspC is tightly regulated. *B.*

burgdorferi sl within the midguts of unfed ticks do not express OspC (6, 7). As ticks

begin to feed on host blood, spirochetes initiate production of OspC, with the large

majority producing detectable quantities of OspC at the time of transfer from tick to

vertebrate (6, 8-10). The function(s) of OspC is not completely understood - it can bind a

tick salivary protein and vertebrate plasminogen and complement C4b - but may have

additional functions (11-21). Infected humans and other animals mount strong antibody

responses against OspC within days of infection (22-25). At the same time, *B. burgdorferi*

sl represses expression of OspC, and spirochetes that do not shut off OspC synthesis are

cleared by immunocompetent hosts (2, 3, 5, 26-30). Thus, regulation of OspC may be an

excellent target for novel therapeutics to prevent and/or clear *B. burgdorferi* infections.

OspC was the first *B. burgdorferi* sl protein to be identified as being regulated

during the natural tick-vertebrate infectious cycle (6). As noted above, OspC is not

produced in unfed ticks, but is highly expressed in feeding ticks and during the earliest

stages of vertebrate infection (6, 8). Regulated production can also be manipulated in

culture, with bacteria grown at 23°C, pH=7.5, or culture media that yield slow

replication rates, producing low to no OspC, while the protein is highly expressed

following transfer from those conditions to 35°C, pH=6.5, or complete culture media,

respectively (6, 31-33). It is not yet known how those culture conditions are interpreted

by *B. burgdorferi* sl to modulate OspC levels, but they provide convenient means to

evaluate borrelial regulatory mechanisms.

An alternative sigma factor, named RpoS, is also affected by the above-described

infection and culture conditions (33-35). RpoS controls production of numerous *B.*

burgdorferi sl proteins that are required for vertebrate infection (36, 37). Deletion of *rpoS*

blocks induction of OspC, which has led to a hypothesis that expression of *ospC* is solely

due to RpoS-RNA polymerase holoenzyme directly transcribing from the *ospC*

promoter (34, 38). Yet, there is evidence that transcriptional control of *ospC* is more

complicated than that model assumes. Previous studies found that a region of DNA 5'

of the *ospC* transcriptional start site, designated the *ospC* operator, is necessary for

induction of *ospC* transcription (39-41). The mode of function of the *ospC* operator had

not been defined prior to the current investigations.

B. burgdorferi sl has been divided into several clades that are often referred to as

genospecies, which include *B. burgdorferi* sensu stricto (*B. burgdorferi* ss) and *B. garinii*.

An earlier study noted that the *ospC* operator of *B. burgdorferi* ss type strain B31 includes

a pair of inverted repeat sequences (Fig. 1) (41). Yet, those inverted repeats are found in

only a subset of Lyme disease borreliae (42). However, the *ospC* operators of all

examined Lyme spirochetes contain two or more copies of a conserved 11bp directly-

repeating sequence (Fig. 1 and (42)). *B. burgdorferi* ss B31 contains two copies of the 11bp

repeat, which overlap the previously discussed inverted repeats (Fig. 1 and (41, 42)). In

contrast, the *ospC* operators of some strains of the *B. garinii* genospecies contain seven

adjacent copies of the 11bp repeat (Fig. 1). All *ospC* promoters also contain a copy of the

11bp repeat, oriented in the opposite direction (Fig. 1). The retention of the 11bp repeat

suggested to us that it may be a key component of *ospC* regulation, with sequence

conservation pointing toward it being a binding site for a regulatory protein. To address

that hypothesis, we performed DNA-binding studies with borrelial cytoplasmic extract,

identified a component protein that has high affinity for the *ospC* operator, and

identified a functional role for that protein as a transcriptional repressor of *ospC*.

Results

Identification of Gac as an *ospC* operator-binding protein. As a first step, an

infectious clone of *B. burgdorferi* ss type strain B31, B31-MI-16, was cultured to mid-

exponential phase at 35˚C. That condition was adopted based on our previous

characterization of the *B. burgdorferi erp* operons, whose expression is also responsive to

culture temperature, in which we found that the *erp* repressing and activating proteins

were all adequately expressed when cultured at 35˚C (43-45).

A fluorescently-tagged DNA sequence corresponding to the *ospC* operator region

of *B. garinii* Ip89 was incubated with the B31-MI-16 cytoplasmic extract, then subjected

to electrophoretic mobility shift electrophoresis (EMSA). The strain Ip89 sequence was

used for these analyses under the hypothesis that a protein that binds to the 11bp

repeated sequence would have an overall greater affinity for the 7 repeats of Ip89 as

opposed to the 2 repeats of B31 (Fig. 1). Addition of the cytoplasmic extract resulted in

retarded mobility of the labeled DNA probe, indicating that a borrelial component

bound to the DNA (Fig. 2A).

Ip89 *ospC* operator DNA was again PCR amplified, but with one of the

oligonucleotide primers containing a 5′ biotin moiety. That DNA was adhered to

streptavidin-coated magnetic beads, incubated with B31 cytoplasmic extract, washed

extensively, then bound proteins eluted with buffers having progressively greater

concentrations of NaCl. Aliquots of each eluted fraction were subjected to SDS-PAGE

and proteins stained with SYPRO-Ruby. A protein band with an approximate molecular

mass of 30-35 kDa eluted at 0.3 and 0.5 M NaCl (Fig. 2B). The bands were excised from

the gel, subjected to trypsin digestion and mass spectrometry, and results compared to

the predicted proteome of B31-MI-16. Only one potential match was obtained, to the

GyrA subunit of DNA gyrase, with 9 unique peptides identified. Notably, all of the

identified peptide fragments mapped to the C-terminal half of GyrA (Fig. 2C).

Bacterial DNA gyrases are comprised of two subunits, GyrA and GyrB (46).

GyrA proteins possess a unique C-terminal domain that bends DNA to facilitate the

topoisomerase activity of the gyrase (47, 48). Curiously, Lyme disease borreliae produce

two separate proteins from the *gyrA* locus, a full length GyrA protein and a second, 35

kDa protein named Gac, which is transcribed from an internal promoter and consists

solely of the C-terminal DNA-binding domain (48-50) (Fig. 2D, and Fig. S1). The

function(s) of Gac was previously unknown. We pursued the possibility that Gac

preferentially binds to *ospC* operator DNA and thereby affects production of OspC.

Gac exhibits high-affinity binding to the *ospC* operator and promoter. The Gac

protein was first detected by methods similar to ours, in a DNA-affinity pull-down

study using a *B. burgdorferi* plasmid telomere sequence (49). That study observed that

recombinant Gac could also bind to other DNA sequences, and so it was hypothesized

that binding of Gac lacked specificity (49). We note, however, that the previous report

did not state the concentrations of recombinant Gac or DNA probes used in their

EMSAs, nor were results shown. The brief description of methods in that publication do

not mention use of unlabeled competitor DNAs, which can also be critical to

determining binding specificity. We therefore conducted comparative analyses to assess

the relative affinity of Gac for *ospC* operator sequences and other DNAs.

To address that question, labeled DNA probes were constructed that consisted of

either the strain Ip89 or B31 *ospC* operators (Fig. 1). Additional unlabeled DNAs were

produced for use as EMSA competitors: each of the *ospC* operators, truncated segments

of the B31 *ospC* operator, and DNA sequences corresponding to the 5' ends of the

borrelial *flaB* and *ospAB* operons. The Ip89 *ospC* operator probe and competitor were

derived by PCR from plasmid pCR2.1 clones, using plasmid-specific M13 Reverse and

Forward primers, so an additional unlabeled competitor was produced by PCR of

empty pCR2.1, to account for vector-derived DNA sequences.

EMSAs of the Ip89 *ospC* operator probe and recombinant Gac found that the

protein bound to the labeled probe (Fig 3A, lane 2). Addition of 10-times excess of

unlabeled Ip89 *ospC* operator partly depleted the DNA-Gac EMSA band, while 25-fold

excess of unlabeled Ip89 *ospC* operator completely eliminated the complex (Fig. 3A,

lanes 3 and 4, respectively). Addition of 25-times excess unlabeled B31 *ospC* operator

had little effect on the Ip89 operator-Gac complex, indicating a greater affinity of Gac

for the Ip89 operator than for the B31 operator (Fig 3A, lane 5). Note that the primary

difference between the Ip89 and B31 DNAs is the number of 11bp repeat sequences in

their *ospC* operators. Inclusion of 25-fold excesses of either empty pCR2.1 amplicon,

flaB, or *ospAB* unlabeled DNAs did not compete Gac away from the labeled probe (Fig

3A, lanes 6, 7, and 8, respectively). Thus, Gac displays substantially greater affinity for

the Ip89 *ospC* operator than it does for any of the other tested sequences.

Similar EMSAs were performed with labeled B31 *ospC* operator as probe.

Unlabeled B31 *ospC* operator DNA effectively competed Gac away from the labeled

probe (Fig 3B, lanes 3-4). Inclusion of 10-fold excesses of the empty pCR2.1 amplicon,

flaB, or *ospAB* unlabeled DNAs exhibited lower levels of competition for Gac binding

(Fig 3B, lanes 5, 6, and 7, respectively). However, 25-fold excess of all of those unlabeled

DNA competed Gac away from the labeled B31 *ospC* labeled probe (data not shown).

These results indicate that Gac binds to the B31 *ospC* operator with a greater affinity

than to the *flaB, ospAB,* or pCR2.1 competitor sequences, but has a lower affinity for the

B31 *ospC* operator than for the Ip89 *ospC* operator.

Using labeled *ospC* operator from strain B31 as the EMSA probe, truncated

segments of that DNA region were used as competitors to narrow down the sequences

that bound Gac (Fig 4). As above, 10- and 25-fold excesses of unlabeled B31 *ospC*

operator DNA competed Gac away from the labeled probe. Since the labeled probe

included vector sequences at each end, competitor A was designed to contain only *ospC*

operator sequences; it also effectively competed for Gac-binding at 10-fold excess. Of

the other competitors, D yielded the greatest level of competition, while B, C, F, and G

competed to lesser extents. These results indicate that Gac also has an elevated affinity

for the *ospC* promoter region. We note that the space between the *ospC* -10 and -35

sequences includes an 11bp sequence that resembles the operator repeat (Figure 1).

Altered *ospC* expression in *gac* mutant *B. burgdorferi*. To assess the effects of

Gac in live *B. burgdorferi*, *gac* mutant strain CKO-1, strain CKO-1 (pBLS820) (CKO-1

complemented with wild-type *gac*), and strain NGR (an isogenic wild-type *gac* clone)

were cultured to mid-exponential phase at 23 ˚C, then diluted into fresh media and

incubated to mid-exponential phase at 35 ˚C (6, 31). Immunoblot analyses of lysates of

the 23 and 35˚C cultures using OspC-specific antibodies found that all three strains

exhibited differential expression of OspC protein, with substantially greater amounts of

OspC being synthesized at 35 ˚C (Fig. 5). However, Gac-deficient strain CKO-1

produced 14- and 1.4-times more OspC than did NGR at 23 and 34˚C, respectively. The

greatest net impact was observed with the 23˚C cultures, consistent with Gac being a

repressor of OspC synthesis. Complementation of CKO-1 with a wild-type *gac* gene on

pBLS820 reduced OspC levels by 32% and 20% at 23 and 34˚C, respectively. The

complemented strain did not fully return to wild-type OspC levels, suggesting that the

minimal *gac* promoter in pBLS820 does not contain all of the cis elements necessary for

normal control of *gac* expression.

B. burgdorferi strains NGR and CKO-1 were next transformed with plasmids that

carry transcriptional fusions between B31 *ospC* operator/promoter sequences and *gfp*.

Plasmid pBLS761 carries the same operator/promoter sequence that was used as the

EMSA probe in the above-described studies, while pBLS764 lacks the 11 bp direct

repeats (equivalent to the sequences of competitors A and G in Fig. 4, respectively).

Each transformant was cultured at 35˚C to mid-exponential phase, then bacterial GFP

content was assessed by flow cytometry (Fig. 6). CKO-1(pBLS761) produced

substantially greater levels of GFP than did NGR(pBLS761), consistent with the

observed levels of OspC protein and transcript described above.

NGR(pBLS764), which lacks the *ospC* operator, did not produce GFP (Fig. 6). In

contrast, CKO-1(pBLS764) produced GFP at levels comparable to CKO-1(pBLS761). We

conclude that absence of Gac relieves transcriptional repression of the *ospC* promoter,

while the *ospC* operator is also required for efficient transcription from the *ospC*

promoter if Gac is present. Those results suggest that one or more additional factors

interact with *ospC* operator DNA to inhibit Gac binding to the *ospC* promoter. Studies

are ongoing to examine that hypothesis.

***B. burgdorferi* regulates transcription of *gac*.** Since *ospC* is induced in *B.*

burgdorferi cultured at 35°C, we assessed *ospC* and *gyrA/gac* transcript levels over time in

cultures of wild-type *B. burgdorferi* (Fig. 7). Levels of *ospC* transcript increased as the

cultures grew through exponential into stationary phase. The two *gyrA/gac* primer sets

yielded different Q-RT-PCR results for each pair. Since the 5′ primers detected mRNA

that only encoded *gyrA*, whereas the 3′ primers detected both *gyrA* and *gac* mRNAs, the

relative concentration of *gac* mRNA was approximated by subtracting the results of 5′

Q-RT-PCR from results from the 3′ primer set. The concentration of Gac-encoding

transcript was greatest during lag and early exponential phases, then declined as the

cultures matured and entered stationary phase. Those changes were opposite of *ospC*,

and are consistent with the above-described results that indicate Gac is a repressor of

ospC.

Gac exhibits affinity for the telomere of *B. burgdorferi* B31 plasmid lp17.

Borrelia species contain both circular and linear DNAs, and there is considerable interest

in the mechanisms by which the spirochetes maintain linear replicons (51-53). The Gac

protein was initially identified through an affinity pull-down method similar to ours,

using a bait sequence derived from a telomere of strain B31's linear plasmid lp17 (49).

Having found that Gac binds to *ospC* 5′ DNAs with high degrees of affinity, we

revisited the previous study. A labeled DNA probe with the same telomere-derived

sequence was produced, then assessed by EMSA for ability to bind Gac in the presence

of excess levels of various unlabeled DNAs. Recombinant Gac produced an EMSA shift

with the labeled telomere probe, which was effectively competed away with 10-fold and

25-fold excesses of unlabeled telomere sequence (Fig. 8, lanes 2-4). Ten-fold excesses of

unlabeled Ip89 or B31 *ospC* operator DNAs also competed Gac away from the labeled

telomere DNA (Fig. 8, lanes 6 and 7, respectively). Unlabeled empty pCR2.1 amplicon

partly competed for Gac binding, while neither the *flaB* nor the *ospAB* DNAs detectably

competed (Fig. 8, lanes 5, 8, and 9, respectively). We conclude that Gac actually does

exhibit specific affinity for the lp17 telomere region, although its relative affinity is

lower than for the Ip89 and B31 *ospC* operators, yet higher than for the examined *ospAB*

or *flaB* DNA sequences.

Discussion

These studies found that Gac binds with high affinity to *ospC* promoter and

operator DNA, and inhibits transcription. Deletion of the *ospC* operator did not alleviate

Gac-dependent repression of transcription from the *ospC* promoter, suggesting that Gac

binding to the promoter itself is sufficient to repress transcription. The *ospC* operator is

evidently necessary for relief of Gac-dependent transcriptional repression, mirroring

results from previous studies that also found that the operator is essential for regulation

of *ospC* transcription (39-41). Thus, it appears that a "factor(s) X" interacts with the *ospC*

operator and somehow prevents repression by Gac (Fig. 9). We are currently pursuing

biochemical and genetic routes to identify the hypothesized factor X.

Gac was initially identified because it bound to a telomere of linear plasmid lp17

(49), although it was not studied further due to perceived lack of binding specificity.

The present analyses indicate that Gac actually has a marked preference for the lp17

telomere. This raises questions about the function(s) of Gac in *B. burgdorferi*. Our studies

found that Gac has elevated affinity for three sites in the *B. burgdorferi* genome – the

ospC operator, the *ospC* promoter, and the telomere of lp17 – yet can also possesses

lesser affinity for other DNA sequences. Gac wraps DNA around itself, forming 180-

220˚ bends (48). Taken together, these data suggest that Gac may be a nucleoid-

associated protein (NAP, or "histone-like protein"). Supporting that model, Gac

complements some features of the *E. coli* NAP HU when expressed in that species (49,

50). It is not uncommon for NAPs to also influence transcription (54-56). Indeed, the

antirepressor of *B. burgdorferi erp* operon transcription, EbfC, meets the criteria of being

an NAP (57).

At this time, it is not known whether production of a separate Gac protein from

the 3′ end of the *gyrA* ORF is unique to *B. burgdorferi*. The possibility that *gyrA* can

encode two distinct proteins has never been addressed for the vast majority of other

bacterial species. It would be worth examining other *gyrA* genes for presence of

potential ribosome binding sites and start codons near the beginning of the C-terminal

domain.

In conclusion, our investigations found that *B. burgdorferi* Gac has elevated

affinity for *ospC* operator and promoter DNAs. Gac repressed *ospC* transcription in vitro

and in live spirochetes. Levels of *ospC* and *gac* transcripts changed in opposite directions

during cultivation, consistent with *ospC* repression by Gac. Analyses of mutants

indicate that the *ospC* operator is required to alleviate Gac repression of the *ospC*

promoter, implying that one or more additional factors interact with the *ospC* operator

to prevent repression by Gac. That protein also bound with elevated affinity to an lp17

telomere, suggesting that Gac might play a role in linear DNA maintenance.

Materials and Methods

Bacteria, plasmids, and culture conditions. *B. burgdorferi* sensu stricto strain B31

is the type strain, and was isolated from an *Ixodes scapularis* tick that was collected on

Shelter Island, New York (58, 59). Infectious clone B31-MI-16 was used to prepare

cytoplasmic extracts for identification of *ospC* operator-binding proteins (60). *B. garinii*

strain Ip89 was isolated from an *I. persulcatus* tick that was collected in central Eurasia

(61, 62).

Borrelia spp. strains were cultured in BSK-II broth at either 23 or 35˚C (63).

Temperature shift cultivation was performed by diluting a 23˚C culture 1:100 into fresh

medium, then incubating at 35˚C (31). Culture densities were determined with a Petroff-

Hausser counting chamber and dark field microscopy.

Analyses of the effects of Gac on OspC expression used a Δ*gac* clone of strain

B31, CKO-1, and an isogenic control strain, NGR (50). The overlap of the *gac* ORF with

the essential *gyrA* ORF precludes outright deletion of the *gac* locus. For that reason,

Knight et al. (50) produced *gac* defective strain CKO-1 by an allelic exchange mutation

that changed the two potential *gac* start codons from ATG to CTT and ATT (leucine and

isoleucine, respectively) and the ribosome binding site from AGGAGG to AGAAGA.

The mutant does not produce a Gac protein, although it does produce normal levels of

GyrA (50). The *gac* mutations were selected by simultaneous introduction of point

mutations into the adjacent *gyrB* gene, which confer resistance to coumermycin (50, 64).

The CKO-1 *gac* mutant exhibits growth patterns that are physiologically

indistinguishable from the wild-type parent or from NGR, an isogenic coumermycin-

resistant *gyrB* strain (50).

Plasmid pBLS820 was custom produced by GenScript (Piscataway, NJ) by

cloning the *B. burgdorferi gac* open reading frame, ribosome-binding site, and promoter

into vector pBSV2-G (65) (Fig. S1). *B. burgdorferi* CKO-1 was transformed by

electroporation, and transformants were selected by culture in 150 µg/ml gentamicin.

Plasmid pBLS761 contains the B31 *ospC* promoter and operator in a

transcriptional fusion with *gfp*. Plasmid pBLS764 is identical to pBLS761 except that it

lacks the *ospC* operator. Both are based on pBLS740, which was produced for this work

by cloning the *gfp* gene of pBLS590 between the BamHI and HindIII sites of vector

pKFSS1 (66, 67). The *ospC*-derived DNAs used to make pBLS761 and pBLS764 were

inserted between the KpnI and BamHI sites of pBLS740, custom synthesized by

GenScript. Two of our previously-described plasmids were used as GFP expression

controls: KS10, which carries a promoterless *gfp* on pBLS590, and KS20, which carries

the constitutively-expressed minimal *erpAB* promoter driving *gfp* on pBLS599 (66).

Plasmids pBLS590 and pBLS599 confer resistance to kanamycin. *B. burgdorferi*

containing pBLS590 or pBLS599 were cultured in media that contained 200 µg/ml

kanamycin, while spirochetes that contain pBLS761 or pBLS764 were cultured in media

with 50 µg/ml streptomycin (66-69).

Cytoplasmic extracts. *B. burgdorferi* ss B31-MI-16 was cultured at 35˚C to mid-

exponential phase (approximately 5×10^7 bacteria/ml). Bacteria were harvested by

centrifugation, washed with phosphate buffered saline (PBS), and resuspended in a

minimal volume of 50mM Tris (pH=7.6), 10mM EDTA, 10% (w/vol) sucrose. The

suspension was frozen in a dry ice/ ethanol bath, then allowed to thaw in an ice-water

bath. NaCl, dithiothreitol, and lysozyme were added to final concentrations of 140 mM,

1 mM, and 0.4 mg/ml, respectively, and incubated on ice for 5 min. The cell mixture

was subjected to four additional freeze-thaw cycles. Cellular debris was pelleted by

centrifugation, and the supernatant was aliquoted and stored at -80°C (70).

Electrophoretic mobility shift assays (EMSAs). Labeled and unlabeled nucleic

acids are described in Supplementary Table 1. Fluorescently tagged IRDye800 labeled

probes were utilized for EMSAs. For each EMSA binding reaction, a labeled probe was

incubated for 15 min at room temperature with protein and unlabeled competitor DNA

as necessary, in EMSA buffer (50mM Tris-HCl, 25mMKCl, 10% glycerol (vol/vol),

0.01% Tween 20, 100nM dithiothreitol, and 1mM phenylmethanesulfonyl fluoride

(PMSF). Protein-nucleic acid mixtures were then loaded into either 6% or 10% Novex

TBE gels (Thermo Fisher). Gels were pre-run for at least 30 min at 100V in 0.5x TBE

buffer before sample loading. Following the addition of the samples, electrophoresis

was conducted for up to 1.5 h at 100V.

Initially, a fragment of *B. garinii* Ip89 DNA, from 5' of the *ospC* operator through

the *ospC* promoter, was amplified from genomic DNA by PCR with oligonucleotide

primers 1021 and 1022 (Supplementary Table 1). The resulting amplicon was cloned

into pCR2.1 (Invitrogen) to create pBLS756. Fluorescently-tagged IP89 *ospC* operator

was produced by PCR of pBLS756 using IRDye800-tagged M13-Reverse and unlabeled

M13-Forward primers (Supplementary Table 1). The probe was incubated with cleared

borrelial cytoplasmic extract in EMSA buffer for 15 min at room temperature with 10ng

poly-dI-dC, then subjected to electrophoresis.

Other EMSAs used purified recombinant Gac protein, labeled *ospC*

operator/promoter probes, and unlabeled DNA competitors.

In addition, a sub-telomeric sequence of strain B31 lp17 was amplified from total

genomic DNA by PCR with oligonucleotides TL16h and TL16g as previously described

(49). The resulting PCR product was cloned into pCR2.1 (Invitrogen) and maintained in

E. coli TOP-10. The amplicon was completely sequenced on both strands to ensure

fidelity. Fluorescently tagged telomere probe was produced by PCR using IRDye800

tagged M13-Reverse and unlabeled M13-Forward primers.

DNA affinity pull-down. Proteins that bind *ospC* operator DNA were purified

from B31-MI-16 cytoplasmic extracts as previously described (71). Briefly, bacteria of a

mid-exponential phase (approximately 5×10^7 bacteria/ml) culture were harvested,

washed, and frozen at -80˚C. Bacterial pellets were then thawed on ice, and subjected to

three additional rounds of freezing at -80˚C / thawing on ice. Bacterial lysates were

resuspended in a minimal volume of BS/THES buffer (44.3% THES buffer, 20% BS

buffer, 35.7% water). THES Buffer consists of 50mM Tri HCl (pH 7.5), 10mM EDTA,

20% sucrose (w/vol), 140mM NaCl, 0.7% (vol/vol) Protease Inhibitor Cocktail II

(Sigma), and 0.1% (vol/vol) Phosphatase Inhibitor Cocktail II (Sigma). BS buffer

consists of 50mM HEPES (pH=7.5), 25mM $CaCl_2$, 250mM KCl, and 60% (vol/vol)

glycerol. Suspended bacteria were lysed by sonication, on ice. Cellular debris was

pelleted by centrifugation, and the supernatant was divided into aliquots.

Recombinant Gac protein. The B31 *gac* ORF was cloned into pET101, then

sequenced to confirm fidelity. C-terminally His-tagged Gac was expressed in *E. coli*

Rosetta II upon induction with 1 mM of isopropyl-β-d-thiogalactopyranoside (IPTG) for

1 hour at 37°C. Bacteria were collected by centrifugation, and the pellet was frozen at -

80°C. The bacterial pellet was then thawed on ice, resuspended in phosphate buffered

saline (PBS) containing 10 mM imidazole, and bacteria lysed by sonication. The sonicate

was subjected to centrifugation, then recombinant protein purified from the soluble

fraction using a HisPur Ni-NTA (Thermo Scientific) column. Purified protein was

dialyzed against EMSA buffer. Protein concentration was determined by Bradford assay

and purity assessed by SDS-PAGE with Coomassie brilliant blue staining.

Immunoblot analyses. Cultures of strains NGR, CKO-1, and CKO-1(pBLS820)

were harvested at mid-exponential phase by centrifugation, then split into aliquots for

immunoblot assays and RNA extraction. Pellets for immunoblot assays were frozen at -

80°C, subsequently thawed at room temperature, then resuspended in 10% SDS, boiled

for 5-minutes, and vigorously vortexed. The resulting lysates were stored at 4°C. Prior

to electrophoresis, samples were preheated to 42° before adding to 2x sodium dodecyl

sulfate (SDS)-page loading dye (63mM Tris pH7.5, 18.75% w/v glycerol, 10% w/v 2-

mercaptoethanol, 4% SDS, and .5% w/v bromophenol blue). Separation by SDS-

polyacrylamide gel electrophoresis was followed by electrotransfer to nitrocellulose

membranes and blocking with 5% nonfat dried milk in Tris-buffered saline-Tween-20

(20 mM Tris [pH 7.5], 150 mM NaCl, 0.05% [vol/vol] Tween-20). The membranes were

then incubated with murine monoclonal antibodies directed against FlaB or OspC (72,

73). After incubation with primary antibodies, membranes were incubated with goat

anti-mouse IgG conjugated to IRDye800 (LI-Cor). Immunoblots were imaged and

analyzed with ChemiDoc MP and Image-Lab software, respectively (Bio-Rad, CA).

415   **Quantitative reverse transcription PCR (Q-RT-PCR).** *B. burgdorferi* was

cultured at 35 ºC, and 2 ml aliquots of bacteria were taken 1-, 3-, 5-, 7-, 9-, and 11-days

post-inoculation. Bacteria were pelleted, then washed two times with PBS and frozen at

-80 ºC. Frozen pellets were later resuspended in 60 ºC TRIzol (Thermo Fisher), and RNA

was extracted using Qiagen Mini RNA kits. Residual genomic DNA was depleted using

on-column Turbo DNase (Thermo Fisher). The quality (RIN > 7) and quantity of the

RNA were determined using an Agilent 2100 Bioanalyzer and Agilent 6000 Nano chips.

Purified RNA stocks were stored at -80 ºC.

423   Complementary DNA (cDNA) for qPCR was produced using the iScript gDNA

Clear cDNA synthesis kits (Bio-Rad). The cDNA stocks were diluted 1:100 into

nuclease-free water for use as template. Quantitative PCR was carried out using iTaq

Universal SYBR Green Supermix (Bio-Rad) and a Bio-Rad CFX96 Touch Real-time PCR

thermocycler. For all reactions, cycles involved melting at 95 ºC for 10s, annealing at 55

ºC for 10 s, and extension at 72 ºC for 30s. Each run was performed with technical

triplicates of each reaction as well as no-template control reactions that lacked template.

Melting curves were performed with each run to validate production of specific

amplicons. Every reaction was tested with a reverse transcriptase-negative template to

validate the absence of contaminating gDNA. Cq values were normalized to *ftsK*, a gene

whose expression is stable across different growth phases (74), using the ΔCq method,

and fold-difference was determined by the function $2^{-\Delta Cq}$.

Acknowledgements

This work was funded by NIH grant 3R01AI144126-03S1. We thank Nerina Jusufovic

and Jessamyn Moore for constructive comments on this project and manuscript; Scott

Samuels for providing *B. burgdorferi* strains NGR and CKO-1, and for helpful comments

on these studies; Haining Zhu and Jing Chen for assistance with mass spectrometric

analyses; Jennifer Strange, Greg Bowman, and Cuiping Zhang for flow cytometry

analyses. Figure 9 was produced using BioRender.com.

References

1.   Grimm D, Tilly K, Byram R, Stewart PE, Krum JG, Bueschel DM, Schwan TG,

Policastro PF, Elias AF, Rosa PA. 2004. Outer-surface protein C of the Lyme

disease spirochete: a protein induced in ticks for infection of mammals. Proc Natl

Acad Sci USA 101:3142-3147.

2.   Stewart PE, Wang X, Bueschel DM, Clifton DR, Grimm D, Tilly K, Carroll JA,

Weis JJ, Rosa PA. 2006. Delineating the requirement for the *Borrelia burgdorferi*

virulence factor OspC in the mammalian host. Infect Immun 74:3547-3553.

3.   Tilly K, Krum JG, Bestor A, Jewett MW, Grimm D, Bueschel D, Byram R,

Dorward D, Vanraden MJ, Stewart P, Rosa P. 2006. *Borrelia burgdorferi* OspC

protein is required exclusively in a crucial early stage of mammalian infection.

Infect Immun 74:3554-3564.

4.   Tilly K, Bestor A, Jewett MW, Rosa P. 2007. Rapid clearance of Lyme disease

spirochetes lacking OspC from skin. Infect Immun 75:1517-1519.

5.   Tilly K, Bestor A, Rosa PA. 2013. Lipoprotein succession in *Borrelia burgdorferi*:

similar but distinct roles for OspC and VlsE at different stages of mammalian

infection. Mol Microbiol 89:216-227.

6.   Schwan TG, Piesman J, Golde WT, Dolan MC, Rosa PA. 1995. Induction of an

outer surface protein on *Borrelia burgdorferi* during tick feeding. Proc Natl Acad

Sci USA 92:2909-2913.

7.   Fingerle V, Hauser U, Liegl G, Petko B, Preac-Mursic V, Wilske B. 1995.

Expression of outer surface proteins A and C of *Borrelia burgdorferi* in *Ixodes*

ricinus. J Clin Microbiol 33:1867-1869.

8.   Schwan TG, Piesman J. 2000. Temporal changes in outer surface proteins A and

C of the Lyme disease-associated spirochete, *Borrelia burgdorferi*, during the chain

of infection in ticks and mice. J Clin Microbiol 38:382-388.

9.   Pal U, Yang X, Chen M, Bockenstedt LK, Anderson JF, Flavell RA, Norgard MV,

Fikrig E. 2004. OspC facilitates *Borrelia burgdorferi* invasion of *Ixodes scapularis*

salivary glands. J Clin Invest 113:220-230.

10. Skare JT, Shaw DK, Trzeciakowski JP, Hyde JA. 2016. In vivo imaging

demonstrates that *Borrelia burgdorferi ospC* is uniquely expressed temporally and

spatially throughout experimental infection. PLoS ONE 11:e0162501.

11. Lagal V, Portnoï D, Faure G, Postic D, Baranton G. 2006. *Borrelia burgdorferi* sensu

stricto invasiveness is correlated with OspC-plasminogen affinity. Microbes

Infect 8:645-652.

12.   Önder Ö, Humphrey PT, McOmber B, Korobova F, Francella N, Greenbaum DC,

482   Brisson D. 2012. OspC is potent plasminogen receptor on surface of *Borrelia*

483   *burgdorferi*. J Biol Chem 287:16860-16868.

484   13.   Ramamoorthi N, Narasimhan S, Pal U, Bao F, Yang XF, Fish D, Anguita J,

485   Norgard MV, Kantor FS, Anderson JF, Koski RA, Fikrig E. 2005. The Lyme

486   disease agent exploits a tick protein to infect the mammalian host. Nature

487   436:573-577.

488   14.   Xu Q, McShan K, Liang FT. 2008. Essential protective role attributed to the

489   surface lipoproteins of *Borrelia burgdorferi* against innate defences. Mol Microbiol

490   69:15-29.

491   15.   Earnhart CG, Leblanc DV, Alix KE, Desrosiers DC, Radolf JD, Marconi RT. 2010.

492   Identification of residues within ligand-binding domain 1 (LBD1) of the *Borrelia*

493   *burgdorferi* OspC protein required for function in the mammalian environment.

494   Mol Microbiol 76:393-408.

495   16.   Antonara S, Ristow P, McCarthy J, Coburn J. 2010. Effect of *Borrelia burgdorferi*

496   OspC at the site of inoculation in mouse skin. Infect Immun 78:4723-4733.

497   17.   Caine JA, Coburn J. 2015. A short-term *Borrelia burgdorferi* infection model

498   identifies tissue tropisms and bloodstream survival conferred by adhesion

499   proteins. Infect Immun 83:3184-3194.

500   18.   Carrasco SE, Troxell B, Yang Y, Brandt SL, Li H, Sandusky GE, Condon KW,

501   Serezani CH, Yang XF. 2015. Outer surface protein OspC is an antiphagocytic

502   factor that protects *Borrelia burgdorferi* from phagocytosis by macrophages. Infect

503   Immun 83:4848-4860.

504   19.   Caine JA, Lin YP, Kessler JR, Sato H, Leong JM, Coburn J. 2017. *Borrelia*

505   *burgdorferi* outer surface protein C (OspC) binds complement component C4b

506   and confers bloodstream survival. Cell Microbiol 19:10.1111/cmi.12786.

507   20.   Bierwagen P, Szpotkowski K, Jaskolski M, Urbanowicz A. 2019. *Borrelia* outer

508   surface protein C is capable of human fibrinogen binding. FEBS J 286:2415-2428.

509   21.   Lin YP, Tan X, Caine JA, Castellanos M, Chaconas G, Coburn J, Leong JM. 2020.

510   Strain-specific joint invasion and colonization by Lyme disease spirochetes is

511   promoted by outer surface protein C. PLoS Pathog 16:e1008516.

512   22.   Fuchs R, Jauris S, Lottspeich F, Preac-Mursic V, Wilske B, Soutschek E. 1992.

513   Molecular analysis and expression of a *Borrelia burgdorferi* gene encoding a 22kDa

514   protein (pC) in *Escherichia coli*. Mol Microbiol 6:503-509.

515   23.   Fung BP, McHugh GL, Leong JM, Steere AC. 1994. Humoral immune response to

516   outer surface protein C of *Borrelia burgdorferi* in Lyme disease: role of the

517   immunoglobulin M response in the serodiagnosis of early infection. Infect

518   Immun 62:3213-3221.

24.　Padula SJ, Dias F, Sampieri A, Craven RB, Ryan RW. 1994. Use of recombinant

OspC from *Borrelia burgdorferi* for serodiagnosis of early Lyme disease. J Clin

Microbiol 32:1733-1738.

25.　Gilmore RD, Kappel KJ, Dolan MC, Burkot TR, Johnson BJB. 1996. Outer surface

protein C (OspC), but not P39, is a protective immunogen against a tick-

transmitted *Borrelia burgdorferi* challenge: evidence for a conformational

protective epitope in OspC. Infect Immun 64:2234-2239.

26.　Liang FT, Jacobs MB, Bowers LC, Philipp MT. 2002. An immune evasion

mechanism for spirochetal persistence in Lyme borreliosis. J Exp Med 195:415-

422.

27.　Crother TR, Champion CI, Whitelegge JP, Aguilera R, Wu XY, Blanco DR, Miller

JN, Lovett MA. 2004. Temporal analysis of the antigenic composition of *Borrelia*

burgdorferi during infection in rabbit skin. Infect Immun 72:5063-5072.

28.　Liang FT, Yan J, Mbow ML, Sviat SL, Gilmore RD, Mamula M, Fikrig E. 2004.

Borrelia burgdorferi changes its surface antigenic expression in response to host

immune reponses. Infect Immun 72:5759-5767.

29.　Xu Q, Seemanapalli SV, McShan K, Liang FT. 2006. Constitutive expression of

outer surface protein C diminishes the ability of *Borrelia burgdorferi* to evade

specific humoral immunity. Infect Immun 74:5177-5184.

30.   Tilly K, Bestor A, Dulebohn DP, Rosa PA. 2009. OspC-independent infection and
539   dissemination by host-adapted *Borrelia burgdorferi*. Infect Immun 77:2672-2682.

540   31.   Stevenson B, Schwan TG, Rosa PA. 1995. Temperature-related differential
541   expression of antigens in the Lyme disease spirochete, *Borrelia burgdorferi*. Infect
542   Immun 63:4535-4539.

543   32.   Carroll JA, Garon CF, Schwan TG. 1999. Effects of environmental pH on
544   membrane proteins in *Borrelia burgdorferi*. Infect Immun 67:3181-3187.

545   33.   Jutras BL, Chenail AM, Stevenson B. 2013. Changes in bacterial growth rate
546   govern expression of the *Borrelia burgdorferi* OspC and Erp infection-associated
547   surface proteins. J Bacteriol 195:757-764.

548   34.   Hübner A, Yang X, Nolen DM, Popova TG, Cabello PC, Norgard MV. 2001.
549   Expression of *Borrelia burgdorferi* OspC and DbpA is controlled by a RpoN-RpoS
550   regulatory pathway. Proc Natl Acad Sci USA 98:12724-12729.

551   35.   Caimano MJ, Groshong AM, Belperron A, Mao J, Hawley KL, Luthra A, Graham
552   DE, Earnhart CG, Marconi RT, Bockenstedt LK, Blevins JS, Radolf JD. 2019. The
553   RpoS gatekeeper in *Borrelia burgdorferi*: an invariant regulatory scheme that
554   promotes spirochete persistence in reservoir hosts and niche diversity. Front
555   Microbiol 10:1923.

556   36.   Burtnick MN, Downey JS, Brett PJ, Boylan JA, Frye JG, Hoover TR, Gherardini

557   FC. 2007. Insights into the complex regulation of *rpoS* in *Borrelia burgdorferi*. Mol

558   Microbiol 65:277-293.

559   37.   Caimano MJ, Iyer R, Eggers CH, Gonzalez C, Morton EA, Gilbert MA, Schwartz

560   I, Radolf JD. 2007. Analysis of the RpoS regulon in *Borrelia burgdorferi* in response

561   to mammalian host signals provides insight into RpoS function during the

562   enzootic cycle. Mol Microbiol 65:1193-1217.

563   38.   Yang XF, Lybecker MC, Pal U, Alani SM, Blevins J, Revel AT, Samuels DS,

564   Norgard MV. 2005. Analysis of the *ospC* regulatory element controlled by the

565   RpoN-RpoS regulatory pathway in *Borrelia burgdorferi*. J Bacteriol 187:4822-4829.

566   39.   Xu Q, McShan K, Liang FT. 2007. Identification of an *ospC* operator critical for

567   immune evasion of *Borrelia burgdorferi*. Mol Microbiol 64:220-231.

568   40.   Xu Q, McShan K, Liang FT. 2008. Verification and dissection of the *ospC* operator

569   by using *flaB* promoter as a reporter in *Borrelia burgdorferi*. Microb Pathog 45:70-

570   78.

571   41.   Drecktrah D, Hall LS, Hoon-Hanks LL, Samuels DS. 2013. An inverted repeat in

572   the ospC operator is required for induction in *Borrelia burgdorferi*. PLoS One

573   8:e68799.

574   42.   Tilly K, Casjens S, Stevenson B, Bono JL, Samuels DS, Hogan D, Rosa P. 1997. The

575   *Borrelia burgdorferi* circular plasmid cp26: conservation of plasmid structure and

576   targeted inactivation of the *ospC* gene. Mol Microbiol 25:361-373.

577   43.   Babb K, Bykowski T, Riley SP, Miller MC, DeMoll E, Stevenson B. 2006. *Borrelia*

578   *burgdorferi* EbfC, a novel, chromosomally-encoded protein, binds specific DNA

579   sequences adjacent to *erp* loci on the spirochete's resident cp32 prophages. J

580   Bacteriol 188:4331-4339.

581   44.   Burns LH, Adams CA, Riley SP, Jutras BL, Bowman A, Chenail AM, Cooley AE,

582   Haselhorst LA, Moore AM, Babb K, Fried MG, Stevenson B. 2010. BpaB, a novel

583   protein encoded by the Lyme disease spirochete's cp32 prophages, binds to *erp*

584   Operator 2 DNA. Nucleic Acids Res 38:5443-5455.

585   45.   Jutras BL, Chenail AM, Carroll DW, Miller MC, Zhu H, Bowman A, Stevenson B.

586   2013. Bpur, the Lyme disease spirochete's PUR-domain protein: identification as

587   a transcriptional modulator and characterization of nucleic acid interactions. J

588   Biol Chem 288:26220-26234.

589   46.   Sugino A, Higgins NO, Cozzarelli NR. 1908. DNA gyrase subunit stoichiometry

590   and the covalent attachment of subunit A to DNA during DNA cleavage. Nucleic

591   Acids Res 8:3865-3874.

592   47.   Reece RJ, Maxwell A. 1991. The C-terminal domain of the *Escherichia coli* DNA

593   gyrase A subunit is a DNA-binding protein. Nucleic Acids Res 19:1399-1405.

594   48.   Corbett KD, Shultzaberger RK, Berger JM. 2004. The C-terminal domain of DNA

595   gyrase A adopts a DNA-bending beta-pinwheel fold. Proc Natl Acad Sci USA

596   101:7293-7298.

597   49.   Knight SW, Samuels DS. 1999. Natural synthesis of a DNA-binding protein from

598   the C-terminal domain of DNA gyrase A in *Borrelia burgdorferi*. EMBO J 18:4875-

599   4881.

600   50.   Knight SW, Kimmel BJ, Eggers CH, Samuels DS. 2000. Disruption of the *Borrelia

601   burgdorferi gac* gene, encoding the naturally synthesized GyrA C-terminal

602   domain. J Bacteriol 182:2048-2051.

603   51.   Barbour AG, Garon CF. 1987. Linear plasmids of the bacterium *Borrelia

604   burgdorferi* have covalently closed ends. Science 237:409-411.

605   52.   Fraser CM, Casjens S, Huang WM, Sutton GG, Clayton R, Lathigra R, White O,

606   Ketchum KA, Dodson R, Hickey EK, Gwinn M, Dougherty B, Tomb J-F,

607   Fleischmann RD, Richardson D, Peterson J, Kerlavage AR, Quackenbush J,

608   Salzberg S, Hanson M, van Vugt R, Palmer N, Adams MD, Gocayne J,

609   Weidmann J, Utterback T, Watthey L, McDonald L, Artiach P, Bowman C,

610   Garland S, Fujii C, Cotton MD, Horst K, Roberts K, Hatch B, Smith HO, Venter

611   JC. 1997. Genomic sequence of a Lyme disease spirochaete, *Borrelia burgdorferi*.

612   Nature 390:580-586.

613   53.   Casjens S, Palmer N, van Vugt R, Huang WM, Stevenson B, Rosa P, Lathigra R,

614   Sutton G, Peterson J, Dodson RJ, Haft D, Hickey E, Gwinn M, White O, Fraser C.

615   2000. A bacterial genome in flux: the twelve linear and nine circular

616   extrachromosomal DNAs of an infectious isolate of the Lyme disease spirochete

617   *Borrelia burgdorferi*. Mol Microbiol 35:490-516.

618   54.   Dorman CJ, Schumacher MA, Bush MJ, Brennan RG, Buttner MJ. 2020. When is a

619   transcription factor a NAP? Curr Opin Microbiol 55:26-33.

620   55.   Amemiya HM, Schroeder J, Freddolino PL. 2021. Nucleoid-associated proteins

621   shape chromatin structure and transcriptional regulation across the bacterial

622   kingdom. Transcription 12:182-218.

623   56.   Muskhelishvili G, Sobetzko P, Mehandziska S, Travers A. 2021. Composition of

624   transcription machinery and its crosstalk with nucleoid-associated proteins and

625   global transcription factors. Biomolecules 11:924.

626   57.   Jutras BL, Bowman A, Brissette CA, Adams CA, Verma A, Chenail AM,

627   Stevenson B. 2012. EbfC (YbaB) is a new type of bacterial nucleoid-associated

628   protein, and a global regulator of gene expression in the Lyme disease

629   spirochete. J Bacteriol 194:3395-3406.

58.   Burgdorfer W, Barbour AG, Hayes SF, Benach JL, Grunwaldt E, Davis JP. 1982.

Lyme disease - a tick-borne spirochetosis? Science 216:1317-1319.

59.   Johnson RC, Schmid GP, Hyde FW, Steigerwalt AG, Brenner DJ. 1984. *Borrelia*

burgdorferi sp. nov.: etiologic agent of Lyme disease. Int J Syst Bacteriol 34:496-

497.

60.   Miller JC, von Lackum K, Babb K, McAlister JD, Stevenson B. 2003. Temporal

analysis of *Borrelia burgdorferi* Erp protein expression throughout the mammal-

tick infectious cycle. Infect Immun 71:6943-6952.

61.   Xu Y, Johnson RC. 1995. Analysis and comparison of plasmid profiles of *Borrelia*

burgdorferi sensu lato strains. J Clin Microbiol 33:2679-2685.

62.   Kraiczy P, Seling A, Brissette CA, Rossmann E, Hunfeld K-P, Bykowski T, Burns

LH, Troese MJ, Cooley AE, Miller JC, Brade V, Wallich R, Casjens S, Stevenson B.

2008. *Borrelia burgdorferi* complement regulator-acquiring surface protein 2

(CspZ) as a serological marker of human Lyme disease. Clin Vaccine Immunol

15:484-491.

63.   Zückert WR. 2007. Laboratory maintenance of *Borrelia burgdorferi*. Curr Protoc

Microbiol 12C:1-10.

64.   Samuels DS, Mach KE, Garon CF. 1994. Genetic transformation of the Lyme

648   disease agent *Borrelia burgdorferi* with coumarin-resistant *gyrB*. J Bacteriol

649   176:6045-6049.

650   65.   Elias AF, Bono JL, Kupko JJ, Stewart PE, Krum JG, Rosa PA. 2003. New antibiotic

651   resistance cassettes suitable for genetic studies in *Borrelia burgdorferi*. J Mol

652   Microbiol Biotechnol 6:29-40.

653   66.   Babb K, McAlister JD, Miller JC, Stevenson B. 2004. Molecular characterization of

654   *Borrelia burgdorferi erp* promoter/operator elements. J Bacteriol 186:2745-2756.

655   67.   Frank KL, Bundle SF, Kresge ME, Eggers CE, Samuels DS. 2003. *aadA* confers

656   streptomycin resistance in *Borrelia burgdorferi*. J Bacteriol 185:6723-6727.

657   68.   Bono JL, Elias AF, Kupko JJ, Stevenson B, Tilly K, Rosa P. 2000. Efficient targeted

658   mutagenesis in *Borrelia burgdorferi*. J Bacteriol 182:2445-2452.

659   69.   Stewart PE, Thalken R, Bono JL, Rosa P. 2001. Isolation of a circular plasmid

660   region sufficient for autonomous replication and transformation of infectious

661   *Borrelia burgdorferi*. Mol Microbiol 39:714-721.

662   70.   Babb K, El-Hage N, Miller JC, Carroll JA, Stevenson B. 2001. Distinct regulatory

663   pathways control the synthesis of *Borrelia burgdorferi* infection-associated OspC

664   and Erp surface proteins. Infect Immun 69:4146-4153.

71.   Jutras BL, Verma A, Stevenson B. 2012. Identification of novel DNA-binding

proteins using DNA-affinity chromatography/pull down. Curr Protocols

Microbiol 1F:1-13.

72.   Barbour AG, Hayes SF, Heiland RA, Schrumpf ME, Tessier SL. 1986. A *Borrelia-*

specific monoclonal antibody binds to a flagellar epitope. Infect Immun 52:549-

554.

73.   Mbow ML, Gilmore RD, Stevenson B, Golde WT, Piesman J, Johnson BJB. 2002.

Borrelia burgdorferi-specific monoclonal antibodies derived from mice primed

with Lyme disease spirochete-infected *Ixodes scapularis* ticks. Hybrid

Hybridomics 21:179-182.

74.   Arnold WK, Savage CR, Brissette CA, Seshu J, Livny J, Stevenson B. 2016. RNA-

Seq of *Borrelia burgdorferi* in Multiple Phases of Growth Reveals Insights into the

Dynamics of Gene Expression, Transcriptome Architecture, and Noncoding

RNAs. PLoS One 11:e0164165.

Figure Legends

Figure 1. Alignment of the DNA sequences 5′ of the *ospC* ORFs of *B. burgdorferi* ss B31

and *B. garinii* Ip89, which includes the *ospC* operator (39-41). Both strains contain

multiple copies of an 11bp repeated sequence (blue arrows), with Ip89 carrying seven

directly repeated copies. Additionally, both strains contain an inverted copy of this

sequence between their -35 and -10 promoter elements. A previous study noted an

inverted repeat sequence in the operator of strain B31 (red arrows), which is not found

in strain Ip89 or many other Lyme disease borreliae (41, 42).

Figure 2. (A) Electrophoretic mobility shift assay (EMSA) of labeled Ip89 *ospC* operator,

without (-) and with (+) cytoplasmic extract from strain B31. The asterisk indicates

shifted probe that was present only when borrelial extract was included, indicating

presence of a DNA-binding component. Unbound DNA is indicated by "free". Some

labeled DNA probe remained in the wells of the gel. **(B)** Cytoplasmic extract from strain

B31 was incubated with Ip89 *ospC* operator DNA affixed to beads, eluted with

increasing concentrations of NaCl, and eluates were separated by SDS-polyacrylamide

gel electrophoresis. Lanes are labeled with the concentration of NaCl used for that

elution. The arrow indicated a protein band of 30-35 kDa that eluted in the 0.3 and 0.5

mM NaCl washes. **(C)** Amino acid sequence of GyrA and Gac (the two potential

initiation methionines of Gac are indicated by boldface and underlining). Mass

spectrometry identified nine unique peptides that all mapped to the Gac ORF. Peptides

identified with high confidence are shaded green, and those with lower confidence in

red. **(D)** Predicted structures of *B. burgdorferi* Gac and GyrA proteins, as determined by

AlphaFold (https://alphafold.ebi.ac.uk). Both are consistent with the solved structures

of *B. burgdorferi* Gac (48) and the N-terminal domain of *E. coli* GyrA

(https://www.rcsb.org/structure/4CKK).

Figure 3. EMSAs of purified recombinant Gac with labeled *ospC* operator DNAs and

unlabeled competitors. Shifted DNAs are indicated with asterisks. Unoccupied DNAs

are labeled "free". **(A)** Labeled Ip89 operator (100 nM) without (lane 1) and with 25 nM

Gac (lanes 2-8). Lane 2: no added competitor. Lane 3: plus 10x excess unlabeled Ip89

ospC operator. Lane 4: plus 25x excess unlabeled Ip89 *ospC* operator. Lane 5: plus 25x

excess unlabeled B31 *ospC* operator. Lane 6: plus 25x unlabeled pCR2.1 amplicon. Lane

7: plus 25x excess unlabeled *flaB* DNA. Lane 8: plus 25x unlabeled *ospAB* DNA. **(B)**

Labeled B31 operator (100 nM) without (lane 1) and with 25 nM Gac (lanes 2-7). Lane 2:

no added competitor. Lane 3: plus 10x excess unlabeled B31 *ospC* operator. Lane 4: plus

25x excess unlabeled B31 *ospC* operator. Lane 5: plus 10x unlabeled pCR2.1 amplicon.

Lane 6: plus 10x excess unlabeled *flaB* DNA. Lane 7: plus 10x unlabeled *ospAB* DNA.

Figure 4. Mapping Gac binding sites in B31 *ospC* 5′ DNA by EMSA with purified

recombinant Gac and unlabeled competitor DNA fragments. Sequences of the probe

and competitors are shown above the EMSA, with the 11bp repeats indicated by blue

arrows. Shifted DNAs are indicated with asterisks. Unoccupied DNAs are labeled

"free". Labeled B31 operator (100 nM) without (lane 1) and with 25 nM Gac (lanes 2-11).

Lane 2: no added competitor. Lane 3: plus 10x excess unlabeled B31 *ospC* operator

probe. Lane 4: plus 25x excess unlabeled B31 *ospC* operator probe. Lanes 5-11: plus 10x

unlabeled competitors A through G, respectively.

Figure 5. Western blot analyses of strains NGR, CKO-1, and CKO-1(pBLS820) that were

cultured to mid-exponential phase at 23 or 35˚C, probed with monospecific antibodies

against OspC and FlaB. In the illustrated experiment, strain CKO-1 produced 14-times

and 1.4-times more OspC than did NGR at 23 and 34˚C, respectively. Complementation

of CKO-1 with a wild-type *gac* gene on pBLS820 led to 32% and 20% reduction in OspC

at 23 and 34˚C, respectively.

Figure 6. Flow cytometric analyses of green fluorescent protein (GFP) levels. Each panel

represent approximately 100,000 bacteria. Isogenic NGR and *gac* mutant CKO-1

contained transcriptional fusions of the *ospC* promoter, with an without the operator

(pBLS761 and pBLS764, respectively), fused to *gfp*. Negative control strain KS10 carries

promoterless *gfp* on pBLS590, and positive control strain KS20 carries the constitutive

erpAB promoter fused to *gfp* on pBLS599 (66). *B. burgdorferi* were cultured to mid-

exponential phase at 35˚C.

Figure 7. Growth curves and quantitative-RT-PCR analyses of selected transcripts of *B.*

burgdorferi B31, cultured at 35˚C. Bacterial cell densities are indicated by black squares

and lines. Relative transcript levels are indicated by red circles and lines. **(A)** Schematic

of the locations of amplicons used to determine relative expression of the 5' and 3'

portions of *gyrA*. The 5' primer pair amplify only transcripts arising from the promoter

5' of *gyrB*, while the 3' primer pair amplify transcripts arising from both the promoter 5'

of *gyrB* and within the *gyrA* ORF. **(B)** Results from the 5' segment of *gyrA*. **(C)** Results from

the 3' segment of *gyrA*. **(D)** Results from the *gyrA* 3' segment minus the *gyrA* 5'. **(E)**

Analyses of *ospC* transcript levels.

Figure 8. EMSAs of purified recombinant Gac with labeled *ospC* lp17 telomere DNA

and competitors. Shifted DNAs are indicated with asterisks. Unoccupied DNA is

labeled "free". All lanes contain 100mM labeled lp17 telomere probe, without (lane 1)

or with 25 nM Gac (lanes 2-9). Lane 2: no added competitor. Lane 3: plus 10x excess

unlabeled lp17 telomere. Lane 4: plus 25x excess unlabeled lp17 telomere. Lane 5: plus

10x unlabeled pCR2.1 amplicon. Lane 6: plus 10x excess unlabeled Ip89 *ospC* operator.

Lane 7: plus 10x excess unlabeled B31 *ospC* operator. Lane 8: plus 10x excess unlabeled

flaB DNA. Lane 9: plus 10x unlabeled *ospAB* DNA.

Figure 9. Graphic summary of these results. Gac binds at or near the 11bp repeated

sequences in the *ospC* promoter and operator, and inhibits transcription from the *ospC*

promoter. Deletion of the *ospC* operator resulted in ablation of transcription from the

ospC promoter, as has also been noted previously (39-41). Those data indicate that

another factor, "X", interacts with the *ospC* operator and is required to activate

transcription. In the absence of Gac, the *ospC* promoter is transcribed at high levels,

indicating that "X" competes against Gac repression. X may be one or more protein or

other factor. The 11bp repeat sequences are indicated by blue arrows beneath the DNA.

The *ospC* promoter -10 and -35 sequences are indicated by black bars above the DNA.

The

Supplemental Material

Figure S1. Sequence of the *gac* 5′ and open reading frame, which was cloned between

the KpnI and BamHI sites of vector pKFSS1 to produce pBLS820. The -10 and -13 sites

of the transcriptional promoter and the ribosome binding site (RBS) are indicated in

purple. The *gac* open reading frame probably begins at the 5′-most ATG (methionine)

codon, although a second, downstream ATG codon could also be used. Both of those

initiation codons, and the RBS, were mutated in *B. burgdorferi* strain CKO-1, which

prevented production of Gac protein (50). The TAA termination codon of *gac* is shown

in red.

Table S1. Oligonucleotide primers used in this study.

B31   GCATATTTGGCTTTGCTTATGTCGATTTTAAAAATCAAAATTAAGA
Ip89   GCATATTTGGCTTTGCTTCTGTCAATCTTATAATCAATTAAGAT

B31   CAATATTTTTCAAAATTCTT CAATATTTATT
Ip89   AGTTCGTTTGTAATATTT CAATAATTATT CAATAATTTTT CAATAATTATT CAATAATTTTT CAATAATTATT GAAAAATTATT CAAATTATTT

-35

-10

B31   CAAGATATTGAAGAATTGAAAAATTATTTTTTTCAAATAAAAAAGGAGG   CAAAATTGTTG   GACTAATAATTCATAAATAAAAAGGAGG
Ip89   --------ATTGAAAAATTATTTTTTTTTATATAAAGAATTGAAAAG   AAAAATTGTTG   GACTAATAATTTATATAAAAAGGAGG

Figure 1

MAVGENKEQILNVRIEDEIKTSYLNYAMSVIVSRALPDVRDGLKPVHRRILYSMYEMGLRSDKAFKKAGRIVGDVLGKYHPH
GDQSIYDALVRLAQDFSLRYPVIRGQGNFGSIDGDPPAAMRYTEAKMEKITEYIVKDIDKETVNFKSNYDDSLSEPEIMPSS
FPFLLVNGSSGIAVGMATNMAPHNLREICDAIVYMLDNENASIFDLLKIVKGPDFPTFGEIVYNDNLIKAYKTGKGSVVIRA
RYHIEERAEDRNAIIVTEIPYTVNKSALLMKVALLAKEEKLEGLLDIRDESDREGIRIVLEVKRGFDPHVIMNLLYEYTEFK
KHFSINNLALVNGIPKQLNLEELLFEFIEHRKNIIERRIEFDLRKAKEKAHVLEGLNIALNNIDEVIKIIKSSKLAKDARER
LVSNFGLSEIQANSVLDMRLQKLTALEIFKLEEELNILLSLIKDYEDILLNPVRIINIIREETINLGLKFGDERRTKIIYDE
EVLKTS**M**SDL**M**QKENIVVMLTKKGFLKRLSQNEYKLQGTGGKGLSSFDLNDGDEIVIALCVNTHDYLFMISNEGKLYLINAY
EIKDSSRASKGQNISELINLGDQEEILTIKNSKDLTDDAYLLLTTASGKIARFESTDFKAVKSRGVIVIKLNDKDFVTSAEI
VFKDEKVICLSKKGSAFIFNSRDVRLTNRGTQGVCGMKLKEGDLFVKVLSVKENPYLLIVSENGYGKRLNMSKISELKRGAT
GYTSYKKSDKKAGSVVDAIAVSEDDEILLVSKRSKALRTVAGKVSEQGKDARGIQVLFLDNDSLVSVSKFIK

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

B

C

D

E

Figure 8

Figure 9