Diversity of the woodland strawberry inflorescences results from heterochrony

antagonistically regulated by *FvTFL1* and *FvFT1*

Short title: Inflorescence development in strawberry

Sergei Lembinen[1], Mikolaj Cieslak[2], Teng Zhang[1], Kathryn Mackenzie[1], Paula

Elomaa[1], Przemyslaw Prusinkiewicz[2*], Timo Hytönen[1*].

[1] Department of Agricultural Sciences, Viikki Plant Science Centre, University of

Helsinki, P.O. Box 27, FIN-00014 Helsinki, Finland

[2] Department of Computer Science, University of Calgary, Calgary, AB T2N 1N4,

Canada

[*]Correspondence:

Timo Hytönen: timo.hytonen@helsinki.fi

Przemyslaw Prusinkiewicz: pwp@ucalgary.ca

The authors responsible for distribution of materials integral to the findings presented

in this article in accordance with the policy described in the Instructions for Authors

(https://academic.oup.com/plcell/pages/General-Instructions) are: Timo Hytönen

(timo.hytonen@helsinki.fi) and Przemyslaw Prusinkiewicz (pwp@ucalgary.ca).

.

Abstract

A vast variety of inflorescence architectures have formed during angiosperm evolution. Here we analyze the diversity and development of the woodland strawberry inflorescence. We show that it is a thyrse: a compound inflorescence in which the primary monopodial axis supports lateral sympodial branches, thus combining features of racemes and cymes. We demonstrate that this architecture is related to differences in the size and shape of the primary and lateral inflorescence meristems. We further show that woodland strawberry homologs of TERMINAL FLOWER 1 (TFL1) and FLOWERING LOCUS T (FT) antagonistically regulate the development of both the racemose and cymose components of the strawberry thyrse: the loss of functional *FvTFL1* and overexpression of *FvFT1* reduce the number and complexity of the cymose components, whereas silencing of *FvFT1* has the opposite effect and can partially rescue the *fvtfl1* mutation. We complement our experimental findings with a computational model, which captures the development of the woodland strawberry inflorescence using a small set of rules, and shows that its phenotypic diversity can be explained in terms of heterochrony resulting from the opposite action of FvTFL1 and FvFT1 on the progression from the branching to flowering state.

Key words: *Fragaria,* inflorescence, thyrse, meristem, branching, modelling, L-system, Petri net, vegetativeness,

INTRODUCTION

The arrangement of individual flowers in time and space is critical for plants' reproductive success (Harder et al., 2004; Harder & Prusinkiwicz, 2013), and ultimately for agricultural yield and fruit uniformity (Eshed & Lippman, 2019). Many angiosperms organize their flowers into clusters called inflorescences. The structural arrangement of inflorescences has long attracted the attention of plant scientists, horticulturalists, and agronomists, who described, explained, and enhanced through breeding the diverse inflorescence architectures.

Based on their branching patterns, a vast variety of inflorescence architectures can be classified as monopodial or sympodial (Weberling, 1989; Prusinkiewicz & Lindenmayer 1990). In monopodial inflorescences, or racemes (Figure 1A), new flowers form at the lateral positions on the indeterminate or determinate primary axis. In contrast, in cymose or sympodial inflorescences (Figure 1B), new flowers form at the terminal positions, and the new growth axes are established laterally. The classification of inflorescence architectures is complicated by the occurrence of compound inflorescences. For instance, panicles, or polypodial inflorescences, consist of repetitively branching determinate or indeterminate monopodial axes (Figure 1C). In contrast, in thyrses, the primary indeterminate or determinate (Prenner et al., 2009) monopodial axis bears sympodial branches (Figure 1C).

The development of inflorescence branching patterns is regulated by the activity of meristems, the small groups of multipotent cells located at the branch tips. Inflorescence meristems (IMs) are capable of initiating new IMs (i.e., branching axes) and eventually transition to flower meristems (FMs), which produce flower organs. Differences in the timing of these transitions – a form of heterochrony – control the diversity of inflorescence architectures (Grimes, 1993; Lemmon et al., 2016; Prusinkiewicz et al., 2007).

84

85 **Figure 1.** Inflorescence branching types. **A.** Monopodial branching of a simple
86 raceme. **B.** Sympodial branching of monochasial and dichasial cymes. **C.** Compound
87 branching of a panicle and a thyrse (dichasial)
88

The molecular processes that regulate developmental decisions in meristems have
been unraveled in model plants with monopodial as well as sympodial
inflorescences. In simple racemes, such as in *Arabidopsis thaliana*, the IM state is
maintained by TERMINAL FLOWER 1 (TFL1). Plants lacking a functional TFL1,
which is normally expressed in the center of the developing IM, produce determinate
inflorescences with a few flowers (Alvarez et al., 1992; Schultz & Haughn, 1993). On
the IM flanks, the expression of *LEAFY* (*LFY*) and *APETALA1* (*AP1*) activates the
floral organ identity genes, promoting FM identity. Inactivation of *LFY* prevents
specification of FM identity and transforms flowers into shoots (Weigel et al., 1992;
Weigel & Meyerowitz, 1993).

Subsequent studies in Arabidopsis revealed that a *TFL1* homolog *FLOWERING*
LOCUS T (*FT*) is also involved in regulating inflorescence architecture (Lee et al.,
2019). FT and TFL1 belong to a family of phosphatidylethanolamine-binding proteins
(PEBPs) and integrate environmental signals such as photoperiod and temperature
to control transition to reproductive growth (reviewed in McGarry & Ayre 2012;
Rantanen et al., 2015). FT is a mobile protein which is produced in leaves and
moves into the shoot apical meristem (SAM) to promote flowering (Corbesier et al.,
2007). In racemose Arabidopsis, the loss of functional FT promotes meristem
indeterminacy and inflorescence branching (Lee et al., 2019). Further experiments
revealed that FT and TFL1 proteins compete for the same binding partner, a bZIP
transcription factor FD, and form complexes via 14-3-3 proteins (Zhu et al., 2021).
When formed, FT-FD and TFL1-FD complexes act as transcriptional activators and

repressors, respectively, affecting multiple target genes including the meristem identity regulators *LFY* and *AP1* (Zhu et al., 2020).

Studies in plants with cymose inflorescences have revealed both similarities and differences in the genetic control of inflorescence development, compared to racemes. For example, in petunia (*Petunia hybrida*) and tomato (*Solanum lycopersicum*), both members of the *Solanaceae* family, the orthologs of *LFY* — *FALSIFLORA* (*FA*) and *ABERRANT LEAVES AND FLOWERS* (*ALF*) — are expressed in the terminal meristems and promote the transition to FM state (Molinero-Rosales et al., 1999; Souer et al., 2008). However, to fully establish FM identity, FA and ALF require a co-factor. In petunia, DOUBLE TOP (DOT), an ortholog of UNUSUAL FLOWER ORGANS (UFO), physically interacts with ALF to establish the FM identity (Souer et al., 2008). Interestingly, inactivation of *SELF-PRUNING* (*SP*)*,* an ortholog of *TFL1* in tomato, does not directly affect inflorescence architecture (Pnueli et al., 1998). In contrast, it was shown that *FT* homolog *SINGLE FLOWER TRUSS* (*SFT*; Molinero-Roslaes et al., 2004) regulates inflorescence branching in tomato (Park et al., 2012; Quinet et al., 2006).

Woodland strawberry (*Fragaria vesca* L.) is a perennial rosette plant and a model for the octoploid cultivated strawberry (*Fragaria × ananassa*) and the Rosaceae family in general (Edger et al. 2018). Historically, inflorescence architecture of *Fragaria* has been classified as pleiochasial cyme (Valleau, 1918), dichasial cyme (Anderson & Guttridge 1982; Guttridge, 1985), cyme (Menzel, 2019) and corymb (Gleason & Cronquist, 1991). This diversity of terms reflects the extensive morphological variability of strawberry inflorescences, which was highlighted almost a century ago (Darrow, 1929). The variability was attributed to the environment, genetic differences between specific cultivars (Darrow, 1929; Foster & Janick 1969; Anderson & Guttridge 1982), or to the sex of the flowers (Ashman & Hitchens, 2000). However, the developmental and genetic basis of inflorescence architecture in strawberry has never been elucidated.

Here we analyze inflorescence architecture and development in woodland strawberry and begin to dissect the underlying molecular mechanism regulating its development. We show that the inflorescences of woodland strawberry have compound architecture, combining a monopodial primary axis with sympodial lateral

branches. We summarize these findings in a computational model of thyrse
development, which supports the diversity of strawberry inflorescence architectures
due to heterochrony controlled by strawberry homologs of *FT* and *TFL1*.

RESULTS

Inflorescence of woodland strawberry is a determinate thyrse

To understand the variability of strawberry inflorescences, we analyzed inflorescence
development in diploid woodland strawberry. A woodland strawberry inflorescence
has a monopodial primary axis terminated by a flower. This axis typically supports
two (Figure 2A) and occasionally three (Figure 2B) secondary branches separated
by long internodes. In many monopodial structures, the main axis is relatively
straight and can be recognized easily. In woodland strawberry, however, secondary
branches often assume a dominant position, continuing approximately in the
direction of their supporting internodes, whereas the primary axis changes direction
at each branching point. To identify the course of the primary axis we thus relied on
the positions of bracts (b; Figure 2) associated with the branches they subtend.

The architecture of lateral branches is different from that of the main axis. Each
secondary axis supports a pair of third-order branches subtended by bracts, then
terminates with a secondary flower. The third-order branches are approximately
opposite each other: the internode between them is practically absent. This
branching pattern repeats with each third-order branch producing a pair of fourth-
order branches and a terminal flower, and typically continues up to fifth- or sixth-
order branches. At high branching orders, only one lateral branch may emerge,
although both bracts are present (Jahn & Dana 1970). In spite of this departure from
symmetry, the structures supported by the primary axis are best characterized as
dichasial cymes (Figure 1B; Weberling, 1989; Prenner et al., 2009). With the
determinate primary monopodial axis supporting sympodial branches, inflorescence
of woodland strawberry qualifies as a determinate thyrse (Figure 1C).

Characterizing these inflorescences further, we observed that they are basitonic, i.e.,
the branches originating near the inflorescence base are more elaborate than those
near the top (Lück et al., 1990). The first lateral branch growing from the axil of the

leaf-like bract b1 is larger and bears more flowers than the branch that originates

from the axil of b2 (Figure 2, A and B). The third lateral branch, if present, is even

smaller and produces fewer flowers than the second or the first lateral branch. The

internode between b2 and b3 is also shorter than between b1 and b2 (Figure 2, A

and B). Moreover, the size of successive bracts is dramatically reduced, and the

shape is simplified from a three-lobed structure resembling a leaf to a bracteole-like

structure with a single lobe (Figure 2C). Within branches, the bracts are more similar

in shape, but also decrease in size with each successive branching iteration.

Two meristem types produce determinate thyrse of woodland strawberry

To investigate the ontogenesis of the strawberry inflorescence, we obtained a

developmental sequence of inflorescence meristems using SEM imaging (Figure 2,

D─M). The transition to flowering is associated with bulging and an almost two-fold

increase in the size of the primary inflorescence meristem (i.e., the meristem that

generates the primary inflorescence axis: IM1), compared to the vegetative shoot

apical meristem (SAM) (Figure 2, D and E). IM1 has an approximately circular cross-

section and produces bracts and lateral meristems (IM2) that are almost as large as

IM1. The first bract (b1) is initiated during the SAM to IM1 transition (Figure 2E), with

a lateral meristem (IM2) then forming in the bract axil (Figure 2F). As the primary

meristem (IM1) continues to grow, an additional one or two bracts with associated

lateral meristems (IM2) arise sequentially (Figure 2, G, H and N), leading to the

monopodial architecture of the main axis. Due to their size, these bracts and

meristems push the primary meristem (IM1) to the side (Figure 2O), which changes

the course of the main axis at each branching point (Figure 2P) as evident in the

mature inflorescences (e.g., Figure 2, A and B). The patterning of the main axis is

terminated by the primary meristem acquiring flower identity and switching to the

production of flower organs (FM1; Figure 2H).

The second and higher-order inflorescence meristems extend along the perimeter of

their parent meristems and have an elongated, crescent-like shape (Figure 2, F and

G). In consequence, the space needed to form the next-order meristems with

associated bracts emerges concurrently near both ends of their parent meristem

(Figure 2, I and J). These primordia become meristems that may produce the next

iteration of the same pattern, while the parent meristem continues to grow without

changing direction and develops into a flower (Figure 2, K–M). The compound dichasial structure of the woodland strawberry inflorescence thus results.

However, one end of each meristem of the third order (IM3) is inevitably closer to the inflorescence center than the other end, which leads to the uneven restriction of space available for the development of the fourth order primordia (IM4; Figure 2Q). A similar asymmetry occurs in higher order meristems, which may be the cause of, or contribute to, the commonly observed departure from symmetry at high branching orders, where only one lateral branch is present.

Figure 2. Architecture and development of thyrse inflorescence in woodland strawberry. **A – B.** Inflorescence architectures of wild-type woodland strawberry plants with two **(A)** and three **(B)** lateral branches on the primary axis. White arrowheads indicate the primary flowers; white lines with arrows indicate the

distances between bracts on the primary branching axes; numbers in circles indicate branching iterations **C.** Typical bract phenotypes at branching points of different orders. **D–H.** Development of the primary inflorescence meristem (IM). **D.** Round shoot apical meristem (SAM) with monopodially produced leaves. **E.** Early IM1 of increased size, compared to the SAM. **F.** Initiation of the first lateral IM2. **G.** Initiation of the second lateral IM2. **H.** IM1 transitioned into flower meristem (FM1). **I–M**. Development of the lateral meristems. **I.** Elongated, crescent-like shape of IM2. **J.** An almost simultaneous initiation of the lateral IM3s at the ends of IM2. **K–M.** Transition of IM2 into FM2 and development of IM3s. **N.** Primary axis with three bracts, each subtending a lateral meristem. **O.** The first lateral branch (white arrow) displaces the primary flower (F1) and assumes a more dominant position. **P.** Schematic representation of thyrse inflorescence with leading lateral branch. The primary axis is highlighted in darker green. **Q.** Schematic representation of relative meristem positions during the development of woodland strawberry thyrse. Relatively more confined IMs are highlighted in red. Scale bars in **D–N** equal 100 µm. b – bract.

FvTFL1 controls inflorescence architecture in woodland strawberry

In many plant species, inflorescence architecture and flowering time are tightly connected (Lee et al., 2019; Lifschitz et al., 2006; Liu et al., 2013). With this in mind, we looked for the genetic regulators that can affect architecture and development of the strawberry thyrse. In woodland strawberry, a mutation in *FvTFL1* accelerates flowering and reverses the photoperiodic requirement for flower induction from short-day to long-day (Koskela et al., 2012). To understand whether FvTFL1 also controls inflorescence development in woodland strawberry, we analyzed the inflorescence architecture and branching in seven *fvtfl1* mutants and nine wild type (WT) genotypes collected across Europe (Supplemental Figure S1, A–C). All *fvtfl1* cultivars used in this study were found to have a 2-bp deletion in the first exon of *FvTFL1* (Supplemental Figure S2) putatively leading to a nonfunctional FvTFL1 protein.

We found that the number of flowers per inflorescence was reduced more than two-fold in the *fvtfl1* mutant plants. On average, WT plants produced 14.2 ± 1.2 flowers per inflorescence, while *fvtfl1* mutants produced only 6.5 ± 1.4 flowers per inflorescence ($p < 0.0001$; Figure 3A). This reduction of flower numbers was associated with changes in the general architecture of inflorescences, particularly with dramatic changes in the number of lateral branches on the primary axis (Figure 3B–D). In plants with functional FvTFL1, we observed a high proportion of inflorescences with two (72%), and three (27%), lateral branches produced on the

primary axis. Only 1% of the inflorescences had a single secondary branch. In
contrast, the majority of inflorescences of *fvtfl1* mutant plants (78%) produced only
one secondary branch (Figure 3C) and the remaining 22% had two secondary
branches. Such a dramatic change in the proportion of lateral branches supported by
the primary axis suggests that in *fvtfl1* mutants IM1 transitions to FM1 before it is
able to produce the second lateral branch on the primary axis.

In the lateral cymose branches, the loss of functional FvTFL1 did not affect the
dichasial branching pattern. However, the total number of branching iterations in the
fvtfl1 mutants (4.3 ± 1.1) was reduced compared to the WT plants (5.1 ± 1.0; p =
0.02; Figure 3D). To understand the reasons behind this variation we compared the
bracts of *fvtfl1* mutants and WT plants (Figure 3E). The first bract (b1) on the primary
axis was the largest and typically had three lobes in plants with functional and non-
functional FvTFL1. Furthermore, in both groups of plants, each successive lateral IM
produced smaller bracts; however, at each corresponding branching point, WT
plants had larger bracts than *fvtfl1* mutants.

271
Figure 3. FvTFL1 controls inflorescence architecture in woodland strawberry. **A.**
Number of flowers on the first inflorescence of seven *fvtfl1* (red) and nine *FvTFL1*
(grey) genotypes. Boxplots and points show the distribution of raw data. Each point
represents an individual inflorescence. **B.** Percentage of observed inflorescences
with one, two, or three branches on the primary branching axis. **C.** Average number
of branching iterations along the longest branching path. Bars and whiskers
represent the mean ± standard deviation. Data in **A** and **C** were analyzed using a
generalized linear mixed model (GLMM), with accessions and cultivars nested within
fvtfl1 and *FvTFL1* groups. P values for significant differences between *fvtfl1* and
FvTFL1 groups of plants are shown (*t*-test). **D.** Inflorescence phenotype of *fvtfl1*

cultivar Reine des Vallées; white arrow indicates the position of the primary flower. b
– bract. **E.** Typical bract phenotypes along the longest branching path of *FvTFL1* and
fvtfl1 genotypes. Numbers in circles indicate branching iterations.
285
FvFT1 regulates inflorescence architecture in *fvtfl1* background

Next, we decided to investigate the role of *FvFT1* in the regulation of inflorescence

architecture. Previously, FvFT1 was found to promote flowering under long-day

conditions in the *fvtfl1* mutant background (Rantanen et al., 2014; Kurokura et al.,

2017). We analyzed the inflorescence architecture and branching in *fvtfl1* plants

(*Hawaii-4*) with silenced (RNAi) and overexpressed (35S) *FvFT1* (Supplemental

Figure S3). Overexpression (OX) of *FvFT1* caused a pleiotropic effect by

dramatically reducing the overall plant and leaf size and quickly transforming all

shoots into inflorescences (Supplemental Figure S4). *Hawaii-4* plants typically

produced inflorescences with 5.7 ± 1.9 flowers, whereas transgenic plants with

constitutive *FvFT1* expression produced inflorescences with only 1 – 3 flowers ($p <$

0.0001; Figure 4, A, D and E). In contrast, silencing of *FvFT1* caused an almost two-

fold increase in the number of flowers per inflorescence (12 ± 0.24; $p = 0.006$; Figure

4A). The main reason for the higher number of flowers per inflorescence was the

formation of the second lateral branch along the primary axis, as in the genotypes

with functional FvTFL1. We observed that about 80% of the inflorescences in *FvFT1-*

RNAi plants initiated two lateral branches on the primary axis, compared to only 20%

in *Hawaii-4* plants (Figure 4B). Furthermore, *FvFT1-RNAi* plants typically produced

more branching iterations than *Hawaii-4* ($p = 0.0001$; Figure 4, C, D and F). The

branching of the primary axis in *fvtfl1* mutant plants (*Hawaii-4*) with silenced *FvFT1*

expression was similar to WT plants with functional FvTFL1, except that we did not

observe inflorescences with three lateral branches on the primary axis in *FvFT1-*

RNAi lines (Figure 4F). Moreover, we found that *FvFT1-RNAi* plants occasionally

produced an intermediate phenotype, where the second bract (b2) on the primary

axis was associated with the sepals of the primary flower (Figure 4G). A similar

phenotype was observed in the *FvFT1-OX* lines, however in these lines the first

bract (b1) became associated with the sepals of the primary flower (Figure 4H).

In the plants with the functional FvTFL1, the overexpression of *FvFT1* did not cause

as severe a reduction in the overall plant size (Supplemental Figure S5A) as in

FvFT1-OX//fvtfl1 plants (Supplemental Figure S4). The number of flowers per

inflorescence of *FvFT1-OX* plants with functional FvTFL1 background was reduced compared to WT and *fvtfl1* mutant plants (Supplemental Figure S3B). However, it was higher than in the *FvFT1-OX* plants with *fvtfl1* background. Altogether our data thus suggests the direct antagonistic functions of FvFT1 and FvTFL1 in inflorescence development of woodland strawberry.

Figure 4. FvFT1 regulates inflorescence architecture in woodland strawberry. **A.** Number of flowers per inflorescence in *FvFT1* overexpression *(FT1-OX)*, *FvFT1* silenced *(FT1-RNAi),* and control *(Hawaii-4)* plants. Boxplots and points show the distribution of raw data. Each point represents an individual inflorescence. Up to five inflorescences per plant were examined. **B.** Maximum number of branching iterations in *FvFT1-RNAi* lines and control *(Hawaii-4)* plants. Bars and whiskers show the mean ± standard deviation of the raw data. Data in **A** and **B** were analyzed by fitting a generalized linear mixed model (GLMM); plants were grouped within construct type (OX, RNAi, or control). P values for significant differences between OX/RNAi and control are shown (*t*-test). **C.** Percentage of inflorescences with one and two branches along the primary axis. **D–F.** Inflorescence phenotypes of control *(Hawaii-4), FT1-OX5* and *FT1-RNAi6* plants. The numbers in circles indicate branching iterations. **G–H.** Intermediate phenotypes of *FvFT1-RNAi* and *FvFT1-OX* plants associating the second (b2) or first (b1) bract of the primary axis with the sepals of the primary flower. b – bract.

FvTFL1 and FvFT1 antagonistically regulate the inflorescence architecture in woodland strawberry

To understand the mechanism of the second lateral branch formation on the primary
axis we analyzed WT (*FIN56*), *fvtfl1* (*Hawaii-4*) and *fvtfl1//FvFT1-RNAi* plants using
SEM imaging (Figure 5, A–C). We compared the WT and *fvtfl1* plants at a stage
when the primary FMs showed a similar developmental phase. The first lateral
meristem (IM2) of *FIN56* with functional FvTFL1 was at a later developmental stage
(Figure 5A) than the first (and single) lateral meristem (IM2) of *Hawaii-4* (Figure 5B).
In *FIN56*, the lateral IM2 already formed a pair of axillary meristems (IM3s) and sepal
primordia, while in *Hawaii-4*, only the bracts could be clearly distinguished on the
single lateral IM2. A similar difference was also evident when *FvFT1* was silenced in
fvtfl1 background (Figure 5C). Overall, these findings suggest that the primary IM1 of
the plants with functional FvTFL1 or silenced *FvFT1* develops into FM1 slower than
in *fvtfl1* mutants, thus allowing formation of additional bracts and lateral IM2s.

In woodland strawberry, *FvTFL1* is highly expressed in the vegetative shoot apex
and gradually downregulated during flowering induction (Koskela et al. 2012).
Molecular antagonism between *TFL1* and *LFY* has been described as the
mechanism controlling inflorescence architecture in Arabidopsis (Schultz & Haughn,
1993). Subsequent studies in other species found that *AP1* may substitute *LFY* as
TFL1 antagonist in inflorescence development (Kobayashi et al., 2012; Ma et al.,
2017). Here we observed that the decrease of *FvTFL1* expression is associated with
the progression of the vegetative SAM towards FM identity (Figure 5D). The
expression of the strawberry homologs of *LFY* and *AP1* showed the opposite pattern
(Figure 5, E and F). The expression of both *FvLFYa* (*FvH4_5g09660*) and *FvAP1*
(*FvH4_6g29600*) began increasing in late IM1s, where *FvTFL1* was still present.
FvTFL1 had the lowest expression in the FM tissues, concordant with the highest
expression of *FvLFYa* and *FvAP1*. Altogether, our data suggests that the role of
TFL1 in maintaining IM identity (Alvarez et al., 1992; Schultz & Haughn, 1993) is
conserved in woodland strawberry.

To further elucidate the mechanism by which FvFT1 regulates inflorescence
architecture in the absence of functional FvTFL1, we analyzed the expression levels
of *FvTFL1, FvLFYa* and *FvAP1* in the SAMs of *FIN56* (*FvTFL1*)*, Hawaii-4* (*fvtfl1*) and

FvFT1-RNAi (*fvtfl1*) plants. We observed that the expression of *FvLFYa* and *FvAP1*

was lower in the SAMs of *FIN56* and *FvFT1-RNAi* plants compared to *fvtfl1* (*Hawaii-*

4) (Figure 5, H and J). In addition, the expression of *FvTFL1* was the lowest in the

fvtfl1 mutant (Figure 5G). Our results thus suggest that FvTFL1 and FvFT1

antagonistically control the transition to FM identity by regulating the expression of

FvLFYa and *FvAP1*.

376
Figure 5. FvTFL1 and FvFT1 antagonistically shape the inflorescence of woodland
strawberry. **A–C.** Primary axis development in *FvTFL1 WT* **(A)**, *fvtfl1* mutant **(B)**, and
FvFT1-RNAi **(C)** plants. White arrows denote the first lateral IM2. Scale bars equal
100µm. **D–F.** The expression pattern of *FvTFL1* **(D)**, *FvLFYa* **(E)***, and FvAP1* **(F)** in
different meristems of *FvTFL1 WT* (*FIN56*) during transition to flowering. **G–J.** The
expression pattern of *FvTFL1* **(G)**, *FvLFYa* **(H)**, and *FvAP1* **(J)** in the SAMs of FIN56
(WT), *Hawaii-4* (*fvtfl1*) and *FvFT1-RNAi* (*Hawaii-4*) plants. Bars and whiskers show
the mean ± standard error (n = 3 – 5). Expression levels were normalized to SAM (D
–F) or *Hawaii-4* (G–J). *FvMSI1* was used as a calibrator gene.
386

A computational model supports heterochrony as the key determinant of the
strawberry inflorescence diversity

We constructed a parametrized computational model of the woodland strawberry
inflorescence to show that the observed diversity of the inflorescence architecture

can be attributed to the regulation of the rate of development by *FvTFL1* and *FvFT1*.

The model is expressed as an L-system, with separate rules capturing the

monopodial development of the primary axis and the sympodial development of

branches. Generic rules for both types of branching (c.f. Prusinkiewicz &

Lindenmayer, 1990) are extended with a mechanism that controls transitions of

individual meristems to flowers and terminates the formation of new meristems at the

end of inflorescence development. These processes are controlled by a synthetic

variable (integrating many influences) called *vegetativeness* (*veg*) (Prusinkiewicz et

al., 2007), which is related to the concepts of a controller of phase switching (Schultz

and Haughn, 1993) and meristem maturation (Park et al., 2012). In the strawberry

model, *veg* decreases monotonically with time (Figure 6A). Its values are compared

to two thresholds, th_m and th_s, which are pertinent to the fate of monopodial and

sympodial meristems, respectively. As long as $veg \geq th_m$, the primary meristem

periodically produces lateral primordia subtended by bracts (Event 1 in Figure 6B[*]).

This process terminates when *veg* drops below th_m, at which point the meristem

produces a terminal flower (Event 2). A lateral primordium develops in turn into a

lateral meristem supported by a bract (Event 3), which subsequently produces two

next-order primordia and a terminal flower (Event 4). This process periodically

iterates as long as $veg \geq th_s$, giving rise to a compound dichasial cyme. Eventually

veg drops below th_s, which results in the production of bracts without associated

meristems (Event 5) and arrests further branching. The distinct development of

branches originating at the same node, in particular the case when only one lateral

branch is present, are captured by introducing a small difference in the value of

threshold th_s between sibling meristems. This difference may be related to the

unequal space available for the development of primordia on opposite sides of an

elongated meristem (Figure 2Q).

In addition to the thresholds controlling branching architecture, the model includes

thresholds bth_1 and bth_2 (Figure 6A), which control the transition of bracts from the 3-

lobed form to 1-lobed form and to narrow scales, respectively. The model also

includes a number of parameters and growth functions that control the geometry of

---

[*] The concept of Petri nets formally underlying Figure 6B is further described by Peterson (1981), and in application to plant modeling by Prusinkiewicz and Remphrey (2000). The use of Petri nets enhances the flowchart representation of inflorescence development proposed by Kellogg (2000) by explicitly representing events that produce multiple organs, which then develop concurrently.

the phenotypes of interest. Parameter values and functions can be chosen such that
the simulated inflorescence development (Supplemental Movie 1) closely
approximates observations (Figure 7). In particular, the deviation of the primary axis
from a straight course substantially affects the overall appearance of the
inflorescence, obscuring its thyrsoid architecture (Figure 6C). For details see the
supplemental Model Code, Model Description, and the parameter values listed in
Supplemental Table S2.

428
Figure 6. Developmental model of a woodland strawberry thyrse. **A**. Control of a
thyrse development by *vegetativeness* (*veg*). Crossing threshold th_m terminates the
production of primordia by the primary (monopodial) meristem; crossing threshold th_s
terminates production of lateral primordia by the sympodial meristems. Times th_m
and th_s at which these thresholds are crossed determine the inflorescence
complexity (extent of branching). Additional thresholds, *bth*1 and *bth*$_2$, control the
transitions of bracts from 3-lobed to 1-lobed to narrow-scale forms. **B.** Petri net
representation of the meristem production and fate. Circles and rectangles represent
plant organs. The term "lateral primordium" denotes an incipient structure yielding a
lateral meristem supported by a bract, or a bract alone. Short black bars represent
events taking place during development. These events are labelled by a number and
are associated with conditions under which they may take place. Arcs with arrows
represent relations between these events and the resulting structures. Two arrows
originating in the same rectangle indicate alternative organ fates. Two or three
arrows emanating from the same bar indicate production of multiple organs. Green-
colored arrows indicate the introduction of an internode. **C**. The impact of the
monopodial axis course on thyrse geometry. In the top model the monopodial axis is
straight; in the bottom it deviates in the direction opposite to the lateral branch, as
commonly observed in woodland strawberry (c.f. Figure 2 A, B, O and P).
448

449
Figure 7. Development of a woodland strawberry inflorescence. **A.** Snapshots of
inflorescence growth in a WT (GER12 x FIN2) woodland strawberry. Photographs
were taken with the inflorescence supported in a vertical position to expose the
branching pattern. Plants were induced to flower under 11°C and a 12h/12h
(day/night) photoperiod for 6 weeks, then moved to 17°C and a 18h/6h photoperiod
during inflorescence growth. Numbers indicate days of observation. **B.** Simulated
development calibrated to sequence (A).
457

We experimented with the model to capture the key inflorescence features of the
plants with different constitutions of *FvTFL1* and *FvFT1*. The most prominent trait
that distinguishes the architectures collected in Figure 8A is the decreasing
complexity (maximum order) of branching, caused by an accelerated cessation of
branching. This decrease is accompanied by the accelerated progression of bracts
from the three-lobed leaf-like form at the most basal position to the one-lobed and
small-scale forms at more distal positions. Both the branching architecture and bract
form are controlled in the model by the initial value and rate of decline of *veg*, and
the thresholds th_m, th_s, bth_1, and bth_2 with respect to *veg* (Figure 6A). We observed
that the sequence of inflorescence architectures (branching topologies) presented in
Figure 8A can be reproduced simply by gradually increasing the rate of decline of
veg, with minimal adjustments to other parameters except for the extreme phenotype
of the *fvtfl1//FvFT1-OX* plants (Figure 8B; Supplemental Movies 2–8; Supplemental
Table S1). A similar effect is caused by gradually reducing the initial value of *veg*.
This result supports the conclusion that the diversity of strawberry inflorescences is a

manifestation of heterochrony, in which *FvTFL1* decelerates, and *FvFT1* accelerates

the rate of *veg* decline, respectively.

Figure 8. Diversity of woodland strawberry inflorescence phenotypes can be
reproduced by altering model parameters. **A.** Observed diversity of inflorescence
phenotypes. Scale bar equals 1 cm. **B.** Phenotypes reproduced by the model. Lines
with text describe the genetic constitution of the plants. The color of the line indicates
the rate of *veg* decline (green – slower, red – faster).

DISCUSSION

Almost a century ago, Darrow (1929) pointed out the extensive variability of

inflorescence architectures in cultivated strawberry *(F. × ananassa)* varieties, which

he attributed to genetic differences between cultivars, environmental factors, and the

vigor of individual plants. As outlined in Darrow's inflorescence diagrams, the

variation in inflorescence architecture of strawberries is mainly due to the variable

number of branches on the primary inflorescence axis and branching iterations in the

lateral branches. Here, using diploid woodland strawberry, we established the

developmental and genetic basis of this variation. We have shown that the

inflorescences of woodland strawberry consist of a monopodial determinate primary

axis supporting sympodial lateral branches, an architecture characterized as a

determinate thyrse (Prenner et al., 2009). This architecture arises from differences

between the primary and lateral inflorescence meristems.

We have further demonstrated that the number of lateral branches on the primary

monopodial axis and the number of iterations in the lateral sympodial branches are

oppositely controlled by the two PEBP family proteins FvTFL1 and FvFT1. These

proteins act by changing the timing of developmental decisions that transition

inflorescence meristems to the flowering state, and eventually terminate the

formation of further meristems. We supported this observation with a computational

model which captures the diversity of studied phenotypes.

503   Two types of IMs produce thyrse inflorescence in woodland strawberry

In the ontogenic view of inflorescence diversity, distinct inflorescence architectures

are produced by different types of meristems (Claßen-Bockhoff & Bull-Hereñu,

2013). Our data shows that the determinate thyrse of woodland strawberry, which

combines monopodial and sympodial branching patterns, is produced by two types

of IMs. The monopodial primary axis is produced by a primary IM (IM1), which

originates from the SAM that has expanded and begun producing leaf-like bracts

upon entering the reproductive phase. Like the SAM, IM1 has a circular cross-

section and produces bracts acropetally. After producing one to three bracts, the

primary IM acquires FM identity, thus forming a closed monopodial axis. Each of the

bracts on the monopodial axis subtends the newly formed and rapidly growing lateral

IM, which, unlike the primary IM, has a crescent-shaped cross-section. According to

the first available space theory (Hofmeister, 1868), new organ primordia are initiated

in the largest available space between the previous ones. The initially elongated

shape of the lateral IMs thus promotes a dichasial branching pattern, by allowing the

pairwise initiation of next-order bracts and branches near opposite ends of the

meristem.

We observed that, despite their morphological differences, both types of IMs

followed a similar developmental progression towards FMs. Moreover, along the

monopodial axis and with each successive branching iteration, the leaf-like bracts

and the internodes were gradually reduced. By associating this reduction with the

monotonic decrease of *veg*, properly choosing thresholds th_m and th_s at which the

activity of each type of IM terminates, and setting additional parameters (e.g.,

branching angles) to agree with the observation of reference plants, we have been

able to model the development of the woodland strawberry thyrse. The model

confirms the usefulness of *veg* as a controller of development and sets the stage for
understanding the diversity of strawberry inflorescences.

FvTFL1 and FvFT1 antagonistically regulate the complexity of woodland
strawberry inflorescences

Previous studies uncovered the role of *FvTFL1* in the regulation of flowering time in
woodland strawberry (Koskela et al., 2012; Koskela et al., 2017). Flowering in WT
strawberry requires downregulation of *FvTFL1*, which triggers the expression of the
FM identity gene *FvAP1* (Koskela et al., 2012). Our results further show that *FvTFL1*
is gradually downregulated as the SAM progresses towards the IM and then to the
FM state. This expression pattern is different from that previously described in
racemose and cymose plants. In monopodial Arabidopsis, *TFL1* is expressed in the
SAM and IM and is specifically required to maintain IM identity (Alvarez et al., 1992;
Schultz & Haughn, 1993). In sympodial tomato, *SP* - the closest homolog of *TFL1* -
is expressed only in vegetative axillary meristems (Pnueli et al., 1998; Thouet et al.,
2012). Concordantly, the *tfl1* mutation in Arabidopsis leads to the formation of a
terminal flower on the otherwise indeterminate IM, whereas *sp* mutants in tomato do
not show alterations of inflorescence architecture. The determinate primary IM of
woodland strawberry is related to Arabidopsis in the sense that the loss of functional
FvTFL1 accelerates the formation of the terminal flower and cessation of the
formation of lateral meristems. As a result, only one lateral branch is formed on the
primary monopodial inflorescence axis of the *fvtfl1* mutant. Furthermore, in the lateral
sympodial branches, the maximum number of branching iterations is reduced in the
fvtfl1 mutant plants, consistent with the accelerated termination of IMs.

Mutations in *FT* homologs were shown to affect the architecture in both monopodial
and sympodial inflorescences. In Arabidopsis, FT, and its close homolog TSF,
promote FM identity transition and act antagonistically to TFL1 (Lee et al., 2019).
Furthermore, mutation in *SFT* was found to promote inflorescence branching in
tomato (Park et al., 2012), suggesting a similar function. In woodland strawberry,
FvFT1 was previously shown to promote flowering in *fvtfl1* mutant background
(Koskela et al., 2012; Rantanen et al., 2014). Here we found that the role of *FvFT1* in
inflorescence development of woodland strawberry is antagonistic to that of *FvTFL1*.
Our data suggests that the faster progression towards FM in the *fvtfl1* mutant plants
may be delayed or further accelerated by altering the expression of *FvFT1*.

Silencing *FvFT1* partially compensated for the lack of functional FvTFL1*,* as we

observed an increased percentage of inflorescences with two lateral branches on the

primary axis and an increased number of branching iterations. In contrast, the

ectopic *FvFT1* expression in the *fvtfl1* mutants caused pleiotropic changes in plant

architecture by inducing flowering in young seedlings, thus reducing overall plant

size, and resulting in inflorescences with only a few flowers. Furthermore, in the

functional FvTFL1 background, the overexpression of *FvFT1* resulted in an

inflorescence phenotype intermediate between *fvtfl1* and *fvtfl1//FvFT1-OX*, in line

with the proposed antagonistic function of these genes.

Recently it was shown that the mutation of *FvLFYa* in woodland strawberry causes

the homeotic conversion of flower organs into bracts and the production of additional

flowers (Zhang et al., 2022), indicating a conserved function for specifying FM

identity. In Arabidopsis and rice, FT and TFL1 proteins are known to competitively

bind to bZIP transcription factor FD and 14-3-3 proteins (Zhu et al., 2021; Kaneko-

Suzuki et al., 2018). When formed, the FT-FD complex was found to act as a

transcriptional activator of *LFY* and *AP1*, whereas TFL1-FD was found to act as a

repressor (Zhu et al., 2021). In our experiments, both *FvLFYa* and *FvAP1* were

downregulated in the plants with either functional FvTFL1 or silenced *FvFT1* in *fvtfl1*

background, suggesting a similar regulatory mechanism.

Altogether, our experimental data indicate that FvTFL1 and FvFT1 antagonistically

regulate the complexity of the woodland strawberry inflorescence architecture. With

the help of a computational model, we have further shown that the observed diversity

of strawberry architectures can be attributed to heterochrony, where FvTFL1 delays*,*

and FvFT1 accelerates the cessation of branching and the production of flowers.

MATERIALS AND METHODS

Plant material

Nine *Fragaria vesca* and seven *F. vesca semperflorens* (*fvtfl1* mutant) accessions

were used in this study (Supplemental Table S2). Previously reported *FvFT1*

overexpression (OX) and RNA silencing (RNAi) lines in *fvtfl1* (*Hawaii-4*) background

were used (Koskela et al., 2012; Rantanen et al., 2014). The specificity of the RNAi

construct to *FvFT1* was confirmed by analyzing the expression of *FvFT2* and *FvFT3*

in the FM tissues (Supplemental Figure S6A – B). To obtain *FvFT1-OX* plants in

FvTFL1 background, we crossed *FvFT1-OX5 (Hawaii-4)* line with *FvTFL1* accession

(FIN56) and self-pollinated the progeny. F_2 plants carrying FIN56 *FvTFL1* allele and

the desired *FvFT1* construct were used in the experiments.

Growth conditions and phenotyping

All plants were grown in greenhouse conditions with controlled temperature and

photoperiod. In the greenhouse, the plants were illuminated with 150µmol m^{-2} s^{-1}

light intensity using high-pressure sodium lamps (Airam 400W, Kerava, Finland).

Germinated from seeds or clonally propagated plants were first potted into 7x7cm

plastic pots filled with peat moss (Kekkilä, Finland), and kept in the greenhouse

under plastic covers for two weeks. Plants were pre-grown at 18^{o}C and an 18h

photoperiod for four - six weeks, and periodically supplemented with fertilizer (NPK

17-4-25; Kekkilä). Plants were then potted into 13cm Ø pots and grown until the

inflorescences were fully formed. Plants were irrigated with tap water supplemented

with fertilizer (NPK 17-4-25; Kekkilä)

For the analysis of FvFT1 and FvTFL1 functions we pre-grew 8 – 13 seed

germinated plants per transgenic line or cultivar, or 6 – 13 clonally propagated plants

per accession with functional FvTFL1, in growth rooms under an 18h/6h (day/night)

photoperiod (AP67, Valoya, Finland) and 25^{o}C temperature. For floral induction, the

plants were then moved to a 12h/12h (day/night) photoperiod at 18^{o}C in the

greenhouse for six weeks. After the treatment, the plants were kept under 18^{o}C and

an 18h/6h (day/night) photoperiod until the inflorescences were fully developed.

Inflorescence architecture was analyzed by counting the total number of flowers and

flower buds, the number of branching iterations in the longest branching path, and

the number of branches on the primary axis. For the functional analysis of FvTFL1,

the first fully developed inflorescence was analyzed from each plant. For the

functional analysis of FvFT1, five fully developed inflorescences were analyzed.

Genotyping

621 DNA was extracted from young, folded leaves of *fvtfl1* mutants and WT plants listed

622 in Supplemental Table S2 as in Koskela et al. (2012). PCR amplified *FvTFL1*

623 fragments were excised from agarose gel, purified, and sequenced using primers

624 listed in Supplemental Table S3.

RNA extraction and qPCR analysis

Meristems were dissected under stereomicroscope (Zeiss Stemi 2000). SAMs, IMs,

and FMs were excised and collected into separate Eppendorf tubes. 5 – 10

meristems were pooled into one tube and used for RNA extraction. Total RNA was

extracted from pooled meristem samples as in Mouhu et al. (2009) and treated with

rDNase (Macherey-Nagel GmbH, Düren, Germany). 500ng of total RNA was used

for cDNA synthesis using ProtoScriptII reverse transcriptase. RT-PCR was

performed using a LightCycler 480 SYBR Green I Master kit (Roche Diagnostics,

Indianapolis, US) and a Roche LightCycler (Roche Diagnostics, Indianapolis, US)

with three technical replicates for each of the tested genes. *FvMSI1 (FvH4_7g08380)*

was used as a reference gene. The qPCR primers used in this study are listed in

Supplemental Table S3.

SEM imaging

Dissected meristem samples were fixed in an FAA buffer (3.7 % formaldehyde 5 %

acetic acid, and 50 % ethanol) overnight and transferred through ethanol series (50

%, 60%, 70 %, 80 %, 90 %, 100%, 100 %) under mild vacuum (~0.6 atm). Critical

point drying was done using a Leica EM CPD300 (Leica Mikrosystems GmbH,

Vienna, Austria). The dried samples were then coated with platinum by a Quorum

Q150TS coater (Quorum Technologies, UK). The samples were examined under a

Quanta 250 FEG (FEI, Hillsboro, OR, USA) scanning electron microscope located at

the Electron Microscopy Unit (Institute of Biotechnology, University of Helsinki).
Pseudo-coloring was done in Adobe Photoshop CC 2019.

Statistical analyses

The phenotypic data (flowering time, number of flowers per inflorescence and
branching iterations) were analyzed using random intercept models, $y = \beta_0 + \beta_1 X +$
$Gt + \varepsilon$, where y = dependent variable, βs = fixed effects, X = design matrices for
FvTFL1 background or *FvFT1* transgenic construct, Gt = random accession or
independent line effect, and ε = an error term. Poisson distribution model was
applied for count data. Statistical analyses were performed using the R/lme4
package (Bates et al., 2015). All analyses were performed in R version 4.1.1 (R Core
Team 2021).

Inflorescence modelling

The models were written in the L-system-based L+C plant modeling language
(Karwowski and Prusinkiewicz, 2003) and executed using the lpfg simulator
incorporated into the Virtual Laboratory (vlab) v5.0 plant modeling software
(algorithmicbotany.org/virtual_laboratory). All simulations were performed on
MacBook Pro computers under macOS High Sierra 10.13.6. The supplemental
materials include three complete vlab objects (versions of the model): the basic
version with a simplified representation of plant organs (bracts and flowers),
convenient for analyzing the model logic; the extended version with full
representation of organs, used in the model calibration shown in Figure 7; and the
same extended version configured to simulate the diverse phenotypes shown in
Figure 8. The key elements of the model code are discussed in the Supplemental
Model Description. Parameters used to simulate the diverse phenotypes were found
by interactively exploring the model parameter space using the tools included in vlab
(control panels and graphically defined timelines and functions). The reference
model (Figure 7) was calibrated using the method described by Cieslak et al. (2022).

Acknowledgements

We thank Elli Koskela for providing seeds; Eija Takala and Marjo Kilpinen for
laboratory assistance; Katriina Palm for maintaining the plant materials; the staff of
the Electron Microscopy Unit at the Institute of Biotechnology, University of Helsinki,

for assistance. This work was supported by Jane and Aatos Erkko Foundation grant
4709012 (to P.E. and T.H.), the Academy of Finland grant 317306 (to T.H), Natural
Sciences and Engineering Research Council of Canada Discovery Grant 2019-
06279 (to P.P.), the Plant Phenotyping and Imaging Research Centre/Canada First
Research Excellence Fund (to P.P. and M.C.), and the Finnish Cultural Foundation
grant 00221168 (to S.L.). S.L. was affiliated with the Doctoral Programme in Plant
Sciences (DPPS, University of Helsinki).

REFERENCES

Alvarez J, Guli CL, Yu X-H, Smyth DR (1992) *terminal flower*: a gene affecting

inflorescence development in *Arabidopsis thaliana*. Plant J **2:** 103–116

Anderson HM, Guttridge CG (1982) Strawberry truss morphology and the fate of high-

order flower buds. Crop Research **22:** 105–122

Ashman TL, Hitchens MS (2000) Dissecting the causes of variation in intra-

inflorescence allocation in a sexually polymorphic species, *Fragaria virginiana*

(Rosaceae). Am J Bot **87:** 197–204

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting linear mixed-effects models

using lme4. J Stat Softw **67:** 1–48

Cieslak M, Khan N, Ferraro P, Soolanayakanahally R, Robinson SJ, Parkin I,

McQuillan I, Prusinkiewicz P (2022) L-system models for image-based phenomics:

case studies of maize and canola. isP **4:** diab039

Claßen-Bockhoff R, Bull-Hereñu K (2013) Towards an ontogenetic understanding of

inflorescence diversity. Ann Bot **112:** 1523–1542

Corbesier L, Vincent C, Jang S, Fornara F, Fan Q, Searle I, Giakountis A, Farrona S,

Gissot L, Turnbull C, Coupland G (2007) FT protein movement contributes to long-

distance signaling in floral induction of *Arabidopsis*. Science **316:** 1030–1033

Darrow GM (1929) Inflorescence types of strawberry varieties. Am J Bot **16:** 571–585

Edger PP, VanBuren R, Colle M, Poorten TJ, Wai CM, Niederhuth CE, Alger EI, Ou

S, Acharya CB, Wang J, Callow P, McKain MR, Shi J, Collier C, Xiong Z, Mower

JP, Slovin JP, Hytönen T, Jiang N, Childs KL, Knapp SJ (2018) Single-molecule

sequencing and optical mapping yields an improved genome of woodland strawberry

(*Fragaria vesca*) with chromosome-scale contiguity. GigaScience **7:** gix124

Eshed Y, Lippman ZB (2019) Revolutions in agriculture chart a course for targeted

breeding of old and new crops. Science **366:** eaax0025

Foster JC, Janick J (1969) Variable branching patterns in the strawberry inflorescence.

J Am Soc Hortic Sci **94:** 440–443

Gleason HA, Cronquist A (1991) *Manual of Vascular Plants of Northeastern United*
States and Adjacent Canada (2nd ed.). New York Botanical Garden

Grimes J (1999) Inflorescence morphology, heterochrony, and phytogeny in the
mimosoid tribes ingeae and acacieae (Leguminosae: Mimosoideae). Bot Rev **65:**
317–347

Guttridge CG (1985) *Fragaria x ananassa*. *In* AH Halevy ed, Handbook of Flowering, v*ol.*
3. CRC Press, Boca Raton, FL, pp 16–33

Harder LD, Jordan CY, Gross WE, Routley MB (2004) Beyond floricentrism: the
pollination function of inflorescences. Plant Species Biol **19:** 137–148

Harder LD, Prusinkiewicz P (2013) The interplay between inflorescence development
and function as the crucible of architectural diversity. Ann Bot **112:** 1477–1493

Hofmeister W (1868) Allgemeine Morphologie der Gewächse. *In* A de Bary, TH Irmisch,
J Sachs, eds. Handbuch der Physiologischen Botanik. Engelmann, Leipzig, pp 405 –
664

Jahn OL, Dana MN (1970) Crown and inflorescence development in the strawberry,
Fragaria ananassa. Am J Bot **57:** 605–612

Kaneko-Suzuki M, Kurihara-Ishikawa R, Okushita-Terakawa C, Kojima C, Nagano-
Fujiwara M, Ohki I, Tsuji H, Shimamoto K, Taoka KI (2018) TFL1-like proteins in
rice antagonize rice FT-like protein in inflorescence development by competition for
complex formation with 14-3-3 and FD. Plant Cell Physiol **59:** 458–468

Karwowski R, Prusinkiewicz P (2003) Design and implementation of the L+C modeling
language. Electron Notes Theor Comput Sci **86:** 134–152

Kellogg EA (2000) A model of inflorescence development. *In* KL Wilson and DA
Morrison eds, Monocots: Systematics and Evolution. CSIRO, Melbourne, pp 84–88.

Kobayashi K, Yasuno N, Sato Y, Yoda M, Yamazaki R, Kimizu M, Yoshida H,
Nagamura Y, Kyozuka J (2012) Inflorescence meristem identity in rice is specified
by overlapping functions of three *AP1/FUL*-Like MADS box genes and *PAP2*, a
SEPALLATA MADS box gene. Plant Cell **24:** 1848–1859

Koskela EA, Mouhu K, Albani MC, Kurokura T, Rantanen M, Sargent DJ, Battey NH,
Coupland G, Elomaa P, Hytönen T (2012) Mutation in *TERMINAL FLOWER1*
reverses the photoperiodic requirement for flowering in the wild strawberry *Fragaria*
vesca. Plant Physiol **159:** 1043–1054

Koskela EA, Kurokura T, Toivainen T, Sønsteby A, Heide OM, Sargent DJ, Isobe S,
Jaakola L, Hilmarsson H, Elomaa P, Hytönen T (2017) Altered regulation of
TERMINAL FLOWER 1 causes the unique vernalisation response in an arctic
woodland strawberry accession. New Phytol **216:** 841–853

Kurokura T, Samad S, Koskela E, Mouhu K, Hytönen T (2017) *Fragaria vesca*
CONSTANS controls photoperiodic flowering and vegetative development. J Exp Bot
68: 4839–4850

Lee C, Kim S-J, Jin S, Susila H, Youn G, Nasim Z, Alavilli H, Chung K-S, Yoo SJ,
Ahn JH (2019) Genetic interactions reveal the antagonistic roles of *FT/TSF* and
TFL1 in the determination of inflorescence meristem identity in Arabidopsis. Plant J
99: 452–464

Lemmon ZH, Park SJ, Jiang K, Van Eck J, Schatz MC, Lippman ZB (2016) The
evolution of inflorescence diversity in the nightshades and heterochrony during
meristem maturation. Genome Res **26:** 1676–1686

Lifschitz E, Eviatar T, Rozman A, Shalit A, Goldshmidt A, Amsellem Z, Alvarez JP,
Eshed Y (2006) The tomato *FT* ortholog triggers systemic signals that regulate
growth and flowering and substitute for diverse environmental stimuli. Proc Natl Acad
Sci USA **103:** 6398–6403

Liu C, Teo ZWN, Bi Y, Song S, Xi W, Yang X, Yin Z, Yu H (2013) A Conserved Genetic
Pathway Determines Inflorescence Architecture in Arabidopsis and Rice. Dev Cell
24: 612–622

Lück J, Lück HB, Bakkali M (1990) A comprehensive model for acrotonic, mesotonic
and basitonic branchings in plants. Acta Biotheor **38:** 257–288

Ma Q, Liu X, Franks RG, Xiang Q-Y (2017) Alterations of *CorTFL1* and *CorAP1*
expression correlate with major evolutionary shifts of inflorescence architecture in

Cornus (Cornaceae) – a proposed model for variation of closed inflorescence forms. New Phytol **216:** 519–535

McGarry RC, Ayre BG (2012) Manipulation plant architecture with members of the CETS gene family. Plant Sci **188–189:** 71–81

Menzel CM (2019) Temperature has a greater effect on fruit growth than defoliation or fruit thinning in strawberries in the subtropics. Agriculture **9***:* 127

Molinero-Rosales N, Jamilena M, Zurita S, Gómez P, Capel J, Lozano R (1999) *FALSIFLORA*, the tomato orthologue of *FLORICAULA* and *LEAFY*, controls flowering time and floral meristem identity. Plant J **20:** 685–693

Molinero-Rosales N, Latorre A, Jamilena M, Lozano R (2004) *SINGLE FLOWER TRUSS* regulates the transition and maintenance of flowering in tomato. Planta **218:** 427–434

Mouhu K, Hytönen T, Folta K, Rantanen M, Paulin L, Auvinen P, Elomaa P (2009) Identification of flowering genes in strawberry, a perennial SD plant. BMC Plant Biol **9:** 122

Nakano Y, Higuchi Y, Yoshida Y, Hisamatsu T (2015) Environmental responses of the FT/TFL1 gene family and their involvement in flower induction in *Fragaria × ananassa*. J Plant Physiol **177:** 60–66

Park SJ, Jiang K, Schatz MC, Lippman ZB (2012) Rate of meristem maturation determines inflorescence architecture in tomato. Proc Natl Acad Sci USA **109:** 639–644

Pnueli L, Carmel-Goren L, Hareven D, Gutfinger T, Alvarez J, Ganal M, Zamir D, Lifschitz E (1998) The *SELF-PRUNING* gene of tomato regulates vegetative to reproductive switching of sympodial meristems and is the ortholog of *CEN* and *TFL1*. Development **125:** 1979–1989

Peterson JL (1981) *Petri Net Theory and the Modeling of Systems*. Prentice-Hall, Englewood Cliffs

Prenner G, Vergara-Silva F, Rudall PJ (2009) The key role of morphology in modelling inflorescence architecture. Trends Plant Sci **14:** 302–309

Prusinkiewicz P, Erasmus Y, Lane B, Harder LD, Coen E (2007). Evolution and

development of inflorescence architectures. Science **316:** 1452–1456

Prusinkiewicz P, Lindenmayer A (1990) *The Algorithmic Beauty of Plants*. With JS

Hanan, FD Fracchia, D Fowler, MJM. de Boer and L Mercer. Springer, New York

Prusinkiewicz P, Remphrey W (2000) Characterization of architectural tree models

using L-systems and Petri nets. In M. Labrecque, editor, *L'arbre — The Tree 2000:*

Papers presented at the 4th International Symposium on the Tree, pp 177–186.

Isabelle Quentin, Montreal

R Core Team (2021) R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Rantanen M, Kurokura T, Mouhu K, Pinho P, Tetri E, Halonen L, Palonen P, Elomaa

P, Hytönen T (2014) Light quality regulates flowering in FvFT1/FvTFL1 dependent

manner in the woodland strawberry *Fragaria vesca*. Front Plant Sci **5:** 271

Rantanen M, Kurokura T, Jiang P, Mouhu K, Hytönen T (2015) Strawberry homologue

of TERMINAL FLOWER1 integrates photoperiod and temperature signals to inhibit

flowering. Plant J **82:** 163 – 173

Quinet M, Dubois C, Goffin M-C, Chao J, Dielen V, Batoko H, Boutry M, Kinet J-M

(2006) Characterization of tomato (*Solanum lycopersicum* L.) mutants affected in

their flowering time and in the morphogenesis of their reproductive structure. J Exp

Bot **57:** 1381–1390

Schultz EA, Haughn GW (1993) Genetic analysis of the floral initiation process (FLIP) in

Arabidopsis. Development **119:** 745–765

Souer E, Rebocho AB, Bliek M, Kusters E, de Bruin RAM, Koes R (2008) Patterning

of inflorescences and flowers by the F-Box protein DOUBLE TOP and the LEAFY

homolog ABERRANT LEAF AND FLOWER of Petunia. Plant Cell **20:** 2033–2048

Thouet J, Quinet M, Lutts S, Kinet J-M, Périlleux C (2012) Repression of floral

meristem fate is crucial in shaping tomato inflorescence. PloS One **7:** e31096–

e31096

Weberling F (1989) *Morphology of flowers and inflorescences.* Cambridge University

Press, Cambridge

Weigel D, Alvarez J, Smyth DR, Yanofsky MF, Meyerowitz EM (1992) LEAFY controls
floral meristem identity in Arabidopsis. Cell **69:** 843–859

Weigel D, Meyerowitz ME (1993) Activation of floral homeotic genes in Arabidopsis.
Science, **261:** 1723–1726

Zhang Y, Kan L, Hu S, Cheng L, Liu Z, Kang C (2022) The roles and evolution of the
four LEAFY homologues in floral patterning and leaf development in woodland
strawberry. BioRxiv, https://doi.org/10.1101/2022.10.06.511044

Zhu Y, Klasfeld S, Jeong CW, Jin R, Goto K, Yamaguchi N, Wagner D (2020)
TERMINAL FLOWER 1-FD complex target genes and competition with
FLOWERING LOCUS T. Nat Commun **11:** 5118

Zhu Y, Klasfeld S, Wagner D (2021) Molecular regulation of plant developmental
transitions and plant architecture via PEPB family proteins: an update on mechanism
of action. J Exp Bot **72:** 2301–2311

Parsed Citations

Alvarez J, Guli CL, Yu X-H, Smyth DR (1992) terminal flower: a gene affecting inflorescence development in Arabidopsis thaliana. Plant J 2: 103–116
Google Scholar: Author Only Title Only Author and Title

Anderson HM, Guttridge CG (1982) Strawberry truss morphology and the fate of high-order flower buds. Crop Research 22: 105–122
Google Scholar: Author Only Title Only Author and Title

Ashman TL, Hitchens MS (2000) Dissecting the causes of variation in intra-inflorescence allocation in a sexually polymorphic species, Fragaria virginiana (Rosaceae). Am J Bot 87: 197–204
Google Scholar: Author Only Title Only Author and Title

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting linear mixed-effects models using lme4. J Stat Softw 67: 1–48
Google Scholar: Author Only Title Only Author and Title

Cieslak M, Khan N, Ferraro P, Soolanayakanahally R, Robinson SJ, Parkin I, McQuillan I, Prusinkiewicz P (2022) L-system models for image-based phenomics: case studies of maize and canola. isP 4: diab039
Google Scholar: Author Only Title Only Author and Title

Claßen-Bockhoff R, Bull-Hereñu K (2013) Towards an ontogenetic understanding of inflorescence diversity. Ann Bot 112: 1523–1542
Google Scholar: Author Only Title Only Author and Title

Corbesier L, Vincent C, Jang S, Fornara F, Fan Q, Searle I, Giakountis A, Farrona S, Gissot L, Turnbull C, Coupland G (2007) FT protein movement contributes to long-distance signaling in floral induction of Arabidopsis. Science 316: 1030–1033
Google Scholar: Author Only Title Only Author and Title

Darrow GM (1929) Inflorescence types of strawberry varieties. Am J Bot 16: 571–585
Google Scholar: Author Only Title Only Author and Title

Edger PP, VanBuren R, Colle M, Poorten TJ, Wai CM, Niederhuth CE, Alger EI, Ou S, Acharya CB, Wang J, Callow P, McKain MR, Shi J, Collier C, Xiong Z, Mower JP, Slovin JP, Hytönen T, Jiang N, Childs KL, Knapp SJ (2018) Single-molecule sequencing and optical mapping yields an improved genome of woodland strawberry (Fragaria vesca) with chromosome-scale contiguity. GigaScience 7: gix124
Google Scholar: Author Only Title Only Author and Title

Eshed Y, Lippman ZB (2019) Revolutions in agriculture chart a course for targeted breeding of old and new crops. Science 366: eaax0025
Google Scholar: Author Only Title Only Author and Title

Foster JC, Janick J (1969) Variable branching patterns in the strawberry inflorescence. J Am Soc Hortic Sci 94: 440–443
Google Scholar: Author Only Title Only Author and Title

Gleason HA, Cronquist A (1991) Manual of Vascular Plants of Northeastern United States and Adjacent Canada (2nd ed.). New York Botanical Garden
Google Scholar: Author Only Title Only Author and Title

Grimes J (1999) Inflorescence morphology, heterochrony, and phytogeny in the mimosoid tribes ingeae and acacieae (Leguminosae: Mimosoideae). Bot Rev 65: 317–347
Google Scholar: Author Only Title Only Author and Title

Guttridge CG (1985) Fragaria x ananassa. In AH Halevy ed, Handbook of Flowering, vol. 3. CRC Press, Boca Raton, FL, pp 16–33
Google Scholar: Author Only Title Only Author and Title

Harder LD, Jordan CY, Gross WE, Routley MB (2004) Beyond floricentrism: the pollination function of inflorescences. Plant Species Biol 19: 137–148
Google Scholar: Author Only Title Only Author and Title

Harder LD, Prusinkiewicz P (2013) The interplay between inflorescence development and function as the crucible of architectural diversity. Ann Bot 112: 1477–1493
Google Scholar: Author Only Title Only Author and Title

Hofmeister W (1868) Allgemeine Morphologie der Gewächse. In A de Bary, TH Irmisch, J Sachs, eds. Handbuch der Physiologischen Botanik. Engelmann, Leipzig, pp 405 – 664
Google Scholar: Author Only Title Only Author and Title

Jahn OL, Dana MN (1970) Crown and inflorescence development in the strawberry, Fragaria ananassa. Am J Bot 57: 605–612
Google Scholar: Author Only Title Only Author and Title

Kaneko-Suzuki M, Kurihara-Ishikawa R, Okushita-Terakawa C, Kojima C, Nagano-Fujiwara M, Ohki I, Tsuji H, Shimamoto K, Taoka KI (2018) TFL1-like proteins in rice antagonize rice FT-like protein in inflorescence development by competition for complex formation with 14-3-3 and FD. Plant Cell Physiol 59: 458–468
  Google Scholar: Author Only Title Only Author and Title

Karwowski R, Prusinkiewicz P (2003) Design and implementation of the L+C modeling language. Electron Notes Theor Comput Sci 86: 134–152
  Google Scholar: Author Only Title Only Author and Title

Kellogg EA (2000) A model of inflorescence development. In KL Wilson and DA Morrison eds, Monocots: Systematics and Evolution. CSIRO, Melbourne, pp 84–88.
  Google Scholar: Author Only Title Only Author and Title

Kobayashi K, Yasuno N, Sato Y, Yoda M, Yamazaki R, Kimizu M, Yoshida H, Nagamura Y, Kyozuka J (2012) Inflorescence meristem identity in rice is specified by overlapping functions of three AP1/FUL-Like MADS box genes and PAP2, a SEPALLATA MADS box gene. Plant Cell 24: 1848–1859
  Google Scholar: Author Only Title Only Author and Title

Koskela EA, Mouhu K, Albani MC, Kurokura T, Rantanen M, Sargent DJ, Battey NH, Coupland G, Elomaa P, Hytönen T (2012) Mutation in TERMINAL FLOWER1 reverses the photoperiodic requirement for flowering in the wild strawberry Fragaria vesca. Plant Physiol 159: 1043–1054
  Google Scholar: Author Only Title Only Author and Title

Koskela EA, Kurokura T, Toivainen T, Sønsteby A, Heide OM, Sargent DJ, Isobe S, Jaakola L, Hilmarsson H, Elomaa P, Hytönen T (2017) Altered regulation of TERMINAL FLOWER 1 causes the unique vernalisation response in an arctic woodland strawberry accession. New Phytol 216: 841–853
  Google Scholar: Author Only Title Only Author and Title

Kurokura T, Samad S, Koskela E, Mouhu K, Hytönen T (2017) Fragaria vesca CONSTANS controls photoperiodic flowering and vegetative development. J Exp Bot 68: 4839–4850
  Google Scholar: Author Only Title Only Author and Title

Lee C, Kim S-J, Jin S, Susila H, Youn G, Nasim Z, Alavilli H, Chung K-S, Yoo SJ, Ahn JH (2019) Genetic interactions reveal the antagonistic roles of FT/TSF and TFL1 in the determination of inflorescence meristem identity in Arabidopsis. Plant J 99: 452–464
  Google Scholar: Author Only Title Only Author and Title

Lemmon ZH, Park SJ, Jiang K, Van Eck J, Schatz MC, Lippman ZB (2016) The evolution of inflorescence diversity in the nightshades and heterochrony during meristem maturation. Genome Res 26: 1676–1686
  Google Scholar: Author Only Title Only Author and Title

Lifschitz E, Eviatar T, Rozman A, Shalit A, Goldshmidt A, Amsellem Z, Alvarez JP, Eshed Y (2006) The tomato FT ortholog triggers systemic signals that regulate growth and flowering and substitute for diverse environmental stimuli. Proc Natl Acad Sci USA 103: 6398–6403
  Google Scholar: Author Only Title Only Author and Title

Liu C, Teo ZWN, Bi Y, Song S, Xi W, Yang X, Yin Z, Yu H (2013) A Conserved Genetic Pathway Determines Inflorescence Architecture in Arabidopsis and Rice. Dev Cell 24: 612–622
  Google Scholar: Author Only Title Only Author and Title

Lück J, Lück HB, Bakkali M (1990) A comprehensive model for acrotonic, mesotonic and basitonic branchings in plants. Acta Biotheor 38: 257–288
  Google Scholar: Author Only Title Only Author and Title

Ma Q, Liu X, Franks RG, Xiang Q-Y (2017) Alterations of CorTFL1 and CorAP1 expression correlate with major evolutionary shifts of inflorescence architecture in Cornus (Cornaceae) – a proposed model for variation of closed inflorescence forms. New Phytol 216: 519–535
  Google Scholar: Author Only Title Only Author and Title

McGarry RC, Ayre BG (2012) Manipulation plant architecture with members of the CETS gene family. Plant Sci 188–189: 71–81
  Google Scholar: Author Only Title Only Author and Title

Menzel CM (2019) Temperature has a greater effect on fruit growth than defoliation or fruit thinning in strawberries in the subtropics. Agriculture 9: 127
  Google Scholar: Author Only Title Only Author and Title

Molinero-Rosales N, Jamilena M, Zurita S, Gómez P, Capel J, Lozano R (1999) FALSIFLORA, the tomato orthologue of FLORICAULA and LEAFY, controls flowering time and floral meristem identity. Plant J 20: 685–693
  Google Scholar: Author Only Title Only Author and Title

Molinero-Rosales N, Latorre A, Jamilena M, Lozano R (2004) SINGLE FLOWER TRUSS regulates the transition and maintenance

of flowering in tomato. Planta 218: 427–434
Google Scholar: Author Only Title Only Author and Title

Mouhu K, Hytönen T, Folta K, Rantanen M, Paulin L, Auvinen P, Elomaa P (2009) Identification of flowering genes in strawberry, a perennial SD plant. BMC Plant Biol 9: 122
Google Scholar: Author Only Title Only Author and Title

Nakano Y, Higuchi Y, Yoshida Y, Hisamatsu T (2015) Environmental responses of the FT/TFL1 gene family and their involvement in flower induction in Fragaria × ananassa. J Plant Physiol 177: 60–66
Google Scholar: Author Only Title Only Author and Title

Park SJ, Jiang K, Schatz MC, Lippman ZB (2012) Rate of meristem maturation determines inflorescence architecture in tomato. Proc Natl Acad Sci USA 109: 639–644
Google Scholar: Author Only Title Only Author and Title

Pnueli L, Carmel-Goren L, Hareven D, Gutfinger T, Alvarez J, Ganal M, Zamir D, Lifschitz E (1998) The SELF-PRUNING gene of tomato regulates vegetative to reproductive switching of sympodial meristems and is the ortholog of CEN and TFL1. Development 125: 1979–1989
Google Scholar: Author Only Title Only Author and Title

Peterson JL (1981) Petri Net Theory and the Modeling of Systems. Prentice-Hall, Englewood Cliffs
Google Scholar: Author Only Title Only Author and Title

Prenner G, Vergara-Silva F, Rudall PJ (2009) The key role of morphology in modelling inflorescence architecture. Trends Plant Sci 14: 302–309
Google Scholar: Author Only Title Only Author and Title

Prusinkiewicz P, Erasmus Y, Lane B, Harder LD, Coen E (2007). Evolution and development of inflorescence architectures. Science 316: 1452–1456
Google Scholar: Author Only Title Only Author and Title

Prusinkiewicz P, Lindenmayer A (1990) The Algorithmic Beauty of Plants. With JS Hanan, FD Fracchia, D Fowler, MJM. de Boer and L Mercer. Springer, New York
Google Scholar: Author Only Title Only Author and Title

Prusinkiewicz P, Remphrey W (2000) Characterization of architectural tree models using L-systems and Petri nets. In M. Labrecque, editor, L'arbre - The Tree 2000: Papers presented at the 4th International Symposium on the Tree, pp 177–186. Isabelle Quentin, Montreal
Google Scholar: Author Only Title Only Author and Title

R Core Team (2021) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/
Google Scholar: Author Only Title Only Author and Title

Rantanen M, Kurokura T, Mouhu K, Pinho P, Tetri E, Halonen L, Palonen P, Elomaa P, Hytönen T (2014) Light quality regulates flowering in FvFT1/FvTFL1 dependent manner in the woodland strawberry Fragaria vesca. Front Plant Sci 5: 271
Google Scholar: Author Only Title Only Author and Title

Rantanen M, Kurokura T, Jiang P, Mouhu K, Hytönen T (2015) Strawberry homologue of TERMINAL FLOWER1 integrates photoperiod and temperature signals to inhibit flowering. Plant J 82: 163 – 173
Google Scholar: Author Only Title Only Author and Title

Quinet M, Dubois C, Goffin M-C, Chao J, Dielen V, Batoko H, Boutry M, Kinet J-M (2006) Characterization of tomato (Solanum lycopersicum L.) mutants affected in their flowering time and in the morphogenesis of their reproductive structure. J Exp Bot 57: 1381–1390
Google Scholar: Author Only Title Only Author and Title

Schultz EA, Haughn GW (1993) Genetic analysis of the floral initiation process (FLIP) in Arabidopsis. Development 119: 745–765
Google Scholar: Author Only Title Only Author and Title

Souer E, Rebocho AB, Bliek M, Kusters E, de Bruin RAM, Koes R (2008) Patterning of inflorescences and flowers by the F-Box protein DOUBLE TOP and the LEAFY homolog ABERRANT LEAF AND FLOWER of Petunia. Plant Cell 20: 2033–2048
Google Scholar: Author Only Title Only Author and Title

Thouet J, Quinet M, Lutts S, Kinet J-M, Périlleux C (2012) Repression of floral meristem fate is crucial in shaping tomato inflorescence. PloS One 7: e31096–e31096
Google Scholar: Author Only Title Only Author and Title

Weberling F (1989) Morphology of flowers and inflorescences. Cambridge University Press, Cambridge
Google Scholar: Author Only Title Only Author and Title

Weigel D, Alvarez J, Smyth DR, Yanofsky MF, Meyerowitz EM (1992) LEAFY controls floral meristem identity in Arabidopsis. Cell 69: 843–859

Google Scholar: Author Only Title Only Author and Title

Weigel D, Meyerowitz ME (1993) Activation of floral homeotic genes in Arabidopsis. Science, 261: 1723–1726

Google Scholar: Author Only Title Only Author and Title

Zhang Y, Kan L, Hu S, Cheng L, Liu Z, Kang C (2022) The roles and evolution of the four LEAFY homologues in floral patterning and leaf development in woodland strawberry. BioRxiv, https://doi.org/10.1101/2022.10.06.511044

Google Scholar: Author Only Title Only Author and Title

Zhu Y, Klasfeld S, Jeong CW, Jin R, Goto K, Yamaguchi N, Wagner D (2020) TERMINAL FLOWER 1-FD complex target genes and competition with FLOWERING LOCUS T. Nat Commun 11: 5118

Google Scholar: Author Only Title Only Author and Title

Zhu Y, Klasfeld S, Wagner D (2021) Molecular regulation of plant developmental transitions and plant architecture via PEPB family proteins: an update on mechanism of action. J Exp Bot 72: 2301–2311

Google Scholar: Author Only Title Only Author and Title