1

Allele surfing causes maladaptation in a Pacific salmon of conservation concern

Quentin Rougemont[1*], Thibault Leroy[2], Eric B. Rondeau[3], Ben Koop[4], Louis Bernatchez[5]

[1.] Centre d'Ecologie Fonctionnelle et Evolutive, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France
[2.]GenPhySE, INRAE, INP, ENVT, Université de Toulouse, 24 chemin de Borde-Rouge, Auzeville-Tolosane, 31326 Castanet Tolosan, France
[3.] Department of Fisheries and Ocean, Pacific Biological Station, 3190 Hammond Bay Road, Nanaimo, British Columbia, Canada Or V9T 6N7
[4] Department of Biology, University of Victoria, Victoria, BC, Canada
[5.] Département de Biologie, Institut de Biologie Intégrative et des Systèmes (IBIS), Université Laval, Québec, Québec, Canada

* **Corresponding Author :** quentinrougemont@orange.fr

Abstract

How various factors, including demography, recombination or genome duplication, may impact the efficacy of natural selection and the burden of deleterious mutations,is a central question in evolutionary biology and genetics. In this study, we show that key evolutionary processes, including variations in *i)* effective population size (N_e) *ii)* recombination rates and *iii)* chromosome inheritance, have influenced the genetic load and efficacy of selection in Coho salmon (*Oncorhynchus kisutch*), a widely distributed salmonid species on the west coast of North America. Using whole genome resequencing data from 14 populations at different migratory distances from their southern glacial refugium, we found evidence supporting gene surfing, wherein reduced N_e at the postglacial recolonization front, leads to a decrease in the efficacy of selection and a surf of deleterious alleles in the northernmost populations. Furthermore, our results indicate that recombination rates play a prime role in shaping the load along the genome. Additionally, we identified variation in polyploidy as a contributing factor to within-genome variation of the load. Overall, our results align remarkably well with expectations under the nearly neutral theory of molecular evolution. We discuss the fundamental and applied implications of these findings for evolutionary and conservation genomics.

Author Summary

Understanding how historical processes, such as past glaciations, may have impacted variations in population size and genetic diversity along the genome is a fundamental question in evolution. In this study, we investigated how recent postglacial demographic expansion has affected the distribution of deleterious genetic variants and the resulting deleterious mutation load in Coho salmon (*Oncorhynchus kisutch*), throughout its native range in North America. By sequencing the entire genome of 71 Coho salmon, we reveal that postglacial expansion has led to allele surfing, a process where alleles increase in frequency in populations that are expanding or colonizing new environments. Here, allele surfing resulted in an increased

deleterious mutation load at the colonization front. Furthermore, we demonstrated that the efficacy of

natural selection scales with variation in effective population size among populations. We showed that the

specific genomic features of Coho salmon, namely variation in local recombination rate and variation in

chromosomal inheritance, strongly impacted the segregation of deleterious mutations.

Introduction

Demographic events, including population contractions, expansions and secondary contacts have profound

impacts on the spatial distribution of genetic diversity within species [1]. Range expansions can be

accompanied by multiple founder events when few individuals colonize new areas. At the wave front, this

process results in increased genetic drift and a reduced effective population size [2,3]. This has multiple

consequences genome-wide including (i) extended linkage disequilibrium [4], (ii) a loss of genetic diversity

and (iii) increased levels of genetic differentiation [2,3,5]. At the wavefront, increased genetic drift favors

allele surfing, a process that will increase the relative proportion of neutral, deleterious or advantageous

mutations that will fix in the population [5–7]. Such allele surfing is expected to have two main negative

consequences at the wave front: *i*) an increase of the deleterious load (called expansion load [8,9]) and *ii*) a

loss of adaptive potential [10].

First, allele surfing through increased genetic drift can overwhelm the effect of weak purifying

selection, resulting in an increase of the deleterious load [11]. Due to the lower efficacy of natural selection

in low N_e populations, a greater fraction of slightly deleterious mutations is expected to be present in

populations at the wavefront (*i.e.*, elevated ratios of non-synonymous to synonymous nucleotide diversity

(π_N/π_S) [8,9]). Moreover, under the nearly neutral theory of molecular evolution, the reduced efficacy of

purifying selection in small populations should increase the fixation rate of slightly deleterious mutations

[12]. On the other hand, given that most deleterious mutations are expected to be recessive, increased

homozygosity of deleterious mutations should enable their removal more efficiently through genetic

purging, leading to a reduction of the recessive and additive load [13]. To take this into account, a possible

strategy is to investigate both the additive (i.*e*,. total) and the recessive (*i.e.*, fixed) deleterious mutation

load. [8]. Evidence for an increased recessive genetic load has been reported in humans which has been

associated with the Out-of-Africa bottleneck [14,15]. Evidence has also been gathered in expanding

populations in plants [16–20]. For instance in *Arabidopsis lyrata*, range expansion has been associated with

an increased mutational load [16] and increased linkage disequilibrium [21]. Similar results were observed in

Escherichia coli[22]. In contrast, a recent study in *A. lyrata* found no evidence of increased load following

range expansion [23].

Second, allele surfing is expected to lead to a loss of adaptive potential due to a reduction in the rate of

adaptive substitutions given that the supply of new mutations is proportional to N_e *[24]*. Unless

compensatory mutations (*i.e.*, favourable mutations whose probability of occurrence increases to

compensate for the fixation of slightly detrimental mutations – themselves with a higher probability of

fixation in small populations–), counteract this effect in populations with a low N_e [25], one expects a lower

proportion of substitutions driven by positive selection [26]. Support for a decreased rate of adaptation

associated with range expansion was documented, particularly in plants [17,20]. For instance, such a pattern

was reported in the North American populations of *A. lyrata*[17] [27], but not in the European ones despite a

slight increase in their genomic burden [23]. Similarly an increased load is observed in human populations, a

result that matches theoretical expectations under the out-of-Africa scenario [28] but with limited evidence

of a decrease in the efficacy of natural selection [29]. Beyond application to plants, humans and bacteria,

there is a lack of empirical studies documenting the consequences of range expansion on the deleterious

mutation load and the efficacy of natural selection.

Geographic variation in the overall mutation load is not the only interesting pattern to investigate; the

genomic scale is also important. Both deleterious load and efficacy of selection are expected to vary along

the genome depending on the local recombination rate [30,31]. Non-recombining regions are expected to

more freely accumulate deleterious mutations, a process called Muller's Ratchet [32]. Moreover, Hill-

Robertson interference, a process whereby competing alleles interfere with each other to become fixed,

increases the fixation rate of deleterious variants linked to positively selected sites. Such a process will be

exacerbated in the absence of recombination [30,33].

In species that have undergone a whole-genome duplication (WGD) with associated residual polyploidy,

variation in chromosomal inheritance and genomic architecture may affect the distribution of deleterious

mutations along the genome [34]. By doubling the chromosome number, WGD has major consequences. In

the short term, WGD alters gene expression and favours polyploid dysfunction (reviewed in [35]). In the

middle to long-term, the progressive return to a diploid state will result in regions of residual tetraploidy.

Such regions should display an increased effective population size as compared to the rest of the genome (4

N_e instead of 2 N_e) and therefore a higher efficacy of selection and a reduced genetic load. Similarly, the

population scale recombination rate, which also depends on N_e, should favour a higher efficacy of selection

and a reduced genetic load. We thus predict that these regions should display a lower genetic load

compared to diploid regions of the genome. Finally, theoretical work has shown that dominance of

mutation, especially in the recessive case, is an important factor influencing the detection of hard sweeps in

polyploid genomes [36]. Dominance may therefore also influence the segregation of deleterious variants in

regions of residual tetraploidy. WGD has been associated with a reduced efficacy of purifying selection which

may favour the accumulation of transposable elements. Such WGD may in turn have contrasted effects, as

observed in plants [37,38]. For instance, a LTR insertion favouring early flowering time (an adaptive trait in

harsh environments), was shown to be present in tetraploid populations, but not in diploid populations of *A.*

arenosa. In contrast, WGS was shown to be associated with small but significant differences in the load and

distribution of fitness effect in the same species.

The Coho salmon (*Oncorhynchus kisutch*) is a Pacific salmon species of major cultural and economical

importance, which has severely declined over the last decades (reviewed in [39]). It has undergone further

recent declines in multiple parts of its native range [40]. In addition to human-induced perturbations, the

species has undergone a series of postglacial expansions from its main glacial refugium (Cascadia/California)

with multiple founder events along its route toward Alaska, leading to a pattern of isolation by distance, a

gradient of population structure and a decrease in genetic diversity [41–43]. Accordingly, these range

expansions may result in an increased expansion load, questioning whether such load may have

consequences on the fitness of populations at the expansion front. The Coho salmon offers a unique

opportunity to test the role of demography and recombination on the deleterious load and on the efficacy of

selection. The Coho salmon belongs to the Salmonidae family, a particularly relevant group of species to

investigate the effect of WGD. The common ancestor of all present-day salmonids has undergone a whole-

genome duplication event ~80-100MyA [44]. The genome of salmonid species is still on its path to

rediploidization. For instance, approximately 8% of the Coho salmon genome displays residual tetraploidy

[41]. Residual tetraploidy offers a unique opportunity to investigate the direct effect of the genomic variation

of N_e on the efficacy of selection.

To address these questions, we aimed at testing the following general ypotheses: i) whether demographic

expansion and bottlenecks have led to an increased load and decreased efficacy of selection, ii) whether

heterogeneous recombination levels shape the within-genome variation of the load as a result of Hill-

Robertson effects and iii) whether residual tetraploidy results in an increased efficacy of selection and

reduced deleterious load through increased recombination and N_e.

Results and Discussion

Strong postglacial population expansion revealed through whole genome sequences

We generated 30X coverage whole genome resequencing data for 71 Coho salmon representing 14

populations distributed from California to some of the most upstream populations of the Porcupine River in

Yukon, Canada (**Fig 1A**). We also included several outgroup species namely the Sockeye salmon (*O. nerka*),

Chinook salmon (*O. tshawytscha*), Pink salmon *(O. Gorbuscha)*, Rainbow trout (*O. mykiss*) and Atlantic

salmon (*Salmo salar*). These genomes were used to identify the ancestral and derived alleles in Coho and/or

testing correlation(s) between genomic load estimates and recombination rate (**Table S1-S3**).

To explore the amount of clustering among individuals a PCA was performed (**Fig S1A, B**). The PCA illustrates

that populations are spatially structured following both latitudinal and longitudinal gradients. The first axis

strongly correlates with both latitude (Pearson r = 0.97, p < 2e^{-16}) and longitude (Pearson r = -0.82, p < 2e^{-16}),

as expected when space drives population structure [45]. Moreover, admixture analyses (**Fig S1B**) revealed that Coho populations were generally structured at the river/watershed level. These observations corroborate our previous inferences based on lower resolution genomic dataset [42,43].

Under a single refugial expansion scenario, we expect a linear decrease in genetic diversity as a function of the distance from the source. We plotted the distribution of observed heterozygosity (H_o) as a function of the distance to the southernmost sample and observed a negative correlation with the distance to the southernmost site ($R^2 = 0.77$, $p<2e^{-16}$, **Fig S2B**). Next, we used the β_{ST} coefficient to identify the most likely ancestral populations from which the expansion could have started. β_{ST} can account for the non-independence among populations and negative values are indicative of ancestral populations [46]. This metric was also positively correlated with distance to the southernmost site ($R^2 = 0.75$, $p < 2e^{-16}$, **Fig S2A**).

The Salmon River population from the Thompson area (dark green point, **Fig 1A** and **Fig S2A**) exhibits lower diversity levels, as compared to all non-Alaskan populations. Excluding this outlier river, revealed even stronger correlations, with a correlation with β_{ST} and H_O respectively raising to $R^2 = 0.88$ and $R^2 = 0.90$ ($p <2e^{-16}$ for both tests), suggesting an overall clear spatial pattern across the populations sampled, with the notable exception of this specific population which had a different demographic trajectory in this population, which seems consistent with a strong bottleneck and inbreeding [41].

Figure 1: Sampling design and population size change inferred through whole genome sequences. A) Sample site along the Pacific coast of North America. Map produced in R using the World data from Maps package with public domain data available at [47]. B) Average Tajima's D values computed along the genome and summarized for each sample locality. Populations are ordered from distance to the river mouth and from the South to the North (i) California, (ii) Cascadia, (iii) British-Columbia (BC), (iv) Haida-Gwaii, (v) Salmon River (Fraser Basin) and (vi) Alaska. Abbreviation provided in Table S2). Tajima's D values vary positively as a function of the distance from the southernmost site ($R^2 = 0.28$, $p <2e^{-16}$) and as a function of the distance to the river mouth ($R^2 = 0.47$, $p <2e^{-16}$) reflecting the history of founder events from downstream to upstream sites.

Another indicator of changes in population sizes is an increase in Tajima's D values at the genome-wide level. We tested the hypothesis of increased Tajima's D values as a function of (i) the traveled distance from the southernmost site (reflecting northward postglacial expansion) and (ii) the distance to the sea only

(reflecting upstream directed founder events). Results are consistent with a signal of population expansion

along the colonization axis from the south to the north based on Tajima's D values (R^2=0.28, p< $2e^{-16}$, **Fig 1B**).

Distance to the sea is also highly significant (R^2 = 0.47, p <$2e^{-16}$ for Tajima's D). This suggests that more

genetic drift occurs in upstream populations e.g. Porcupine (POR), Mile Slough (MSL), Thompson (SAL) and

Deschutes (DES).

Another hallmark of population bottlenecks and genetic drift associated with founder events is an increase in

linkage disequilibrium across the genome [4]. We tested this hypothesis by measuring linkage disequilibrium

(r^2) following Hill & Robertson [48]. In line with the above observations, we observed extended linkage

disequilibrium (LD) along the genome in remote Alaskan/Yukon populations (Porcupine and Mile Slough

Rivers) and in the Salmon R. population (Thompson R. Watershed, **Fig S2C**). Accordingly, the LD decay - here

defined as a r^2 reaching half of its maximum value - is observed at 25 , 12 and 6 kbp in these three

populations, contrasting with values around 0.5 kbp observed in the other populations(**Fig S2C**). These

results show a large among population variance in effective population sizes, likely associated with the

strength of the founder events.

In addition, we constructed a population phylogeny based on a matrix of pairwise F_{ST} values computed

between each pair of populations(**Fig 2A-B**) and on a matrix of variance-covariance in allele frequencies

implemented in Treemix [49] (**Fig S3**). Both analyses confirmed that populations located in the north

underwent more genetic drift, in line with the postglacial expansion from the south to the north and the

subsequent founder effects associated with upstream colonisation of the rivers from the west to the east.

To consolidate the above observations, we investigated historical changes in N_e using the Sequentially

Markovian Coalescent in SMC++ [50]. This method takes advantage of both the SMC and of the site

frequency spectrum to infer change of effective population size though time. We inferred such changed for

each population separately (**Fig S2D)**, testing a mutation rate of $8e^{-9}$ and $1.25e^{-8}$ mutations/bp/generation,

which respectively corresponds to the median and the mean mutation rate estimated in Atlantic salmon

(*salmo* salar), a closely related salmonid (J. Wang, personal communication). This analysis revealed: *i)* an

expansion of most populations approximately 12-20 KyA ago, concomitant with postglacial recolonization, *ii)*

a slow and steady demographic decline in the Thompson R. (**Fig S2D**), and *iii)* a split time between all

pairwise combinations of populations (median = 16,3 KyA, range = 6,7KyA - 61KyA, **Fig S4**) compatible with

the onset of postglacial population expansion and colonisation of different rivers following glacial retreat

[41,42]. Using the mean mutation rate yielded similar results with a more recent estimates of split times

(median = 9,6 KyA, **Fig S4**) (min = 5 KyA – max = 18 KyA). Overall, SMC++ results indicate that all populations

shared a similar demographic history until they began to diverge following the end of the last glacial cycle

(see **Note S1** and associated Supplementary Tables).

Range expansion explained variation in the mutation load

We estimated the mutation load of the populations using different metrics : *i)* π_N/π_S, *ii)* count of derived

homozygous mutations (including both missense and Loss-of-function (LoF) mutations), and *iii)* total count of

derived mutations (including both missense and LoF mutations). We observed a significant positive

correlation of the π_N/π_S ratio of each local population and the distance to the southernmost site (see

methods), corresponding to the most-likely refugia as discussed in our previous work [42] (R^2 = 0.73,

p<0.0001, **Fig 2D**). Similarly, using the tree branch length to the root of our tree (**Fig 2B**) as a proxy of

expansion route revealed a significant correlation with the π_N/π_S ratio (R^2 = 0.66, p<0.0001, F**ig 2C**). This

result was also supported by an analysis considering tree branch lengths inferred with Treemix (**Table S4**).

Accordingly, there was a significant correlation between π_N/π_S and two different proxies of the effective

population size, namely levels of nucleotide diversity at synonymous sites (π_S) and SMC++ N_e, expected to be

reduced at the colonization front and in bottlenecked populations (π_S in **Fig S5A** and SMC++ N_e in **Fig S5B**).

These results support expectations pertaining to variation in mutation load under a model of expansion load

(e.g. [28]).

Figure 2: Expansion routes inferred from phylogenetic analyses revealed the extent of the mutation load. A) population phylogenetic tree inferred from the pairwise F_{ST} values (see also Figure S3) superimposed on the map. Map produced with R using the worldHires data [51]. B) Population specific branches are coloured according to the π_N/π_S values, used as a proxy of the genetic load. C) Correlation between tree branch lengths to the root and the π_N/π_S ratio. D) Correlation between the travelled distance to the southernmost site and the π_N/π_S ratio. The blue line represents the slope of the linear model between the two variables. The grey area in panel C and D represents the 95% confidence interval levels around the regression lines.

The interpretation of π_N/π_S in populations that are not at mutation-drift equilibrium (say after a bottleneck)

is however difficult, since π_N will reach an equilibrium value faster than π_S because selected alleles will be

subject to higher negative selection and undergo a faster turnover [14,52,53]. Thus, the π_N/π_S may not be the

best predictor of the total burden of deleterious mutations, as it is potentially affected by demography and

selection [14]. To circumvent this limitation, we counted and plotted the distribution of non-synonymous

mutations classified according to i) their impact and expected consequences on fitness (*i.e.* missense and LoF

mutations) and; ii) segregation patterns (*i.e.* additive (total) load composed of heterozygous and derived

homozygous genotypes or recessive (fixed) load composed of homozygous derived genotype) [54]. Just as

the π_N/π_S, we found a significant association between the derived load and the distance to the southernmost

sites both for missense mutations (R^2 = 0.790, p < 0.001, **Fig 3A**) and for LoF mutations (R^2 = 0.777, p < 0.001,

Fig 3B). Similarly, we observed a significant association between the tree branch length to the root and the

derived load of missense and LoF mutations (R^2 = 0.39, p = 0.0096 and R^2 = 0.40, p = 0.009, respectively,

Table S4).

The most deleterious mutations are efficiently purged across the range

The total number of missense mutations was significantly more abundant in the more southern populations

(*i.e* California, Cascadia, **Table S5** for all Tukey-HSD *p-values*) than in northern ones (**Fig 3A**), while it was

nearly constant from the south to the north regarding LoF mutations (**Fig 3B**), with no significant differences

among populations (**Table S5**). The southernmost populations displayed a higher load of mutations in

heterozygous state (**Fig 3**), as expected due to their higher historical effective population size, favouring the

segregation of recessive mutations hidden in a heterozygous state [42,55].

The fixed load increases with population expansion

The fixed load (i.e. count of derived homozygous sites) increased from south to north for both missense and

LoF with the most extremes samples, including Mile Slough (MSL), Porcupine (POR) and Salmon (SAL) rivers

being the most significantly loaded (**Table S5**), as expected due to founder events and allele surfing for

Alaskan/Yukon samples and due to bottlenecks for the Salmon River (**Fig 3** right panel).

Figure 3: *Number of deleterious alleles per riverper individual for all populations sorted from the South to the North. The plot* shows the additive load (left panel) and recessive derived homozygous load (right panel) for A) missense mutations and B) Loss-Of-Function (LoF) mutations. No strong differences are observed in the additive load among populations. In contrast, significant differences were observed for the recessive load in populations at the expansion front both for missense mutations and LoF mutations and this was significantly correlated with the distance to the southernmost site. Each color represents a major regional group. Abbreviation for each site is provided in Table S01.

Range expansion have reduced the efficacy of selection at range margin

We predicted that selection efficacy will be reduced at the expansion front and in populations with a lower N_e. To test this, we measured three parameters, namely the rate of non-adaptive and adaptive amino-acid substitutions relative to neutral divergence (ω_{NA} and ω_A, respectively; with $\omega_A = d_N/d_S - \omega_{NA}$) and the proportion of non-synonymous amino-acid substitution that result from positive selection ($\alpha = \omega_A/(d_N/d_S)$) using the software Grapes [56] using maximum likelihood estimation under a population genetic model. A general expectation is that populations at the expansion margin will exhibit a higher rate of non-adaptive substitutions (ω_{NA}), a lower rate of adaptive substitutions (ω_A) and a lower proportion of amino-acid substitutions (α).

We observed a reduced efficacy of purifying selection likely associated with the multiple founder

events and bottlenecks which resulted in a decrease rate of adaptive substitutions (ω_A, R^2 = 0.37, p = 0.013)

and increased ω_{NA} as a function of the distance to the South (R^2 = 0.57, p = 0.0011, **Fig 4A**). We also found a

significant positive correlation between ω_{NA} and the distance to the ocean (R^2 = 0.52, p = 0.002, **Fig S6A**) and

conversely a negative correlation between ω_A and the distance to the ocean (R^2 = 0.40, p = 0.00895, **Fig S6B**),

suggesting that upstream populations displays a lower adaptive potential. This observation thus suggests

that the population expansion had impacts on both the adaptive and non-adaptive substitution rates.

Figure 4: Efficacy of selection decreases at the expansion front and as a function of long-term effective population
size. A) increased rate of nonsynonymous non-adaptive substitutions (ω_{NA}) in northernmost populations of Coho
salmon (Alaska) and in the bottlenecked Thompson River. B) Correlation between α and historical variation in the
coalescent effective population size estimated from SMC++. See also **Fig S7-S8** for correlations based on tree branch
lengths. Note that the distance from the southernmost site was computed considering a Californian sample, from a
previous study [42], for which no WGS data were available.

Similarly strong correlations were observed when considering the tree branch length to the root (details in

Fig S7-S8, Table S4). The Salmon R. population (Thompson R. watershed), which suffered a recent decline in

abundance, supposedly due to the sustained release of hatchery fish derived from few individuals and the

Ryman-Laikre effect [57] (see discussion below), also displayed a high ω_{NA}.

We also predicted a higher α value in populations with higher N_e and, conversely, a decreased value

of α in populations with lower N_e. We observed a strong positive correlation between α and the synonymous

nucleotide diversity (π_S) of each local population, which represents a good predictor of the long term

(coalescent) N_e (R^2 = 0.63, p = 0.0004, **Fig S9**). To more directly test the association with N_e, we correlated

SMC++ based estimates of N_e (averaged over a 200KyA window) with α and recovered significant

correlations (R^2 = 0.49, p = 0.004, **Fig 4B**). It is noteworthy that a good correlation between proxies of N_e

obtained from SMC++ and π_S was also observed (R^2 = 0.68, p = 0.00017, results not shown). Altogether, the

above results provide empirical support pertaining to the evolutionary consequences of allele surfing at

expanding range margins, in particular regarding the loss of adaptive potential and the mutation burden.

Recombination rates shape the deleterious mutational landscape

In addition to the spatial structure associated with the postglacial recolonization, we investigated the genome-wide variation in mutation load. Such variation could be associated with the occurrence of structural variants (e.g. chromosomal inversions) which may incur a significant load because deleterious recessive mutations may freely accumulate in the absence of recombination, as observed for instance in sex chromosome and related supergene-like architecture [58]. To test this hypothesis, we used the GC content at third codon position (GC3 hereafter) as a proxy of the rate of recombination [59,60] (see Sup. Note S2 for an explanation). We observed strong correlations between levels of GC3 and π_N/π_S ratio for all Coho salmon populations (R^2 range = 0.938 – 0.955, p<0.001) except for population MSL (R^2 = 0.252; p = 0.0038) (**Fig 5A**). An analysis focused on GC-conservative sites, which are not affected by GC-biased Gene Conversion (gBGC), revealed similarly strong patterns across all Coho salmon populations (**Fig 5B**, R^2 range = 0.665 - 0.876, p < 0.01). π_N/π_S ratios estimated based on all sites or GC-conservative-sites only are highly correlated (Pearson r = 0.909, p <0.001). We tested the generality of this relationship using four other closely related Pacific salmon species and found strikingly similar pattern in Chinook salmon (R^2 GC3 ~ π_N/π_S = 0.9338, p = 1e-11) Sockeye and Kokanee salmon (R^2 range = 0.81 - 0.96, p <0.001), as well as Rainbow trout populations (R^2 range = 0.95 - 0.96, p <0.001, **Fig 5C**).

Figure 5) The deleterious load is determined by variation in GC3 content in multiple salmonids
A) Correlations between the deleterious load (π_N/π_S) and GC content at third codon position in multiple Coho salmon populations. Each line represents the values colored by major regional groups (See **Fig S10A** for detail). B) Correlations between the deleterious load (π_N/π_S) and the GC content at third codon position considering an independent method based on the site frequency spectrum at GC-conservative sites. C) Correlation between the deleterious load (π_N/π_S) and GC content at third codon position in sockeye salmon ecotypes (sockeye and kokanee), chinook, pink salmon and in rainbow trout. Averages are provided for rainbow trout, salmon and kokanee at the species level, see **Fig S10B-C** for detail by population.

We then analyzed the difference in the correlations between levels of GC3 and π_N/π_S ratio among all Coho salmon populations. Indeed, the intensity of this correlation highlights the effect of recombination rate

variation on the efficiency of purifying selection [61]. The populations at the expansion front (Mile Slough R.,

Porcupine R., Snake R.) in Alaska exhibit the lowest correlations between GC3 and π_N/π_S. If both

demographic factors (*i.e.* distance to the source here) and genomic factors (*i.e.* recombination) interact to

shape the load, we may predict a positive relationship between these two factors because the northernmost

populations should have an even higher load in regions of low recombination. We tested this hypothesis

using a linear model between the slope of the regression between π_N/π_S and GC3 and the distance to the

ancestral source populations did not reveal such a relationship (p = 0.133, **Fig S11**).

Regions of residual tetraploidy revealed drivers of the load in Coho salmon

We tested the hypothesis that regions with residual tetraploidy exhibit a reduced load due to increased

efficacy of purifying selection due to higher population sizes (4*Ne* rather 2*Ne*). Contrary to this expectation,

increased deleterious load was observed in regions of residual tetraploidy (3,700 genes) as compared to

diploid regions (**Fig 6A**, red dotted line mean $\pi_N/\pi_S > 0.35$, diploid region $\pi_N/\pi_S < 0.30$). Given that we also

observed lower levels of recombination in regions with residual tetraploidy compared to re-diploidized

genomic regions (**Fig S12,** p < 0.0001, W Mann-Whitney = $3.04e^7$, see also **Fig S13** and **table S6** for

differences in recombination among populations), our results suggest that this higher π_N/π_S could be mostly

due to lower recombination rates. Another expected genomic consequence of this lower recombination rate

is a higher load of transposable elements [62]. When computing the relative length of TE, *i.e.* the length of

TE corrected by the chromosome length (see methods), we found a significant enrichment of TE in the

regions of residual tetraploidy as compared to diploid chromosomes (**Fig 6B**, p <0.0001, WMann-Whitney =

$1.36e^{10}$). This tendency was also observed across the different TE categories (**Fig S14)**. To more directly test

the N_e-effect hypothesis, we eliminated the effect of the recombination rate by comparing the load across

similar bins of GC in diploid vs. tetraploid regions. The π_N/π_S was systematically higher in diploid regions than

in the tetraploid ones after excluding the class with the lowest GC content (**Table S7, Fig 6C**) indicating that

the load was significantly higher in diploid compared to tetraploid regions, with the notable exception of

regions with extremely reduced recombination. Therefore, it is still possible that increased efficacy of

selection is at play in recombining regions of residual tetraploidy, following the hypothesis of higher effective

size in these regions.

Figure 6: Increased load in region of residual tetraploidy. A) Distribution of π_N/π_S ratio when considering all genes in regions of residual tetraploidy (red lines) for each population compared to a set of 200 randomly generated samples of 4,000 genes in the rest of the genome (gray histogram). **B**) Violin plot showing that regions of residual tetraploidy displayed significantly longer transposable elements when compared to diploid regions. Orange = diploid chromosome. Gray = chromosome with residual tetraploidy. Red point = mean +/- 1*sd. **C**) Difference in the distribution of $\pi N/\pi S$ ratio in different genomic regions of the Coho salmon. Each dot represents the value observed for a given population in either diploid or tetraploid region with the "lowest" GC3 value (corresponding to low recombination) compared to the rest of GC3 values (labelle "other GC3", corresponding to intermediate to high recombination). The GC values were computed in 4mb windows separately for diploid and tetraploid regions.

Conclusion

The role of genetic drift, recombination, selection and variation in inheritance in affecting the deleterious load and efficacy of natural selection is a central question with fundamental and applied consequences for biodiversity.

Using population genomics analyses, we investigated the evolutionary consequences of recent demographic events in Coho salmon. Our results supported gene surfing across North America [42]. Results also supported lower N_e at the northern range expansion front, which induced two main evolutionary consequences: a surf of slightly deleterious mutations and a putative reduction of the adaptive potential, as expected under the nearly neutral hypothesis of molecular evolution [12].

We further demonstrated that one population, the Salmon River (Thompson R. watershed) displayed

an increased fixed load and decreased selection efficacy compared to southern populations. Previous studies

showed that this population is genetically isolated from others [42,43,63] and displayed genomic footprints

of a bottlenecked population such as long runs of homozygosity [41]. This population has been subjected to

extensive hatchery enhancement (from a single population) to circumvent the decline of Coho salmon in the

Thompson drainage [64]. From a theoretical standpoint, if a few captive individuals were used as parents for

subsequent releases, then an increase in inbreeding and a reduction in effective population size are

expected, a phenomenon called the Ryman-Laikre effect [57] [65]. Consequently it is possible that the

increased π_N/π_S, ω_{NA}, and fixed deleterious load could be explained by both long-term evolution at low N_e

and by the Ryman-Laikre effect associated with hatchery enhancements. Regardless, this suggests that

careful enhancement needs to be performed with a diversity of parents to maximize genetic diversity of

supplemented populations. This result has implications for fisheries enhancement, as well as genetic rescue

programs aiming at reducing the inbreeding load of declining populations and restoring the fitness of these

populations.

It is noteworthy that the southernmost population (Klamath R, California) was not characterized by the

highest α value. Several non-exclusive hypotheses may explain this pattern: first, some Californian

populations are known to have undergone strong recent (human-driven) decrease in abundance due to both

habitat degradation and climate change, which may leave a detectable footprint in the genome [66]. The

recent demography could have an impact on the accuracy of the fit of the synonymous and nonsynonymous

SFS under Grapes. Second, the Klamath R. population is located in the far upstream part of that river, and

may have undergone a founder event when reaching this upper part, a hypothesis that is supported by the

correlations we observed between several metrics and distance to the river mouth. Finally, we did not

include the southernmost populations from California, which are more likely to be more ancestral, as we

previously reported [42].

Since the use of π_N/π_S ratio can be criticized [14,67,68], especially when used to quantify the load within

species, we computed additional metrics [56] to more directly estimate mutation load, as advocated by

others [14,69]. In this case, we found small differences in the additive load among Coho salmon populations,

but a linear increase in the recessive load as a function of the distance to the southernmost sites and as a

function of the tree branch length, both are used as proxies of the expansion route.

We demonstrated that the π_N/π_S ratio was negatively correlated with the GC content at third codon

position, which represents a good proxy of the local recombination rate (see **note S2**). The negative

correlation between π_N/π_S and GC3 was repeatedly observed across different salmonid species using both a

sequence based estimate of π_N/π_S and an estimate based on GC-conservative site (non-affected by gBGC)

[60]. These results indicated that recombination plays a key role in explaining the variation in the mutation

load along the genome in salmonids. Our study empirically supports theoretical work about the

accumulation of slightly deleterious mutations in non-recombining regions [30,32]. Similarly, increased

prevalence of deleterious mutations in low recombining regions have been reported in plants [70] and in

human populations [71–73]. In particular, recent work in human populations have shown that both variation

in demographic history (i.e. change in effective population size) and recombination rates are affecting allele-

specific expression of harmful mutations [71]. The authors showed that allele specific expression causes

underexpression of harmful mutations more efficiently in normally and highly recombining regions

compared to low recombining regions. They further documented variation of this process among

populations with varying demographic histories.

In line with the key role of recombination rate, we observed that only regions of residual tetraploidy with

extremely low recombination rate (lower GC content) displayed an increased load in Coho salmon,

accumulating more transposable elements, suggesting efficient selection in more "normally" recombining

region of residual tetraploidy. In particular, our results indicate that i) the regions of residual tetraploidy

(4Ne) with a normal recombination rate do not display an increased load, which may be expected under the

nearly neutral theory if higher Ne is associated with the efficacy of selection; ii) only region of extremely low

recombination rate (lower GC content) displays the highest load, highlighting once again the primary role of

recombination in shaping the load, with a higher contribution than Ne. A third factor is the variation in

dominance of deleterious mutation. Indeed, a recent simulation study found that in the case of hard sweeps,

dominance of recessive mutation was a central aspect determining the signal of sweeps in polyploid

genomes [39]. A detailed investigation of this aspect in regions of residual tetraploidy was beyond the scope

of our study but would be worthy of further investigation.

The detailed consequences of ongoing rediploidization have been extensively studied at the regulatory levels

elsewhere, with support for an increased load (higher d_N/d_S and TE load) in duplicated genes undergoing

lower expression [74]. Recent studies in polyploid plant species have also documented various evolutionary

consequences of such duplication on local variation in the load and efficacy of selection [35]. For instance in

A. arenosa, a reduced efficacy of purifying selection was suggested in tetraploids (4X) genome compared to

diploids (2X), and the authors suggest that this is because deleterious alleles are better masked in

autotetraploids [38]. Similarly, a recent work in the allopolyploid cotton *(Gossypium)* demonstrated that this

species accumulates more deleterious mutations than the diploid species [75]. This observation supports a

theory proposed by Haldane that recessive deleterious mutations accumulate faster in allopolyploids

because of the masking effect of duplicated genes [76]. To sum up, our results concur with predictions from

the nearly neutral theory of molecular evolution [12], in which slightly deleterious mutations are effectively

neutral and purged effectively in regions of higher recombination, except perhaps in populations at the

extreme of the expansion front. Interestingly, these results indicate that regions of low-recombination re-

diploidize later than other genomic regions. We are not aware of any paper having documented such

patterns so far. Further studies of this process across more species would be welcome to validate the

generality of this observation.

Finally, our results have implications for conservation practices. We showed that the additive load is

approximately constant, indicative of efficient purging across populations for both missense and LoF, similar

to some recent studies on several plant and animal models [77–80]. However, results indicate that

population at higher latitude from the Yukon watershed (Mile Slough R., Porcupine R.) or from the

bottlenecked Salmon R. have not entirely purged the most deleterious mutations, including missense and

LoF mutations, which may impose a fitness cost to these populations. Further empirical evidence for a causal

link between the putative fitness cost of LoF mutations and adaptive phenotypic variation will be necessary

to validate our observations. With this caveat in mind, our results for the Salmon R could guide practices in

supplementation programs. For instance, choosing a diversity of parents from moderately differentiated

populations of modest size may help increase the levels of heterozygosity and mask the expression of

recessive deleterious mutations [55]. This strategy could reduce the occurrence of deleterious alleles and

counteract the Ryman-Laikre effect described above. Moreover, populations (e.g. most upstream

Alaskan/Yukon populations) for which our results suggest a reduced adaptive potential are also the most

strongly exposed to rapid climate change, as the rate of temperature increase is most rapid at higher

latitudes [81]. In all cases, maximizing the connectivity among populations and limiting habitat degradation

appears as fundamental strategies to maintain high effective population size and increase the adaptive

potential of Coho salmon to the multiple ongoing anthropogenic pressures [82]. More generally, how best to

manage declining populations and guide conservation policies in these conditions to minimize the load

and/or maximize genetic diversity is another debated issue [55,83–85]. In the meantime, while conservation

genomics undoubtedly has a major role for the short-term preservation of endangered species, this should

not override the crucial need for reducing human impacts on natural ecosystems to preserve biodiversity

over long time scales [86].

Methods

Sampling design for Coho salmon

We sampled and sequenced 71 individuals representing 14 populations distributed from California to Alaska

(**Table S1**). A set of 55 individuals was sequenced on an Illumina HiSeq2500 platform [38] and the other 16

individuals were sequenced on a NovaSeq6000 S4 platform using paired-end 150 bp reads. Reads were

processed using fastp for trimming [87], and mapped to the most recent Coho reference genome

(https://www.ncbi.nlm.nih.gov/assembly/GCF_002021735.2/) using bwa mem v2 [88]. Reads with a

minimum quality below 20 were discarded with samtools v1.7. Duplicates were removed with picard

(http://broadinstitute.github.io/picard/). SNP calling was performed using GATK v4.2 [89] using our pipeline

available at github.com/quentinrougemont/gatk_haplotype/. We generated a Haplotype gVCF for each

sample individually, combined all gVCF and then performed a joint genotyping. We checked the variants

quality score of our data and filtered our genotypes according to their quality following GATK best practices

and based on quantiles distributions of quality metrics. We excluded all sites that did not match the

following criterion: MQ < 30, QD < 2, FS > 60, MQRankSum < -20, ReadPosRankSum < 10, ReadPosRankSum >

10. We also excluded multiallelic SNPs, as well as indels. Genotypes with a depth lower than 6 or higher than

100 reads were also excluded to remove low confidence genotypes potentially associated with paralogs.

Finally, we also generated a separate vcf file using the --all-sites option to obtain a file with invariant position

to reconstruct sequence data (see the **Genetic load estimation in Coho salmon and related species** section).

A total of 14,701,439 SNPs were identified without missing data. Population structure was evaluated using a

principal component analysis (PCA) performed using Ade4 [90] on a set of LD-pruned SNPs without missing

data (1,739,037 SNPs) identified stringently with plink1.9 [91] (command indep-pairwise 100 50 0.1).

Outgroup dataset.

In order to test the generality of the relationship observed between π_N/π_S and GC3 we took advantage of the

newly assembled reference genomes and resequencing data from other closely related Pacific salmonid

species with similar demographic histories[92–94]. Sockeye salmon published by [92] were retrieved from

NCBI PRJNA530256. Samples with a high number of individuals per ecotype (Sockeye and Kokanee) were

chosen from the NCBI table. We retained a total of 5 Kokanee populations and 5 Sockeye populations (from

Fraser & Columbia watershed described in T**able S1 and Table S3**). Three Chinook salmon samples were

provided by B. Koop (also available at NCBI PRJNA694998). Additionally, 11 samples of "even" and 10

samples of "odd" pink salmon (*O. gorbuscha*) were downloaded from PRJNA 556728 and included in the

analysis. Here "even" and "odd" refers to Salmon returning to their natal rivers in "even" and "odd" years to

spawn, leading to a temporal isolation of these ecotypes [93]. These were indeed clearly separated based on

a PCA. Finally, a number of rainbow trout available from NCBI PRJNA386519 were used (n = 19 from 3

random populations showing genomic differentiation based on a PCA).

Each sample was downloaded and mapped onto its species' reference genome downloaded from NCBI and

using the exact same procedure as described above relying on fastp, bwa-mem2, picard and GATK 4.2.5.0 to

generate a final vcf filter based on usual GATK quality criteria and variance in sequencing depth. For each

species, we then quantified the load using the π_N/π_s ratio with the procedure described below for Coho

salmon. For the Sockeye/Kokanee ecotypes, the Sockeye is a fully migratory ecotype, whereas the Kokanee

is a resident (non-migratory) ecotype that typically comprises more isolated populations. The two alternative

ecotypes are sometimes found in similar locations, with three rivers from our sampling design including both

ecotypes (Table S03). For all other species we tested the relationship between π_N/π_S and GC3 (see below).

Ancestral and derived alleles identification

To accurately recover the ancestral allelic states, we used three outgroup species; including the chinook salmon and rainbow trout sample (see above, n = 5 for rainbow trout) plus data from Atlantic salmon (n = 5, SRP059652). Each individual was aligned against the Coho salmon V2 genome (GCF_002021745.2) using GATK with the same procedure as above and calling every SNP using the –all-site mode to obtain invariant positions. We then determined the ancestral state of each SNP if 1) the SNP was homozygous in at least 90% of the individuals from our three outgroups, and 2) matched one of the two alleles identified in Coho salmon. Otherwise, the site was inferred as missing and was not used in subsequent analyses of the load. In addition, we reconstructed a consensus ancestral fasta sequence using the doFasta option from angsd [95]. This was used for demographic reconstruction detailed in **Note S1.**

Demographic reconstruction

We first tested our prediction that genetic diversity (observed heterozygosity H_o) decreases towards the North following our «out of Cascadia» model previously inferred [42,43]. Conversely, we verified that the β_{ST} coefficient, a measure of both genetic differentiation and ancestrality [46] increases northwards the North, as expected due to isolation by distance. β_{ST} and observed heterozygosity (H_o) were measured using the hierfstat R package [96] Details about β_{ST} computation are provided in Weir & Goudet [46] and H_o was computed following Nei (1987)[97] as $1 - \sum_k \sum_i Pkii / np$ with *Pkii* the proportion of homozygotes *i* in sample *k* and *np* the number of samples. Oceanic coastal distances were computed using the marmap package [98] and waterway distance was computed using ArcGIS. Then we summed the 2 distances to obtain the distance to the most likely refugia that we identified in our previous studies [42,43]. A shapefile of the rivers used here is available at https://github.com/QuentinRougemont/selection_efficacy/tree/main/00.raw_data.

Under postglacial expansion from a single refugium, the general hypothesis would be that all sampled populations should follow a common temporal trajectory of a population decline (bottleneck due to founder events by few individuals) followed by a (strong) increase in N_e corresponding to the expansion phase. To test this hypothesis, we inferred temporal changes in N_e using SMC++ [50]. SMC++ works similarly to the PSMC model but takes into account the Site Frequency Spectrum (SFS) and is better suited to large sample sizes. Estimates of changes in population size were performed for all populations. To validate the fact that expansions are indeed postglacial, splitting time was estimated between all pairs of samples from different geographic areas based on the joint SFS (n = 75 pairwise comparisons). A generation time of 3.5 years, well documented for Coho, and a mutation rate of $8e^{-9}$ mutation/bp/generation were applied (corresponding to the median substitution rate inferred in Atlantic salmon, J. Wang, Personal communication). We also compared the results to the mean substitution rate of $1.25e^{-8}$ mutation/bp/generation also inferred by Wang.

In addition, pairwise linkage disequilibrium provides valuable information regarding population size and

inbreeding. We computed the squared correlation coefficient between genotypes in each sample and all

populations separately using popLDdecay [99]. We used a MAF of 5% in each population, keeping between

3.7 and 6.4 million SNPs with populations having undergone stronger bottleneck/founding events displaying

the lowest amount of variation. We estimated LD decay by plotting LD (R^2) against physical distance

measured in base pairs in ggplot2 [100] package in R.

We observed slight discrepancies between our SMC++ estimates of divergence time and our previous work

based on the site frequency spectrum [42]. To investigate this, we performed a new set of inference based

on the unfolded joint site frequency spectrum (jSFS) using ∂a∂i [101] and a new set of refined models as

detailed in **supp. Note S1, Table S8-S9**.

To reconstruct the population expansion routes we computed Weir and Cockerham [102] F_{ST} values among

populations using Vcftools [103] and constructed a phylogenetic population tree using the R package ape

[104]. We then computed the tree branch lengths to the root using the function distroot from adephylo

[105]. We also explored broad relationships among populations with treemix [106]. We fitted a model with

an increasing number of migration edges. We chose a model with K = 3 migration edges as all edges

displayed significant p-value and captured a high proportion of explained variance. Fitting more edges

decreased the p-value without really improving the fit to our data. Similarly, we computed the tree branch

length to the root using the function adephylo and compared our results to those from F_{ST}-based values. The

resulting tree branch length values were used as a proxy for the expansion routes, and we tested their

correlation using linear models in R, with all our metrics of deleterious load (π_N/π_S, total number of putative

homozygous derived deleterious alleles (recessive load) and total number of deleterious alleles (both in

homozygous and heterozygous states, additive load) for derived missense and LoF mutations) and metrics of

selection efficacy (ω_{NA}, ω_A, α detailed below). Similar inference was performed, but using the reconstructed

distances to the southernmost site included in our previous studies [42,43]. This distance was computed by

summing the oceanographic distance and river distances described above.

Population-scaled recombination rate

Statistical phasing of the Coho whole genome sequences was performed using the Shapeit software [107],

considering all individuals at once. We then estimated effective recombination rates ($\rho=4.N_e.r$ where r

represents the recombination rate per generation and N_e is the effective population size) along the genome

using LDHat [108]. Phased genotypes were converted into LDHat format using vcftools after excluding SNPs

with MAF < 10% since rare variants are not informative for such inferences. Following the guidelines of the

LDHat manual, the genome was split into fragments of 2,000 SNPs with overlapping windows of 500 SNPs.

We measured recombination rates independently for each population. Differences in the distribution of

population-scaled recombination was visualized using violin plot (**Fig S13**) and statistically tested using

Tuckey HSD tests in R (**Table S5**) .

Genetic load estimation

π_N/π_S **estimates**

The approach developed in [59] was used to reconstruct fasta sequences for each individual. For

each species (Coho and the other salmonids), coding sequences (CDS) from each reconstructed sequence

were extracted using the gff files available with the reference genome to estimate the nucleotide diversity

(π). We also concatenated the CDS sequences into different classes according to their length and computed

π_N and π_S over 4-Mb concatenated gene space windows. Such large windows reduce the stochasticity due to

the low π_S values in Coho salmon.

Identifying potential deleterious non-synonymous alleles

We tested the difference in count of non-synonymous mutations in each local population of Coho

salmon across non-synonymous missense mutations (putatively deleterious) and Loss of Function (LoF)

mutations (likely to be strongly deleterious) identified with SNPeff. We analyzed data in two ways: first, we

counted the total number of putative homozygous derived deleterious alleles (recessive load) per individual

as well as the total number of deleterious alleles (both in homozygous and heterozygous states, additive

load) using: Ntotal = 2 X N_{homo} + N_{hetero} [28]. These individual values were then averaged per population. We

tested for differences in the distribution of these counts among populations using an ANOVA followed by a

TukeyHSD test. The p-values were corrected using a Bonferroni correction. We also computed mean derived

allele frequencies (DAF) in all sampling locations and across mutation categories (synonymous, non-

synonymous missense and LoF). We also applied the commonly used Provean [109] software based on a

random set of non-synonymous mutations (but see **Note S4** for a brief discussion regarding the

limitations).These results were then compared with results from non-synonymous mutations **(Fig S15)**.

Correlation between π_N/π_S and recombination

We computed the GC content at third-codon positions of protein coding genes (GC3) which has been

shown to be an accurate proxy of local recombination rate in other species and these positions are generally

silent [85,86]. To compute π_N/π_S values we sorted genes by ascending GC3 values, which enabled us to

obtain a ratio based on genes with similar GC3 values. Moreover, we also used the site frequency-based

approach proposed by Rousselle et al. [60] to estimate the π_N/π_S ratios. This approach enabled us to

compute SFS separately for GC conservative sites (A<->T and C<->G mutations), that is, not affected by GC-

biased Gene Conversion (gBGC).

Finally, we measured the correlation between GC3 and π_N/π_S using linear models. We replicated these

analyses considering only genes (n = 3,500) and SNPs in regions of residual tetraploidy (8% of the genome).

DFE estimation and rate of adaptation in Coho salmon

We estimated the rate of non-adaptive and adaptive synonymous substitutions (ω_{NA} and ω_A, respectively;

with $\omega_A = d_N/d_S - \omega_{NA}$) and the proportion of amino-acid substitution potentially resulting from positive

selection ($\alpha = \omega_A/(d_N/d_S)$, with d_N being the rate of non-synonymous substitutions, d_S being the rate of

synonymous substitutions). To do so, we used the method implemented in Grapes v1.0 which builds upon

the approach of [110]. Grapes models the effect of favorable mutations on the non-synonymous site

frequency spectrum (SFS) while accounting for various confounding factors distorting the SFS (demographic

change, linked selection, genotyping errors, SNP misorientation).

We used the following parameters in Grapes v1.0: we assumed a negative Gamma distribution to the

synonymous and non-synoymous SFS (parameters GammaExpo in Grapes and parameter "unfolded" site

frequency spectrum). To obtain suitable data, we converted the quality filtered whole genome vcf file into a

fasta file for each population containing sequence information for each individual in the population (using

vcf2fasta available at https://github.com/QuentinRougemont/selection_efficacy/tree/main/08.piNpiS/). This

file was then converted into a site-frequency spectrum using bppml [111]. We required at least 10 sites

without gap to keep a given site (-gapN_site), with a maximum proportion of gap of 0.5 (-gapN_seq), a

minimum number of complete codon of 6 (-min_nb_codon), we did not allow frameshift (remove_frameshift

parameter), we also required a sample size of 8 (-sample_size), the number of initial/terminal position in

which Stop codons or FrameShift codons are tolerated was set to 20 (-tolerance_zone), we did not allow

Stop or FrameShift codons between the two tolerance zones (-allow_internal_bc parameter). The kappa

value (i.e Ts/Tv ratio) was set to 1.6 and we used an unfolded SFS using the three species as an outgroup.

Fitted parameters of the DFE were used to compute the expected d_N/d_S under near neutrality, which was

compared to the observed d_N/d_S to estimate α, $\omega_{NA,}$ ωa.

Differences in load for region of residual tetraploidy in Coho salmon

π_N/π_S **comparison**

In the Coho reference genome, Chromosomes 1 to 30 are considered to represent generally diploid

chromosomes, whereas chromosomes 31 to 38 represent those with a clear signal of residual tetraploid *[41]*.

We took advantage of this specificity regarding chromosome evolution to contrast the load for the 3700

genes in regions of residual tetraploidy (averaged across all genes for each population). For diploid regions,

we generated 200 datasets of 4,000 genes randomly sampled and then estimated the load for each of these

datasets.

TE annotations

We used the TEs annotation file from repeatmasker [112] (made available on NCBI for the reference genome [41]) and tested for difference in the length of TEs between diploid and tetraploid regions, after correcting for the difference in chromosome length.

Data availability

Acknowledgements

We are grateful to Alysse Perreault-Payette and Bérénice Bougas for help in DNA extraction of additional individuals for whole genome sequencing. We thank Benoit Nabholz for the initial development of the π_N/π_S C++ app we used here. We are grateful to the genomic platform at Compute Canada (Canada), IBIS (Canada), and MBB (Montpellier, France) for providing computational capacity as well as Bird (Nantes, France) for data storage. This research was carried out in conjunction with EPIC4 (Enhanced Production in Coho: Culture, Community, Catch), a project supported in part by University Laval, Ressources Aquatiques Québec (RAQ), and the Government of Canada through Genome Canada, Genome British Columbia, and Genome Québec. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References:

1. Hewitt GM. Genetic consequences of climatic oscillations in the Quaternary. Philos Trans R Soc Lond, B, Biol Sci. 2004;359: 183–195; discussion 195. doi:10.1098/rstb.2003.1388

2. Excoffier L, Foll M, Petit RJ. Genetic Consequences of Range Expansions. Annual Review of Ecology, Evolution, and Systematics. 2009;40: 481.

3. Slatkin M, Excoffier L. Serial Founder Effects During Range Expansion: A Spatial Analog of Genetic Drift. Genetics. 2012;191: 171–181. doi:10.1534/genetics.112.139022

4. Ohta T. Linkage disequilibrium due to random genetic drift in finite subdivided populations. Proc Natl Acad Sci U S A. 1982;79: 1940–1944. doi:10.1073/pnas.79.6.1940

5. Hallatschek O, Hersen P, Ramanathan S, Nelson DR. Genetic drift at expanding frontiers promotes gene segregation. Proc Natl Acad Sci U S A. 2007;104: 19926–19930. doi:10.1073/pnas.0710150104

6. Klopfstein S, Currat M, Excoffier L. The Fate of Mutations Surfing on the Wave of a Range Expansion. Molecular Biology and Evolution. 2006;23: 482–490. doi:10.1093/molbev/msj057

7. Travis JMJ, Münkemüller T, Burton OJ, Best A, Dytham C, Johst K. Deleterious Mutations Can Surf to High Densities on the Wave Front of an Expanding Population. Molecular Biology and Evolution. 2007;24: 2334–2343. doi:10.1093/molbev/msm167

8. Peischl S, Dupanloup I, Kirkpatrick M, Excoffier L. On the accumulation of deleterious mutations during range expansions. Mol Ecol. 2013;22: 5972–5982. doi:10.1111/mec.12524

9. Peischl S, Excoffier L. Expansion load: recessive mutations and the role of standing genetic variation. Molecular Ecology. 2015;24: 2084–2094. doi:https://doi.org/10.1111/mec.13154

10. Peischl S, Dupanloup I, Foucal A, Jomphe M, Bruat V, Grenier J-C, et al. Relaxed Selection During a Recent Human Expansion. Genetics. 2018;208: 763–777. doi:10.1534/genetics.117.300551

11. Haag CR, Roze D. Genetic Load in Sexual and Asexual Diploids: Segregation, Dominance and Genetic Drift. Genetics. 2007;176: 1663–1678. doi:10.1534/genetics.107.073080

12. Ohta T. Slightly Deleterious Mutant Substitutions in Evolution. Nature. 1973;246: 96–98. doi:10.1038/246096a0

13. Hedrick PW, Garcia-Dorado A. Understanding Inbreeding Depression, Purging, and Genetic Rescue. Trends in Ecology & Evolution. 2016;31: 940–952. doi:10.1016/j.tree.2016.09.005

14. Simons YB, Sella G. The impact of recent population history on the deleterious mutation load in humans and close evolutionary relatives. Curr Opin Genet Dev. 2016;41: 150–158. doi:10.1016/j.gde.2016.09.006

15. Lohmueller KE. The Impact of Population Demography and Selection on the Genetic Architecture of Complex Traits. Williams SM, editor. PLoS Genet. 2014;10: e1004379. doi:10.1371/journal.pgen.1004379

16. Willi Y, Fracassetti M, Zoller S, Van Buskirk J. Accumulation of Mutational Load at the Edges of a Species Range. Mol Biol Evol. 2018;35: 781–791. doi:10.1093/molbev/msy003

17. Willi Y, Fracassetti M, Bachmann O, Van Buskirk J. Demographic Processes Linked to Genetic Diversity and Positive Selection across a Species' Range. Plant Communications. 2020;1: 100111. doi:10.1016/j.xplc.2020.100111

18. Perrier A, Sánchez-Castro D, Willi Y. Expressed mutational load increases toward the edge of a species' geographic range. Evolution. 2020;74: 1711–1723. doi:10.1111/evo.14042

19. de Pedro M, Riba M, González-Martínez SC, Seoane P, Bautista R, Claros MG, et al. Demography, genetic diversity and expansion load in the colonizing species Leontodon longirostris (Asteraceae) throughout its native range. Molecular Ecology. 2021;30: 1190–1205. doi:10.1111/mec.15802

20. González-Martínez SC, Ridout K, Pannell JR. Range Expansion Compromises Adaptive Evolution in an Outcrossing Plant. Current Biology. 2017;27: 2544-2551.e4. doi:10.1016/j.cub.2017.07.007

21. Lucek K, Willi Y. Drivers of linkage disequilibrium across a species' geographic range. PLOS Genetics. 2021;17: e1009477. doi:10.1371/journal.pgen.1009477

22. Bosshard L, Dupanloup I, Tenaillon O, Bruggmann R, Ackermann M, Peischl S, et al. Accumulation of Deleterious Mutations During Bacterial Range Expansions. Genetics. 2017;207: 669–684. doi:10.1534/genetics.117.300144

23. Takou M, Hämälä T, Koch EM, Steige KA, Dittberner H, Yant L, et al. Maintenance of Adaptive Dynamics and No Detectable Load in a Range-Edge Outcrossing Plant Population. Molecular Biology and Evolution. 2021;38: 1820–1836. doi:10.1093/molbev/msaa322

24. Rousselle M, Simion P, Tilak M-K, Figuet E, Nabholz B, Galtier N. Is adaptation limited by mutation? A timescale-dependent effect of genetic diversity on the adaptive substitution rate in animals. PLoS Genet. 2020;16: e1008668. doi:10.1371/journal.pgen.1008668

25. Lanfear R, Kokko H, Eyre-Walker A. Population size and the rate of evolution. Trends Ecol Evol. 2014;29: 33–41. doi:10.1016/j.tree.2013.09.009

26. McDonald JH, Kreitman M. Adaptive protein evolution at the Adh locus in Drosophila. Nature. 1991;351: 652–

654. doi:10.1038/351652a0

27. Laenen B, Tedder A, Nowak MD, Toräng P, Wunder J, Wötzel S, et al. Demography and mating system shape the genome-wide impact of purifying selection in Arabis alpina. PNAS. 2018;115: 816–821. doi:10.1073/pnas.1707492115

28. Henn BM, Botigué LR, Peischl S, Dupanloup I, Lipatov M, Maples BK, et al. Distance from sub-Saharan Africa predicts mutational load in diverse human genomes. PNAS. 2016;113: E440–E449. doi:10.1073/pnas.1510805112

29. Do R, Balick D, Li H, Adzhubei I, Sunyaev S, Reich D. No evidence that selection has been less effective at removing deleterious mutations in Europeans than in Africans. Nat Genet. 2015;47: 126–131. doi:10.1038/ng.3186

30. Felsenstein J. The evolutionary advantage of recombination. Genetics. 1974;78: 737–756. doi:10.1093/genetics/78.2.737

31. Nordborg M, Charlesworth B, Charlesworth D. The effect of recombination on background selection. Genet Res. 1996;67: 159–174.

32. Muller HJ. The relation of recombination to mutational advance. Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis. 1964;1: 2–9. doi:10.1016/0027-5107(64)90047-8

33. Hill WG, Robertson A. The effect of linkage on limits to artificial selection. Genetics Research. 1966;8: 269–294. doi:10.1017/S0016672300010156

34. Hill RR. Selection in autotetraploids. Theor Appl Genet. 1970;41: 181–186. doi:10.1007/BF00277621

35. Baduel P, Bray S, Vallejo-Marin M, Kolář F, Yant L. The "Polyploid Hop": Shifting Challenges and Opportunities Over the Evolutionary Lifespan of Genome Duplications. Frontiers in Ecology and Evolution. 2018;6. Available: https://www.frontiersin.org/articles/10.3389/fevo.2018.00117

36. Monnahan P, Brandvain Y. The effect of autopolyploidy on population genetic signals of hard sweeps. Biology Letters. 2020;16: 20190796. doi:10.1098/rsbl.2019.0796

37. Baduel P, Quadrana L, Hunter B, Bomblies K, Colot V. Relaxed purifying selection in autopolyploids drives transposable element over-accumulation which provides variants for local adaptation. Nat Commun. 2019;10: 5818. doi:10.1038/s41467-019-13730-0

38. Monnahan P, Kolář F, Baduel P, Sailer C, Koch J, Horvath R, et al. Pervasive population genomic consequences of genome duplication in Arabidopsis arenosa. Nat Ecol Evol. 2019;3: 457–468. doi:10.1038/s41559-019-0807-4

39. Sergeant CJ, Sexton EK, Moore JW, Westwood AR, Nagorski SA, Ebersole JL, et al. Risks of mining to salmonid-bearing watersheds. Science Advances. 2022;8: eabn0929. doi:10.1126/sciadv.abn0929

40. Gustafson RG, Waples RS, Myers JM, Weitkamp LA, Bryant GJ, Johnson OW, et al. Pacific Salmon Extinctions: Quantifying Lost and Remaining Diversity. Conservation Biology. 2007;21: 1009–1020. doi:10.1111/j.1523-1739.2007.00693.x

41. Rondeau EB, Christensen KA, Minkley DR, Leong JS, Chan MTT, Despins CA, et al. Population-size history inferences from the coho salmon (Oncorhynchus kisutch) genome. G3 Genes|Genomes|Genetics. 2023;13: jkad033. doi:10.1093/g3journal/jkad033

42. Rougemont Q, Moore J-S, Leroy T, Normandeau E, Rondeau EB, Withler RE, et al. Demographic history shaped geographical patterns of deleterious mutation load in a broadly distributed Pacific Salmon. PLOS Genetics. 2020;16: e1008348. doi:10.1371/journal.pgen.1008348

43. Rougemont Q, Xuereb A, Dallaire X, Moore J-S, Normandeau E, Perreault-Payette A, et al. Long-distance migration is a major factor driving local adaptation at continental scale in Coho salmon. Molecular Ecology.

2023;32: 542–559.

44. Macqueen DJ, Johnston IA. A well-constrained estimate for the timing of the salmonid whole genome duplication reveals major decoupling from species diversification. Proc Biol Sci. 2014;281. doi:10.1098/rspb.2013.2881

45. Novembre J, Stephens M. Interpreting principal component analyses of spatial population genetic variation. Nat Genet. 2008;40: 646–649. doi:10.1038/ng.139

46. Weir BS, Goudet J. A Unified Characterization of Population Structure and Relatedness. Genetics. 2017;206: 2085–2103. doi:10.1534/genetics.116.198424

47. natural earth data. https://www.naturalearthdata.com/http//www.naturalearthdata.com/download/110m/physical/110m_physical.zip.

48. Hill WG, Robertson A. Linkage disequilibrium in finite populations. Theor Appl Genet. 1968;38: 226–231. doi:10.1007/BF01245622

49. Pickrell JK, Pritchard JK. Inference of Population Splits and Mixtures from Genome-Wide Allele Frequency Data. PLOS Genetics. 2012;8: e1002967. doi:10.1371/journal.pgen.1002967

50. Terhorst J, Kamm JA, Song YS. Robust and scalable inference of population history from hundreds of unphased whole genomes. Nature Genetics. 2017;49: 303–309. doi:10.1038/ng.3748

51. world Hire data. worldHires data public domain https://www.evl.uic.edu/pape/data/WDB/namer.tar.gz).

52. Gravel S. When Is Selection Effective? Genetics. 2016;203: 451–462. doi:10.1534/genetics.115.184630

53. Brandvain Y, Wright SI. The Limits of Natural Selection in a Nonequilibrium World. Trends in Genetics. 2016;32: 201–210. doi:10.1016/j.tig.2016.01.004

54. Bertorelle G, Raffini F, Bosse M, Bortoluzzi C, Iannucci A, Trucchi E, et al. Genetic load: genomic estimates and applications in non-model animals. Nat Rev Genet. 2022;23: 492–503. doi:10.1038/s41576-022-00448-x

55. Kyriazis C, Wayne RK, Lohmueller KE. High genetic diversity can contribute to extinction in small populations | bioRxiv. 10 Mar 2020 [cited 10 Mar 2020]. Available: https://www.biorxiv.org/content/10.1101/678524v1

56. Galtier N. Adaptive Protein Evolution in Animals and the Effective Population Size Hypothesis. PLOS Genetics. 2016;12: e1005774. doi:10.1371/journal.pgen.1005774

57. Ryman N, Laikre L. Effects of Supportive Breeding on the Genetically Effective Population Size. Conservation Biology. 1991;5: 325–329. doi:https://doi.org/10.1111/j.1523-1739.1991.tb00144.x

58. Berdan EL, Blanckaert A, Butlin RK, Bank C. Deleterious mutation accumulation and the long-term fate of chromosomal inversions. PLOS Genetics. 2021;17: e1009411. doi:10.1371/journal.pgen.1009411

59. Leroy T, Rousselle M, Tilak M-K, Caizergues AE, Scornavacca C, Recuerda M, et al. Island songbirds as windows into evolution in small populations. Current Biology. 2021 [cited 3 Mar 2021]. doi:10.1016/j.cub.2020.12.040

60. Rousselle M, Laverré A, Figuet E, Nabholz B, Galtier N. Influence of Recombination and GC-biased Gene Conversion on the Adaptive and Nonadaptive Substitution Rate in Mammals versus Birds. Mol Biol Evol. 2019;36: 458–471. doi:10.1093/molbev/msy243

61. MacPherson B, Scott R, Gras R. Sex and recombination purge the genome of deleterious alleles: An Individual Based Modeling Approach. Ecological Complexity. 2021;45: 100910. doi:10.1016/j.ecocom.2021.100910

62. Kent TV, Uzunović J, Wright SI. Coevolution between transposable elements and recombination. Philosophical Transactions of the Royal Society B: Biological Sciences. 2017;372: 20160458. doi:10.1098/rstb.2016.0458

63. Xuereb A, Rougemont Q, Dallaire X, Moore J-S, Normandeau E, Bougas B, et al. Re-evaluating Coho salmon (Oncorhynchus kisutch) conservation units in Canada using genomic data. Evolutionary Applications. 2022;15: 1925–1944. doi:10.1111/eva.13489

64. Government of Canada PS and PC. Stock assessment of Thompson River/upper Fraser River coho salmon / J.R. Irvine ... [et al.].: Fs70-1/1999-28E-PDF - Government of Canada Publications - Canada.ca. 1 Jul 2002 [cited 11 Aug 2022]. Available: https://publications.gc.ca/site/eng/9.804890/publication.html

65. Waples RS, Hindar K, Karlsson S, Hard JJ. Evaluating the Ryman–Laikre effect for marine stock enhancement and aquaculture. Curr Zool. 2016;62: 617–627. doi:10.1093/cz/zow060

66. Gilbert-Horvath EA, Pipal KA, Spence BC, Williams TH, Garza JC. Hierarchical Phylogeographic Structure of Coho Salmon in California. Transactions of the American Fisheries Society. 2016;145: 1122–1138. doi:10.1080/00028487.2016.1201003

67. Kratochvíl L, Rovatsos M. Ratios can be misleading for detecting selection. Curr Biol. 2022;32: R28–R30. doi:10.1016/j.cub.2021.11.066

68. Leroy T, Nabholz B. Response to Kratochvíl and Rovatsos. Curr Biol. 2022;32: R30–R31. doi:10.1016/j.cub.2021.11.067

69. Simons YB, Turchin MC, Pritchard JK, Sella G. The deleterious mutation load is insensitive to recent population history. Nature Genetics. 2014;46: 220–224. doi:10.1038/ng.2896

70. Ramu P, Esuma W, Kawuki R, Rabbi IY, Egesi C, Bredeson JV, et al. Cassava haplotype map highlights fixation of deleterious mutations during clonal propagation. Nat Genet. 2017;49: 959–963. doi:10.1038/ng.3845

71. Harwood MP, Alves I, Edgington H, Agbessi M, Bruat V, Soave D, et al. Recombination affects allele-specific expression of deleterious variants in human populations. Sci Adv. 8: eabl3819. doi:10.1126/sciadv.abl3819

72. Hussin JG, Hodgkinson A, Idaghdour Y, Grenier J-C, Goulet J-P, Gbeha E, et al. Recombination affects accumulation of damaging and disease-associated mutations in human populations. Nat Genet. 2015;47: 400–404. doi:10.1038/ng.3216

73. Alves I, Houle AA, Hussin JG, Awadalla P. The impact of recombination on human mutation load and disease. Philos Trans R Soc Lond B Biol Sci. 2017;372: 20160465. doi:10.1098/rstb.2016.0465

74. Gillard GB, Grønvold L, Røsæg LL, Holen MM, Monsen Ø, Koop BF, et al. Comparative regulomics supports pervasive selection on gene dosage following whole genome duplication. Genome Biology. 2021;22: 103. doi:10.1186/s13059-021-02323-0

75. Conover JL, Wendel JF. Deleterious Mutations Accumulate Faster in Allopolyploid Than Diploid Cotton (Gossypium) and Unequally between Subgenomes. Molecular Biology and Evolution. 2022;39: msac024. doi:10.1093/molbev/msac024

76. Haldane J. The Causes of Evolution. 1932 [cited 9 Jun 2023]. Available: https://jbshaldane.org/books/1932-Causes-of-Evolution/index.html

77. Khan A, Patel K, Shukla H, Viswanathan A, van der Valk T, Borthakur U, et al. Genomic evidence for inbreeding depression and purging of deleterious genetic variation in Indian tigers. Proceedings of the National Academy of Sciences. 2021;118: e2023018118. doi:10.1073/pnas.2023018118

78. Robinson JA, Räikkönen J, Vucetich LM, Vucetich JA, Peterson RO, Lohmueller KE, et al. Genomic signatures of extensive inbreeding in Isle Royale wolves, a population on the threshold of extinction. Sci Adv. 2019;5: eaau0757. doi:10.1126/sciadv.aau0757

79. Grossen C, Guillaume F, Keller LF, Croll D. Purging of highly deleterious mutations through severe bottlenecks in

Alpine ibex. Nat Commun. 2020;11: 1001. doi:10.1038/s41467-020-14803-1

80. Kleinman-Ruiz D, Lucena-Perez M, Villanueva B, Fernández J, Saveljev AP, Ratkiewicz M, et al. Purging of deleterious burden in the endangered Iberian lynx. Proceedings of the National Academy of Sciences. 2022;119: e2110614119. doi:10.1073/pnas.2110614119

81. Parmesan C, Yohe G. A globally coherent fingerprint of climate change impacts across natural systems. Nature. 2003;421: 37–42. doi:10.1038/nature01286

82. Irvine JR, Fukuwaka M. Pacific salmon abundance trends and climate change. ICES J Mar Sci. 2011;68: 1122–1130. doi:10.1093/icesjms/fsq199

83. Teixeira JC, Huber CD. The inflated significance of neutral genetic diversity in conservation genetics. PNAS. 2021;118. doi:10.1073/pnas.2015096118

84. García-Dorado A, Caballero A. Neutral genetic diversity as a useful tool for conservation biology. Conserv Genet. 2021;22: 541–545. doi:10.1007/s10592-021-01384-9

85. Kardos M, Armstrong EE, Fitzpatrick SW, Hauser S, Hedrick PW, Miller JM, et al. The crucial role of genome-wide genetic variation in conservation. Proceedings of the National Academy of Sciences. 2021;118: e2104642118. doi:10.1073/pnas.2104642118

86. Ripple WJ, Wolf C, Newsome TM, Gregg JW, Lenton TM, Palomo I, et al. World Scientists' Warning of a Climate Emergency 2021. BioScience. 2021;71: 894–898. doi:10.1093/biosci/biab079

87. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 2018;34: i884–i890. doi:10.1093/bioinformatics/bty560

88. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv:13033997 [q-bio]. 2013 [cited 3 Jul 2019]. Available: http://arxiv.org/abs/1303.3997

89. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A framework for variation discovery and genotyping using next-generation DNA sequencing data. Nature Genetics. 2011;43: 491. doi:10.1038/ng.806

90. Dray S, Dufour A-B. The ade4 Package: Implementing the Duality Diagram for Ecologists. Journal of Statistical Software. 2007;22: 1–20.

91. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. Am J Hum Genet. 2007;81: 559–575.

92. Christensen KA, Rondeau EB, Minkley DR, Sakhrani D, Biagi CA, Flores A-M, et al. The sockeye salmon genome, transcriptome, and analyses identifying population defining regions of the genome. PLOS ONE. 2020;15: e0240935. doi:10.1371/journal.pone.0240935

93. Christensen KA, Rondeau EB, Sakhrani D, Biagi CA, Johnson H, Joshi J, et al. The pink salmon genome: Uncovering the genomic consequences of a two-year life cycle. PLoS One. 2021;16: e0255752. doi:10.1371/journal.pone.0255752

94. Christensen KA, Leong JS, Sakhrani D, Biagi CA, Minkley DR, Withler RE, et al. Chinook salmon (Oncorhynchus tshawytscha) genome and transcriptome. PLOS ONE. 2018;13: e0195461. doi:10.1371/journal.pone.0195461

95. Korneliussen TS, Albrechtsen A, Nielsen R. ANGSD: Analysis of Next Generation Sequencing Data. BMC Bioinformatics. 2014;15: 356. doi:10.1186/s12859-014-0356-4

96. Goudet J. hierfstat, a package for r to compute and test hierarchical F-statistics. Molecular Ecology Notes. 2005;5: 184–186. doi:10.1111/j.1471-8286.2004.00828.x

97.   Nei M. Molecular Evolutionary Genetics. Molecular Evolutionary Genetics. Columbia University Press; 2019. doi:10.7312/nei-92038

98.   Pante E, Simon-Bouhet B. marmap: A Package for Importing, Plotting and Analyzing Bathymetric and Topographic Data in R. PLOS ONE. 2013;8: e73051. doi:10.1371/journal.pone.0073051

99.   Zhang C, Dong S-S, Xu J-Y, He W-M, Yang T-L. PopLDdecay: a fast and effective tool for linkage disequilibrium decay analysis based on variant call format files. Bioinformatics. 2019;35: 1786–1788. doi:10.1093/bioinformatics/bty875

100.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York: Springer-Verlag; 2009. Available: https://www.springer.com/gp/book/9780387981413

101.   Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD. Inferring the Joint Demographic History of Multiple Populations from Multidimensional SNP Frequency Data. PLOS Genetics. 2009;5: e1000695. doi:10.1371/journal.pgen.1000695

102.   Weir BS, Cockerham CC. Estimating F-Statistics for the Analysis of Population Structure. Evolution. 1984;38: 1358–1370. doi:10.2307/2408641

103.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call format and VCFtools. Bioinformatics. 2011;27: 2156–2158. doi:10.1093/bioinformatics/btr330

104.   Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. Bioinformatics. 2019;35: 526–528. doi:10.1093/bioinformatics/bty633

105.   Jombart T, Balloux F, Dray S. adephylo: new tools for investigating the phylogenetic signal in biological traits. Bioinformatics. 2010;26: 1907–1909. doi:10.1093/bioinformatics/btq292

106.   Pickrell JK, Pritchard JK. Inference of Population Splits and Mixtures from Genome-Wide Allele Frequency Data. PLOS Genetics. 2012;8: e1002967. doi:10.1371/journal.pgen.1002967

107.   Delaneau O, Zagury J-F, Robinson MR, Marchini JL, Dermitzakis ET. Accurate, scalable and integrative haplotype estimation. Nat Commun. 2019;10: 5436. doi:10.1038/s41467-019-13225-y

108.   McVean G, Awadalla P, Fearnhead P. A Coalescent-Based Method for Detecting and Estimating Recombination From Gene Sequences. Genetics. 2002;160: 1231–1241.

109.   Choi Y, Sims GE, Murphy S, Miller JR, Chan AP. Predicting the functional effect of amino acid substitutions and indels. PLoS ONE. 2012;7: e46688. doi:10.1371/journal.pone.0046688

110.   Eyre-Walker A, Keightley PD. Estimating the Rate of Adaptive Molecular Evolution in the Presence of Slightly Deleterious Mutations and Population Size Change. Molecular Biology and Evolution. 2009;26: 2097–2108. doi:10.1093/molbev/msp119

111.   Guéguen L, Gaillard S, Boussau B, Gouy M, Groussin M, Rochette NC, et al. Bio++: Efficient Extensible Libraries and Tools for Computational Molecular Evolution. Molecular Biology and Evolution. 2013;30: 1745–1750. doi:10.1093/molbev/mst097

112.   Tarailo-Graovac M, Chen N. Using RepeatMasker to identify repetitive elements in genomic sequences. Curr Protoc Bioinformatics. 2009;Chapter 4: Unit 4.10. doi:10.1002/0471250953.bi0410s25

Supporting Information Legend

Table S1: sampling strategy, including various outgroup species.

Table S2: Summary Statistics across coho populations, including geographic coordinates, tree branch length, genetic
diversity and load statistics. Figure 1 to Figure 4 can be reconstructed from these.

Table S3: Outgroup summary statistics by species, with a focus on the load

Table S4: Results of linear models. These provide tests of the relationship between branch length of the population
phylogenies based on (a) pairwise Fst value and b) treemix tree inferred with 3 migrations event. c) treemix results
obtained after removing the "salmon" river with aberrant branch length

Table S5: Results of Mann-Whitney test for difference in load among population based on missense mutation and
Loss of Function mutation. Mutation effect identified with SNPeff.

Table S06: Results of Tukey-HSD test for difference in recombination rate among populations based on Ldhat
estimates. "Pair" is the compared population pair with the initial described in Table S01. "Diff" is the difference in
recombination between the pair Lower and Upper CI are the 95% confidence level

Table S7: difference in load among diploid region, compared to region of residual tetraploidy with load computed in
500kb windows for each kind of regions

Tabble S8: AIC of each model for each pairwise comparison. AM = Ancient Migration, IM = Isolation With Migration, SI
= Strict Isolation, SC = Secondary Contact, G = prefix indicating Growth in the descending population, 2N= prefix
indicating linked selection, 2m = prefix indicating reduced effective migration along the genome, the A prefix after each
model (e.g. AM<u>A</u>) indicated a Growth in the ancestral population

Table S9: parameter estimates for each of the major regional group where $\partial a \partial i$ was fitted. Na = effective size of the
ancestral population, Ne pop1 = effective size of the first population, Nepop2 =effective size of the second population,
m12 = migration rate from 2 into 1, m21 being the reverse, me12 = effective migration rate in barriers regions, me21
being the same in the reverse, Tsplit = Split time, Tam = time of migration stop, P = proportion of the genome being
neutrally exchanged, Q = proportion the genome undergoing linked selection, O = proportion of the genome correctly
oriented, hrf = Hill-Roberston factor, indicating the extent of reduction in Ne in region affected by linked selection, b1,
b2 = extend of population growth in current population1 and 2 respectively

Supplementary Note S1: Reconstruction of demographic history from RADseq data

Supplementary Note S2: Relationship between GC3 and recombination

Supplementary Note S3 Comparison of Provean and non-synonymous results.

Figure S1: PCA and admixture plots among population.

A   Result of a principal component analysis obtained from a set of high quality biallelic SNPs without missing data
showing both a clusterization along latitude and longitude as well as discrete clusters corresponding broadly to
each river. Each label represents a given individual from a given river (labelled following table S01) and is
coloured according to its region of sampling.
B   LEA results for admixture inference for K = 14. Each bar represents an individual and is coloured according to
its membership probability. Each name corresponds to a river. The TsooYes appear as a mixture of different
individuals. Results must be interpreted with caution given the small sample sizes.

Figure S2: Summary statistics revealed the demography of Coho salmon

A   Positive correlation between the β_{ST} and distance to the southernmost site showing that differentiation
increased linearly from the south to the north. In all panels each point represents a sampling site and is
coloured according to the region in which it was sampled. The most negative values display likely ancestral
samples. The Thompson sample displays high inbreeding and is bottlenecked. Displayed is the adjusted R^2 of a
linear model along with its *p-value*. The grey area represents the 95% confidence interval levels around the
regression lines obtained with the predict function in R.

B   Negative correlation between genetic diversity (observed heterozygosity) and distance to the south.The Thompson sample displays high inbreeding and is bottlenecked. Displayed is the adjusted R^2 of a linear model along with its *p-value*. The grey area represents the 95% confidence interval levels around the regression lines obtained with the predict function in R.

C   Rates of LD decay as a function of distance along the genome. The higher LD indicates a history of inbreeding or bottleneck.

D   SMC++ inference of population size change with whole genome sequences for each local population of Coho salmon. Recent times should be interpreted carefully.

Figure S3: Inference of population split and mixture from Treemix.   A) Tree with three migration arrows. Each name describes a river sample site. Each river is color coded following the color scheme provided elsewhere (e.g. Fig 1). 3 significant migration arrows are displayed. Each migration arrow is colored according to the weight it received (from yellow to red) in Treemix. The weights are related to the fraction of alleles in the descendant population that originated in each donor population. Each node was highly supported based on 500 bootstrap. B) proportion of variance in the covariance of allele frequency explained as a function of the number of migration edges. C) Same tree colored according to the values of π_N/π_S

Figure S4: SMC++ split time.   Estimates of population split time from SMC++ under a model without gene flow among populations. Shown are estimates obtained when comparing split time between pairs of samples from different major regional groups. Two different mutation rates were used the: mean and median values based on *Salmo salar* orthologues mapped on the pike *Esox lucius* genome (Wang J. personal communication).

Figure S5: Correlation between π_N/π_S and π_S and π_N/π_S and N_e from smc++ A) Distribution of π_N/π_S as a function of π_S in each coho salmon populations from the study. B) Distribution of π_N/π_S as a function of N_e from SMC++ for each coho salmon populations from the study. Results of linear models are displayed. In all panels each point represents a sampling site and is coloured according to the region in which it was sampled. Displayed is the adjusted R^2 of a linear model along with its *p-value*. The grey area represents the 95% confidence interval levels around the regression lines obtained with the predict function in R.+

Figure S6: Correlation between distance to the ocean of each sample location (i.e. corresponding to the spawning migration) and the inferred rate of **A**) non-adaptive substitution (ω_{NA}) and **B**) adaptive substitution (ω_A). In all panels each point represents a sampling site and is coloured according to the region in which it was sampled. . Displayed is the adjusted R^2 of a linear model along with its p-value. The grey area represents the 95% confidence interval levels around the regression lines obtained with the predict function in R.

Figure S7: Results of linear models testing the effect of tree branch length to the root extracted from a F_{ST}-based population phylogeny on different metrics of selection efficacy   **A:** relationship between ω_{NA} and tree branch length; **B:** relationship between ω_A and tree branch length; **C:** relationship between α and tree branch length. See text for a definition of each metrics. Sample sites are coloured by region. The blue line represents the value of the regression line. In all panels each point represents a sampling site and is coloured according to the region in which it was sampled. Displayed is the adjusted R^2 along with its p-value.

Figure S8: Results of linear models testing the effect of tree branch length to the root extracted from a treemix population phylogeny on the load (π_N/π_S, panel A) and different metrics of selection efficacy B: relationship between ω_{NA} and tree branch length; **C:** relationship between ω_A and tree branch length; **D:** relationship between α and tree branch length. See text for a definition of each metrics). Sample sites are coloured by region. The blue line represents the value of the regression line. In all panels each point represents a sampling site and is coloured according to the region in which it was sampled. Displayed is the adjusted R^2 along with its p-value.

Figure S9: Correlation between the proportion of amino-acid substitution that results from positive selection (α) and the synonymous diversity π_S used as a proxy of effective population size. Each point represents a sampling site and is coloured according to the region in which it was sampled. Displayed is the adjusted R^2 along with its p-value. The grey area represents the 95% confidence interval levels around the regression lines obtained with the predict function in R.

Figure S10: Relationship between GC3 and pN/pS for all populations and all outgroups
A) Correlation for each population of coho salmon. Each point represents a sampling site and is coloured according to the region in which it was sampled; **B**) correlation within populations of rainbow trout. Each point represent a

population as infered using a PCA and corresponds to different rivers of sampling. **C)** correlation for each population of
Sockeye and Kokanee ecotype. Each point corresponds to differents rivers.
All correlations are significant. The x-axis displays the median GC3 and y-axis the π_N/π_S ratio. Abbreviation for each site
is available in Table S01. Displayed is the adjusted R^2 of a linear model along with its *p-value*. The grey area represents
the 95% confidence interval levels around the regression lines obtained with the predict function in R.

Figure S11: Relationship between recombination (GC3) and demographic factors (distance to the southernmost site).
A) relationship between the slope of the linear model between GC3 ~ and π_N/π_S and the distance to the southernmost
site. **B)** Correlation between the lowest recombining GC3 classes (expected to display the highest load) and the distance
to the southernmost sites. In all panels each point represents a sampling site and is coloured according to the region in
which it was sampled. Displayed is the adjusted R^2 of a linear model along with its *p-value*. The grey area represents the
95% confidence interval levels around the regression lines obtained with the predict function in R.

Figure S12: region of residual tetraploidy vs diploid region of the genome display different recombination landscapes.
Combined Violin plot and boxplot showing the distribution of population scale recombination ($\rho = 4*N_e*r$) inferred
from LDhat in Diploid chromosomes (orange) *vs* the 8 Regions of residual tetraploidy (gray).

Figure S13: Distribution of recombination rate among all populations. Each point represents the observed value of
population scale recombination rate ($\rho = 4N_e*\mu$) computed in 1 mb windows over the whole genome in each
population. The harmonic mean is plotted by a red dot along with its value. For each population a violin plot embedded
within a boxplot is shown.

Figure S14: TE length differs between diploid chromosome versus chromosome displaying residual tetraploidy.
Violin plot displaying the difference in TEs relative length (i.e. length corrected by the total length of each chromosome)
for each major TE category and each type of chromosome. Red point = mean +/- 1*sd.

Figure S15: Provean analysis of potentially deleterious mutation and the resulting load
Boxplot showing the number of deleterious alleles per river (sorted from the south to to north)
left panel = additive load, right panel= recessive load
top = missense deleterious mutations according to Provean predictions, bottom = missense tolerated mutations
according to Provean predictions
No strong differences are observed in the additive load among populations.
Significant differences were observed for the recessive load in populations at the expansion front which is qualitatively
similar to our inferences from missense and LoF mutations. Each color represents a major regional group. Results were
obtained for a random subset of mutations only given the strong computational burden of Provean.