Thermal phenotypic plasticity of pre- and post-copulatory male harm buffers sexual

conflict in wild *Drosophila melanogaster*

Claudia Londoño-Nieto[1*], Roberto García-Roa[1], Clara Garcia-Co[1], Paula González[1], and

Pau Carazo[1]

[1]Etology Lab, Cavanilles Institute of Biodiversity and Evolutionary Biology, University of

Valencia, Valencia, Spain.

ABSTRACT

Strong sexual selection frequently leads to sexual conflict and ensuing male harm, whereby males increase their reproductive success at the expense of harming females. Male harm is a widespread evolutionary phenomenon with a strong bearing on population viability. Thus, understanding how it unfolds in the wild is a current priority. Here, we sampled a wild *Drosophila melanogaster* population and studied male harm across the normal range of temperatures under which it reproduces optimally in nature by comparing female lifetime reproductive success and underlying male harm mechanisms under monogamy (i.e., low male competition/harm) vs. polyandry (i.e., high male competition/harm). While females had equal lifetime reproductive success across temperatures under monogamy, polygamy resulted in a maximum decrease of female fitness at 24°C (relative harm; H = 0.36), reducing its impact at both 20°C (H = 0.22), and 28°C (H = 0.10). Furthermore, female fitness components and pre- (i.e., harassment) and post-copulatory (i.e., ejaculate "toxicity") mechanisms of male harm were asymmetrically affected by temperature. At 20ºC, male harassment of females was reduced, and high male competition accelerated female actuarial senescence. In contrast, at 28ºC male ejaculate effects on female reproduction were modulated, and high male competition mostly resulted in accelerated reproductive ageing. We thus show that, across a natural thermal range, sexual conflict processes and their effects on female fitness components are plastic and complex. As a result, the net effect of male harm on overall population viability is likely to be lower than previously surmised. We discuss how such plasticity may affect selection, adaptation and, ultimately, evolutionary rescue under a warming climate.

1 INTRODUCTION

Females and males share the common goal of siring offspring. This central tenet of sexual
reproduction enforces a certain degree of cooperation between the sexes. However, product
of anisogamy, general asymmetries in the evolutionary interests around reproduction across
the sexes lead to distinct sex roles and thus diverging intensity and form of sexual selection
(Arnqvist & Rowe, 2005; Chapman, Arnqvist, Bangham, & Rowe, 2003). This phenomenon,
termed sexual conflict, favours traits in one sex that might be costly for the other (Parker,
1979), and can thus lead to antagonistic male-female coevolution (Arnqvist & Rowe, 2005).
Sexually antagonistic co-evolution has received much attention, and is recognized as a
fundamental process in evolution due to its role in shaping male and female adaptations
(Bonduriansky, Maklakov, Zajitschek, & Brooks, 2008), in contributing to drive reproductive
isolation and speciation (Arnqvist, Edvardsson, Friberg, & Nilsson, 2000; Bonduriansky,
2011; Bonduriansky & Chenoweth, 2009; Gavrilets, 2014), and as a major determinant of
population demography (Kokko & Brooks, 2003; Bonduriansky & Chenoweth, 2009; Berger
et al., 2016). Specifically, sexual conflict has been shown to have profound consequences
for female fitness and population growth when it favours male reproductive traits that
increase male intra-sexual competitive ability at the expense of harming females (i.e., male
harm, Crudgington & Siva-Jothy, 2000; Wigby & Chapman, 2004).

Harmful male adaptations are widespread and incredibly diverse and sophisticated across
the tree of life (Arnqvist & Rowe, 2005). For example, male harassment of females during
pre-copulatory competition for mating has been documented in myriad vertebrate and
invertebrate species (García-Roa et al., 2021), driving antagonistic male-female co-
evolution in a host of behavioural and morphological traits (Arnqvist & Rowe, 2005). Male
harm adaptations in the context of post-copulatory competition are similarly widespread in
invertebrates, featuring (amongst others) toxic ejaculates (Wigby & Chapman, 2005), love
darts (Koene & Schulenburg, 2005), and a range of male adaptations for traumatic
insemination that range from genital ablation to spiny penises (Crudgington & Siva-Jothy,
2000; Lange, Reinhardt, Michiels, & Anthes, 2013). Importantly, beyond driving male-female
phenotypes and associated diversification processes, male harm generally leads to a
"reproductive tragedy of the commons" that can substantially impact population demography
by depressing net female productivity (Arnqvist & Tuda, 2010; Berger et al., 2016; Holland
& Rice, 1999; Rankin et al., 2011), and even facilitate population extinction (Le Galliard,
Fitze, Ferriere, & Clobert, 2005). Unsurprisingly, understanding what factors underlie male

harm evolution, its diversity in form, strength, and outcomes, is a main concern in
evolutionary biology.

Despite a growing number of studies in the field of sexual conflict, most have been
conducted under uniform laboratory conditions, frequently in populations adapted to stable
environments for thousands of generations (Chapman et al., 2003b; Hopkins et al., 2020;
Wigby & Chapman, 2004, 2005). In contrast, recent research has highlighted the role of
ecology in shaping the evolution of traits under sexual conflict (Arbuthnott et al., 2014;
García-Roa et al., 2019; MacPherson et al., 2018; Perry & Rowe, 2018; Yun et al., 2017),
including habitat complexity (Malek & Long, 2019; Miller & Svensson, 2014; Myhre,
Forsgren, & Amundsen, 2013), nutritional status (Fricke, Bretman, & Chapman, 2010), or
sex ratio and population density (Chapman et al., 2003a). For example, Gomez-Llano and
colleagues (2018) recently showed that conspecific densities and the presence of
heterospecifics modify the intensity and outcome of sexual conflict in the banded demoiselle
(*Calopteryx splendens*), and the spatial complexity of the environment in which mate
competition occurs mediates how sexual conflict operates in fruit flies (Yun et al., 2018). The
incorporation of more realistic ecological scenarios in the study of sexual conflict is thus a
key avenue to disentangle the evolution of male harm, and its consequences for population
viability (Cornwallis & Uller, 2010; Fricke, Perry, Chapman, & Rowe, 2009; Plesnar-Bielak &
Łukasiewicz, 2021).

Temperature is recognized as a crucial abiotic ecological factor due to its impact on life
history traits and physiological and behavioural responses (De Lisle, Goedert, Reedy, &
Svensson, 2018; Kim, Jang, & Lee, 2020; Miler, Stec, & Czarnoleski, 2020; Monteiro et al.,
2017). Furthermore, in nature temperature varies widely both within and across spatio-
temporal scales (e.g., daily, inter-seasonal and intra-seasonal variation), and may
consequently have short, medium, and long-term effects on organism phenotypes that can
impact many different aspects of its reproductive behaviour (e.g., sex-specific potential
reproductive rates, operational sex ratios, density, etc.; García-Roa et al., 2020). In fact, a
recent meta-analysis suggests that temperature may have a sizeable effect on sexual
selection processes even when fluctuations occur well within the normal range of
temperature variation for the studied species (García-Roa et al., 2020). This latter finding is
particularly relevant given that we know almost nothing about how average temperature
fluctuations, such as those experienced by wild populations during their reproductive
season, affect male harm and sexual selection at large.

Our aim was to contribute to fill this gap in knowledge by studying how male harm responds
to temperature variation that mimics average fluctuations that are normal during the
reproductive season, using *Drosophila melanogaster* flies sampled from a wild population.
D. melanogaster is an ideal subject for this study for two main reasons. First, it exhibits high
levels of male- male competition and well characterized pre- and post-copulatory male harm
mechanisms, and is perhaps the main model species in the study of sexual conflict
(Arbuthnott et al., 2014; Chapman et al., 2003a; MacPherson et al., 2018; Malek & Long,
2019; Wigby & Chapman, 2005). During male-male pre-copulatory competition over access
to mating, males harm females via intense harassment that causes physical injuries,
interferes with female behaviours such as egg-laying or feeding, and results in energetically
costly resistance to males (Bretman & Fricke, 2019; Partridge & Fowler, 1990; Teseo,
Veerus, Moreno, & Mery, 2016). In the context of sperm competition over fertilizations, males
can also harm females via "toxic" ejaculates, whereby certain male seminal fluid proteins
manipulate female re-mating and egg-laying rates to the male advantage, but at a costs to
females in terms of lifespan and lifetime reproductive success (Chapman et al., 2003b; Sirot
et al., 2009; Wigby & Chapman, 2005). These proteins are secreted by male accessory
glands, have been well characterised, and are strategically allocated by males in response
to variation in the socio-sexual context (Hopkins et al., 2019; Sirot, Wolfner, & Wigby, 2011).
Second, despite the fact that almost all work on male harm has, to date, been conducted in
laboratory strains adapted to stable temperature conditions, *D. melanogaster* reproduces in
the wild under a wide range of temperatures (Dukas, 2020; Kapun et al., 2018).

We collected flies from a continental wild population in Requena (Spain) that experiences
significant fluctuations in temperature even during the mildest months when it is
reproductively active (e.g., July: average: 24.9°C, average min: 19.8°C, average max:
30.1°C). After acclimation of the resulting population to the laboratory conditions under a
fluctuating temperature regime mimicking natural conditions, we conducted a series of
behavioural and fitness assays to gauge how temperature variation within a normal range
affects: a) the overall impact of male-male competition on female lifetime reproductive
success (i.e., male harm), b) how the net effects of harm are accrued in terms of different
female fitness components (i.e., reproductive rate, actuarial ageing and reproductive
ageing), and c) underlying male pre-copulatory (i.e., harassment) and post-copulatory (i.e.,
ejaculate "toxicity") harm mechanisms.

RESULTS

<u>Lifetime reproductive success and related fitness components</u>

LRS- We detected a significant temperature x mating system interaction for female lifetime
reproductive success (χ^2_1= 16.931; p < 0.001, Fig. 1), and a marginally non-significant
temperature effect (χ^2_1= 3.712; p= 0.054). Running models separately for each temperature
level, mating system affected lifetime reproductive success at 20°C (χ^2_1= 4.575; p= 0.034),
and at 24°C (χ^2_1=12.569; p < 0.001), but not at 28°C (χ^2_1= 2.424; p= 0.122). The decrease
in LRS in polyandry vs monogamy peaked at 24°C (H = 0.36) and was 1,6 (H = 0.22) and
3,4 times smaller (H = 0.10) at 20 and 28°C, respectively. Rate-sensitive fitness estimates
show that estimated population costs are dependent on background growth rates (Fig. 2),
and in general particularly accused in decreasing populations.

Early reproductive rate- We did not detect a significant temperature x mating system
interaction (χ^2_2= 0.690; p= 0.7080), or a significant main mating system effect (χ^2_1= 3.287;
p= 0.069), but as expected we did detect a significant main temperature effect (χ^2_2= 84.347;
p < 0.001; Fig. SI1).

Figure 1. Female lifetime reproductive success (mean ± SEM) across temperature and mating
system treatments. Second axis represents the model coefficient estimates (95% CI) for the
estimated variation in lifetime reproductive success that ensue with polyandry vs. monogamy, across
the different temperature treatments. Estimates are coefficients from GLMs fitted separately for the
three temperature treatments.

Figure 2. a) Lifetime reproductive success (mean ± SEM) for different population growth rates across temperature and mating system treatments, b) relative population fitness costs of polyandry vs. monogamy for different population growth rates for experimental temperatures. Rate-sensitive fitness estimates were modelled based on our LRS measures.

Reproductive senescence- We detected a marginally nonsignificant temperature x mating
system effect for reproductive ageing (i.e., number of offspring during weeks 1 and 2 minus
number of offspring during weeks 3 and 4) (χ^2_2= 5.599; p= 0.060; Fig. SI1), and clear main
significant effects for both temperature (χ^2_2= 76.015; $p < 0.001$) and mating system (χ^2_1=
58.760; $p < 0.001$). Running models separately for each temperature level, mating system
affected reproductive ageing at 20°C (χ^2_1= 12.107; $p < 0.001$; coefficient = 7.9 with a 95%
CI of 3.49 to 12.49), at 24°C (χ^2_1= 35.324; $p < 0.001$; coefficient = 17.2 with a 95% CI of
11.53 to 22.88), and at 28°C (χ^2_1= 14.082; $p < 0.001$; coefficient = 11.8 with a 95% CI of
5.67 to 18.08).

Actuarial senescence- Survival analysis by means of a Cox proportional hazards model
detected a significant temperature x mating system interaction for female survival (χ^2_2=
10.303; p= 0.005; Fig. 3), as well as main temperature (χ^2_2= 76.801; $p < 0.001$) and mating
system effects (χ^2_1= 98.740; $p < 0.001$). Running models separately for each temperature
level, mating system affected female survival at 20°C (χ^2_1= 43.426; $p < 0.001$; exponentiated
coefficient = 1.87 with a 95% CI of 1.55 to 2.26), at 24°C (χ^2_1= 40.068; $p < 0.001$;
exponentiated coefficient = 1.83 with a 95% CI of 1.52 to 2.22), and at 28°C (χ^2_1= 18.216; p
< 0.001; exponentiated coefficient = 1.46 with a 95% CI of 1.23 to 1.74).

144
Figure 3. Male harm effect on female lifespan across temperature and mating system treatments.

<u>Reproductive behaviour</u>

The interaction between temperature and mating system was significant for aggression rate

(χ^2_2= 15.043; p < 0.001), and we also detected main temperature (χ^2_2= 15.062; p < 0.001)

and mating system effects (χ^2_1= 101.946; p < 0.001; Fig. 4a). Running models separately

for each temperature level, mating system affected aggression rate at 20° (χ^2_1= 6.912; p=

0.008), at 24° (χ^2_1= 28.858; p < 0.001) and at 28° (χ^2_2= 32.311; p < 0.001). Likewise, we

detected a significant temperature x mating system effect for courtship rate (i.e., number of

courtships experienced by the female per hour) (χ^2_2= 15.915; p < 0.001), and clear main

effects for both temperature (χ^2_2= 19.887; p < 0.001) and mating system (χ^2_1= 41.284; p <

0.001; Fig. 4b). Running models separately for each temperature level, mating system did

not affect courtship rate at 20° (χ^2_1= 0.367; p= 0.545) but did at 24° (χ^2_1= 21.787; p < 0.001)

and at 28° (χ^2_1= 40.136; p < 0.001). We also detected a significant interaction for female

rejection rate (χ^2_2= 6.422; p= 0.040), as well as main temperature (χ^2_2= 18.791; p < 0.001)

and mating system effects (χ^2_1= 25.561; p < 0.001; Fig. 4c). Running models separately for

each temperature level, mating system did not affect female rejection rate at 20° (χ^2_1= 0.201;

p= 0.653) but did at 24° (χ^2_1= 10.917; p < 0.001) and at 28° (χ^2_1= 19.27; p < 0.001).

<u>Ejaculate effects on female reproduction</u>

Female receptivity- The duration of the first mating in our female receptivity assays exhibited

a clear sperm competition risk level x treatment duration interaction (χ^2_1= 5.947; p = 0.015),

sperm competition risk level x temperature interaction (χ^2_2= 8.544; p = 0.014, Fig. 5a), and

main effects for treatment duration (χ^2_1= 6.057; p = 0.014) and sperm competition risk level

(χ^2_1= 36.738; p < 0.001). We found no significant main effect of temperature (χ^2_2= 4.260; p

= 0.119). Running models separately for each temperature level, sperm competition risk

level affected the duration of the first mating at 20°C (χ^2_1= 3.993; p = 0.045), at 24°C

(χ^2_1=10.653; p = 0.001), and at 28°C (χ^2_1= 26.529; p < 0.001). Similarly, we detected a

significant temperature x treatment duration interaction (χ^2_2= 6.593; p = 0.037), and a

marginally non-significant sperm competition risk level x temperature interaction (χ^2_1= 5.423;

p = 0.066; Fig. 5b) in female remating latency. We also detected significant main effects for

treatment duration (χ^2_1= 4.026; p = 0.044), sperm competition risk level (χ^2_1= 4.494; p =

0.034) and temperature (χ^2_2= 9.408; p = 0.009). Running models separately for each

temperature level, sperm competition risk level only affected remating latency at 28°C

(χ^2_1=8.052; p = 0.004). At 20°C we found no significant sperm competition risk level effect

(χ^2_1=0.950; p = 0.329), and the same for 24°C (χ^2_1=0.078; p = 0.779).

Female fecundity and survival- For the number of eggs produced by females during the
three first days, we did not detect any significant interaction between factors: sperm
competition risk level x treatment duration (χ^2_1= 0.039; *p*= 0.843), sperm competition risk
level x temperature (χ^2_2= 1.911; *p*= 0.384), temperature x treatment duration (χ^2_2= 2.787; *p*=
0.248), neither main significant effect for temperature (χ^2_2= 2.742; *p*= 0.253) nor for sperm
competition risk level (χ^2_1= 0.001; *p*= 0.968), but we did for treatment duration (χ^2_1= 9.009;
p= 0.002). Likewise, for the total number of offspring produced by females during the days
1, 2, 3, 4, 5, and 8 after mating, we did not find any significant interaction between factors:
male competition level x treatment duration (χ^2_1= 0.034; *p* = 0.852), male competition level
x temperature (χ^2_2= 0.520; *p* = 0.770), temperature x treatment duration (χ^2_2= 0.972; *p* =
0.614). Total offspring did not differ significantly between temperatures (χ^2_2= 2.142; *p* =
0.342), nor between sperm competition risk levels (χ^2_1= 0.326; *p* = 0.567), but did differ
between treatment durations (χ^2_1= 5.490; *p* = 0.019). Finally, survival analysis by means of
a Cox proportional hazards model did not detect any significant interaction between factors:
sperm competition risk level x treatment duration (χ^2_1= 0.012; *p* = 0.912), sperm competition
risk level x temperature (χ^2_2= 0.463; *p* = 0.793), temperature x treatment duration (χ^2_2=
2.190; *p* = 0.334). However, we detected a significant main temperature (χ^2_2=10.189; *p* =
0.006) and treatment duration effects (χ^2_1= 114.030; *p* < 0.001). We did not detect a
significant main sperm competition risk level effect (χ^2_1= 3.255; *p* = 0.071).

DISCUSSION

We show that male harm exhibits complex plasticity in response to temperature changes
within an optimal reproductive range (20ºC-28ºC) for a wild *D. melanogaster* population, with
several implications for our understanding of how sexual conflict unfolds in nature. First, we
show that net harm to females varies markedly within this range, such that relative harm is
maximal at 24ºC (H = 0.36), decreases at 20ºC (H = 0.22) and is not statistically detectable
at 28ºC (H = 0.10). Second, rate-sensitive fitness estimates predict that population costs are
dependent on population demography, so that male harm impacts population growth
differently at cold vs. warm temperatures depending on whether populations are decreasing
or growing. Third, our results strongly suggest that the male harm effect on population
growth has to do with the fact that different mechanisms exhibit qualitatively different
reaction norms in response to temperature, with distinct effects on different female fitness
components. Thus, cold temperatures decreased courtship intensity (i.e., male harassment;
pre-copulatory mechanism) and impacted female fitness mainly via accelerated actuarial

senescence. In contrast, warm temperatures modulated male ejaculate effects on female
reproduction (i.e., receptivity), and impacted female fitness mainly via accelerated
reproductive senescence. We discuss how such plasticity may affect how male harm
impacts populations, as well as selection, adaptation and, ultimately, evolutionary rescue
under a warming climate.

Temperature effects on male harm and its consequences for populations

We found that temperature variation within the optimal reproductive range for our study
population in the wild had a strong effect on male harm levels. To gauge male harm, we
used the standard procedure of comparing female LRS in monogamy, which imposes low
male-male competition and limits sexual conflict, vs. polyandry (i.e., a female with three
males), which imposes high male-male competition and intensifies sexual conflict between
the sexes (Yun et al., 2021). Importantly, a recent study suggests that these sex ratios are
common in mating patches in the wild, and are actually representative of the extremes in
natural levels of male-male competition (Dukas, 2020). In monogamy, temperature did not
affect female fitness (Fig. 1), showing that female reproduction is indeed optimal within this
range. In contrast, the net decrease in female LRS in polyandry vs. monogamy was highly
dependent on the thermal environment, with an average decrease of H = 0.36 at 24ºC, H =
0.22 at 20ºC and H = 0.10 at 28ºC, at which temperature we did not find a statistically
significant effect of mating system on female LRS (Fig. 1).

Male harm effects are expected to be cumulative over time, so that their impact on female
survival and reproductive output is likely not constant across female lifespan (Bonduriansky
et al., 2008; Filice et al., 2020). At the same time, early vs. late life reproduction weigh
differently on both individual fitness and how this impacts background population growth,
depending on whether such population is decreasing, stable or growing (Edward et al., 2011;
Priest et al., 2008). Thus, in order to evaluate how male harm is likely to impact populations
across different temperatures, we calculated rate-sensitive fitness estimates for individual
(Fig. 2a) and population (Fig. 2b) fitness across a range of demographic scenarios (i.e.,
decreasing, stable and growing populations; Edward et al., 2011). Overall, the impact of
male harm was higher in decreasing populations, where late-life reproduction gains
importance, which is consistent with the idea that male harm effects are cumulative. Above
and beyond this general effect, the observed interaction between temperature and female
fitness is maintained irrespective of population demography. That is, male harm decreases

female individual fitness more at 24ºC than at 20ºC and 28ºC. Interestingly, though,
temperature also has a clear effect on how demography affects population-level costs. At
hot temperatures the relative population costs of male harm vary considerably less with
demography (i.e., background population growth) compared to colder temperatures (Fig.
2b). For a decreasing population, the relative population costs of male harm are significantly
higher at 20ºC than at 28ºC, but this difference wanes as population growth rate (r)
increases, to the point of reverting in a rapidly growing population (Fig. 2b). This suggests
that male harm has more impact on late-life female fitness at cold temperatures, and hints
at the possibility that cold vs. hot temperature affect qualitatively distinct parameters of
female fitness, and underlying mechanisms of male harm.

Looking at effects on separate female fitness components yields results largely in agreement
with the above ideas. Again, consistent with the fact that harm needs to accumulate in time
to impact female fitness, temperature had no effect on how or whether male competition
impacted early reproductive rate. Temperature did, however, modulate how male harm
impacted on female actuarial vs. reproductive ageing. We found clear differences in female
lifespan across temperature and mating systems treatments. Male harm effects on actuarial
ageing (an increase in mortality rate with age) were more severe at 20°C (35% decrease in
female lifespan) than at 24°C (31% decrease) and at 28°C (22% decrease; Fig. 3). In
contrast, while male harm accelerated reproductive senescence at all temperatures, this
decrease was more marked at 24°C (coefficient = 17.2 with a 95% CI of 11.53 to 22.88) and
at 28°C (coefficient = 11.8 with a 95% CI of 5.67 to 18.08) than at 20°C (coefficient = 7.9
with a 95% CI of 3.49 to 12.49; Fig. SI1). In accordance with available evidence in lab-
adapted flies (García-Roa et al., 2019), these results show that temperature does not seem
to have a linear effect on ageing processes.

The above results open up the intriguing possibility that warm climates may lessen the
impact of sexual conflict on population viability, perhaps facilitating evolutionary rescue.
Male harm effects were found to be relatively lower in warmer temperatures and in
decreasing populations, precisely the type of context that would be typical of a climate-
change scenario. Similar results have been reported previously but in response to stressful
temperatures. For instance, temperature has been used to induce environmental stress in
natural populations of seed beetles (*Callosobruchus maculatus*), showing that a stressful
thermal regime reduces intra-locus sexual conflict by aligning selection in males and females
(Berger et al., 2014, but see Martinossi-Allibert et al., 2019). A previous study in a lab-

adapted population of *D. melanogaster* suggests male harm levels decrease when subject

to maladaptive warm temperatures (García-Roa et al., 2019). A further consequence of the

temperature sensitivity of male harm is that its net fitness consequences might be

considerably lower than expected from past research when considered in its natural thermal

setting. Our work thus joins an increasing number of recent studies in highlighting the

importance of evaluating more ecologically realistic scenarios in sexual conflict research,

particularly how natural fluctuations in the ecology of the socio-sexual context may affect

sexual conflict processes (García-Roa et al., 2019, 2020; Gomez-Llano et al., 2018; Yun et

al., 2017).

Temperature effects on sexual conflict mechanisms in Drosophila melanogaster

Prior to mating, *D. melanogaster* males harm females via sexual harassment, due to

protracted courtship of one or several males that results in physical damage, interference

with other behaviours (e.g., foraging or egg-laying) and costly energetic investment into male

avoidance (e.g., female rejection) (Bretman & Fricke, 2019; Partridge & Fowler, 1990; Teseo

et al., 2016). Importantly, previous studies in this species have shown that male harm is

directly related to courtship intensity and female rejection, and indirectly to male-male

aggression as a direct measure of male intrasexual competition (e.g., Bretman & Fricke,

2019; Carazo et al., 2014; Partridge & Fowler, 1990). In our study, we found a clear increase

in both courtship intensity and female rejection in polyandry vs. monogamy, but this was

largely dependent on the thermal environment (Fig. 4b-c). While we found clear evidence

that harassment increases at polygamy at 28°C, and 24°C, we did not find a similar effect

at 20°C (Fig. 4b). Female rejection behaviours exhibited the exact same trend (Fig. 4c).

Thus, our results suggest that male harassment decreases drastically at cold temperatures

and is perhaps maximal at warm temperatures, at which temperatures we also detected the

highest level of male-male aggression (Fig. 4a). The above results seem to suggest that the

decrease in male harm to females that we detected at 20ºC vs. 24ºC (Fig. 1) is likely

explained by a drastic drop in male harassment of females at this temperature.

Figure 4. Reproductive behaviours (mean ± SEM) across temperature and mating system treatments. a) aggressions male-male per hour, b) courtships per female per hour, and c) female rejections per hour. Second axis represents the model coefficient estimates (95% CI) for the estimated variation in aggression, courtship and female rejection rate that ensue with polyandry vs. monogamy, across the different temperature treatments. Estimates are coefficients from GLMs fitted separately for the three temperature treatments.

During mating, *D. melanogaster* males transfer seminal fluid proteins that manipulate female re-mating and egg-laying rates to their advantage, but at a costs to females in terms of lifespan and lifetime reproductive success (Chapman, Bangham, et al., 2003b; Chapman et al., 1995; Sirot et al., 2009; Wigby & Chapman, 2005). Furthermore, these effects are known to be modulated by the socio-sexual context, such that males strategically adjust their investment in SFPs depending on expected sperm competition risk (SCR) levels (Bretman, Fricke, & Chapman, 2009; Hopkins et al., 2019; Wigby et al., 2009). We run a series of assays to investigate whether temperature modulates any of the known phenotypic effects that SFPs have on females following a single mating (note these assays are standard for *D. melanogaster*; e.g., Chapman et al., 1995; Perry, Sirot, & Wigby, 2013; Wigby & Chapman, 2005; Wigby et al., 2009) with males that were exposed to low (i.e., kept alone in their vial) or high sperm competition risk (i.e., kept with 7 other males in a vial). The temperature males were kept at prior to mating did not modulate how mating affected short-term female fecundity or lifespan, but we did detect a clear effect of temperature with respect female receptivity. Generally, reception of a male ejaculate resulted in a sharper decrease in female receptivity (i.e., longer remating latency) when males perceived a high SCR, which is to be expected and is in accordance with previous studies (Bretman et al., 2009; Bretman, Fricke, Hetherington, Stone, & Chapman, 2010; Denis et al., 2017). However, this effect was largely modulated by temperature, such that it was only clearly detected at 28ºC (Fig. 5b). Interestingly, this effect was consistent for males treated for 48h and 13 days and was paralleled by mating duration (Fig. 5a). Namely, males exposed to high SCR generally mated for longer than males exposed to low SCR, again in line with previous findings (Bretman et al., 2010; Bretman, Westmancoat, Gage, & Chapman, 2013), but this difference was clearly larger at 28ºC (Fig. 5a). It is worth restating that, under monogamous conditions, we did not detect a decrease in female productivity at 28ºC vs 24ºC and 20ºC, so that general differences in sperm viability are unlikely to account for the aforementioned differences. All in all, our results suggest that at least some post-copulatory harm mechanisms are sensitive to temperature, and that receipt of a male ejaculate results in a sharper decrease of female receptivity at warm temperatures. We speculate that this may contribute to explain why male harm drops so sharply at 28ºC despite the fact that male harassment and male-male competition seems to be maximal at this temperature. In *D. melanogaster*, as in many other species, repeatedly mating is costly for females in terms of lifetime reproductive success (Arnqvist & Nilsson, 2000; Chapman et al., 1995; Fowler &

344 Partridge, 1989). An intriguing possibility is thus that SFPs are more effective at lowering
345 female re-mating rates at warm temperatures, thereby buffering these costs. We suggest
346 future studies should look into this by examining in detail how temperature affects the
347 composition and transfer of SFPs to females.

349
350

Figure 5. a) Mating duration of males exposed to a high (8 males per vial) or low sperm competition
risk (1 male per vial) level 48 hours and 13 days before mating across temperature treatments, b)
females remating latency following a single mating with either a male from a high or low sperm
competition risk level 48 hours and 13 days before mating across temperature treatments. Second
axis represents the model coefficient estimates (95% CI) for the estimated variation in mating duration
and remating latency that ensue with high vs. low sperm competition risk, across the different
temperature treatments. Estimates are coefficients from GLMs fitted separately for the three
temperature treatments.

Conclusions

Our findings may have implications for our understanding of how sexual conflict unfolds in
nature, and its consequences for populations. First, they add to growing evidence (Gomez-
Llano et al., 2018; MacPherson et al., 2018; Malek & Long, 2019; Perry & Rowe, 2018; Yun
et al., 2017) indicating that ecological context is key in shaping sexually antagonistic
coevolution and, in particular, suggest that temperature may be a particularly salient
ecological factor to understand how sexual conflict evolves and operates in nature (García-
Roa et al., 2020). Second, they highlight that male harm mechanisms can be highly plastic
even in response to relatively minor fluctuations in temperature well within the optimal

reproductive range, and suggest that different harm mechanisms are differently affected by
temperature. Third, they suggest that male harm effects on female life-history and fitness
components are asymmetrically modulated by temperature; male harm particularly
decreased survival at cold and moderate temperatures, and reproductive ageing at
moderate and hot temperatures. In conjunction, these phenomena may have a bearing on
evolutionary rescue and local adaptation processes. For example, in maintaining genetic
variation in sexually selected traits in males, and/or in ameliorating the demographic impact
of sexual conflict in populations facing environmental change. In this sense, our results hint
at the intriguing possibility that rapid warming, such as human-induced climate change, may
have unexpected positive consequences via its effect on male harm, potentially facilitating
evolutionary rescue. We thus suggest two priorities for future research on this area are: a)
studying the effect of temperature on post-copulatory mechanisms (e.g., seminal fluid
accessory glands proteins and their effects in *Drosophila*), and b) developing formal models
to explore under what contexts might temperature effects on male harm facilitate
evolutionary rescue.

MATERIALS AND METHODS

Field Collection

In October 2018, we used banana traps to sample *Drosophila melanogaster* flies from three
wineries in Requena (Spain): "Hispano-Suizas" (39.466128, -1.149642), "Pago de Tharsys"
(39.497834, -1.122781) and "Dominio de la Vega" (39.515079, -1.143757). Traps were
setup within the wineries, but in premises that were open to the exterior and to ample
surrounding vineyards and/or in the vineyards themselves. After collection, we
anaesthetized flies using mild CO_2 exposure. We then separated and individually distributed
field-collected females in vials with standard food, left them to lay eggs for a period of 48
hours and then incubated their eggs at 24ºC, 60% humidity and a 12:12 dark-light cycle for
14 days to allow adult flies to emerge. We inspected the genital arch of F1 males of each of
these female isolines to distinguish *D. melanogaster* isolines from *D. simulans*. We then
collected three males and three females from each *D. melanogaster* isoline (total of 276 flies
from 46 isolines) and released them into a population cage with a surplus of food medium
supplemented with live yeast, setting up the "Vegalibre (VG)" population. In November 2019
and October 2020, to maintain natural variation, we re-sampled the wineries and added 348
and 756 flies from 58 and 126 isolines, respectively, following the same procedure (27

isolines from "Pago de Tharsys" and 31 isolines from "Dominio de la Vega" in November
2019 and 33 isolines from "Pago de Tharsys", 50 isolines from "Dominio de la Vega" and 43
isolines from "Hipano-Suizas" in October 2020).

Stock maintenance and acclimation

We carried out all experiments between March 2020 and April 2021, using individuals from
the VG field population kept in the laboratory with overlapping generations at an average
temperature of 24°C with daily fluctuations (±4°C) that mimic natural daily temperature
conditions during the reproductively active season, at ~60% humidity and on a 12:12-hr light:
dark cycle. We used maize-malt medium (7g of agar, 72 g of malt, 72g of maize, 9g of soya,
15g of live yeast, 20g of molasses, and 33ml of Nipagin mix (3g of methyl 4-hydroxy-
benzoate and 28ml of ethanol 90%) per 1000ml of water) throughout. We used the following
procedure to collect experimental flies. We introduced yeasted grape juice agar plates into
stock populations to induce female oviposition. We then collected eggs and placed them in
bottles containing ~75ml of medium to be incubated at 24 ± 4°C at a controlled density
(Clancy & Kennington, 2001). We collected virgin flies within 6h of eclosion, under ice
anaesthesia, and then we sexed and kept them in vials with food until their use (~3 days),
at 24 ± 4°C.

Fitness assays: quantifying male harm

To study whether male harm is affected by temperature, we established a factorial design
to measure survival and lifetime reproduction success (LRS) of female flies under
monogamy (i.e., one male and one female per vial) vs. polyandry (i.e., three males and one
female per vial), across three stable temperature treatments typical of this population during
their reproductively active period in the wild: 20, 24, and 28°C. Comparison of female fitness
at monogamy vs. polyandry is a common way to gauge male harm in *Drosophila* and other
organisms (Yun, Agrawal, & Rundle, 2021), and our treatments reflect the low and high-end
of the spectrum of sex ratios that are typical of *D. melanogaster* at mating patches in the
wild (Dukas, 2020).

We first randomly divided virgin flies into three groups that we allocated to the three different
stable temperature treatments 48 hours before starting the experiment. Flies remained at
those temperatures until the end of the experiment. To estimate LRS, we transferred flies to
fresh vials twice a week using mild CO_2 exposure anaesthesia. We incubated the vials

containing female eggs at 24 ± 4°C for 15-20 days (~15 days for 28°C, ~17 days for 24°C

and ~20 days for 20°C) to allow F1 offspring emergence, after which we froze them at -21°C

for later counting. The differences in incubation time are due to differences in developmental

time because of temperature differences in the first 3-4 days of each vial (i.e., before flipping

females to new fresh vials). We discarded and replaced males by young (2 to 4 days old)

virgin males three weeks after starting (at the same time for all treatments). We kept a stock

of replacement males maintained at each of the three temperatures (20, 24 and 28°C) to

replace dead male flies if needed. We kept flies under these conditions during six weeks,

after which we discarded males and followed females until they died. We started the

experiment with 468 females (78 per each temperature x mating system treatment) and 936

males (234 per each temperature x polyandry and 78 per each temperature x monogamy).

Final sample sizes were: a) at 20°C: 74 females for polyandry and 76 females for

monogamy; b) at 24°C: 72 females for polyandry and 77 females for monogamy; c) at 28°C:

70 females for polyandry and 75 females for monogamy. Differences between start and final

sample sizes across treatments are due to discarded/escaped flies since the start of the

experiment. We estimated the overall degree of male harm by calculating the relative harm

(H) following Yun et al. (2021):

$$H = \frac{W_{monogamy} - W_{polygamy}}{W_{monogamy}}$$

Finally, and using the data collected above, we partitioned overall effects on LRS into effects

of temperature x mating system (i.e., male-male competition level) on early reproductive rate

(i.e., offspring produced during the first two weeks of age) and reproductive (i.e., offspring

produced over weeks 1-2 vs. 3-4) and actuarial senescence (i.e., survival).

Behavioural assays: quantifying male-male aggression and harassment of females

Immediately after the fitness experiment started, we conducted behavioural observations on

the first day of the experiment across all treatments, to investigate the behavioural

mechanisms that might underlie the potential fitness effects evaluated above. Due to logistic

limitations, we conducted behavioural observations in the same temperature control room,

so we had to conduct trials at 20, 24 and 28°C in three consecutive days (with both

monogamy and polyandry treatments evaluated at the same time for each temperature), in

randomized order (i.e., 20, 28 and 24°C). However, all flies were 5 days-old at the start of

the experiment. We measured the following behaviours: (a) courtship intensity (number of

courtships experienced by a female per hour), (b) male-male aggression behaviours and (c)
female rejection behaviours (Bastock & Manning, 1955; Connolly & Cook, 1973). We also
recorded the number of total matings during the observation period.

Observations started at lights-on (10 a.m.) and lasted for 8 hr, during which time we
continuously recorded reproductive behaviours using scan sampling of vials. Each complete
scan lasted approximately 8 min, so that we always conducted one complete scan every 10
min to ensure recording all matings (see below). Scans consisted in observing all vials in
succession for ca. 3s each and recording all occurrences of the behaviours listed above.
We interspersed these behavioural scans with very quick (<1 min) mating scans where we
rapidly swept all vials for copulas at the beginning, in the middle and at the end of each
behavioural scan. This strategy ensured that we recorded all successful matings (>10min),
which typically last between 15 and 25min in our population of *D. melanogaster*, during our
8-hr observation. We obtained a total of 49 scans per vial. Behavioural observations were
conducted only once, on day 1 of the fitness experiment, as prior experiments have shown
that courtship, aggressive and female rejection behaviours are stable over time in *D.*
melanogaster, so that our behavioural indexes are representative of long-term treatment
differences (e.g., Carazo, Perry, Johnson, Pizzari, & Wigby, 2015; Carazo, Tan, Allen,
Wigby, & Pizzari, 2014). In contrast to courtship and aggression indexes, note that total
mating frequency over the first day cannot be taken as a reliable measure of mating rate
(Wolfner, 1997), and thus our rationale in recording this variable was just to ensure that early
mating ensued normally across treatments (which was the case, see SI2).

Female reproduction and survival assays: quantifying male ejaculate "toxicity"

To examine post-mating mechanisms that might underlie the fitness effects observed in our
first experiment, we conducted four additional assays to test whether temperature modulates
the previously well-documented effects of male ejaculates on female reproduction and
survival in *D. melanogaster*. Briefly, males of this species transfer seminal fluid proteins
(SFPs) produced by their accessory glands that increase short-term female fecundity, as
well as decrease female receptivity and survival (Chapman, Liddle, Kalb, Wolfner, &
Partridge, 1995; Wigby & Chapman, 2005). In addition, prior studies have shown that males
are able to tailor investment into SFPs according to the expected sperm competition risk and
intensity (Hopkins et al., 2019). Thus, we set-up a factorial design where we manipulated
the temperature (i.e., 20, 24 and 28ºC) and perceived sperm competition risk levels (i.e.,

males kept alone vs. with 7 more males in a vial) at which adult males were kept prior to mating, for two temperature treatment durations (i.e., 48 hours or 13 days), and then measured how reception of their ejaculate in a common garden environment (i.e., 24ºC) affected female fecundity, survival and reproduction.

Receptivity assays- We first collected experimental males as virgins (i.e., within 6 h of eclosion) under ice anesthesia and randomly placed them either individually or in a same-sex group of 8 in medium containing plastic vials. Next, we randomly divided them into three groups that we allocated to the different stable temperature treatments for either 48 hours (experiment 1) or 13 days (experiment 2) immediately before the beginning of each experiment. In experiment 2 (treatment duration of 13 days), we emptied the seminal fluid of experimental males before we allocated them to the different temperature / competition treatments. To do so, we housed individually experimental males with four standard virgin females for 24 hours. This strategy ensured that spermatogenesis took place over the 13 days under the temperature / competition treatments.

We collected all females and competitor males used in receptivity assays as virgins and held them in groups of 15 to 20 flies at 24 ± 4°C. Experiments started by exposing all virgin females to single experimental males for 2.5 h at 24°C. After a successful copulation, we separated the mated females from the males and kept them individually in medium containing vials until the next mating trial. We discarded unmated females and experimental males. 72 h after the first mating, we individually exposed females to single virgin competitor males for 12 h. After each trial, we transferred unmated females into a new medium containing vial, until the next mating assay 24 h later. We repeated remating trials every 24 h for three consecutive days. We calculated the cumulative percentage of remated females for each of the three days of each experiment. We conducted the experiments in two blocks each: with n = 390 females for each batch in experiment 1 (n = 436 rematings) and n = 420 females for each batch in experiment 2 (n = 676 rematings). We also recorded mating duration for the first mating and mating latency (i.e., time between males being introduced into the female-containing vial and copulation) and mating duration for re-matings. Females that did not remate within those three days were right-censored. Females and all experimental males were 4 days old at the start of experiment 1. In experiment 2, females and "re-mating" competitor males were 4 days old, while experimental males were 18 days old.

Fecundity and survival assays- To gauge effects on female short-term fecundity and long-
term survival, we performed two experiments (experiments 3 and 4) where we compared
the oviposition and egg fertility of females mated with male flies subject to the same factorial
design imposed in receptivity experiments. We collected and treated all experimental males
as in the receptivity assays described above, and then proceeded to mate ~4 day-old virgin
females in single pairs to either 4- (experiment 3, 48h temperature treatment duration) or
18-day old experimental males (experiment 4, 13d temperature treatment duration) for 2.5h
at 24°C. After copulation, we separated mated females from males and kept them
individually in medium containing vials. We discarded unmated females. We then transferred
females to fresh vials every 24h for 4 days, and then every 3 days twice. Later, we
transferred female flies to fresh vials once a week, and we combined vials to maintain
density at ten flies per vial. We kept female flies under these conditions until they died. We
removed dead flies at each transfer and recorded female deaths. We counted eggs laid the
first 3 days and vials from days 1, 2, 3, 4, 5, and 8 after mating were retained to count
progeny to determine egg viability. We tested 545 females at the starting point of the
experiment 3, and 480 females at the starting point of the experiment 4.

Statistical analyses

We performed all statistical analyses using R statistical software (version 3.5.2). In all
models, we assessed fit by visual inspections of diagnostic plots on raw and residual data
(Zuur, Ieno, & Elphick, 2010). To examine temperature effects on male harm, we evaluated
the interaction between mating system and temperature on female fitness (LRS), survival
and male and female reproductive behaviours. We fitted generalized linear models (GLMs)
with temperature, mating system and their interaction as fixed effects. We also used Cox
proportional hazards survival model to analyse potential differences in mortality risk across
treatments, using the *survival* and *survminer* packages, including the females lost during
manipulations as "right censored" individuals (i.e., individuals that are taken into account for
demographic analysis until the day they disappear, Kleinbaum & Klein, 2012). We also
censored any remaining females after the six weeks of the experiment. Graphical inspection
of LRS and reproductive behaviours (courtship intensity, intersexual aggression behaviours
and female rejection) revealed that the normality assumption was apparently violated, as
well as the independence assumption for LRS. Box–Cox transformation (Quinn & Keough,
2002) solved these problems and allowed us to run a GLM with a Gaussian error distribution.
In all these afore mentioned cases, fitted models were subsequently validated. In addition,

we compared GLMs with their corresponding null GLMs using likelihood ratio test. In all
models, we used ANOVA type III to compute p-values. We detected a problem of collinearity
between mating system and the interaction in the LRS model. We fitted the model again
without the main mating system effect (which was not our main interest) and also ran models
separately for each temperature level. As a complementary analysis, we ran a model with
temperature as factor with a quadratic contrast table predetermined (note that the relation
between LRS and temperature does not seem to be linear, Fig 1), obtaining similar results.
For the total matings recorded across the 8h observation period (i.e., mating rate), we initially
used a GLM with Poisson errors. However, the dataset contained many zero values. We
therefore analysed the data using a Hurdle model, in which the zero values are modelled
separately from the nonzero values (Zuur et al., 2010). Additionally, we compared GLM and
Hurdle models using an information-theoretic approach based on Akaike's Information
Criterion (AIC). The minimum AICc value indicates the best supported model given the
trade-off between fit to the data and model complexity (Konishi & Kitagawa, 2008). To
estimate how male harm impact estimated population growth across varying demographic
scenarios (i.e., decreasing, stable and growing populations), based on our LRS measures,
we calculated rate-sensitive fitness estimates (Edward, Fricke, Gerrard, & Chapman, 2011)
against different population background growth rates of r = -0.1, r = -0.05, r = 0, r = 0.05 and
r = 0.1.

For receptivity (mating duration and remating latency) and fecundity (total adults) analyses,
we fitted generalized linear models (GLMs) with temperature, sperm competition risk level,
treatment duration and their interaction as fixed effects. For egg production data, we fitted a
generalized linear mixed model (GLMM) with temperature, sperm competition risk level,
treatment duration and their interaction as fixed effects and day number as a random effect,
with a zero inflated distribution, in which the zero values are modelled separately from the
nonzero values (Zuur et al., 2010). We assessed significance of factors by dropping
individual terms from the full model using the "drop1" function, refitting where the interaction
was nonsignificant. We also used Cox proportional hazards survival model to analyse
potential differences in mortality risk across treatments, using the *survival* and *survminer*
packages, including the females lost during manipulations as "right censored" individuals.
Finally, we also ran models separately for each temperature level for the cases in which a
significant interaction between sperm competition risk level and temperature treatment was
detected.

ACKNOWLEDGEMENTS

We thank Alejandro Hita for assistance with the experimental procedure.

COMPETING INTERESTS

PC was supported by a research grant (PID2020-118027GB-I00) funded by MCIN/AEI/ 10.13039/501100011033. RGR was supported by a Ministerio de Educación Cultura y Deportes postdoc grant grant (Juan de la Cierva Formación FJC2018-037058-I). CLN was supported by a MINECO predoctoral (PRE2018-084009) grant from the Spanish Government.

REFERENCES

Arbuthnott, D., Dutton, E. M., Agrawal, A. F., & Rundle, H. D. (2014). The ecology of sexual conflict: Ecologically dependent parallel evolution of male harm and female resistance in Drosophila melanogaster. *Ecology Letters*, *17*(2), 221–228. https://doi.org/10.1111/ele.12222

Arnqvist, G., Edvardsson, M., Friberg, U., & Nilsson, T. (2000). Sexual conflict promotes speciation in insects. *Proceedings of the National Academy of Sciences of the United States of America*, *97*(19), 10460–10464. https://doi.org/10.1073/pnas.97.19.10460

Arnqvist, G., & Nilsson, T. (2000). The evolution of polyandry: Multiple mating and female fitness in insects. *Animal Behaviour*, *60*(2), 145–164. https://doi.org/10.1006/anbe.2000.1446

Arnqvist, G., & Rowe, L. (2005). *Sexual Conflict*. Princeton University Press, Princeton, NJ.

Arnqvist, G., & Tuda, M. (2010). Sexual conflict and the gender load: Correlated evolution between population fitness and sexual dimorphism in seed beetles. *Proceedings of the Royal Society B: Biological Sciences*, *277*(1686), 1345–1352. https://doi.org/10.1098/rspb.2009.2026

Bastock, M., & Manning, A. (1955). The courtship behaviour of Drosophila melanogaster. *Behaviour*, *8*, 85–111.

Berger, D., Grieshop, K., Lind, M. I., Goenaga, J., Maklakov, A. A., & Arnqvist, G. (2014). Intralocus sexual conflict and environmental stress. *Evolution*, *68*(8), 2184–2196. https://doi.org/10.1111/evo.12439

Berger, D., Martinossi-Allibert, I., Grieshop, K., Lind, M. I., Maklakov, A. A., & Arnqvist, G. (2016). Intralocus sexual conflict and the tragedy of the commons in seed beetles. *American Naturalist*, *188*(4), E98–E112. https://doi.org/10.1086/687963

Bonduriansky, R. (2011). Sexual selection and conflict as engines of ecological diversification. *American Naturalist*, *178*(6), 729–745. https://doi.org/10.1086/662665

Bonduriansky, R., & Chenoweth, S. F. (2009). Intralocus sexual conflict. *Trends in Ecology and Evolution*, *24*(5), 280–288. https://doi.org/10.1016/j.tree.2008.12.005

Bonduriansky, R., Maklakov, A., Zajitschek, F., & Brooks, R. (2008). Sexual selection,
sexual conflict and the evolution of ageing and life span. *Functional Ecology*, *22*(3),
443–453. https://doi.org/10.1111/j.1365-2435.2008.01417.x

Bretman, A., & Fricke, C. (2019). Exposure to males, but not receipt of sex peptide,
accelerates functional ageing in female fruit flies. *Functional Ecology*, *33*(8), 1459–
1468. https://doi.org/10.1111/1365-2435.13339

Bretman, A., Fricke, C., & Chapman, T. (2009). Plastic responses of male Drosophila
melanogaster to the level of spermcompetition increase male reproductive fitness.
Proceedings of the Royal Society B: Biological Sciences, *276*(1662), 1705–1711.
https://doi.org/10.1098/rspb.2008.1878

Bretman, A., Fricke, C., Hetherington, P., Stone, R., & Chapman, T. (2010). Exposure to
rivals and plastic responses to sperm competition in Drosophila melanogaster.
Behavioral Ecology, *21*(2), 317–321. https://doi.org/10.1093/beheco/arp189

Bretman, A., Westmancoat, J. D., Gage, M. J. G., & Chapman, T. (2013). Costs and
benefits of lifetime exposure to mating rivals in male drosophila melanogaster.
Evolution, *67*(8), 2413–2422. https://doi.org/10.1111/evo.12125

Carazo, P., Perry, J. C., Johnson, F., Pizzari, T., & Wigby, S. (2015). Related male
Drosophila melanogaster reared together as larvae fight less and sire longer lived
daughters. *Ecology and Evolution*, *5*(14), 2787–2797.
https://doi.org/10.1002/ece3.1549

Carazo, P., Tan, C. K. W., Allen, F., Wigby, S., & Pizzari, T. (2014). Within-group male
relatedness reduces harm to females in Drosophila. *Nature*, *505*(7485), 672–675.
https://doi.org/10.1038/nature12949

Chapman, T., Arnqvist, G., Bangham, J., & Rowe, L. (2003a). Sexual conflict. *Trends in
Ecology and Evolution*, *18*(1), 41–47. https://doi.org/10.4161/fly.5.1.14459

Chapman, T., Bangham, J., Vinti, G., Seifried, B., Lung, O., Wolfner, M. F., … Partridge, L.
(2003b). The sex peptide of Drosophila melanogaster: Female post-mating responses
analyzed by using RNA interference. *Proceedings of the National Academy of
Sciences of the United States of America*, *100*(17), 9923–9928.
https://doi.org/10.1073/pnas.1631635100

Chapman, T., Liddle, L., Kalb, J., Wolfner, M. F., & Partridge, L. (1995). *Cost of mating in
Drosophila melanogaster females is mediated by male accessory gland products.
Nature* (Vol. 373). https://doi.org/10.1016/S0003-3472(05)81016-4

Clancy, D. J., & Kennington, J. (2001). A simple method to achieve consistent larval
density in bottle cultures. *Drosophila Information Service*, *84*(January), 168–169.

Connolly, K., & Cook, R. (1973). Rejection Responses By Female Drosophila
Melanogaster : Their Ontogeny, Causality and Effects Upon the Behaviour of the
Courting Male. *Behaviour*, *44*(1–2), 142–165.
https://doi.org/10.1163/156853973x00364

Cornwallis, C. K., & Uller, T. (2010). Towards an evolutionary ecology of sexual traits.
Trends in Ecology and Evolution, *25*(3), 145–152.
https://doi.org/10.1016/j.tree.2009.09.008

Crudgington, H. S., & Siva-Jothy, M. T. (2000). Genital damage, kicking and early death.
Nature, *407*(6806), 855–856. https://doi.org/10.1038/35038154

De Lisle, S. P., Goedert, D., Reedy, A. M., & Svensson, E. I. (2018). Climatic factors and
species range position predict sexually antagonistic selection across taxa.
Philosophical Transactions of the Royal Society B: Biological Sciences, *373*(1757).
https://doi.org/10.1098/rstb.2017.0415

Denis, B., Claisse, G., Le Rouzic, A., Wicker-Thomas, C., Lepennetier, G., & Joly, D.
(2017). Male accessory gland proteins affect differentially female sexual receptivity
and remating in closely related Drosophila species. *Journal of Insect Physiology*,
99(December 2016), 67–77. https://doi.org/10.1016/j.jinsphys.2017.03.008

Depeursinge, A., Racoceanu, D., Iavindrasana, J., Cohen, G., Platon, A., Poletti, P.-A., …
Chapman, T. (1997). Seminal Fluid Protein Allocation and Male Reproductive
Success. *Insect Biochemistry and Molecular Biology*, *27*(March 2005), 751–757.
https://doi.org/10.1016/S0965-1748(96)00084-7

Dukas, R. (2020). Natural history of social and sexual behavior in fruit flies. *Scientific
Reports*, *10*(1), 21932. https://doi.org/10.1038/s41598-020-79075-7

Edward, D. A., Fricke, C., Gerrard, D. T., & Chapman, T. (2011). QUANTIFYING THE
LIFE-HISTORY RESPONSE TO INCREASED MALE EXPOSURE IN FEMALE
DROSOPHILA MELANOGASTER. *Evolution*, *65*(2), 564–573.

Filice, D. C. S., Bhargava, R., & Dukas, R. (2020). Plasticity in male mating behavior
modulates female life history in fruit flies. *Evolution*, *74*(2), 365–376.
https://doi.org/10.1111/evo.13926

Fowler, K., & Partridge, L. (1989). A cost of mating in female fruitflies. *Nature*, *338*, 760–
761.

Fricke, C., Bretman, A., & Chapman, T. (2010). Female nutritional status determines the
magnitude and sign of responses to a male ejaculate signal in Drosophila
melanogaster. *Journal of Evolutionary Biology*, *23*(1), 157–165.
https://doi.org/10.1111/j.1420-9101.2009.01882.x

Fricke, C., Perry, J., Chapman, T., & Rowe, L. (2009). The conditional economics of
sexual conflict. *Biology Letters*, *5*(5), 671–674. https://doi.org/10.1098/rsbl.2009.0433

García-Roa, R., Garcia-Gonzalez, F., Noble, D. W. A., & Carazo, P. (2020). Temperature
as a modulator of sexual selection. *Biological Reviews*, *95*(6), 1607–1629.
https://doi.org/10.1111/brv.12632

García-Roa, R., García-González, F., Noble, D. W. A., & Carazo, P. (2021). Motherly love
curbs harm: maternal effects can modulate sexual conflict. Retrieved from
https://doi.org/10.32942/osf.io/nfmz8

García-Roa, Roberto, Chirinos, V., & Carazo, P. (2019). The ecology of sexual conflict:
Temperature variation in the social environment can drastically modulate male harm
to females. *Functional Ecology*, *33*(4), 681–692. https://doi.org/10.1111/1365-
2435.13275

Gavrilets, S. (2014). Is sexual conflict an "Engine of speciation"? *Cold Spring Harbor
Perspectives in Biology*, *6*(12), 1–13. https://doi.org/10.1101/cshperspect.a017723

Gomez-Llano, M. A., Bensch, H. M., & Svensson, E. I. (2018). Sexual conflict and ecology:
Species composition and male density interact to reduce male mating harassment
and increase female survival. *Evolution*, *72*(4), 906–915.
https://doi.org/10.1111/evo.13457

Holland, B., & Rice, W. R. (1999). Experimental removal of sexual selection reverses
intersexual antagonistic coevolution and removes a reproductive load. *Proceedings of*
the National Academy of Sciences of the United States of America, *96*(9), 5083–
5088. https://doi.org/10.1073/pnas.96.9.5083

Hopkins, B. R., Sepil, I., Thézénas, M. L., Craig, J. F., Miller, T., Charles, P. D., … Wigby,
S. (2019). Divergent allocation of sperm and the seminal proteome along a
competition gradient in Drosophila melanogaster. *Proceedings of the National*
Academy of Sciences of the United States of America, *116*(36), 17925–17933.
https://doi.org/10.1073/pnas.1906149116

Hopkins, Sepil, I., & Wigby, S. (2020). Structural variation in Drosophila melanogaster
spermathecal ducts and its association with sperm competition dynamics. *Royal*
Society Open Science, *7*(3). https://doi.org/10.1098/rsos.200130

Kapun, M., Barrón, M. G., Staubach, F., Vieira, J., & Darren, J. (2018). Genomic analysis
of European Drosophila populations reveals longitudinal structure and continent-wide
selection.

Kim, K. E., Jang, T., & Lee, K. P. (2020). Combined effects of temperature and
macronutrient balance on life-history traits in Drosophila melanogaster: implications
for life-history trade-offs and fundamental niche. *Oecologia*, *193*(2), 299–309.
https://doi.org/10.1007/s00442-020-04666-0

Kleinbaum, D. G., & Klein, M. (2012). *Survival Analysis: A Self-Learning Text* (Third Edit).
Springer.

Koene, J. M., & Schulenburg, H. (2005). Shooting darts: Co-evolution and counter-
adaptation in hermaphroditic snails. *BMC Evolutionary Biology*, *5*(February).
https://doi.org/10.1186/1471-2148-5-25

Kokko, H., & Brooks, R. (2003). Sexy to die for? Sexual selection and the risk of extinction.
Annales Zoologici Fennici, *40*(2), 207–219.

Konishi, S., & Kitagawa, G. (2008). Information Criterion. In *Information Criteria and*
Statistical Modeling (pp. 29–74). Springer New York.

Lange, R., Reinhardt, K., Michiels, N. K., & Anthes, N. (2013). Functions, diversity, and
evolution of traumatic mating. *Biological Reviews*, *88*(3), 585–601.
https://doi.org/10.1111/brv.12018

Le Galliard, J. ., Fitze, P. ., Ferriere, R., & Clobert, J. (2005). Sex ratio bias , male
aggression , and population collapse in lizards, *102*(50), 18231–18236.

MacPherson, A., Yun, L., Barrera, T. S., Agrawal, A. F., & Rundle, H. D. (2018). The
effects of male harm vary with female quality and environmental complexity in
Drosophila melanogaster. *Biology Letters*, *14*(8).
https://doi.org/10.1098/rsbl.2018.0443

Malek, H., & Long, T. (2019). Spatial environmental complexity mediates sexual conflict
and sexual selection in Drosophila melanogaster. *Ecology and Evolution*, *9*(5).
https://doi.org/10.1002/ece3.4932

Martinossi-Allibert, I., Thilliez, E., Arnqvist, G., & Berger, D. (2019). Sexual selection,
environmental robustness, and evolutionary demography of maladapted populations:
A test using experimental evolution in seed beetles. *Evolutionary Applications*, *12*(7),
1371–1384. https://doi.org/10.1111/eva.12758

Miler, K., Stec, D., & Czarnoleski, M. (2020). Heat wave effects on the behavior and life-
history traits of sedentary antlions. *Behavioral Ecology*, *31*, 1326–1333.
https://doi.org/10.1093/beheco/araa085

Miller, C. W., & Svensson, E. I. (2014). Sexual Selection in Complex Environments.
Annual Review of Entomology, (59), 427–445. https://doi.org/10.1146/annurev-ento-
011613-162044

Monteiro, N., Cunha, M., Ferreira, L., Vieira, N., Antunes, A., Lyons, D., & Jones, A. G.
(2017). Parabolic variation in sexual selection intensity across the range of a cold-
water pipefish: implications for susceptibility to climate change. *Global Change
Biology*, *23*(9), 3600–3609. https://doi.org/10.1111/gcb.13630

Myhre, L. C., Forsgren, E., & Amundsen, T. (2013). Effects of habitat complexity on mating
behavior and mating success in a marine fish. *Behavioral Ecology*, *24*(2), 553–563.
https://doi.org/10.1093/beheco/ars197

Parker, G. A. (1979). Sexual selection and sexual conflict. In M. . Blum & N. A. Blum
(Eds.), *Sexual Selection and Reproductive Competition in Insects* (Academic P, pp.
123–166).

Partridge, L., & Fowler, K. (1990). Non-mating costs of exposure to males in female
Drosophila melanogaster. *Journal of Insect Physiology*, *36*(6), 419–425.

Perry, J. C., & Rowe, L. (2018). Sexual conflict in its ecological setting. *Philosophical
Transactions of the Royal Society B: Biological Sciences*, *373*(1757).
https://doi.org/10.1098/rstb.2017.0418

Perry, J. C., Sirot, L., & Wigby, S. (2013). The seminal symphony: How to compose an
ejaculate. *Trends in Ecology and Evolution*, *28*(7), 414–422.
https://doi.org/10.1016/j.tree.2013.03.005

Plesnar-Bielak, A., & Łukasiewicz, A. (2021). Sexual conflict in a changing environment.
Biological Reviews. https://doi.org/10.1111/brv.12728

Priest, N. K., Galloway, L. F., & Roach, D. A. (2008). Mating frequency and inclusive
fitness in Drosophila melanogaster. *American Naturalist*, *171*(1), 10–21.
https://doi.org/10.1086/523944

Quinn, G., & Keough, M. (2002). *Experimental Design and Data Anal- ysis for Biologists.*
(C. University Press, Ed.). Cambridge.

Rankin, D. J., Dieckmann, U., & Kokko, H. (2011). Sexual conflict and the tragedy of the
commons. *American Naturalist*, *177*(6), 780–791. https://doi.org/10.1086/659947

Sirot, L. K., LaFlamme, B. A., Sitnik, J. L., Rubinstein, C. D., Avila, F. W., Chow, C. Y., &
Wolfner, M. F. (2009). *Molecular social interactions: Drosophila melanogaster seminal
fluid proteins as a case study. Advances in genetics* (1st ed., Vol. 68). Elsevier Inc.
https://doi.org/10.1016/S0065-2660(09)68002-0

Sirot, L. K., Wolfner, M. F., & Wigby, S. (2011). Protein-specific manipulation of ejaculate
composition in response to female mating status in Drosophila melanogaster.
Proceedings of the National Academy of Sciences of the United States of America,
108(24), 9922–9926. https://doi.org/10.1073/pnas.1100905108

Teseo, S., Veerus, L., Moreno, C., & Mery, F. (2016). Sexual harassment induces a
temporary fitness cost but does not constrain the acquisition of environmental
information in fruit flies. *Biology Letters*, *12*(1). https://doi.org/10.1098/rsbl.2015.0917

Wigby, S., & Chapman, T. (2004). Female resistance to male harm evolves in response to manipulation of sexual conflict. *Evolution*, *58*(5), 1028–1037. https://doi.org/10.1111/j.0014-3820.2004.tb00436.x

Wigby, S., & Chapman, T. (2005). Sex Peptide Causes Mating Costs in Female Drosophila melanogaster. *Current Biology*, (March 2005), 1118. https://doi.org/10.1016/j

Wigby, S., Sirot, L. K., Linklater, J. R., Buehner, N., Calboli, F. C. F., Bretman, A., … Chapman, T. (2009). Seminal Fluid Protein Allocation and Male Reproductive Success. *Current Biology*, *19*(9), 751–757. https://doi.org/10.1016/j.cub.2009.03.036

Wolfner, M. F. (1997). Tokens of love: Functions and regulation of Drosophila male accessory gland products. *Insect Biochemistry and Molecular Biology*, *27*(3), 179–192. https://doi.org/10.1016/S0965-1748(96)00084-7

Yun, L., Agrawal, A. F., & Rundle, H. D. (2021). On Male Harm: How it is Measured and How it Evolves in Different Environments. *The American Naturalist*, *81*(1), 1–37.

Yun, L., Chen, P. J., Kwok, K. E., Angell, C. S., Rundle, H. D., & Agrawal, A. F. (2018). Competition for mates and the improvement of nonsexual fitness. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(26), 6762–6767. https://doi.org/10.1073/pnas.1805435115

Yun, L., Chen, P. J., Singh, A., Agrawal, A. F., & Rundle, H. D. (2017). The physical environment mediates male harm and its effect on selection in females. *Proceedings of the Royal Society B: Biological Sciences*. https://doi.org/10.1098/rspb.2017.0424

Zuur, A. F., Ieno, E. N., & Elphick, C. S. (2010). A protocol for data exploration to avoid common statistical problems. *Methods in Ecology and Evolution*, *1*(1), 3–14. https://doi.org/10.1111/j.2041-210x.2009.00001.x