A machine-learning tool to identify bistable states from calcium imaging data

Aalok Varma[1], Sathvik Udupa[2], Mohini Sengupta[1#], Prasanta Kumar Ghosh[2] and Vatsala Thirumalai[1*]

[1] National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bellary Road, Bangalore 560065.

[2] Department of Electrical Engineering, Indian Institute of Science, Bangalore - 560 012.

[#]Current Address:
Washington University School of Medicine in St Louis, 660 S Euclid Ave, St. Louis, MO 63110.

*Correspondence:
vatsala@ncbs.res.in

Keywords:
Purkinje neuron, bistability, zebrafish, machine learning, convolutional neural network, recurrent neural networks, cerebellum.

Key Points Summary

- Calcium imaging – the gold standard of inferring neuronal activity – does not report cellular state in neurons that are bistable, such as Purkinje neurons in the cerebellum of larval zebrafish. We model the relationship between Purkinje neuron electrical activity and its corresponding calcium signal to compile a dataset of state-labelled simulated calcium signals.

- We apply machine-learning methods to this dataset to develop a tool that can classify the state of a Purkinje neuron using only its calcium signal, which works well on real data even though it was trained only on simulated data.

- CaMLsort also generalizes well to bistable neurons in a different brain region (ventral tegmental area) in a different model organism (mouse).
This tool offers a new way of analyzing calcium imaging data from populations of bistable neurons, thereby facilitating our understanding of how these neurons carry out their functions in a circuit.

Abstract

Mapping neuronal activation using calcium imaging *in vivo* during behavioral tasks has advanced our understanding of nervous system function. In almost all of these studies, calcium imaging is used to infer spike probabilities since action potentials activate voltage-gated calcium channels and increase intracellular calcium levels. However, neurons not only fire action potentials, but also convey information via intrinsic dynamics such as by generating bistable membrane potential states. While a number of tools for spike inference have been developed and are currently being used, no tool exists for converting calcium imaging signals to maps of cellular state in bistable neurons.

Purkinje neurons (PNs) in the larval zebrafish cerebellum exhibit membrane potential bistability, firing either tonically or in bursts. Several studies have implicated the role of a population code in cerebellar function, with bistability adding an extra layer of complexity to this code. In this manuscript we develop a tool, CaMLSort which uses convolutional recurrent neural networks to classify calcium imaging traces as arising from either tonic or bursting cells. We validate this classifier using a number of different methods and find that it performs well on simulated event rasters as well as real biological data that it had not previously seen. Moreover, we find that CaMLsort generalizes to other bistable neurons, such as dopaminergic neurons in the ventral tegmental area of mice. Thus, this tool offers a new way of analyzing calcium imaging data from bistable neurons to understand how they participate in network computation and natural behaviors.

Introduction

Calcium imaging enables us to sample the activity of populations of neurons and even whole brains during animal behaviour (Ahrens et al., 2013; Grewe and Helmchen, 2009; Grienberger and Konnerth, 2012), providing us a window into the functioning of the nervous system. When a neuron fires an action potential, somatic calcium levels increase dramatically due to the opening of voltage-dependent calcium channels. Simultaneous electrophysiology and calcium imaging recordings of neuronal activity demonstrate the correspondence of somatic calcium imaging signals with action potential firing (Chen et al., 2013; Kerr et al., 2005; Kwan and Dan, 2012; Tian et al., 2009). Nevertheless, the calcium imaging signal is at best a delayed, low-pass filtered and non-linearly transformed proxy for neuronal activity. Non-linearities are introduced due to the dynamics of calcium release, and the kinetics of calcium binding and unbinding by the sensor molecules (Ali and Kwan, 2020). Yet, this technique is useful in cases where the neurons being studied have low baseline firing rate. However, this is not often the case and several neuronal classes, such as cerebellar Purkinje neurons and cortical interneurons are known to fire at very high spike rates spontaneously (Armstrong and Edgley, 1984; Markram et al., 2004). Further, many neuronal types have interesting intracellular dynamics apart from action potential firing that might strongly modulate calcium imaging signals (Lin et al., 2007; Moreaux and Laurent, 2007). Lastly, confounds may be introduced when more than one type of electrical event leads to somatic calcium increase (Lev-Ram et al., 1992; Ramirez and Stell, 2016).

Purkinje neurons (PNs) show all of the features mentioned above that make the interpretation of calcium signals in these neurons especially difficult. PNs are principal neurons of the cerebellum, integrate thousands of synaptic inputs and are indispensable for cerebellar function (Narayanan and Thirumalai, 2019). Therefore, studying PN population activity during motor behaviours is critical to deciphering cerebellar computation. PNs fire simple spikes and complex spikes - in mammals, climbing fiber (CF) input from the inferior olive leads to calcium entry within the dendrites and several spikelets that ride the depolarization wave (Kitamura and Häusser, 2011; Miyakawa et al., 1992); in fish, the olivary input causes an all-or-none AMPAR-mediated giant synaptic potential with concomitant calcium entry but without any spikelets (Sengupta and Thirumalai, 2015). While the CF input causes a relatively large calcium signal, bursts of simple spikes can also generate calcium signals via the activation of somatic voltage-dependent calcium channels, albeit smaller in amplitude (Knogler et al., 2019; Sengupta, 2015). This means that optically recorded calcium dF/F signals cannot be uniquely assigned to simple spikes or CF inputs.

In addition to the above, larval zebrafish PNs exhibit membrane potential bistability – these neurons can exist in one of two stable membrane potentials. At a depolarized state, they generate action potentials at a more or less regular rate and are said to be tonic. When hyperpolarized, they generate bursts of action potentials (Sengupta and Thirumalai, 2015). Simple spikes in the bursting state are highly correlated with motor bouts whereas those in the tonic state are not (Sengupta, 2015). This implies that bistability has important ramifications for how motor bouts are represented across the entire PN population. We wondered if calcium imaging can be used to deduce the state of PNs. Since both CF inputs and simple spikes contribute to

122 calcium signals, assigning optically recorded dF/F signals is not straight-forward and
123 deduction of spike rate using existing spike inference algorithms is not possible.
124
Here, we first performed simultaneous electrophysiological recordings and calcium
imaging in larval zebrafish PNs to obtain ground truth data. Unsurprisingly, we found
that none of the usually measured parameters such as peak amplitude, area under
the curve etc., of the calcium signal uniquely correlated with cellular state. We
therefore built a sorter using convolutional recurrent neural networks to identify
cellular state from dF/F data. We have tested our sorter with data that was not part of
the training set and validated that it performs at above 90% accuracy and with F1
scores greater than 0.8 in most cases. Our method demonstrates that calcium
imaging datasets can be mined using such tools to glean greater insights into
neuronal activity states.
135

Materials and Methods

137
Ethical approval
139 All experimental procedures were approved by the institutional animal ethics
140 committee (IAEC) and institutional biosafety committee (IBSC) at the National Centre
141 for Biological Sciences.
142
Experimental Animals
Experiments were performed on Indian wild type (Ind WT) zebrafish (*Danio rerio*).
Embryos were obtained by setting up an in-cross between adult Ind WT zebrafish
raised and housed in a ZebTec multi linking system (Tecniplast, Italy) with a pH
setting of 7.8 and conductivity of 1200µS. Larvae were kept in a MIR-154 incubator
(Sanyo, Japan) with 14:10h light-dark cycle maintained at 28°C in E3 medium
(composition in mM: 5 NaCl, 0.17 KCl, 0.33 $CaCl_2$,and 0.33 $MgSO_4$, pH 7.8) in
standard 90mm Petri dishes (Tarsons, Kolkata, India). The medium was routinely
replaced, and starting at 5 dpf, larvae were fed Zeigler Larval diet AP100 (<100
microns) (Pentair AES, FL, United States). Larvae have not yet undergone sex
specification by these stages, so experiments performed were agnostic to the sex of
the animal.
155
Transient transgenesis by microinjection
Thin-walled borosilicate capillaries (1.0mm OD, 0.56mm ID; Warner Instruments,
Hamden, CT, United States) were pulled using a Flaming-Brown P-97 pipette puller
(Sutter Instruments, Novato, CA, United States) to produce long and thin pipettes for
microinjection. Indian WT embryos at the 1-2 cell stage were injected with ~10nL of
a mixture of tagRFP-T:PC:GCaMP5G plasmid (Matsui et al., 2014) (50 ng/µL) and
Tol2 transposase mRNA (50 ng/µl) (Urasaki et al., 2006), using a Picospritzer III
(Parker Hannifin, NH, United States) pressure injection apparatus, and raised as
described above. Larvae were screened at 5 dpf at an epifluorescence dissection
microscope (Olympus SZX-16) and those showing mosaic labelling of Purkinje
neurons were selected for experiments, which were performed at 7 dpf at room
temperature (25-28°C).
168
169
170

Electrophysiology

Larvae were anaesthetised in 0.01% MS-222 (Sigma-Aldrich; Missouri, USA) and transferred to a custom-made acrylic recording chamber. Fine tungsten wire (California Fine Wire, CA, USA) was used to pin the larva onto a piece of Sylgard (Dow Corning, Midland, MI, United States) glued to the recording chamber. Two pins were put through the notochord at the level of the swim bladder and the tail, respectively. A third pin was then used to position the larval head dorsal-up. The MS222 was replaced by external solution (composition in mM: 134 NaCl, 2.9 KCl, 1.2 $MgCl_2$, 10 HEPES, 10 glucose, 2.1 $CaCl_2$, 0.01 D-tubocurarine; pH 7.8; 290 mOsm) and skin on the dorsal surface of the larva was carefully removed using fine no. 5 forceps (Fine Science Tools, Foster City, USA) to expose the brain.

A Flaming-Brown P-97 pipette puller (Sutter Instruments, Novato, CA, United States) was used to pull patch pipettes made of standard-walled borosilicate capillaries with a filament (1.5 mm OD; 0.86 mm ID; Warner Instruments, Hamden, CT, United States), pulled to a final tip diameter of 1-1.5 µm. For loose patch recordings, pipettes were backfilled with external solution, while for whole-cell recordings, pipettes were backfilled with K-gluconate internal solution (composition in mM: 115 K gluconate, 15 KCl, 2 $MgCl_2$, 10 HEPES, 10 EGTA, 4 Mg-ATP; pH 7.2; 290 mOsm). Solutions were filtered using a 0.2µm filter (Millipore, Merck, Germany) before backfilling pipettes, which typically had resistances between 8 and 10 MΩ. Sulforhodamine (Sigma-Aldrich, St.Louis, MO, United States) was added to the patch internal solution at a final concentration of 5µg/mL to facilitate visualization of cellular morphology post-recording.

Recordings were acquired with a Multiclamp 700B amplifier, Digidata 1440A digitizer and pCLAMP 10 software (Molecular Devices, Sunnyvale, CA, United States). Data were low pass filtered at 2 kHz and sampled at 20-50 kHz with a gain of 1 (whole-cell recordings) or between 20-1000 (loose patch recordings), so as to optimally use the dynamic range of the digitizer.

Simultaneous calcium imaging and electrophysiology

Indian WT larvae microinjected with the Tol2-tagRFP-T:PC:GCaMP5G construct (gift from Dr. Hideaki Matsui, Niigata University, Japan) and screened for sparse labelling were used for this experiment. Purkinje neurons expressing GCaMP5G were targeted for the recordings. Widefield imaging was performed using a 60x water-immersion objective (1.0 NA) at an Olympus BX61WI microscope. Excitation light was provided using a pE300-ultra (CoolLED, Andover, England). Imaging was performed using an EVOLV EM-CCD camera (Photometrics, Tucson, AZ, United States) at the rig, with an exposure of 30 ms, and an on-chip gain of 170. An ROI was drawn around the targeted cell, so that only an isolated cell was imaged. Imaging was synchronized with electrophysiology by interfacing with the Digidata 1440A.

Calcium Imaging Analysis

Image analysis was performed in Fiji/ImageJ (Schindelin et al., 2012). In order to identify cell boundaries, the averaged time series was used as a reference and an ROI drawn around the cell. Bleach correction was performed using the Exponential Fitting Method in the Bleach Correction plugin in Fiji. The average raw pixel intensity values within the ROI for each time point was extracted. The baseline intensity, F0,

was taken to be the 5th percentile of these values. dF/F was then calculated for each
frame using the formula (F - F0)/F0, where F represents the fluorescence in that
frame and F0 the baseline fluorescence. Following this calculation, a Gaussian filter
was applied to the time series in MATLAB R2019b (Mathworks, Natick, MA), using
its function '*smoothdata*', with a window size of 15.

To detect transients in the calcium signal, dF/F traces were first normalised to a
range of [0, 1] using min-max feature scaling. Peaks in the normalised trace were
then detected using the '*findpeaks*' function in MATLAB, with a minimum peak height
of 0.1, a minimum peak prominence of 0.05 and a minimum interval of 500ms
between consecutive peaks.

Spike detection and analysis
All electrophysiological analyses were done using custom scripts written in MATLAB,
using functions defined in a publicly available repository
(https://github.com/wagenadl/mbl-nsb-toolbox) to read electrophysiology data. For
extracellular recordings, spikes were identified as events that showed large
fluctuations relative to a 25ms rolling window baseline and the peak timing and
amplitude were extracted. Events were sorted based on amplitude, with small
amplitude events being classified as simple spikes and large amplitude events as
climbing fiber (CF) inputs, as per (Sengupta and Thirumalai, 2015). For intracellular
recordings, peak detection algorithms inbuilt in MATLAB were used to identify
spikes. Once again, events were sorted as simple spikes or CF inputs based on their
amplitude relative to baseline.

Calcium imaging reconstruction from electrophysiology data
Simultaneous calcium imaging and extracellular electrophysiology data was used to
infer optimal GCaMP5G kernels, which are of the general form

$$A1 * e^{-\frac{t}{\tau_{decay}}} - A2 * e^{-\frac{t}{\tau rise}}$$

for simple spikes and CF inputs of zebrafish PNs. Simple spike and CF input timings
were first extracted from the electrophysiology data for a single cell. Simple spike
timings were convolved with a simple spike kernel, with amplitudes A1s and A2s and
CF input timings were convolved with a similar CF input kernel, with amplitudes A1c
and A2c. The sum of both these convolutions was taken to be the putative calcium
imaging reconstruction, down sampled to match the sampling rate of calcium
imaging. The mean squared error (MSE) between the reconstruction and the true
calcium imaging trace was calculated. The values for A1s, A2s, A1c and A2c were
obtained by minimizing the MSE between the true calcium imaging trace and the
reconstructed trace using gradient descent. To ensure that local minima were
avoided, the gradient descent algorithm was repeated with randomized starting
points. The kernels with these coefficients were taken to be the optimal simple spike
and CF input kernels, respectively, for that cell.

The 'optimal' kernel would be one that generalized best. This was tested by using
the kernels from one cell to reconstruct calcium imaging for other cells for which
ground truth data was available. As before, the MSE and also the Pearson
correlation coefficient between the reconstructed trace and the true trace were
calculated. The final 'optimal' kernel was obtained by taking the kernel which

generalized the best, i.e., had minimum MSE and maximum correlation across all
cells. To test whether the correlation was real or not, the Pearson correlation
coefficient between a scrambled reconstructed trace and the true trace was also
calculated, for reference.
273
274 This process of convolving simple spike and CF timings with their optimal GCaMP5G
275 kernels and summating the resultant traces was then used to generate a calcium
276 imaging trace from both extracellular and intracellular Purkinje cell recordings.
277
278
Building a state-labelled calcium imaging dataset
Intracellular and extracellular recordings from larval zebrafish Purkinje neurons
acquired during the period 2011-2020 by various members of the lab were compiled.
Good quality recordings were those in which the resting membrane potential of the
cell was not below -80mV or not above -20mV, were uninterrupted gap-free
recordings with no current injections and had two clearly distinguishable event types.
Besides poor quality recordings, those recordings with an ambiguous, indeterminate
state were also excluded. Only those recordings which met the aforementioned
criteria were retained. Then, recordings were renumbered randomly, to remove
grouping of data collected by the same individual. Recordings were then classified
as tonic or bursting, first by visual inspection. These classifications were then further
verified by inspecting the inter-event intervals of simple spikes. While inter-event
intervals from tonic cells form a single cluster, bursting cells have inter-event
intervals that trail to large values, corresponding to the inter-burst intervals. Lastly,
the coefficient of variation of the inter-spike intervals was used as an extra parameter
of interest. These three pieces of information were used in conjunction (**Fig. 1D-G**),
but the final classification relied on visual inspection. If there was a switch in state
within the recording, the recordingswere labelled as "switch", with the appropriate
state switch timing and kind (tonic-to- bursting, or bursting-to-tonic) marked. A
summary of the recording information is in Table 1.
299
Here, data was split into two groups - one that contained traces only with a single
state throughout the recording, and another that contained traces with switches in
state. Once split, the spike detection protocol described above was applied to each
group of recordings to detect the simple spikes and CF inputs. These event timings
were then used to generate simulated calcium imaging traces as described above.
Combining the reconstruction with the assigned state resulted in the state-labelled
calcium imaging dataset.
307
Principal Components Analysis
Principal Components Analysis (PCA) was performed using the PCA module that is
part of the *sklearn* package in Python (Pedregosa et al., 2011). The input to the
model was a vector with the trace features for each 10-second-long trace in the
reconstructions dataset (number of peaks, peak amplitude, area under the curve,
mean, standard deviation and coefficient of variation). A 2-component PCA was
performed, and the resulting fit (shown in Figure 4E) explained 88% of the variance
in the data.
316
317

Training the CNN-LSTM model for classification

The dataset of reconstructed calcium imaging time series was first split into 5 folds to perform cross validation (Fig. 5A). The experiments are repeated by considering each fold as a test set, with one of the folds being the validation set and remaining 3 folds to be training sets such that the data split for each experiment is as follows - 60% for training, 20% for validation, 20% for testing. This splitting was done at the level of entire traces, such that the training, testing and cross-validation sets all had a balanced representation of traces from both the tonic and bursting classes. We chose to do a 5-fold cross-validation, i.e. the entire training dataset is split into 5 parts, each containing 20% of the data. Four of the folds are used for training and the remaining one is used for cross-validation.

The architecture of the network is shown in Fig. 5B. The input to the network is the entire trace sequence at 30Hz, normalized using min-max normalization for every non-overlapping 10 second period, thus setting the minimum to 0, and maximum to 1. Note that if the sampling frequency of a calcium trace was not 30Hz, it was first linearly interpolated to a 30Hz sampling frequency before being seen by the neural network. The first layer of processing is a CNN with kernel size 30 and stride size 30. The CNN down samples the input trace from 30Hz to 1Hz, and produces 4 vector outputs,which act as the input to the LSTM. The LSTM is bidirectional, so it looks at the sequence in both the forward and reverse order to produce an output. The LSTM has 24 hidden layers, so it produces 48 numbers per time step (Fig. 5B). The feature dimensions are collapsed by a Linear layer by taking their weighted average. This provides the classification logits at each position, which is passed through a softmax layer. The network is optimized using a cross entropy loss and the Adam optimizer (Kingma and Ba, 2017). The output of a trained neural network, then, is the probability of each timestep (at 1 Hz) belonging to the bursting class, from which the probability of it belonging to the tonic class can be derived. The class with the higher probability is assigned to each time step.

Since the model acts on the entire trace sequence, the amount of independent data for training the model is significantly reduced. To overcome this issue, we used a data augmentation strategy, which allows the network to learn features on arbitrary length sequences. For a sequence of length N, we determine a random segment size n, with n between [k, N], where k is the minimum sequence length. We randomly determine a starting position x, with a constraint that x < N - n. In our case, k=300 samples, and n varies in multiples of k (i.e. n = 300, 600, 900…). In each iteration of the network pass during the training phase, this random sequential segment was taken as the input trace to the model and gradient descent was performed to update weights. This was done for all the traces in the training set, so that the network saw entire traces at least once and in a non-overlapping fashion during the 500 epochs of training. The training is stopped before the full 500 epochs, based on early stopping criteria - if the validation loss does not improve after few epochs. All models were trained using the PyTorch framework in Python 3 (Paszke et al., 2019).

For all evaluations - cross-validation, test and other independent validations, the network had to make predictions for the entire trace duration and not just the n-sample long chunks. Networks were evaluated using two metrics - classification accuracies and F1-scores. Classification accuracies are simply the fraction of predictions that were correct. F1-scores, on the other hand, are adjusted for class

imbalances and account for the kind of errors made. By definition, an F1-score is the harmonic mean of the precision and recall metrics, where precision is the fraction of all predicted positives that were true positives, and recall is the fraction of all ground truth positives that were true positives.

To improve/smoothen predictions made by CaMLsort, we used a majority voting strategy. To do this, we re-sampled the time series using rolling window sampling with a window length N. For each time step of prediction, t_n, we found all the m windows (where $m{\le}N$) that contained t_n (i.e. $[t_{n-N+1}, t_{n-N+2}, \ldots t_n, t_{n+1}]$, $[t_{n-N}, t_{n-N+1}, \ldots t_{n-1}, t_n]$, and so on until $[t_n, t_{n1}, \ldots t_{n+N-1}, t_{n+N}]$). For each of these windows, an average posterior score was calculated for each state, and the "average class" taken to be the state which had the higher score. The "majority class", then, was the one which was the most frequent average class across all m windows. A "confident call" was made if and only if the average posterior scores for the majority class across all m windows was strictly higher than the average posterior scores for the minority class.

Generating artificial Purkinje neuron spike trains
Synthetic spike trains of the tonic class were generated using a Poisson statistical model. The only parameter needed for this is an average event rate, λ, which we took to be a random number between 4-10 for simple spikes. Spike trains for bursting neurons do not follow Poisson statistics. Hence, we used an inverse sampling method to generate spike trains of the bursting class. For this, we used the compiled electrophysiology data to obtain the real cumulative distribution of interevent intervals for simple spikes across all bursting cells. A random number, y, is generated from the uniform distribution and it is used to find a value x from the reference cumulative interspike interval (ISI) distribution such that P(ISI <= x) = y. Thus, by generating events sequentially with a given next interevent interval, we obtained artificial simple spike trains for the bursting class.

CF input event trains were generated using a Poisson statistics model with an average event rate between 0.2-1. The distributions of interevent intervals in the synthetic datasets were compared against the real distributions using histograms and QQ plots. Independent reviewers were asked to classify unlabelled spike trains as real or artificial and managed to do so with an accuracy of ~42% on average.

Statistical Analysis
We used a one-sided Mann-Whitney U test to compare the distributions of interspike interval coefficient of variation between tonic and bursting cells, for both Purkinje neurons (Fig 1G) and dopaminergic VTA neurons (Fig. 9C).

To compare the quality of reconstructions obtained from the generative forward model, we used a Kruskal-Wallis test to detect any overall group differences and followed it up with a post-hoc Dunn's test for individual comparisons (Fig. 3C and Fig. 9E).

For comparisons involving a nested data structure, such as those involving trace features between tonic and bursting cells (Fig. 4D), where there are multiple cells and multiple measurements per cell, we used a linear mixed-effects model with a maximum likelihood estimator to calculate p-values.

Results

Zebrafish Purkinje neurons exhibit membrane potential bistability

Purkinje neurons (PNs) are the principal neurons in the cerebellar circuit (Fig. 1A). In larval zebrafish, they exhibit two kinds of electrical events that can be distinguished solely based on their amplitude in both extracellular and intracellular recordings (Sengupta and Thirumalai, 2015) (Fig. 1 B-E). The smaller amplitude events are action potentials and are called simple spikes (Fig. 1B,D,E, cyan), while the larger amplitude events are all-or-none synaptic input from climbing fibers (Fig. 1C,D,E, magenta). Individual simple spike and CF input waveforms are shown in Fig. 1B and 1C, respectively, for both whole-cell and juxtacellular recordings.

These neurons exhibit membrane potential bistability, i.e. they are found to be at one of two stable membrane potentials, depending on which they fire either tonically or in bursts. Cells can spontaneously switch between the two states, but can also be taken from one state to the other by current injection (Sengupta and Thirumalai, 2015).

The two states can be distinguished visually from extracellular loose-patch recordings based on (i) the firing frequency of simple spikes and (ii) the presence of long intervals of inactivity in the burst mode (Fig. 1D, left) which are not present in the tonic mode (Fig. 1D, right). The same features can be used to distinguish between the two modes in whole-cell patch- clamp recordings, along with an extra feature – membrane potentials are more hyperpolarized in the bursting mode (Fig. 1E, left) as opposed to the tonic mode (Fig. 1E, right). Additionally, in tonic cells, inter-spike intervals form a single cluster, while bursting cells have inter-event intervals that trail to large values, corresponding to the inter-burst intervals (Fig. 1F). This is also reflected in the values of coefficient of variation of the inter-spike intervals, which are larger in bursting cells than the more regularly spiking tonic cells (Fig. 1G). Given that there is a lot of variability across cells and recordings, no single cut-off value could be determined for these parameters (Fig. 1G). Instead, all the aforementioned features were taken together to determine the final state classification.

Calcium imaging reports both simple spike bursts and individual CF inputs, independent of state

Given that PNs manifest two distinct activity patterns (tonic and bursting) and two types of electrical events (simple spikes and CF events), we wondered if calcium imaging signals can reliably report these distinct patterns of activity. To do this, we performed simultaneous wide-field calcium imaging and targeted patch-clamp electrophysiology from zebrafish PNs. We used the carbonic anhydrase (Ca8) enhancer element and a minimal promoter (*cfos*) to drive expression of the genetically encoded calcium sensor GCaMP5G in PNs (Matsui et al., 2014).

Embryos were microinjected at the 1-2 cell stage and those larvae that showed sparse labelling in PNs were selected for targeted patch-clamp electrophysiology and imaging (Fig. 2A). We chose to perform extracellular, loose-patch recordings so as to be minimally invasive and to avoid the dialysis of cellular contents including GCaMP. We obtained such recordings from 15 cells in 12 larvae, out of which 9 cells were naturally in the tonic state, and 6 cells in the bursting state.

Representative traces from this experiment for cells in the bursting and the tonic
states are shown in Fig. 2B and 2C, respectively. The first observation we made was
that many of the calcium transients (marked with a hash symbol in Fig. 2B and 2C)
had an almost one-to-one correspondence with CF inputs (magenta lines in the
raster plot) in cells of both states. In tonic cells, most simple spikes were not reported
as transients, as one would expect from cells that fire at high frequencies. However,
in some cases, calcium transient peaks occurred in the absence of CF inputs, during
times of intense simple spike firing (Fig. 2C, see asterisk and cyan lines in raster
plot). Similarly, in bursting cells, we found some calcium transients associated only
with a burst of simple spikes (Fig. 2B, asterisk). Naturally, then, one would expect to
see calcium transients when CF inputs co-occurred with a burst of simple spikes,
which is what we observed (marked with open circles in Fig. 2B and 2C). Thus,
regardless of the state of a cell, we found calcium transients that corresponded to
individual CF inputs, bursts of simple spikes or a combination of both.
480
While this method confirms that a calcium transient occurs after a CF input,
suggesting a causal link between the CF input's arrival and the measured calcium
influx into the cell, it is not useful in the case of simple spikes. Firstly, there are
multiple simple spikes per frame - approximately 2-3 spikes per frame when imaged
at 30Hz. This is further worsened by the timescales of GCaMP5G's rise and decay,
which occurs on the order of 100s of milliseconds (Akerboom et al., 2012; Chen et
al., 2013). Moreover, one cannot separate out the individual effects of simple spikes
and CF inputs because the two events are interspersed and thus, when calculating
the triggered average response for one event, there is bleed-through from the other.
490
To ascertain whether CFs and simple spikes indeed trigger calcium signals, we
decided to compare the number of events in the time leading up to the calcium peak
(Fig. 2D, region I) and during the decay period (Fig. 2D, region II). The expectation
was that if an event triggered calcium activity, there would be more of them in region
I, which includes the time of initiation of the calcium transient (Fig. 2D), than in
region II, when the signal is decaying. GCaMP5G has a rise time constant of
~100ms, which means that the time it takes to reach a peak value from the baseline
is ~201ms (modelled using a difference of single exponentials, see Methods).
Hence, to be sure that the time of initiation was included in region I, we considered a
300ms interval before the peak of the calcium signal. An interval of the same 300ms
duration after the peak was taken to be region II.
502
We first detected calcium transient peaks in the dF/F signal (see Methods), of which
there were a total of n=171 detected across the N=15 cells. The average number of
simple spikes (Fig. 2E) and CF inputs (Fig. 2F) are significantly different in region I
and II (linear mixed effect model; p<0.0001 for simple spikes and p = 0.0025 for CF
inputs). There are more CF inputs in region I compared to region II indicating that
these events can trigger calcium signals. However, for simple spikes, there is a
small, but significant increase in region II compared to region I. While this is opposite
of what one might have expected, it can be explained by the high rate of simple
spikes lasting over several hundreds of milliseconds in both bursting and tonic cells.
In both regions I and II, there are approximately 2.5 simple spikes within the 300 ms
window. These results imply that both simple spikes and CF inputs are associated
with dF/F transients.
515

516 In summary, it is difficult to infer the underlying event train or even the cellular state
517 of a Purkinje neuron directly from its calcium signal, as there is no one-to-one
518 correspondence between calcium transients and their underlying electrophysiological
519 events, as has also been reported previously (Knogler et al., 2019; Sengupta, 2015).
520
The calcium signal for a Purkinje neuron can be reconstructed from its
electrophysiological recording
Though the calcium signal in PNs could arise from more than one electrical event,
we hypothesized that there might be sufficient information in the calcium signal time
series to allow us to determine whether it is tonically firing or bursting. One possible
approach to this problem could be the application of machine learning algorithms to
build a classifier that labels cells as tonic or bursting given a time series of calcium
transients. For this, it was first necessary to generate a large ground truth dataset
(100's of cells) comprising the electrical activity patterns and their corresponding
calcium transients for training the classifier. Ideally, this dataset would take the form
of what is shown in Fig. 2B,C, where both electrophysiology and imaging data are
simultaneously acquired. Our ground truth dataset has only 15 cells, however, given
that the simultaneous imaging and electrophysiology experiments are low yield and
technically challenging. Hence, we needed a strategy to augment our dataset.
Mathematically, the calcium signal obtained by imaging can be thought of as the
result of the convolution of spike trains with a calcium sensor kernel having a fast
rise and a slow decay and described as a difference of exponentials. This
transformation has been shown to work well in cells that are mostly quiescent and
where every single action potential produces a large increase in intracellular calcium
concentration (Smetters et al., 1999; Rupprecht et al., 2021; Vogelstein et al., 2010;
Yaksi and Friedrich, 2006). We hypothesized that a similar transformation could be
used to reconstruct the calcium signal profiles of PNs from their electrophysiological
recordings alone. Such reconstructed calcium imaging traces with their
corresponding electrical recording traces may then be applied to build the classifier.
545
However, given that these neurons have not one, but two electrophysiological
events, each contributing differently to the calcium signal obtained from imaging, we
must assign each event a separate kernel (Fig. 3A), which we assume to be
independent of each other. This is a valid assumption, since simple spikes and CF
events arise from distinct intrinsic mechanisms (Sengupta and Thirumalai, 2015). We
used simultaneous imaging and electrophysiology recordings to find optimal kernels
for simple spikes and CF events. For each PN recording, we obtained simple spike
and CF event rasters. Each event raster was then independently convolved with its
own exponential response kernel, and the resulting "event-specific signals" were
linearly summed to produce the final reconstruction (Fig. 3A). We then calculated
the correlation coefficient and the mean squared error (MSE) between the
reconstructed calcium signal time series and the experimentally obtained calcium
imaging time series.
559
With this description in place, the problem then becomes one of finding optimal
coefficients for each response kernel. This optimization was done by gradient
descent to minimize the MSE between the ground truth and the reconstruction (Fig.
3A). The process was performed independently for each cell in the dataset. Then,
these optimal kernel combinations were used to make reconstructions for other cells
in the dataset, and the kernel that generalized best across all cells, as quantified by

least total MSE and highest total cross-correlation between the ground truth and
reconstruction, was taken to be the best kernel for the nextsteps.
568
Although this is a simple forward model with calcium dynamics assumed to be linear
and time-invariant, we found that this method to approximate the calcium signal
works well, no matter the state of the neuron (Fig. 3B). The quality of the
reconstructions was quantified by measuring the correlation coefficient between the
ground truth and the reconstruction (Fig. 3C). To benchmark the correlation
coefficient values obtained against those obtained simply by chance, we also
calculated the correlation coefficient between the ground truth and a scrambled
version of the reconstruction, as well as that between the ground truth and a
reconstruction generated after shuffling all the event times in the recording (Fig. 3C).
In both the latter cases, the correlation disappears, suggesting that the
reconstructions generated by this forward model are, indeed, faithful reconstructions.
We next calculated the residuals between ground truth and reconstructions, the
distribution for which is shown in Fig. 3D. It is not normally distributed, and has a
positive skew, which means that our reconstructions tend to underestimate the true
signal. Nevertheless, the small values of the residuals further lend support to the
idea that the forward model of calcium signal reconstruction works well.
585
Given that only groups/bursts of simple spikes produce a calcium transient amplitude
comparable to that generated by an individual CF input (Knogler et al., 2019; Fig.
2B), one would expect a much smaller response kernel for simple spikes than for the
CF input. Indeed, the optimal kernels found by this method show that the CF input
response kernel is approximately 7.5 times larger than the simple spike response
kernel (Fig. 3E).
592
A state-labelled calcium signal dataset was generated by reconstructing
the calcium signal for a compilation of electrophysiological traces
Having a method of converting any given PN electrophysiology trace to its
corresponding calcium imaging trace allowed us to generate simulated calcium
imaging traces for a compilation of all electrophysiological recordings acquired in our
lab (Fig. 4A). To do so, we first pooled all the electrophysiological recordings, both
extracellular and whole-cell, made during the period of 2011-2020, for a total of
N=173 recordings. Recordings of poor quality (N=37), i.e., those with interruptions,
or noisy baselines, unusually broad/filtered events, or indistinguishable simple spikes
and CF inputs were first excluded from the dataset. Retained recordings then had a
state assigned to them (using criteria described above; also see Fig 1D-G), including
state switches, if there were any. This collection was split into two sets, one with
those recordings which had only one state throughout (N=131), and another with
those which had a state switch (N=5). The entire complement of recordings and their
corresponding states is summarized in Table 1. Events detected in all the recordings
were used to reconstruct a putative calcium signal using the kernels described in Fig.
3E. This finally produced two labelled datasets - a state-labelled reconstructed
calcium signal dataset and an independent dataset with switches (Fig. 4A). The
recordings used to generate this dataset were of different durations with a median
duration of 60 seconds. In order to get multiple samples per recording, we decided to
split the traces into 10- second-long non-overlapping chunks, representative samples
from which are shown in Fig 4B. This subsampling yielded a dataset of state-labelled
calcium imaging traces (n=928 traces of which n=580 bursting, n=348 tonic) that is

much larger than could have been obtained experimentally. The entire complement
of non-overlapping samples is shown as a heatmap in Fig. 4C.
618
To test whether there were properties of these traces that could be used to
unambiguously classify their source state, we compared the number of peaks,
average peak amplitude, area under the curve, mean, standard deviation and
coefficient of variation of the reconstructed calcium traces (Fig. 4D). Except peak
amplitude, all other properties were significantly different from each other for the
tonic and bursting groups. Nevertheless, given any of these properties, it was not
possible to unambiguously assign state as the distributions are overlapping (Fig.
4D). Further, a principal component analysis of these trace features also failed to
separate tonic and bursting cells based on the above-mentioned features of their
calcium traces (Fig. 4E).
629
630   Machine learning methods have successfully been used to solve such problems
631   when the data points are not linearly separable (He et al., 2016). Thus, we turned to
632   developing a machine learning model to identify the state of a cell from a calcium
633   signal time series.
634
Training a convolutional recurrent neural network to distinguish state
from calcium imaging data alone
To find good machine learning models to solve this classification problem, we
performed 5-fold cross validation, which is used to ensure that models generalize
well and do not overfit the training data (Fig. 5A). On each fold, only 60% of the
training traces were shown to the network in the training phase (Fig. 5A, white
boxes), the network optimized on 20% of data used as a validation set (Fig. 5A,
yellow boxes), and the generalizability of the trained network assessed using the
remaining 20% (Fig. 5A, blue boxes). Each of the 5 folds had a different 60-20-20
split, such that the training and cross-validation datasets had a balance of traces
from both classes.
646
We tested several different machine learning models on this classification problem -
Support Vector Machines (SVMs), Deep Neural Networks (DNN), Convolutional
Neural Networks (CNNs) and convolutional recurrent neural networks (CNN-LSTM,
i.e. a Convolutional Neural Network with Long Short-Term Memory). The prediction
accuracies and F1 scores (a metric that adjusts for class imbalances as well as
whether a prediction was a true/false positive/negative, defined in the methods
section) for the various models tested are shown in Table 2. Of these models, the
1D-CNN and CNN-LSTM models performed best across all folds. However, the
former only predicts from snapshots of 10s-long snippets of the calcium trace, and
the networks had more erroneous performance when predicting on the entire time
series (data not shown). The CNN-LSTM model, however, also incorporates a
recurrent neural network module (Hochreiter and Schmidhuber, 1997), which has the
ability to learn features across an entire sequence and not just across neighbouring
positions in a trace sequence, like CNNs do.
661
662   The architecture of the network is shown in Fig. 5B. The input to the network is a
663   trace sampled at 30Hz, which is some multiple of 10 seconds in duration. It is first
664   remapped to values between 0 and 1 using min-max normalization and then

processed by a CNN, which acts as a local feature extractor that also down samples the input data, such that it produces 4 vector outputs at a 1Hz sampling frequency. These outputs are passed to a bidirectional LSTM, one time-step at a time. A forward LSTM cell looks at trends in the forward direction (Fig. 5B, blue dotted box and arrow), while a reverse LSTM cell goes through the data in the reverse direction (Fig. 5B, red dotted box and arrow). Each of the two cells produces a "hidden state" as its output. Given that the LSTM is a recurrent neural network, this output is fed back into the network along with the next appropriate time step from the CNN's output, a process which is repeated 9 more times (for every 10 second trace). In essence, the LSTM is detecting trends across the entire sequence. The final hidden states from the forward and reverse LSTMs, produced after passing through the entire trace, are retained and concatenated. These outputs are combined by passing them through a linear layer, which performs a matrix multiplication to produce an output that is 10 samples long. At this step, each sample represents the likelihood that the trace in question comes from a cell in the bursting mode, predicted at 1 second intervals.

The likelihood of each time step having come from the tonic state can then be inferred (Fig. 5B, grayscale boxes on the right). The state with the larger likelihood score is taken to be the final classification at each time step.

Five networks were produced, one for each fold of cross-validation. The median accuracy was 80.15% for the cross-validation phase and 82.02% for the test phase (Fig. 5C). The median F1-score was 0.7302 for the cross-validation phase and 0.8071 for the test phase (Fig. 5C). Next, we looked at the Receiver Operating Characteristic (ROC) curve for each network fold. This curve plots the false positive rate against the true positive rate of the classifier for various classification threshold values, and the area under this curve is an indicator of the quality of the classifier. The area under the ROC curve (AUC) for each of the five network folds is plotted in Fig 5E. The AUC values for all folds on both cross-validation and test data are much higher than 0.5, indicating that these networks have been trained well and make accurate predictions.
Of all the network folds, Fold 2 looked particularly promising, because it (1) consistently outperformed the other folds with respect to classification accuracy and F1-scores (Fig 5C, D, purple), and (2) had higher AUC values than the other network folds (Fig 5E,F). Thus, we decided to retain this network for future use, and have dubbed it CaMLsort (**Ca**lcium imaging and **M**achine **L**earning based tool to **sort** intracellular state).

CaMLsort performs well on previously unseen calcium signal reconstructions
To be sure that networks weren't performing well just because of peculiarities in the dataset used for training and testing, we presented an independent set of calcium signal reconstructions to the network that were not used in any phase of the training. These reconstructions (n=19) also had state switches in them. The ground truth class labels for each of the 19 cells (Fig. 6A, top) were compared with the raw predictions from the network (Fig. 6A, middle).

At first glance, it appears that the network isn't making accurate calls, because each cell's predicted state alternates between both states. However, these fluctuations are

transient and the overall F1-scores for the raw predictions are >0.5 for 13 out of the
19 cells (median score 0.777). The distribution of F1-scores is shown in Fig. 6C. On
closer observation, it is clear that the 6 cells for which F1-scores were low (< 0.5)
had a switch in state during the recording. The poor prediction accuracy for these
can be explained by the fact that the training dataset didn't have any cases with state
switches.
721
Given our prior knowledge that the state of a neuron doesn't flicker rapidly between
tonic and bursting, we decided to smoothen out the raw predictions using a majority
voting strategy. For this, we took a rolling window of length N around each time step
of the prediction and asked what the majority of the network calls were across all the
windows that included the time step. If the average posterior scores for the majority
class were higher than the average posterior scores for the minority class, we would
confidently assign the majority class to that time step. If not, no confident call would
be made for that time step. For example, to make a confident call at time t_0 with a
window length of 5, we first extract all the 5-step windows that include time t_0 (there
will be 5 such windows, except at the start and end, where there will be fewer). Next,
we average the posterior scores within each of the windows and assign the 'average
state' as the state with the higher posterior score. In this case, let's say these were
2:3 in favour of bursting. If the average bursting posterior score for the 3 "bursting"
windows was greater than the average tonic posterior score for the 2 "tonic"
windows, the state for time t_0 would be confidently assigned as "bursting". If not, the
prediction at time t_0 would be left blank.
738
Using this majority voting strategy with a window length of 7, we were able to make
confident calls for >90% of the duration for all 19 cells. As seen in Fig. 6B, the jitter in
the raw CaMLsort predictions was considerably smoothened out by this process, and
the F1-scores showed a marginal improvement as well, with a median F1-score of
0.8194 (Fig. 6C). As before, the cases where F1-scores were <0.5 corresponded to
recordings with state switches. As an extra control, we obtained CaMLsort
predictions from signals that were scrambled. These predictions had much lower F1-
scores (Fig. 6C). In fact, when presented with scrambled traces, CaMLsort almost
always (99.76% of the time) predicted the state to be tonic, with a very large,
consistent differential between the posterior scores for the tonic (mean ± stdev:
0.7810 ± 0.0565) and bursting (mean ± stdev: 0.2190 ± 0.0565) classes. This implies
that CaMLsort has, indeed, captured a latent trend present only in the true signal and
not in a collection of random numbers of comparable magnitude.
752
753 **CaMLsort performs well on calcium signal reconstructions from simulated**
754 **spike trains**
755
As a further challenge to the trained networks, we decided to generate artificial spike
trains that represent canonical cases of the tonic or bursting class, reconstruct the
calcium signal for these spike trains and ask the network to classify these completely
synthetic data (Fig. 7). We simulated Purkinje neuron simple spiking using either a
Poisson model (tonic state) or by randomly sampling from the distribution of inter-
event intervals from experiments (bursting state; also see Methods). The CF input
spike train was considered to be independent and was generated using a Poisson
model.The ground truth states for these simulated spike trains is shown in Fig. 7A
(top trace). The raw CaMLsort predictions obtained for these data had a median F1-

score 0.8571 (Fig. 7A, middle trace and 7C), although these too, had several
reversals in the state predictions. On applying the majority voting strategy described
above, however, these reversals were smoothened out (Fig. 7B), yielding a median
F1-score of 0.9189 (Fig. 7C) and a confident prediction for at least 80% of the
recording duration in every cell. As was observed in the previous case, scrambling
the signal prior to prediction by the neural network were all of one class (tonic),
yielding F1-scores which were either 1 (all right) or 0 (all wrong) (Fig. 7C).
772
773

Neural networks trained only on simulated calcium signals correctly
identify Purkinje neuron state from real data
Thus far, all the inputs and challenges to the network were made using calcium
signal reconstructions. To test the applicability of these networks on data from
experiments, we went back to our original simultaneous electrophysiology-imaging
data and asked if the network could call the state of those cells correctly based on
the dF/F traces alone.
781
We found that CaMLsort predicted the state of a cell from imaging data well (Fig.
8A), with a median F1-score of 0.7654 across all cells (Fig. 8C). This outcome is
interesting, because the networks have only been exposed to simulated calcium
signals generated using a first-order generative model in the training and test
phases, and yet are able to reasonably assess the state from real data, which have
several challenging features - variable sensor expression levels, shot noise and
signal bleaching, to name a few.
789
790 Majority voting did not substantially improve the overall F1-score across all cells
791 (median 0.75). However, by retaining only confident calls, the flip-flopping of network
792 predictions was markedly reduced (Fig. 8B), as was observed before in the case of
793 classifications using reconstructed calcium signals.
794
To benchmark the quality of these predictions, we obtained CaMLsort predictions
from calcium signals reconstructed from the spike trains of these cells (since
CaMLsort was trained only on simulated calcium signals). The prediction quality from
reconstructions was comparable to that obtained from raw imaging data (median F1-
score of 0.9011; Fig. 8C). Therefore, while CaMLsort is clearly able to classify
cellular state from experimentally-obtained imaging data, its performance has room
for improvement. Also consistent with our previous results were the predictions from
scrambled dF/F traces, which were almost all (96.83%) tonic, thereby yielding F1-
scores that were mostly either 0 or 1 (Fig. 8C).
804
805 To sum up, we showed that CaMLsort has learnt to sort the cellular state of a
806 Purkinje neuron based only on its calcium signal and is capable of producing
807 confident calls of high quality.
808
809

CaMLsort generalizes well to calcium imaging data from other systems
Having developed a tool that can reasonably predict the bistable state of a Purkinje
neuron given its calcium signal, we asked whether it could be broadly applicable to
other neurons, too. Dopaminergic neurons of the ventral tegmental area (VTA) in
mice also fire either tonically or phasically (Grace and Bunney, 1984a, b). Fleming et

al., 2021 performed simultaneous electrophysiology and calcium imaging from these
neurons, which expressed either GCaMP6f or GCaMP6m. This dataset is available
publicly (https://github.com/jewellsean/spike_tools) and we tested CaMLsort on this
dataset.
819
We first compiled the dataset and sorted cells as tonic or bursting, using a
combination of visual inspection of spike rasters (Fig. 9A, B) and the coefficient of
variation of inter-spike intervals (Fig. 9C), as we did for Purkinje neurons. Three
independent reviewers used this method to classify states, and a consensus was
taken across reviewers, while also taking into consideration the state labels provided
by the authors in their metadata. Data were included only if at least 2 out of the 4
labels (3 reviewers plus original metadata) agreed. Using this method, 78/84 cells
(92.86%) of the original dataset were included for further analysis. Of these, 16 cells
were in the tonic state, while the remaining 62 cells were in the bursting state.
829
Before trying to use CaMLsort on this data, we asked whether the convolution-based
forward model of reconstructing calcium signals from electrophysiology would work
on this dataset. Since there is only a single kind of spike in these neurons, we used
only one kernel, the amplitude, half-rise time constant and half-decay time constant
of which were taken to be the mean values for GCaMP6f or GCaMP5G (Chen et al.,
2013). No optimization of kernels was performed.
836
Visual inspection shows that reconstructions obtained from both GCaMP6f and
GCaMP5G kernels were comparable to the signals imaged from VTA DA neurons,
with the only difference between them being their amplitudes (Fig. 9D). To quantify
the similarity between the ground truth and the two simulated calcium signals, we
used the Pearson correlation coefficient. The correlation coefficients between the
ground truth and reconstructions from both GCaMP6f and GCaMP5G kernels were
nearly identical, with a value of 0.6400 ± 0.3189, on average, for both (Fig. 9E).
Reconstructions that were either scrambled or generated after shuffling the DA
neuron's spike times had no correlation with the ground truth signal, as expected
(Fig. 9E).
847
Having confirmed that the Fleming et al. dataset had both tonic and bursting data,
and that the mapping from electrophysiology to imaging was also possible with this
data, we were ready to test CaMLsort on it. As before, we first interpolated the
calcium signals to get a sampling frequency of 30Hz before inputting them to
CaMLsort. Raw predictions from CaMLsort and their corresponding ground truth
labels for a subset of 12 of the 78 cells are shown in Fig. 10A. The distribution of F1-
scores obtained from all 78 cells, the median for which is 0.6841, is shown in Fig.
10B.
856
We also calculated F1-scores for predictions made on calcium signals reconstructed
from true electrophysiology data using the GCaMP5G kernel (Fig. 10B). We decided
to use the 5G kernel since the reconstructions that CaMLsort was trained on
followed kinetics of GCaMP5G. The predictions from signal reconstructions were
comparable to those from raw imaging data, with a median of 0.8889. Lastly, if the
imaging signal was scrambled before prediction, CaMLsort predicted a default "tonic"
state 89.03% of the time, yielding F1-scores that were mostly either 0 or 1 (Fig. 10B).
864

865 In all, our results from VTA DA neurons in mice were similar to those from zebrafish
866 Purkinje neurons, which means that CaMLsort generalizes well and is likely to find
867 broad use in classifying cellular state across systems. In order to facilitate this use,
868 we have made CaMLsort publicly available as a GitHub package
869 (https://github.com/vatsalat/CaMLsort23).
870

Discussion
In this paper, we describe the development of CaMLsort, a machine learning-based
tool to identify the bistable state of Purkinje neurons from calcium imaging data. We
develop this network and train it only on a simulated calcium signal dataset, but find
that it can be used on experimental data, too. Furthermore, we demonstrate that
CaMLsort is able to classify bistable states in a completely different cell type
(dopaminergic neurons) from a different brain region (ventral tegmental area) in a
different model system (mice), and using a different version of the calcium sensor
(GCaMP6) than it was originally trained on. While this demonstrates that CaMLsort
could have wider application, more broader testing on other datasets will be
essential to prove generalizability.
882

Bistability in Purkinje neuron populations
The ability of neurons to exist in two stable states and exhibit distinct firing patterns/
excitabilities in those states has been documented in invertebrate (Dickinson and
Nagy, 1983; Lechner et al., 1996; Marder et al., 1996; Russell and Hartline, 1982)
and vertebrate nervous systems (Cowan and Wilson, 1994; Hounsgaard et al., 1984;
Loewenstein et al., 2005; Misgeld et al., 1986; Schwindt and Crill, 1980).
890

891 We showed that larval zebrafish Purkinje neurons are bistable and that they switch
892 between firing either tonically or in bursts (Sengupta and Thirumalai, 2015). Such
893 behavior has also been reported in many other cell types such as thalamic (Jahnsen
894 and Llinás, 1984) and cortical neurons (McCormick et al., 1985).
895

Though it is appreciated that bistability increases the information processing capacity
of neurons, such as by allowing gating of synaptic inputs (Thirumalai and Jha, 2022),
its implication for network computation or behavioural output is less well understood.
This is true of Purkinje neurons as well - while it is established that these neurons
are bistable and are important for motor behaviours (Engbers et al., 2013; Popa et
al., 2018; Sengupta and Thirumalai, 2015), the role of bistability in Purkinje neuron
computation or motor behaviour output has thus far not been clear.
903

We propose that having more than one stable state allows a cell to toggle in and out
of participating in circuit computations. In the cerebellum, specifically, bistability may
have significant consequences by modulating PN neuronal synchrony with other
PNs. PNs make weak inhibitory synapses onto deep cerebellar nuclear (DCN) cells
in mammals and eurydendroid cells in fish, with a high degree of convergence (Bae
et al., 2009; De Zeeuw et al., 2011; Harmon et al., 2020; Ikenaga et al., 2006;
Person and Raman, 2012). DCN cell spike times could be effectively controlled via
inhibition from several PNs firing synchronously (Person and Raman, 2012). Since
PNs fire tonically when synaptically-isolated and CF inputs can trigger bursts
(Sengupta and Thirumalai, 2015), the bursting state is a network-embedded state

while the tonic state is independent of network drive. By toggling between these two
states PNs can increase or decrease the levels of population synchrony and thereby
affect DCN firing differentially. Mapping state distribution across the PN population
during various tasks will lead us to an understanding of how PNs synchronize or
desynchronize during these tasks and how that in turn shapes cerebellar output and
motor behaviors.
920
Inferring long-term dynamics from calcium imaging data
Calcium imaging has been one of the core experimental techniques in neuroscience
for more than three decades and has seen many major developments in sensor
quality, microscopy and analytical algorithms and software (Chen et al., 2013; Göbel
and Helmchen, 2007; Peron et al., 2015; Renninger and Orger, 2013; Tian et al.,
2009; Yang and Yuste, 2017). Yet, the technique is largely used to identify regions of
the brain that respond to presented stimuli or correlate with behavioral output under
well-defined experimental conditions. This is resultant from the fact that the
predominant mode of transient calcium elevation occurs due to the opening of
voltage-gated calcium channels during action potential firing (Borst and Helmchen,
1998; Göbel and Helmchen, 2007). Thus, these calcium transients are quite brief,
typically lasting only tens of milliseconds and report spiking activity of neuronal
populations during sensory stimulation and/or motor output. However, it is clear that
neurons also exhibit longer time scale dynamics such as bursting, which carries
information for downstream neurons (Zeldenrust et al., 2018). As the currently
available calcium sensors themselves are much slower compared to the calcium
transients, it is obvious that the recorded dF/F signals must manifest signatures of
such long-term dynamics. CaMLsort addresses this issue by looking at longer term
signatures in the calcium imaging time series data to identify cellular state.
940
Broader Applications of CaMLsort
The development of CaMLsort has opened up the possibility of a lot of future
research in understanding PN bistability and its role in the function of the cerebellum
as a whole. We can now experimentally address, for instance, what causes a switch
in a PN's state by doing a high-throughput screen for chemical agents that affect
bistability. We know that while bistability is a cell-intrinsic property, state switches
could be a result of modulation of network activity. Neuromodulators are key
candidates for this, as they can alter cell-intrinsic properties as well as network
activity, and could thereby affect the bistability of a neuron, as has been
demonstrated in multiple experimental systems (Abbinanti et al., 2012; Conway et
al., 1988; Hounsgaard et al., 1988; Lechner et al., 1996; Williams et al., 2002).
Another question we can now directly address is how the state of a cell changes
over development. We hypothesize that the tonic state could be a developmentally
early state, with the burst-like phenotype developing as they get integrated into the
developing cerebellar circuit. By birth-dating PNs using a photoconvertible protein
like Kaede and imaging their activity with GCaMP, we can use CaMLsort to ask
whether younger cells tend to be more tonic or vice versa. While this question could
potentially be addressed using electrophysiology, CaMLsort allows us to perform this
experiment in a more high-throughput and minimally invasive fashion.
960
A number of cell types exhibit bistability including cortical neurons (Cowan and
Wilson, 1994; Egorov et al., 2002), striatal neurons (Misgeld et al., 1986), thalamic
neurons (Fuentealba et al., 2005) and spinal motoneurons (Hounsgaard et al.,

1984). In all of these cases, cellular state dictates how synaptic inputs are integrated.
Therefore, predicting cellular state is essential to understanding circuit computation
and here CaMLsort will be immensely useful. It is natural to wonder whether
CaMLsort will be equally effective in predicting cellular state in different cell types
and in different model organisms. We have found that CaMLsort generalizes well to
a completely different system than it was trained on - neurons from a different brain
region (ventral tegmental area) in a different model organism (mouse) imaged using
a later generation of GCaMP. We would caution against using CaMLsort right out of
the box without validating its efficacy in the new system of interest using a small
ground truth dataset.
974
975   Taken together, CaMLsort is a powerful tool that opens up new avenues of research
976   into the functioning of any circuit with bistable neurons.
977

Limitations of CaMLsort

Having said the above, CaMLsort is not free of limitations. For instance, the
generative model we use to train the neural network is a linear, time-invariant model
and results may improve with the use of more accurate models, such as ones that
estimate the true increase in intracellular calcium concentration resulting from simple
spikes and CF inputs. Furthermore, it could be that CaMLsort may not work
independent of the imaging method used. In both the cases we tested CaMLsort,
widefield epifluorescence microscopy was used. The signal and noise characteristics
of other, more modern and popular, imaging methods like multiphoton and light
sheet microscopy are different from traditional widefield imaging. However, we posit
that this should also be easy to resolve, either by the acquisition of a small ground
truth dataset, or by comparing the traces obtained from imaging to those in the
reconstructions dataset, or by denoising the data prior to prediction using CaMLsort.
991
992

References

Abbinanti MD, Zhong G, Harris-Warrick RM. 2012. Postnatal emergence of serotonin-induced plateau potentials in commissural interneurons of themouse spinal cord. *Journal of Neurophysiology* **108**:2191–2202. doi:10.1152/jn.00336.2012

Ahrens MB, Orger MB, Robson DN, Li JM, Keller PJ. 2013. Whole-brain functional imaging at cellular resolution using light-sheet microscopy. *Nat Methods* **10**:413–420. doi:10.1038/nmeth.2434

Ali F, Kwan AC. 2020. Interpreting in vivo calcium signals from neuronal cell bodies, axons, and dendrites: a review. *Neurophotonics* **7**:011402. doi:10.1117/1.NPh.7.1.011402

Armstrong DM, Edgley SA. 1984. Discharges of Purkinje cells in the paravermal part of the cerebellar anterior lobe during locomotion in the cat. *J Physiol (Lond)* **352**:403–424.

Bae Y-K, Kani S, Shimizu T, Tanabe K, Nojima H, Kimura Y, Higashijima S, Hibi M. 2009. Anatomy of zebrafish cerebellum and screen for mutations affecting its development. *Dev Biol* **330**:406–426. doi:10.1016/j.ydbio.2009.04.013

Borst JG, Helmchen F. 1998. Calcium influx during an action potential. *Methods Enzymol* **293**:352–371. doi:10.1016/s0076-6879(98)93023-3

Chen T-W, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, Schreiter ER, Kerr RA, Orger MB, Jayaraman V, Looger LL, Svoboda K, Kim DS. 2013. Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* **499**:295–300. doi:10.1038/nature12354

Conway BA, Hultborn H, Kiehn O, Mintz I. 1988. Plateau potentials in alpha-motoneurones induced by intravenous injection of L-dopa and clonidine in the spinal cat. *J Physiol (Lond)* **405**:369–384. doi:10.1113/jphysiol.1988.sp017337

Cowan RL, Wilson CJ. 1994. Spontaneous firing patterns and axonal projections of single corticostriatal neurons in the rat medial agranular cortex. *Journal of Neurophysiology* **71**:17–32. doi:10.1152/jn.1994.71.1.17

De Zeeuw CI, Hoebeek FE, Bosman LWJ, Schonewille M, Witter L, Koekkoek SK. 2011. Spatiotemporal firing patterns in the cerebellum. *Nature ReviewsNeuroscience* **12**:327–344. doi:10.1038/nrn3011

Dickinson PS, Nagy F. 1983. Control of a central pattern generator by an identified modulatory interneurone in crustacea. II. Induction and modification of plateau properties in pyloric neurones. *Journal of Experimental Biology* **105**:59–82. doi:10.1242/jeb.105.1.59

Egorov AV, Hamam BN, Fransén E, Hasselmo ME, Alonso AA. 2002. Graded persistent activity in entorhinal cortex neurons. *Nature* **420**:173–178. doi:10.1038/nature01171

Engbers JDT, Fernandez FR, Turner RW. 2013. Bistability in Purkinje neurons: Ups and downs in cerebellar research. *Neural Networks*, Computation in the Cerebellum **47**:18–31. doi:10.1016/j.neunet.2012.09.006

Fleming W, Jewell S, Engelhard B, Witten DM, Witten IB (2021) Inferring spikes from calcium imaging in dopamine neurons. PLOS ONE 16:e0252345.

Fuentealba P, Timofeev I, Bazhenov M, Sejnowski TJ, Steriade M. 2005. Membrane
Bistability in Thalamic Reticular Neurons During Spindle Oscillations. *Journal of*
Neurophysiology **93**:294–304. doi:10.1152/jn.00552.2004
Göbel W, Helmchen F. 2007. In Vivo Calcium Imaging of Neural Network Function.
Physiology **22**:358–365. doi:10.1152/physiol.00032.2007
Grewe BF, Helmchen F. 2009. Optical probing of neuronal ensemble activity. *Current*
Opinion in Neurobiology, Neuronal and glial cell biology ● New technologies
19:520–529. doi:10.1016/j.conb.2009.09.003
Grienberger C, Konnerth A. 2012. Imaging Calcium in Neurons. *Neuron* **73**:862–885.
doi:10.1016/j.neuron.2012.02.011
Harmon TC, McLean DL, Raman IM. 2020. Integration of Swimming-Related
Synaptic Excitation and Inhibition by olig2+ Eurydendroid Neurons in Larval
Zebrafish Cerebellum. *J Neurosci* **40**:3063–3074.
doi:10.1523/JNEUROSCI.2322-19.2020
He K, Zhang X, Ren S, Sun J. 2016. Deep Residual Learning for Image
Recognition2016 IEEE Conference on Computer Vision and Pattern Recognition
(CVPR). Presented at the 2016 IEEE Conference on ComputerVision and Pattern
Recognition (CVPR). pp. 770–778. doi:10.1109/CVPR.2016.90
Hochreiter S, Schmidhuber J. 1997. Long Short-Term Memory. *Neural Computation*
9:1735–1780. doi:10.1162/neco.1997.9.8.1735
Hounsgaard J, Hultborn H, Jespersen B, Kiehn O. 1988. Bistability of alpha-
motoneurones in the decerebrate cat and in the acute spinal cat after intravenous
5-hydroxytryptophan. *The Journal of Physiology* **405**:345–367.
doi:10.1113/jphysiol.1988.sp017336
Hounsgaard J, Hultborn H, Jespersen B, Kiehn O. 1984. Intrinsic membrane
properties causing a bistable behaviour of alpha-motoneurones. *Exp BrainRes*
55:391–394. doi:10.1007/BF00237290
Ikenaga T, Yoshida M, Uematsu K. 2006. Cerebellar efferent neurons in teleost fish.
Cerebellum **5**:268–274. doi:10.1080/14734220600930588
Jahnsen H, Llinás R. 1984. Electrophysiological properties of guinea-pig thalamic
neurones: an in vitro study. *The Journal of Physiology* **349**:205–226.
doi:10.1113/jphysiol.1984.sp015153
Kerr JND, Greenberg D, Helmchen F. 2005. Imaging input and output of neocortical
networks in vivo. *Proceedings of the National Academy of Sciences* **102**:14063–
14068. doi:10.1073/pnas.0506029102
Kingma DP, Ba J. 2017. Adam: A Method for Stochastic Optimization.
doi:10.48550/arXiv.1412.6980
Kitamura K, Häusser M. 2011. Dendritic calcium signaling triggered by spontaneous
and sensory-evoked climbing fiber input to cerebellar Purkinje cells in vivo. *J*
Neurosci **31**:10847–10858. doi:10.1523/JNEUROSCI.2525-10.2011
Knogler LD, Kist AM, Portugues R. 2019. Motor context dominates output from
purkinje cell functional regions during reflexive visuomotor behaviours. *eLife*
8:e42138. doi:10.7554/eLife.42138
Kwan AC, Dan Y. 2012. Dissection of Cortical Microcircuits by Single-Neuron
Stimulation In Vivo. *Current Biology* **22**:1459–1467.
doi:10.1016/j.cub.2012.06.007
Lechner HA, Baxter DA, Clark JW, Byrne JH. 1996. Bistability and its regulation by
serotonin in the endogenously bursting neuron R15 in Aplysia. *J Neurophysiol*
75:957–962.

Lev-Ram V, Miyakawa H, Lasser-Ross N, Ross WN. 1992. Calcium transients in cerebellar Purkinje neurons evoked by intracellular stimulation. *Journal of Neurophysiology* **68**:1167–1177. doi:10.1152/jn.1992.68.4.1167

Lin B-J, Chen T-W, Schild D. 2007. Cell type-specific relationships between spiking and [Ca2+]i in neurons of the Xenopus tadpole olfactory bulb. *The Journal of Physiology* **582**:163–175. doi:10.1113/jphysiol.2006.125963

Loewenstein Y, Mahon S, Chadderton P, Kitamura K, Sompolinsky H, Yarom Y, Häusser M. 2005. Bistability of cerebellar Purkinje cells modulated by sensory stimulation. *Nat Neurosci* **8**:202–211. doi:10.1038/nn1393

Marder E, Abbott LF, Turrigiano GG, Liu Z, Golowasch J. 1996. Memory from the dynamics of intrinsic membranecurrents. *Proceedings of the National Academy of Sciences* **93**:13481–13486. doi:10.1073/pnas.93.24.13481

Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C. 2004. Interneurons of the neocortical inhibitory system. *Nat Rev Neurosci* **5**:793–807. doi:10.1038/nrn1519

Matsui H, Namikawa K, Babaryka A, Köster RW. 2014. Functional regionalization of the teleost cerebellum analyzed in vivo. *Proc Natl Acad Sci USA* **111**:11846–11851. doi:10.1073/pnas.1403105111

McCormick DA, Connors BW, Lighthall JW, Prince DA. 1985. Comparative electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. *Journal of Neurophysiology* **54**:782–806. doi:10.1152/jn.1985.54.4.782

Misgeld U, Calabresi P, Dodt HU. 1986. Muscarinic modulation of calcium dependent plateau potentials in rat neostriatal neurons. *Pflugers Arch* **407**:482–487. doi:10.1007/BF00657504

Miyakawa H, Lev-Ram V, Lasser-Ross N, Ross WN. 1992. Calcium transients evoked by climbing fiber and parallel fiber synaptic inputs in guinea pig cerebellar Purkinje neurons. *J Neurophysiol* **68**:1178–1189. doi:10.1152/jn.1992.68.4.1178

Moreaux L, Laurent G. 2007. Estimating firing rates from calcium signals in locust projection neurons in vivo. *Frontiers in Neural Circuits* **1**.

Narayanan S, Thirumalai V. 2019. Contributions of the Cerebellum for Predictive and Instructional Control of Movement. *Curr Opin Physiol* **8**:146–151. doi:10.1016/j.cophys.2019.01.011

Paszke A, Gross S, Massa F, Lerer A, Bradbury J, Chanan G, Killeen T, Lin Z, Gimelshein N, Antiga L, Desmaison A, Kopf A, Yang E, DeVito Z, Raison M, Tejani A, Chilamkurthy S, Steiner B, Fang L, Bai J, Chintala S. 2019. PyTorch: An Imperative Style, High-Performance Deep Learning LibraryAdvances in Neural Information Processing Systems. CurranAssociates, Inc.

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Prettenhofer P, Weiss R, Dubourg V, Vanderplas J, Passos A, Cournapeau D, Brucher M, Perrot M, Duchesnay É. 2011. Scikit-learn: Machine Learning in Python. *Journal of Machine Learning Research* **12**:2825–2830.

Peron S, Chen T-W, Svoboda K. 2015. Comprehensive imaging of cortical networks. *Current Opinion in Neurobiology*, Large-Scale Recording Technology (32) **32**:115–123. doi:10.1016/j.conb.2015.03.016

Person AL, Raman IM. 2012. Purkinje neuron synchrony elicits time-locked spikingin the cerebellar nuclei. *Nature* **481**:502–505. doi:10.1038/nature10732

Popa LS, Streng ML, Ebner TJ. 2018. Purkinje Cell Representations of Behavior: Diary of a Busy Neuron. *Neuroscientist* 1073858418785628. doi:10.1177/1073858418785628

Ramirez JE, Stell BM. 2016. Calcium Imaging Reveals Coordinated Simple Spike Pauses in Populations of Cerebellar Purkinje Cells. *Cell Reports* **17**:3125–3132. doi:10.1016/j.celrep.2016.11.075

Renninger SL, Orger MB. 2013. Two-photon imaging of neural population activity in zebrafish. *Methods*, Zebrafish Methods **62**:255–267. doi:10.1016/j.ymeth.2013.05.016

Rupprecht P, Carta S, Hoffmann A, Echizen M, Blot A, Kwan AC, Dan Y, Hofer SB, Kitamura K, Helmchen F, Friedrich RW. 2021. A database and deep learning toolbox for noise-optimized, generalized spike inference from calcium imaging. *Nat Neurosci* **24**:1324–1337. doi:10.1038/s41593-021-00895-5

Russell DF, Hartline DK. 1982. Slow active potentials and bursting motor patterns in pyloric network of the lobster, Panulirus interruptus. *Journal of Neurophysiology* **48**:914–937. doi:10.1152/jn.1982.48.4.914

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**:676–682. doi:10.1038/nmeth.2019

Schwindt P, Crill W. 1980. Role of a persistent inward current in motoneuron bursting during spinal seizures. *Journal of Neurophysiology* **43**:1296–1318. doi:10.1152/jn.1980.43.5.1296

Sengupta M. 2015. Purkinje neurons in action: From single cells to ensembles. Mumbai: Tata Institute of Fundamental Research.

Sengupta M, Thirumalai V. 2015. AMPA receptor mediated synaptic excitation drives state-dependent bursting in Purkinje neurons of zebrafish larvae. *Elife* **4**. doi:10.7554/eLife.09158

Smetters D, Majewska A, Yuste R (1999) Detecting action potentials in neuronal populations with calcium imaging. Methods San Diego Calif 18:215–221.

Thirumalai V, Jha U. 2022. RECRUITMENT OF MOTONEURONS Vertebrate Motoneurons, In Press. Springer-Verlag.

Tian L, Hires SA, Mao T, Huber D, Chiappe ME, Chalasani SH, Petreanu L, Akerboom J, McKinney SA, Schreiter ER, Bargmann CI, Jayaraman V, Svoboda K, Looger LL. 2009. Imaging neural activity in worms, flies and micewith improved GCaMP calcium indicators. *Nat Methods* **6**:875–881. doi:10.1038/nmeth.1398

Urasaki A, Morvan G, Kawakami K. 2006. Functional dissection of the Tol2 transposable element identified the minimal cis-sequence and a highly repetitive sequence in the subterminal region essential for transposition.*Genetics* **174**:639–649. doi:10.1534/genetics.106.060244

Vogelstein JT, Packer AM, Machado TA, Sippy T, Babadi B, Yuste R, Paninski L. 2010. Fast Nonnegative Deconvolution for Spike Train Inference From Population Calcium Imaging. *Journal of Neurophysiology* **104**:3691–3704. doi:10.1152/jn.01073.2009

Williams SR, Christensen SR, Stuart GJ, Häusser M. 2002. Membrane potential bistability is controlled by the hyperpolarization-activated current I(H) in rat cerebellar Purkinje neurons in vitro. *J Physiol (Lond)* **539**:469–483.

Yaksi E, Friedrich RW. 2006. Reconstruction of firing rate changes across neuronal populations by temporally deconvolved Ca2+ imaging. *Nat Methods* **3**:377–383. doi:10.1038/nmeth874

Yang W, Yuste R. 2017. In vivo imaging of neural activity. *Nat Methods* **14**:349–359. doi:10.1038/nmeth.4230

Zeldenrust F, Wadman WJ, Englitz B. 2018. Neural Coding With Bursts—Current State and Future Perspectives. *Frontiers in Computational Neuroscience* **12**.

Additional Information:

Data Availability:
All data are available in the main text and figures. The CaMLsort package codes are available on Github: https://github.com/vatsalat/CaMLsort23

Competing Interests:
The authors declare no competing interests.

Author Contributions:
Conceptualization: AV and VT
Investigation: AV, SU, MS, PKG and VT
Analysis: AV, SU, MS, PKG and VT
Writing: AV and VT

Funding:
The authors would like to thank the following sources of funding support: Wellcome Trust-DBT India Alliance Intermediate (VT; 500040/Z/09/Z) and Senior fellowships (VT; IA/S/17/2/503297), Department of Biotechnology (VT; BT/PR4983/MED/30/790/2012), Science and Engineering Research Board, Department of Science and Technology (VT; EMR/2015/000595), Department of Atomic Energy (VT) and Department of Science and Technology, Govt. of India (PKG).

Acknowledgements:
The authors would also like to thank P.T. Jagadeesh for maintaining the fish facility and members of the Fishfolk lab at NCBS for their inputs at various stages of this project. In particular, we would like to thank Ms. Meha Jadhav and Ms. Shivangi Verma for their help with independently verifying parts of this manuscript. The authors would also like to thank Prof. Rohini Balakrishnan for initiating this collaboration.

1225 **Table 1 - Number of recordings compiled for the state-labelled imaging dataset**

	Extracellular	Intracellular	**Total**
Bursting	39	39	**78**
Tonic	11	42	**53**
Switch	0	5	**5**
Ambiguous	0	25	**25**
Poor Quality	0	12	**12**
Total	**50**	**123**	**173**

1226
1227

1228 **Table 2 - Prediction accuracies and F1-scores for various models trained**
1229 **tosolve the classification problem.**

Model	Fold	Validation		Test	
		Accuracy (%)	F1-score	Accuracy(%)	F1-score
Support Vector Machine (SVM)	1	81.41	0.7206	70.19	0.6077
	2	72.68	0.593	79.66	0.6842
	3	75.84	0.703	72.55	0.6055
	4	73.10	0.6422	73.65	0.671
	5	70.67	0.629	74.89	0.6755
	Mean ± Stdev	74.74 ± 4.16	0.6576 ± 0.0531	74.19 ± 3.51	0.6488 ± 0.0388
Deep Neural Network	1	83.91	0.7893	72.99	0.6649
	2	77.71	0.7205	82.89	0.7849
	3	74.85	0.7099	76.10	0.6788
	4	73.31	0.6535	75.12	0.7143
	5	77.82	0.7060	75.24	0.6629
	Mean ± Stdev	77.52 ± 4.06	0.7158 ± 0.0486	76.47 ± 3.77	0.7012 ± 0.0511
1D CNN	1	87.20	0.8163	84.00	0.7933
	2	80.67	0.7541	88.33	0.8336
	3	82.66	0.7880	80.09	0.7263
	4	78.28	0.7185	81.07	0.7719
	5	82.60	0.7604	78.34	0.7004

1230

	Mean ± Stdev	82.28 ± 3.28	0.7675 ± 0.0369	82.37 ± 3.91	0.7651 ± 0.0530
1D CNN (3x)	1	87.54	0.8226	84.00	0.7893
	2	80.61	0.7535	86.97	0.8142
	3	80.61	0.7650	78.41	0.7089
	4	78.62	0.73	73.66	0.6592
	5	82.49	0.7592	76.83	0.6775
	Mean ± Stdev	81.97 ± 3.40	0.7661 ± 0.0343	79.97 ± 5.42	0.7298 ± 0.0686
CNN-LSTM	1	80.15	0.7227	79.7	0.7570
	2	**91.55**	**0.8788**	**92.75**	**0.8937**
	3	79.06	0.7233	92.12	0.887
	4	83.25	0.7878	75.9	0.7219
	5	79.9	0.7302	82.02	0.8071
	Mean ± Stdev	82.78 ± 5.15	0.7686 ± 0.0674	84.50 ± 7.57	0.8133 ± 0.0766

1231
1232

 Figures and Figure Legends

1234

Figure 1 - Cerebellar Purkinje neurons (PNs) in larval zebrafish exhibit
membrane potential bistability (A) Schematic of the cerebellar circuitry in larval
zebrafish (*Danio rerio*), showing the location of the cerebellum in the brain (left,
dashed line box) and the cell types and circuit architecture (right). Purkinje neurons

(green) are the principal neurons in this circuit and receive multiple inputs from
afferent sensorimotor nuclei outside the cerebellum via parallel fibers (black) and
climbing fibers (magenta). **(B)** Simple spikes, as observed intracellularly (left) and
extracellularly (right). Individual events (n=100, intracellular and n=100,
extracellular) are shown in cyan and have been aligned and superimposed on one
another. The average of these events is shown in black. **(C)** Climbing fiber (CF)
inputs, as observed intracellularly (left) and extracellularly (right). Individual events
(n=55 intracellular and n=10 extracellular) are shown in magenta, and their
respective averages in black. **(D)** Representative loose patch recordings from larval
zebrafish PNs showing the two modes of firing. The bursting mode recording is
shown on the left, and the tonic mode is shown on the right. In each mode,
electrical events can be distinguished by their amplitude, with small-amplitude
simple spikes marked in cyan and the large-amplitude CF inputs marked in
magenta. **(E)** Representative whole-cell current clamp recordings showing the two
modes of firing in PNs, with the same colour scheme for events as before. Cells
shown in D and E are different. **(F)** Example plots of simple spike inter-event
intervals from individual recordings. Five example cells are shown from each class
(Bursting and Tonic). The coefficient of variation of this distribution is marked under
each cell as "C.V.". **(G)** The distribution of inter-spike interval coefficients of
variation (n=131 cells), sorted by state (n=53 for tonic, and n=78 for bursting). The
distributions are significantly different, tested using a Mann-Whitney U-Test (p =
6.0733e-16).
1261

1262
Figure 2 – Both simple spikes and CF EPSPs contribute to the calcium signal
in PNs, irrespective of cellular state. (A) Schematic of the experimental design
and setup of simultaneous calcium imaging and electrophysiology in larval
zebrafish. **(B)** and **(C)** Representative traces obtained during simultaneous imaging
and electrophysiology for cells in the bursting (B) and tonic (C) mode. Simple
spikes and CF inputs in the trace are identified and marked in the raster above the

trace in cyan and magenta, respectively. The simultaneously-recorded change in
GCaMP fluorescence (dF/F) is shown below in green. Calcium transients
corresponding to simple spikes only ('*'), CF-events only ('#'), and both events ('o')
are marked. **(D)** The typical GCaMP5G response to a single action potential,
assuming it follows the profile of a difference of single exponentials with a half-rise
time of 100ms and half-decay time of 500ms. The vertical dashed lines indicate the
time of initiation of the calcium transient and the time when the transient peaks,
which corresponds to an interval of 201ms. Region I is the period up to 300ms
before the peak of the calcium signal, and region II is the period up to 300ms after
the peak of the transient. There were a total of n=171 transient peaks in the dF/F
signal detected across the N=15 cells. **(E)** Average simple spike counts in the
periods corresponding to regions I and II as defined in (D). Each cyan line
corresponds to an individual cell (N=15) and the average of all these lines is shown
in black. **(F)** Average CF input event counts in the periods corresponding to regions
I and II as defined in (D). Each magenta line corresponds to an individual cell
(N=15) and the average of all these lines is shown in black.
1285

1286

1287

**Figure 3 – Reconstruction of the calcium signal from electrophysiology
traces. (A)** Schematic showing the method used to reconstruct the calcium signal
from an electrophysiological trace. Simple spike (cyan) and CF input (magenta)
rasters (left) are independently convolved with calcium sensor kernels (cyan for

simple spikes, magenta for CF inputs) to generate an event-specific calcium signal
time series (middle). These are then simply added together to get the final
reconstruction (right).
(B) Comparison of ground truth (green) and reconstructed (purple) calcium signals
for 3 randomly chosen bursting cells (Bi) and 3 randomly chose tonic cells (Bii). The
Pearson's correlation coefficient between the two traces is marked above the plot for
each pair. **(C)** Correlation coefficients for all cells calculated for either the true
reconstruction, a scrambled version of the reconstruction, or the reconstruction for
when simple spike and CF input event times were shuffled. A Kruskal-Wallis test
yielded a p-value of 2.108×10^{-7}, which was followed by a post-hoc Dunn's test for
individual comparisons, the latter of which is marked on the figure (n=15 cells; 6
bursting, 9 tonic). **(E)** Distribution of residuals between the ground truth and
reconstructed calcium signal, pooled across all cells (n=15 cells). **(F)** The optimal
calcium sensor kernels for simple spikes (cyan) and CF inputs (magenta). The
formula for each kernel is mentioned above the plots, and the values for rise and
decay time constants are mentioned below the plot.
1308

1309
1310

1311
Figure 4 - Generating a labelled dataset of reconstructed calcium signal traces
(A) Flowchart showing how electrophysiological recordings from PNs were
processed to generate the state-labelled calcium signal database. **(B)**
Representative randomly sampled traces from the state-labelled reconstructed
calcium signal dataset. The left column has reconstructions from tonic cells and the

right column has reconstructions from bursting cells. Above each trace is the raster
showing the events in the source recording, with simple spikes in cyan, and CF
inputs in magenta. **(C)** Heatmaps representing all the reconstructions in the state-
labelled calcium signal dataset, for each state, after splitting all traces into 10-
second-long non-overlapping chunks. (n=348 tonic traces; n=580 bursting traces).
(D) Comparing the distribution of trace properties for both states. The top three plots
show (from left to right) the number of peaks, peak amplitude and area under the
curve of the reconstructions in the dataset. The bottom three plots show (from left to
right) the mean, standard deviation and the coefficient of variation of $\Delta F/F$ values. n's
are the number of 10s-long samples. N's are the number of cells used to generate
the samples. p-values were computed using linear mixed-effects models. **(E)**
Principal Components Analysis of trace properties, with data points marked by state
- tonic (red crosses) and bursting (black open circles).
1330
1331

1332

Figure 5 - Design and training of CaMLsort (A) Flowchart showing how the
state- labelled calcium signal dataset was split into training and test datasets
using a 5-fold 60-20-20 split into training (white boxes), test (blue boxes) and
cross-validation (yellow boxes) samples, respectively. The networks that
performed well in the training phase, as assessed by performance on the cross-
validation samples, were then challenged with the unseen test data (blue

boxes) in each fold. An independent dataset with state switches was also used
to further validate the networks obtained. **(B)** The architecture of the
convolutional recurrent neural networks used to solve the classification
problem. The network takes an input time series trace sampled at 30Hz
(purple) and that is a multiple of 10 seconds long in duration (here just 10
seconds long). This trace is first normalized using min-max normalization,
following which it is passed through a 1-dimensional convolutional neural
network (1D CNN), which has 4 kernels with a step size and stride length of 30
each. The resulting 4 traces (grey boxes) are down-sampled, local feature-
extracted versions of the original trace. All 4 outputs are passed to the LSTM
module (central black box) for trend identification. The LSTM is a bidirectional
one, which processes traces in both the forward (blue arrow) and reverse (red
arrow) directions. A single time step is taken from all the CNN outputs (red and
blue dotted boxes) and passed to the respective LSTM cell, which then
processes the input to produce a resultant "hidden state" output (red and blue
filled boxes). This output is passed back into the LSTM cell along with the next
sample from the CNN's output traces, until every time step from it has been
processed (here 10 rounds). Once this is done, the final "hidden state" from
each LSTM cell is retained and concatenated before their weighted average is
taken by a Linear layer. The resulting average is converted to a "posterior
score" using an activation function. These posterior scores represent the
likelihood of the cell being in the bursting state at each time step. The likelihood
of it being in the tonic state can be calculated from this. The final call is taken to
be the state which has a higher posterior score. The text in italics represents
the net effect of each phase of the neural network. The numbers in parentheses
indicate the sampling frequency and/or the duration of the resultant vectors at
various stages of processing by the neural network. Similarly, the numbers in
bold and italics within square brackets indicate the size of the vectors/matrices
at various stages of passing through the network. **(C)** Average classification
accuracy of trained networks at the cross-validation and the test phases. Each
fold is represented in a different color, marked in the legend at the bottom of
panel (D). **(D)** Average F1-scores for each of the trained networks at the cross-
validation and the test phases. **(E)** Area under the ROC (Receiver Operating
Characteristic) curve for each of the five trained CNN-LSTM networks for both
cross-validation (black) and test (gray) data. **(F)** ROC curve for Fold 2 of the
CNN-LSTM for cross-validation (black) and test (gray) data.
1375

1376

Figure 6 - CaMLsort predictions from previously unseen calcium signal
reconstructions (A) Raw CaMLsort class label predictions from previously
unseen calcium signal reconstructions (n=19 recordings). Predictions from
CaMLsort (middle) are compared against the ground truth state (top) at each
time step. Predictions for each recording were made independently, but have

been stitched here for the purposes of representation. Each recording is
separated by a vertical dashed line. The number above each prediction is the
F1-score for that recording. The bottommost plot has posterior scores for each
time step, with those for the "bursting" class in blue, and those for the "tonic"
class in light orange. **(B)** CaMLsort predictions (top) that remain after majority
voting with a window length of 7 steps has been used to retain only confident
classifications (see Results and Methods for details). The posterior score
(bottom) for each class at each time step, taken as the average class posterior
score across all 7-step windows that included said time step. As in (A), the
posterior scores for the "bursting" class are in blue, whereas those for the
"tonic" class are in light orange, and the numbers above each prediction
represent the F1-scores for that recording. **(C)** The distribution of F1-scores
between the ground truth state and various predictions from CaMLsort - either
the raw predictions (left), predictions obtained after majority voting (middle), or
the raw predictions obtained by passing a scrambled version of the calcium
signal as an input.
1398

1399

Figure 7 - CaMLsort predictions from calcium signal reconstructions of simulated spike trains (A) Raw CaMLsort class label predictions from previously unseen calcium signal reconstructions that were generated from simulated spike trains (n=20 recordings). Predictions from CaMLsort (middle) are compared against the ground truth state (top) at each time step. Predictions

for each recording were made independently, but have been stitched here for
the purposes of representation. Each recording is separated by a vertical
dashed line. The number above each prediction is the F1-score for that
recording. The bottommost plot has posterior scores for each time step, with
those for the "bursting" class in blue, and those for the "tonic" class in light
orange. **(B)** CaMLsort predictions (top) that remain after majority voting with a
window length of 7 steps has been used to retain only confident classifications
(see Results and Methods for details). The posterior score (bottom) for each
class at each time step, taken as the average class posterior score across all 7-
step windows that included said time step. As in (A), the posterior scores for the
"bursting" class are in blue, whereas those for the "tonic" class are in light
orange, and the numbers above each prediction represent the F1-scores for
that recording. **(C)** The distribution of F1-scores between the ground truth state
and various predictions from CaMLsort - either the raw predictions (left),
predictions obtained after majority voting (middle), or the raw predictions
obtained by passing a scrambled version of the calcium signal as an input.
1421

Figure 8 - CaMLsort predictions on imaging data from zebrafish Purkinje neurons (A) Raw CaMLsort predictions of cellular state inferred from experimental data acquired using widefield imaging (n=15; also see Chapter 4). Predictions from CaMLsort (middle) are compared against the ground truth state (top) at each time step. Each recording was treated independently, but

predictions across cells have been stitched here for ease of visualisation. Each
recording is separated by a vertical dashed line, and the number above each
prediction is the F1-score for that recording. The bottommost plot has posterior
scores for every time step, with those for the "bursting" class in blue, and those
for the "tonic" class in light orange. **(B)** CaMLsort predictions (top) that remain
after majority voting with a window length of 7 steps has been used to retain
only confident classifications (see Results and Methods for details). The
posterior score (bottom) for each class at each time step, taken as the average
class posterior score across all 7-step windows that included said time step. As
in (A), the posterior scores for the "bursting" class are in blue, whereas those
for the "tonic" class are in light orange. **(C)** The distribution of F1-scores
between the ground truth state and various predictions from CaMLsort - raw
predictions from the imaging data (far left), predictions obtained after majority
voting (second from left), raw predictions obtained from calcium signals
reconstructed from spike times (third from left) and those predictions from
imaging dF/F traces that were scrambled prior to prediction.
1444
1445

1446

Figure 9 - Reconstructing calcium signals for VTA DA neurons using a
convolution-based forward model. (A-B) Spike rasters of two representative

VTA DA neurons. **(C)** The distributions of C.V.s of interspike intervals for cells

classified as either bursting (n=62) or tonic (n=16 cells). The p-value was

calculated using a one-sided Mann-Whitney U-test. **(D)** Electrophysiological

activity and associated calcium signals from two representative VTA DA

neurons. For each cell, spike timings are shown as rasters (black) and the raw

simultaneously-acquired dF/F signals aligned below (green). Two calcium

signal reconstructions were generated, using either a GCaMP6f kernel (gold) or

a GCaMP5G kernel (blue). Numbers alongside each reconstruction indicate the

Pearson correlation coefficient between the ground truth and the corresponding

calcium signal reconstruction. **(E)** Quantification of the reconstruction quality, as

measured using the Pearson correlation coefficient between the ground truth

and reconstructed signals ("Reconstruction"). For each cell (n=78), the

correlation between ground truth and a scrambled version of the reconstructed

signal ("Scrambled"), as well as the correlation between ground truth and a

reconstruction generated after shuffling spike timings ("Events Shuffled") were

measured. Individual data points are plotted alongside each boxplot. p-values

were calculated using a Kruskal-Wallis test yielded a p-value of 4.6130×10^{-49},

and then using the post-hoc Dunn's test for individual comparisons, the latter of

which are marked on the figure.

1468

1469

1470
Figure 10 - CaMLsort predictions of cellular state from calcium signals of
VTA DA neurons (A) A subset of 12 (of 78) raw CaMLsort predictions from
imaging data acquired from DA neurons of the VTA in mice. Each prediction

was independently obtained but has been stitched together to produce a single
time series with cells separated by vertical dashed lines. The ground truth state
(top) is used as reference to assess the quality of CaMLsort's predictions
(middle) using the F1-score, which is printed above the prediction for each cell.
The posterior scores for each class - bursting (blue) and tonic (light orange) -
are shown in the lowest plot. **(B)** The entire distribution of F1-scores (n=78)
between the ground truth state and the binary classification from CaMLsort
obtained from various inputs - (from left to right) experimentally-acquired dF/F
traces, calcium signals reconstructed from true spike trains using the
GCaMP5G kernel, calcium signals reconstructed from shuffled spike trains
using the GCaMP5G kernel, or a scrambled version of the raw dF/F data.
Individual datapoints are shown alongside each boxplot. The horizontal dashed
line corresponds to an F1-score of 0.5.
1487