1

CryoEM reveals unprecedented binding site for Na$_V$1.7 inhibitors enabling rational design of potent hybrid inhibitors

Marc Kschonsak[1*], Christine C. Jao[1*], Christopher P. Arthur[1], Alexis L. Rohou[1], Philippe Bergeron[2], Daniel Ortwine[2], Steven J. McKerrall[2], David H. Hackos[3], Lunbin Deng[3], Jun Chen[4], Peter S. Dragovich[2], Matthew Volgraf[2], Matthew R. Wright[5], Jian Payandeh[1,#], Claudio Ciferri[1,#], John C. Tellis[2,#]

[1]Genentech Inc., Structural Biology; South San Francisco, CA, United States.
[2]Genentech Inc., Discovery Chemistry; South San Francisco, CA, United States.
[3]Genentech Inc., Neuroscience; South San Francisco, CA, United States.
[4]Genentech Inc., Biochemical and Cellular Pharmacology; South San Francisco, CA, United States.
[5]Genentech Inc., Drug Metabolism and Pharmacokinetics; South San Francisco, CA, United States.

[#]Corresponding authors. Email: jpayandeh@exelixis.com (J.P); ciferri.claudio@gene.com (C.C.); tellis.john@gene.com (J.C.T.).

Author Contributions: J.P., C.C., and J.C.T. designed research, M.K., C.J., C.P.A., A.R., D.H., and J.C., performed research, J.C.T, P.B., D.O., S.M., P.S.D., and M.V. analyzed data and contributed to compound design. J.C.T., C.C., J.P., and M.K. wrote the paper.

Abstract

The voltage-gated sodium (Na$_V$) channel Na$_V$1.7 has been identified as a potential novel pain target due to its striking human genetics. However, clinically available drugs (e.g. lidocaine, carbamazepine, etc.) are not selective among the nine Na$_V$ channel subtypes, Na$_V$1.1-Na$_V$1.9, and the two currently known classes of Na$_V$1.7 subtype-selective inhibitors (aryl- and acylsulfonamides) have undesirable characteristics that may limit their development. Moreover, understanding of the structure-activity relationships of the acylsulfonamide class of Na$_V$1.7 inhibitors, exemplified by the clinical development candidate **GDC-0310**, has been based solely on a single co-crystal structure of an arylsulfonamide inhibitor series. To advance inhibitor design targeting the Na$_V$1.7 channel, we established an iterative system to routinely obtain high-resolution ligand-bound Na$_V$1.7 structures using cryogenic electron microscopy (cryo-EM). We report that **GDC-0310** engages the Na$_V$1.7 voltage-sensing domain 4 (VSD4) through an unexpected binding mode orthogonal to the arylsulfonamide class binding pose, which identifies a previously unknown ligand binding site in Na$_V$ channels. This finding enabled the design of a novel hybrid inhibitor series that bridges the aryl and acylsulfonamide binding pockets and allows for the generation of molecules with substantially differentiated structures and properties. Overall, this study highlights the power of cryo-EM methods to pursue challenging drug targets using iterative and high-resolution structure-guided inhibitor design. It also underscores an important role of the membrane bilayer in the discovery of selective Na$_V$ channel modulators.

Introduction

Voltage-gated sodium (Na$_V$) channels initiate and propagate action potentials in excitable tissues and play important roles in health and disease.[1,2] The Na$_V$1.7 channel is expressed predominantly in the peripheral nervous system and genetic studies have identified compelling loss-of-function and gain-of-function phenotypes in human pain syndromes, prompting significant efforts to develop Na$_V$1.7-selective inhibitors as potential novel analgesic drugs.[3-5] Na$_V$ channels contain 24-transmembrane segments linked in four homologous domains (DI-DIV), where four peripheral voltage-sensor domains (VSD1-4) surround a central ion-conducting pore module that houses the ion selectivity filter and key ligand and toxin receptor sites. Traditionally, all clinically available Na$_V$ channel inhibitors lack significant molecular selectivity among the Na$_V$1.1–1.9 subtypes owing to the high sequence conservation found at the ligand binding site within the central cavity of the ion conducting pore module.[5,6]

A breakthrough study in 2013 by McCormack and colleagues reported the discovery of an arylsulfonamide antagonist, **PF-04856264**, that bound to an unprecedented receptor site in VSD4 with demonstrated molecular selectivity for human Na$_V$1.7 over other subtypes.[7] However, the related development candidate **PF-05089771** did not meet clinical endpoints in human subjects with painful diabetic peripheral neuropathy, possibly due to poor target coverage and the intolerable doses required to achieve efficacy.[8-10] Additionally, an alternate acylsulfonamide inhibitor that also targeted the VSD4 receptor

site in Na$_V$1.7, **GDC-0276**, was halted in Phase I clinical trials due to safety concerns and potential off-target effects likely attributed to the high lipophilicity of the compound.[11,12] To date, selective Na$_V$1.7 inhibitors with an improved therapeutic index relative to these clinical-stage compounds have not yet been identified, and this absence underscores the need to optimize such molecules using modern, structure-guided design approaches.

The inherent complexity and dynamic nature of human Na$_V$ channels have historically presented significant barriers to obtaining high-resolution experimental structural information, especially of inhibitor-bound complexes.[13] Using an engineered human VSD4 Na$_V$1.7-Na$_V$Ab bacterial channel chimera and X-ray crystallography, the binding mode of the **GX-936** arylsulfonamide inhibitor (Supplementary Figure 1) revealed that the anionic sulfonamide group engages the fourth arginine gating charge (R4) to trap VSD4 in an activated conformation, which stabilizes a non-conductive, inactivated state of the channel.[14] While determinants of subtype selectivity and structure-activity relationships (SAR) of the arylsulfonamide inhibitor class could be rationalized by the **GX-936** co-crystal structure, additional ligand-bound structures of suitable resolution were not returned by the Na$_V$1.7-Na$_V$Ab chimeric channel crystallography system. This shortcoming led molecular docking studies to presume that the acylsulfonamide inhibitors also bound VSD4 in an analogous manner, despite points of inexplicable SAR.[15] The overall paucity of direct structural information has made the optimization of Na$_V$1.7 inhibitors challenging, with many questions about the determinants of potency, selectivity, and the relationship between the acylsulfonamide and arylsulfonamide inhibitor classes remaining unanswered.

Over the last three decades, protein crystallography and structure-based drug design (SBDD) have become gold standards across the pharmaceutical industry for the identification of ligand binding pockets and the optimization of drug candidates for clinical development. While SBDD has proven successful for many important targets, including G protein-coupled receptors (GPCRs),[16] its applicability to several membrane protein targets, such as Na$_V$ channels, has been limited due to the extreme difficulties with their iterative crystallization and structure determination.[13] Cryogenic electron microscopy (cryo-EM) has recently emerged as a transformative technique to determine the high-resolution structures of diverse protein targets and has proven particularly effective for membrane proteins that are recalcitrant to crystallization.[17] Despite this breakthrough, cryo-EM structure determination of membrane targets in complex with small or large molecule therapeutics frequently remains retrospective and is often enabled after the advancement of key molecules into the clinic.[18,19] Sluggish turn-around times and modest resolutions typically afforded by cryo-EM have limited its application for real-time SBDD efforts. Here, we describe a system for the iterative determination of high-resolution Na$_V$1.7 small molecule co-structures via cryo-EM that has led to the development of a novel class of inhibitors. Critical to this advancement has been establishment of a robust protocol for sample preparation and structure determination and the enablement of the first structure of an acylsulfonamide bound to Na$_V$1.7 VSD4, revealing a previously unknown pocket between the S3 and S4 helices. Consequently, a novel hybrid inhibitor series bridging the aryl and acylsulfonamide pockets was generated. Our work exemplifies the deployment of cryo-EM as a workhorse structural biology tool in an active medicinal

121 chemistry campaign and thus represents an important milestone toward Na$_V$ channel drug
122 discovery.
123
124
125 **Results**
126
Structure of the G-3565-VSD4 Na$_V$1.7-Na$_V$Pas channel complex in lipid nanodiscs
Our approach to establish an iterative, high-resolution system to enable Na$_V$1.7
SBDD was ultimately guided by three observations: our repeated failure to generate well-
diffracting crystals of the Na$_V$1.7-Na$_V$Ab bacterial channel chimera system,[14] our difficulty
of reproducibly expressing a suitable amount of full-length human Na$_V$1.7 channel,[20,21] and
the very limited local resolution observed at VSD4 in all available human Nav1.7 cryo-EM
structures.[22,23] Thus, we sought to exploit an engineered human VSD4-Na$_V$1.7-Na$_V$Pas
cockroach channel chimeric construct that had been previously shown to complex with
small molecule inhibitors and peptide toxins known to target VSD4.[21] The robust protein
yields following expression and purification allowed us to readily pursue reconstitution of
the VSD4-Na$_V$1.7-Na$_V$Pas channel into lipid nanodisc with high recovery (Supplementary
Figure 2). Small molecule inhibitors were added prior to sample vitrification, followed by
cryo-EM data collection and processing procedures, which allowed us to routinely obtain
3D-reconstructions in the 2.2-3.0 Å resolution range (Figure 1, Supplementary Figure 3).
141
The VSD4-Na$_V$1.7-Na$_V$Pas chimera displays the expected domain-swapped
arrangement with numerous densities assigned as phospholipids bound to the channel,
confirming the maintenance of a native membrane-like environment (Figure 1). As seen in
previous Na$_V$Pas structures,[21,24,25] the ion-conducting pore module of the VSD4-Na$_V$1.7-
Na$_V$Pas chimeric channel is closed, consistent with a nonconductive or inactivated state.
The quality and resolution of our structure allowed us to assign 112 water molecules for
the highest resolution structure (Figure 1, Video 1). The visualization of many well-
resolved water molecules bound within the VSDs and coordinated within the ion selectivity
filter provides new insights into the interactions that might occur during gating charge
transfer and ion conduction, respectively.
152
Although we were able to determine the crystal structure of VSD4-Na$_V$Ab bound
to the arylsulfonamide inhibitor **GX-936**,[14] no well-diffracting crystals were obtained for
any other arylsulfonamide compounds despite years of continued effort. Arylsulfonamide
GNE-3565 is representative of an advanced series of arylsulfonamide class Na$_V$1.7
inhibitors that demonstrates channel blockage at subnanomolar concentrations with mixed
subtype selectivity (Figure 2A). We complexed **GNE-3565** with the VSD4 Na$_V$1.7-Na$_V$Pas
channel-nanodisc system and employed cryo-EM to assess its overall binding pose (Figure
2B, C). The 2.9 Å resolution cryo-EM map revealed that the S1-S2 and S3-S4 helices from
VSD4 form a clamshell-like structure that closes over **GNE-3565** (Figure 2D, E,
Supplementary Data Figure 4A-D, Supplementary Data Table 1), similar to the binding
pose reported for **GX-936**.[14] Specifically, the ionized arylsulfonamide group of **GNE-
3565** (measured pKa = 5.8) bisects VSD4 to salt-bridge directly to R4 on the S4 helix,
while the central phenyl ring extends perpendicular between the S2 and S3 helices to
directly contact established subtype-selectivity determinants Tyr1537 and Trp1538 on the

S2 helix (Figure 2A-D). As for **GX-936**, the **GNE-3565** arylsulfonamide VSD4 receptor
binding site can be divided into three regions: an anion-binding pocket, a selectivity pocket,
and a lipid-exposed pocket (Figure 2D). The root-mean-square deviation (RMSD) between
the **GNE-3565**-VSD4 and the **GX-936**-VSD4 structures[14] is 0.667 (714 to 714 atoms),
(Supplementary Figure 5A), supporting long-held assumptions by medicinal chemistry
teams that all structurally-related arylsulfonamides should be expected to complex to
VSD4 through similar determinants.[8,26-31]
174
Structure of the acylsulfonamide GDC-0310 in complex with the VSD4 Na$_V$1.7-
Na$_V$Pas channel
To investigate the binding pose of a representative Na$_V$1.7 selective
acylsulfonamide, **GDC-0310** (Figure 3A) was complexed with the VSD4 Na$_V$1.7-Na$_V$Pas
channel in nanodiscs and a cryo-EM structure was determined to 2.5 Å resolution (Figure
3B-D, Supplementary Data Figure 3G-J, Supplementary Data Table 1). Remarkably, the
binding mode of **GDC-0310** revealed an unexpected and previously unknown pocket that
formed between the S3 and S4 helices (Figure 3B-E). While the aryl and acyl moieties of
GDC-0310 and **GNE-3565** bind in the same region, the remainder of the acyl receptor site
is orthogonal to the corresponding binding domain reported for the aryl class (Figure 3B-
D). In detail, the anionic acylsulfonamide of **GDC-0310** participates in salt-bridge bonding
interactions with R4 and R3 through the carbonyl and sulfonamide oxygen atoms,
respectively (Figure 3F). Moreover, the aryl ring of the **GDC-0310** splits the S3 and S4
helices to occupy a lipophilic pocket, displacing the S3 helix laterally by ~3 Å relative to
the **GNE-3565**-VSD4 structure (Figure 3G). Our cryo-EM inhibitor bound structures serve
to highlight a dynamic environment in VSD4 where the S1-S4 helixes can be differentially
separated by distinct small molecule binding poses (Figure 2D, Figure 3D).
192
It is notable that the cyclopropyl substituent of **GDC-0310** makes effective van der
Waals interactions with I1574 and I1588 from S3 and I1601 and I1604 on S4 (Figure 3G),
while the alpha-methylbenzylamine "tail" region sits almost entirely within the lipid
bilayer (Figure 3A, C), where density for this portion of the inhibitor is only poorly
resolved (Figure 3E). Considering the relative depth of the **GDC-0310** binding site in
relation to the membrane-water interface (~8 Å), our structure suggest a membrane access
pathway for acylsulfonamides to the VSD4 receptor site, which has important implications
for understanding the pharmacology, SAR, and potential development liabilities of the
series.[32] Notably, this structure also reveals that the Tyr1537 side-chain on S2 exists in a
down rotamer and does not contact **GDC-0310** directly, which provides the first direct
structural rationale for why the selectivity profiles between the acyl and arylsulfonamide
classes differ substantially (Figure 3D, *vide infra*).[33]
205
Structure-based rational design of hybrid Na$_V$1.7 inhibitors
Upon inspection of the superimposed **GNE-3565**-aryl and **GDC-0310**-acyl
structures, we could immediately envision a novel class of hybrid inhibitors that would
simultaneously occupy both the aryl- and acylsulfonamide binding pockets (Figure 4A, B).
Such an enlarged hybrid binding pocket would offer unique opportunities to gain potency
directly from ligand-protein interactions, which might enable removal of the hydrophobic
lipid-facing tail groups seen in all previous generations of VSD4-targeting Na$_V$ channel

inhibitors. These groups have typically been critical for maintaining potency, but use of
the plasma membrane to drive membrane-bound target occupancy is largely nonspecific,
and potentially introduces off-target liabilities that can contribute to toxicity. However, it
was unclear if this hybrid conformation of the VSD4 domain could accommodate the
described inhibitors and/or whether it was accessible during any stage of $Na_V1.7$ channel
activation. We therefore set out to demonstrate proof-of-concept that appropriately
designed small molecules could induce this hypothesized receptor state while inhibiting
channel function with meaningful potency.
221
GDC-0310 was a relatively unattractive starting point for hybridization because of
its high molecular weight (543 g/mol) and lipophilicity (cLogP = 5.2). Accordingly,
analysis of historical data identified progenitor compound **1**, which offers reasonable
potency against $Na_V1.7$ (IC_{50} = 25 nM, n = 8; Fig 4C) and reduced molecular weight (350
g/mol) and lipophilicity (cLogP = 2.7). Seeking to leverage existing knowledge of the
arylsulfonamide binding pocket, molecule **2** was synthesized in an attempt to directly graft
on a significant portion of the **GNE-3565** scaffold. Gratifyingly, this molecule retained
substantial potency, affording channel blockade at an IC_{50} of 47 nM (n = 8).
230
A cryo-EM structure of **2** bound to the VSD4 $Na_V1.7$-NaVPas channel confirmed
that the molecule adopts the desired hybrid binding mode (Figure 4D). Here, the anionic
group of the novel hybrid compound binds in the same position as in the aryl and acyl
poses, interacting closely with R4, as seen for the aryl-like binding (Figure 4E). The aryl-
and acylsulfonamide-derived structural motifs of **2** occupy their expected positions in the
pockets formed between S2/S3 and S3/S4, respectively. Notably, in this hybrid
configuration, Y1537 adopts a similar arrangement as observed for the arylsulfonamides
(Figure 4F).
239
Encouraged by this result, we hoped to draw on previously established knowledge
of acylsulfonamide SAR to improve the potency of this compound. Established SAR within
the acylsulfonamide series suggested that replacement of the chloro substituent on the
benzamide fragment of the molecule with a cyclopropane would typically improve
potency. This SAR proved translatable to the hybrid class molecules, resulting in inhibitor
3 (IC_{50} = 16 nM, n = 12; Figure 4C).
246
Further analysis of the cryo-EM structure of **2** revealed that the aryl ring in the
S2/S3 binding pocket was shifted ~1 Å away from the VSD4 core when compared to
arylsulfonamides such as **GNE-3565** (data not shown). The aryl ring in the S3/S4 pocket
exhibited better overlap with the analogous substituents in the **GDC-0310** structure. We
hypothesized that the acylsulfonamide moiety of **2**, which places the two aryl substituents
4.7 Å apart, might not be fully optimized for use in hybrid inhibitors. In comparison, an
arylsulfonamide moiety would position these two groups at a distance of ~3.3 Å. On this
basis we designed *N*-benzothiazolyl sulfonamide **4**, which demonstrated potent $Na_V1.7$
inhibition (IC_{50} = 9 nM, n = 8, Figure 4C).
256
Our last hypothesis was again validated with a cryo-EM co-structure of **4** bound to
the VSD4 $Na_V1.7$-NaVPas channel (Figure 4G). Here, we observed the typical face-on pi-

stacking orientation of the S2/S3 substituent with Y1537 while maintaining efficient space
filling in the S3/S4 pocket. Moreover, the efficient occupation of the non-membrane
exposed regions of the binding pocket permitted the removal of the lipophilic tail of the
molecule in **5**, where a CF_3 group replaces the cyclopentylmethyl substituent. Although
this molecule shows reduced potency (IC_{50} = 74 nM, n = 8), lack of substantial small
molecule contact with the lipid bilayer is an uncommon feature in all other known $Na_V1.7$
VSD4 domain inhibitors. These molecules serve as a key proof of concept, highlighting
the power of cryoEM to yield new structural insights that can inspire the design of novel,
structurally differentiated small molecule inhibitors.
268
269
270   **Discussion**
271
Iterative structure-based design is among the most powerful techniques to facilitate
the development of small molecule drug candidates. Although a variety of technologies are
capable of providing information about small molecule ligand-protein interactions, SBDD
has traditionally been synonymous with the use of X-ray crystallography, with other
techniques failing to rival its high resolution and fast cycle times. Here, we have
demonstrated that a robust cryo-EM structure pipeline is sufficient to quickly provide co-
structures at resolutions suitable for use in an active medicinal chemistry optimization
campaign. Moreover, we have highlighted the power of structural methods, and cryo-EM
in particular, to provide new and important insights into inhibitor design through the
discovery of a previously unknown binding mode for acylsulfonamide $Na_V1.7$ inhibitors.
We further note that our use of an engineered VSD4-$Na_V1.7$-Na_VPas chimeric channel
construct was essential to produce high yields of recombinant protein capable of routinely
returning multiple cryo-EM samples per preparation. This system repeatedly delivered
high-resolution depictions of the VSD4 inhibitor binding site and thus strongly contrasts
with all prior accounts of cryo-EM structures of full-length human $Na_V1.7$ channels.[22,23]
Importantly, high quality cryo-EM maps at VSD4 in complex with ICA121431 have
recently been reported from full-length human $Na_V1.3$ channel protein,[37] indicating that
SBDD efforts targeting a particular Na_V channel subtype would benefit from exploring
both engineered and native channel sample options at early project stages.
291
The co-structure of **GDC-0310** was critical to contextualize historical differences
between aryl- and acylsulfonamide $Na_V1.7$ inhibitors, which include divergent SAR in the
linker phenyl ring and tail regions, disparate Na_V family selectivity patterns, and
differences in on-rates and off-rates.[12,15,33-36] Previous efforts to model acylsulfonamide
binding have attempted to dock molecules into the **GX-936** co-structure and have proposed
subtle changes in molecular register and/or pose to explain the non-translatable SAR.[15,38]
Here, we demonstrate that the molecular features of aryl- and acylsulfonamide inhibitors
diverge because they occupy distinct, but overlapping, pockets within the VSD4 domain.
300
The selectivity determinants of arylsulfonamide $Na_V1.7$ inhibitors have been well
studied. Mutation of key non-conserved residues proximal to the binding pocket observed
in the **GX-936** co-structure have been shown to alter $Na_V1.7$ potency consistent with
selectivity patterns observed for other Na_V family members.[14] Typically, arylsulfonamides

have shown high levels of selectivity against Na$_V$1.5, Na$_V$1.1, and Na$_V$1.4, but lower
selectivity against Na$_V$1.2 and Na$_V$1.6. In comparison, several acylsulfonamides have been
characterized with very high selectivity over Na$_V$1.2 and Na$_V$1.6.[33] Interestingly, residues
forming the contours of the acylsulfonamide binding pocket show high sequence homology
across Na$_V$ isoforms, suggesting that selectivity for this class is likely an allosteric
phenomenon. Nonetheless, the disparate binding pockets for these two inhibitor classes
provide a clear rationale for their historically divergent patterns of selectivity.
312
Our structure of **GDC-0310** provides a structural rationale for the relative slow
dissociation kinetics of acylsulfonamide inhibitors compared to arylsulfonamides.[33]
Specifically, the acylsulfonamide pocket is buried more deeply into the plasma membrane
than the arylsulfonamide pocket, and may be available only through a membrane-access
pathway that first involves partitioning of the molecule into the intramembrane space. In
comparison, the arylsulfonamide pocket is open to the outer membrane interface, offering
access from the solvent compartment. Slow dissociation kinetics are known to be
associated with membrane-access pathways for small molecules binding to transmembrane
proteins.[32,39] For example, the long duration of action of salmeterol (a long-acting β$_2$
adrenergic receptor agonist) is proposed to be a result of its membrane partitioning
properties. In comparison, salbutamol, which binds to the same target and site but through
a solvent-access pathway, shows much faster dissociation from the receptor.[40-46]
325
In addition to offering valuable information for retrospective analysis of historical
Na$_V$1.7 VSD4 domain inhibitors, the structure of **GDC-0310** inspired the development of
a class of structurally differentiated molecules that bind to a novel conformation of the
receptor. These hybrid molecules bridge both the S2/S3 and S3/S4 helical gaps, with the
anionic moiety situated centrally, interacting with the arginine network at the core of the
domain. We have demonstrated proof-of-concept that potent hybrid inhibitors can be
designed with fewer membrane-associated elements than prior generation Na$_V$1.7
inhibitors, and anticipate that this will offer opportunities to develop molecules with
improved lipophilic ligand efficiency and/or differentiated pharmacokinetic profiles.
Hybrid inhibitors can also access selectivity determinants from both the S2/S3 and S3/S4
pockets simultaneously, offering a tantalizing opportunity to develop exquisite selectivity
over other Na$_V$ isoforms.
338
Beyond the scope of Na$_V$ channel inhibitors, our experiences may also offer lessons
translatable to small molecule modulators of other transmembrane proteins. Specifically,
it is noteworthy to observe that both aryl- and acylsulfonamide inhibitors have access to
cryptic pockets that are not observable in the co-structures of the parent proteins. Designing
small molecules to "push" on a structural protein element is a common strategy in
medicinal chemistry, but it is often challenging to predict when such efforts will afford
new, well-defined binding pockets. As such, these designs are often met with failure and
quickly abandoned. However, some scenarios exist in which a cryptic pocket is more likely
to be found. In these cases, medicinal chemists are more often emboldened to continue
experimenting with designs that push into new areas of a protein binding pocket, even
when met with initial failure. For example, type II kinase inhibitors can be rationally
designed starting from a type I inhibitor structure by appending groups that push through

the DFG into the cryptic back pocket.[47] This pocket is not visible from the type I inhibitor
structure, but knowledge of the protein dynamics associated with the DFG-in/DFG-out
transition provides a rationale for exploring these designs. We speculate that
transmembrane intrahelical domains, particularly those in ion channels, transporters, and
GPCRs, may present similarly privileged protein structural motifs that are rich with
opportunities for the discovery of druggable cryptic pockets.
357
In summary, we have developed a robust pipeline for high-resolution structural
determination of small-molecule bound Nav1.7 channels suitable for use in iterative
structure-based drug design campaigns. We also report the first co-structure of an
acylsulfonamide inhibitor bound to the VSD4 domain of $Na_V1.7$, revealing an unexpected
and unique binding mode. This finding provides clear rationale for previously
unexplainable divergence in the in vitro pharmacological behavior between the two
inhibitor classes, and inspired the development of hybrid inhibitors that merge structural
features from both inhibitor classes, engaging the receptor in a novel conformation. Our
findings highlight the power of cryoEM as an enabling drug discovery technology and offer
interesting learnings regarding ion channel structural dynamics that are potentially
applicable to other related targets and target classes.
369
370 **Materials and Methods**
371
Generation of VSD4-Na$_V$1.7-Na$_V$Pas channel constructs
VSD4-Na$_V$1.7-Na$_V$Pas chimeric constructs were used as described previously.[21] In brief,
optimized coding DNA for Nav1.7 VSD4-Na$_V$PaS chimeras with N-terminal tandem
StrepII and FLAG tag was cloned into a pRK vector with CMV promoter. HEK293 cells
in suspension were cultured in SMM 293T-I medium under 5% CO_2 at 37°C and
transfected using PEI when the cell density reached 4×10^6 cells per ml. Transfected cells
were harvested 48 hours after transfection. The Dc1a toxin coding DNA from *Diguetia*
canities was cloned into a modified pAcGP67A vector downstream of the polyhedron
promoter and an N-terminal 6x HIS tag. Recombinant baculovirus was generated using the
Baculogold system (BD Biosciences) and *Trichoplusia ni* cells were infected for protein
production. The supernatant was harvested 48 h post-infection.
383
Protein expression and purification:
One hundred fifty grams of cell pellet was resuspended in 500mL of 25mM Hepes pH 7.5,
200mM NaCl, 1ug/mL benzonase, 1mM PMSF, and Roche protease inhibitor tablets. Cells
were lysed by dounce homogenization. Proteins were solubilized by addition of 2% GDN
(Avanti) with 0.3% cholesteryl hemisuccinate (Avanti) for 2 hours at 4°C. Cell debris was
separated by ultracentrifugation at 40,000 rpm at 4°C. Affinity purification using anti-Flag
resin was performed by batch binding for 1 hour at 4°C. The resin was washed with 5CV
Purification Buffer (25mM Hepes pH 7.5, 200mM NaCl, 0.01% GDN). Another 5CV
wash was performed using Purification Buffer supplemented with 5mM ATP and 10mM
$MgCl_2$. Proteins were eluted in 6 CV Purification Buffer with 300ug/mL Flag peptide.
Proteins were subjected to another round of affinity purification using StrepTactin resin
(IBA). Proteins were eluted, and concentrated to ~ 5 mg/mL. For the compound **2**
containing sample, instead of nanodisc reconstitution VSD4-Na$_V$1.7-Na$_V$Pas was

incubated overnight with Dc1a toxin at a 2:1 molar ration of toxin:VSD4-Na$_V$1.7-Na$_V$Pas.
The eluted sample was concentrated to 100uL, and separated on Superose 6 3.2/300.
399
For nanodisc reconstitution used for the samples containing **GNE-3565, GDC-0310** and
compound **4**, a 200-molar excess of lipid mix (3POPC:1POPE:1POPG resuspended in
50mM Hepes pH7.5, 100mM NaCl, 5mM MgCl$_2$, 1% CHAPS) was added to detergent-
solubilized protein and incubated on ice for 30 minutes. A four-molar excess of scaffold
protein (MSP1E3D1, Sigma) was added to the protein-lipid mix, and incubated on ice for
another 30 minutes. To remove detergent, BioBeads (Bio-Rad) were added to 0.25 mg/mL,
and incubated overnight at 4°C. To remove the empty nanodiscs, the sample was subjected
to a round of affinity purification using Strep-Tactin (IBA). The eluted sample was
concentrated to 100uL, and separated on Superose 6 3.2/300. Peak fractions were
combined, and split into four samples. Fifty uM of the small molecule of interest was
added to each sample, and incubated at 22°C for 10 minutes. The nanodisc reconstituted
samples were crosslinked with 0.05% glutaraldehyde (Electron Microscopy Sciences) then
quenched with 1M Tris pH 7.0. Samples at 2 mg/mL were used for grid freezing.
413
414
CryoEM sample preparation and data acquisition
CryoEM grids for the small molecule VSD4-Na$_V$1.7-Na$_V$Pas chimeric complexes shown
in this study were prepared as follows: For the **GDC-0310** and compound **4** VSD4-Na$_V$1.7-
Na$_V$Pas complexes, holey carbon grids (Ultrafoil 25 nM Au R 0.6/1 300 mesh; Quantifoil)
were incubated with a thiol reactive, self-assembling reaction mixture of 4mM
monothiolalkane(C11)PEG6-OH (11-mercaptoundecyl) hexaethyleneglycol (SPT-
0011P6, SensoPath Technologies, Inc., Bozeman, MT). Grids were incubated with this
self-assembled monolayer (SAM) solution for 24 hours and afterwards rinsed with EtOH.
3 μl of the sample was applied to the grid and blotted with Vitrobot Mark IV
(Thermofisher) using 3.5 s blotting time with 100% humidity and plunge-frozen in liquid
ethane cooled by liquid nitrogen. The **GNE-3565** VSD4-Na$_V$1.7-Na$_V$Pas complex was,
similarly as described above, applied to a holey carbon grid (Ultrafoil 25 nM Au R 1.2/1,3
300 mesh; Quantifoil) pretreated with SAM solution. The grid was blotted single-sided
with a Leica EM GP (Leica) using 3 s blotting time with 100% humidity and plunge-frozen
in liquid ethane cooled by liquid nitrogen. For compound **2** VSD4-Na$_V$1.7-Na$_V$Pas-DC1a
complex, holey carbon grids (Ultrafoil 25 nM Au R 2/2 200 mesh; Quantifoil) were glow-
discharged for 10 s using the Solarus plasma cleaner (Gatan). 3 μl of the sample was applied
to the grid and blotted with Vitrobot Mark IV (Thermofisher) using 2.5 s blotting time with
100% humidity and plunge-frozen in liquid ethane cooled by liquid nitrogen.
434
Movie stacks for compound **2** VSD4-Na$_V$1.7-Na$_V$Pas-DC1a complex were collected using
SerialEM[48] on a Titan Krios operated at 300 keV with bioquantum energy filter equipped
with a K2 Summit direct electron detector camera (Gatan). Images were recorded at
165,000 × magnification corresponding to 0.824 Å per pixel, using a 20 eV energy slit.
Each image stack contains 50 frames recorded every 0.2 s for an accumulated dose of ~50
e Å−2 and a total exposure time of 10 s. Images were recorded with a set defocus range of
0.5 to 1.5 μm.
442

Movie stacks for **GNE-3565, GDC-0310** and compound **4** VSD4-Na_V1.7-Na_VPas
complexes were collected using SerialEM on a Titan Krios G3i (ThermoFisher Scientific,
Waltham, MA) operated at 300 keV with bioquantum energy filter equipped with a K3
Summit direct electron detector camera (Gatan Inc, Pleasanton, CA). Images were recorded
in EFTEM mode at 105,000 × magnification corresponding to 0.838 Å per pixel, using a
20 eV energy slit. Each image stack contains 60 frames recorded every 0.05 s for an
accumulated dose of ~60 e Å−2 and a total exposure time of 3 s. Images were recorded
with a set defocus range of 0.5 to 1.5 μm.
451
452 **CryoEM data processing**
453 CryoEM data were processed using a combination of the RELION[49] and cisTEM[50]
454 software packages, similarly as described previously[51] and as illustrated in Suppl. Figure 2
455 and 3.
456
Movies were corrected for frame motion using the MotionCor2[52] implementation in
RELION and contrast-transfer function parameters were fit using the 30-4.5 Å band of the
spectrum with CTFFIND-4.[53] CTF fitted images were filtered on the basis of the detected
fit resolution better than 6-10 Å. Particles were picked by template-matching with
gautomatch   (https://www2.mrc-lmb.cam.ac.uk/research/locally-developed-
software/zhang-software/#gauto) using a 30 Å low-pass filtered apo VSD4-Na_V1.7-
Na_VPas reference structure. Particles were sorted during RELION 2D classification and
selected particles were imported into cisTEM for 3D refinements. 3D reconstructions were
obtained after auto-refine and manual refinements with a mask around the channel
(excluding the detergent micelle) and by applying low-pass filter outside the mask (filter
resolution 20 Å) and a score threshold of 0.10-0.30, so that only the best-scoring 10-30%
of particle images would be included in the 3D reconstruction at each cycle. The weight
outside of the mask was set to 0.8. No data beyond 3.0 Å for compound **4**, 3.4 Å for **GDC-**
0310, 3.7Å **GNE-3565** and 4.0 Å for compound **2** were used in the refinements. Phenix
ResolveCryoEM[54] density modification was applied to each of the reconstructions to
obtained the final map used for model building. Local resolution was determined in
cisTEM using a reimplementation of the blocres algorithm.
474
Model building and structure analysis
The previously determined VSD4-Na_V1.7-Na_VPas model (PDB: 6NT3)[21] was fit as a rigid
body into the cryoEM map. After manual adjustments, multiple rounds of real space
refinement using the phenix.real_space_refinement tool[55] was used to correct structural
differences between the initial model and the map. The small molecule coordinates were
generated with eLBOW,[56] placed and manually adjusted in Coot,[57] energy minimized with
MOE (Chemical Computing Group ULC,) and refined with real space refinements in
Phenix[55]. The model was validated using phenix.validation_cryoem[58] with built-in
MolProbity scoring.[59] Figures were made using PyMOL (The PyMOL Molecular Graphics
System, v.2.07 Schrödinger, LLC), UCSF Chimera,[60] and UCSF ChimeraX.[61]
485
486 **Chemical synthesis**

487 The synthesis of **GNE-3565**, **GDC-0310**, and compound **1** have been previously
488 reported.[12,15,62] The synthesis and characterization of compounds **2-5** are described in the
489 supporting information.
490
Na$_V$1.7 Syncropatch electrophysiology potency
cDNAs for Na$_V$1.7 (NM_002977) were stably expressed in Chinese Hamster Ovary (CHO)
cells. Sodium currents were measured in the whole-cell configuration using Syncropatch
384PE (NanIon Technologies, Germany). 1NPC®-384 chips with custom medium
resistance and single hole mode were used. Internal solution consisted of (in mM): 110
CsCl, 10 CsCl, 20 EGTA, and 10 Hepes (pH adjusted to 7.2); and external solution contains
(in mM): 60 NMDG, 80 NaCl, 4 KCl, 1 MgCl$_2$, 2 CaCl$_2$, 2 D-Glucose monohydrate, 10
Hepes (pH adjusted to 7.4 with NaOH).
499
After system flushing, testing compounds were dissolved in external solution containing
0.1% Pluronic F-127. 10µl cells were added to the chip from a cell hotel, and a negative
pressure of −50 mBar was applied to form a seal. Following treatment with seal enhancer
solution and wash-off with external solution, negative pressure of −250 mbar was applied
for 1 second to achieve the whole-cell configuration, followed by three washing steps in
external solution. 20µl of compounds were added to 40µl in each well (1:3 dilution of
compounds), and after mixing, 20µl was removed so the volume was retained at 40 ul.
After approximately 13 minutes recording, 20µl/well of 2 uM TTX was added to achieve
full block.
509
For voltage protocol, a holding potential of -50 mV was applied during the whole
experiment. A depolarizing step was applied to -10mV for 10ms, followed by a
hyperpolarization step to -150 mV for 20 ms to allow channel recovery from inactivation.
A second depolarizing step was applied from -150mV to -10mV for 10 ms, where currents
were measured to derive blocking effects of compounds. Inhibition was determined based
on 7.5 min of compound incubation.
516
517 **Figure Legends:**
518
Figure 1: (A) Schematic of the VSD4-Na$_V$1.7-Na$_V$Pas channel. The portions humanized
to the Na$_V$1.7 sequence are shown in green. N-terminal domain (NTD) and CTD are
indicated. **(B)** Side view of the single-particle cryo-EM reconstruction of VSD4- Na$_V$1.7-
Na$_V$Pas channel **(C-D)** Cartoon representations of the Top and Side views of VSD4-
Na$_V$1.7-Na$_V$Pas channel. Individual VSD domains are indicated. VSD4 is highlighted in
green. **(E)** Localization of water molecules (in red) in the VSD4-Na$_V$1.7-Na$_V$Pas channel
structure. VSD4 is highlighted in green.
526
Figure 2: (A) Chemical structures of arylsulfonamide **GNE-3565**. **(B,C)** Top and Side
views of VSD4-Na$_V$1.7-Na$_V$Pas channel bound to **GNE-3565**. VSD4 is highlighted in
green, **GNE-3565** in magenta. **(D)** Extracellular view of VSD4-Na$_V$1.7-Na$_V$Pas
arylsulfonamide receptor site is shown with select side chains rendered as sticks. **(E)** The
cryoEM map surrounding the ligand **GNE-3565** is shown in mesh representation. **(F)** View
towards the membrane highlighting key interactions with the **GNE-3565** warhead. **(G)**

View from the membrane highlighting key interactions with the **GNE-3565** central phenyl ring.

Figure 3: (A) Chemical structures of acylsulfonamide **GDC-0310**. **(B)** Top and Side views of VSD4-Na$_V$1.7-Na$_V$Pas channel bound to **GDC-0310**. VSD4 is highlighted in green, **GDC-0310** in magenta. **(D)** Extracellular view of VSD4-Na$_V$1.7-Na$_V$Pas acylsulfonamide receptor site is shown with select side chains rendered as sticks. **(E)** The cryoEM map surrounding the ligand **GDC-0310** is shown in mesh representation. **(F)** View towards the membrane highlighting key interactions with the **GDC-0310** warhead. **(G)** View from the membrane highlighting van der Waals interactions with the **GDC-0310** cyclopropyl substituent. The alpha-methylbenzylamine tail region sits almost entirely within the lipid bilayer.

Figure 4: (A) Illustration of hybrid molecule design approach. **(B)** Arylsulfonamide, acylsulfonamide and hybrid molecule poses **(C)** Hybridization strategy and molecule optimization. Highlighted are molecule **2** and molecule **4 (D)** Extracellular view of VSD4-Na$_V$1.7-Na$_V$Pas bound to the hybrid molecule **2**. **(E)** View towards the membrane highlighting key interactions with the anionic group. **(F)** View from the membrane highlighting the lack of a stacking interaction between Y1537 and the phenyl ring. **(G)** Extracellular view of VSD4-Na$_V$1.7-Na$_V$Pas bound to the hybrid molecule **4**. **(H)** View towards the membrane highlighting key interactions with the anionic group. **(I)** View from the membrane highlighting the π-stacking interaction between Y1537 and the phenyl ring.

Acknowledgments

We are grateful to BMR and RMG for construct generation and protein scale-up. We are thankful to Cameron Noland for help with the biochemistry of the VSD4-Na$_V$1.7-Na$_V$Pas channel system. We thank Tommy Lai and Wenfeng Liu for chemistry support.

References

1. W. A. Catterall, A. L. Goldin, S. G. Waxman, International Union of Pharmacology. XLVII. Nomenclature and structure-function relationships of voltage-gated sodium channels. *Pharmacol. Rev.* **57**, 397 (2005).
2. C. A. Ahern, J. Payandeh, F. Bosmans, B. Chanda, The hitchhiker's guide to the voltage-gated sodium channel galaxy. *J. Gen. Physiol.* **147**, 1 (2016).
3. S. D. Dib-Hajj, Y. Yang, J. A. Black, S. G. Waxman, The Na(V)1.7 sodium channel: from molecule to man. *Nat. Rev. Neurosci.* **14**, 49 (2013).
4. J. Payandeh, D. H. Hackos, Selective ligands and drug discovery targeting the voltage-gated sodium channel Na$_V$1.7. *Handb. Exp. Pharmacol.* **246**, 271 (2018).
5. S. J. McKerrall, D. P. Sutherlin, Na$_V$1.7 inhibitors for the treatment of chronic pain. *Bioorg. Med. Chem. Lett.* **28**, 3141 (2018).
6. M. de Lera Ruiz, R. L. Kraus, Voltage-gated sodium channels: structure, function, pharmacology, and clinical indications. *J. Med. Chem.* **58***, 7093 (2015).

7. K. McCormack, *et al.*, Voltage sensor interaction site for selective small molecule
inhibitors of voltage-gated sodium channels. *Proc. Natl. Acad. Sci. USA* **110**,
E2724 (2013).
8. N. A. Swain, *et al.,* Discovery of clinical candidate 4-[2-(5-amino-1H-pyrazol-4-
yl)-4-chlorophenoxy]-5-chloro-2-fluoro-N-1,3-thiazol-4-ylbenzenesulfonamide
(PF-05089771): design and optimization of diaryl ether aryl sulfonamides as
selective inhibitors of Na$_V$1.7. *J. Med. Chem.* **60**, 7029 (2017).
9. A. McDonnell, *et al.,* Efficacy of the Na$_V$1.7 blocker PF-05089771 in a randomised,
placebo-controlled, double-blind clinical study in subjects with painful diabetic
peripheral neuropathy. *Pain* **159**, 1465 (2018).
10. P. Siebenga, *et al.*, Lack of detection of the analgesic properties of PF-05089771, a
selective Na$_V$1.7 inhibitor, using a battery of pain models in healthy subjects. *Clin.*
Transl. Sci. **13**, 318 (2020).
11. M. E. Rothenberg, *et al.*, Safety, tolerability, and pharmacokinetics of GDC-0276,
a novel Na$_V$1.7 inhibitor, in a first-in-human, single- and multiple-dose study in
healthy volunteers. *Clin. Drug Investig.* **39**, 873 (2019).
12. B. S. Safina, *et al.*, Discovery of acyl-sulfonamide Na$_V$1.7 inhibitors GDC-0276
and GDC-0310. *J. Med. Chem.* **64**, 2953 (2021).
13. S. Noreng, T. Li, J. Payandeh, Structural pharmacology of voltage-gated sodium
channels. *J. Mol. Bio.* **433**, 166967 (2021).
14. S. Ahuja, *et al.*, Structural basis of Nav1.7 inhibition by an isoform-selective small-
molecule antagonist. *Science* **350**, aac5464 (2015).
15. S. Sun, *et al.*, Identification of selective acyl sulfonamide-cycloalkylether inhibitors
of the voltage-gated sodium channel (Na$_V$)1.7 with potent analgesic activity. *J.*
Med. Chem. **62**, 908 (2019).
16. M. Congreve, J. M. Dias, F. H. Marshall, Structure-based drug design for G protein-
coupled receptors. *Prog. Med. Chem.* **53**, 1 (2014).
17. Y. Cheng, Membrane protein structural biology in the era of single particle cryo-
EM. *Curr. Opin. Struct. Biol.* **52**, 58 (2018).
18. A. Balestrini, *et al.*, A TRPA1 inhibitor suppresses neurogenic inflammation and
airway contraction for asthma treatment. *J. Exp. Med.* **218**, e20201637 (2021).
19. L. Rougé, *et al.*, Structure of CD20 in complex with the therapeutic monoclonal
antibody rituximab. *Science* **367**, 1224 (2020).
20. H. Xu, *et al.*, Structural basis of Na$_V$1.7 inhibition by a gating-modifier spider toxin.
Cell **176**, 702 (2019).
21. T. Clairfeuille, et al., Structural basis of α-scorpion toxin action on Na$_V$ channels.
Science **363**, eaav8537 (2019).
22. H. Shen, D. Liu, K. Wu, J. Lei, N. Yan, Structures of human Na$_V$ 1.7 channel in
complex with auxiliary subunits and animal toxins. *Science* **363**, 1303 (2019).
23. G. Huang, *et al.*, High-resolution structures of human Na$_V$1.7 reveal gating
modulation through α-π helical transition of S6$_{IV}$. *Cell Rep.* **39**, 110735 (2022).
24. H. Shen, *et al.*, Structure of a eukaryotic voltage-gated sodium channel at near-
atomic resolution. *Science* **355**, eaal4326 (2017).
25. H. Shen, *et al.*, Structural basis for the modulation of voltage-gated sodium
channels by animal toxins. *Science* **362**, eaau2596 (2018).

26. M. Weiss, *et al.*, Sulfonamides as selective Na$_V$1.7 inhibitors: optimizing potency and pharmacokinetics while mitigating metabolic liabilities, *J. Med. Chem.* **60**, 5969 (2017).

27. R. F. Graceffa, *et al.*, Sulfonamides as selective Na$_V$1.7 inhibitors: optimizing potency, pharmacokinetics, and metabolic properties to obtain atropisomeric quinolinone (AM-0466) that affords robust in vivo activity, *J. Med. Chem.* **60**, 5990 (2017).

28. S. J. McKerrall, *et al.*, Structure- and ligand-based discovery of chromane arylsulfonamide Na$_V$1.7 inhibitors for the treatment of chronic pain, *J. Med. Chem.* **62**, 4091 (2019).

29. A. J. Roecker, *et al.*, Discovery of arylsulfonamide Na$_V$1.7 inhibitors: IVIVC, MPO methods, and optimization of selectivity profile, *ACS Med. Chem. Lett.* **12**, 1038 (2021).

30. T. Focken, *et al.*, Discovery of aryl sulfonamides as isoform-selective inhibitors of Na$_V$1.7 with efficacy in rodent pain models, *ACS Med. Chem. Lett.* **7**, 277 (2016)

31. Y-J. W., *et al.*, Development of new benzenesulfonamides as potent and selective Na$_V$1.7 inhibitors for the treatment of pain, *J. Med. Chem.* **60**, 2513 (2017).

32. J. Payandeh, M. Volgraf, Ligand binding at the protein-lipid interface: strategic considerations for drug design. *Nat. Rev. Drug Discov.* **20**, 710 (2021).

33. G. Bankar, et al., Selective Na$_V$1.7 antagonists with long residence time show improved efficacy against inflammatory and neuropathic pain. *Cell Rep.* **24**, 3133 (2018).

34. G. Luo, *et al.*, Discovery of indole- and indazole-acylsulfonamides as potent and selective Na$_V$1.7 inhibitors for the treatment of pain, *J. Med. Chem.* **62**, 831 (2019).

35. E. F. DiMauro, *et al.*, Application of a parallel synthetic strategy in the discovery of biaryl acyl sulfonamides as efficient and selective Na$_V$1.7 inhibitors, *J. Med. Chem.* **59**, 7818 (2016).

36. T. Focken, *et al.*, Design of conformationally constrained acyl sulfonamide isosteres: identification of N-([1,2,4]triazolo[4,3-a]pyridin-3-yl)methane-sulfonamides as potent and selective hNa$_V$1.7 inhibitors for the treatment of pain, *J. Med. Chem.* **61**, 4810 (2018).

37. X. Li, et al., Structural basis for modulation of human Na$_V$1.3 by clinical drug and selective antagonist. *Nat. Commun.* **13**, 1286 (2022).

38. N. K. Kotla, K. A. Kumari, S. K. Gunda, M. Shaik, 3D-QSAR and molecular docking studies of acylsulfonamides as Na$_V$1.7 inhibitors in chronic pain. *Res. J. Life Sci. Bioinform. Pharm. Chem. Sci.* **5**, 120 (2019).

39. R. P. Mason, D. G. Rhodes, L. G. Herbette, Reevaluating equilibrium and kinetic binding parameters for lipophilic drugs based on a structural model for drug interactions with biological membranes. *J. Med. Chem.* **34**, 869 (1991).

40. R. A. Coleman, M. Johnson, A. T. Nials, C. J. Vardey, Exosites: their current status, and their relevance to the duration of action of long- acting beta 2-adrenoceptor agonists. *Trends Pharmacol. Sci.* **17**, 324 (1996).

41. G. P. Anderson, A. Lindén, K. F. Rabe, Why are long-acting beta-adrenoceptor agonists long-acting? *Eur. Respir. J.* **7**, 569 (1994).

42. M. Masureel, *et al.*, Structural insights into binding specificity, efficacy and bias of a β2 AR partial agonist. *Nat. Chem. Biol.* **14**, 1059 (2018).

43. R. P. Austin, *et al*., QSAR and the rational design of long- acting dual D2-receptor/beta 2-adrenoceptor agonists. *J. Med. Chem*. **46**, 3210 (2003).

44. D. G. Rhodes, R. Newton, R. Butler, L. Herbette, Equilibrium and kinetic studies of the interactions of salmeterol with membrane bilayers. *Mol. Pharmacol.* **42**, 596 (1992).

45. D. A. Sykes, *et al.*, Observed drug- receptor association rates are governed by membrane affinity: the importance of establishing "micro- pharmacokinetic/ pharmacodynamic relationships" at the β2-adrenoceptor. *Mol. Pharmacol.* **85**, 608 (2014).

46. C. J. Dickson, *et al.*, Uncoupling the structure−activity relationships of β2 adrenergic receptor ligands from membrane binding. *J. Med. Chem.* **59**, 5780 (2016).

47. Z. Zhao, *et al.*, Exploration of type II binding mode: a privileged approach for kinase inhibitor focused drug discovery? *ACS Chem. Biol.* **9**, 1230 (2014).

48. D. N. Mastronarde, Automated electron microscope tomography using robust prediction of specimen movements. *J. Struct. Biol.* **152**, 36 (2005).

49. S. H. W. Scheres, RELION: implementation of a Bayesian approach to cryo-EM structure determination. *J. Struct. Biol.* **180**, 519 (2012).

50. T. Grant, A. Rohou, N. Grigorieff, *cis*TEM, user-friendly software for single-particle image processing. *Elife* **7**, e35383 (2018).

51. M. Kschonsak, *et al.* Structural architecture of the human NALCN channelosome. *Nature* **603**, 180 (2022).

52. S. Q. Zheng *et al.* MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nature Methods* **14**, 331 (2017).

53. A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron micrographs. *J. Struct. Biol.* **192**, 216 (2015).

54. T. C. Terwilliger, S. J. Ludtke, R. J. Read, P. D. Adams, P. V. Afonine, Improvement of cryo-EM maps by density modification. *Nature Methods* **17**, 923 (2020).

55. P. V. Afonine, *et al*. Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta Crystallogr. D. Struct. Biol.* **74**, 531 (2018).

56. N. W. Moriarty, R. W. Grosse-Kunstleve, P. D. Adams, electronic Ligand Builder and Optimization Workbench (eLBOW): a tool for ligand coordinate and restraint generation. *Acta Crystallogr. D. Biol. Crystallogr.* **65**, 1074 (2009).

57. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta Crystallogr. D. Biol. Crystallogr.* **66**, 486 (2010).

58. P. V. Afonine, *et al*. New tools for the analysis and validation of cryo-EM maps and atomic models. *Acta Crystallogr. D Struct. Biol.* **74**, 814 (2018).

59. C. J. Williams, *et al*. MolProbity: More and better reference data for improved all-atom structure validation. *Protein Sci.* **27**, 293 (2018).

60. E. F. Pettersen, *et al*. UCSF Chimera--a visualization system for exploratory research and analysis. *J. Comput. Chem.* **25**, 1605 (2004).

61. T. D. Goddard, *et al*. UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Sci.* **27**, 14 (2018).

62. D. Sutherlin *et al*. Therapeutic compounds and methods of use thereof. US11028075B2, June 8, 2021.

716
717
718
719

Figure 1

A

VSD4-Na$_V$1.7-Na$_V$Pas chimera

VSD4
S1 S2 S3 S4

NTD

CTD

B

C

VSD1

VSD2

VSD3

VSD4

top view

D

Extracellular

Intracellular

front view

E

Figure 2

A

GNE-3565

B VSD4

top view

C side view

D extracellular view

S2

S1

E1524

W1538

Y1537

N1540

R2

R3

A1585

S3

S4

lipid exposed pocket   selectivity pocket   anion binding pocket

E

F view towards membrane

S3   Y1537   S2

A1585

N1540

3.4

3.3

3.1

R4

G view from membrane

S2   Y1537   S3

W1538

D1586

3.0

F1583

Figure 3

A GDC-0310

B VSD4 — top view

C side view

D extracellular view
S2, S1, E1524, W1538, N1540, Y1537, R3, A1585, R2, S3, S4

E

F view towards membrane
S3, S2, A1585, Y1537, N1540, 3.1, 3.2, 2.9, R4

G view from membrane
S3, S4, I1588, I1601, I1574, I1604

Figure 4

Arylsulfonamide

Acylsulfonamide

Hybrid