1
2  **CryoEM reveals unprecedented binding site for Na$_V$1.7 inhibitors enabling rational**
3  **design of potent hybrid inhibitors**
4
5

6  Marc Kschonsak[1*], Christine C. Jao[1*], Christopher P. Arthur[1], Alexis L. Rohou[1], Philippe
7  Bergeron[2], Daniel F. Ortwine[2], Steven J. McKerrall[2], David H. Hackos[3], Lunbin Deng[3],
8  Jun Chen[4], Peter S. Dragovich[2], Matthew Volgraf[2], Matthew R. Wright[5], Jian
9  Payandeh[1,#], Claudio Ciferri[1,#], John C. Tellis[2,#]
10
[1]Genentech Inc., Structural Biology; South San Francisco, CA, United States.
[2]Genentech Inc., Discovery Chemistry; South San Francisco, CA, United States.
[3]Genentech Inc., Neuroscience; South San Francisco, CA, United States.
[4]Genentech Inc., Biochemical and Cellular Pharmacology; South San Francisco, CA,
United States.
[5]Genentech Inc., Drug Metabolism and Pharmacokinetics; South San Francisco, CA,
United States.
18
[#]Corresponding       authors.        Email:       jpayandeh@exelixis.com        (J.P);
ciferri.claudio@gene.com (C.C.); tellis.john@gene.com (J.C.T.).

**Author Contributions:** J.P., C.C., and J.C.T. designed research, M.K., C.J., C.P.A.,
A.R., D.H., and J.C., performed research, J.C.T, P.B., D.O., S.M., P.S.D., and M.V.
analyzed data and contributed to compound design. J.C.T., C.C., J.P., and M.K. wrote the
paper.

**Competing Interest Statement:** M.K., C.C.J, A.L.R, D.F.O, D.H.H, L.D, J.C, P.S.D,
M.W, C.C and J.C.T are Genentech employees. All authors declare no competing
financial interest.

30
31
32
33

**Abstract**

The voltage-gated sodium ($Na_V$) channel $Na_V1.7$ has been identified as a potential novel
pain target due to its striking human genetics. However, clinically available drugs (e.g.
lidocaine, carbamazepine, etc.) are not selective among the nine $Na_V$ channel subtypes,
$Na_V1.1$-$Na_V1.9$, and the two currently known classes of $Na_V1.7$ subtype-selective
inhibitors (aryl- and acylsulfonamides) have undesirable characteristics that may limit
their development. Moreover, understanding of the structure-activity relationships of the
acylsulfonamide class of $Na_V1.7$ inhibitors, exemplified by the clinical development
candidate **GDC-0310**, has been based solely on a single co-crystal structure of an
arylsulfonamide inhibitor series. To advance inhibitor design targeting the $Na_V1.7$
channel, we established an iterative system to routinely obtain high-resolution ligand-
bound $Na_V1.7$ structures using cryogenic electron microscopy (cryo-EM). We report that
**GDC-0310** engages the $Na_V1.7$ voltage-sensing domain 4 (VSD4) through an unexpected
binding mode orthogonal to the arylsulfonamide class binding pose, which identifies a
previously unknown ligand binding site in $Na_V$ channels. This finding enabled the design
of a novel hybrid inhibitor series that bridges the aryl and acylsulfonamide binding
pockets and allows for the generation of molecules with substantially differentiated
structures and properties. Overall, this study highlights the power of cryo-EM methods
to pursue challenging drug targets using iterative and high-resolution structure-guided
inhibitor design. It also underscores an important role of the membrane bilayer in the
discovery of selective $Na_V$ channel modulators.

55
56
57    **Introduction**
58
Voltage-gated sodium ($Na_V$) channels initiate and propagate action potentials in
excitable tissues and play important roles in health and disease.[1,2] The $Na_V1.7$ channel is
expressed predominantly in the peripheral nervous system and genetic studies have
identified compelling loss-of-function and gain-of-function phenotypes in human pain
syndromes, prompting significant efforts to develop $Na_V1.7$-selective inhibitors as
potential novel analgesic drugs.[3-5] $Na_V$ channels contain 24-transmembrane segments
linked in four homologous domains (DI-DIV), where four peripheral voltage-sensor
domains (VSD1-4) surround a central ion-conducting pore module that houses the ion
selectivity filter and key ligand and toxin receptor sites. Traditionally, all clinically
available $Na_V$ channel inhibitors lack significant molecular selectivity among the
$Na_V1.1$–1.9 subtypes owing to the high sequence conservation found at the ligand
binding site within the central cavity of the ion conducting pore module.[5,6]
71
A breakthrough study in 2013 by McCormack and colleagues reported the
discovery of an arylsulfonamide antagonist, **PF-04856264**, that bound to an
unprecedented receptor site in VSD4 with demonstrated molecular selectivity for human
$Na_V1.7$ over other subtypes.[7] However, the related development candidate **PF-05089771**
did not meet clinical endpoints in human subjects with painful diabetic peripheral
neuropathy, possibly due to poor target coverage and the intolerable doses required to
achieve efficacy.[8-10] Additionally, an alternate acylsulfonamide inhibitor that also

targeted the VSD4 receptor site in $Na_V1.7$, **GDC-0276**, was halted in Phase I clinical
trials due to safety concerns and potential off-target effects likely attributed to the high
lipophilicity of the compound.[11,12] To date, selective $Na_V1.7$ inhibitors with an improved
therapeutic index relative to these clinical-stage compounds have not yet been identified,
and this absence underscores the need to optimize such molecules using modern,
structure-guided design approaches.
85
The inherent complexity and dynamic nature of human $Na_V$ channels have
historically presented significant barriers to obtaining high-resolution experimental
structural information, especially of inhibitor-bound complexes.[13] Using an engineered
human VSD4 $Na_V1.7$-$Na_VAb$ bacterial channel chimera and X-ray crystallography, the
binding mode of the **GX-936** arylsulfonamide inhibitor (Supplementary Figure 1)
revealed that the anionic sulfonamide group engages the fourth arginine gating charge
(R4) to trap VSD4 in an activated conformation, which stabilizes a non-conductive,
inactivated state of the channel.[14] While determinants of subtype selectivity and
structure-activity relationships (SAR) of the arylsulfonamide inhibitor class could be
rationalized by the **GX-936** co-crystal structure, additional ligand-bound structures of
suitable resolution were not returned by the $Na_V1.7$-$Na_VAb$ chimeric channel
crystallography system. This shortcoming led molecular docking studies to presume that
the acylsulfonamide inhibitors also bound VSD4 in an analogous manner, despite points
of inexplicable SAR.[15] The overall paucity of direct structural information has made the
optimization of $Na_V1.7$ inhibitors challenging, with many questions about the
determinants of potency, selectivity, and the relationship between the acylsulfonamide
and arylsulfonamide inhibitor classes remaining unanswered.
103
Over the last three decades, protein crystallography and structure-based drug
design (SBDD) have become gold standards across the pharmaceutical industry for the
identification of ligand binding pockets and the optimization of drug candidates for
clinical development. While SBDD has proven successful for many important targets,
including G protein-coupled receptors (GPCRs),[16] its applicability to several membrane
protein targets, such as $Na_V$ channels, has been limited due to the extreme difficulties
with their iterative crystallization and structure determination.[13] Cryogenic electron
microscopy (cryo-EM) has recently emerged as a transformative technique to determine
the high-resolution structures of diverse protein targets and has proven particularly
effective for membrane proteins that are recalcitrant to crystallization.[17] Despite this
breakthrough, cryo-EM structure determination of membrane targets in complex with
small or large molecule therapeutics frequently remains retrospective and is often enabled
after the advancement of key molecules into the clinic.[18,19] Sluggish turn-around times
and modest resolutions typically afforded by cryo-EM have limited its application for
real-time SBDD efforts. Here, we describe a system for the iterative determination of
high-resolution $Na_V1.7$ small molecule co-structures via cryo-EM that has led to the
development of a novel class of inhibitors. Critical to this advancement has been
establishment of a robust protocol for sample preparation and structure determination and
the enablement of the first structure of an acylsulfonamide bound to $Na_V1.7$ VSD4,
revealing a previously unknown pocket between the S3 and S4 helices. Consequently, a
novel hybrid inhibitor series bridging the aryl and acylsulfonamide pockets was

generated. Our work exemplifies the deployment of cryo-EM as a workhorse structural
biology tool in an active medicinal chemistry campaign and thus represents an important
milestone toward Na$_V$ channel drug discovery.
128
129
130  **Results**
131
**Structure of the G-3565-VSD4 Na$_V$1.7-Na$_V$Pas channel complex in lipid nanodiscs**
Our approach to establish an iterative, high-resolution system to enable Na$_V$1.7
SBDD was ultimately guided by three observations: our repeated failure to generate well-
diffracting crystals of the Na$_V$1.7-Na$_V$Ab bacterial channel chimera system,[14] our
difficulty of reproducibly expressing a suitable amount of full-length human Na$_V$1.7
channel,[20,21] and the very limited local resolution observed at VSD4 in all available
human Nav1.7 cryo-EM structures.[22,23]  Thus, we sought to exploit an engineered human
VSD4-Na$_V$1.7-Na$_V$Pas cockroach channel chimeric construct that had been previously
shown to complex with small molecule inhibitors and peptide toxins known to target
VSD4.[21]  The robust protein yields following expression and purification allowed us to
readily pursue reconstitution of the VSD4-Na$_V$1.7-Na$_V$Pas channel into lipid nanodisc
with high recovery (Supplementary Figure 2).  Small molecule inhibitors were added
prior to sample vitrification, followed by cryo-EM data collection and processing
procedures, which allowed us to routinely obtain 3D-reconstructions in the 2.2-3.0 Å
resolution range (Figure 1, Supplementary Figure 3).
147
The VSD4-Na$_V$1.7-Na$_V$Pas chimera displays the expected domain-swapped
arrangement with numerous densities assigned as phospholipids bound to the channel,
confirming the maintenance of a native membrane-like environment (Figure 1). As seen
in previous Na$_V$Pas structures,[21,24,25] the ion-conducting pore module of the VSD4-
Na$_V$1.7-Na$_V$Pas chimeric channel is closed, consistent with a nonconductive or
inactivated state. The quality and resolution of our structure allowed us to assign 112
water molecules for the highest resolution structure (Figure 1, Video 1). The visualization
of many well-resolved water molecules bound within the VSDs and coordinated within
the ion selectivity filter provides new insights into the interactions that might occur
during gating charge transfer and ion conduction, respectively.
158
Although we were able to determine the crystal structure of VSD4-Na$_V$Ab bound
to the arylsulfonamide inhibitor **GX-936**,[14] no well-diffracting crystals were obtained for
any other arylsulfonamide compounds despite years of continued effort. Arylsulfonamide
**GNE-3565** is representative of an advanced series of arylsulfonamide class Na$_V$1.7
inhibitors that demonstrates channel blockage at subnanomolar concentrations with
mixed subtype selectivity (Figure 2A). We complexed **GNE-3565** with the VSD4
Na$_V$1.7-Na$_V$Pas channel-nanodisc system and employed cryo-EM to assess its overall
binding pose (Figure 2B, C).  The 2.9 Å resolution cryo-EM map revealed that the S1-S2
and S3-S4 helices from VSD4 form a clamshell-like structure that closes over **GNE-3565**
(Figure 2D, E, Supplementary Data Figure 4A-D, Supplementary Data Table 1), similar
to the binding pose reported for **GX-936**.[14]  Specifically, the ionized arylsulfonamide
group of **GNE-3565** (measured pKa = 5.8) bisects VSD4 to salt-bridge directly to R4 on

the S4 helix, while the central phenyl ring extends perpendicular between the S2 and S3
helices to directly contact established subtype-selectivity determinants Tyr1537 and
Trp1538 on the S2 helix (Figure 2A-D).  As for **GX-936**, the **GNE-3565** arylsulfonamide
VSD4 receptor binding site can be divided into three regions: an anion-binding pocket, a
selectivity pocket, and a lipid-exposed pocket (Figure 2D).   The root-mean-square
deviation (RMSD) between the **GNE-3565**-VSD4 and the **GX-936**-VSD4 structures[14] is
0.667 (714 to 714 atoms), (Supplementary Figure 5A), supporting long-held assumptions
by medicinal chemistry teams that all structurally-related arylsulfonamides should be
expected to complex to VSD4 through similar determinants.[8,26-31]
180
**Structure of the acylsulfonamide GDC-0310 in complex with the VSD4 Na$_V$1.7-**
**Na$_V$Pas channel**
To investigate the binding pose of a representative Na$_V$1.7 selective
acylsulfonamide, **GDC-0310** (Figure 3A) was complexed with the VSD4 Na$_V$1.7-Na$_V$Pas
channel in nanodiscs and a cryo-EM structure was determined to 2.5 Å resolution (Figure
3B-D, Supplementary Data Figure 3G-J, Supplementary Data Table 1).  Remarkably, the
binding mode of **GDC-0310** revealed an unexpected and previously unknown pocket that
formed between the S3 and S4 helices (Figure 3B-E).  While the aryl and acyl moieties of
**GDC-0310** and **GNE-3565** bind in the same region, the remainder of the acyl receptor
site is orthogonal to the corresponding binding domain reported for the aryl class (Figure
3B-D).  In detail, the anionic acylsulfonamide of **GDC-0310** participates in salt-bridge
bonding interactions with R4 and R3 through the carbonyl and sulfonamide oxygen
atoms, respectively (Figure 3F).  Moreover, the aryl ring of the **GDC-0310** splits the S3
and S4 helices to occupy a lipophilic pocket, displacing the S3 helix laterally by ~3 Å
relative to the **GNE-3565**-VSD4 structure (Figure 3G). Our cryo-EM inhibitor bound
structures serve to highlight a dynamic environment in VSD4 where the S1-S4 helices
can be differentially separated by distinct small molecule binding poses (Figure 2D,
Figure 3D).
199
It is notable that the cyclopropyl substituent of **GDC-0310** makes effective van
der Waals interactions with I1574 and I1588 from S3 and I1601 and I1604 on S4 (Figure
3G), while the alpha-methylbenzylamine "tail" region sits almost entirely within the lipid
bilayer (Figure 3A, C), where density for this portion of the inhibitor is only poorly
resolved (Figure 3E).  Considering the relative depth of the **GDC-0310** binding site in
relation to the membrane-water interface (~8 Å), our structure suggest a membrane
access pathway for acylsulfonamides to the VSD4 receptor site, which has important
implications for understanding the pharmacology, SAR, and potential development
liabilities of the series.[32]  Notably, this structure also reveals that the Tyr1537 side-chain
on S2 exists in a down rotamer and does not contact **GDC-0310** directly, which provides
the first direct structural rationale for why the selectivity profiles between the acyl and
arylsulfonamide classes differ substantially (Figure 3D, *vide infra*).[33]
212
**Structure-based rational design of hybrid Na$_V$1.7 inhibitors**
Upon inspection of the superimposed **GNE-3565**-aryl and **GDC-0310**-acyl
structures, we could immediately envision a novel class of hybrid inhibitors that would
simultaneously occupy both the aryl- and acylsulfonamide binding pockets (Figure 4A,

B). Such an enlarged hybrid binding pocket would offer unique opportunities to gain
potency directly from ligand-protein interactions, which might enable removal of the
hydrophobic lipid-facing tail groups seen in all previous generations of VSD4-targeting
$Na_V$ channel inhibitors. These groups have typically been critical for maintaining
potency, but use of the plasma membrane to drive membrane-bound target occupancy is
largely nonspecific, and potentially introduces off-target liabilities that can contribute to
toxicity. However, it was unclear if this hybrid conformation of the VSD4 domain could
accommodate the described inhibitors and/or whether it was accessible during any stage
of $Na_V1.7$ channel activation. We therefore set out to demonstrate proof-of-concept that
appropriately designed small molecules could induce this hypothesized receptor state
while inhibiting channel function with meaningful potency.
228
**GDC-0310** was a relatively unattractive starting point for hybridization because
of its high molecular weight (543 g/mol) and lipophilicity (cLogP = 5.2). Accordingly,
analysis of historical data identified progenitor compound **1**, which offers reasonable
potency against $Na_V1.7$ ($IC_{50}$ = 25 nM, n = 8; Fig 4C) and reduced molecular weight (350
g/mol) and lipophilicity (cLogP = 2.7). Seeking to leverage existing knowledge of the
arylsulfonamide binding pocket, molecule **2** was synthesized in an attempt to directly
graft on a significant portion of the **GNE-3565** scaffold. Gratifyingly, this molecule
retained substantial potency, affording channel blockade at an $IC_{50}$ of 47 nM (n = 8).
237
A cryo-EM structure of **2** bound to the VSD4 $Na_V1.7$-$Na_V$Pas channel confirmed
that the molecule adopts the desired hybrid binding mode (Figure 4D). Here, the anionic
group of the novel hybrid compound binds in the same position as in the aryl and acyl
poses, interacting closely with R4, as seen for the aryl-like binding (Figure 4E). The aryl-
and acylsulfonamide-derived structural motifs of **2** occupy their expected positions in the
pockets formed between S2/S3 and S3/S4, respectively. Notably, in this hybrid
configuration, Y1537 adopts a similar arrangement as observed for the arylsulfonamides
(Figure 4F).
246
Encouraged by this result, we hoped to draw on previously established knowledge
of acylsulfonamide SAR to improve the potency of this compound. Established SAR
within the acylsulfonamide series suggested that replacement of the chloro substituent on
the benzamide fragment of the molecule with a cyclopropane would typically improve
potency. This SAR proved translatable to the hybrid class molecules, resulting in
inhibitor **3** ($IC_{50}$ = 16 nM, n = 12; Figure 4C).
253
Further analysis of the cryo-EM structure of **2** revealed that the aryl ring in the
S2/S3 binding pocket was shifted ~1 Å away from the VSD4 core when compared to
arylsulfonamides such as **GNE-3565** (data not shown). The aryl ring in the S3/S4 pocket
exhibited better overlap with the analogous substituents in the **GDC-0310** structure. We
hypothesized that the acylsulfonamide moiety of **2**, which places the two aryl substituents
4.7 Å apart, might not be fully optimized for use in hybrid inhibitors. In comparison, an
arylsulfonamide moiety would position these two groups at a distance of ~3.3 Å. On this
basis we designed *N*-benzothiazolyl sulfonamide **4**, which demonstrated potent $Na_V1.7$
inhibition ($IC_{50}$ = 9 nM, n = 8, Figure 4C).

263
Our last hypothesis was again validated with a cryo-EM co-structure of **4** bound
to the VSD4 $Na_V1.7$-$Na_V$Pas channel (Figure 4G). Here, we observed the typical face-on
pi-stacking orientation of the S2/S3 substituent with Y1537 while maintaining efficient
space filling in the S3/S4 pocket. Moreover, the efficient occupation of the non-
membrane exposed regions of the binding pocket permitted the removal of the lipophilic
tail of the molecule in **5**, where a $CF_3$ group replaces the cyclopentylmethyl substituent.
Although this molecule shows reduced potency ($IC_{50}$ = 74 nM, n = 8), lack of substantial
small molecule contact with the lipid bilayer is an uncommon feature in all other known
$Na_V1.7$ VSD4 domain inhibitors. These molecules serve as a key proof of concept,
highlighting the power of cryoEM to yield new structural insights that can inspire the
design of novel, structurally differentiated small molecule inhibitors.
275
276
277  **Discussion**
278
Iterative structure-based design is among the most powerful techniques to
facilitate the development of small molecule drug candidates. Although a variety of
technologies are capable of providing information about small molecule ligand-protein
interactions, SBDD has traditionally been synonymous with the use of X-ray
crystallography, with other techniques failing to rival its high resolution and fast cycle
times. Here, we have demonstrated that a robust cryo-EM structure pipeline is sufficient
to quickly provide co-structures at resolutions suitable for use in an active medicinal
chemistry optimization campaign. Moreover, we have highlighted the power of structural
methods, and cryo-EM in particular, to provide new and important insights into inhibitor
design through the discovery of a previously unknown binding mode for acylsulfonamide
$Na_V1.7$ inhibitors. We further note that our use of an engineered VSD4-$Na_V1.7$-$Na_V$Pas
chimeric channel construct was essential to produce high yields of recombinant protein
capable of routinely returning multiple cryo-EM samples per preparation. This system
repeatedly delivered high-resolution depictions of the VSD4 inhibitor binding site and
thus strongly contrasts with all prior accounts of cryo-EM structures of full-length human
$Na_V1.7$ channels.[22,23] Importantly, high quality cryo-EM maps at VSD4 in complex with
ICA121431 have recently been reported from full-length human $Na_V1.3$ channel
protein,[37] indicating that SBDD efforts targeting a particular $Na_V$ channel subtype would
benefit from exploring both engineered and native channel sample options at early project
stages.
299
The co-structure of **GDC-0310** was critical to contextualize historical differences
between aryl- and acylsulfonamide $Na_V1.7$ inhibitors, which include divergent SAR in
the linker phenyl ring and tail regions, disparate $Na_V$ family selectivity patterns, and
differences in on-rates and off-rates.[12,15,33-36] Previous efforts to model acylsulfonamide
binding have attempted to dock molecules into the **GX-936** co-structure and have
proposed subtle changes in molecular register and/or pose to explain the non-translatable
SAR.[15,38] Here, we demonstrate that the molecular features of aryl- and acylsulfonamide
inhibitors diverge because they occupy distinct, but overlapping, pockets within the
VSD4 domain.

The selectivity determinants of arylsulfonamide Na$_V$1.7 inhibitors have been well
studied. Mutation of key non-conserved residues proximal to the binding pocket observed
in the **GX-936** co-structure have been shown to alter Na$_V$1.7 potency consistent with
selectivity patterns observed for other Na$_V$ family members.[14] Typically,
arylsulfonamides have shown high levels of selectivity against Na$_V$1.5, Na$_V$1.1, and
Na$_V$1.4, but lower selectivity against Na$_V$1.2 and Na$_V$1.6. In comparison, several
acylsulfonamides have been characterized with very high selectivity over Na$_V$1.2 and
Na$_V$1.6.[33] Interestingly, residues forming the contours of the acylsulfonamide binding
pocket show high sequence homology across Na$_V$ isoforms, suggesting that selectivity for
this class is likely an allosteric phenomenon. Nonetheless, the disparate binding pockets
for these two inhibitor classes provide a clear rationale for their historically divergent
patterns of selectivity.

Our structure of **GDC-0310** provides a structural rationale for the relative slow
dissociation kinetics of acylsulfonamide inhibitors compared to arylsulfonamides.[33]
Specifically, the acylsulfonamide pocket is buried more deeply into the plasma
membrane than the arylsulfonamide pocket, and may be available only through a
membrane-access pathway that first involves partitioning of the molecule into the
intramembrane space. In comparison, the arylsulfonamide pocket is open to the outer
membrane interface, offering access from the solvent compartment. Slow dissociation
kinetics are known to be associated with membrane-access pathways for small molecules
binding to transmembrane proteins.[32,39] For example, the long duration of action of
salmeterol (a long-acting $\beta_2$ adrenergic receptor agonist) is proposed to be a result of its
membrane partitioning properties. In comparison, salbutamol, which binds to the same
target and site but through a solvent-access pathway, shows much faster dissociation
from the receptor.[40-46]

In addition to offering valuable information for retrospective analysis of historical
Na$_V$1.7 VSD4 domain inhibitors, the structure of **GDC-0310** inspired the development of
a class of structurally differentiated molecules that bind to a novel conformation of the
receptor. These hybrid molecules bridge both the S2/S3 and S3/S4 helical gaps, with the
anionic moiety situated centrally, interacting with the arginine network at the core of the
domain. We have demonstrated proof-of-concept that potent hybrid inhibitors can be
designed with fewer membrane-associated elements than prior generation Na$_V$1.7
inhibitors, and anticipate that this will offer opportunities to develop molecules with
improved lipophilic ligand efficiency and/or differentiated pharmacokinetic profiles.
Hybrid inhibitors can also access selectivity determinants from both the S2/S3 and S3/S4
pockets simultaneously, offering a tantalizing opportunity to develop exquisite selectivity
over other Na$_V$ isoforms.

Beyond the scope of Na$_V$ channel inhibitors, our experiences may also offer
lessons translatable to small molecule modulators of other transmembrane proteins.
Specifically, it is noteworthy to observe that both aryl- and acylsulfonamide inhibitors
have access to cryptic pockets that are not observable in the co-structures of the parent
proteins. Designing small molecules to "push" on a structural protein element is a

common strategy in medicinal chemistry, but it is often challenging to predict when such
efforts will afford new, well-defined binding pockets. As such, these designs are often
met with failure and quickly abandoned. However, some scenarios exist in which a
cryptic pocket is more likely to be found. In these cases, medicinal chemists are more
often emboldened to continue experimenting with designs that push into new areas of a
protein binding pocket, even when met with initial failure. For example, type II kinase
inhibitors can be rationally designed starting from a type I inhibitor structure by
appending groups that push through the DFG into the cryptic back pocket.[47] This pocket
is not visible from the type I inhibitor structure, but knowledge of the protein dynamics
associated with the DFG-in/DFG-out transition provides a rationale for exploring these
designs. We speculate that transmembrane intrahelical domains, particularly those in ion
channels, transporters, and GPCRs, may present similarly privileged protein structural
motifs that are rich with opportunities for the discovery of druggable cryptic pockets.
368
In summary, we have developed a robust pipeline for high-resolution structural
determination of small-molecule bound $Na_V1.7$ channels suitable for use in iterative
structure-based drug design campaigns. We also report the first co-structure of an
acylsulfonamide inhibitor bound to the VSD4 domain of $Na_V1.7$, revealing an
unexpected and unique binding mode. This finding provides clear rationale for previously
unexplainable divergence in the in vitro pharmacological behavior between the two
inhibitor classes, and inspired the development of hybrid inhibitors that merge structural
features from both inhibitor classes, engaging the receptor in a novel conformation. Our
findings highlight the power of cryoEM as an enabling drug discovery technology and
offer interesting learnings regarding ion channel structural dynamics that are potentially
applicable to other related targets and target classes.
380
381  **Materials and Methods**
382
**Generation of** VSD4-$Na_V1.7$-$Na_V$Pas channel constructs
VSD4-$Na_V1.7$-$Na_V$Pas chimeric constructs were used as described previously.[21] In brief,
optimized coding DNA for $Na_V1.7$ VSD4-$Na_V$PaS chimeras with N-terminal tandem
StrepII and FLAG tag was cloned into a pRK vector with CMV promoter. HEK293 cells
in suspension were cultured in SMM 293T-I medium under 5% $CO_2$ at 37°C and
transfected using PEI when the cell density reached $4 \times 10^6$ cells per ml. Transfected cells
were harvested 48 hours after transfection. The Dc1a toxin coding DNA from *Diguetia*
*canities* was cloned into a modified pAcGP67A vector downstream of the polyhedron
promoter and an N-terminal 6x HIS tag. Recombinant baculovirus was generated using
the Baculogold system (BD Biosciences) and *Trichoplusia ni* cells were infected for
protein production. The supernatant was harvested 48 h post-infection.
394
**Protein expression and purification:**
One hundred fifty grams of cell pellet was resuspended in 500mL of 25mM Hepes pH
7.5, 200mM NaCl, 1ug/mL benzonase, 1mM PMSF, and Roche protease inhibitor tablets.
Cells were lysed by dounce homogenization. Proteins were solubilized by addition of 2%
GDN (Avanti) with 0.3% cholesteryl hemisuccinate (Avanti) for 2 hours at 4°C. Cell
debris was separated by ultracentrifugation at 40,000 rpm at 4°C. Affinity purification

using anti-Flag resin was performed by batch binding for 1 hour at $4^{\circ}$C. The resin was
washed with 5CV Purification Buffer (25mM Hepes pH 7.5, 200mM NaCl, 0.01%
GDN). Another 5CV wash was performed using Purification Buffer supplemented with
5mM ATP and 10mM $MgCl_2$. Proteins were eluted in 6 CV Purification Buffer with
300ug/mL Flag peptide. Proteins were subjected to another round of affinity purification
using StrepTactin resin (IBA). Proteins were eluted, and concentrated to ~ 5 mg/mL. For
the compound **2** containing sample, instead of nanodisc reconstitution VSD4-$Na_V$1.7-
$Na_V$Pas was incubated overnight with Dc1a toxin at a 2:1 molar ration of toxin:VSD4-
$Na_V$1.7-$Na_V$Pas. The eluted sample was concentrated to 100uL, and separated on
Superose 6 3.2/300.
411
For nanodisc reconstitution used for the samples containing **GNE-3565, GDC-0310** and
compound **4**, a 200-molar excess of lipid mix (3POPC:1POPE:1POPG resuspended in
50mM Hepes pH7.5, 100mM NaCl, 5mM $MgCl_2$, 1% CHAPS) was added to detergent-
solubilized protein and incubated on ice for 30 minutes. A four-molar excess of scaffold
protein (MSP1E3D1, Sigma) was added to the protein-lipid mix, and incubated on ice for
another 30 minutes. To remove detergent, BioBeads (Bio-Rad) were added to 0.25
mg/mL, and incubated overnight at $4^{\circ}$C. To remove the empty nanodiscs, the sample was
subjected to a round of affinity purification using Strep-Tactin (IBA). The eluted sample
was concentrated to 100uL, and separated on Superose 6 3.2/300. Peak fractions were
combined, and split into four samples. Fifty uM of the small molecule of interest was
added to each sample, and incubated at $22^{\circ}$C for 10 minutes. Samples were crosslinked
with 0.05% glutaraldehyde (Electron Microscopy Sciences) then quenched with 1M Tris
pH 7.0. Samples at 2 mg/mL were used for grid freezing.
425
426
**CryoEM sample preparation and data acquisition**
CryoEM grids for the small molecule VSD4-$Na_V$1.7-$Na_V$Pas chimeric complexes shown
in this study were prepared as follows: For the **GDC-0310** and compound **4** VSD4-
$Na_V$1.7-$Na_V$Pas complexes, holey carbon grids (Ultrafoil 25 nM Au R 0.6/1 300 mesh;
Quantifoil) were incubated with a thiol reactive, self-assembling reaction mixture of
4mM monothiolalkane(C11)PEG6-OH (11-mercaptoundecyl) hexaethyleneglycol (SPT-
0011P6, SensoPath Technologies, Inc., Bozeman, MT). Grids were incubated with this
self-assembled monolayer (SAM) solution for 24 hours and afterwards rinsed with EtOH.
3 μl of the sample was applied to the grid and blotted with Vitrobot Mark IV
(Thermofisher) using 3.5 s blotting time with 100% humidity and plunge-frozen in liquid
ethane cooled by liquid nitrogen. The **GNE-3565** VSD4-$Na_V$1.7-$Na_V$Pas complex was,
similarly as described above, applied to a holey carbon grid (Ultrafoil 25 nM Au R
1.2/1,3 300 mesh; Quantifoil) pretreated with SAM solution. The grid was blotted single-
sided with a Leica EM GP (Leica) using 3 s blotting time with 100% humidity and
plunge-frozen in liquid ethane cooled by liquid nitrogen. For compound **2** VSD4-$Na_V$1.7-
$Na_V$Pas-DC1a complex, holey carbon grids (Ultrafoil 25 nM Au R 2/2 200 mesh;
Quantifoil) were glow-discharged for 10 s using the Solarus plasma cleaner (Gatan). 3 μl
of the sample was applied to the grid and blotted with Vitrobot Mark IV (Thermofisher)
using 2.5 s blotting time with 100% humidity and plunge-frozen in liquid ethane cooled
by liquid nitrogen.

Movie stacks for compound **2** VSD4-Na$_V$1.7-Na$_V$Pas-DC1a complex were collected
using SerialEM[48] on a Titan Krios operated at 300 keV with bioquantum energy filter
equipped with a K2 Summit direct electron detector camera (Gatan). Images were
recorded at 165,000 × magnification corresponding to 0.824 Å per pixel, using a 20 eV
energy slit. Each image stack contains 50 frames recorded every 0.2 s for an accumulated
dose of ~50 e Å−2 and a total exposure time of 10 s. Images were recorded with a set
defocus range of 0.5 to 1.5 μm.

Movie stacks for **GNE-3565, GDC-0310** and compound **4** VSD4-Na$_V$1.7-Na$_V$Pas
complexes were collected using SerialEM on a Titan Krios G3i (ThermoFisher Scientific,
Waltham, MA) operated at 300 keV with bioquantum energy filter equipped with a K3
Summit direct electron detector camera (Gatan Inc, Pleasanton, CA). Images were
recorded in EFTEM mode at 105,000 × magnification corresponding to 0.838 Å per
pixel, using a 20 eV energy slit. Each image stack contains 60 frames recorded every 0.05
s for an accumulated dose of ~60 e Å−2 and a total exposure time of 3 s. Images were
recorded with a set defocus range of 0.5 to 1.5 μm.

**CryoEM data processing**
CryoEM data were processed using a combination of the RELION[49] and cisTEM[50]
software packages, similarly as described previously[51] and as illustrated in Suppl. Figure
2 and 3.

Movies were corrected for frame motion using the MotionCor2[52] implementation in
RELION and contrast-transfer function parameters were fit using the 30-4.5 Å band of
the spectrum with CTFFIND-4.[53] CTF fitted images were filtered on the basis of the
detected fit resolution better than 6-10 Å. Particles were picked by template-matching
with gautomatch (https://www2.mrc-lmb.cam.ac.uk/research/locally-developed-
software/zhang-software/#gauto) using a 30 Å low-pass filtered apo VSD4-Na$_V$1.7-
Na$_V$Pas reference structure. Particles were sorted during RELION 2D classification and
selected particles were imported into cisTEM for 3D refinements. 3D reconstructions
were obtained after auto-refine and manual refinements with a mask around the channel
(excluding the detergent micelle) and by applying low-pass filter outside the mask (filter
resolution 20 Å) and a score threshold of 0.10-0.30, so that only the best-scoring 10-30%
of particle images would be included in the 3D reconstruction at each cycle. The weight
outside of the mask was set to 0.8. No data beyond 3.0 Å for compound **4**, 3.4 Å for
**GDC-0310**, 3.7Å **GNE-3565** and 4.0 Å for compound **2** were used in the refinements.
Phenix ResolveCryoEM[54] density modification was applied to each of the reconstructions
to obtained the final map used for model building. Local resolution was determined in
cisTEM using a reimplementation of the blocres algorithm.

**Model building and structure analysis**
The previously determined VSD4-Na$_V$1.7-Na$_V$Pas model (PDB: 6NT3)[21] was fit as a
rigid body into the cryoEM map. After manual adjustments, multiple rounds of real space
refinement using the phenix.real_space_refinement tool[55] was used to correct structural
differences between the initial model and the map. The small molecule coordinates were

generated with eLBOW,[56] placed and manually adjusted in Coot,[57] energy minimized
with MOE (Chemical Computing Group ULC,) and refined with real space refinements
in Phenix[55]. The model was validated using phenix.validation_cryoem[58] with built-in
MolProbity scoring.[59] Figures were made using PyMOL (The PyMOL Molecular
Graphics System, v.2.07 Schrödinger, LLC), UCSF Chimera,[60] and UCSF ChimeraX.[61]
498
499 **Chemical synthesis**
500 The synthesis of **GNE-3565**, **GDC-0310**, and compound **1** have been previously
501 reported.[12,15,62] The synthesis and characterization of compounds **2-5** are described in the
502 supporting information.
503
**Na$_V$1.7 Syncropatch electrophysiology potency**
cDNAs for Na$_V$1.7 (NM_002977) were stably expressed in Chinese Hamster Ovary
(CHO) cells. Sodium currents were measured in the whole-cell configuration using
Syncropatch 384PE (NanIon Technologies, Germany). 1NPC®-384 chips with custom
medium resistance and single hole mode were used. Internal solution consisted of (in
mM): 110 CsCl, 10 CsCl, 20 EGTA, and 10 Hepes (pH adjusted to 7.2); and external
solution contains (in mM): 60 NMDG, 80 NaCl, 4 KCl, 1 MgCl$_2$, 2 CaCl$_2$, 2 D-Glucose
monohydrate, 10 Hepes (pH adjusted to 7.4 with NaOH).
512
After system flushing, testing compounds were dissolved in external solution containing
0.1% Pluronic F-127. 10µl cells were added to the chip from a cell hotel, and a negative
pressure of −50 mBar was applied to form a seal. Following treatment with seal enhancer
solution and wash-off with external solution, negative pressure of −250 mbar was applied
for 1 second to achieve the whole-cell configuration, followed by three washing steps in
external solution. 20µl of compounds were added to 40µl in each well (1:3 dilution of
compounds), and after mixing, 20µl was removed so the volume was retained at 40 ul.
After approximately 13 minutes recording, 20µl/well of 2 uM TTX was added to achieve
full block.
522
For voltage protocol, a holding potential of -50 mV was applied during the whole
experiment. A depolarizing step was applied to -10mV for 10ms, followed by a
hyperpolarization step to -150 mV for 20 ms to allow channel recovery from inactivation.
A second depolarizing step was applied from -150mV to -10mV for 10 ms, where
currents were measured to derive blocking effects of compounds. Inhibition was
determined based on 7.5 min of compound incubation.
529
530 **Figure Legends:**
531
**Figure 1: (A)** Schematic of the VSD4-Na$_V$1.7-Na$_V$Pas channel. The portions humanized
to the Na$_V$1.7 sequence are shown in green. N-terminal domain (NTD) and CTD are
indicated. **(B)** Side view of the single-particle cryo-EM reconstruction of VSD4- Na$_V$1.7-
Na$_V$Pas channel **(C-D)** Cartoon representations of the Top and Side views of VSD4-
Na$_V$1.7-Na$_V$Pas channel. Individual VSD domains are indicated. VSD4 is highlighted in
green. **(E)** Localization of water molecules (in red) in the VSD4-Na$_V$1.7-Na$_V$Pas channel
structure. VSD4 is highlighted in green.

539
**Figure 2: (A)** Chemical structures of arylsulfonamide **GNE-3565**. **(B,C)** Top and Side
views of VSD4-Na$_V$1.7-Na$_V$Pas channel bound to **GNE-3565**. VSD4 is highlighted in
green, **GNE-3565** in magenta. **(D)** Extracellular view of VSD4-Na$_V$1.7-Na$_V$Pas
arylsulfonamide receptor site is shown with select side chains rendered as sticks. **(E)** The
cryoEM map surrounding the ligand **GNE-3565** is shown in mesh representation. **(F)**
View towards the membrane highlighting key interactions with the **GNE-3565** warhead.
**(G)** View from the membrane highlighting key interactions with the **GNE-3565** central
phenyl ring.
548
**Figure 3: (A)** Chemical structures of acylsulfonamide **GDC-0310**. **(B)** Top and Side
views of VSD4-Na$_V$1.7-Na$_V$Pas channel bound to **GDC-0310**. VSD4 is highlighted in
green, **GDC-0310** in magenta. **(D)** Extracellular view of VSD4-Na$_V$1.7-Na$_V$Pas
acylsulfonamide receptor site is shown with select side chains rendered as sticks. **(E)**
The cryoEM map surrounding the ligand **GDC-0310** is shown in mesh representation.
**(F)** View towards the membrane highlighting key interactions with the **GDC-0310**
warhead. **(G)** View from the membrane highlighting van der Waals interactions with the
**GDC-0310** cyclopropyl substituent. The alpha-methylbenzylamine tail region sits almost
entirely within the lipid bilayer.
558
**Figure 4: (A)** Illustration of hybrid molecule design approach. **(B)** Arylsulfonamide,
acylsulfonamide and hybrid molecule poses **(C)** Hybridization strategy and molecule
optimization. Highlighted are molecule **2** and molecule **4** **(D)** Extracellular view of
VSD4-Na$_V$1.7-Na$_V$Pas bound to the hybrid molecule **2**. **(E)** View towards the membrane
highlighting key interactions with the anionic group. **(F)** View from the membrane
highlighting the lack of a stacking interaction between Y1537 and the phenyl ring. **(G)**
Extracellular view of VSD4-Na$_V$1.7-Na$_V$Pas bound to the hybrid molecule **4**. **(H)** View
towards the membrane highlighting key interactions with the anionic group. **(I)** View
from the membrane highlighting the π-stacking interaction between Y1537 and the
phenyl ring.
569
570

571 **Acknowledgments**
572
573 We are grateful to BMR and RMG for construct generation and protein scale-up. We are
574 thankful to Cameron Noland for help with the biochemistry of the VSD4-Na$_V$1.7-Na$_V$Pas
575 channel system. We thank Tommy Lai and Wenfeng Liu for chemistry support.

576
577
**Material and Data Availability:** All data generated or analyzed during this study are
included in the manuscript and supporting files. The Na$_V$1.7-Na$_V$Pas/GNE-3565
coordinates and cryo-EM maps were deposited in the PDB: XXX and EMDB: EMD-
XXX, respectively. The Na$_V$1.7-Na$_V$Pas/GDC-0310 coordinates and cryo-EM maps were
deposited in the PDB: XXX and EMDB: EMD-XXX, respectively. The Na$_V$1.7-Na$_V$Pas
bound to the hybrid molecule **2** coordinates and cryo-EM maps were deposited in the
PDB: XXX and EMDB: EMD-XXX, respectively. The Na$_V$1.7-Na$_V$Pas bound to the

hybrid molecule **4** coordinates and cryo-EM maps were deposited in the PDB: XXX and EMDB: EMD-XXX, respectively. Requests for materials should be addressed to J.C.T.

**References**

1. W. A. Catterall, A. L. Goldin, S. G. Waxman, International Union of Pharmacology. XLVII. Nomenclature and structure-function relationships of voltage-gated sodium channels. *Pharmacol. Rev.* **57**, 397 (2005).
2. C. A. Ahern, J. Payandeh, F. Bosmans, B. Chanda, The hitchhiker's guide to the voltage-gated sodium channel galaxy. *J. Gen. Physiol.* **147**, 1 (2016).
3. S. D. Dib-Hajj, Y. Yang, J. A. Black, S. G. Waxman, The Na(V)1.7 sodium channel: from molecule to man. *Nat. Rev. Neurosci.* **14**, 49 (2013).
4. J. Payandeh, D. H. Hackos, Selective ligands and drug discovery targeting the voltage-gated sodium channel $Na_V1.7$. *Handb. Exp. Pharmacol.* **246**, 271 (2018).
5. S. J. McKerrall, D. P. Sutherlin, $Na_v1.7$ inhibitors for the treatment of chronic pain. *Bioorg. Med. Chem. Lett.* **28**, 3141 (2018).
6. M. de Lera Ruiz, R. L. Kraus, Voltage-gated sodium channels: structure, function, pharmacology, and clinical indications. *J. Med. Chem.* **58**, 7093 (2015).
7. K. McCormack, *et al.*, Voltage sensor interaction site for selective small molecule inhibitors of voltage-gated sodium channels. *Proc. Natl. Acad. Sci. USA* **110**, E2724 (2013).
8. N. A. Swain, *et al.,* Discovery of clinical candidate 4-[2-(5-amino-1H-pyrazol-4-yl)-4-chlorophenoxy]-5-chloro-2-fluoro-N-1,3-thiazol-4-ylbenzenesulfonamide (PF-05089771): design and optimization of diaryl ether aryl sulfonamides as selective inhibitors of $Na_V1.7$. *J. Med. Chem.* **60**, 7029 (2017).
9. A. McDonnell, *et al.,* Efficacy of the $Na_V1.7$ blocker PF-05089771 in a randomised, placebo-controlled, double-blind clinical study in subjects with painful diabetic peripheral neuropathy. *Pain* **159**, 1465 (2018).
10. P. Siebenga, *et al.*, Lack of detection of the analgesic properties of PF-05089771, a selective $Na_V1.7$ inhibitor, using a battery of pain models in healthy subjects. *Clin. Transl. Sci.* **13**, 318 (2020).
11. M. E. Rothenberg, *et al.*, Safety, tolerability, and pharmacokinetics of GDC-0276, a novel $Na_V1.7$ inhibitor, in a first-in-human, single- and multiple-dose study in healthy volunteers. *Clin. Drug Investig.* **39**, 873 (2019).
12. B. S. Safina, *et al.*, Discovery of acyl-sulfonamide $Na_V1.7$ inhibitors GDC-0276 and GDC-0310. *J. Med. Chem.* **64**, 2953 (2021).
13. S. Noreng, T. Li, J. Payandeh, Structural pharmacology of voltage-gated sodium channels. *J. Mol. Bio.* **433**, 166967 (2021).
14. S. Ahuja, *et al.*, Structural basis of Nav1.7 inhibition by an isoform-selective small-molecule antagonist. *Science* **350**, aac5464 (2015).
15. S. Sun, *et al.*, Identification of selective acyl sulfonamide-cycloalkylether inhibitors of the voltage-gated sodium channel $(Na_V)1.7$ with potent analgesic activity. *J. Med. Chem.* **62**, 908 (2019).
16. M. Congreve, J. M. Dias, F. H. Marshall, Structure-based drug design for G protein-coupled receptors. *Prog. Med. Chem.* **53**, 1 (2014).

17. Y. Cheng, Membrane protein structural biology in the era of single particle cryo-EM. *Curr. Opin. Struct. Biol.* **52**, 58 (2018).

18. A. Balestrini, *et al.*, A TRPA1 inhibitor suppresses neurogenic inflammation and airway contraction for asthma treatment. *J. Exp. Med.* **218**, e20201637 (2021).

19. L. Rougé, *et al.*, Structure of CD20 in complex with the therapeutic monoclonal antibody rituximab. *Science* **367**, 1224 (2020).

20. H. Xu, *et al.*, Structural basis of $Na_V1.7$ inhibition by a gating-modifier spider toxin. *Cell* **176**, 702 (2019).

21. T. Clairfeuille, et al., Structural basis of α-scorpion toxin action on $Na_V$ channels. *Science* **363**, eaav8537 (2019).

22. H. Shen, D. Liu, K. Wu, J. Lei, N. Yan, Structures of human $Na_V 1.7$ channel in complex with auxiliary subunits and animal toxins. *Science* **363**, 1303 (2019).

23. G. Huang, *et al.*, High-resolution structures of human $Na_V1.7$ reveal gating modulation through α-π helical transition of $S6_{IV}$. *Cell Rep.* **39**, 110735 (2022).

24. H. Shen, *et al.*, Structure of a eukaryotic voltage-gated sodium channel at near-atomic resolution. *Science* **355**, eaal4326 (2017).

25. H. Shen, *et al.*, Structural basis for the modulation of voltage-gated sodium channels by animal toxins. *Science* **362**, eaau2596 (2018).

26. M. Weiss, *et al.*, Sulfonamides as selective $Na_V1.7$ inhibitors: optimizing potency and pharmacokinetics while mitigating metabolic liabilities, *J. Med. Chem.* **60**, 5969 (2017).

27. R. F. Graceffa, *et al.*, Sulfonamides as selective $Na_V1.7$ inhibitors: optimizing potency, pharmacokinetics, and metabolic properties to obtain atropisomeric quinolinone (AM-0466) that affords robust in vivo activity, *J. Med. Chem.* **60**, 5990 (2017).

28. S. J. McKerrall, *et al.*, Structure- and ligand-based discovery of chromane arylsulfonamide $Na_V1.7$ inhibitors for the treatment of chronic pain, *J. Med. Chem.* **62**, 4091 (2019).

29. A. J. Roecker, *et al.*, Discovery of arylsulfonamide $Na_V1.7$ inhibitors: IVIVC, MPO methods, and optimization of selectivity profile, *ACS Med. Chem. Lett.* **12**, 1038 (2021).

30. T. Focken, *et al.*, Discovery of aryl sulfonamides as isoform-selective inhibitors of $Na_V1.7$ with efficacy in rodent pain models, *ACS Med. Chem. Lett.* **7**, 277 (2016)

31. Y-J. W., *et al.*, Development of new benzenesulfonamides as potent and selective $Na_V1.7$ inhibitors for the treatment of pain, *J. Med. Chem.* **60**, 2513 (2017).

32. J. Payandeh, M. Volgraf, Ligand binding at the protein-lipid interface: strategic considerations for drug design. *Nat. Rev. Drug Discov.* **20**, 710 (2021).

33. G. Bankar, et al., Selective $Na_V1.7$ antagonists with long residence time show improved efficacy against inflammatory and neuropathic pain. *Cell Rep.* **24**, 3133 (2018).

34. G. Luo, *et al.*, Discovery of indole- and indazole-acylsulfonamides as potent and selective $Na_V1.7$ inhibitors for the treatment of pain, *J. Med. Chem.* **62**, 831 (2019).

35. E. F. DiMauro, *et al.*, Application of a parallel synthetic strategy in the discovery
of biaryl acyl sulfonamides as efficient and selective Na$_V$1.7 inhibitors, *J. Med.*
*Chem.* **59**, 7818 (2016).
36. T. Focken, *et al.*, Design of conformationally constrained acyl sulfonamide
isosteres: identification of N-([1,2,4]triazolo[4,3-a]pyridin-3-yl)methane-
sulfonamides as potent and selective hNa$_V$1.7 inhibitors for the treatment of pain,
*J. Med. Chem.* **61**, 4810 (2018).
37. X. Li, et al., Structural basis for modulation of human Na$_V$1.3 by clinical drug and
selective antagonist. *Nat. Commun.* **13**, 1286 (2022).
38. N. K. Kotla, K. A. Kumari, S. K. Gunda, M. Shaik, 3D-QSAR and molecular
docking studies of acylsulfonamides as Na$_V$1.7 inhibitors in chronic pain. *Res. J.*
*Life Sci. Bioinform. Pharm. Chem. Sci.* **5**, 120 (2019).
39. R. P. Mason, D. G. Rhodes, L. G. Herbette, Reevaluating equilibrium and kinetic
binding parameters for lipophilic drugs based on a structural model for drug
interactions with biological membranes. *J. Med. Chem.* **34**, 869 (1991).
40. R. A. Coleman, M. Johnson, A. T. Nials, C. J. Vardey, Exosites: their current
status, and their relevance to the duration of action of long- acting beta 2-
adrenoceptor agonists. *Trends Pharmacol. Sci.* **17**, 324 (1996).
41. G. P. Anderson, A. Lindén, K. F. Rabe, Why are long-acting beta-adrenoceptor
agonists long-acting? *Eur. Respir. J.* **7**, 569 (1994).
42. M. Masureel, *et al.*, Structural insights into binding specificity, efficacy and bias
of a β2 AR partial agonist. *Nat. Chem. Biol.* **14**, 1059 (2018).
43. R. P. Austin, *et al.*, QSAR and the rational design of long- acting dual D2-
receptor/beta 2-adrenoceptor agonists. *J. Med. Chem.* **46**, 3210 (2003).
44. D. G. Rhodes, R. Newton, R. Butler, L. Herbette, Equilibrium and kinetic studies
of the interactions of salmeterol with membrane bilayers. *Mol. Pharmacol.* **42**,
596 (1992).
45. D. A. Sykes, *et al.*, Observed drug- receptor association rates are governed by
membrane affinity: the importance of establishing "micro- pharmacokinetic/
pharmacodynamic relationships" at the β2-adrenoceptor. *Mol. Pharmacol.* **85**, 608
(2014).
46. C. J. Dickson, *et al.*, Uncoupling the structure−activity relationships of β2
adrenergic receptor ligands from membrane binding. *J. Med. Chem.* **59**, 5780
(2016).
47. Z. Zhao, *et al.*, Exploration of type II binding mode: a privileged approach for
kinase inhibitor focused drug discovery? *ACS Chem. Biol.* **9**, 1230 (2014).
48. D. N. Mastronarde, Automated electron microscope tomography using robust
prediction of specimen movements. *J. Struct. Biol.* **152**, 36 (2005).
49. S. H. W. Scheres, RELION: implementation of a Bayesian approach to cryo-EM
structure determination. *J. Struct. Biol.* **180**, 519 (2012).
50. T. Grant, A. Rohou, N. Grigorieff, *cis*TEM, user-friendly software for single-
particle image processing. *Elife* **7**, e35383 (2018).
51. M. Kschonsak, *et al.* Structural architecture of the human NALCN channelosome.
*Nature* **603**, 180 (2022).
52. S. Q. Zheng *et al.* MotionCor2: anisotropic correction of beam-induced motion
for improved cryo-electron microscopy. *Nature Methods* **14**, 331 (2017).

53. A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from
electron micrographs. *J. Struct. Biol.* **192**, 216 (2015).
54. T. C. Terwilliger, S. J. Ludtke, R. J. Read, P. D. Adams, P. V. Afonine,
Improvement of cryo-EM maps by density modification. *Nature Methods* **17**, 923
(2020).
55. P. V. Afonine, *et al*. Real-space refinement in PHENIX for cryo-EM and
crystallography. *Acta Crystallogr. D. Struct. Biol.* **74**, 531 (2018).
56. N. W. Moriarty, R. W. Grosse-Kunstleve, P. D. Adams, electronic Ligand Builder
and Optimization Workbench (eLBOW): a tool for ligand coordinate and restraint
generation. *Acta Crystallogr. D. Biol. Crystallogr.* **65**, 1074 (2009).
57. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of
Coot. *Acta Crystallogr. D. Biol. Crystallogr.* **66**, 486 (2010).
58. P. V. Afonine, *et al*. New tools for the analysis and validation of cryo-EM maps
and atomic models. *Acta Crystallogr. D Struct. Biol.* **74**, 814 (2018).
59. C. J. Williams, *et al*. MolProbity: More and better reference data for improved all-
atom structure validation. *Protein Sci.* **27**, 293 (2018).
60. E. F. Pettersen, *et al*. UCSF Chimera--a visualization system for exploratory
research and analysis. *J. Comput. Chem.* **25**, 1605 (2004).
61. T. D. Goddard, *et al*. UCSF ChimeraX: Meeting modern challenges in
visualization and analysis. *Protein Sci.* **27**, 14 (2018).
62. D. Sutherlin *et al*. Therapeutic compounds and methods of use thereof.
US11028075B2, June 8, 2021.

743
744
745
746

1 **Supporting Information**
2
3 **CryoEM reveals unprecedented binding site for Na$_V$1.7 inhibitors enabling rational design of potent hybrid**
4 **inhibitors**
5

6 Marc Kschonsak[1*], Christine C. Jao[1*], Christopher P. Arthur[1], Alexis L. Rohou[1], Philippe Bergeron[2], Daniel
7 Ortwine[2], Steven J. McKerrall[2], David H. Hackos[3], Lunbin Deng[3], Jun Chen[4], Peter S. Dragovich[2], Matthew
8 Volgraf[2], Matthew R. Wright[5], Jian Payandeh[1,#], Claudio Ciferri[1,#], John C. Tellis[2,#]
9
10 [1]Genentech Inc., Structural Biology; South San Francisco, CA, United States.
11 [2]Genentech Inc., Discovery Chemistry; South San Francisco, CA, United States.
12 [3]Genentech Inc., Neuroscience; South San Francisco, CA, United States.
13 [4]Genentech Inc., Biochemical and Cellular Pharmacology; South San Francisco, CA, United States.
14 [5]Genentech Inc., Drug Metabolism and Pharmacokinetics; South San Francisco, CA, United States.
15
16
17 [#]Corresponding authors. Email: jpayandeh@exelixis.com (J.P); ciferri.claudio@gene.com (C.C.);
18 tellis.john@gene.com (J.C.T.).

41    **Table S1. Cryo-EM data collection, refinement and validation statistics.**

| | **GNE-3565** VSD4-Na$_V$1.7-Na$_V$Pas complex | **GDC-0310** VSD4-Na$_V$1.7-Na$_V$Pas complex | **GNE-1305** VSD4-Na$_V$1.7-Na$_V$Pas complex | **GNE-9296** VSD4-Na$_V$1.7-Na$_V$Pas-DC1a complex |
|---|---|---|---|---|
| **Data Collection** | | | | |
| Magnification | 105,000x | 105,000x | 165,000x | 165,000x |
| Voltage (kV) | 300 | 300 | 300 | 300 |
| Electron exposure (e/Å$^2$) | 64 | 64 | 64 | 48 |
| Defocus range (μm) | 0.5-1.5 | 0.5-1.5 | 0.5-1.5 | 0.5-1.5 |
| Pixel size (Å) | 0.838 | 0.838 | 0.838 | 0.824 |
| Symmetry imposed | C1 | C1 | C1 | C1 |
| Initial particle images | 3,686,837 | 2,253.983 | 2,741,402 | 1,774,062 |
| Final particle images | 964,572 | 795,792 | 1,201,168 | 826,525 |
| Map resolution (Å) overall | 2.9 | 2.5 | 2.2 | 3.1 |
|     FSC threshold | 0.143 | 0.143 | 0.143 | 0.143 |
| Map resolution range (Å) | 2.7-33.5 | 2.3-33.5 | 2.1-33.5 | 3.3-33.0 |
| **Refinement** | | | | |
| Initial models used (PDB code) | 6NT3 | 6NT3 | 6NT3 | 6NT3 |
| Model resolution (Å) | 2.9 | 1.9 | 1.7 | 2.5 |
|     FSC threshold | 0.5 | 0.5 | 0.5 | 0.5 |
| Model composition | | | | |
|     Non-hydrogen atoms | 9396 | 9557 | 9616 | 9734 |
|     Protein residues | 1115 | 1115 | 1117 | 1159 |
|     Waters | 0 | 78 | 111 | 0 |
|     Ligands | 13 | 13 | 14 | 13 |
| B factors (Å$^2$) | | | | |
|     Protein | 25.62 | 12.42 | 6.27 | 24.05 |
|     Ligand | 35.81 | 15.79 | 10.33 | 40.62 |
|     Water | N/A | 8.11 | 6.54 | N/A |
| R.m.s. deviations | | | | |
|     Bond lengths (Å) | 0.004 | 0.005 | 0.005 | 0.005 |
|     Bond angles (°) | 0.931 | 1.018 | 0.994 | 0.941 |
| Validation | | | | |
|     MolProbity score | 1.30 | 1.19 | 1.14 | 1.38 |
|     Clashscore | 5.47 | 4.09 | 3.51 | 6.42 |
|     Poor rotamers (%) | 0.10 | 0.20 | 0.61 | 0.00 |
| Ramachandran Plot | | | | |
|     Favored (%) | 98.45 | 98.27 | 98.19 | 97.90 |
|     Allowed (%) | 1.55 | 1.73 | 1.81 | 2.10 |
|     Disallowed (%) | 0.00 | 0.00 | 0.00 | 0.00 |
| | | | | |

GX-936

**Supplementary Figure 1:** Structure of arylsulfonamide inhibitor **GX-936**

**Chemical compound synthesis and characterization**

**General Considerations**

All commercial solvents and reagents were used without additional purification unless indicated otherwise. [1]H NMR spectra were measured on Bruker Avance III 300, 400, or 500 MHz spectrometers. Chemical shifts (in ppm) were referenced to tetramethylsilane as an internal standard ($\delta = 0$ ppm). Reaction progress was monitored by either a Shimadzu LCMS/UV system with an LC-30 AD solvent pump, Sil-30 AC autosampler, 2020 MS, SPDM30A UV detector, and CTO-20A column oven, using 2−98% acetonitrile/0.1% formic acid (or 0.01% ammonia) over 2.5 min or a Waters Acquity LCMS system using 2−98% acetonitrile/0.1% formic acid (or 0.1% ammonia) over 2 min. Flash column chromatography purifications were performed using a Teledyne Isco Combiflash Rf and Silicycle HP columns. Reverse-phase purification was done on a Phenomenex Gemini-NX C18 (30 × 100 mm, 5 μm) with a gradient of 5−95% acetonitrile/water (with 0.1% NH4OH or 0.1% formic acid) at 60 mL/min over 10 min. Preparative SFC separations were carried out on a PIC Solutions instrument, with conditions specified in the Experimental Section. High-resolution mass spectrometry (HRMS) of final compounds was obtained on a Thermo UHPLC/ QE with a Thermo-Q Exactive mass spectrometry detector using ESI ionization, following elution on an Acquity BEH C18 stationary phase (2.1 mm × 50 mm; 1.7 μm particle size) using a gradient of water/ acetonitrile (3−97% over 7 min with 0.1% formic acid in both phases). Unless stated otherwise, analytical purity was >95% as determined by LCMS using UV 254 nm detection

**Compound 2**

**5-chloro-4-(cyclopentylmethoxy)-*N*-((4-(((1*S*,2*S*)-2-(dimethylamino)cyclohexyl)amino)-2-fluorophenyl)sulfonyl)-2-fluorobenzamide**

**Step 1:**

**4-(((1*S*,2*S*)-2-(dimethylamino)cyclohexyl)amino)-2-fluorobenzenesulfonamide**

A solution of (1*S*,2*S*)-*N*[1],*N*[1]-dimethylcyclohexane-1,2-diamine (162 mg, 1.1 mmol), 2,4-difluorobenzenesulfonamide (200 mg, 1.0 mmol) and DIPEA (0.3 mL, 1.7 mmol) in DMSO (3 mL) was stirred at

80 °C for 16 h. After cooling to room temperature, the reaction was diluted with EtOAc (100 mL), and washed
with brine (50 mL x 5). The organic lawyer was dried over anhydrous $Na_2SO_4$, filtered and concentrated in vacuo.
The crude residue was purified by silica gel chromatography (solvent gradient: 0 - 10% EtOAc in petroleum
ether) to afford the title compound (60 mg, 0.19 mmol) as yellow oil.
LCMS (ESI) m/z: 316.1 [M+H]$^+$.

**Step 2:**

**tert-butyl 5-chloro-4-(cyclopentylmethoxy)-2-fluorobenzoate**
A mixture of tert-butyl 5-chloro-2,4-difluorobenzoate (548 g, 2.20 mol), cyclopentylmethanol (200 g, 2.00 mol)
and $Cs_2CO_3$ (1.31 kg, 4.00 mol) in DMSO (3.28 L) was stirred at 80 °C for 5 hrs. The mixture was filtered and
filtrate was diluted with EtOAc (3.00 L), washed with water (3.00 L x 3) and brine (1000 mL). The resultant
organic layer was dried over $Na_2SO_4$, filtered, concentrated in vacuo. The residue was purified by column
chromatography (SiO$_2$, Petroleum ether/Ethyl acetate = 100/1 to 50/1). The title compound (0.50 kg, crude) was
obtained as light yellow oil.

**Step 3:**

**5-chloro-4-(cyclopentylmethoxy)-2-fluorobenzoic acid**
To a solution of *tert*-butyl 5-chloro-4-(cyclopentylmethoxy)-2-fluorobenzoate (327 g, 994 mmol) in DCM (654
mL) was added TFA (654 mL) at 25 °C. The mixture was stirred at 25 °C for 24 hrs. The reaction mixture was
filtered and concentrated under reduced pressure to give a residue that was triturated with 1/1 MTBE/Petroleum
ether at 25 °C for 30 mins. The title compound was obtained by filtration as a white solid (83.7 g, 30.5% yield).
$^1$H NMR (400 MHz, CDCl$_3$) $\delta$ 8.04 (d, $J$ = 7.6 Hz, 1H), 6.70 (d, $J$ = 12.4 Hz, 1H), 3.96 (d, $J$ = 6.8 Hz, 2H), 2.42-
2.50 (m 1H), 1.86-1.93 (m, 2H), 1.75-1.58 (m, 4H), 1.48-1.35 (m, 2H).

**Step 4:**

**5-chloro-4-(cyclopentylmethoxy)-*N*-((4-(((1*S*,2*S*)-2-(dimethylamino)cyclohexyl)amino)-2-**
**fluorophenyl)sulfonyl)-2-fluorobenzamide**
To a solution of 4-(((1*S*,2*S*)-2-(dimethylamino)cyclohexyl)amino)-2-fluorobenzenesulfonamide (80 mg, 0.24
mmol) and DMAP (59 mg, 0.48 mmol) in DCM (2 mL) was added EDCI (51 mg, 0.26 mmol) and 5-chloro-4-
(cyclopentylmethoxy)-2-fluoro-benzoic acid (72 mg, 0.26 mmol). The reaction was stirred at room temperature
for 2 h. The reaction was quenched with 10% aqueous citric acid (5 mL). The reaction was diluted with water (10
mL) and extracted with DCM (10 mL x 3). The combined organic layers were dried over anhydrous $Na_2SO_4$,
filtered and concentrated in vacuo. The crude residue was purified by silica gel chromatography (solvent gradient:
0 - 5% MeOH in DCM) to afford the title compound (40 mg, 28%) as a white solid.

$^1$H NMR (400 MHz, DMSO-$d_6$) δ 7.75 (d, $J$ = 7.6 Hz, 1H), 7.55 - 7.48 (m, 1H), 6.94 (d, $J$ = 12.4 Hz, 1H), 6.50 (s,

1H), 6.47 (s, 1H), 6.07 (d, $J$ = 10.0 Hz, 1H), 3.94 (d, $J$ = 6.8 Hz, 2H), 3.79 - 3.64 (m, 1H), 3.15 - 3.05 (m, 1H),

2.66 (s, 6H), 2.35 - 2.25 (m, 1H), 2.09 - 1.94 (m, 2H), 1.84 - 1.69 (m, 3H), 1.63 - 1.14 (m, 11H).

HRMS $m/z$ calcd for $C_{27}H_{34}ClF_2N_3O_4S$ [M+H]$^+$: 570.1999. Found 570.2003.

**Compound 3**

**4-(cyclopentylmethoxy)-5-cyclopropyl-N-((4-(((1S,2S)-2-(dimethylamino)cyclohexyl)amino)-2-**

**fluorophenyl)sulfonyl)-2-fluorobenzamide**

Following the procedure described in compound **3** and replacing 5-chloro-4-(cyclopentylmethoxy)-2-

fluorobenzoic acid with 4-(cyclopentylmethoxy)-5-cyclopropyl-2-fluorobenzoic acid, the title compound was

obtained as a white solid.

$^1$H NMR (400 MHz, DMSO-$d_6$) δ 7.50 (t, $J$ = 8.8 Hz, 1H), 7.20 (d, $J$ = 8.8 Hz, 1H), 6.65 (d, $J$ = 12.0 Hz, 1H),

6.52 - 6.34 (m, 2H), 5.99 (s, 1H), 3.87 (d, $J$ = 6.8 Hz, 2H), 3.72 - 3.52 (m, 1H), 3.05 - 2.81 (m, 1H), 2.61 - 2.52

(m, 6H), 2.37 - 2.26 (m, 1H), 2.07 - 1.90 (m, 3H), 1.87 - 1.71 (m, 3H), 1.66 - 1.52 (m, 5H), 1.45 - 1.30 (m, 4H),

1.29 - 1.20 (m, 1H), 1.15 - 1.05 (m, 1H), 0.91 - 0.80 (m, 2H), 0.60 - 0.45 (m, 2H).

HRMS $m/z$ calcd for $C_{30}H_{39}F_2N_3O_4S$ [M+H]$^+$: 576.2702. Found 576.2713.

**Compound 4**

**N-(6-(cyclopentylmethoxy)benzo[d]thiazol-2-yl)-4-(((1S,2S)-2-(dimethylamino)cyclohexyl)amino)-2-**

**fluorobenzenesulfonamide**

**Step 1:**

**4-(cyclopentylmethoxy)aniline**

To a stirred solution of NaH (1.32 g, 54.98 mmol, 60% in mineral oil) in DMF (20 mL) was added 4-aminophenol

(2 g, 18.33 mmol) at 0 $^o$C under nitrogen atmosphere. After stirring at 0 $^o$C for 10 min,

(bromomethyl)cyclopentane (4.48 g, 27.49 mmol) was added at 0 $^o$C. The reaction was stirred at room

temperature for 16 h. The reaction was quenched with water (100 mL) and extracted with EtOAc (100 mL x 3).

The combined organic lawyers were washed with brine (100 mL), dried over anhydrous $Na_2SO_4$, filtered and

concentrated in vacuo. The crude residue was purified by silica gel chromatography (solvent gradient: 10 - 20%

EtOAc in petroleum ether) to afford the title compound (1.35 g, 39%) as black oil.

$^1$H NMR (400 MHz, CDCl$_3$) δ 6.77 - 6.74 (m, 2H), 6.66 - 6.63 (m, 2H), 3.76 (d, $J$ = 7.2 Hz, 2H), 3.42 (s, 2H),

2.38 - 2.27 (m, 1H), 1.86 - 1.78 (m, 2H), 1.64 - 1.58 (m, 4H), 1.39 - 1.30 (m, 2H).

LCMS (ESI) m/z: 192.2 [M+H]$^+$.

**Step 2:**

**6-(cyclopentylmethoxy)benzo[*d*]thiazol-2-amine**

The solution of 4-(cyclopentylmethoxy)aniline (1.30 g, 7.06 mmol) and potassium thiocyanate (685 mg, 7.06
mmol) in acetic acid (7.5 mL) was stirred at 0 °C for 20 min. Bromine (0.36 mL, 7.06 mmol) in acetic acid (3.5
mL) was added slowly and keep the temperature below 10 °C. Then, the mixture was stirred at room temperature
for 18 h. The reaction was filtered and the filter cake was washed with acetic acid (5 mL). The filtrate was
concentrated in vacuo and the crude reside was diluted with hot water (5 mL) and basified to pH>11 with
NH$_3$•H$_2$O. The resulting precipitate was filtered and the filter cake was washed with water (5 mL). The filter cake
was diluted with DCM (20 mL), dried over anhydrous Na$_2$SO$_4$, filtered and concentrated in vacuo. The crude
residue was purified by silica gel chromatography (solvent gradient: 0 - 14% EtOAc in petroleum ether) to afford
the title compound (800 mg, 46%) as a gray solid.
$^1$H NMR (400 MHz, CDCl$_3$) δ 7.43 (d, *J* = 8.8 Hz, 1H), 7.13 (d, *J* = 6.0 Hz, 1H), 6.93 - 6.90 (m, 1H), 5.22 (s, 2H),
3.84 (d, *J* = 7.2 Hz, 2H), 2.41 - 2.33 (m, 1H), 1.89 - 1.81 (m, 2H), 1.68 - 1.58 (m, 4H), 1.41 - 1.33 (m, 2H).
LCMS (ESI) m/z: 249.0 [M+H]$^+$.

**Step 3:**

**6-(cyclopentylmethoxy)-N-(2,4-dimethoxybenzyl)benzo[d]thiazol-2-amine**

To a stirred solution of 6-(cyclopentylmethoxy)benzo[d]thiazol-2-amine (0.71 g, 2.65 mmol) and 2,4-
dimethoxybenzaldehyde (0.4 g, 2.41 mmol) in DCM (12 mL) was added TiCl(O*i*-Pr)$_3$ (1.86 mL, 5.54 mmol) in
one portion under nitrogen atmosphere. The solution was stirred for 10 min before the portion wise addition of
NaBH(OAc)$_3$ (1.53 g, 7.22 mmol) at 0 °C. The reaction was stirred at room temperature for 16 h. The reaction
was quenched with saturated aqueous NaHCO$_3$ solution (50 mL), extracted with DCM (50 mL x 3). The
combined organic lawyers were washed with brine (50 mL), dried over anhydrous Na$_2$SO$_4$, filtered and
concentrated in vacuo. The crude residue was purified by silica gel chromatography (solvent gradient: 0 - 25%
EtOAc in petroleum ether) to afford the title compound (0.53 g, 53%) as a white solid.

**Step 4:**

**N-(6-(cyclopentylmethoxy)benzo[d]thiazol-2-yl)-N-(2,4-dimethoxybenzyl)-2,4-difluorobenzenesulfonamide**

To a solution of 6-(cyclopentylmethoxy)-N-(2,4-dimethoxybenzyl)benzo[d]thiazol-2-amine (170 mg, 0.52 mmol)
in THF (2 mL) was added LiHMDS (0.62 mL, 0.62 mmol, 1 M) at -78 °C. The reaction was stirred for 30 min at
0 °C and a solution of 2,4-difluorobenzenesulfonylchloride (0.22 g, 1.03 mmol) in THF (2 mL) was added
dropwise at -78 °C. After the addition was complete, the cooling bath was removed. The reaction mixture was
stirred at room temperature for 3 h. The reaction was diluted with water (30 mL) and extracted with EtOAc (50
mL x 3). The combined organic lawyers were washed with brine (50 mL), dried over anhydrous Na$_2$SO$_4$, filtered
and concentrated in vacuo. The crude residue was purified by silica gel chromatography (solvent gradient: 0 -
30% EtOAc in petroleum ether) to afford the title compound (160 mg, 52%) as a white solid.

**Step 5:**

**N-(6-(cyclopentylmethoxy)benzo[d]thiazol-2-yl)-N-(2,4-dimethoxybenzyl)-4-(((1S,2S)-2-**

**(dimethylamino)cyclohexyl)amino)-2-fluorobenzenesulfonamide**

To a solution of *N*-(6-bromothiazolo[4,5-*b*]pyridin-2-yl)-4-(((1*S*,2*S*)-2-(dimethylamino)cyclohexyl)amino)-2-

fluorobenzenesulfonamide (40 mg, 0.07 mmol) in DMSO (1 mL) was added DIPEA (14 mg, 0.11 mmol) and

(1*S*,2*S*)-$N_1$,$N_1$-dimethylcyclohexane-1,2-diamine (15 mg, 0.11 mmol) at room temperature. The reaction mixture

was stirred at room temperature for 20 h. The reaction was quenched with saturated aqueous $NH_4Cl$ (20 mL),

extracted with EtOAc (30 mL x 3). The combined organic layers were washed with brine (30 mL), dried over

anhydrous $Na_2SO_4$, filtered and concentrated in vacuo to afford the title compound (48 mg, crude) as yellow oil

that required no further purification.

**Step 6:**

**N-(6-(cyclopentylmethoxy)benzo[d]thiazol-2-yl)-4-(((1S,2S)-2-(dimethylamino)cyclohexyl)amino)-2-**

**fluorobenzenesulfonamide**

A solution of *N*-(6-(cyclopentylmethoxy)benzo[d]thiazol-2-yl)-*N*-(2,4-dimethoxybenzyl)-4-(((1*S*,2*S*)-2-

(dimethylamino)cyclohexyl)amino)-2-fluorobenzenesulfonamide (60 mg, 0.08 mmol) in HCOOH (3 mL) was

stirred at room temperature for 16 h. The mixture was concentrated in vacuo and the crude residue was purified by

reverse phase chromatography (acetonitrile 35 - 65% / 0.2% HCOOH in water) to afford the title compound (2

mg, 4%) as a white solid

[1]H NMR (400 MHz, DMSO-$d_6$) δ 7.49 (t, *J* = 8.8 Hz, 1H), 7.26 (d, *J* = 2.4 Hz, 1H), 7.16 (d, *J* = 8.8 Hz, 1H), 6.84

- 6.81 (m, 1H), 6.42 - 6.36 (m, 3H), 3.79 (d, *J* = 6.8 Hz, 2H), 3.47 - 3.40 (m, 1H), 2.82 - 2.75 (m, 1H), 2.42 (s,

6H), 2.33 - 2.24 (m, 1H), 2.02 - 1.92 (m, 2H), 1.79 - 1.71 (m, 3H), 1.61 - 1.48 (m, 5H), 1.34 - 1.20 (m, 6H).

HRMS (ESI) m/z: 547.2196 [M+H]+

**Compound 5**

**4-(((1S,2S)-2-(dimethylamino)cyclohexyl)amino)-2-fluoro-N-(6-(trifluoromethoxy)benzo[d]thiazol-2-**

**yl)benzenesulfonamide**

Following the procedure described in compound **6** and replacing 4-(cyclopentylmethoxy)aniline with 4-

(trifluoromethoxy)aniline, the title compound was obtained as a white solid.

[1]H NMR (400 MHz, DMSO-$d_6$) δ 7.64 (m, 1H), 7.52 (m, 1H), 7.30 (d, *J* = 8.8 Hz, 1H), 7.11 (m, 1H), 6.44-6.37

(m, 3 H), 3.68 (m, 1H), 3.06 (m, 1H), 2.60 (s, 6 H), 2.00 (m, 2H), 1.79 (m, 1H), 1.61 (m, 1H), 1.40- 1.14 (m, 4 H).

HRMS (ESI) m/z: 533.1286 [M+H]+.

**Supplementary Figure Legends:**

**Supplementary Figure 2: (A)** VSD4-Na$_V$1.7-Na$_V$Pas channel protein expression and purification scheme **(B)** Example size exclusion chromatogram and SDS-PAGE of nanodisc-reconstituted VSD4-Na$_V$1.7-Na$_V$Pas channel sample.

**Supplementary Figure 3: (A)** Example cryo-EM micrograph image of the VSD4-Na$_V$1.7-Na$_V$Pas channel. **(B)** Representative 2D-class averages of selected particles. **(C)** Data collection and processing workflow for VSD4-Na$_V$1.7-Na$_V$Pas channel bound to compound **4**. **(D)** Heat map representation of the distribution of assigned particle orientations. **(E)** FSC between two half datasets yields a global resolution estimate of approximately 2.2 Å resolution from the refinement using an overall mask of the VSD4-Na$_V$1.7-Na$_V$Pas channel bound to compound **4 (F)** Local resolution of maps. **(G)** Data collection and processing workflow for VSD4-Na$_V$1.7-Na$_V$Pas channel bound to **GDC-0310**. **(H)** Heat map representation of the distribution of assigned particle orientations. **(I)** FSC between two half datasets yields a global resolution estimate of approximately 2.5 Å resolution from the refinement using an overall mask of the VSD4-Na$_V$1.7-Na$_V$Pas channel bound to **GDC-0310**. **(J)** Local resolution of maps.

**Supplementary Figure 4: (A)** Data collection and processing workflow for VSD4-Na$_V$1.7-Na$_V$Pas channel bound to compound **2**. **(B)** Heat map representation of the distribution of assigned particle orientations. **(C)** FSC between two half datasets yields a global resolution estimate of approximately 2.9 Å resolution from the refinement using an overall mask of the VSD4-Na$_V$1.7-Na$_V$Pas channel bound to compound **2 (D)** Local resolution of maps. **(E)** Data collection and processing workflow for VSD4-Na$_V$1.7-Na$_V$Pas channel bound to compound **2**. **(F)** Heat map representation of the distribution of assigned particle orientations. **(G)** FSC between two half datasets yields a global resolution estimate of approximately 2.5 Å resolution from the refinement using an overall mask of the VSD4-Na$_V$1.7-Na$_V$Pas channel bound to compound **2**. **(H)** Local resolution of maps.

**Supplementary Figure 5: (A)** The cryoEM map surrounding the ligand compound **2** is shown in mesh representation. **(B)** The cryoEM map surrounding the ligand compound **4** is shown in mesh representation.

260        HPLC chromatogram of compound **2**

Peak Table

PDA Ch1 220nm

| Peak# | Ret. Time | Height | Height% | USP Width | Area | Area% |
|---|---|---|---|---|---|---|
| 1 | 1.058 | 2105 | 0.384 | 0.071 | 6000 | 0.342 |
| 2 | 4.129 | 546761 | 99.616 | 0.084 | 1750524 | 99.658 |

Chromatogram

mAU

1 PDA Multi 2 254nm,4nm

1 PDA Multi 2 / 254nm,4nm

Peak Table

PDA Ch2 254nm

| Peak# | Ret. Time | Height | Height% | USP Width | Area | Area% |
|---|---|---|---|---|---|---|
| 1 | 4.129 | 484289 | 100.000 | 0.084 | 1542826 | 100.000 |

Chromatogram

mAU

1 PDA Multi 3 215nm,4nm

1 PDA Multi 3 / 215nm,4nm

Peak Table

PDA Ch3 215nm

| Peak# | Ret. Time | Height | Height% | USP Width | Area | Area% |
|---|---|---|---|---|---|---|
| 1 | 1.059 | 3033 | 0.461 | 0.071 | 8450 | 0.401 |
| 2 | 4.129 | 655545 | 99.539 | 0.084 | 2099153 | 99.599 |

263
264
265

266
267      HPLC chromatogram of compound **3**

**1: 220 nm, 8 nm**

| Retention Time | Height | Area | Area Percent |
|---|---|---|---|
| 0.88 | 2006 | 14264 | 0.23 |
| 2.71 | 17825 | 94685 | 1.52 |
| 4.62 | 1574 | 5119 | 0.08 |
| 4.79 | 4288 | 13870 | 0.22 |
| 5.23 | 1156786 | 6087427 | 97.94 |

**2: 215 nm, 8 nm**

| Retention Time | Height | Area | Area Percent |
|---|---|---|---|
| 0.55 | 3225 | 12600 | 0.18 |
| 0.87 | 3232 | 23433 | 0.33 |
| 2.71 | 29397 | 156299 | 2.23 |
| 4.62 | 2484 | 7826 | 0.11 |
| 4.79 | 4768 | 15348 | 0.22 |
| 5.23 | 1320766 | 6789788 | 96.69 |
| 5.65 | 1640 | 5918 | 0.08 |
| 5.97 | 1527 | 5675 | 0.08 |
| 7.06 | 1670 | 5605 | 0.08 |

**3: 254 nm, 8 nm**

| Retention Time | Height | Area | Area Percent |
|---|---|---|---|
| 5.23 | 897527 | 4692528 | 100.00 |

270
271
272
273

274
275     HPLC chromatogram of compound **4**

**1: 220 nm, 8 nm**

| Retention Time | Height | Area | Area Percent |
|---|---|---|---|
| 0.88 | 5778 | 48393 | 0.91 |
| 4.70 | 1288689 | 5232399 | 98.25 |
| 5.67 | 1644 | 5558 | 0.10 |
| 6.01 | 10759 | 39356 | 0.74 |

**2: 215 nm, 8 nm**

| Retention Time | Height | Area | Area Percent |
|---|---|---|---|
| 0.56 | 2946 | 9287 | 0.14 |
| 0.88 | 9264 | 78075 | 1.22 |
| 4.70 | 1604066 | 6253624 | 97.59 |
| 5.10 | 1747 | 8500 | 0.13 |
| 5.67 | 1685 | 5554 | 0.09 |
| 6.01 | 11358 | 52842 | 0.82 |

**3: 254 nm, 8 nm**

| Retention Time | Height | Area | Area Percent |
|---|---|---|---|
| 4.70 | 665091 | 2626651 | 99.73 |
| 6.01 | 1968 | 7007 | 0.27 |

278
279
280

281      HPLC chromatogram of compound **5**

**1: 220 nm, 8 nm**

| Retention Time | Height | Area | Area Percent |
|---|---|---|---|
| 0.83 | 5245 | 47427 | 1.85 |
| 1.08 | 2110 | 6804 | 0.27 |
| 4.00 | 753425 | 2510118 | 97.89 |

**2: 215 nm, 8 nm**

| Retention Time | Height | Area | Area Percent |
|---|---|---|---|
| 0.83 | 8794 | 83696 | 2.90 |
| 1.08 | 3010 | 9655 | 0.33 |
| 4.00 | 842279 | 2784306 | 96.38 |
| 5.04 | 1857 | 11275 | 0.39 |

**3: 254 nm, 8 nm**

| Retention Time | Height | Area | Area Percent |
|---|---|---|---|
| 4.00 | 393524 | 1296194 | 100.00 |

284
285
286

# Figure 1

**A**

change color of linker and of S5

**B**

**C** Na$_V$Pas-Na$_V$1.7 VSD4

top view

**D**

front view

**E**

Water molecules

**F** Selectivity filter

overlay density

**G** Activation gate

overlay density

# Figure 2

**A**
should we highlight the warhead?

G'3565

**B** VSD4

top view

**C**

side view

**D** extracellular view

S2    S1

E1524

W1538    Y1537    N1540

R2

R3

A1585

S3    S4

**E**

contour 0.5
(need to flip)

**F** view towards membrane

S3    S2

A1585    Y1537

**need to flip**    3.6

3.5

3.6    N1540

R4

**G** view from membrane

S2    Y1537

W1538    D1586    S3

3.1

F1583

# Figure 3

should we highlight the warhead?

**A**

G'1310

**B**

VSD4

top view

**C**

side view

**D**

extracellular view

S2    S1

E1524

W1538

N1540

Y1537    R3

A1585

R2

S3

S4

**E**

contour 1.5

**F**

view towards membrane

S3    S2

A1585    Y1537

3.1

3.2    N1540

2.7

R4

**G**

S3    S4

I1588    Ixxxx

I1574    Axxxx

# Figure 4

# Supplementary Figure 1

**A**

Cell lysis

↓

Detergent solubilization

↓

Dual affinity purification

↓

Nanodisc reconstitution

↓

SEC

↓

Compound incubation

↓

CryoEM

**B**

Volume

NavPas

MSP

# Supplementary Figure 2

**A**

scale 10 nm

**B**

scale 10 nm

**C**

Na$_V$Pas G03451305

12,339 movies collected from 1 grid → motion correction / CTF fit resolution 6.0 Å or better → 12,299 images → template-matching / particle picking (gautomatch) → 2,741,402 particle coordinates → 2D classification / select 18 of 100 classes → 1,201,168 particles | RELION

export particle stack from RELION and import to cisTEM | cisTEM

2.2 Å map ← man. refine (masked) hi.res limit: 3.0 Å / low-pass filter outside channel, CTF refinement ← 2.4 Å map ← auto-refine (auto-masked) / initial res. limit: 8.0 Å

phenix density modification

**D**

φ=90° φ=135° θ=60° φ=45° θ=30° φ=180° φ=0° φ=225° φ=315° φ=270°

176+ / 0

**E**

FSC vs Spatial Frequency (1/Å), ref. limit, FSC= 0.143

**F**

resolution (Å) 2.0 3.0 4.0 5.0 >6.0

**G**

Na$_V$Pas G02851310

7803 movies collected from 1 grid → motion correction / CTF fit resolution 6.0 Å or better → 7786 images → template-matching / particle picking (gautomatch) → 2,253,983 particle coordinates → 2D classification / select 24 of 100 classes → 795,792 particles | RELION

export particle stack from RELION and import to cisTEM | cisTEM

2.5 Å map ← man. refine (masked) hi.res limit: 3.4 Å / low-pass filter outside channel, CTF refinement ← 3.1 Å map ← auto-refine (auto-masked) / initial res. limit: 8.0 Å

phenix density modification

**H**

φ=90° φ=135° θ=60° φ=45° θ=30° φ=180° φ=0° φ=225° φ=315° φ=270°

117+ / 0

**I**

FSC vs Spatial Frequency (1/Å), ref. limit, FSC= 0.143

**J**

resolution (Å) 2.0 3.0 4.0 5.0 >6.0

# Supplementary Figure 3

**A**                                                          **GNE-3565**

**B**

**C**

**D**

resolution (Å)

2.0  3.0  4.0  5.0  >6.0

**E**                                                          **GNE-9296**

**F**

**G**

**H**

resolution (Å)

2.0  3.0  4.0  5.0  >6.0

Supplementary Figure 4

**A**

G'9296

**B**

G'1305