Title: Intranasal self-amplifying RNA SARS-CoV-2 vaccine produces protective

respiratory and systemic immunity and prevents viral transmission

Authors: Madeleine F. Jennewein[1†], Michael D. Schultz[2†], Samuel Beaver[1], Peter Battisti[1], Julie

Bakken[1], Derek Hanson[1], Jobaida Akther[3], Raodoh Mohamath[1], Jasneet Singh[1], Noah Cross[1],

Sierra Reed[1], Davies Kalange[3], Jeremy B. Foote[2], R. Glenn King[2], Aaron Silva-Sanchez[3],

Davide Botta[2], Alana Gerhardt[1], Corey Casper[1,4,5,6], Troy D. Randall[3], Frances E. Lund[2], Emily

A. Voigt[1*]

Affiliations:

[1]Access to Advanced Health Institute (formerly Infectious Disease Research Institute); Seattle,

WA 98102, USA.

[2]Department of Microbiology, University of Alabama at Birmingham (UAB), Birmingham, AL

35294, USA

[3]Department of Medicine, Division of Clinical Immunology and Rheumatology, UAB,

Birmingham, AL 35294, USA

[4]Department of Global Health, University of Washington; Seattle, WA 98195, USA.

[5]Department of Medicine, University of Washington; Seattle, WA 98195, USA.

[6]Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Center; Seattle, WA 98109,

USA.

†These authors contributed equally to this work.

*Corresponding author: emily.voigt@aahi.org

Short title: An intranasal saRNA vaccine for SARS-CoV-2

Abstract:

While mRNA vaccines have been effective in combating SARS-CoV-2, waning of vaccine-induced antibody responses and lack of vaccine-induced respiratory tract immunity contribute to ongoing infection and transmission. In this work, we compare and contrast intranasal (i.n.) and intramuscular (i.m.) administration of a SARS-CoV-2 self-amplifying RNA (saRNA) vaccine delivered by a nanostructured lipid carrier (NLC). Both i.m. and i.n. vaccines induce potent systemic serum neutralizing antibodies, bone marrow-resident IgG-secreting cells, and robust lymphoid tissue T cell immune responses. The i.n. vaccine additionally induces robust respiratory mucosal immune responses, including SARS-CoV-2-reactive lung-resident memory and lung-homing T cell populations. As a booster following previous i.m. vaccination, the i.n. vaccine also elicits the development of mucosal virus-specific T cells. Both the i.m. and i.n. administered vaccines protect hamsters from infection-associated morbidity upon viral challenge, significantly reducing viral loads and preventing challenged hamsters from transmitting virus to naive cagemates. This saRNA vaccine's potent systemic immunogenicity, and additional mucosal immunogenicity when delivered i.n., may be key for combating SARS-CoV-2 and other respiratory pathogens.

One Sentence Summary:

Intranasal SARS-CoV-2 saRNA vaccination induces systemic and mucosal immunity in mice, and prevents morbidity and blocks viral transmission in hamsters.

Main Text:

INTRODUCTION

The ongoing severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic has led

to the rapid advancement and development of novel vaccines for coronavirus disease 2019

(COVID-19). Since their introduction in late 2020, two now FDA-approved mRNA vaccines

(Comirnaty and Spikevax) have had a remarkable impact on the trajectory of the pandemic and

were developed at unprecedented speeds (*1*). However, while these vaccines have displayed

remarkable efficacy at preventing severe COVID-19 and slowing the worldwide pandemic, the

lack of vaccine-induced local immunity in the respiratory tract allows for mild infection of

vaccinated individuals, resulting in transmission of the virus, despite prior vaccination (*2–4*).

Between lack of vaccine induction of mucosal immunity, newly emerging variants (*5*), waning

efficacy of currently approved mRNA vaccines (*6*), need for frequent booster doses (*7*), vaccine

hesitancy in part fueled by a fear of needles (*8*), and limited access to cold chain storage (*9*),

global susceptibility to SARS-CoV-2 remains high. Improved vaccines and associated

technologies are therefore necessary to fully combat the epidemic.

We previously developed an intramuscularly (i.m.) administered nanostructured lipid carrier

(NLC)-delivered self-amplifying RNA (saRNA) vaccine against SARS-CoV-2 (*10, 11*), which

induces both high serum neutralizing responses (*12–14*) and robust activation of systemic CD4$^+$

and CD8$^+$ T cells, responses that may be critical to sustained potent immunity (*13*). This saRNA-

NLC vaccine platform additionally has unique thermostability relative to other mRNA delivery

systems (*11, 15–17*), as it may be readily lyophilized and stored at room temperature for months

or under refrigeration likely for years (*11, 17*).

However, we wanted to examine the potential for this vaccine to be administered intranasally

(i.n.) to stimulate mucosal immune responses (*18*), which may be key to improving sustained,

potent immunity to SARS-CoV-2 and other respiratory pathogens. Unlike i.m. administered

vaccines, i.n. vaccines induce critical immune cells within the mucosa and respiratory tract,

generating IgA-secreting cells as well as specific effector cell subsets, including tissue-resident

memory (T_{RM}) T cells that are maintained within the lung (*19–26*). Such mucosal immunity,

mediated by IgA antibodies and lung-resident memory T and B cells (*27*), can prevent early-

stage infection and viral replication and shedding, minimizing chances for viral transmission (*20*,

21, *26*, *28*, *29*). Effective i.n. vaccines can also induce cross-reactive antibodies and thus provide

expanded protection to multiple strains (*30*). Intranasal vaccination also presents many other

advantages, such as easier needle-free administration, higher patient compliance, simpler

pediatric dosing, and easier storage and transport (*31*), and thus potentially increased vaccine

uptake. However, while several i.n. vaccines are currently under development for SARS-CoV-2,

most of these rely on older vaccine technologies, i.e., live-attenuated viruses and viral vectors

(*1*), creating a slow pathway to manufacturing, testing, and licensing. Fortunately, bridging RNA

vaccine technologies to i.n. delivery is possible, combining the production speed of RNA

vaccines with the mucosal immune stimulation of i.n. vaccines.

We now demonstrate that both i.n. and i.m. administrations of our saRNA SARS-CoV-2 vaccine

strongly induce systemic immunity in preclinical models and prevent viral transmission by

vaccinated, virally challenged hamsters. Intranasal dosing uniquely induces strong mucosal T

cell immunity, and a heterologous dosing strategy—i.m. prime followed by i.n. boosting—

appears to show the greatest potential to maximize mucosal and systemic immunity, leading to

prevention of viral transmission. This saRNA-NLC vaccine candidate combines the flexibility

and manufacturability of RNA vaccines with potent induction of mucosal immunity using a

thermostable formulation that represents a significant new tool for current and future pandemic

responses.

RESULTS

Fig. 1. Design of saRNA-NLC SARS-CoV-2 vaccine AAHI-SC2. **(A)** saRNA SARS-CoV-2

construct diagram. **(B)** NLC-RNA complex diagram.

Intranasal saRNA-NLC vaccine stimulates robust systemic immune responses

For development of an intranasally delivered saRNA vaccine against SARS-CoV-2, we

harnessed our previously developed and published long-term thermostable SARS-CoV-2

saRNA-NLC vaccine (*11*). This vaccine, currently in clinical trials using intramuscularly

administered material, contains a codon-optimized Wuhan-strain D614G SARS-CoV-2 spike

protein sequence driven by a Venezuelan equine encephalitis virus (VEEV)-based replicon

(Figure 1A) (*11*). This saRNA is complexed to the exterior of AAHI's proprietary NLC

nanoparticle to create the final vaccine product AAHI-SC2 (Figure 1B).

To investigate the ability of the saRNA-NLC vaccine to generate effective immunogenicity when

delivered intranasally, 6–8-week-old C57BL/6J mice were given prime and boost vaccine doses

of 1, 5, or 10 µg, 3 weeks apart. Control mice were i.n. dosed with 10 µg of a non-immunogenic

saRNA-NLC expressing the reporter protein secreted alkaline phosphatase 2 (SEAP). Systemic

immunogenicity was compared between groups. Serum, bone marrow, lung, and spleen tissues

were harvested after vaccination and tested for humoral (serum IgG and pseudovirus

neutralization titers; bone marrow antibody-secreting cells [ASCs] by enzyme-linked

immunospot [ELISpot]) and cellular (systemic T cell responses by intracellular cytokine staining

and flow cytometry [ICS-flow] and T cell ELISpot of splenocytes; mucosal T cell responses by

ICS-flow of lungs) vaccine-induced SARS-CoV-2-directed immune responses.

The i.n. saRNA-NLC vaccine elicited high serum spike-specific IgG and SARS-CoV-2 Wuhan-

strain neutralization titers that would be predicted to be protective (>1000 IC_{50}) (*32, 33*) at all i.n.

doses after prime and boost (Figure 2A, B). Strikingly, at the 5 and10-µg doses, no significant

differences were observed between i.n. and i.m. vaccines post-boost, suggesting equivalent

induction of serum responses – the key correlates of protection for SARS-CoV-2 (*32*, *33*) – by

the i.n. vaccine. IgG1 and IgG2a serum antibody titers indicated that i.n. vaccination with the

saRNA-NLC vaccine maintains the desired Th1 bias seen with our i.m. vaccine (Figure 2C).

Next, bone marrow-resident ASCs were assessed post-boost by ELISpot to determine if there

was establishment of a memory niche for IgG-secreting ASCs. No differences were seen

between the i.n. and i.m. vaccines at either the 5- or 10-µg dose i.n. as compared to the 10-µg

dose i.m. (Figure 2D), further suggesting effective systemic induction by both delivery routes

and suggesting durability of response established by both vaccines.

Cell-mediated immunity, particularly T cell immunity, is also thought to be critical to sustained,

long-term protection against SARS-CoV-2 (*13*, *33*). We therefore investigated systemic cellular

responses in i.n. and i.m. vaccinated mouse spleens after boost, both by ELISpot and ICS.

Significant populations of IFNγ-secreting T cells were observed in both the i.n. and i.m. dosed

(10-µg dose) mouse groups as measured by ELISpot (Figure 2E), indicating effective induction

of effector T cells primed to respond to SARS-CoV-2. While the cell numbers were significantly

higher in the i.m. dosed group, both groups showed robust T cell responses. IL-5- and IL-17-

secreting cells were seen at negligible populations in either group (Figure 2F-G), suggesting a

strong Th1-biased response (*34*, *35*) induced by both the i.n. and i.m. vaccines with very little

induction of a potentially harmful Th2 response associated with immunopathology and natural

infection (*34*, *36*, *37*).

These robust, strongly Th1-skewed systemic $CD4^+$ and $CD8^+$ T cell responses were verified

using ICS-flow performed on mouse splenocytes collected post-vaccination. After prime-boost

immunization, significant populations of spike-reactive polyfunctional ($IFN\gamma^+$, $IL-2^+$, and

$TNF\alpha^+$) $CD8^+$ and $CD4^+$ T cells were observed in mice immunized with the 5- and 10-μg i.n.

vaccine doses and the 10-μg i.m. dose (Figure 2H, I), demonstrating strong T cell-mediated

responses induced by both i.m. and i.n. vaccines. The spike-reactive polyfunctional T cell

populations of $CD4^+$ T cells induced by the i.n. vaccine were lower than those induced by the

i.m. vaccine (20-30% reduction between the i.n. and i.m. vaccines at a 10-μg dose, $p = 0.0191$);

however, spike-reactive polyfunctional $CD8^+$ T cell populations were not significantly different

between the i.n. and i.m. vaccinated mice, and $CD4^+$ T cell responses induced by the i.n. vaccine

were still high, demonstrating effective seeding of systemic T cell immunity across all i.n. and

i.m. vaccinated mice. As seen with vaccine-induced antibody titers, a stronger induction of a

Th1-specific response (secretion of $IFN\gamma^+$, $IL-2^+$, or $TNF\alpha^+$) compared to a Th2 response

(secretion of $IL-5^+$ or $IL-10^+$) or a Th17 response (secretion of $IL-17^+$) occurred within the T cell

compartment in both i.n. and i.m. vaccinated mice and across all doses (Figure S2A), suggesting

an overall systemic functional response to the virus. These data therefore indicate that the i.n.

saRNA-NLC vaccine elicits humoral and cellular responses at levels predicted to be protective.

Fig. 2. Systemic immunogenicity of intranasally administered SARS-CoV-2 saRNA-NLC vaccine. Systemic immunogenicity was assessed 3 weeks post-prime or post-boost in C57BL/6J mice immunized with AAHI-SC2 i.m. at 10 µg or i.n. at 1, 5, or 10 µg or a vector control (SEAP) i.n. at 10 µg at day 0 and day 21. $n = 8$ mice per group, evenly split male-female. **(A)** Serum SARS-CoV-2 spike-specific IgG responses measured post-prime and post-boost. LOD = limit of detection. **(B)** Serum SARS-CoV-2 spike Wuhan pseudovirus neutralizing titers measured post-prime and post-boost. **(A-B)** Data were log transformed and analyzed by a mixed-effects model with Tukey's multiple comparisons test. **(C)** Serum spike-specific IgG1 and IgG2a responses measured post-boost. **(D)** Induction of bone marrow-resident spike-specific IgG-secreting ASCs as measured by ELISpot 3 weeks post-boost. **(E-G)** SARS-CoV-2 spike-

responsive secretion of IFNγ, IL-5, or IL-17a by splenocytes post-boost as measured by ELISpot

in mice dosed with 10 µg of saRNA. All bars on graphs show geometric mean ± geometric SD.

(H-I) Splenic spike-responsive polyfunctional T cells (IFNγ$^+$, IL-2$^+$, and TNFα$^+$ CD4$^+$ or CD8$^+$

T cells) post-boost as measured by intracellular cytokine staining and flow cytometry. Horizontal

lines show mean ± SD. See Figure S1 for flow cytometry gating strategies. **(D-I)** Data were

analyzed by one-way ANOVA with Tukey's multiple comparisons test. *$p < 0.05$ and ****$p <$

0.0001.

Intranasal vaccine administration induces specialized lung T cell populations

While both i.m. and i.n. vaccines induce excellent systemic immunity, a key objective to

assessing i.n. immunization was to determine whether effective mucosal responses could be

generated with i.n. administration of an RNA vaccine. To that end, the lung T cell compartment

was analyzed from C57BL/6J mice vaccinated i.n. or i.m. with 10 μg of AAHI-SC2 3 weeks

post-prime and 3 weeks post-boost; the lung T cell compartment was analyzed by ICS-flow for

the presence of specialized T cell subsets that would indicate the development of long-lasting

respiratory-specific immunity.

First, we examined post-vaccination lung infiltration of antigen-experienced memory T cells.

These T_{RM} cells are characterized by CD69 expression as well as CD103 expression in the CD8

compartment (*38*). The percentage of T_{RM} cells were significantly increased in the CD8

compartment with i.n. vaccination (median of CD69$^+$ cells >20% of CD8$^+$) compared to i.m.

vaccination (median of CD69$^+$ cells <10% of CD8$^+$) following boost (CD69$^+$ CD8$^+$ cells, *p* =

0.0029; CD69$^+$ CD103$^+$ CD8$^+$ cells, *p* < 0.0001) (Figure 3A, B). While the fractions of CD8$^+$

T_{RM} cells were equivalent post-prime, lung expansion of T_{RM} cells followed boost administration

in the i.n. dosed group only, suggesting seeding of these durable populations followed by recall

and expansion upon boost dosing.

We then assessed CD154$^+$ CD4$^+$ T cells. CD154 (CD40 ligand [CD40L]) is often expressed on

activated T cells and T follicular helper cells and is critical for licensing dendritic cells and

engaging with B cells to enhance mucosal antibody production (*39*). Here, we found similar

patterns to those seen with T_{RM} cells; significantly increased populations of CD154$^+$ CD4$^+$ cells

were observed in the lungs after i.n. saRNA vaccination post-boost compared to i.m. vaccination

($p = 0.0029$) (Figure 3C). Collectively, these data show that i.n. administration appears to elicit

an expansion of T_{RM} and $CD40L^+$ activated $CD4^+$ T cells within the lung. In contrast, i.m.

vaccination elicits minimal levels of these cell subsets, suggesting that these specialized

respiratory cells are not inducible by i.m. vaccination.

Fig. 3. Mucosal T cell populations following intranasal vaccination with SARS-CoV-2

saRNA-NLC candidate. T cell populations in the lungs were assessed 3 weeks post-prime and

post-boost in C57BL/6J mice immunized with 10 µg of AAHI-SC2 either i.m. or i.n. *n* = 8 mice

per group, evenly split male-female. Vector control-dosed mice were also included in the

experiment and are shown in Figure 4. See Figure S3 for sample flow cytometry gating

strategies. **(A-B)** CD69$^+$ lung-resident memory (A) CD8$^+$ and (B) CD103$^+$ CD8$^+$ T cells. **(C)**

CD40L$^+$ (CD154$^+$)-expressing CD4$^+$ T cells. Percentage of cells in total live lung cells per mouse

indicated by dots with horizontal line at median. Data were analyzed by two-way ANOVA with

Sidak's multiple comparisons test. *******p* < 0.01 and *********p* < 0.0001.

Intranasal administration elicits polyfunctional SARS-CoV-2 reactive mucosal T cell responses

While the lung T cell subsets were present in large populations post-i.n. administration, we

wanted to verify that these cell subsets were functional effector cells and confirm that they

responded to SARS-CoV-2. Here, we took mice that received the vector control or had been

immunized and boosted with AAHI-SC2 i.m. (10 µg) or i.n. (1 µg, 5 µg, or 10 µg) and used ICS-

flow cytometry to determine the frequency of polyfunctional lung $IFN\gamma^+IL-2^+TNF\alpha^+$ $CD4^+$ and

$CD8^+$ T cells following re-stimulation for 6 hours with overlapping spike peptides.

Polyfunctional $IFN\gamma^+IL-2^+TNF\alpha^+$-secreting SARS-CoV-2 spike-responsive T cells composed a

large fraction of the key T_{RM}, $CD69^+$ population in the 5- and 10-µg i.n. immunized mice (12.8%

and 9.11%, respectively), which was significant in the $CD69^+$ $CD8^+$ subset of these T_{RM} cells

compared to the i.m. immunized mice ($p = 0.0005$ for the 5-µg dose and $p = 0.0258$ for the 10-

µg dose), as well as in the $CD69^+$ $CD103^+$ $CD8^+$ subset ($p = 0.0181$) from 5-µg i.n. immunized

mice (Figure 4A-B). Intramuscularly immunized groups again induced consistently negligible

levels of polyfunctional $CD8^+$ cells. We also examined the activated $CD154^+$/$CD40L^+$ $CD4^+$ T

cells and the lung-homing $CD194^+$/$CCR4^+$ $CD4^+$ and $CD8^+$ cells. Similar to the T_{RM} cells,

$CD154^+$ $CD4^+$ cells showed significantly higher percentages of polyfunctionally responding cells

at the 5-µg i.n. dose compared to the i.m. dose ($p = 0.0006$) (Figure 4C). Likewise, $CD194^+$

$CD4^+$ and $CD8^+$ cells were also induced by the i.n. vaccine at significantly higher percentages at

the 5-µg dose ($p = 0.0441$ and $p = 0.0071$, respectively) compared to the i.m. vaccine (Figure

4D-E).

Finally, we examined the quality of the response by lung T cells to confirm that i.n.

administration elicited Th1-biased lung T cell responses, like what was observed in the spleen.

The majority of the responses to both the i.n. and i.m. vaccines were Th1-specific responses,

with negligible Th2 and Th17 responses (Figure S2B), similar to what was observed in the

spleen. Th2 responses were not significantly different between i.m. vaccinated mice and mice

receiving the vaccine via the i.n. route, did not scale with increasing vaccine dose, and were not

significantly increased from buffer-only groups. Collectively, these data demonstrate that i.n.

administration seeds specialized polyfunctional CD4$^+$ and CD8$^+$ T cell populations in the lung –

populations which are not elicited following i.m. administration.

CD69⁺ lung-resident memory T cells (T$_{RM}$)

CD40L⁺ lung cells (activated T cells) **CCR4⁺ lung-homing T cells**

Fig. 4. Mucosal SARS-CoV-2 spike polyfunctionally reactive T cell populations following immunization with the intranasal or intramuscular SARS-CoV-2 saRNA-NLC vaccine. T cell populations in the lungs were stimulated with SARS-CoV-2 spike peptides for 6 hours and stained for T cell markers and cytokine expression 3 weeks post-boost in C57BL/6J mice immunized with 1, 5, or 10 µg of the i.n. saRNA-NLC vaccine, relative to 10 µg of the i.m. vaccine or an i.n. vector control (SEAP) saRNA-NLC. n = 8 mice for all dosing groups. Percentages of polyfunctional cells (expressing IFNγ, IL-2, and TNFα) were then measured by flow cytometry. **(A-B)** Polyfunctional CD69⁺ lung-resident memory (A) CD8⁺ and (B) CD103⁺ CD8⁺ T cells. **(C)** Polyfunctional CD40L-expressing (CD154⁺) CD4⁺ T cells. **(D-E)** Polyfunctional CCR4-expressing (CD194⁺) lung-homing (D) CD4⁺ and (E) CD8⁺ T cells. Dots

represent values measured from individual mice. Horizontal lines indicate data median.

Significance was determined by one-way ANOVA with Tukey's multiple comparisons test: *$p <$

0.05, **$p < 0.01$, and *** $p < 0.001$.

Intranasal boosting of previous i.m. vaccination induces both systemic and mucosal immune

responses

To understand whether i.n. administration could be viable as a boosting strategy for individuals

previously vaccinated with an i.m. SARS-CoV-2 vaccine, we examined mice that were primed

with an i.m. vaccine and then boosted with a medium dose (5 µg) either i.m. or i.n. Intranasally

boosted mice showed equivalent serum anti-spike IgG levels and similar systemic neutralizing

antibody responses to the i.m. boosted mice (Figure 5A, B). Within the splenic $CD4^+$ and $CD8^+$

T cell compartment, i.n. boost immunization induced equivalent or increased spleen-resident

polyfunctional T cell responses in $CD4^+$ or $CD8^+$ ($p = 0.0454$) cells, respectively, compared to an

i.m. boost immunization (Figure 5C, D), indicating that an i.n. booster may effectively boost the

systemic immunity developed in response to an original i.m. vaccination. With respect to

mucosal immunity, the i.n. booster induced a higher but not significant percentage of $CD40L^+$

($CD154^+$) activated $CD4^+$ T cells within the lungs than the i.m. booster (Figure 5E). Most

importantly, the i.n. booster elicited significantly more polyfunctional lung-resident $CD8^+$ $CD69^+$

T_{RM} cells ($p = 0.0236$) than the i.m. booster (Figure 5F), indicating that just boosting with an i.n.

dose may be sufficient to elicit the full mucosal immune-stimulating effect of i.n. vaccination.

These data are consistent with other i.n. vaccines that suggest i.n. boosting can have enhanced

efficacy systemically as compared to i.m. vaccines (*21*). This suggests that an i.n. boost

vaccination strategy may be ideal to boost systemic and induce respiratory immunity in

previously vaccinated individuals, reflecting the real-world scenario of second-generation SARS-

CoV-2 vaccine rollouts.

Fig. 5. Systemic and mucosal immune responses following boost immunization with the

intranasal or intramuscular SARS-CoV-2 saRNA-NLC vaccines. 10-week-old C57BL/6J

mice were injected i.m. with a 10-μg prime dose (N:P of 15) and then boosted with a 5-μg dose

of either the i.n. or i.m. saRNA-NLC vaccine 7 weeks later. n = female 5 mice per group. **(A)**

SARS-CoV-2 spike-specific IgG titers as measured by ELISA 3 weeks post-boost. Horizontal

lines show median. **(B)** Serum SARS-CoV-2 pseudovirus neutralizing titers 3 weeks post-boost.

Horizontal lines show median. **(C-D)** Spleen polyfunctional (D) CD4[+] and (E) CD8[+] T cell

responses. **(E)** Polyfunctional CD40L-expressing (CD154[+]) CD4[+] T cells. **(F)** Polyfunctional

lung-resident memory CD69[+] CD8[+] T cell responses. Statistics evaluated as unpaired two-sided

homoscedastic t tests, *$p < 0.05$. Bars on graphs show mean with SD.

saRNA vaccination protects against SARS-CoV-2-induced disease in hamsters

Given our data showing that i.m. prime and i.n. boost with our saRNA-NLC SARS-CoV-2

vaccine elicited strong systemic and pulmonary immunity in mice, we hypothesized that

vaccination, particularly via the i.m./i.n. route, would be sufficient to protect animals from

morbidity due to SARS-CoV-2 infection and would prevent these animals from transmitting the

virus to naive bystander animals. To test this hypothesis, we co-housed hamster pairs (1:1) for

several weeks to acclimate them to one another. Then, paired animals were separated and one

animal in each pair was either sham-vaccinated or vaccinated with 5 µg of the saRNA-NLC

SARS-CoV-2 vaccine given either i.m. or i.n (Figure 6A). After 24 hours, the sham-vaccinated

and vaccinated animals were re-paired with their naive (non-vaccinated) cagemates. On day 21,

peripheral blood was collected from all animals. On day 28, all pairs of animals were separated,

and the sham-vaccinated and vaccinated animals were boosted via the i.m. or i.n. route. All

animals were re-paired with their naive cagemates after 24 hours and blood was collected from

all animals five days later. As expected, vaccinated hamsters, particularly i.m.-dosed hamsters,

demonstrated strong induction of spike-specific serum IgG post-prime, which was absent in the

sham-vaccinated group (Figure 6B) and in all the naive cagemates (Figure S4). Intranasally

primed animals also showed development of significant serum IgG titers, albeit at titers

significantly lower than those observed in i.m.-primed animals ($p < 0.0001$). Boosting on day 28

did not result in a significant increase in systemic anti-spike IgG responses in any of the

vaccinated groups (Figure 6C), perhaps due to the still ongoing robust primary response.

At 9 days post-boost (day 37), the vaccinated and sham-vaccinated hamsters were separated from

their naive cagemates (Figure 6A). Some sham-vaccinated hamsters were set aside to serve as

uninfected negative controls (sham-vaccinated-uninfected), and all other vaccinated or sham-

vaccinated hamsters were experimentally infected with 1 x 10^6 pfu SARS-CoV-2 USA/WA-1

strain delivered intranasally. We then compared morbidity of the vaccinated-infected and sham-

vaccinated-infected animals to the sham-vaccinated-uninfected controls. As expected, the sham-

vaccinated-uninfected hamsters exhibited a modest increase in weight over the next 14 days

(Figure 6D). In contrast, the sham-vaccinated-infected animals had increased morbidity with

peak body weight loss of 9-10% by day 6 post-infection and minimal body weight recovery by

termination of study. The vaccinated-infected animals (i.n.-i.n., i.m.-i.n., and i.m.-i.m.

vaccination), on the other hand, showed minimal weight loss and exhibited rapid recovery in

weight by day 3-4 post-infection, with statistically significantly reduced virus infection-induced

morbidity when compared to the sham-vaccinated-infected group ($p = 0.0091$, $p = 0.0074$, and p

$= 0.0206$ for i.n.-i.n., i.m.-i.n., and i.m.-i.m. vaccinated-infected animals, respectively). Body

weight loss after challenge did not correlate with systemic anti-spike IgG titers, suggesting that

either all three vaccination regimens induced sufficient systemic IgG to protect against SARS-

CoV-2-induced weight loss and/or cellular or mucosal responses played a role in protection

(Figure S5).

To further understand the dynamics of infection and transmission, total and subgenomic viral

nucleocapsid (N) RNA was measured in nasal swipes from all infected hamsters across the first

10 days after infection. Total viral N RNA was detected in all infected hamsters as early as 1

hour post infection (Figure 6E), which confirmed successful intranasal instillation of SARS-

CoV-2. Total viral copies peaked between 24-48 hours in all infected groups. However, total

viral N RNA copies were significantly lower in the vaccinated-infected groups compared to the

sham-vaccinated-infected group ($p < 0.0001$, $p < 0.0001$, and $p = 0.0003$ for i.n.-i.n., i.m.-i.n.,

and i.m.-i.m., respectively). Next, subgenomic viral N RNA was quantified in the nasal discharge

to compare infectious, actively replicating viral load between the sham-vaccinated-infected and

vaccinated-infected hamsters. Subgenomic viral N RNA was detected in the nasal swipes from

all infected hamsters. However, significantly diminished subgenomic viral N RNA copies were

observed in all vaccinated-infected animals relative to the sham-vaccinated-infected animals

(Figure 6F) ($p < 0.0001$, $p = 0.0002$, and $p < 0.0001$ for i.n.-i.n., i.m.-i.n., and i.m.-i.m. vaccine

regimens, respectively). Moreover, the peak copies of subgenomic N RNA in nasal swipes were

observed within 24 hours of infection in the vaccinated-infected hamsters, while subgenomic N

RNA peaked in the sham-vaccinated-infected hamsters between days 1-2 (Figure 6F).

Importantly, the vaccinated-infected hamsters had increased clearance of viral subgenomic N

RNA that declined to undetectable levels by day 6 (i.m.-i.n. and i.m.-i.m. groups) and day 8 (i.n.-

i.n. group) post-infection. In contrast, the sham-vaccinated-infected control animals had

detectable viral subgenomic N RNA until day 10, indicating that vaccination promoted more

rapid reduction in replicating virus.

The viral RNA measurements from nasal swipes suggested that none of the saRNA-NLC

vaccination regimens induced sterilizing immunity following direct intranasal instillation with

1×10^6 pfu of SARS-CoV-2. To address whether the vaccinated animals made an immune

response to experimental infection, we measured systemic anti-N IgG antibodies on day 35 post-

infection. Indeed, anti-N IgG antibody was detected in the serum from all three groups of

vaccinated-infected animals (Figure 6G), indicating that the amount of N protein made by the

infectious virus was sufficient to elicit a *de novo* N-specific antibody response in these hamsters.

However, compared to the sham-vaccinated-infected animals, the amount of anti-N IgG antibody

was significantly lower in the i.m.-i.m. and i.m.-i.n. vaccinated-infected hamsters ($p < 0.0001$

and $p = 0.0470$, respectively) and modestly lower in the i.n.-i.n. vaccinated-infected hamsters,

again consistent with a decreased viral load in all vaccinated animals.

Next, to assess SARS-CoV-2-induced pulmonary pathology, cross sections of the hamster lungs

were examined and scored for histology by a board-certified pathologist. As expected, the sham-

vaccinated-infected hamsters had mild to moderate histology scores confirming that the SARS-

CoV-2 viral inoculum used to infect the hamsters caused detectable pulmonary

immunopathology as late as day 35 post-infection (Figure 6H). These sham-vaccinated-infected

hamsters exhibited signs of inflammation that consisted primarily of accumulation of

mononuclear and polymorphonuclear infiltrates in the alveolar interstitium and lumen, and

adjacent inflammation around blood vessels and bronchioles (Figure S6A). Conversely, all three

groups of vaccinated-infected hamsters had significantly decreased total histology scores (Figure

6H) and exhibited only sporadic interstitial, perivascular, and peribronchiolar inflammation

(Figure S6B) that were comparable to sham-vaccinated-uninfected negative control animals

(Figure S6C), suggesting that vaccination reduced the extent and severity of lung pathology

following infection with a high dose of SARS-CoV-2. Therefore, while vaccination did not

prevent infection following a robust viral challenge, all vaccine regimens effectively prevented

disease-associated morbidity as measured by weight loss, decreased peak viral copies in the nasal

discharge, suppression of viral replication, increased viral clearance, and protection against

SARS-CoV-2-induced lung damage.

Fig. 6. Protection from SARS-CoV-2 infection-induced morbidity, viral load, and lung damage in vaccinated hamsters. (A) Study design for experiment in vaccinated and sham-vaccinated co-housed hamsters. Acclimated co-housed hamster pairs were separated on day 0. One animal in each pair was vaccinated i.n. or i.m. or was sham-vaccinated. The other animal in each pair was left untouched (naive, see Figure 7). Sham-vaccinated and vaccinated animals were re-paired with their naive cagemates on day 1. Blood was collected from all animals on day 21. Animals were separated again on day 28, and sham-vaccinated and vaccinated animals were

406 boosted either i.n. or i.m. Animals were re-paired with their naive cagemates on day 29, and

407 blood was collected on day 33. (**B-C**) Vaccine antibody responses in sham-vaccinated (open

408 black circles), i.n.-prime/i.n.-boost vaccinated (blue diamonds), i.m.-prime/i.m.-boost vaccinated

409 (yellow squares), or i.m.-prime/i.n.-boost vaccinated (green triangles) animals. *n* = 5-6/group.

410 Mean fluorescence intensity (MFI) of serum anti-spike IgG on day 21 (**B**) and day 33 (**C**). Data

411 were log-transformed and assessed by two-way ANOVA with Tukey's multiple comparison test.

412 Horizontal lines show geometric mean. (**D-H**) Experimental infection in sham-vaccinated and

413 vaccinated hamsters. Vaccinated and sham-vaccinated hamsters were separated from their naive

414 cagemates on day 37 (see panel A). Two sham-vaccinated hamsters were left uninfected (grey

415 triangles, sham-vaccinated-uninfected), and all remaining sham-vaccinated and vaccinated

416 animals were experimentally infected with SARS-CoV-2 via the intranasal route. (**D**) Body

417 weight of vaccinated-infected, sham-vaccinated-infected, and sham-vaccinated-uninfected

418 hamsters. Body weight line plots show group weight mean and SEM. Body weight data was

419 analyzed on area under the curve measures, assessed statistically by one-way ANOVA with

420 Tukey's multiple comparison test. (**E-F**) Total N RNA (**E**) and subgenomic N RNA (**F**) copy

421 number in nasal swipes from vaccinated-infected and sham-vaccinated-infected hamsters. Viral

422 RNA measures were log-transformed and shown as mean and SEM. Area under the curve

423 measures were assessed with one-way ANOVA with Tukey's multiple comparison test. (**G**)

424 Anti-N IgG in serum of vaccinated-infected and sham-vaccinated-infected animals is shown as

425 MFI after log-transformation. Data statistically analyzed using one-way ANOVA with Dunnett's

426 multiple comparisons test. (**H**) Lung pathology scores were calculated for sham-vaccinated-

427 infected and vaccinated-infected animals. Horizontal lines and error bars represent mean and SD,

respectively. See Figure S6 for representative histopathology images. * $p < 0.05$, ** $p < 0.01$,

*** $p < 0.001$, **** $p < 0.0001$.

saRNA vaccination prevents viral transmission by virally challenged hamsters

Although all three groups of vaccinated-infected hamsters appeared to have been productively

infected following experimental SARS-CoV-2 exposure, the amount of subgenomic viral N

RNA in the nasal secretions of these hamsters had already declined significantly by day 2 post-

infection relative to the sham-vaccinated-infected group (Figure 6F). To assess whether this

reduction in infectious viral load was sufficient to prevent transmission, we re-paired the sham-

vaccinated-infected and vaccinated-infected hamsters with their naive cagemates on day 2 post-

infection (day 39 on timeline in Figure 6A; day 0 on timeline in Figure 7A) for a period of 24

hours (Figure 7A) and assessed morbidity, nasal swipe viral RNA, serum IgG antibodies, and

lung immunopathology in the naturally virally challenged cagemates. As expected, naive

cagemates of the sham-vaccinated-infected animals began losing weight within 24 hours of re-

pairing, reaching maximum weight loss of 6-7% by day 7 post-re-pairing and exhibiting minimal

body weight recovery by termination of study (Figure 7B). In striking contrast, the naive

cagemates of the vaccinated-infected hamsters exhibited no body weight loss following re-

pairing ($p = 0.0412$, $p = 0.0470$, and $p = 0.0478$ for i.n.-i.n., i.m.-i.n., and i.m.-i.m., respectively).

Given the little morbidity observed in the naive cagemates co-housed with the vaccinated-

infected hamsters, we hypothesized that viral transmission between the vaccinated-infected

animals and their naive cagemates would also be diminished. To test this, we measured total

(Figure 7C) and subgenomic (Figure 7D) viral N RNA in the nasal swipes from the naive

cagemates that were paired for 24 hours with the sham-vaccinated-infected and vaccinated-

infected groups. Total viral N RNA, which reflects both replicating and non-replicating virus,

was present in nasal swipes from all the naive cagemates as early as 24 hours post-re-pairing

with infected hamsters (Figure 7C). Interestingly, the naive cagemates paired with vaccinated-

infected hamsters had on average 70-fold lower total viral N RNA at 24 hours post-re-pairing

compared to naive hamsters that were co-housed with sham-vaccinated-infected animals (Figure

7C). Hence, immediately following re-pairing, the viral load transmitted to naive cagemates by

the vaccinated-infected animals was lower than that transmitted by the sham-vaccinated-infected

animals. Importantly, total viral N RNA in nasal secretions from the naive cagemates paired with

vaccinated-infected hamsters never surpassed 10^4 copies/mL (Figure 7C) and remained just

above the limit of detection starting at day 4 post-re-pairing ($p < 0.0001$ for all groups).

Conversely, the naive cagemates paired with sham-vaccinated-infected hamsters had total viral N

RNA nasal discharge titers that peaked at 10^7 copies/mL by day 2 post-re-pairing, with minimal

clearance by end of study at day 8 post-re-pairing (Figure 7C). Similarly, subgenomic viral N

RNA in the nasal discharge, which is indicative of viral replication/infection, was detected at 24

hours post-re-pairing in the naive cagemates co-housed with sham-vaccinated-infected hamsters,

peaked at 10^5 copies/mL by day 2-3 post-re-pairing and remained detectable even out to day 8

post-re-pairing (Figure 7D). This was in striking contrast to the naive cagemates of the

vaccinated-infected animals, which had minimal to non-detectable levels of subgenomic viral N

RNA in the nasal discharge (Figure 7D) ($p < 0.0001$ for all three groups).

To assess whether the naive cagemates exposed to the vaccinated-infected animals were

protected from active infection, we measured *de novo* anti-N serum IgG responses. As expected,

the naive cagemates co-housed with sham-vaccinated-infected hamsters developed high serum

N-specific IgG responses (Figure 7E). The naive cagemates co-housed with vaccinated-infected

hamsters, on the other hand, exhibited negligible responses ($p < 0.0001$ for all vaccine regimen

groups). Indeed, no anti-N IgG response was detected in the naive cagemates of the i.n.-i.n.

vaccinated-infected group, and only one naive cagemate from the i.m.-i.n and the i.m.-i.m.

vaccinated-infected groups developed a measurable anti-N IgG serum antibody response. Thus,

all three vaccine regimens were effective at preventing significant transmission of infectious

virus capable of inducing an immune response to naive cagemates.

Next, we compared the total histology score of each group of naive cagemates (Figure 7F).

Naive hamsters co-housed with the sham-vaccinated-infected hamsters exhibited mild to

moderate histology scores, primarily due to mild inflammation in the alveolar interstitium and

lumen that was similar in severity to that observed in the sham-vaccinated-infected hamsters

(Figure S6A). Naive hamsters co-housed with the vaccinated-infected hamsters, regardless of

vaccination regimen, exhibited a significantly decreased histology score ($p < 0.0001$ for all

vaccine regimen groups), with minimal lung pathology that was very similar to that observed in

the vaccinated-infected cagemates (Figure S6B). Taken together, the data demonstrate that i.m.-

i.m., i.n.-i.n., or i.m.-i.n. dosing of the saRNA-NLC spike vaccine was not only effective in

preventing disease-associated morbidity in hamsters but also prevented the vaccinated and

subsequently experimentally infected hamsters from transmitting infectious virus to naive

cagemates, even at a time when viral load peaked in the nasal discharge.

Fig. 7. Vaccination prevents transmission of SARS-CoV-2 to naive hamster cagemates. (A)
Study design for transmission experiment in naive cagemates. Vaccinated and sham-vaccinated
hamsters were separated from their naive cagemates on day -2 (day 37 in Figure 6A) and then
experimentally infected with SARS-CoV-2. On day 0 (day 39 in Figure 6A), the naive animals
were re-paired with their vaccinated-infected or sham-vaccinated-infected cagemates. On day 1
(day 40 in Figure 6A), the animals were separated and the naive cagemates were monitored for
SARS-CoV-2 infection and infection-induced morbidity. **(B)** Body weight of the naive
cagemates that were re-paired for 24 hours with vaccinated-infected or sham-vaccinated-infected
hamsters. Body weight line plots show group weight mean and SEM. Body weight data was
analyzed on area under the curve measures, assessed statistically by one-way ANOVA with
Tukey's multiple comparison test. **(C-D)** Total N RNA **(C)** and subgenomic N RNA **(D)** copy

number in nasal swipes from naive cagemates that were re-paired for 24 hours with vaccinated-

infected and sham-vaccinated-infected hamsters. Viral RNA copy numbers were log-

transformed, and area under the curve measures are shown as mean and SEM. Data were

assessed with one-way ANOVA with Tukey's multiple comparison test. **(E)** Anti-N IgG in

serum of naive cagemates that were re-paired for 24 hours with vaccinated-infected and sham-

vaccinated-infected groups. Antibody levels are reported as mean fluorescence intensity (MFI)

following log-transformation. Data statistically analyzed using one-way ANOVA with Dunnett's

multiple comparisons test. **(F)** Lung pathology scores were calculated for naive cagemates that

were re-paired for 24 hours with sham-vaccinated-infected and vaccinated-infected animals.

Horizontal lines and error bars represent mean and SD, respectively. See Figure S6 for

representative histopathology images. * $p < 0.05$, **** $p < 0.0001$.

DISCUSSION

Here we extend previous work on this saRNA-NLC vaccine platform to show delivery of an intranasal RNA vaccine that induces key respiratory mucosal immune responses. This i.n. administered saRNA vaccine induced systemic immunity in both the humoral and cellular compartments that rivaled i.m. administration. Additionally, i.n. administration elicited not only robust populations of spike-responsive polyfunctional T cells in the spleen but also in the lung, demonstrating induction of key respiratory mucosal T cell populations by an RNA vaccine (*41–43*). Furthermore, we were able to demonstrate potential for this product as an i.n. booster of prior i.m. vaccine-induced immunity, a common real-life scenario given the largely COVID-experienced population at this point in the pandemic. This i.n. boost of prior i.m. vaccine-induced immunity significantly boosted systemic responses in addition to inducing mucosal T cell populations not stimulated by i.m. vaccine boosting. Lastly, we demonstrated that both i.n.-i.n., i.m.-i.m., and heterologous i.m.-i.n. vaccination regimens prevented virus-associated morbidity in the hamster model and, most importantly, prevented viral transmission and development of disease in naive cagemates.

Now in the third year of the COVID-19 pandemic, second-generation SARS-CoV-2 vaccines are urgently needed. While several vaccines have been developed and approved in the last 2.5 years, many utilizing novel mRNA vaccine technologies (*44–46*), which have admirably contributed to the waning of disease mortality globally (*47*), emerging trends highlight the insufficiencies of current vaccine approaches. Vaccine efficacy has waned rapidly (*48–51*), and frequent boosters have been recommended for individuals that received the Pfizer and/or Moderna mRNA vaccines (*49, 50, 52*). Increasing evidence of reinfection, both post-infection and post-

vaccination (*53*), particularly with the Omicron variant (*54*), as well as ongoing viral

transmission post-vaccination (*55–57*), has negatively impacted vaccine credibility. With the

ongoing pandemic predicted to remain a present threat, new variants will continually place

pressure on the efficacy of these first-generation vaccines (*58*, *59*).

While more research is necessary as to why current vaccines are waning, emerging research is

pointing to insufficient nasal and mucosal immunity (*56*, *60*). Increasingly, data are

demonstrating that i.m. vaccination alone may not generate potent mucosal immunity that is

predicted to be critical for early control and clearance of SARS-CoV-2 (*61*). Vaccinees still show

evidence of viral RNA in the upper respiratory tract after viral exposure, suggesting ongoing

viral replication and transmission (*20*, *22*, *56*, *62*) even by vaccinated populations. However,

breakthrough infection with SARS-CoV-2 allows vaccinees to elicit potent mucosal and nasal

immunity (*41–43*). Thus, improving the induction of nasal and mucosal immunity following

vaccination is a critical component for second-generation vaccines to finally achieve durable and

flexible protection. However, data also show that generating this immunity is generally not

possible via standard i.m. vaccination and likely requires administration of an i.n. vaccine (*60*,

63).

Intranasal vaccines to date have primarily been developed as live-attenuated flu vaccines,

including FluMist (*64*). Recently, four new i.n. or inhalable vaccines have been approved for

SARS-CoV-2 in China, India, Iran, and Russia, including two adenovirus-vectored vaccines (*65–*

67). Further development of nasal or airway administration of vaccines for influenza and other

respiratory viruses has remained an attractive target given the potential for development of

critical airway T$_{RM}$ cells at this first line of defense (*3*, *68*). Several other i.n. vaccines are in

development for SARS-CoV-2, but most rely on standard protein or adenovirus vector vaccine

technologies (*20*, *21*, *43*, *69*, *70*). An RNA-based vaccine, in contrast, can be developed more

rapidly, and allows cross-over i.m. to i.n. vaccination without developing a new vaccine,

allowing for significantly greater utility in pandemic situations. The thermostability and

stockpiling ability of our saRNA-NLC vaccine platform additionally provide key attributes for

rapid and widespread pandemic response (*11*, *17*). Validating an RNA-based, i.n. administered

vaccine technology would be a significant development not only for the SARS-CoV-2 pandemic

but for other respiratory pathogens, combining the rapidity and reliability of RNA vaccine

adaptation to new targets with the key mucosal immune responses induced by mucosal-specific

dosing.

AAHI's SARS-CoV-2 saRNA-NLC vaccine candidate elicits both robust systemic humoral

immunity and spike-responsive systemic polyfunctional T cells as an i.m. or i.n. vaccine.

Delivered i.n., it also elicits tissue-resident T cell populations within the lungs, suggesting broad

T cell presence across the respiratory tract – an effect that i.m. delivered mRNA vaccines have

been unable to induce to date (*41–43*). While CD4 and CD8 T cell populations are now being

recognized as a major determinant of disease progression and critical to long-lasting immunity

(*13*), the main vaccines on the market elicit minimal peripheral T cell responses (*71*, *72*) and

negligible mucosal/respiratory T cell responses (*41–43*). The thermostability and ability to

administer i.n. may make this vaccine substantially more desirable in the fight against respiratory

viruses. These data demonstrate the potential for this i.n. saRNA-NLC vaccine candidate as an

effective booster to pre-existing SARS-CoV-2 immunity, outperforming an i.m. booster,

showing that the additive effect of this i.n. vaccine to pre-existing systemic immune responses

generated the greatest benefit of all vaccination regimens tested.

These data demonstrate that all immunization methods elicited full protection against virus-

associated morbidity and viral transmission. While our original hypothesis was that only i.n.

vaccination would prevent transmission, we were pleasantly surprised that all three delivery

methods prevented transmission. These data bolster the overall strength of the vaccine by

demonstrating that i.m. vaccination is protective as well. While it is unclear the specific cellular

or serum components mediating this protection in i.m. vaccinated animals, which lack substantial

lung T_{RM} cell populations and significant mucosal IgA, it may be that sufficiently high systemic

IgG so early after boost (9 days post-boost) may circulate and prevent transmission (*73*, *74*).

Future studies examining protection and transmission by vaccinated hamsters at longer

timepoints post-vaccination may more fully demonstrate the advantages of i.n. vaccine

administration and induction of mucosal immunity on protection and transmission prevention

relative to i.m.-i.m. or heterologous i.m.-prime, i.n.-boost vaccine regimens.

The i.n. saRNA vaccine may be most effective as a booster dose following an i.m. prime. This

aligns with the results of other studies (*14*, *62*, *75*) where priming an immune response with a

strong i.m. vaccine followed by i.n. vaccination is able to optimally stimulate the T_{RM}

compartment while still boosting the extant i.m. induced immunity. Our data demonstrate the

clear potential of using an identical or similar RNA vaccine product for both i.m. and i.n.

vaccination, potentially eliminating many manufacturing hurdles and using an already clinically

tested vaccine platform technology (*76*) for multiple routes of administration. In our studies, i.n.

boosting outperformed i.m. boosting particularly in mucosal immune measures, hinting at the

potential for i.n. dosing to access novel cell compartments and mechanisms. Further work

remains to fully optimize the i.n. saRNA vaccine formulation specifically for i.n. administration,

potentially including use of nasal spray devices, with the goal of enhancing induction of

immunity while avoiding toxicity and reducing heterogeneity in nasal delivery. As a first step in

this further development process, this i.n. SARS-CoV-2 vaccine candidate is being tested in

nonhuman primate studies to enable future human testing.

While these studies have been carefully designed and carried out to maximize scientific rigor and

reproducibility, they do have certain limitations. While we showed robust systemic and mucosal

immunity, protection, and transmission prevention, these data are developed in mice and

hamsters, and these responses may not translate fully to primates and humans. Ongoing studies

have been designed to investigate this issue. Our data are also limited by the heterogeneity in

mouse immune responses when vaccinated i.n., related to a lack of consistent dosing devices for

i.n. vaccination. Similarly, because mice are obligate nasal breathers (*77*) and vaccine volumes

are difficult to appropriately scale to mouse size, some potential exists for delivery of a fraction

of the vaccine volume beyond the nasal tract to the lungs and digestive tract. These issues are

likely only resolvable with larger animal models better representing the nasal passages and

respiratory tracts of humans, and use of appropriate delivery devices for improvement in vaccine

dose delivery. Similarly, some toxicity was noted in the highest i.n. delivered saRNA-NLC

vaccine dose (10 µg) in mice, resulting in notable transient mouse weight loss. While these

effects are not expected to scale to primates due to non-linear dose scaling of saRNA vaccines

relative to animal size and nasal tissue surface area, any potential toxicity noted with an i.n.

vaccine will be closely monitored in future studies, and further optimization of the NLC

formulation specifically for i.n. delivery will also be conducted to minimize any such effects.

Finally, we have been unable to test this saRNA-NLC vaccine candidate head-to-head against

any commercially available SARS-CoV-2 vaccines for comparison of either vaccine

immunogenicity or efficacy due to restrictions on vaccine sale and research use. Should such

comparisons become more feasible in the future, we anticipate them to greatly benefit future

work, including enabling further exploration of the use of this i.n. saRNA vaccine as an effective

mucosal immunity-inducing booster of prior approved SARS-CoV-2 vaccination.

This saRNA-NLC vaccine platform represents a promising technology to combat both the

existing SARS-CoV-2 pandemic and other respiratory illnesses, such as seasonal and pandemic

influenza. For respiratory pathogens, i.n. vaccination or i.n. vaccine boosting of i.m. vaccination

may help raise RNA vaccine technologies to a new level, through stimulating respiratory T cell

and B cell populations (*27*, *78*, *79*) and generating effector subsets within the lungs and nasal

mucosa primed to respond rapidly at the first point of infection (*62*). Beyond its efficacy at

stimulating T_{RM} subsets and bolstering systemic immunity, i.n. administration may also be

desirable as a needle-less delivery method with potentially higher uptake, particularly in

pediatric and vaccine-hesitant populations. Altogether, this novel demonstration of a unique i.n.

delivered RNA vaccine that elicits strong systemic and mucosal immunity is an indication of a

promising new technology ripe for future development and application.

MATERIALS AND METHODS

Study design

<u>Research objectives</u>

This study was designed to test the immunogenicity and efficacy of i.n. and i.m. delivered

SARS-CoV-2 saRNA-NLC vaccines in mice and hamsters. Our hypothesis prior to the study

initiation was that i.n. delivery of this saRNA vaccine would elicit superior respiratory immunity

in rodents, specifically eliciting long-lived memory T cells, while possibly driving systemic

immune responses, even if not at the level induced by standard i.m. delivery of the vaccine.

Furthermore, we believed that the i.n. delivered vaccine would therefore elicit protection from

morbidity in the hamster model and superior reduction in viral transmission relative to the i.m.

delivered vaccine. As the data developed, we further hypothesized that heterologous dosing of

the vaccine, with an i.m. prime and i.n. boost, would combine the best attributes of both vaccine

delivery methods and elicit a combination of systemic and respiratory immune responses that

would optimally protect against viral challenge and transmission.

<u>Research subjects or units of investigation</u>

Research was conducted on C57BL/6J mice obtained from The Jackson Laboratory. Lakeview

Golden (LVG) Syrian hamsters were obtained from Charles River Laboratories.

<u>Experimental design</u>

We conducted several mouse vaccine immunogenicity experiments to address these questions,

vaccinating mice i.m and/or i.n. with our SARS-CoV-2 saRNA vaccine. Mice were vaccinated

using different doses and dosing strategies (homologous vs. heterologous prime-boost vaccine

regimens), and immunogenicity was assessed by ELISA and pseudoneutralization assay on

serum samples, T cell intracellular cytokine staining (ICS) on spleen and lung samples, bone

marrow B cell ELISpot, and T cell ELISpot to assess systemic antibody responses and induction

of specific vaccine-induced T cell and B cell populations.

<u>Randomization</u>

Rodents were obtained commercially and randomized into study groups prior to study onset.

<u>Blinding</u>

Investigators were not blinded to the study groups during data collection and analysis as this is

typically unnecessary for this type of inbred animal study and introduces more systemic and

human error than it prevents. Quantitative, unbiased assays were developed with highly regulated

SOPs in order to minimize data biases.

<u>Sample size</u>

Statistical analyses have been the guiding strategy to maximize statistical power while reducing

the number of animals used to a minimum. We have previously determined that 6-8 animals per

group (3-4 male, 3-4 female) is the lowest estimate to achieve meaningful comparisons between

vaccine candidates with subtle differences in immunogenicity. 10 mice/group, 5 male and 5

female, is required to provide statistical power of 90% with an alpha (p-value) of 0.05 to detect a

statistically significant difference in pseudovirus neutralizing antibody titers between vaccinated

groups with two-fold differences in mean antibody titers with expected statistical variance. We

use both male and female mice in most study groups to avoid sex-driven data biases and allow

for identification of any sex-specific differences in vaccine immunogenicity. For one select study

where mouse numbers were limited (Figure 5), $n = 5$ all-female mice were used per group to

provide statistical power of 80% to detect two-fold differences in serum antibody titers. The use

of all female mice in this study was conducted in order to reduce data variability sufficiently to

minimize animal use without compromising the main study immunogenicity readouts.

Data inclusion/exclusion criteria

No data were excluded from any plots or analyses except two cases: (1) for flow cytometry after

ICS where <50,000 cells were collected in the well (this was established as a guideline

prospectively) and (2) one i.n. vaccinated hamster was removed from Figure 6 where the

complete lack of anti-spike IgG antibody established that it had not been effectively vaccinated.

Outliers

Our policy prior to study initiation was not to exclude any outliers. Therefore, we have not

excluded any outliers. Assays were repeated on any suspected outliers in order to verify data

prior to inclusion.

Replicates

Replication of data was conducted whenever possible. All experiments were conducted with a

minimum of biological triplicates. ELISAs and pseudoneutralization assays were conducted with

technical duplicates for each study sample. T cell immunogenicity was only assessed in technical

singlicate due to limiting tissue quantity and cell number. Bridging groups were used between

independent animal studies to provide replication of key study groups across multiple studies

offset by months. All attempts at replication verified the robustness of the scientific approach.

saRNA expression plasmid design, cloning, and production

The saRNA plasmid was designed and selected based on immunogenicity measures from i.m.

immunization (*11*). Briefly, the saRNA encodes a SARS-CoV-2 spike sequence based on a

GenBank sequence (MT246667.1) containing the D614G mutation, a diproline at sites 987-988,

and a QQAQ substitution for RRAR in the furin cleavage site (683-686). The sequence was

codon-optimized for expression in humans, synthesized by BioXp, and inserted via Gibson

cloning into AAHI's backbone saRNA expression vector. A SEAP-expressing plasmid was also

used as a control, created similarly with the SEAP sequence in the place of the vaccine antigen.

Sanger sequencing was used to confirm plasmid sequences. These were then amplified in

Escherichia coli and extracted using Qiagen maxi- or gigaprep kits, linearized with NotI (New

England Biolabs), and purified using a Qiagen DNA purification kit.

RNA manufacture

NotI-linearized DNA plasmids containing T7 promoters were used as templates for *in vitro*

transcription (IVT) of saRNA for vaccination. Our in-house optimized IVT protocol with T7

polymerase, RNase inhibitor, and pyrophosphatase (Aldevron) was used. Next, a Dnase step was

used to digest the templates, and Cap0 structures were added to the RNA transcripts with

guanylyltransferase (Aldevron), GTP, and S-adenosylmethionine (New England Biolabs).

CaptoCore 700 resin (GE Healthcare) was used to chromatographically purify the RNA followed

by diafiltration and concentration. saRNA material was filtered through a 0.22-μm

polyethersulfone filter and stored at −80°C until further use. saRNA size and integrity was

characterized by gel electrophoresis, and RNA concentration was quantified by UV absorbance

(NanoDrop 1000) and RiboGreen assay (Thermo Fisher).

NLC manufacture

A mixture of trimyristin (IOI Oleochemical), squalene (Sigma-Aldrich), sorbitan monostearate

(Sigma-Aldrich), and the cationic lipid 1,2-dioleoyl-3-trimethylammonium-propane (DOTAP;

Corden) was heated at 70°C in a bath sonicator. Separately, polysorbate 80 (Fisher Scientific)

was diluted in 10 mM sodium citrate trihydrate and heated to 70°C in a bath sonicator. After all

components were in suspension, a high-speed laboratory emulsifier (Silverson Machines)

running at 7,000 rpm was used to blend the oil and aqueous phases. Particle size of the mixture

was further decreased through high-shear homogenization. The colloid mixture was then

processed at 30,000 psi for 11 discrete microfluidization passes using an M-110P microfluidizer

(Microfluidics). The NLC product was then filtered through a 0.22-μm polyethersulfone filter

and stored at 2°C–8°C until use (*11*, *17*, *80*).

Vaccine complexing and characterization

Vaccine complexes were produced by combining aqueous RNA at a 1:1 volume ratio with NLC

that was pre-diluted in a 10 mM sodium citrate and 20% w/v sucrose buffer. Vaccines were

prepared at a nitrogen:phosphate (N:P) ratio of 5-15, representing the ratio of amine groups on

the NLC DOTAP to the RNA's backbone phosphate groups. This complexing resulted in vaccine

containing the intended dose of complexed saRNA-NLC in an isotonic 10% w/v sucrose, 5 mM

sodium citrate solution. The vaccine solution was incubated on ice for 30 minutes after mixing to

ensure complete complexing.

Mouse studies

All mouse work was done according to the Bloodworks Northwest Research Institute's

Institutional Animal Care and Use Committee. All mouse work was in compliance with all

applicable sections of the Final Rules of the Animal Welfare Act regulations (9 CFR Parts 1, 2,

and 3) and the *Guide for the Care and Use of Laboratory Animals* (*81*).

C57BL/6J mice purchased from The Jackson Laboratory were used for all mouse studies. Mice

were between 6 and 8 weeks of age at study onset and evenly split between male and female. At

arrival, mice were simultaneously randomized into study groups, half male and half female,

housed separately 3-5 mice per cage with irradiated bedding in Allentown individually ventilated

cages. Sterile water gel packs were given to all newly arrived animals during the acclimation

period, and animals had access to fresh potable water *ad libitum* via Edstrom automatic reverse-

osmosis and chlorinated water system. Cages were changed every 2 weeks inside a laminar flow

cage change station. A spot check for dirty cages was conducted on the non-change weeks.

Mice were observed daily for 3 days post-injection after both prime and boost injections by

trained personnel. Isoflurane delivered by inhalation was used for temporary anesthesia

immediately prior to retro-orbital blood collection and i.n. immunization to mitigate animal pain

and distress. At study completion, animals were humanely euthanized by CO_2 overdose.

Animal handling technicians were not blinded to study groups. After 1 week of acclimatization

to the facility, mice were then primed with vaccine. Mice were then boosted 3 weeks later and

harvested 3 weeks after boost. Mouse weights were monitored for 4 days following prime and

boost vaccinations. Mice were immunized by i.m. injection in both rear quadriceps muscles (50

µL/leg, 100 µL total) or by i.n. inoculation in two 25-µL doses. Intranasal inoculation was done

under full anesthesia, and mice were allowed to recover under a heat lamp. During ongoing

studies, serum samples were taken by retro-orbital bleed. Three-weeks post-boost, mice were

harvested, terminal bleeds were taken, spleens and lungs were dissected, and femurs were

harvested for bone marrow purification.

Mouse serum IgG, IgG1, and IgG2a titers by ELISA

SARS-CoV-2 spike-specific IgG was measured by ELISA. Plates (Corning 384-well high-

binding #CLS3700) were coated with Recombinant SARS-Cov-2 Spike His Protein, Carrier Free

(R&D Systems #10549-CV), at 1 µg/mL and incubated overnight at 4°C. Plates were incubated

in blocking buffer (2% dry milk, 0.05% Tween 20, 1% goat serum) for over 1 hour. Samples

were plated on low-binding plates, starting at a dilution of 1:40 and then serially diluting 1:2

across the plate. High-binding plates were washed, and samples were then transferred from the

low-binding plates to the coated, high-binding plates. Naive mice serum at 1:40 was used as a

negative control. A SARS-Cov-2 neutralizing monoclonal antibody (GenScript #A02057) was

used as a positive control at a starting concentration of 3.2 ng/µL. Following a 1-hour incubation,

plates were washed, and a secondary antibody, Anti-Mouse IgG (Fc Specific)-Alkaline

Phosphatase antibody (Sigma-Aldrich #A2429), at a 1:4000 dilution was added. Following the

secondary antibody and another wash step, phosphatase substrate tablets (Sigma-Aldrich

#S0942) were dissolved in diethanolamine substrate buffer (Fisher Scientific #PI34064) at a final

concentration of 1 mg/mL and added to the plates. Plates were incubated for 30 minutes and then

read spectrophotometrically at 405 nm. A 4-point logistic (4PL) curve was used to fit the

antibody standard titration curve, and sample concentrations were interpolated off the linear

region of the curve.

For IgG1 and IgG2a isotype-specific ELISAs, plates were coated and blocked as described

above. For IgG1 and IgG2a, the standard curve was run using SARS-CoV-2 neutralizing

antibodies (GenScript #A02055 and #BS-M0220, respectively). Sample dilution and incubation

were identical to the total IgG curve, and plates were probed with IgG1- and IgG2a-specific

secondary alkaline phosphatase (AP)-conjugated detection antibodies (Sigma-Aldrich

#SAB3701172 and #SAB3701179, respectively) prior to development, reading, and

quantification as described above.

Pseudovirus neutralization assay

SARS-CoV-2 pseudovirus neutralizing antibody titers in mouse sera were measured via a

pseudoneutralization assay (*82*). Lentiviral SARS-CoV-2 spike protein pseudotyped particles

were prepared by co-transfecting HEK-293 cells (American Type Culture Collection CRL

#11268) with plasmids containing a lentiviral backbone-expressing luciferase and ZsGreen (BEI

Resources #NR-52516), lentiviral helper genes (BEI Resources #NR-52517, NR-52518, and NR-

52519), or a delta19 cytoplasmic tail-truncated SARS-CoV-2 spike protein (Wuhan strain

plasmids from Jesse Bloom of Fred Hutchinson Cancer Center). Cells were incubated for 72

hours at standard cell culture conditions (37°C, 5% CO_2), and cell culture media were harvested

and filtered through a 0.2-μm filter. Pseudovirus-containing supernatant was frozen until titering

and use.

To perform the assay, plates were seeded with Human Angiotensin-Converting Enzyme 2

(hACE2)-expressing HEK-293 cells (BEI Resources #NR52511) and incubated overnight.

Serum samples were pre-diluted 1:10 in media (Gibco Dulbecco's Modified Eagle Medium +

GlutaMAX + 10% fetal bovine serum [FBS]) and then serially diluted 1:2 for 11 total dilutions.

These were then incubated with polybrene (Sigma-Aldrich #TR-1003-G) and pseudovirus for 1

hour. Then, the serum samples were added onto the hACE2 cells in duplicate and incubated at

37°C and 5% CO_2 for 72 hours. To determine 50% inhibitory concentration (IC_{50}) values, plates

were scanned on a fluorescent imager (Molecular Devices ImageXpress Pico Automated Cell

Imaging System) for ZsGreen expression. Total integrated intensity per well was used to

calculate the percent of pseudovirus inhibition per well. Neutralization data for each sample were

fit with a 4-parameter sigmoidal curve, which was used to interpolate IC_{50} values.

Spleen, lung, and bone marrow cell harvest

Spleens were prepared in 4 mL of RPMI medium by manual maceration through a filter using

the back of a syringe. Dissociated splenocyte samples were centrifuged briefly at 400 x g to

pellet fat cells, and the supernatants containing lymphocytes were either transferred to 5-mL

mesh-cap tubes to strain out any remining tissue debris or lysed with ammonium-chloride-

potassium (ACK) lysing buffer. Cells were counted on a Guava easyCyte cytometer (Luminex)

and plated in round-bottom 96-well plates at 1-2 x 10^6 cells per well in complete RPMI (cRPMI)

medium containing CD28 costimulatory antibody (BD Biosciences #553294) and brefeldin A.

One of three stimulation treatments was added to each well: 0.0475% dimethyl sulphoxide

[DMSO], 2 μg/μL spike peptide pool (JPT peptides #PM-WCPV-S-1), or phorbol myristate

acetate (PMA)-ionomycin. Plates were incubated for 6 hours at 37 C with 5% CO_2.

Lung cells were isolated via enzymatic digestion using a gentleMACS Dissociator (Miltenyi

Biotec). Lungs were dissociated in 4 mL of Hanks' Balanced Salt Solution (HBSS)

supplemented with 10% Liberase (MilliporeSigma), 10% aminoguanidine, 0.1% KN-62, and

1.25% Dnase. Lungs and enzymatic mix were added to a gentleMACS M tube (Miltneyi Biotec),

and the m_lung_01.02 program was run. Samples were then incubated at 37°C and 5% CO_2 for

30 minutes. Directly after, samples were run again on the gentleMACS Dissociator using the

m_lung_02.01 program. The resulting slurry was added to 10 mL of RPMI medium and

centrifuged for 5 minutes. The supernatant was discarded, and the cells were either filtered

through a 5-mL snap-top tube or washed again in RPMI medium prior to counting on a Guava

easyCyte cytometer and plated in round-bottom 96-well plates at 1-2 x 10^6 cells per well in RPMI

medium containing 10% FBS, 50 μM beta-mercaptoethanol, CD28 costimulatory antibody,

brefeldin A, and one of three stimulation treatments: 0.0475% DMSO, 2 μg/μL spike peptide

pool (JPT peptides #PM-WCPV-S-1), or PMA-ionomycin. Plates were incubated for 6 hours at

37 C with 5% CO_2.

Intracellular cytokine staining and flow cytometry

Following incubation, plates were centrifuged at 400 x g for 3 minutes, the supernatants were

removed, and cells were resuspended in phosphate-buffered saline (PBS). Splenocytes were

stained for viability with Zombie Green (Biolegend) in 50 μL of PBS. Cells were washed twice,

and then spleen samples were incubated with CD16/CD32 antibody (Invitrogen #14-0161-86) to

block Fc receptors. Next, the cells were surface stained in an additional 50 μL of staining buffer

(PBS with 0.5% bovine serum albumin and 0.1% sodium azide). Spleen samples were stained

with fluorochrome-labeled monoclonal antibody (mAb) specific for mouse CD4 (eBioscience

#45-0042-82), CD8 (BD Biosciences #563068), CD44 (BD Biosciences #560568), and CD107a

(BioLegend #121614), while lung samples were stained for CD4 (BioLegend #100526), CD8

(BD Biosciences #563068), CD44 (BD Biosciences #562464), CD69 (BioLegend #104508),

CD103 (BioLegend #121435), CD194 (BioLegend #131220), and CD154 (BD Biosciences

#745242). Cells were washed twice, permeabilized using the Fixation/Permeabilization Kit (BD

Biosciences #554714), and stained intracellularly. Spleens were stained for TNFα (BioLegend

#506327), IL-2 (BioLegend #503824), IFNγ (Invitrogen #25731182), IL-5 (eBioscience #12-

7052-82), IL-10 (BD Biosciences #564081), and IL-17a (BD Biosciences #560820), while lungs

were stained for TNFα (BioLegend #506327), IL-2 (BioLegend #503824), IFNγ (Invitrogen

#25731182), IL-5 (BioLegend #504306), and IL-17a (BD Biosciences #560820). After two

washes in staining buffer, cells were resuspended in 100 μL of staining buffer and analyzed

using an LSRFortessa flow cytometer (BD Biosciences). After initial gating for live CD4$^+$ or

CD8$^+$ lymphocytes, cells gating positive for all three activation markers IL-2, IFNγ, and TNFα

were counted as activated polyfunctional T cells.

Bone marrow harvest and ASC ELISpot

Antibody-secreting bone marrow-resident cell counts were measured by ELISpot.

MultiScreen$_{HTS}$ IP Filter plates (0.45 μm, MilliporeSigma) were treated with 15 μL of 35%

ethanol for 30 seconds. Recombinant SARS-CoV-2 Spike His Protein, Carrier Free (R&D

Systems #10549-CV-100), was diluted at 2 µg/mL in ELISpot coating buffer (eBioscience), and

plates were coated with 100 µL. Plates were incubated overnight at 4°C. The next day, plates

were washed with PBS with 0.1% Tween 20 and blocked with cRPMI medium for 2 hours.

Femurs were removed from mice and inserted into a snipped-end 0.6-mL Eppendorf tube and

then inserted into a 1.5-mL Eppendorf tube with 1 mL of cRPMI medium. Femurs were

centrifuged for 15 seconds at 10,000 rpm, and supernatant was discarded. Cell pellets were

vortexed, resuspended in 200 µL of RBC Lysis Buffer (Invitrogen), and then incubated on ice for

30 seconds. An additional 800 µL of RPMI medium was added, and cells were centrifuged for 5

minutes at 400 x g before the supernatant was decanted. Cells were resuspended in 1 mL of

cRPMI medium, counted, and transferred to prepared filter plates at 1 million cells per well

followed by a 3-fold dilution across five adjacent wells.

Plates were incubated for 3 hours, then washed three times with PBS plus 0.1% Tween 20.

Secondary antibody (Goat Anti-Mouse IgG-HRP or IgA-HRP [SouthernBiotech #1030-05 and

#1040-05, respectively]) was added at a 1:1000 dilution in PBS with 0.1% Tween and 5% FBS

overnight at 4°C. Plates were then washed three times in PBS with 0.1% Tween 20 and two

times in PBS. 100 µL of Vector NovaRED Substrate Peroxidase (Vector Laboratories #SK-

4800) was applied for 7 minutes to develop the plates. The reaction was quenched by rinsing

plates with distilled water for 2 minutes, and plates were dried in the dark. Spots were counted

and data were analyzed using ImmunoSpot v7 software (Cellular Technology Limited).

T cell ELISpot assay

IFNγ (BD Biosciences #51-2525KZ), IL-17A (Invitrogen #88-7371-88), or IL-5 (BD

Biosciences #51-1805KZ) at a 1:200 dilution in Dulbecco's PBS (DPBS; Gibco) was used to

coat ELISpot plates (MilliporeSigma). Plates were incubated overnight at 4°C and then washed

and blocked with cRPMI medium for at least 2 hours. Previously harvested splenocytes (see

above) were added at 2 x 10^5 cells per well. Samples were then stimulated with 1 µg/mL PepMix

SARS-CoV-2 (JPT Peptide Technologies #PM-WCPV-S-1). Plates were incubated at 37°C and

5% CO_2 for 48 hours. Then plates were washed with PBS with 0.1% Tween 20. Next, detection

antibody (IFNγ, BD Biosciences #51-1818KA; IL-17A, Invitrogen #88-7371-88; and IL-5, BD

Biosciences #51-1806KZ) was diluted in ELISpot diluent (eBiosciences) at 1:250 and added

overnight at 4°C. Plates were washed and then developed using Vector NovaRED Substrate

Peroxidase for 15 minutes. The reaction was quenched with deionized water. Plates were then

dried in the dark. Spots were counted and data were analyzed using ImmunoSpot.

Hamster studies

Nine-week-old male Lakeview Golden (LVG) Syrian hamsters were purchased from Charles River

Laboratories and maintained in the University of Alabama at Birmingham (UAB) animal facilities,

which include the UAB Southeastern Biosafety Lab (SEBLAB) ABSL3 facilities. All hamster

procedures were approved by the UAB Institutional Animal Care and Use Committee (IACUC

protocol 22628) and the UAB Biosafety Committee (Protocol 22-160 and 22-161). All procedures

were performed in accordance with National Resource Council guidelines.

Hamster vaccination and infection

Adult male hamsters were housed in pairs and separated at the time of vaccination. One hamster

from each pair was vaccinated with 5 mg of saRNA-NLC delivered i.m. (50 mL per rear quadriceps

muscle, 100 mL total) or i.n. (25 mL/nare, 50 mL total). Sham-vaccinated hamsters received dPBS

i.m. Vaccinated and sham-vaccinated animals were re-paired with naive cagemates after 24 hours.

Cagemates were separated on day 28, and vaccinated animals were boosted with the same vaccine

given via the i.m. or i.n. route. Animals were re-paired after 24 hours. Nine days after the boost,

animals were separated, and the vaccinated and sham-vaccinated hamsters were infected i.n. with

1×10^6 pfu SARS-CoV-2 USA/WA-1 variant (50 mL/nare, 100 mL total). Uninfected controls

received 100 mL (50 mL/nare) of dPBS (vehicle). The infected animals were re-paired with the

uninfected, unvaccinated naive cagemates at 48 hours post-infection. Cagemates were separated

24 hours later and remained apart until the end of the study. Hamster weights were measured before

infection and daily until day 15 post-infection.

Hamster blood collection and processing

Blood samples from sedated hamsters were collected from the lateral saphenous vein on day 21

post-prime and 5 days post-boost vaccinations. Blood samples following euthanasia were collected

from terminal bleed via the posterior vena cava. Blood samples were transferred into BD

Microtainer blood collection tubes (BD Biosciences), and serum was separated by centrifugation

at 10,000 rpm at RT for 10 minutes. Serum samples collected post-infection were heated at 60°C

for 20 minutes to inactivate SARS-CoV-2 virus. All samples were aliquoted and frozen at -80°C

until analyzed by cytometric bead array (CBA).

Hamster nasal viral load samples

Nasal swipes were collected immediately prior to i.n. infection to establish baseline, at 1-hour

post-infection, and then daily until day 10 post-infection. Awake hamsters were transferred into

an open container to minimize movement, and the exterior surface of their nose was swiped for 5-

10 seconds with a polyester-tipped swab (Medical Packaging #SP-7D) that had been dipped in a

screw-cap tube containing 300 mL of viral transport medium (VTM; HBSS ($+Ca^{2+}$ $+Mg^{2+}$)

containing 2% FBS, 100 mg/mL gentamicin, and 0.5 mg/mL amphotericin B). The tip of the swab

was cut and placed inside the VTM-containing tube. The tubes were vortexed, and swab tips were

discarded. Samples were then processed for viral RNA quantitation.

Hamster nasal wash samples

Hamsters were sedated with isoflurane and euthanized with an intraperitoneal injection of

tribromoethanol (500 mg/kg, 3 mL/animal). The trachea was exposed, an incision was made, and

a blunt 19-gauge needle attached to a 1-mL insulin syringe was inserted into the trachea. While

holding the mouth of the hamster shut with one hand, 400 mL of VTM was squeezed through the

nasal passages, out of the nose, and into a screw-cap tube. Samples were heated at 60°C for 20

minutes to inactivate SARS-CoV-2 virus, then aliquoted, and stored at -80°C until analyzed by

CBA.

Recombinant SARS-CoV-2 spike, RBD, and N protein production

Recombinant SARS-CoV-2 spike ectodomain trimeric proteins were prepared as previously

described (*83*). Briefly, two human codon-optimized constructs were generated with a human IgG

leader sequence, the SARS-CoV-2 spike ectodomain (amino acids 14-1211), a GGSG linker, T4

fibritin foldon sequence, a GS linker, and a 15 amino acid biotinylation consensus site (AviTag,

construct 1) or 6X-HisTag (construct 2). Each construct was engineered with two sets of mutations

to stabilize the protein in a pre-fusion conformation (*83*). Recombinant SARS-CoV-2 spike RBD

domain monomers (human codon-optimized) were generated with a human IgG leader sequence,

the spike RBD (amino acids 319-541 from SARS-CoV-2 Wuhan-Hu-1 strain), a GS linker, an

AviTag, and a 6X-HisTag. The coding sequence of N from SARS-CoV-2 Wuhan-Hu-1 strain was

synthesized in frame with the coding sequence for an AviTag and was cloned in frame to the 6X-

HisTag in the pTrcHis2C expression vector (Invitrogen). Biotinylated recombinant N protein was

produced by co-transforming Rosetta cells with the N expression plasmid and an inducible BirA

expression plasmid. Bacteria were grown in the presence of chloramphenicol, ampicillin, and

streptomycin, induced with IPTG, and supplemented with biotin. Recombinant SARS-CoV-2

Avi/His-tagged spike trimeric proteins were produced by co-transfecting each plasmid construct

(1AVI:2HIS construct ratio) into FreeStyle 293-F cells. The Avi-His-tagged spike RBD protein

was produced by transfecting the single dual-tagged RBD construct into FreeStyle 293-F cells. All

recombinant proteins were purified by FPLC using a nickel-affinity column and subsequent size

exclusion chromatography. After buffer exchange, purified spike ectodomain trimers and RBD

domain protein were biotinylated by the addition of biotin-protein ligase (Avidity). Biotinylated

proteins were buffer exchanged into PBS, sterile filtered, aliquoted, and stored at -80ºC until used.

1015

SARS-CoV-2 spike cytometric bead array

CBAs with recombinant SARS-CoV-2 proteins were prepared as previously described (*83*).

Briefly, recombinant proteins were passively absorbed onto streptavidin functionalized 4-µm

fluorescent microparticles (Carboxy Blue Particle Array Kit, Spherotech). 500 µg of biotinylated

recombinant protein was incubated with 2 x 10^7 streptavidin functionalized fluorescent

microparticles in 400 µL of 1% BSA in PBS. Protein coupled beads were washed and resuspended

at 1 x 10^8 beads/mL and stored at 4°C.

1023

Cytometric bead array measurement of spike-specific IgG responses

50 µL of serum (diluted to 1/4000 in PBS) or nasal wash (diluted 1/4 in PBS) was arrayed in 96-

well u-bottom polystyrene plates. 5 µL of a suspension containing 5 x 10^5 spike beads, RBD beads,

and N beads was added to the samples. Suspensions were mixed, incubated for 15 minutes at RT,

washed in PBS, stained with FITC-conjugated polyclonal anti-hamster IgG (Southern Biotech) for

15 minutes at RT, washed in PBS, and then resuspended in 100 µL of 1% paraformaldehyde in

PBS. Samples were collected on a BD CytoFLEX flow cytometer in plate mode at a sample rate

of 100 µL/minute for 1 minute. Following acquisition, the FCS files were analyzed in FlowJo

(TreeStar). Briefly, the beads were identified by gating on singlet 4-µm particles in log scale in the

forward scatter and side scatter parameters. APC-Cy7 channel fluorescence gates were used to

segregate the particles by bead identity. Geometric mean fluorescent intensity was calculated in

the FITC channel.

1036

1037 *Propagation and titer determination of SARS-CoV-2*

1038 The original SARS-CoV-2 isolate USA-WA1/2020 was obtained from BEI Resources (#NR-

1039 52281), propagated in Vero E6 cells (ATCC #C1008), and titered as previously described (*84*).

1040

Viral RNA quantitation by qRT-PCR

SARS-CoV-2 viral N RNA and viral subgenomic N RNA was measured by PCR in nasal swipe

(300 µL) samples as previously described (*84*). AccuPlex SARS-CoV-2 Reference Material

(SeraCare Life Sciences, Inc., # 0505-0126) was extracted and amplified in parallel to generate a

standard curve enabling viral N quantitation. The subgenomic RNA standard was constructed

and validated in-house (*84*).

1047

Histopathology

Hamster lungs were inflated with 10% neutral-buffered formalin (Thermo Fisher) using a blunt

19-gauge needle. Each lung was excised *en bloc* and fixed for 7 days at RT in 15-fold volume of

10% neutral-buffered formalin. The fixed tissues were then embedded dorsal side down in

paraffin using standard procedures. Samples were sectioned at 5 µm, and resulting slides were

stained with hematoxylin and eosin (H&E). All tissue slides were evaluated by light microscopy

by a board-certified veterinary pathologist blinded to study group allocations. Representative

photo images were collected using a Nikon Eclipse Ci microscope (Nikon Inc) and analyzed with

NIS-Elements software (Nikon Inc). Scoring was performed following a published algorithm

(*85*, *86*). Briefly, the airway, alveolar, and vascular pathology in affected areas were assessed

and scored on a scale of 0-4 (*84*), and then added together to generate a total histopathological

score of the whole lung.

1060

1061 *Statistical analysis*

1062 The effect of dosing strategy on CD4 and CD8 cell polyfunctionality was assessed by one-way

1063 ANOVA with Tukey's multiple comparisons test. The effect of dosing strategy on IgG titer and

pseudovirus neutralizing antibody titers was assessed in log-transformed data using a mixed-

effects model with Tukey's multiple comparisons test. The effect of immunization strategy on

IgG titer was assessed on log-transformed data using unpaired t tests or two-way ANOVA with

Tukey's or Sidak's (if any sample data were missing) multiple comparisons test. The effect of

immunization strategy on pseudovirus neutralizing antibody titers was assessed on log-

transformed data using unpaired t tests or a mixed-effects model with Sidak's multiple

comparisons test. The effect of immunization strategy on polyfunctional CD4 and CD8 cell

responses was assessed in untransformed data using unpaired t tests or one-way ANOVA with

Holm-Sidak's multiple comparisons test. Lung, bone marrow, and splenocyte ELISpot data were

analyzed using log-transformed data by one-way ANOVA with Tukey's multiple comparisons

test. Hamster body weight data was analyzed using area under the curve measures assessed

statistically by one-way ANOVA with Tukey's multiple comparison test. Hamster viral load

measures were analyzed using log-transformed area under the curve measures, assessed with

one-way ANOVA with Tukey's multiple comparison test. All statistical analyses were conducted

using Prism version 9 (GraphPad Software).

References and Notes

1. V. Mouro, A. Fischer, Dealing with a mucosal viral pandemic: lessons from COVID-19
    vaccines. *Mucosal Immunol*, 1–11 (2022).

2. S. M. Kissler, J. R. Fauver, C. Mack, C. G. Tai, M. I. Breban, A. E. Watkins, R. M. Samant,
    D. J. Anderson, J. Metti, G. Khullar, R. Baits, M. MacKay, D. Salgado, T. Baker, J. T.
    Dudley, C. E. Mason, D. D. Ho, N. D. Grubaugh, Y. H. Grad, Viral dynamics of SARS-

1086  CoV-2 variants in vaccinated and unvaccinated persons. *New England Journal of Medicine*

1087  (2021), doi:10.1056/NEJMc2102507.

1088 3. M. Tiboni, L. Casettari, L. Illum, Nasal vaccination against SARS-CoV-2: synergistic or

1089  alternative to intramuscular vaccines? *Int J Pharm*. **603**, 120686 (2021).

1090 4. M. Garcia-Knight, K. Anglin, M. Tassetto, S. Lu, A. Zhang, S. A. Goldberg, A. Catching,

1091  M. C. Davidson, J. R. Shak, M. Romero, J. Pineda-Ramirez, R. Diaz-Sanchez, P. Rugart, K.

1092  Donohue, J. Massachi, H. M. Sans, M. Djomaleu, S. Mathur, V. Servellita, D. McIlwain, B.

1093  Gaudiliere, J. Chen, E. O. Martinez, J. M. Tavs, G. Bronstone, J. Weiss, J. T. Watson, M.

1094  Briggs-Hagen, G. R. Abedi, G. W. Rutherford, S. G. Deeks, C. Chiu, S. Saydah, M. J.

1095  Peluso, C. M. Midgley, J. N. Martin, R. Andino, J. D. Kelly, Infectious viral shedding of

1096  SARS-CoV-2 Delta following vaccination: A longitudinal cohort study. *PLOS Pathogens*.

1097  **18**, e1010802 (2022).

1098 5. J. Chen, R. Wang, N. B. Gilby, G.-W. Wei, Omicron variant (B.1.1.529): infectivity,

1099  vaccine breakthrough, and antibody resistance. *J. Chem. Inf. Model.* **62**, 412–422 (2022).

1100 6. D. R. Feikin, M. M. Higdon, L. J. Abu-Raddad, N. Andrews, R. Araos, Y. Goldberg, M. J.

1101  Groome, A. Huppert, K. L. O'Brien, P. G. Smith, A. Wilder-Smith, S. Zeger, M. D. Knoll,

1102  M. K. Patel, Duration of effectiveness of vaccines against SARS-CoV-2 infection and

1103  COVID-19 disease: results of a systematic review and meta-regression. *The Lancet*. **399**,

1104  924–944 (2022).

1105 7. R. Link-Gelles, M. E. Levy, M. Gaglani, S. A. Irving, M. Stockwell, K. Dascomb, M. B.

1106  DeSilva, S. E. Reese, I.-C. Liao, T. C. Ong, Effectiveness of 2, 3, and 4 COVID-19 mRNA

vaccine doses among immunocompetent adults during periods when SARS-CoV-2

Omicron BA.1 and BA.2/BA.2.12.1 sublineages predominated — VISION network, 10

states, December 2021–June 2022. *MMWR Morb Mortal Wkly Rep*. **71** (2022),

doi:10.15585/mmwr.mm7129e1.

8. D. Freeman, S. Lambe, L.-M. Yu, J. Freeman, A. Chadwick, C. Vaccari, F. Waite, L.

Rosebrock, A. Petit, S. Vanderslott, S. Lewandowsky, M. Larkin, S. Innocenti, H.

McShane, A. J. Pollard, B. S. Loe, Injection fears and COVID-19 vaccine hesitancy.

Psychol Med, 1–11.

9. S. Kar, P. Devnath, T. B. Emran, T. E. Tallei, S. Mitra, K. Dhama, Oral and intranasal

vaccines against SARS-CoV-2: current progress, prospects, advantages, and challenges.

Immunity, Inflammation and Disease. **10**, e604 (2022).

10. A. Rice, M. Verma, E. Voigt, P. Battisti, S. Beaver, S. Reed, K. Dinkins, S. Mody, L.

Zakin, S. Tanaka, B. Morimoto, C. A. Olson, E. Gabitzsch, J. T. Safrit, P. Spilman, C.

Casper, P. Soon-Shiong, Heterologous saRNA Prime, DNA Dual-Antigen Boost SARS-

CoV-2 Vaccination Elicits Robust Cellular Immunogenicity and Cross-Variant Neutralizing

Antibodies. *Front Immunol*. **13**, 910136 (2022).

11. E. A. Voigt, A. Gerhardt, D. Hanson, M. F. Jennewein, P. Battisti, S. Reed, J. Singh, R.

Mohamath, J. Bakken, S. Beaver, C. Press, P. Soon-Shiong, C. J. Paddon, C. B. Fox, C.

Casper, A self-amplifying RNA vaccine against COVID-19 with long-term room-

temperature stability. *npj Vaccines*. **7**, 1–13 (2022).

12.   Y. Peng, A. J. Mentzer, G. Liu, X. Yao, Z. Yin, D. Dong, W. Dejnirattisai, T. Rostron, P.

1128   Supasa, C. Liu, C. López-Camacho, J. Slon-Campos, Y. Zhao, D. I. Stuart, G. C. Paesen, J.

1129   M. Grimes, A. A. Antson, O. W. Bayfield, D. E. D. P. Hawkins, D.-S. Ker, B. Wang, L.

1130   Turtle, K. Subramaniam, P. Thomson, P. Zhang, C. Dold, J. Ratcliff, P. Simmonds, T. de

1131   Silva, P. Sopp, D. Wellington, U. Rajapaksa, Y.-L. Chen, M. Salio, G. Napolitani, W. Paes,

1132   P. Borrow, B. M. Kessler, J. W. Fry, N. F. Schwabe, M. G. Semple, J. K. Baillie, S. C.

1133   Moore, P. J. M. Openshaw, M. A. Ansari, S. Dunachie, E. Barnes, J. Frater, G. Kerr, P.

1134   Goulder, T. Lockett, R. Levin, Y. Zhang, R. Jing, L.-P. Ho, Oxford Immunology Network

1135   Covid-19 Response T cell Consortium, E. Barnes, D. Dong, T. Dong, S. Dunachie, J.

1136   Frater, P. Goulder, G. Kerr, P. Klenerman, G. Liu, A. McMichael, G. Napolitani, G. Ogg,

1137   Y. Peng, M. Salio, X. Yao, Z. Yin, ISARIC4C Investigators, J. Kenneth Baillie, P.

1138   Klenerman, A. J. Mentzer, S. C. Moore, P. J. M. Openshaw, M. G. Semple, D. I. Stuart, L.

1139   Turtle, R. J. Cornall, C. P. Conlon, P. Klenerman, G. R. Screaton, J. Mongkolsapaya, A.

1140   McMichael, J. C. Knight, G. Ogg, T. Dong, Broad and strong memory CD4+ and CD8+ T

1141   cells induced by SARS-CoV-2 in UK convalescent individuals following COVID-19. *Nat*

1142   *Immunol*. **21**, 1336–1345 (2020).

1143   13.   C. Rydyznski Moderbacher, S. I. Ramirez, J. M. Dan, A. Grifoni, K. M. Hastie, D.

1144   Weiskopf, S. Belanger, R. K. Abbott, C. Kim, J. Choi, Y. Kato, E. G. Crotty, C. Kim, S. A.

1145   Rawlings, J. Mateus, L. P. V. Tse, A. Frazier, R. Baric, B. Peters, J. Greenbaum, E.

1146   Ollmann Saphire, D. M. Smith, A. Sette, S. Crotty, Antigen-specific adaptive immunity to

1147   SARS-COV-2 in acute COVID-19 and associations with age and disease severity. *Cell*.

1148   **183**, 996-1012.e19 (2020).

1149   14.   Y. Sui, J. Li, H. Andersen, R. Zhang, S. K. Prabhu, T. Hoang, D. Venzon, A. Cook, R.

1150   Brown, E. Teow, J. Velasco, L. Pessaint, I. N. Moore, L. Lagenaur, J. Talton, M. W. Breed,

1151   J. Kramer, K. W. Bock, M. Minai, B. M. Nagata, H. Choo-Wosoba, M. G. Lewis, L.-X.

1152   Wang, J. A. Berzofsky, An intranasally administered SARS-CoV-2 beta variant subunit

1153   booster vaccine prevents beta variant replication in rhesus macaques. *PNAS Nexus*. **1**,

1154   pgac091 (2022).

1155   15.   P. F. McKay, K. Hu, A. K. Blakney, K. Samnuan, J. C. Brown, R. Penn, J. Zhou, C. R.

1156   Bouton, P. Rogers, K. Polra, P. J. C. Lin, C. Barbosa, Y. K. Tam, W. S. Barclay, R. J.

1157   Shattock, Self-amplifying RNA SARS-CoV-2 lipid nanoparticle vaccine candidate induces

1158   high neutralizing antibody titers in mice. *Nat Commun*. **11**, 3523 (2020).

1159   16.   R. de Alwis, E. S. Gan, S. Chen, Y. S. Leong, H. C. Tan, S. L. Zhang, C. Yau, J. G. H.

1160   Low, S. Kalimuddin, D. Matsuda, E. C. Allen, P. Hartman, K. J. Park, M. Alayyoubi, H.

1161   Bhaskaran, A. Dukanovic, Y. Bao, B. Clemente, J. Vega, S. Roberts, J. A. Gonzalez, M.

1162   Sablad, R. Yelin, W. Taylor, K. Tachikawa, S. Parker, P. Karmali, J. Davis, B. M. Sullivan,

1163   S. M. Sullivan, S. G. Hughes, P. Chivukula, E. E. Ooi, A single dose of self-transcribing

1164   and replicating RNA-based SARS-CoV-2 vaccine produces protective adaptive immunity

1165   in mice. *Mol Ther*. **29**, 1970–1983 (2021).

1166   17.   A. Gerhardt, E. Voigt, M. Archer, S. Reed, E. Larson, N. Van Hoeven, R. Kramer, C. Fox,

1167   C. Casper, A flexible, thermostable nanostructured lipid carrier platform for RNA vaccine

1168   delivery. *Mol Ther Methods Clin Dev*. **25**, 205–214 (2022).

1169   18.   H. Xu, L. Cai, S. Hufnagel, Z. Cui, Intranasal vaccine: Factors to consider in research and

1170   development. *Int J Pharm*. **609**, 121180 (2021).

19. L. Feng, Q. Wang, C. Shan, C. Yang, Y. Feng, J. Wu, X. Liu, Y. Zhou, R. Jiang, P. Hu, X.

Liu, F. Zhang, P. Li, X. Niu, Y. Liu, X. Zheng, J. Luo, J. Sun, Y. Gu, B. Liu, Y. Xu, C. Li,

W. Pan, J. Zhao, C. Ke, X. Chen, T. Xu, N. Zhong, S. Guan, Z. Yuan, L. Chen, An

adenovirus-vectored COVID-19 vaccine confers protection from SARS-COV-2 challenge

in rhesus macaques. *Nat Commun*. **11**, 4207 (2020).

20. A. O. Hassan, N. M. Kafai, I. P. Dmitriev, J. M. Fox, B. K. Smith, I. B. Harvey, R. E. Chen,

E. S. Winkler, A. W. Wessel, J. B. Case, E. Kashentseva, B. T. McCune, A. L. Bailey, H.

Zhao, L. A. VanBlargan, Y.-N. Dai, M. Ma, L. J. Adams, S. Shrihari, J. E. Danis, L. E.

Gralinski, Y. J. Hou, A. Schäfer, A. S. Kim, S. P. Keeler, D. Weiskopf, R. S. Baric, M. J.

Holtzman, D. H. Fremont, D. T. Curiel, M. S. Diamond, A single-dose intranasal ChAd

vaccine protects upper and lower respiratory tracts against SARS-CoV-2. *Cell*. **183**, 169-

184.e13 (2020).

21. M.-W. Ku, M. Bourgine, P. Authié, J. Lopez, K. Nemirov, F. Moncoq, A. Noirat, B. Vesin,

F. Nevo, C. Blanc, P. Souque, H. Tabbal, E. Simon, D. Hardy, M. Le Dudal, F. Guinet, L.

Fiette, H. Mouquet, F. Anna, A. Martin, N. Escriou, L. Majlessi, P. Charneau, Intranasal

vaccination with a lentiviral vector protects against SARS-CoV-2 in preclinical animal

models. *Cell Host Microbe*. **29**, 236-249.e6 (2021).

22. N. van Doremalen, J. N. Purushotham, J. E. Schulz, M. G. Holbrook, T. Bushmaker, A.

Carmody, J. R. Port, C. K. Yinda, A. Okumura, G. Saturday, F. Amanat, F. Krammer, P.

W. Hanley, B. J. Smith, J. Lovaglio, S. L. Anzick, K. Barbian, C. Martens, S. C. Gilbert, T.

Lambe, V. J. Munster, Intranasal ChAdOx1 nCoV-19/AZD1222 vaccination reduces viral

shedding after SARS-CoV-2 D614G challenge in preclinical models. *Science Translational*

Medicine. **13**, eabh0755 (2021).

23.   S. Afkhami, M. R. D'Agostino, A. Zhang, H. D. Stacey, A. Marzok, A. Kang, R. Singh, J.

Bavananthasivam, G. Ye, X. Luo, F. Wang, J. C. Ang, A. Zganiacz, U. Sankar, N.

Kazhdan, J. F. E. Koenig, A. Phelps, S. F. Gameiro, S. Tang, M. Jordana, Y. Wan, K. L.

Mossman, M. Jeyanathan, A. Gillgrass, M. F. C. Medina, F. Smaill, B. D. Lichty, M. S.

Miller, Z. Xing, Respiratory mucosal delivery of next-generation COVID-19 vaccine

provides robust protection against both ancestral and variant strains of SARS-CoV-2. *Cell*.

185, 896-915.e19 (2022).

24.   M. Cokarić Brdovčak, J. Materljan, M. Šustić, S. Ravlić, T. Ružić, B. Lisnić, K. Miklić, I.

Brizić, M. Pribanić Matešić, V. Juranić Lisnić, B. Halassy, D. Rončević, Z. Knežević, L.

Štefan, F. Bertoglio, M. Schubert, L. Čičin-Šain, A. Markotić, S. Jonjić, A. Krmpotić,

ChAdOx1-S adenoviral vector vaccine applied intranasally elicits superior mucosal

immunity compared to the intramuscular route of vaccination. *European Journal of*

Immunology. **52**, 936–945 (2022).

25.   H. Lei, A. Alu, J. Yang, W. Ren, C. He, T. Lan, X. He, L. Yang, J. Li, Z. Wang, X. Song,

W. Wang, G. Lu, X. Wei, Intranasal administration of a recombinant RBD vaccine induces

long-term immunity against Omicron-included SARS-CoV-2 variants. *Sig Transduct*

Target Ther. **7**, 1–13 (2022).

26.   T. Y. Wong, K. S. Lee, B. P. Russ, A. M. Horspool, J. Kang, M. T. Winters, M. Allison

Wolf, N. A. Rader, O. A. Miller, M. Shiflett, J. Izac, D. Varisco, E. Sen-Kilic, C.

Cunningham, M. Cooper, H. A. Cyphert, M. Barbier, I. Martinez, J. R. Bevere, R. K. Ernst,

F. H. Damron, Intranasal administration of BReC-CoV-2 COVID-19 vaccine protects K18-

hACE2 mice against lethal SARS-CoV-2 challenge. *NPJ Vaccines*. **7**, 36 (2022).

27. S. R. Allie, J. E. Bradley, U. Mudunuru, M. D. Schultz, B. A. Graf, F. E. Lund, T. D.

Randall, The establishment of resident memory B cells in the lung requires local antigen

encounter. *Nat Immunol*. **20**, 97–108 (2019).

28. X. An, M. Martinez-Paniagua, A. Rezvan, S. R. Sefat, M. Fathi, S. Singh, S. Biswas, M.

Pourpak, C. Yee, X. Liu, N. Varadarajan, Single-dose intranasal vaccination elicits

systemic and mucosal immunity against SARS-CoV-2. *iScience*. **24**, 103037 (2021).

29. X. F. Li, Z. Cui, H. Fan, Q. Chen, L. Cao, H. Y. Qiu, N. N. Zhang, Y. P. Xu, R. R. Zhang,

C. Zhou, Q. Ye, Y. Q. Deng, Y. Guo, S. Qin, K. Fan, L. Wang, Z. Jia, Y. Cui, X. Wang, C.

F. Qin, A highly immunogenic live-attenuated vaccine candidate prevents SARS-CoV-2

infection and transmission in hamsters. *Innovation (N Y)*, 100221 (2022).

30. A. Ainai, S. Tamura, T. Suzuki, E. van Riet, R. Ito, T. Odagiri, M. Tashiro, T. Kurata, H.

Hasegawa, Intranasal vaccination with an inactivated whole influenza virus vaccine induces

strong antibody responses in serum and nasal mucus of healthy adults. *Hum Vaccin*

Immunother. **9**, 1962–70 (2013).

31. K. Dhama, M. Dhawan, R. Tiwari, T. B. Emran, S. Mitra, A. A. Rabaan, S. Alhumaid, Z.

A. Alawi, A. Al Mutair, COVID-19 intranasal vaccines: current progress, advantages,

prospects, and challenges. *Human Vaccines & Immunotherapeutics*, 1–11 (2022).

32. K. McMahan, J. Yu, N. B. Mercado, C. Loos, L. H. Tostanoski, A. Chandrashekar, J. Liu,

L. Peter, C. Atyeo, A. Zhu, E. A. Bondzie, G. Dagotto, M. S. Gebre, C. Jacob-Dolan, Z. Li,

F. Nampanya, S. Patel, L. Pessaint, A. Van Ry, K. Blade, J. Yalley-Ogunro, M. Cabus, R.

Brown, A. Cook, E. Teow, H. Andersen, M. G. Lewis, D. A. Lauffenburger, G. Alter, D. H.

Barouch, Correlates of protection against SARS-CoV-2 in rhesus macaques. *Nature*. **590**,

630–634 (2021).

33. P. B. Gilbert, D. C. Montefiori, A. B. McDermott, Y. Fong, D. Benkeser, W. Deng, H.

Zhou, C. R. Houchens, K. Martins, L. Jayashankar, F. Castellino, B. Flach, B. C. Lin, S.

O'Connell, C. McDanal, A. Eaton, M. Sarzotti-Kelsoe, Y. Lu, C. Yu, B. Borate, L. W. P.

van der Laan, N. S. Hejazi, C. Huynh, J. Miller, H. M. El Sahly, L. R. Baden, M. Baron, L.

De La Cruz, C. Gay, S. Kalams, C. F. Kelley, M. P. Andrasik, J. G. Kublin, L. Corey, K.

M. Neuzil, L. N. Carpp, R. Pajon, D. Follmann, R. O. Donis, R. A. Koup, Immune Assays

Team§, Moderna, Inc. Team§, Coronavirus Vaccine Prevention Network

(CoVPN)/Coronavirus Efficacy (COVE) Team§, United States Government (USG)/CoVPN

Biostatistics Team§, Immune correlates analysis of the mRNA-1273 COVID-19 vaccine

efficacy clinical trial. *Science*. **375**, 43–50 (2022).

34. B. S. Graham, Rapid COVID-19 vaccine development. *Science*. **368**, 945–946 (2020).

35. D. Weiskopf, K. S. Schmitz, M. P. Raadsen, A. Grifoni, N. M. A. Okba, H. Endeman, J. P.

C. van den Akker, R. Molenkamp, M. P. G. Koopmans, E. C. M. van Gorp, B. L.

Haagmans, R. L. de Swart, A. Sette, R. D. de Vries, Phenotype and kinetics of SARS-CoV-

2-specific T cells in COVID-19 patients with acute respiratory distress syndrome. *Sci

Immunol*. **5**, eabd2071 (2020).

36. L. A. Jackson, E. J. Anderson, N. G. Rouphael, P. C. Roberts, M. Makhene, R. N. Coler, M.

P. McCullough, J. D. Chappell, M. R. Denison, L. J. Stevens, A. J. Pruijssers, A.

1257 McDermott, B. Flach, N. A. Doria-Rose, K. S. Corbett, K. M. Morabito, S. O'Dell, S. D.

1258 Schmidt, P. A. Swanson, M. Padilla, J. R. Mascola, K. M. Neuzil, H. Bennett, W. Sun, E.

1259 Peters, M. Makowski, J. Albert, K. Cross, W. Buchanan, R. Pikaart-Tautges, J. E.

1260 Ledgerwood, B. S. Graham, J. H. Beigel, R. N. A. S. G. m, An mRNA vaccine against

1261 SARS-CoV-2 - preliminary report. *N Engl J Med*. **383**, 1920–1931 (2020).

1262 37. N. Lurie, M. Saville, R. Hatchett, J. Halton, Developing Covid-19 vaccines at pandemic

1263   speed. *N Engl J Med*. **382**, 1969–1973 (2020).

1264 38. R. Yuan, J. Yu, Z. Jiao, J. Li, F. Wu, R. Yan, X. Huang, C. Chen, The roles of tissue-

1265   resident memory T cells in lung diseases. *Frontiers in Immunology*. **12**, 710375 (2021).

1266 39. R. Elgueta, M. J. Benson, V. C. de Vries, A. Wasiuk, Y. Guo, R. J. Noelle, *Immunol Rev*, in

1267   press, doi:10.1111/j.1600-065X.2009.00782.x.

1268 40. A. Ballesteros-Tato, B. León, F. E. Lund, T. D. Randall, CD4+ T helper cells use CD154–

1269   CD40 interactions to counteract T reg cell–mediated suppression of CD8+ T cell responses

1270   to influenza. *Journal of Experimental Medicine*. **210**, 1591–1601 (2013).

1271 41. J. M. E. Lim, A. T. Tan, N. Le Bert, S. K. Hang, J. G. H. Low, A. Bertoletti, SARS-CoV-2

1272   breakthrough infection in vaccinees induces virus-specific nasal-resident CD8+ and CD4+

1273   T cells of broad specificity. *J Exp Med*. **219**, e20220780 (2022).

1274 42. Y.-J. Park, D. Pinto, A. C. Walls, Z. Liu, A. De Marco, F. Benigni, F. Zatta, C. Silacci-

1275   Fregni, J. Bassi, K. R. Sprouse, A. Addetia, J. E. Bowen, C. Stewart, M. Giurdanella, C.

1276   Saliba, B. Guarino, M. A. Schmid, N. Franko, J. Logue, H. V. Dang, K. Hauser, J. di Iulio,

1277   W. Rivera, G. Schnell, F. A. Lempp, J. Janer, R. Abdelnabi, P. Maes, P. Ferrari, A. Ceschi,

O. Giannini, G. D. de Melo, L. Kergoat, H. Bourhy, J. Neyts, L. Soriaga, L. A. Purcell, G.

Snell, S. P. J. Whelan, A. Lanzavecchia, H. W. Virgin, L. Piccoli, H. Chu, M. S. Pizzuto, D.

Corti, D. Veesler, *bioRxiv*, in press, doi:10.1101/2022.05.08.491108.

43.   J. Tang, C. Zeng, T. M. Cox, C. Li, Y. M. Son, I. S. Cheon, Y. Wu, S. Behl, J. J. Taylor, R.

Chakraborty, A. J. Johnson, D. N. Schiavo, J. P. Utz, J. S. Reisenauer, D. E. Midthun, J. J.

Mullon, E. S. Edell, M. G. Alameh, L. Borish, W. G. Teague, M. H. Kaplan, D. Weissman,

R. Kern, H. Hu, R. Vassallo, S.-L. Liu, J. Sun, Respiratory mucosal immunity against

SARS-CoV-2 following mRNA vaccination. *Science Immunology*. **0**, eadd4853 (2022).

44.   F. P. Polack, S. J. Thomas, N. Kitchin, J. Absalon, A. Gurtman, S. Lockhart, J. L. Perez, G.

Perez Marc, E. D. Moreira, C. Zerbini, R. Bailey, K. A. Swanson, S. Roychoudhury, K.

Koury, P. Li, W. V. Kalina, D. Cooper, R. W. Frenck, L. L. Hammitt, O. Tureci, H. Nell,

A. Schaefer, S. Unal, D. B. Tresnan, S. Mather, P. R. Dormitzer, U. Sahin, K. U. Jansen,

W. C. Gruber, C. Clinical Trial Group, Safety and efficacy of the BNT162b2 mRNA

Covid-19 vaccine. *N Engl J Med*. **383**, 2603–2615 (2020).

45.   L. R. Baden, H. M. El Sahly, B. Essink, K. Kotloff, S. Frey, R. Novak, D. Diemert, S. A.

Spector, N. Rouphael, C. B. Creech, J. McGettigan, S. Khetan, N. Segall, J. Solis, A. Brosz,

C. Fierro, H. Schwartz, K. Neuzil, L. Corey, P. Gilbert, H. Janes, D. Follmann, M.

Marovich, J. Mascola, L. Polakowski, J. Ledgerwood, B. S. Graham, H. Bennett, R. Pajon,

C. Knightly, B. Leav, W. Deng, H. Zhou, S. Han, M. Ivarsson, J. Miller, T. Zaks, Cove

Study Group, Efficacy and safety of the mRNA-1273 SARS-CoV-2 vaccine. *N Engl J Med*.

384, 403–416 (2021).

46. H. M. El Sahly, L. R. Baden, B. Essink, S. Doblecki-Lewis, J. M. Martin, E. J. Anderson,

T. B. Campbell, J. Clark, L. A. Jackson, C. J. Fichtenbaum, M. Zervos, B. Rankin, F. Eder,

G. Feldman, C. Kennelly, L. Han-Conrad, M. Levin, K. M. Neuzil, L. Corey, P. Gilbert, H.

Janes, D. Follmann, M. Marovich, L. Polakowski, J. R. Mascola, J. E. Ledgerwood, B. S.

Graham, A. August, H. Clouting, W. Deng, S. Han, B. Leav, D. Manzo, R. Pajon, F.

Schodel, J. E. Tomassini, H. Zhou, J. Miller, Cove Study Group, Efficacy of the mRNA-

1273 SARS-CoV-2 vaccine at completion of blinded phase. *N Engl J Med*. **385**, 1774–1785

(2021).

47. O. J. Watson, G. Barnsley, J. Toor, A. B. Hogan, P. Winskill, A. C. Ghani, Global impact

of the first year of COVID-19 vaccination: a mathematical modelling study. *The Lancet*

Infectious Diseases. **0** (2022), doi:10.1016/S1473-3099(22)00320-6.

48. G. Giannotta, N. Giannotta, mRNA COVID-19 vaccines and long-lived plasma cells: a

complicated relationship. *Vaccines (Basel)*. **9**, 1503 (2021).

49. F. J. Ibarrondo, C. Hofmann, J. A. Fulcher, D. Goodman-Meza, W. Mu, M. A. Hausner, A.

Ali, A. Balamurugan, E. Taus, J. Elliott, P. Krogstad, N. H. Tobin, K. G. Ferbas, S. G.

Kitchen, G. M. Aldrovandi, A. W. Rimoin, O. O. Yang, Primary, recall, and decay kinetics

of SARS-CoV-2 vaccine antibody responses. *ACS Nano*. **15**, 11180–11191 (2021).

50. J. Keehner, L. E. Horton, N. J. Binkin, L. C. Laurent, D. Pride, C. A. Longhurst, S. R.

Abeles, F. J. Torriani, Resurgence of SARS-CoV-2 infection in a highly vaccinated health

system workforce. *N Engl J Med*. **385**, 1330–1332 (2021).

51. C. Menni, A. May, L. Polidori, P. Louca, J. Wolf, J. Capdevila, C. Hu, S. Ourselin, C. J.

Steves, A. M. Valdes, T. D. Spector, COVID-19 vaccine waning and effectiveness and

side-effects of boosters: a prospective community study from the ZOE COVID Study. *The*

Lancet Infectious Diseases. **22**, 1002–1010 (2022).

52. S. Rele, COVID-19 vaccine development during pandemic: gap analysis, opportunities, and

impact on future emerging infectious disease development strategies. *Hum Vaccin*

Immunother. **17**, 1122–1127 (2021).

53. X. Ren, J. Zhou, J. Guo, C. Hao, M. Zheng, R. Zhang, Q. Huang, X. Yao, R. Li, Y. Jin,

Reinfection in patients with COVID-19: a systematic review. *Glob Health Res Policy*. **7**, 12

(2022).

54. Y. Araf, F. Akter, Y.-D. Tang, R. Fatemi, M. S. A. Parvez, C. Zheng, M. G. Hossain,

Omicron variant of SARS-CoV-2: Genomics, transmissibility, and responses to current

COVID-19 vaccines. *J Med Virol*. **94**, 1825–1832 (2022).

55. D. W. Eyre, D. Taylor, M. Purver, D. Chapman, T. Fowler, K. B. Pouwels, A. S. Walker, T.

E. A. Peto, Effect of Covid-19 Vaccination on Transmission of Alpha and Delta Variants.

New England Journal of Medicine. **386**, 744–756 (2022).

56. C. Franco-Paredes, Transmissibility of SARS-CoV-2 among fully vaccinated individuals.

The Lancet Infectious Diseases. **22**, 16 (2022).

57. A. Singanayagam, S. Hakki, J. Dunning, K. J. Madon, M. A. Crone, A. Koycheva, N.

Derqui-Fernandez, J. L. Barnett, M. G. Whitfield, R. Varro, A. Charlett, R. Kundu, J. Fenn,

J. Cutajar, V. Quinn, E. Conibear, W. Barclay, P. S. Freemont, G. P. Taylor, S. Ahmad, M.

Zambon, N. M. Ferguson, A. Lalvani, A. Badhan, S. Dustan, C. Tejpal, A. V. Ketkar, J. S. Narean, S. Hammett, E. McDermott, T. Pillay, H. Houston, C. Luca, J. Samuel, S. Bremang, S. Evetts, J. Poh, C. Anderson, D. Jackson, S. Miah, J. Ellis, A. Lackenby, Community transmission and viral load kinetics of the SARS-CoV-2 delta (B.1.617.2) variant in vaccinated and unvaccinated individuals in the UK: a prospective, longitudinal, cohort study. *The Lancet Infectious Diseases*. **22**, 183–195 (2022).

58. M. Cevik, N. D. Grubaugh, A. Iwasaki, P. Openshaw, COVID-19 vaccines: Keeping pace with SARS-CoV-2 variants. *Cell*. **184**, 5077–5081 (2021).

59. M. C. Maher, I. Bartha, S. Weaver, J. di Iulio, E. Ferri, L. Soriaga, F. A. Lempp, B. L. Hie, B. Bryson, B. Berger, D. L. Robertson, G. Snell, D. Corti, H. W. Virgin, S. L. Kosakovsky Pond, A. Telenti, Predicting the mutational drivers of future SARS-CoV-2 variants of concern. *Science Translational Medicine*. **14**, eabk3445 (2022).

60. C. Zhong, H. Xia, A. Adam, B. Wang, R. L. Hajnik, Y. Liang, G. H. Rafael, J. Zou, X. Wang, J. Sun, L. Soong, A. D. T. Barrett, S. C. Weaver, P.-Y. Shi, T. Wang, H. Hu, Mucosal vaccination induces protection against SARS-CoV-2 in the absence of detectable neutralizing antibodies. *NPJ Vaccines*. **6**, 139 (2021).

61. P. F. Wright, A. C. Prevost-Reilly, H. Natarajan, E. B. Brickley, R. I. Connor, W. F. Wieland-Alter, A. S. Miele, J. A. Weiner, R. D. Nerenz, M. E. Ackerman, Longitudinal Systemic and Mucosal Immune Responses to SARS-CoV-2 Infection. *The Journal of Infectious Diseases*, jiac065 (2022).

62. A. Alu, L. Chen, H. Lei, Y. Wei, X. Tian, X. Wei, Intranasal COVID-19 vaccines: From

bench to bed. *eBioMedicine*. **76** (2022), doi:10.1016/j.ebiom.2022.103841.

63. H. E. Mudrick, E. B. McGlinch, B. J. Parrett, J. R. Hemsath, M. E. Barry, J. D. Rubin, C.

Uzendu, M. J. Hansen, C. L. Erskine, V. P. VanKeulen, A. Drelich, C.-T. K. Tseng, S.

Massey, M. Fida, G. A. Suh, T. Peikert, M. S. Block, G. R. Olivier, M. A. Barry,

Comparison of Mucosal and Intramuscular Immunization against SARS-CoV-2 with

Replication-Defective and Replicating Single-cycle Adenovirus Vaccines (2021), p.

2021.04.20.440651, , doi:10.1101/2021.04.20.440651.

64. N. J. Carter, M. P. Curran, Live attenuated influenza vaccine (FluMist®; Fluenz™): a

review of its use in the prevention of seasonal influenza in children and adults. *Drugs*. **71**,

1591–1622 (2011).

65. S. R. Banihashemi, A. Es-haghi, M. H. Fallah Mehrabadi, M. Nofeli, A. R. Mokarram, A.

Ranjbar, M. Salman, M. Hajimoradi, S. H. Razaz, M. Taghdiri, M. Bagheri, M. Dadar, Z.

M. Hassan, M. Eslampanah, Z. Salehi Najafabadi, M. Lotfi, A. Khorasani, F. Rahmani,

Safety and Efficacy of Combined Intramuscular/Intranasal RAZI-COV PARS Vaccine

Candidate Against SARS-CoV-2: A Preclinical Study in Several Animal Models. *Front*

Immunol. **13**, 836745 (2022).

66. E. Waltz, China and India approve nasal COVID vaccines — are they a game changer?

Nature. **609**, 450–450 (2022).

1379 67. F. Xu, S. Wu, L. Yi, S. Peng, F. Wang, W. Si, L. Hou, T. Zhu, Safety, mucosal and

1380   systemic immunopotency of an aerosolized adenovirus-vectored vaccine against SARS-

1381   CoV-2 in rhesus macaques. *Emerging Microbes & Infections*. **11**, 439–442 (2022).

1382 68. B. Slütter, L. L. Pewe, S. M. Kaech, J. T. Harty, Lung Airway-Surveilling CXCR3hi

1383   Memory CD8+ T Cells Are Critical for Protection against Influenza A Virus. *Immunity*. **39**,

1384   939–948 (2013).

1385 69. A. Rice, M. Verma, A. Shin, L. Zakin, P. Sieling, S. Tanaka, J. Balint, K. Dinkins, H.

1386   Adisetiyo, B. Morimoto, W. Higashide, C. Anders Olson, S. Mody, P. Spilman, E.

1387   Gabitzsch, J. T. Safrit, S. Rabizadeh, K. Niazi, P. Soon-Shiong, Intranasal plus

1388   subcutaneous prime vaccination with a dual antigen COVID-19 vaccine elicits T-cell and

1389   antibody responses in mice. *Sci Rep*. **11**, 14917 (2021).

1390 70. T. Y. Wong, K. S. Lee, B. P. Russ, A. M. Horspool, J. Kang, M. T. Winters, M. Allison

1391   Wolf, N. A. Rader, O. A. Miller, M. Shiflett, J. Izac, D. Varisco, E. Sen-Kilic, C.

1392   Cunningham, M. Cooper, H. A. Cyphert, M. Barbier, I. Martinez, J. R. Bevere, R. K. Ernst,

1393   F. H. Damron, Intranasal administration of BReC-CoV-2 COVID-19 vaccine protects K18-

1394   hACE2 mice against lethal SARS-CoV-2 challenge. *NPJ Vaccines*. **7**, 36 (2022).

1395 71. K. S. Corbett, D. K. Edwards, S. R. Leist, O. M. Abiona, S. Boyoglu-Barnum, R. A.

1396   Gillespie, S. Himansu, A. Schäfer, C. T. Ziwawo, A. T. DiPiazza, K. H. Dinnon, S. M.

1397   Elbashir, C. A. Shaw, A. Woods, E. J. Fritch, D. R. Martinez, K. W. Bock, M. Minai, B. M.

1398   Nagata, G. B. Hutchinson, K. Wu, C. Henry, K. Bahl, D. Garcia-Dominguez, L. Ma, I.

1399   Renzi, W.-P. Kong, S. D. Schmidt, L. Wang, Y. Zhang, E. Phung, L. A. Chang, R. J.

1400   Loomis, N. E. Altaras, E. Narayanan, M. Metkar, V. Presnyak, C. Liu, M. K. Louder, W.

Shi, K. Leung, E. S. Yang, A. West, K. L. Gully, L. J. Stevens, N. Wang, D. Wrapp, N. A. Doria-Rose, G. Stewart-Jones, H. Bennett, G. S. Alvarado, M. C. Nason, T. J. Ruckwardt, J. S. McLellan, M. R. Denison, J. D. Chappell, I. N. Moore, K. M. Morabito, J. R. Mascola, R. S. Baric, A. Carfi, B. S. Graham, SARS-CoV-2 mRNA vaccine design enabled by prototype pathogen preparedness. *Nature*. **586**, 567–571 (2020).

72. P. Naaber, L. Tserel, K. Kangro, E. Sepp, V. Jürjenson, A. Adamson, L. Haljasmägi, A. P. Rumm, R. Maruste, J. Kärner, J. M. Gerhold, A. Planken, M. Ustav, K. Kisand, P. Peterson, Dynamics of antibody response to BNT162b2 vaccine after six months: a longitudinal prospective study. *The Lancet Regional Health - Europe*. **10**, 100208 (2021).

73. J. L. Americo, C. A. Cotter, P. L. Earl, R. Liu, B. Moss, Intranasal inoculation of an MVA-based vaccine induces IgA and protects the respiratory tract of hACE2 mice from SARS-CoV-2 infection. *Proc Natl Acad Sci U S A*. **119**, e2202069119 (2022).

74. J. R. Francica, B. J. Flynn, K. E. Foulds, A. T. Noe, A. P. Werner, I. N. Moore, M. Gagne, T. S. Johnston, C. Tucker, R. L. Davis, B. Flach, S. O'Connell, S. F. Andrew, E. Lamb, D. R. Flebbe, S. T. Nurmukhambetova, M. M. Donaldson, J.-P. M. Todd, A. L. Zhu, C. Atyeo, S. Fischinger, M. J. Gorman, S. Shin, V. V. Edara, K. Floyd, L. Lai, S. Boyoglu-Barnum, R. Van De Wetering, A. Tylor, E. McCarthy, V. Lecouturier, S. Ruiz, C. Berry, T. Tibbitts, H. Andersen, A. Cook, A. Dodson, L. Pessaint, A. Van Ry, M. Koutsoukos, C. Gutzeit, I.-T. Teng, T. Zhou, D. Li, B. F. Haynes, P. D. Kwong, A. McDermott, M. G. Lewis, T. M. Fu, R. Chicz, R. van der Most, K. S. Corbett, M. S. Suthar, G. Alter, M. Roederer, N. J. Sullivan, D. C. Douek, B. S. Graham, D. Casimiro, R. A. Seder, Protective antibodies

elicited by SARS-CoV-2 spike protein vaccination are boosted in the lung after challenge in

nonhuman primates. *Science Translational Medicine*. **13**, eabi4547 (2021).

75. T. Mao, B. Israelow, A. Suberi, L. Zhou, M. Reschke, M. A. Peña-Hernández, H. Dong, R.

J. Homer, W. M. Saltzman, A. Iwasaki, *bioRxiv*, in press, doi:10.1101/2022.01.24.477597.

76. ImmunityBio, Inc., "THEMBA II T-CELL Vaccine: A Phase 1/2 Study of the Safety,

Reactogenicity, and Immunogenicity of Vaccination With saRNA COVID-19 Vaccines"

(Clinical trial registration NCT05370040, clinicaltrials.gov, 2022), (available at

https://clinicaltrials.gov/ct2/show/NCT05370040).

77. J. Grimaud, V. N. Murthy, How to monitor breathing in laboratory rodents: a review of the

current methods. *Journal of Neurophysiology*. **120**, 624–632 (2018).

78. Y. M. Son, I. S. Cheon, Y. Wu, C. Li, Z. Wang, X. Gao, Y. Chen, Y. Takahashi, Y.-X. Fu,

A. L. Dent, M. H. Kaplan, J. J. Taylor, W. Cui, J. Sun, Tissue-resident CD4+ T helper cells

assist the development of protective respiratory B and CD8+ T cell memory responses. *Sci*

Immunol. **6**, eabb6852 (2021).

79. N. Swarnalekha, D. Schreiner, L. C. Litzler, S. Iftikhar, D. Kirchmeier, M. Künzli, Y. M.

Son, J. Sun, E. A. Moreira, C. G. King, T resident helper cells promote humoral responses

in the lung. *Sci Immunol*. **6**, eabb6808 (2021).

80. J. H. Erasmus, A. P. Khandhar, J. Guderian, B. Granger, J. Archer, M. Archer, E. Gage, J.

Fuerte-Stone, E. Larson, S. Lin, R. Kramer, R. N. Coler, C. B. Fox, D. T. Stinchcomb, S. G.

Reed, N. Van Hoeven, A Nanostructured Lipid Carrier for Delivery of a Replicating Viral

RNA Provides Single, Low-Dose Protection against Zika. *Mol Ther*. **26**, 2507–2522 (2018).

81. Committee for the Update of the Guide for the Care and Use of Laboratory Animals, *Guide*

for the Care and Use of Laboratory Animals (The National Academies Press, Washington,

DC, USA, Eighth., 2011).

82. K. H. D. Crawford, R. Eguia, A. S. Dingens, A. N. Loes, K. D. Malone, C. R. Wolf, H. Y.

Chu, M. A. Tortorici, D. Veesler, M. Murphy, D. Pettie, N. P. King, A. B. Balazs, J. D.

Bloom, Protocol and reagents for pseudotyping lentiviral particles with SARS-CoV-2 spike

protein for neutralization assays. *Viruses*. **12**, 513 (2020).

83. R. G. King, A. Silva-Sanchez, J. N. Peel, D. Botta, A. M. Dickson, A. K. Pinto, S. Meza-

Perez, S. R. Allie, M. D. Schultz, M. Liu, J. E. Bradley, S. Qiu, G. Yang, F. Zhou, E.

Zumaquero, T. S. Simpler, B. Mousseau, J. T. Killian, B. Dean, Q. Shang, J. L. Tipper, C.

A. Risley, K. S. Harrod, T. Feng, Y. Lee, B. Shiberu, V. Krishnan, I. Peguillet, J. Zhang, T.

J. Green, T. D. Randall, J. J. Suschak, B. Georges, J. D. Brien, F. E. Lund, M. S. Roberts,

Single-Dose Intranasal Administration of AdCOVID Elicits Systemic and Mucosal

Immunity against SARS-CoV-2 and Fully Protects Mice from Lethal Challenge. *Vaccines*

(Basel). **9**, 881 (2021).

84. M. D. Schultz, J. J. Suschak, D. Botta, A. Silva-Sanchez, R. G. King, T. W. Detchemendy,

C. D. Meshram, J. B. Foote, F. Zhou, J. L. Tipper, J. Zhang, K. S. Harrod, S. M. Leal, T. D.

Randall, M. S. Roberts, B. Georges, F. E. Lund, A single intranasal administration of

AdCOVID protects against SARS-CoV-2 infection in the upper and lower respiratory

tracts. *Human Vaccines & Immunotherapeutics*. **0**, 2127292 (2022).

85. T. Carroll, D. Fox, N. van Doremalen, E. Ball, M. K. Morris, A. Sotomayor-Gonzalez, V.

Servellita, A. Rustagi, C. K. Yinda, L. Fritts, J. R. Port, Z.-M. Ma, M. G. Holbrook, J.

Schulz, C. A. Blish, C. Hanson, C. Y. Chiu, V. Munster, S. Stanley, C. J. Miller, The

B.1.427/1.429 (epsilon) SARS-CoV-2 variants are more virulent than ancestral B.1 (614G)

in Syrian hamsters. *PLOS Pathogens*. **18**, e1009914 (2022).

86. A. D. Gruber, N. Osterrieder, L. D. Bertzbach, D. Vladimirova, S. Greuel, J. Ihlow, D.

Horst, J. Trimpert, K. Dietert, Standardization of Reporting Criteria for Lung Pathology in

SARS-CoV-2-infected Hamsters: What Matters? *Am J Respir Cell Mol Biol*. **63**, 856–859

(2020).

1472

Acknowledgments:

We thank Jesse Bloom (Fred Hutchinson Cancer Center) for sharing the SARS-CoV-2 spike

protein plasmids used in pseudovirus production. We are additionally grateful to Robert Kinsey,

who conducted the manufacture of the NLC used in these studies; Christopher Press, who

manufactured some RNA products used in this study; Jeff Guderian; and Ethan Lo. We also

thank Theresa Britschgi, who was instrumental as project manager for this work, and Valerie

Soza, for editorial assistance. We also gratefully acknowledge Thomas 'Scott' Simpler and

Rebecca Burnham for ABSL2 hamster husbandry, Donovan J. Murphy for ABSL3 hamster

husbandry, Jennifer L. Tipper and Kevin Harrod (UAB Department of Anesthesiology and

Perioperative Medicine) for providing SARS-CoV-2 viral stock, Levi Schaefers and Sixto M.

Leal Jr. (UAB Fungal Reference Lab) for viral RNA quantitation, and Emily E. Helmen and

Sherri Coffman (UAB Comparative Pathology Lab) for histology.

1485

Funding:

This work was funded in part by sponsored research from ImmunityBio, Inc. Development and

validation of the CBA array was supported by NIH U19 AI142737. The UAB SEBLAB was

supported by NIAID Regional Biocontainment Lab (RBL) awards UC6AI058599,

1G20AI167409-01, and 1G20AI167409-S1. The content is solely the responsibility of the

authors and does not necessarily represent the views of the funders.

Author Contributions

Conceptualization: MFJ, EAV, FEL

Supervision: MFJ, EAV, AG, TDR, FEL

Project administration: MFJ, EAV, AG, DB, FEL

Funding acquisition: EAV, CC, FEL

Investigation: MFJ, SB, DH, PB, JB, DH, RM, JS, NC, SR, MDS, DB, AS-S, DK, JA, GRK

Data curation: MFJ

Formal analysis: MFJ, SB, PB, NC, EAV, MDS

Writing – original draft: MFJ, EAV

Writing – review & editing: all authors

Competing Interests

AG and EAV declare no Competing Non-Financial Interests but the following Competing

Financial Interests. AG and EAV are co-inventors on PCT patent application PCT/US21/40388,

"Co-lyophilized RNA and Nanostructured Lipid Carrier," and related national filings, as well as

U.S provisional patent application 63/345,345, "Intranasal Administration of Thermostable RNA

1509   Vaccines," and 63/144,169, "A thermostable, flexible RNA vaccine delivery platform for

1510   pandemic response." All other authors declare that they have no competing interests.

1511

Data and Materials Availability

The datasets generated and/or analyzed during the current study are available from the

corresponding author on reasonable request. AAHI's unique saRNA constructs and nanostructured

lipid carriers are available for noncommercial research, academic non-clinical research, or other

not-for-profit scholarly purposes undertaken at a non-profit or government institution, under the

terms and conditions of an appropriate Material Transfer Agreement. Correspondence and material

requests should be addressed to the corresponding author.

1519

Supplementary Materials:

Figure S1. Spleen flow cytometry gating strategy.

Figure S2. Th1-Th2 balance of SARS-CoV-2 spike-responsive CD4$^+$ cells after prime-boost

immunization with SARS-CoV-2 saRNA-NLC vaccine.

Figure S3. Lung flow cytometry gating strategy.

Figure S4. Vaccine antibody responses in naive cagemates of sham-vaccinated and vaccinated

hamsters.

Figure S5. Correlation between body weight and serum anti-spike IgG titer in mice vaccinated

with SARS-CoV-2 saRNA-NLC vaccine and experimentally infected with SARS-CoV-2.

Figure S6. Vaccination with SARS-CoV-2 saRNA-NLC vaccine reduces lung pathology in both

vaccinated hamsters and naive cagemates irrespective of route of priming and boosting

immunization.