A bi-functional PARP-HDAC inhibitor with activity in Ewing sarcoma

Louise Ramos[1,2,3,*], Sarah Truong[2,3,*], Beibei Zhai[1,2,3], Jay Joshi[2,3], Fariba Ghaidi[2,3], Michael M. Lizardo[4], Taras Shyp[4,5], Hans Adomat[2], Stephane Le Bihan[2], Colin Collins[1,2], Jeffrey Bacha[3], Dennis Brown[3], John Langlands[3], Wang Shen[3], Nada Lallous[1,2], Poul H. Sorensen[4,5], and Mads Daugaard[1,2,3,5,#]

[1]Department of Urologic Sciences, University of British Columbia, Vancouver, BC, Canada
[2]Vancouver Prostate Centre, Vancouver, BC, Canada
[3]Rakovina Therapeutics Inc., Vancouver, BC, Canada
[4]BC Cancer Agency, Vancouver, BC, Canada
[5]Department of Pathology and Laboratory Medicine, University of British Columbia, Vancouver, Canada

[*]These two authors contributed equally to this work
[#]Correspondence: Mads Daugaard, Email: mads.daugaard@ubc.ca

Running title

Bi-functional PARP-HDAC inhibitor in Ewing sarcoma

Keywords

PARP inhibition, HDAC inhibition, Ewing sarcoma, DNA repair, bi-functional inhibitor

COI statement

L.R., S.T., B.Z., J.J., F.G. hold stock options and are employees of Rakovina Therapeutics Inc. and J.B., D.B., J.L., W.S. and M.D., are co-founders and/or employees of Rakovina Therapeutics Inc. The remaining authors declare no conflict of interest.

Abstract

HDAC inhibition has been shown to induce pharmacological BRCAness in cancer cells with

proficient DNA repair activity. This provides a rationale for exploring combination treatments

with HDAC and PARP inhibition in cancer types that are insensitive to single-agent PARP

inhibitors. Here, we report the concept and characterization of a novel bifunctional PARP

inhibitor (kt-3283) with dual activity towards PARP1/2 and HDAC enzymes in Ewing

sarcoma cells. Compared to the FDA-approved PARP (olaparib) and HDAC (vorinostat)

inhibitors, kt-3283 displayed enhanced cytotoxicity in Ewing sarcoma models. The kt-3283-

induced cytotoxicity was associated with a strong S and G2/M cell cycle arrest in the

nanomolar concentration range and elevated DNA damage as assessed by γH2AX tracking

and comet assays. In three-dimensional spheroid models of Ewing sarcoma, kt-3283

showed efficacy in lower concentrations than olaparib and vorinostat and kt-3283 inhibited

colonization of Ewing sarcoma cells in an *ex vivo* lung metastasis model. In summary, our

data demonstrates the preclinical justification for studying the benefit of dual PARP and

HDAC inhibition in the treatment of Ewing sarcoma in a clinical trial and provides proof-of-

concept for a bi-functional single-molecule therapeutic strategy.

Introduction

Poly-(ADP-ribose)-polymerase (PARP) proteins catalyze PARylation of cellular proteins

using an ADP (adenosine diphosphate)-ribose subunit of nicotinamide adenine dinucleotide

(NAD+) as the donor (1,2). The human genome encodes 17 PARP enzymes where at least

PARP1-3 has critical functions in DNA repair, PARP1 being the most characterized (3).

PARP1 is essential for the repair of single-strand DNA breaks (SSBs), which is the most

common type of breakpoint lesion in cellular DNA (4). When cells encounter SSBs, PARP1

binds the lesion and initiates a PARylation cascade of itself and histones embedded in the

chromatin surrounding the SSB lesion. This PARylation event serves as a signal to recruit

the SSB repair machinery to patch the lesions before and during DNA replication in the S-

phase of the cell cycle (5). Efficient SSB repair is important to prevent replication stress and

the more severe double-strand break (DSB) lesions that occur in S-phase when unrepaired

SSB lesions collide with the replication forks (4). DSB lesions in S-phase are mainly repaired

by homologous recombination (HR) that relies on proteins such as BRCA1 and BRCA2 (6).

Deleterious mutations in BRCA1/2 are found in subsets of breast, ovarian, and prostate

tumors, and sporadically in other solid tumor indications (7,8). These HR-deficient tumors

are indirectly dependent on proficient PARP enzyme activity to avoid accumulation of

catastrophic DSBs in S-phase and initiation of cell death (5). This dependency has paved

the way for PARP inhibition as a therapeutic strategy to create synthetic lethality in tumor

cells with BRCA1/2-deficiencies (9,10).

Currently, there are four approved PARP inhibitors in the clinic: olaparib (approved in 2014),

rucaparib (approved in 2016), niraparib (approved in 2017), and talazoparib (approved in

2018). These PARP inhibitors have been widely deployed in cancers with defects in HR

DNA repair activity caused by BRCA1/2 mutations (11). Encouraged by the success of

PARP inhibitors in BRCA1/2-mutated cancers, research attention has been expanded

towards cancer sub-types where HR repair is compromised due to molecular events other

than BRCA1/2 mutations. For example, tumors with mutations in RAD51, an enzyme acting

downstream of BRCA1/2 in the HR repair pathway, are also sensitive to PARP inhibition

(12). This concept is commonly referred to as 'BRCAness' and includes all events that mimic

BRCA1/2 loss in the context of HR repair (13).

In HR-proficient cancers, the state of BRCAness can be mimicked pharmacologically by

inhibition of proteins that impact BRCA1/2 expression. This potentially invites opportunities

to broaden the use of PARP inhibitors beyond current clinical practice. For example,

impairing dynamic chromatin events related to DNA replication and repair such as histone

acetylation can induce pharmacological BRCAness through indirect regulation of HR

components (14,15). Recent studies in leukemia, breast cancer, liver cancer, glioblastoma,

prostate cancer, and anaplastic thyroid cancer models demonstrated suppression of HR

activity with histone deacetylase (HDAC) inhibition that further supports the synergistic

potential of HDAC and PARP inhibition (16-23).

Ewing Sarcoma is a highly metastatic bone and soft tissue tumor affecting mainly children

and young adults, with a dismal 5-year survival rate of 15-30% for metastatic disease (24).

Ewing sarcoma is defined by the presence of specific gene fusion events involving *EWSR1*

and the erythroblast transformation specific (ETS) transcription factor *FLI1* (85%) or other

ETS-family transcription factors (15%), most often *ERG* (25). These gene fusions encode

chimeric oncoproteins (*e.g.,* EWS-FLI1 or EWS-ERG) that drive Ewing sarcoma initiation

and progression.

Ewing sarcoma cells are sensitive to PARP inhibitors *in vitro* and this sensitivity depends on

EWS-FLI1 (26,27). Ewing sarcoma cell line-derived xenografts in mice display sensitivity to

FDA-approved PARP inhibitors similar to the responses seen with the standard-of-care

chemotherapy temozolomide (26). These observations prompted a phase II single-agent

trial in Ewing sarcoma with olaparib, but despite encouraging pre-clinical data, these patients

failed to produce durable responses to single agent PARP inhibition (28). The

underwhelming response to PARP inhibitors in Ewing sarcoma patients is most likely due to

insufficient synthetic lethality and Ewing sarcoma is therefore a prime candidate for exploring

pharmacological BRCAness in the context of PARP inhibitors. Here, HDAC inhibition seems

attractive based on previously established sensitivity of Ewing sarcoma cells to HDAC

inhibitors (29-31). In this study, we have characterized and evaluated kt-3283, a novel dual-

specificity single-molecule inhibitor of PARP1/2 and HDAC enzymes in Ewing sarcoma

model systems.

Materials and Methods

Cell culture

All human cell lines were confirmed free of mycoplasma and maintained at 37°C with 5%

CO_2 and 95% humidity. CHLA10 cells were maintained in Iscove's Modified Dulbecco's

Medium (Hyclone cat# SH30228.01) containing 1x Insulin-Transferrin-Selenium (ITS)

(Thermo Fisher Scientific cat# 41400045) and 20% fetal bovine serum (FBS) (Gibco cat#

A3160401). TC32 cells were maintained in RPMI-1640 (Gibco cat# 11875119) with 10%

FBS and 1x GlutaMAX supplement (Thermo Fisher Scientific cat# 35050061).

HDAC activity assay

In vitro HDAC activity was measured using the FLUOR DE LYS® HDAC fluorometric activity

assay kit (Enzo Life Sciences cat# BML-AK500-0001) following the manufacturer's protocol.

IC_{50} values were calculated using a four-parameter variable slope non-linear regression in

GraphPad Prism 8 (GraphPad Software Inc.).

PARP1 and PARP2 activity assay

In vitro PARP1 activity was measured using the HT Universal Colorimetric PARP assay kit

(R&D Systems cat# 4677-096-K) and PARP2 activity was measured using the PARP2

colorimetric assay kit (BPS Bioscience cat# 80581) following the manufacturer's protocol.

IC$_{50}$ values were calculated using a four-parameter variable slope non-linear regression in
GraphPad Prism 8 (GraphPad Software Inc.).

PAR formation assay
Cellular PAR formation assays were used to measure the ability of a test compound to inhibit
polymerization of PAR. CHLA10 cells were plated on a black, clear-bottom 96-well plate and
allowed to attach overnight. Cells were pre-treated with increasing concentrations of test
inhibitors for 30 minutes at 37°C before H$_2$O$_2$ was added to a final concentration of 25 mM
and incubated for 5 minutes at RT. After 2x washes with 0.1% Tween-20 in PBS (PBS-T)
and 2x washes with PBS, cells were fixed with pre-chilled 70:30 methanol:acetone for 15
minutes at -20°C. Cells were washed with PBS, 2x with 3% BSA in PBS (BSA-PBS) and
again with PBS and then blocked with 3% BSA-PBS for 30 minutes at RT. Following 2x
washes with PBS and 1x wash with 3% BSA-PBS, cells were incubated for 1 hour at RT
with anti-PAR/pADPr monoclonal antibody (R&D Systems cat# 4335-MC-100) diluted 1:250
in 3% BSA-PBS. Plates were washed 2x with 3% BSA-PBS, 1x with PBS, 2x with PBS-T,
2x with PBS and 1x with 3% BSA-PBS, then incubated with goat anti-mouse IgG-FITC
(Thermo Scientific cat# F-2761) diluted 1:1000 in 3% BSA-PBS for 1 hour at RT. After
washing 2x with 3% BSA-PBS, 1x with PBS, 2x with PBS-T and 3x with PBS, 100 μL PBS
per well was added and plates were imaged on an IncuCyte® S3 system (Sartorius).
Fluorescence was quantified using the IncuCyte analysis software. Values were normalized
to a no primary antibody control and then % PAR formation was calculated by normalizing
to a DMSO control. IC$_{50}$ values were then calculated using a four parameter variable slope
non-linear regression in GraphPad Prism 8 (GraphPad Software Inc.).

Cell viability assay
Cells were plated on a 96-well plate (1000-5000 cells per well) in 100 μL appropriate medium
and allowed to attach overnight. 100 μL of medium containing dimethylsulfoxide (DMSO) or
increasing concentration of test compound was added to each well. Cells were maintained
at 37°C with 5% CO$_2$ and 95% humidity for 10 days for CHLA10 and 3 days for TC32 and
A673.Cell-Titer-Glo® viability assay was carried out for CHLA10.150 μL of media per well
was removed and plates were equilibrated at room temperature for 30 minutes, then
CellTiter-Glo® assay reagent was added to the wells. The plates were gently shaken on an
orbital shaker for 2 minutes and incubated at room temperature for 10 minutes in the dark.
Luminescence was measured using a Tecan Infinite M200Pro microplate reader. All

measurements were carried out in triplicate. For TC32 and A673, the plates were imaged

on an Incucyte S3 live cell imaging system after the treatment period was complete and %

confluency was measured using the Incucyte software. Values were normalized to a media-

only control and DMSO control to calculate % cell survival. EC_{50} values were calculated

using a four-parameter variable slope non-linear regression in GraphPad Prism 8

(GraphPad Software Inc.).

Cell cycle analysis

Cell cycle arrest profiles were evaluated via propidium iodide staining and flow cytometry.

CHLA10 and TC32 cells were plated in 10 cm plates with a cell density of 1.5×10^6 cells/plate

and 2.0×10^6 cells/plate, respectively. The media was replaced the following day with serum-

free media for 24 hours. Cells were treated with olaparib, vorinostat and kt-3283 in a dose

escalating manner for 24 hours for TC32 and 48 hours for CHLA10. Combination treatments

of olaparib with vorinostat, olaparib with belinostat and olaparib with panobinostat were

evaluated at the highest concentration of kt-3283 for both TC32 and CHLA10. Cells were

harvested and 1.0×10^6 cells for each treatment were fixed in 70% ethanol overnight at -

30°C. The suspension was then washed with cold 1X PBS and stained with a propidium

iodide solution in 1X PBS (50 µg/mL PI from 1mg/mL stock solution, 0.1mg/mL RNase A,

0.05% Triton X-100) and incubated at 37°C for 40min. Cells were then washed with 1X PBS,

filtered through a 40 µm strainer and resuspended with 500 µL of 1X PBS. Samples were

then run on FACS and analyzed in FlowJo v10 relative to the DMSO control.

Alkaline comet assay

To assess single and double-stranded DNA breaks, cells were plated in a 6-well plate at a

density of 1×10^6 cells/well and allowed to settle overnight. Cell media was replaced with

serum-free media for 24 hours, then treated with DMSO or increasing concentration of test

compound for 24 hours at 37°C with 5% CO_2 and 95% humidity. Cells were harvested as

instructed in Trevigen's CometAssay® protocol and combined with molten LMAgarose at a

ratio of 1:10 and pipetted onto CometSlides®. Cells were placed in the dark for 30 minutes

and immersed in lysis solution overnight at 4°C. Slides were immersed in Alkaline Unwinding

solution for 1 hour at 4°C in the dark then placed in a gel electrophoresis tray and immersed

in Alkaline Electrophoresis solution with an applied voltage of 25V for 30 minutes. Samples

were washed with dH_2O and 70% ethanol before staining with SYBR® Gold. Samples were

then viewed using a fluorescence microscope.

Acquired images were analyzed using the OpenComet tool on ImageJ (NIH). Further

statistics were performed using GraphPad Prism 8.

Immunofluorescence

CHLA10 cells were plated in a 24-well plate at a density of 3.5×10^5 cells/well and allowed to

settle overnight. Media was replaced with serum-free media for 24 hours, then cells were

treated with DMSO or test compound. After 24 hours, cells were fixed with 4%

paraformaldehyde (PFA) and stored at 4°C with 1X PBS. Cells were permeabilized with

0.5% Triton-X in PBS before probing with anti-phospho-histone H2AX (Ser 139) rabbit

monoclonal antibody (Cell Signaling Technology cat# 9718) and anti-cyclin A2 (EPR17351)

mouse monoclonal antibody (AbCam cat# 181591) overnight at 4°C. Cells were then

washed with 1X PBS and probed with goat anti rabbit IgG Alexa Fluor® 488 (Abcam cat#

ab150077), and mounted with VECTASHIELD Antifade Mounting medium with DAPI

solution onto microscope slides. Cells were then visualized on a confocal microscope

(Olympus FV3000).

Acquired images were analyzed by quantifying the foci using ImageJ (NIH). Further statistics

were performed using GraphPad Prism 8.

Western blot analysis

Cells were plated in a 6-well plate to 70-80% confluency. After allowing cells to settle

overnight, media was replaced with serum-free media for 24 hours. Cells were treated with

DMSO or test compound for 48 hours. Cells were harvested in Radioimmunoprecipitation

Assay (RIPA) lysis buffer combined with protease and phosphatase inhibitors. Cells were

stored at -80°C before running the western blot analysis. Protein yield was assessed using

Pierce™ BCA Protein Assay Kit (Thermofisher cat# 23225) and quantified using a

spectrophotometer plate reader (TECAN) at 562 nm. A total of 20 μg of protein extracts were

loaded per well in 4-15% Mini-PROTEAN® TGX™ Precast Protein gels (Bio-Rad cat#

4561084). After electrophoresis, proteins were transferred onto a 0.2 μm nitrocellulose

membrane. The membrane was blocked with LICOR® Odyssey Blocking Buffer in PBS.

After blocking, the membrane was incubated with antiphospho-histone H2AX (Ser 139)

rabbit monoclonal antibody (Cell Signaling Technology cat# 2577) and GAPDH (D16H11)

XP® rabbit monoclonal antibody (Cell Signaling Technology cat# 5174) at 4°C overnight and

then incubated with donkey anti-rabbit IRDye®800CW secondary antibody (LI-COR cat#

926-32213) for 1 hour at room temperature. The membrane was washed with 1X tris-

buffered saline with 1% tween-20 (TBS-T) before being scanned with Odyssey scanner (LI-

COR). Band intensities were analyzed using the ImageJ software and plotted using

GraphPad Prism 8.

Spheroid formation assay

CHLA10-tdTomato or TC32-tdTomato cells (2500 per well) were added to a 96-well clear

round bottom ultra-low attachment microplate (Corning cat# 7007) and allowed to form

spheroids for 24 hours or until they reached 200-300 μm in diameter. Medium containing

DMSO or increasing concentration of test compound was added to each well and spheroid

growth was monitored for 4 days post-treatment using the IncuCyte® Spheroid Analysis

system (Sartorius). Images from 4 days post-treatment and day 0 were analyzed using the

IncuCyte® Spheroid Analysis software module. Day 4 values were normalized using a

normalization factor from day 0 values and EC_{50} values were calculated using a four-

parameter variable slope non-linear regression in GraphPad Prism 8 (GraphPad Software

Inc.).

Pulmonary metastasis assay (PuMA)

Procedures involving mice were approved by local animal care committee, University of

British Columbia. EGFP-expressing A673 cells (1×10^6 cells/100 μl saline) were injected tail-

vein into 6-8-week-old immune-compromised NSG female mice (Jax Laboratories). After

injection, mice were euthanized via isoflurane and CO_2 asphyxiation according to local

animal care standard operating procedures. Lungs were insufflated via gravity perfusion with

a pre-warmed (37°C) 1:1 mixture of fully supplemented PneumaCult™-ALI media

(STEMCell cat #05001) and 1.2% low-melting point agarose (Lonza) as previously

described (32). The pluck (heart & lungs) was carefully removed and placed in ice-cold PBS

(supplemented with 1X penicillin/streptomycin) for 20 min to allow for agarose solidification.

Small lung slices (~2 mm x 4 mm) were obtained by manual cutting with sterile surgical

scissors, and 6 slices per condition were chosen for serial imaging at 0 and 14 days post-

injection/treatment. Lung slices were maintained in vitro on gelatine sponges partially

submerged in 2 mL of PneumaCult™ media +/- compound in a 6-well plate; media +/-

compound was refreshed every 3 days. On the day of imaging, lung slices from each group

were transferred to a small 35 mm petri dish with a glass coverslip bottom (IBIDI) to permit

aseptic widefield fluorescence imaging. The lung slices were imaged on an inverted Zeiss

Observer.Z1 Colibri microscope using 2.5X objective. Lung tumour burden (percent tumor

burden) for a lung slice is calculated as the quotient of the summed area of eGFP lesions

and total area of lung slice, multiplied by 100, as previously described (33). ImageJ software

was used for image processing. This calculation was performed for all lung slices (n = 5) per

experimental group. Average values of percent lung tumour burden per group were

compared and analyzed in Graph Prism 8 (GraphPad Software Inc.).

Results

A bi-specific small molecule with dual activity against PARP1/2 and HDAC enzymes

Through medicinal chemistry cycles, we developed a small-molecule inhibitor prototype (kt-

3283) with dual activity against PARP1/2 and HDACs. *In vitro* activity assay kits were used

to determine inhibition of PARP1, PARP2 and HDAC by kt-3283 compared to FDA-approved

PARP inhibitor olaparib and HDAC inhibitor vorinostat. A wide concentration range of each

compound was used to determine IC_{50} values. kt-3283 had an IC_{50} value of 1.89 µM for

inhibition of HDACs while vorinostat was about 40-fold lower at 0.05 µM (**Fig. 1A and**

Supplementary Fig. S1A). The PARP1 and PARP2 inhibitory activities of kt-3283 were

comparable to olaparib, with IC_{50} values for kt-3283 at 0.338 nM and 2.19 nM, respectively

(**Fig. 1B, 1C, Supplementary Fig. S1B and Supplementary Fig. S1C**). To further validate

the ability of kt-3283 to inhibit PARP1/2 activity, we used a cellular PAR synthesis assay to

determine the level of PAR formation. Comparable to olaparib, we determined an IC_{50} of

1.38 nM for the inhibition of PAR formation in cells treated with kt-3283 (**Fig. 1D and**

Supplementary Fig. S1D). These data indicate that kt-3283 is able to inhibit both PARP1/2

and HDAC enzymes.

Ewing sarcoma cells are highly sensitive to dual PARP1/2 and HDAC inhibition

In order to investigate the effect of kt-3283 on cell growth, we performed cell viability assays

in three Ewing sarcoma cell lines. Using an IncuCyte S3 live cell imaging system to evaluate

cell viability, we determined the EC_{50} values after three-day treatment with increasing

concentrations of kt-3283, three FDA-approved PARP inhibitors, or three FDA-approved

HDAC inhibitors in TC32 and A673 cells. kt-3283 demonstrated higher efficacy in

suppression of cell viability than olaparib, niraparib, vorinostat, and belinostat with EC_{50} of

0.0163 µM in TC32 cells (**Fig. 2A and Supplementary Fig. S2A)**. We also detected a similar

effect of kt-3283 in A673 cells with a much lower EC_{50} value of 0.0342 µM compared to

olaparib, niraparib, vorinostat, or belinostat treatment alone (**Fig. 2B and Supplementary**

Fig. S2B). However, treatment with talazoparib or panobinostat showed a more potent

inhibitory effect in both cell lines compared with kt-3283 (**Fig. 2A, 2B, Supplementary Fig.**

S2A and S2B). To further validate our findings, we also performed CellTiter-Glo® viability

assay to determine the EC_{50} values of these tested compounds after 10 days of treatment

in CHLA10 cells. Consistent with the IncuCyte assay, the EC_{50} value (0.053 μM) of kt-3283

treatment in CHLA10 cells was also much lower than olaparib, niraparib, vorinostat, and

belinostat (**Fig. 2C and Supplementary Fig. S2C**). Taken together, our data demonstrates

potent inhibitory effect of kt-3283 in Ewing sarcoma cells compared to most FDA-approved

PARP or HDAC inhibitors.

kt-3283 induces S and G2/M cell cycle arrest in Ewing sarcoma cells

PARP inhibitors and HDAC inhibitors constantly induce S/G2/M and G0/G1 cell cycle arrest,

respectively as PARP regulates replication fork progression and HDACs play a major role

in regulating the expression of cell cycle checkpoint proteins including cyclin-dependent

kinases, cyclin D1 and p21 (34,35). Here we examined the cell cycle profiles of CHLA10

and TC32 cells upon treatment with kt-3283 (3283), olaparib (OLA), and vorinostat (VOR)

in both single and combination regimens. Serum-starved cells were treated with increasing

concentrations of kt-3283 in complete medium for 24 or 48 h, and displayed strong S and

G2/M arrest at and above 0.175 μM in TC32 and 0.25 μM in CHLA10 cells. Similar cell cycle

arrest was only observed with olaparib treatment at concentrations higher than 3 μM in TC32

and 14.7 μM in CHLA10, respectively (**Fig. 3A, 3B, Supplementary Fig. S3A and S3B**).

Combination treatment with olaparib and vorinostat/belinostat (BEL) at equimolar

concentrations of kt-3283 (1 μM and 0.7 μM for CHLA10 and TC32 cells, respectively) had

almost no influence on cell cycle phases compared to control (**Fig. 3A, 3B, Supplementary**

Fig. S3A and S3B). These data demonstrate kt-3283 has stronger potency to induce S and

G2/M cell cycle arrest than olaparib alone or in combination with vorinostat or belinostat in

Ewing sarcoma cells.

kt-3283 treatment induces DNA damage in Ewing sarcoma cells

PARP inhibitors and HDAC inhibitors have been reported to induce DNA damage in cells

(36). We investigated the effect of kt-3283 on DNA damage compared to olaparib and

vorinostat treatment in Ewing sarcoma cells using western blot, immunofluorescence, and

comet assay. Phosphorylated histone variant H2AX (γH2AX) is a surrogate marker for DSBs

in DNA (37). Dianhydrogalactitol (DAG) was included as a positive control because previous

studies in our group showed that DAG induces replication-dependent DNA damage in a
variety of cancer cell lines (38,39). Treatment with kt-3283, olaparib or vorinostat induced
γH2AX expression in a dose-dependent manner in both CHLA10 and TC32 cells. In
comparison to olaparib and vorinostat, kt-3283 was able to induce γH2AX expression at a
much lower concentration range (**Fig. 4A, 4B, and Supplementary Fig. S4A and S4B**).
Moreover, CHLA10 cells treated with kt-3283 or olaparib also showed dose-dependent
γH2AX foci formation in immunofluorescence followed by confocal microscopy imaging with
kt-3283 at a much lower concentration range (**Fig. 4C, 4D, Supplementary Fig. S4C and
S4D**). However, vorinostat that induced G0/G1 cell cycle arrest (**Fig. 3A, 3B,
Supplementary Fig. S3A, and S3B**) demonstrated milder DNA damage foci formation in
CHLA10 cells (**Fig. 4E and Supplementary Fig. S4E**). To further consolidate our data, we
also employed alkaline comet assay as it can detect both SSBs and DSBs in cells. There is
significant DNA damage in CHLA10 cells treated with 1 μM kt-3283 but not with 1 μM
olaparib or vorinostat (**Fig. 4F and Supplementary Fig. S4F).** In summary, our data show
kt-3283 is able to induce DNA damage in Ewing sarcoma cells at a much lower concentration
range than olaparib or vorinostat.

kt-3283 inhibits 3D spheroid growth and metastasis of Ewing sarcoma cells
Spheroids are three-dimensional (3D) cell aggregates that can mimic tumor behavior more
accurately compared to two-dimensional cell cultures (40). To further validate our data, we
investigated the effect of kt-3283 on the 3D spheroid model with Ewing sarcoma cells.
CHLA10 and TC32 spheroids were established to 200-300 μm followed by treatment with
increasing concentrations of kt-3283, olaparib, or vorinostat. The growth of the spheroids
was monitored and quantified for four days using the IncuCyte S3 imaging system. The EC_{50}
values of kt-3283 in suppression of spheroid growth were much lower than olaparib and
vorinostat in both TC32 and CHLA10 cell models (**Fig. 5A and 5B Supplementary Fig.
S5A and S5B**). We also examined the effect of kt-3283 on metastatic growth of Ewing
sarcoma cells in an *ex vivo* pulmonary metastasis assay (PuMA). Here, colonization of A673
cells in mouse lungs was inhibited by 10 nM or 20 nM of kt-3283 (**Fig. 5C**). These data
suggest that kt-3283 is a potent inhibitor of Ewing sarcoma lung metastasis and that it is
more effective than olaparib or vorinostat alone in inhibiting 3D growth in Ewing sarcoma
spheroids.

Discussion

Pharmacological BRCAness can potentially offer a path to take PARP inhibition beyond the

BRCA1/2 mutation space and counter potential resistance to PARPi therapy. Epigenetic

modifiers such as HDACs, as well as DNA and histone methyltransferases are attractive

targets for induced BRCAness in BRCA1/2 proficient cancer scenarios (15,17,19,41-45). At

present, four clinical trials using PARPi in combination with the HDACi vorinostat

(NCT03259503 and NCT03742245), the DNA methyltransferase inhibitor decitabine

(NCT02878785), and the EZH2 histone methyltransferase inhibitor SHR2554

(NCT04355858), are currently ongoing.

FDA has approved three pan-HDACi drugs (vorinostat, belinostat, and Panobinostat) and

one HDAC1/2-selective HDACi (romidepsin) for treatment of hematological cancers (46).

Histone acetylation attenuates chromatin structure and plays a critical role in recognition and

repair of DNA lesions (47). HDACi-induced downregulation of key HR proteins including

BRCA1, BRCA2, and RAD51 has been established in a variety of cancers types, and HDACi

treatment sensitizes cancer cells to PARPi (15,17-21). This corroborative activity of HDAC

and PARP inhibition is particularly interesting in the context of HR-proficient cancer types

where PARPi therapy has limited effect on its own, such as Ewing sarcoma. However, dose-

limiting toxicity with HDACi therapy is not uncommon in solid tumors (e.g. breast cancer and

sarcomas) and has been preventing some therapeutic effects as stand-alone and in

treatment combinations (48-50). The HDACi ingredient must be carefully adjusted to prevent

over-lapping toxicity events arising from the combination with other therapeutic moieties,

which can be challenging when working with different pharmacokinetics profiles.

We have characterized a novel bifunctional PARP-HDAC single-molecule inhibitor, kt-3283,

in Ewing sarcoma models to evaluate the potential benefit of combined PARP-HDAC

inhibition over stand-alone PARPi or HDACi treatments. kt-3283 has similar PARPi activity

as olaparib and slightly lower HDACi activity than vorinostat. However, the dual activity of

kt-3283 is 30-to-80 times more cytotoxic to Ewing sarcoma cells than olaparib, and 30-to-60

times more cytotoxic than vorinostat alone. Similarly, kt-3283 induces cell cycle arrest and

DNA damage in Ewing sarcoma cells in much lower concentrations than olaparib and

vorinostat. When compared in 3D spheroid models, kt-3283 showed efficacy at 30-to-40

times lower concentrations than olaparib and at 5-to-10 times lower concentrations than

vorinostat. kt-3283 also hinders metastatic growth of Ewing sarcoma cells in an *ex vivo*

pulmonary metastasis model, with a strong inhibitory effect using as little as 10 nM of

inhibitor. Combining PARP and HDAC inhibition into one single molecule may also offer a
convenient way to prevent resistance to PARPi therapy. For example, Ewing sarcoma and
many other solid tumor indications epigenetically suppress expression of the tumor
suppressor gene Schlafen 11 (*SLFN11*), which leads to resistance to DNA damage-inducing
agents, including PARPi therapy (51-53). Important here, HDACi treatment prompts re-
expression of SLFN11 and re-sensitization to PARPi (54-57). In summary, our work provides
preclinical justification for studying a novel single-molecule PARP-HDAC inhibitor in Ewing
sarcoma with improved cytotoxicity and DNA damage activity as compared to PARPi and
HDACi alone. This concept will likely be relevant in cancer indications beyond Ewing
sarcoma and potentially offers an opportunity to suppress therapeutic resistance.
383

Acknowledgments

This work was supported by a Robert J. Arceci Innovation Award from the St. Baldrick's
Foundation; a St. Baldrick's Foundation/Martha's Better Ewing Sarcoma Treatment (BEST)
Grant (#663113); an Accelerate grant awarded by the Mathematics of Information
Technology and Complex Systems (Mitacs); and Rakovina Therapeutics Inc.

References
391

1. Bock FJ, Chang P. New directions in poly(ADP-ribose) polymerase biology. *Febs j*
2016;**283**(22):4017-31 doi 10.1111/febs.13737.
2. Kim MY, Zhang T, Kraus WL. Poly(ADP-ribosyl)ation by PARP-1: 'PAR-laying' NAD+
into a nuclear signal. *Genes Dev* 2005;**19**(17):1951-67 doi 10.1101/gad.1331805.
3. Curtin NJ, Szabo C. Poly(ADP-ribose) polymerase inhibition: past, present and
future. *Nat Rev Drug Discov* 2020;**19**(10):711-36 doi 10.1038/s41573-020-0076-6.
4. Caldecott KW. Single-strand break repair and genetic disease. *Nat Rev Genet*
2008;**9**(8):619-31 doi 10.1038/nrg2380.
5. Caldecott KW. Protein ADP-ribosylation and the cellular response to DNA strand
breaks. *DNA Repair (Amst)* 2014;**19**:108-13 doi 10.1016/j.dnarep.2014.03.021.
6. McKinnon PJ, Caldecott KW. DNA strand break repair and human genetic disease.
Annu Rev Genomics Hum Genet 2007;**8**:37-55 doi
10.1146/annurev.genom.7.080505.115648.
7. Maxwell KN, Domchek SM. Cancer treatment according to BRCA1 and BRCA2
mutations. *Nat Rev Clin Oncol* 2012;**9**(9):520-8 doi 10.1038/nrclinonc.2012.123.

8.   de Bono J, Mateo J, Fizazi K, Saad F, Shore N, Sandhu S*, et al.* Olaparib for Metastatic Castration-Resistant Prostate Cancer. *N Engl J Med* 2020;**382**(22):2091-102 doi 10.1056/NEJMoa1911440.

9.   Farmer H, McCabe N, Lord CJ, Tutt AN, Johnson DA, Richardson TB*, et al.* Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature* 2005;**434**(7035):917-21 doi 10.1038/nature03445.

10.   Bryant HE, Schultz N, Thomas HD, Parker KM, Flower D, Lopez E*, et al.* Specific killing of BRCA2-deficient tumours with inhibitors of poly(ADP-ribose) polymerase. *Nature* 2005;**434**(7035):913-7 doi 10.1038/nature03443.

11.   Cai Z, Liu C, Chang C, Shen C, Yin Y, Yin X*, et al.* Comparative safety and tolerability of approved PARP inhibitors in cancer: A systematic review and network meta-analysis. *Pharmacological Research* 2021;**172**:105808 doi https://doi.org/10.1016/j.phrs.2021.105808.

12.   Yang H, Wei Y, Zhang Q, Yang Y, Bi X, Yang L*, et al.* CRISPR/Cas9-induced saturated mutagenesis identifies Rad51 haplotype as a marker of PARP inhibitor sensitivity in breast cancer. *Mol Med Rep* 2022;**26**(2) doi 10.3892/mmr.2022.12774.

13.   Abbotts R, Dellomo AJ, Rassool FV. Pharmacologic Induction of BRCAness in BRCA-Proficient Cancers: Expanding PARP Inhibitor Use. *Cancers (Basel)* 2022;**14**(11) doi 10.3390/cancers14112640.

14.   Lord CJ, Ashworth A. BRCAness revisited. *Nat Rev Cancer* 2016;**16**(2):110-20 doi 10.1038/nrc.2015.21.

15.   Ladd B, Ackroyd JJ, Hicks JK, Canman CE, Flanagan SA, Shewach DS. Inhibition of homologous recombination with vorinostat synergistically enhances ganciclovir cytotoxicity. *DNA Repair (Amst)* 2013;**12**(12):1114-21 doi 10.1016/j.dnarep.2013.10.008.

16.   Valdez BC, Nieto Y, Yuan B, Murray D, Andersson BS. HDAC inhibitors suppress protein poly(ADP-ribosyl)ation and DNA repair protein levels and phosphorylation status in hematologic cancer cells: implications for their use in combination with PARP inhibitors and chemotherapeutic drugs. *Oncotarget* 2022;**13**:1122-35 doi 10.18632/oncotarget.28278.

17.   Wiegmans AP, Yap PY, Ward A, Lim YC, Khanna KK. Differences in Expression of Key DNA Damage Repair Genes after Epigenetic-Induced BRCAness Dictate Synthetic Lethality with PARP1 Inhibition. *Mol Cancer Ther* 2015;**14**(10):2321-31 doi 10.1158/1535-7163.mct-15-0374.

18.   Liang BY, Xiong M, Ji GB, Zhang EL, Zhang ZY, Dong KS, *et al.* Synergistic
suppressive effect of PARP-1 inhibitor PJ34 and HDAC inhibitor SAHA on
proliferation of liver cancer cells. *J Huazhong Univ Sci Technolog Med Sci*
2015;**35**(4):535-40 doi 10.1007/s11596-015-1466-6.
19.   Rasmussen RD, Gajjar MK, Jensen KE, Hamerlik P. Enhanced efficacy of combined
HDAC and PARP targeting in glioblastoma. *Mol Oncol* 2016;**10**(5):751-63 doi
10.1016/j.molonc.2015.12.014.
20.   Xiao W, Graham PH, Hao J, Chang L, Ni J, Power CA, *et al.* Combination therapy
with the histone deacetylase inhibitor LBH589 and radiation is an effective regimen
for prostate cancer cells. *PLoS One* 2013;**8**(8):e74253 doi
10.1371/journal.pone.0074253.
21.   Yin L, Liu Y, Peng Y, Yu X, Gao Y, Yuan B, *et al.* PARP inhibitor veliparib and HDAC
inhibitor SAHA synergistically co-target the UHRF1/BRCA1 DNA damage repair
complex in prostate cancer cells. *J Exp Clin Cancer Res* 2018;**37**(1):153 doi
10.1186/s13046-018-0810-7.
22.   Ha K, Fiskus W, Choi DS, Bhaskara S, Cerchietti L, Devaraj SG, *et al.* Histone
deacetylase inhibitor treatment induces 'BRCAness' and synergistic lethality with
PARP inhibitor and cisplatin against human triple negative breast cancer cells.
Oncotarget 2014;**5**(14):5637-50 doi 10.18632/oncotarget.2154.
23.   Baldan F, Mio C, Allegri L, Puppin C, Russo D, Filetti S, *et al.* Synergy between HDAC
and PARP Inhibitors on Proliferation of a Human Anaplastic Thyroid Cancer-Derived
Cell Line. *Int J Endocrinol* 2015;**2015**:978371 doi 10.1155/2015/978371.
24.   Zöllner SK, Amatruda JF, Bauer S, Collaud S, de Álava E, DuBois SG, *et al.* Ewing
Sarcoma-Diagnosis, Treatment, Clinical Challenges and Future Perspectives. *J Clin*
Med 2021;**10**(8) doi 10.3390/jcm10081685.
25.   Balamuth NJ, Womer RB. Ewing's sarcoma. *Lancet Oncol* 2010;**11**(2):184-92 doi
10.1016/s1470-2045(09)70286-4.
26.   Brenner JC, Ateeq B, Li Y, Yocum AK, Cao Q, Asangani IA, *et al.* Mechanistic
rationale for inhibition of poly(ADP-ribose) polymerase in ETS gene fusion-positive
prostate cancer. *Cancer Cell* 2011;**19**(5):664-78 doi 10.1016/j.ccr.2011.04.010.
27.   Garnett MJ, Edelman EJ, Heidorn SJ, Greenman CD, Dastur A, Lau KW, *et al.*
Systematic identification of genomic markers of drug sensitivity in cancer cells.
Nature 2012;**483**(7391):570-5 doi 10.1038/nature11005.

28. Choy E, Butrynski JE, Harmon DC, Morgan JA, George S, Wagner AJ*, et al.* Phase II study of olaparib in patients with refractory Ewing sarcoma following failure of standard chemotherapy. *BMC Cancer* 2014;**14**(1):813 doi 10.1186/1471-2407-14-813.

29. Sakimura R, Tanaka K, Nakatani F, Matsunobu T, Li X, Hanada M*, et al.* Antitumor effects of histone deacetylase inhibitor on Ewing's family tumors. *Int J Cancer* 2005;**116**(5):784-92 doi 10.1002/ijc.21069.

30. Schmidt O, Nehls N, Prexler C, von Heyking K, Groll T, Pardon K*, et al.* Class I histone deacetylases (HDAC) critically contribute to Ewing sarcoma pathogenesis. *J Exp Clin Cancer Res* 2021;**40**(1):322 doi 10.1186/s13046-021-02125-z.

31. Souza BK, da Costa Lopez PL, Menegotto PR, Vieira IA, Kersting N, Abujamra AL*, et al.* Targeting Histone Deacetylase Activity to Arrest Cell Growth and Promote Neural Differentiation in Ewing Sarcoma. *Mol Neurobiol* 2018;**55**(9):7242-58 doi 10.1007/s12035-018-0874-6.

32. Scopim-Ribeiro R, Lizardo MM, Zhang HF, Dhez AC, Hughes CS, Sorensen PH. NSG Mice Facilitate ex vivo Characterization of Ewing Sarcoma Lung Metastasis Using the PuMA Model. *Front Oncol* 2021;**11**:645757 doi 10.3389/fonc.2021.645757.

33. Lizardo MM, Sorensen PH. Practical Considerations in Studying Metastatic Lung Colonization in Osteosarcoma Using the Pulmonary Metastasis Assay. *J Vis Exp* 2018(133) doi 10.3791/56332.

34. Ramaiah MJ, Tangutur AD, Manyam RR. Epigenetic modulation and understanding of HDAC inhibitors in cancer therapy. *Life Sciences* 2021;**277**:119504 doi https://doi.org/10.1016/j.lfs.2021.119504.

35. da Costa A, Chowdhury D, Shapiro GI, D'Andrea AD, Konstantinopoulos PA. Targeting replication stress in cancer therapy. *Nat Rev Drug Discov* 2022 doi 10.1038/s41573-022-00558-5.

36. Robert C, Rassool FV. HDAC inhibitors: roles of DNA damage and repair. *Adv Cancer Res* 2012;**116**:87-129 doi 10.1016/b978-0-12-394387-3.00003-3.

37. Rahmanian N, Shokrzadeh M, Eskandani M. Recent advances in γH2AX biomarker-based genotoxicity assays: A marker of DNA damage and repair. *DNA Repair* 2021;**108**:103243 doi https://doi.org/10.1016/j.dnarep.2021.103243.

38. Zhai B, Steinø A, Bacha J, Brown D, Daugaard M. Dianhydrogalactitol induces replication-dependent DNA damage in tumor cells preferentially resolved by

homologous recombination. *Cell Death Dis* 2018;**9**(10):1016 doi 10.1038/s41419-
018-1069-9.

39.   Zhai B, Li Y, Kotapalli SS, Bacha J, Brown D, Steinø A*, et al.* Dianhydrogalactitol
synergizes with topoisomerase poisons to overcome DNA repair activity in tumor
cells. *Cell Death Dis* 2020;**11**(7):577 doi 10.1038/s41419-020-02780-8.

40.   Jubelin C, Muñoz-Garcia J, Griscom L, Cochonneau D, Ollivier E, Heymann MF*, et*
al. Three-dimensional in vitro culture models in oncology research. *Cell Biosci*
2022;**12**(1):155 doi 10.1186/s13578-022-00887-3.

41.   Jones PA, Issa JP, Baylin S. Targeting the cancer epigenome for therapy. *Nat Rev*
Genet 2016;**17**(10):630-41 doi 10.1038/nrg.2016.93.

42.   Muvarak NE, Chowdhury K, Xia L, Robert C, Choi EY, Cai Y*, et al.* Enhancing the
Cytotoxic Effects of PARP Inhibitors with DNA Demethylating Agents - A Potential
Therapy for Cancer. *Cancer Cell* 2016;**30**(4):637-50 doi 10.1016/j.ccell.2016.09.002.

43.   Yamaguchi H, Du Y, Nakai K, Ding M, Chang SS, Hsu JL*, et al.* EZH2 contributes to
the response to PARP inhibitors through its PARP-mediated poly-ADP ribosylation in
breast cancer. *Oncogene* 2018;**37**(2):208-17 doi 10.1038/onc.2017.311.

44.   Abbotts R, Topper MJ, Biondi C, Fontaine D, Goswami R, Stojanovic L*, et al.* DNA
methyltransferase inhibitors induce a BRCAness phenotype that sensitizes NSCLC
to PARP inhibitor and ionizing radiation. *Proc Natl Acad Sci U S A*
2019;**116**(45):22609-18 doi 10.1073/pnas.1903765116.

45.   Robert C, Nagaria PK, Pawar N, Adewuyi A, Gojo I, Meyers DJ*, et al.* Histone
deacetylase inhibitors decrease NHEJ both by acetylation of repair factors and
trapping of PARP1 at DNA double-strand breaks in chromatin. *Leuk Res* 2016;**45**:14-
23 doi 10.1016/j.leukres.2016.03.007.

46.   Ho TCS, Chan AHY, Ganesan A. Thirty Years of HDAC Inhibitors: 2020 Insight and
Hindsight. *J Med Chem* 2020;**63**(21):12460-84 doi 10.1021/acs.jmedchem.0c00830.

47.   Misteli T, Soutoglou E. The emerging role of nuclear architecture in DNA repair and
genome   maintenance.   *Nat   Rev   Mol   Cell   Biol*   2009;**10**(4):243-54   doi
10.1038/nrm2651.

48.   Bondarev AD, Attwood MM, Jonsson J, Chubarev VN, Tarasov VV, Schiöth HB.
Recent developments of HDAC inhibitors: Emerging indications and novel molecules.
Br J Clin Pharmacol 2021;**87**(12):4577-97 doi 10.1111/bcp.14889.

49. Wawruszak A, Borkiewicz L, Okon E, Kukula-Koch W, Afshan S, Halasa M. Vorinostat (SAHA) and Breast Cancer: An Overview. *Cancers (Basel)* 2021;**13**(18) doi 10.3390/cancers13184700.

50. Lanzi C, Cassinelli G. Combinatorial strategies to potentiate the efficacy of HDAC inhibitors in fusion-positive sarcomas. *Biochem Pharmacol* 2022;**198**:114944 doi 10.1016/j.bcp.2022.114944.

51. Tang SW, Bilke S, Cao L, Murai J, Sousa FG, Yamade M*, et al.* SLFN11 Is a Transcriptional Target of EWS-FLI1 and a Determinant of Drug Response in Ewing Sarcoma. *Clin Cancer Res* 2015;**21**(18):4184-93 doi 10.1158/1078-0432.ccr-14-2112.

52. Lok BH, Gardner EE, Schneeberger VE, Ni A, Desmeules P, Rekhtman N*, et al.* PARP Inhibitor Activity Correlates with SLFN11 Expression and Demonstrates Synergy with Temozolomide in Small Cell Lung Cancer. *Clin Cancer Res* 2017;**23**(2):523-35 doi 10.1158/1078-0432.ccr-16-1040.

53. Knelson EH, Patel SA, Sands JM. PARP Inhibitors in Small-Cell Lung Cancer: Rational Combinations to Improve Responses. *Cancers (Basel)* 2021;**13**(4) doi 10.3390/cancers13040727.

54. Yin YP, Ma LY, Cao GZ, Hua JH, Lv XT, Lin WC. FK228 potentiates topotecan activity against small cell lung cancer cells via induction of SLFN11. *Acta Pharmacol Sin* 2022;**43**(8):2119-27 doi 10.1038/s41401-021-00817-y.

55. Tang SW, Thomas A, Murai J, Trepel JB, Bates SE, Rajapakse VN*, et al.* Overcoming Resistance to DNA-Targeted Agents by Epigenetic Activation of Schlafen 11 (SLFN11) Expression with Class I Histone Deacetylase Inhibitors. *Clin Cancer Res* 2018;**24**(8):1944-53 doi 10.1158/1078-0432.ccr-17-0443.

56. Moribe F, Nishikori M, Takashima T, Taniyama D, Onishi N, Arima H*, et al.* Epigenetic suppression of SLFN11 in germinal center B-cells during B-cell development. *PLoS One* 2021;**16**(1):e0237554 doi 10.1371/journal.pone.0237554.

57. Murai J, Thomas A, Miettinen M, Pommier Y. Schlafen 11 (SLFN11), a restriction factor for replicative stress induced by DNA-targeting anti-cancer therapies. *Pharmacol Ther* 2019;**201**:94-102 doi 10.1016/j.pharmthera.2019.05.009.

573 **Data Availability Statement**

574 The data generated in this study are available upon request from the corresponding author.

Figure 1

Characterization of a bi-specific small molecule with dual activity against PARP1/2 and HDAC enzymes.
(A) In vitro HDAC activity in HeLa nuclear extracts treated with kt-3283 or vorinostat. **(B)** PARP1 activity in vitro after treatment with kt-3283 and olaparib. **(C)** PARP2 activity in vitro after treatment with kt-3283 and olaparib. **(D)** PAR formation in CHLA10 Ewing sarcoma cells treated with kt-3283 and olaparib. Values were normalized to control and IC50 was calculated as the concentration required to produce 50% inhibition of activity from non-linear regression plots using GraphPad Prism8 software. Data shown are the mean values of n=3 replicates with representative graphs.

Figure 2

A

TC32	
	Mean EC_{50} ± SD (µM)
kt-3283	0.0163 ± 0.0030
olaparib	1.17 ± 0.41
niraparib	0.136 ± 0.056
talazoparib	0.00361 ± 0.00095
vorinostat	0.827 ± 0.107
belinostat	0.188 ± 0.051
panobinostat	0.00576 ± 0.0016

B

A673	
	Mean EC_{50} ± SD (µM)
kt-3283	0.0342 ± 0.0077
olaparib	1.06 ± 0.26
niraparib	0.708 ± 0.83
talazoparib	0.00534 ± 0.0022
vorinostat	1.30 ± 0.20
belinostat	0.290 ± 0.10
panobinostat	0.00641 ± 0.0022

C

CHLA10	
	Mean EC_{50} ± SD (µM)
kt-3283	0.053 ± 0.003
olaparib	4.41 ± 1.13
niraparib	1.22 ± 0.32
talazoparib	0.027 ± 0.006
vorinostat	3.18 ± 0.96
belinostat	0.17 ± 0.016
panobinostat	0.0058 ± 0.0024

Ewing sarcoma cells are highly sensitive to dual PARP1/2 and HDAC inhibition.
(A) In TC32 cells, cell viability was examined by IncuCyte S3 live cell imaging system following three-day treatments with increasing concentrations of indicated compounds. EC50 values were calculated as the concentration required for 50% cell viability. n=3 **(B)** EC50 values of tested compounds were also determined in A673 cells using the same experimental condition as in (A). **(C)** EC50 values of indicated inhibitors were determined using CellTiter-Glo® cell viability assays in CHLA10 cells. Cells were exposed to ten-day treatments of increasing concentrations of the inhibitors, and the EC50 values were calculated as the concentration required for 50% cell viability. n=3.

Figure 3

kt-3283 induces S and G2/M cell cycle arrest in Ewing sarcoma cells.
(A) CHLA10 cells were synchronized at G0/G1 phase by serum starvation for 24 h before treatments with kt-3283, olaparib, or vorinostat as indicated in complete medium for 48 h. Then, cell cycle profiles were examined by propidium iodide (PI) staining followed by flow cytometric analysis as described in "Materials and Methods".
(B) Cell cycle analysis was also performed in TC32 cells treated with kt-3283, olaparib, or vorinostat as indicated for 24 h using the same experimental procedures as in (A).

Figure 4

kt-3283 treatment induces DNA damage in Ewing sarcoma cells.
(A) CHLA10 cells treated with 5 µM dianhydrogalactitol (DAG) or increasing doses of olaparib (1-37 µM), vorinostat (1-20 µM), or kt-3283 (0.05-1 µM) for 48 h and analyzed for γH2AX expression by western blot.
(B) TC32 cells treated with 2.5 µM DAG or increasing doses of olaparib (0.35-13 µM), vorinostat (0.35-8 µM), or kt-3283 (0.018-0.35 µM) for 48 h and analyzed as in (A). **(C-E)** CHLA10 cells treated with DAG (2.5 µM) or increasing doses of kt-3283 (0-1 µM) **(C)**, olaparib (1-37.5 µM) **(D)**, or vorinostat (1-18 µM) **(E)** for 24 h were analyzed for γH2AX foci by immunofluorescence and confocal microscopy imaging. Scale bar represents 10 µm.
(F) CHLA10 cells treated with 1 µM kt-3283, olaparib, or vorinostat followed by comet assay as described in "Materials and Methods". 5 µM DAG was included as positive control. ****p<0.0001

Figure 5

A

TC32

TC32	
	EC$_{50}$ ± SD (µM)
kt-3283	0.166 ± 0.00907
olaparib	4.69 ± 0.950
vorinostat	1.53 ± 0.541

B

CHLA10

CHLA10	
	EC$_{50}$ ± SD (µM)
kt-3283	0.255 ± 0.0457
olaparib	14.3 ± 2.51
vorinostat	2.95 ± 1.09

C

Lung tumour burden (%)

eGFP-expressing A673 cells

kt-3283 inhibits 3D spheroid growth and metastasis of Ewing sarcoma cells.
(A) TC32 spheroid growth following four days of treatment with increasing concentrations of kt-3283, olaparib, or vorinostat was monitored using the IncuCyte® Spheroid Analysis system. The EC50 values were calculated as the concentration required for 50% inhibition of growth from non-linear regression plots using GraphPad Prism8 software. Representative images of TC32 spheroids at day 0 and day 4 with DMSO,1 µM kt-3283, 1 µM olaparib, or 1 µM vorinostat are shown with scale bars representing 400 µm. (B) CHLA10 spheroid growth following four-day treatment with increasing concentrations of kt-3283, olaparib, or vorinostat was also examined using the same experimental procedures as mentioned in (A). Representative images of CHLA10 3D spheroids at day 0 and day 4 with DMSO,1 µM kt-3283, 1 µM olaparib, or 1 µM vorinostat are shown with scale bars representing 400 µm.
(C) Lung tumour burden following 14 days of treatment with vehicle, 5,10 or 20 nM of kt-3283. n=5. Representative fluorescence images of A673 cells in lung slices following 14 days of treatment with 5 or 10 nM of kt-3283. Scale bar represents 1 mm. *p<0.05 **p<0.