Transcriptome dynamics of *Pseudomonas aeruginosa* during transition

from replication-uncoupled to -coupled growth

Kathrin Alpers[1], Elisabeth Vatareck[1], Lothar Gröbe[2], Mathias Müsken[3], Maren Scharfe[4],

Susanne Häussler[1,5,6,7*], Jürgen Tomasch[1,8*]

[1]Department of Molecular Bacteriology, Helmholtz Centre for Infection Research,
Braunschweig, Germany
[2]Platform Flow Cytometry and Cell Sorting, Department of Experimental Immunology,
Helmholtz Centre for Infection Research, Braunschweig, Germany
[3]Central Facility for Microscopy, Helmholtz Centre for Infection Research, Braunschweig,
Germany
[4]Platform Genome Analytics, Helmholtz Centre for Infection Research, Braunschweig,
Germany
[5]Institute for Molecular Bacteriology, Twincore, Centre for Clinical and Experimental
Infection Research, Hannover, Germany
[6]Department of Clinical Microbiology, Copenhagen University Hospital – Rigshospitalet,
2100 Copenhagen, Denmark
[7]Cluster of Excellence RESIST (EXC 2155), Hannover Medical School, 30265 Hannover,
Germany
[8]Institute of Microbiology of the Czech Academy of Science, Center Algatech, Třeboň,
Czech Republic
22
23 *correspondence: Susanne.Haeussler@helmholtz-hzi.de
24 *correspondence: tomasch@alga.cz

Abstract

In bacteria, either chromosome duplication is coupled to cell division with only one replication round per cell cycle or DNA is replicated faster than the cells divide thus both processes are uncoupled. Here, we show that the opportunistic pathogen *Pseudomonas aeruginosa* switches from fast uncoupled to sustained coupled growth when cultivated under standard laboratory conditions. The transition was characterized by fast-paced, sequential changes in transcriptional activity along the *ori-ter* axis of the chromosome reflecting adaptation to the metabolic needs during both growth phases. Quorum sensing (QS) activity was highest at the onset of the coupled growth phase during which only a quarter of the cells keeps replicating. RNA sequencing of subpopulations of these cultures sorted based on their DNA content, revealed a strong gene dosage effect as well as specific expression patterns for replicating and non-replicating cells. Expression of flagella and *mexE*, involved in multi drug efflux was restricted to cells that did not replicate, while those that did showed a high activity of the cell division locus and recombination genes. A possible role of QS in the formation of these subpopulations upon switching to coupled growth could be a subject of further research.

Significance statement

The coordination of gene expression with the cell cycle has so far been studied only in a handful of bacteria, the bottleneck being the need for synchronized cultures. Here, we determined replication-associated effects on transcription by comparing *Pseudomonas aeruginosa* cultures that differ in their growth mode and number of replicating chromosomes. We further show that cell cycle-specific gene regulation can be principally identified by RNA sequencing of subpopulations from cultures that replicate only once per cell division and that are sorted according to their DNA content. Our approach opens the possibility to study asynchronously growing bacteria from a wide phylogenetic range and thereby enhance our understanding of the evolution of cell-cycle control on the transcriptional level.

Introduction

Bacteria differ in the ways replication is coordinated with cell growth and division[1]. In fast-growing representatives, such as the model organisms *Escherichia coli* or *Bacillus subtilis*, the speed of DNA replication exceeds that of cell division. This uncoupling of both processes results in a gene dosage gradient along the origin(*ori*)-terminus(*ter*) axis of the chromosome. The higher gene copy number closer to *ori* can be exploited to maximize expression of traits needed during rapid growth and to control gene expression[2,3]. It has been shown that moving an *ori*-located *Vibrio cholerae* gene cluster coding for ribosomal proteins close to *ter* reduced the growth rate of the culture, while the wild-type growth level could be restored by placing two copies of this cluster at *ter*[4]. Furthermore, the timing of spore formation in *B. subtilis* is an example for dosage imbalances triggering regulatory events between genes located on opposite ends of the replicating chromosomes[5]. In slow-growing bacteria the chromosome is duplicated only once per cell division, thus both processes are coupled[6]. In several bacterial phyla, a differentiation program is triggered during this eukaryote-like cell cycle. The best studied model is the bi-phasic lifestyle of *Caulobacter crescentus*. In this bacterium a complex gene regulatory network precisely times the development of a flagellated from a stalked cell during replication and cell division[7].

Pseudomonas aeruginosa is a ubiquitous environmental bacterium, but also an opportunistic pathogen frequently causing nosocomial infections of various body sites, such as the lung, bloodstream, urinary tract and burn wounds[8]. Furthermore, *P. aeruginosa* poses a particular threat to patients suffering from cystic fibrosis (CF)[9]. During live-long chronic infections of the CF lung, the bacterium adapts and evolves towards a slow growing phenotype[10]. Doubling times are estimated to be around 30 min under laboratory conditions in lysogeny broth (LB) medium and 1.9 to 4.6 h in the CF lung[11]. The cell cycle dynamics of *P. aeruginosa* has been extensively studied. Its chromosome is oriented with *ori* close to the center of the cell and *ter* located at the cell pole where the division plane forms. During replication both *ori* move to the poles of the elongated pre-divisional cell where another round of replication can be started[12,13]. Despite the huge body of comparative transcriptome data available for this important pathogen[14–16], the effect of replication on gene expression has not explicitly been studied yet.

Here we monitored growth and cell division and recorded a time-resolved transcriptome of *P.*

aeruginosa PA14 in LB medium over 10 hours at 1 hour intervals. We show that the culture

switches from fast replication-uncoupled to sustained coupled growth. The transition is

characterized by fast-paced, sequential changes in transcriptional activity along the *ori-ter*

axis. Furthermore, we identified replication- and non-replication-associated gene expression

in cells showing coupled growth using a newly developed protocol based on fluorescence-

activated cell sorting (FACS).

Results

Growth and replication dynamics of *P. aeruginosa* in LB medium.

In accordance with previous reports[11], *P. aeruginosa* cultures reached an OD_{600} of 1.8±0.24

from a starting OD_{600} of 0.05 within 4 h and a maximum doubling time of 34±1 min when

grown under standard laboratory conditions (Figure 1A). This exponential growth phase was

followed by slower growth to a maximum OD_{600} of 3.17±0.11 after 9 h with an OD value

doubling time of 410±110 min. Cell numbers, too, increased exponentially in the first four

hours from $4.5*10^7$ to $7.3*10^8±1.8*10^8$ cells/ml with a doubling time of 30±9 min, followed

by decreased growth to a maximum count of $3.7*10^9±5.8*10^7$ cells/ml after 9 h with a

doubling time of 168±6 min. The notably slower increase of OD_{600} values compared to cell

numbers in the last 6 h of cultivation could be explained by a decrease in cell size at later

growth stages that is indicative for reductive cell division (Figure 1B, Supplementary Figure

S1A).

The chromosome content of cells was monitored by stoichiometric staining with SYBR

Green (Figure 1C, Supplementary Figure S1B). In the over-night grown pre-cultures that

were used for inoculation, 80% of the cells contained one chromosome (C1). One hour after

the transfer into fresh medium already 62±7% and 30±2% of the cells contained two (C2) and

three chromosome equivalents, respectively, and a smaller fraction even more. This clearly

indicates that the cultures had moved to a phase of uncoupled growth with replication being

faster than cell division. After 4 to 5 h of growth, the chromosome content shifted back and

two distinct peaks for cells containing one or two chromosomes became visible again. The

presence of cells with a DNA content between the major peaks indicates actively replicating

cells (R). The proportion of C2 and R cells was only slowly reduced from 38±3% to 23±2%

between 6 h and 10 h of growth. The presence of replicating cells after 10 h of cultivation

was also visible on electron micrographs (Figure 1D). Our data strongly suggest that *P.*

aeruginosa shifts from fast uncoupled to sustained coupled growth cells during the course of

cultivation in LB medium with a short transition phase in between (Figure 1C). If the

doubling time of the coupled growing culture is converted to the approximately 25% of cells

that actually replicate, the individual division times are around 42 min, thus only slightly

lower than in the uncoupled growth phase.

Transcriptome dynamics of *P. aeruginosa* during different growth phases.

We monitored transcriptional changes for the full growth period in one-hour intervals. Two

independent experiments with two and three replicates each were carried out. The

transcriptomes clustered according to the growth phases except for the 6 h samples. For these

samples, the transcriptomes of the first experiment were closer to the transition phase, while

the transcriptomes of the second experiment were closer to the coupled growth phase samples

(Supplementary Figure S2A). The 1736 genes, which showed a significant differential

expression during the course of cultivation, could be assigned to eight clusters

(Supplementary Figure S2B and C and Supplementary Table S1).

The transition between growth-phases was characterized by fast-paced waves of transient

transcriptional activity (Figure 2A). Genes in clusters 1 to 3 showed a comparable high

expression during the first 2 h of uncoupled growth, but with a different timing of maximum

expression and the decline afterwards. During this growth phase, in particular transcription

and translation-related processes were expressed (Figure 2B), including biosynthesis of

tRNAs, RNA polymerase and ribosomes as well as chaperones. A high activity was seen for

oxidative phosphorylation and also for biosynthesis of the vitamins folate (B9) and cobalamin

(B12), in accordance with their respective roles in DNA and methionine synthesis.

Expression of the type III secretion system gene clusters *psc* and *pcr* and the *exoT* effector [17]

peaked at 2 h of cultivation followed by a steep decline.

Cluster 4 to 6 contained genes that were transiently activated at the end of exponential

growth. The high number of sugar and amino acid transporters as well as genes of the

pyruvate metabolism indicated a shift in the metabolic preferences. In particular transporters

for branched-chain amino acids were found to be upregulated in this transition phase, in

accordance with their late utilization as a carbon source observed before[10]. Activation of the

urea cycle and denitrification, and the glycogen metabolism pathway indicate changes of

nitrogen and carbon utilization at this stage. Cluster 7 harbored genes, which were activated

late in the transition phase and exhibited a stable expression throughout the coupled growth

phase. Denitrification genes were among them as well as genes encoding subunits of a sulfate

transporter and the MexHIG antibiotic efflux pump[18]. Finally, expression of the late

responding genes in cluster 8 increased between 5 and 8 h before reaching a stable level. In

particular, activation of the pyoverdine biosynthesis machinery, the heme acquisition protein

HasA and the sulfonate transport and metabolism pathway indicate a response to iron and

sulfur limitation in the medium, respectively.

The three components of the quorum sensing system showed different activation dynamics

consistent with previous data[19,20]. While the primary QS activator *lasR*/*rsaL* pair was not

among the significantly regulated genes, its expression showed a small but consistent

gradually increase from 2 h cultivation on (Supplementary Table S1). The *pqsABCDE*-operon

was transiently activated with a peak between 4 and 5 h followed by a decline and therefore

found in cluster 5. The pyocyanin biosynthesis gene clusters showed the same pattern, but

with a much more pronounced peak. The chemotaxis operon was also activated transiently

during in the transition phase (cluster 6), while flagella genes were not differentially

regulated. The QS regulator RhlR was assigned to cluster 7 with an activation delay but

stable expression throughout coupled growth. The QS target genes coding for alkaline

protease, cyanide production and lectin B were found in the same cluster.

Influence of gene dosage on the transcriptome during uncoupled growth.

Next, we analyzed the distribution of genes in the determined clusters along the *ori-ter*-axis

of the chromosome. Genes active during uncoupled growth (clusters 1 to 3) were

predominantly located close to *ori* while those that were activated during the transition phase

(clusters 4 and 5) were more equally distributed along the chromosome. Genes in cluster 6,

activated at the end of the transition phase, already showed a tendency towards *ter*, a trend

that became even more pronounced for the genes in clusters 7 and 8 that increased expression

during coupled growth (Figure 3A). Furthermore, the average expression levels of genes in

the *ori*-proximal half exceeded those in the *ter*-proximal half of the chromosome during the

first three hours of uncoupled growth. At later time-points a balanced expression of both

halves of the chromosome was observed (Figure 3B). These data are in accordance with the

predicted gene dosage effect in cells with high replication rates.

The gene dosage effect became also visible when a general additive model was fitted to the

$\log_2$ fold-change transcriptome data along the chromosome in order to identify local trends in

expression dynamics that go beyond the regulation of single genes or operons. When

comparing subsequent time-points, with a gradual change in chromosome content, a slightly

lower expression was found around the terminus when transcriptomes from 3 h versus 4 h

were compared (and to a lesser extent for 2 h vs. 3 h, Supplementary Figure S3). This

comparison marks the beginning of the transition from uncoupled to coupled growth and also

showed the strongest shift in chromosome content during cultivation.

The position-specific differences in gene expression became more pronounced when we

compared transcriptomes of time-points with a higher difference in chromosome content

(Figure 3C). A clearly lower transcription of genes in the region surrounding the terminus of

replication was visible when the different growth phases were compared, in particular seen

for uncoupled versus coupled growth. To a lesser extent this trend was also seen for the

comparison of uncoupled growth to transition and transition to coupled growth phase. In

accordance with the analysis above, this specific reduction of gene expression proximal to,

and also increasing towards *ter*, can be parsimoniously explained by a change in mRNA

composition as a result of a higher transcriptional activity of *ori*-proximal genes, thus a gene-

dosage effect (indicated by the orange line in Figure 3C).

Replication-associated transcriptome changes during coupled growth.

The coupled growth with only one replication per cell division in the last 6 hours of

cultivation should allow to discriminate the transcriptomes of non-replicating, replicating and

pre-divisional *P. aeruginosa* cells. To this end, we developed a protocol employing FACS to

separate cells based on their chromosome content (Supplementary Text S1). In order to

determine the influence of fixation with formaldehyde (FA), and FACS on RNA

composition, we compared samples obtained during different steps of the protocol to a

sample fixed with RNAprotect (RP) (Figure 4A). Across the three replicates, the different

samples showed a consistently high correlation (Figure 4B, Supplementary Figure 4A). We

only found 15 genes as well as the chromosomal region of 32 phage-related genes, which

were higher expressed in the RP- than in the FA-treated samples (Supplementary Table S2).

Only two genes found to be regulated during the cell cycle were also influenced by the

fixation method, thus rendering the protocol suitable for the intended purpose.

Next we compared the transcriptomes of the cell populations with one (C1) or two (C2)

chromosomes and those replicating (R). The R and C2 fractions differed from the C1

fraction, but were highly similar to each other (Figure 4C). Only eleven genes were found to

be differentially expressed exclusively when these two fractions were compared. This

included the *gnyDBHAL* gene cluster coding for enzymes of the acyclic isoprenoid

degradation pathway[21], which showed the strongest downregulation in the R versus C2

fraction. The *nrdAB* genes coding for both subunits of the ribonucleotide-diphosphate

reductase were downregulated in the C2 fraction compared to C1 and R. This enzyme

catalyzes the last step in the formation of deoxyribonucleotides. In *E. coli*, it is activity has

been linked to controlling the rate of DNA synthesis[22]. Furthermore, it has been shown that

gene expression peaks at initiation and declines towards the end of replication which is in

accordance with our data for *P. aeruginosa*.

Between the actively replicating R and the C1 fraction, a clear dosage effect was visible with

gene expression decreasing from *ori* to *ter* (Figure 4D). The same was seen for the

comparison of R and C2, but not when the fractions with only completely replicated

chromosomes, C1 and C2 were compared (Supplementary Figure S4B). The differential

expression of several chromosomal loci exceeded this trend dependent on the chromosomal

position. In the R (and C2) fraction, the genes encoding the divisome showed the strongest

activation compared to C1. These comprise of the *mur* and *mra* operons, encoding the

enzymes for remodeling the peptidoglycan layer at the division plane and the *fts* genes,

encoding the components responsible for septum formation[23]. The recombination genes *lexA*

and *recG* were upregulated, too. Of note was also the transcriptional activation of one

genomic island, the region of genomic plasticity RGP41[24], consisting of only uncharacterized

genes. In the C1 fraction, the flagella gene clusters and chemotaxis operons, as well as the *glg*

genes encoding the enzymes of the glycogen metabolic pathway showed the strongest

activation compared to R and C2. Notably, the *mexE* gene, completely inactive in the other

fractions, also showed a more than 64-fold higher expression in the C1 population, by far the

strongest regulation in the dataset (Supplementary Table S2). It encodes the transmembrane

protein part of an efflux-transporter for norfloxacin and imipenem[18].

Discussion

Here we showed that *P. aeruginosa* switches from replication-uncoupled to -coupled growth when cultivated in LB medium, thus allowing to study the effect of replication on the transcriptome. Hereby, the chromosomal gene order reflects the expression maxima during both growth-phases with the genes important for fast uncoupled growth being located closer to *ori* and the stationary phase genes located closer to *ter*. It has been demonstrated before that the *E. coli* sigma 70 factor and its targets, which are mostly active in the exponential phase, are located closer to *ori,* while the sigma S factor and its mostly stationary phase active targets are located closer to *ter*[25]. Thus, while the sigma factors transcriptionally regulate downstream genes, regulon expression is additionally enhanced by a gene dosage effect acting on the regulators and their target genes. Our data show the potential of combining identification of different growth phases by flow cytometry with the comparison of the respective transcriptomes. The gained knowledge could generally be used to identify replication-associated effects on gene expression for the vast number of strains with existing transcriptome data[14,16,26], and integrated into existing gene regulatory models[15,27]. It could further help to better understand chromosomal architecture and to explain gene order evolution[2,25,28,29].

In the coupled growth phase, *P. aeruginosa* displays a distinct transcriptome between the approx. 25% dividing and 75% non-dividing cells. Expression of flagella genes is restricted to cells that are not replicating, while those that replicate differ mainly in the activity of a cell division locus. Furthermore, we found that expression of *mexE*, involved in the expression of an important antibiotic resistance trait is restricted to the non-dividing cells. This induction of subpopulations during the switch in growth phases is coincidental with the activation of the *rhl* QS system. Cell communication induced population heterogeneity has been shown for *P. aeruginosa* before[30] and is also common in other bacteria[31–33]. It might also be the trigger switching the replication mode and restricting activity of the flagella gene clusters to the non-dividing cells. In contrast to chemotaxis, flagella gene expression has not been described to be controlled directly by QS before[19,20]. However, we also did not find them differentially expressed in the culture as a whole, but only in a subpopulation. Thus, a possible connection between communication and development of motility in a fraction of cells might have been overlooked and is worth a closer investigation. Furthermore, slow-growing QS-defective mutants frequently evolve during CF infections[34,35] It would be interesting to determine if

these strains reproduce by coupled growth only and how the transcriptome is affected by this

change.

The highly similar transcriptomes of replicating and pre-divisional cells indicate that in *P.*

aeruginosa no distinct phases of a differentiation program are coupled to progressing

replication. This is in stark contrast to the precisely timed cell cycle of *C. crescentus* with a

defined order of gene activity as cells replicate[36]. Transcriptome dynamics during replication

has so far only be determined for a couple of model bacteria[36–39]. Key to these studies was the

ability to synchronize the cell cycle within the cultures. Our newly developed method based

on cell sorting according to DNA content allows for identification of replication-specific gene

expression without the need for synchronization, as long as the cells grow slowly with

coupled replication and cell division. Not only cell sorting, but also complementary recent

advances in single cell sequencing[40] open up the path to comparative analysis of larger

groups of bacteria, thus contributing to a better understanding of the evolution of cell-cycle

control at the transcriptional level[41].

Material and Methods

Strains and growth conditions.

Pseudomonas aeruginosa PA14[42] was grown in Lysogeny Broth (10 g/L tryptone, 5 g/L yeast

extract, 10 g/L NaCl) at 37˚C and 160 rpm shaking. The growth of cultures inoculated to a

starting OD_{600} of 0.05 was followed for 10h and samples for determination of OD_{600}, cell

count, DNA content, and RNAseq were withdrawn every hour. For FACS-based sorting,

cultures were inoculated to a starting OD_{600} of 0.2 and samples were prepared after 5 h when

the coupled-growth mode was stably reached.

Flow cytometric determination of cell number and chromosome content.

100 μL of culture were mixed with 80 μL of 25% glutaraldehyde in H2O and incubated for 5

min. 820 μL of PBS were added and a dilution series up to 1:1000 was prepared. 10 μL of

SYBR Green (100x) was added to 1 mL of diluted culture. After an incubation time of 20

min, the sample was measured on a BD FACS Canto flow cytometer (BD Biosciences,

Heidelberg, Germany). After gating based on centered forward and sideward scatter, cells

were identified and chromosome content quantified by fluorescence detection in the FITC

channel (excitation 488 and emission 535 nm). Data processing and analysis were performed
with the R package ggcyto[43].

Electron microscopy.

Bacteria were fixed by addition of glutaraldehyde (final concentration 2%) for 30 minutes,

and addition of formaldehyde (final concentration 5%) into the culture medium. EM sample

preparation was performed as previously described[44] with slight modifications. In brief,

samples were washed twice with TE-buffer and fixed to poly-l-lysine coated round cover

slips. After additional washing steps, the samples were dehydrated in a gradient series of

acetone (10%, 30%, 50%, 70%, 90%) on ice and two steps with 100% acetone at room

temperature (each step for 10 minutes). Afterwards, samples were critically point dried with

the CPD300 (Leica Microsystems, Wetzlar, Germany), mounted to aluminum pads and

sputter coated with gold-palladium. Images were acquired with a field emission scanning

electron microscope Merlin (Zeiss, Jena, Germany) equipped with an Everhart Thornley and

an inlens detector and operating at an acceleration voltage of 5kV.

RNAseq library preparation from whole cultures.

Depending on the density, 1 to 2 mL of culture were mixed with the same volume

RNAprotect™ Bacteria Reagent (Qiagen, Hilden, Germany) incubated for 10 min and

centrifuged. The pellets were flash-frozen and stored at -70°C. RNA extraction was carried

out with the RNeasy Plus Kit in combination with QIAshredder™ columns (Qiagen, Hilden,

Germany). Treatment with DNase I was performed in solution. Multiplexed libraries were

generated from directly barcoded fragmented RNA according to a previously published

custom protocol[45], including rRNA removal with the RiboZero Kit (Illumina, San Diego,

USA).

Fluorescence-activated cell sorting for RNAseq of subpopulations.

The method was developed based on a previously published study[46]. A step-by-step protocol

for sample preparation, sorting and RNA isolation is provided in Supplementary Text S1.

Key to successful RNA recovery is the gentle formaldehyde fixation at 4°C. Aliquots of fixed

samples were adjusted to approx. $1.8*10^7$ cells/mL in 30 ml volume each and stained with

SYBR Green. Sorting of $5.4*10^8$ cells based on the FITC-signal (see above) directly into

RNAprotect was performed with the BD FACSAria Fusion (BD Biosciences, Heidelberg,

Germany). The sorted cells were collected on a filter from which RNA was extracted using a

combination of Lysozyme and Proteinase K digestion with bead beating, and purified with

NucleoZOL (Takara Bio, Göteborg, Sweden). Ribosomal RNA depletion was performed with

the NEBNext Bacteria kit (NEB, Frankfurt, Germany). The libraries were prepared with the

TruSeq kit (Illumina, San Diego USA).

Transcriptome analysis

Sequencing of all libraries was performed on a NovaSeq 6000 (Illumina, San Diego, USA) in

paired-end mode with 100 cycles in total. Reads were filtered with fastQC-mcf

(https://github.com/ExpressionAnalysis/ea-utils) and mapped to the *P. aeruginosa* PA14

genome (RefSeq accession GCF_000404265.1) using bowtie2[47]. FeatureCounts was used to

assess the number of reads per gene[48]. Normalization and identification of significantly

differentially regulated genes (FDR < 0.05, absolute $\log_2$ fold change (FC) > 1) was

performed in R using the glmTreat-function of edgeR[49]. Cluster assignment of differentially

expressed genes was performed with the package mfuzz[50].

Data availability

RNAseq raw data have been deposited at the NCBI gene expression omnibus database under

accessions GSE159698 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE159698)

and GSE217100 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE217100).

Acknowledgements

S.H. was funded by the Lower Saxony Ministry for Science and Culture (Bacdata ZN3428),

the European Union (EU, ERC Consolidator Grant COMBAT 724290) and by the Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence

Strategy (EXC 2155 390874280). Furthermore, S.H. received additional funding from the

DFG (DFG SPP 1879) and the Novo Nordisk Foundation (NNF 18OC0033946). J.T.

received funding from the mobility grant of the Czech Ministry of Education

(CZ.02.2.69/0.0/0.0/18_053/0017705). The authors thank Astrid Dröge and Ina Schleicher

for technical support.

References

1. Reyes-Lamothe R, Sherratt DJ. The bacterial cell cycle, chromosome inheritance and cell
growth. *Nat Rev Microbiol*. 2019;17(8):467-478. doi:10.1038/s41579-019-0212-7

2. Couturier E, Rocha EPC. Replication-associated gene dosage effects shape the genomes
of fast-growing bacteria but only for transcription and translation genes. *Molecular*
Microbiology. 2006;59(5):1506-1518. doi:10.1111/j.1365-2958.2006.05046.x

3. Slager J, Veening JW. Hard-Wired Control of Bacterial Processes by Chromosomal Gene
Location. *Trends in Microbiology*. 2016;24(10):788-800. doi:10.1016/j.tim.2016.06.003

4. Soler-Bistué A, Mondotte JA, Bland MJ, Val ME, Saleh MC, Mazel D. Genomic
Location of the Major Ribosomal Protein Gene Locus Determines Vibrio cholerae Global
Growth and Infectivity. *PLOS Genetics*. 2015;11(4):e1005156.
doi:10.1371/journal.pgen.1005156

5. Narula J, Kuchina A, Lee D yeon D, Fujita M, Süel GM, Igoshin OA. Chromosomal
Arrangement of Phosphorelay Genes Couples Sporulation and DNA Replication. *Cell*.
2015;162(2):328-337. doi:10.1016/j.cell.2015.06.012

6. Stokke C, Waldminghaus T, Skarstad K 2011. Replication patterns and organization of
replication forks in Vibrio cholerae. *Microbiology*. 157(3):695-708.
doi:10.1099/mic.0.045112-0

7. Quardokus EM, Brun YV. Cell cycle timing and developmental checkpoints in
Caulobacter crescentus. *Current Opinion in Microbiology*. 2003;6(6):541-549.
doi:10.1016/j.mib.2003.10.013

8. Nathwani D, Raman G, Sulham K, Gavaghan M, Menon V. Clinical and economic
consequences of hospital-acquired resistant and multidrug-resistant Pseudomonas
aeruginosa infections: a systematic review and meta-analysis. *Antimicrobial Resistance*
and Infection Control. 2014;3(1):32. doi:10.1186/2047-2994-3-32

9. Rossi E, La Rosa R, Bartell JA, et al. Pseudomonas aeruginosa adaptation and evolution
in patients with cystic fibrosis. *Nat Rev Microbiol*. 2021;19(5):331-342.
doi:10.1038/s41579-020-00477-5

10. La Rosa R, Johansen HK, Molin S. Convergent Metabolic Specialization through Distinct
Evolutionary Paths in Pseudomonas aeruginosa. *mBio*. 2018;9(2):e00269-18.
doi:10.1128/mBio.00269-18

11. Gibson B, Wilson DJ, Feil E, Eyre-Walker A. The distribution of bacterial doubling
times in the wild. *Proceedings of the Royal Society B: Biological Sciences*.
285(1880):20180789. doi:10.1098/rspb.2018.0789

12. Vallet-Gely I, Boccard F. Chromosomal Organization and Segregation in Pseudomonas
aeruginosa. *PLOS Genetics*. 2013;9(5):e1003492. doi:10.1371/journal.pgen.1003492

13. Bhowmik BK, Clevenger AL, Zhao H, Rybenkov VV. Segregation but Not Replication
of the Pseudomonas aeruginosa Chromosome Terminates at Dif. *mBio*.
2018;9(5):e01088-18. doi:10.1128/mBio.01088-18

14. Dötsch A, Schniederjans M, Khaledi A, et al. The Pseudomonas aeruginosa
Transcriptional Landscape Is Shaped by Environmental Heterogeneity and Genetic
Variation. *mBio*. 2015;6(4):e00749-15. doi:10.1128/mBio.00749-15

15. Tan J, Hammond JH, Hogan DA, Greene CS. ADAGE-Based Integration of Publicly
Available Pseudomonas aeruginosa Gene Expression Data with Denoising Autoencoders
Illuminates Microbe-Host Interactions. *mSystems*. 2016;1(1):e00025-15.
doi:10.1128/mSystems.00025-15

16. Hornischer K, Khaledi A, Pohl S, et al. BACTOME—a reference database to explore the
sequence- and gene expression-variation landscape of Pseudomonas aeruginosa clinical
isolates. *Nucleic Acids Research*. 2019;47(D1):D716-D720. doi:10.1093/nar/gky895

17. Selim H, Radwan TEE, Reyad AM. Regulation of T3SS synthesis, assembly and
secretion in Pseudomonas aeruginosa. *Arch Microbiol*. 2022;204(8):468.
doi:10.1007/s00203-022-03068-5
18. Mesaros N, Glupczynski Y, Avrain L, Caceres NE, Tulkens PM, Van Bambeke F. A
combined phenotypic and genotypic method for the detection of Mex efflux pumps in
Pseudomonas aeruginosa. *Journal of Antimicrobial Chemotherapy*. 2007;59(3):378-386.
doi:10.1093/jac/dkl504
19. Schuster M, Lostroh CP, Ogi T, Greenberg EP. Identification, Timing, and Signal
Specificity of Pseudomonas aeruginosa Quorum-Controlled Genes: a Transcriptome
Analysis. *Journal of Bacteriology*. 2003;185(7):2066-2079. doi:10.1128/JB.185.7.2066-
2079.2003
20. Chadha J, Harjai K, Chhibber S. Revisiting the virulence hallmarks of Pseudomonas
aeruginosa: a chronicle through the perspective of quorum sensing. *Environmental*
Microbiology. 2022;24(6):2630-2656. doi:10.1111/1462-2920.15784
21. Díaz-Pérez AL, Zavala-Hernández AN, Cervantes C, Campos-García J. The
gnyRDBHAL cluster is involved in acyclic isoprenoid degradation in Pseudomonas
aeruginosa. *Appl Environ Microbiol*. 2004;70(9):5102-5110.
doi:10.1128/AEM.70.9.5102-5110.2004
22. Herrick J, Sclavi B. Ribonucleotide reductase and the regulation of DNA replication: an
old story and an ancient heritage. *Molecular Microbiology*. 2007;63(1):22-34.
doi:10.1111/j.1365-2958.2006.05493.x
23. Margolin W. FtsZ and the division of prokaryotic cells and organelles. *Nat Rev Mol Cell*
Biol. 2005;6(11):862-871. doi:10.1038/nrm1745
24. Mathee K, Narasimhan G, Valdes C, et al. Dynamics of Pseudomonas aeruginosa
genome evolution. *Proceedings of the National Academy of Sciences*. 2008;105(8):3100-
3105. doi:10.1073/pnas.0711982105
25. Sobetzko P, Travers A, Muskhelishvili G. Gene order and chromosome dynamics
coordinate spatiotemporal gene expression during the bacterial growth cycle.
Proceedings of the National Academy of Sciences. 2012;109(2):E42-E50.
doi:10.1073/pnas.1108229109
26. Rossi E, Falcone M, Molin S, Johansen HK. High-resolution in situ transcriptomics of
Pseudomonas aeruginosa unveils genotype independent patho-phenotypes in cystic
fibrosis lungs. *Nat Commun*. 2018;9(1):3459. doi:10.1038/s41467-018-05944-5
27. Schulz S, Eckweiler D, Bielecka A, et al. Elucidation of Sigma Factor-Associated
Networks in Pseudomonas aeruginosa Reveals a Modular Architecture with Limited and
Function-Specific Crosstalk. *PLOS Pathogens*. 2015;11(3):e1004744.
doi:10.1371/journal.ppat.1004744
28. Lato DF, Golding GB. Spatial Patterns of Gene Expression in Bacterial Genomes. *J Mol*
Evol. 2020;88(6):510-520. doi:10.1007/s00239-020-09951-3
29. Tomasch J, Koppenhöfer S, Lang AS. Connection Between Chromosomal Location and
Function of CtrA Phosphorelay Genes in Alphaproteobacteria. *Frontiers in*
Microbiology. 2021;12. Accessed October 2, 2022.
https://www.frontiersin.org/articles/10.3389/fmicb.2021.662907
30. Rattray JB, Thomas SA, Wang Y, et al. Bacterial Quorum Sensing Allows Graded and
Bimodal Cellular Responses to Variations in Population Density. *mBio*.
2022;13(3):e00745-22. doi:10.1128/mbio.00745-22
31. Anetzberger C, Pirch T, Jung K. Heterogeneity in quorum sensing-regulated
bioluminescence of Vibrio harveyi. *Molecular Microbiology*. 2009;73(2):267-277.
doi:10.1111/j.1365-2958.2009.06768.x

32. Patzelt D, Wang H, Buchholz I, et al. You are what you talk: quorum sensing induces
individual morphologies and cell division modes in Dinoroseobacter shibae. *ISME J*.
2013;7(12):2274-2286. doi:10.1038/ismej.2013.107
33. Reck M, Tomasch J, Wagner-Döbler I. The Alternative Sigma Factor SigX Controls
Bacteriocin Synthesis and Competence, the Two Quorum Sensing Regulated Traits in
Streptococcus mutans. *PLOS Genetics*. 2015;11(7):e1005353.
doi:10.1371/journal.pgen.1005353
34. Kordes A, Preusse M, Willger SD, et al. Genetically diverse Pseudomonas aeruginosa
populations display similar transcriptomic profiles in a cystic fibrosis explanted lung. *Nat*
Commun. 2019;10(1):3397. doi:10.1038/s41467-019-11414-3
35. Jeske A, Arce-Rodriguez A, Thöming JG, Tomasch J, Häussler S. Evolution of biofilm-
adapted gene expression profiles in lasR-deficient clinical Pseudomonas aeruginosa
isolates. *npj Biofilms Microbiomes*. 2022;8(1):1-14. doi:10.1038/s41522-022-00268-1
36. Laub MT, McAdams HH, Feldblyum T, Fraser CM, Shapiro L. Global Analysis of the
Genetic Network Controlling a Bacterial Cell Cycle. *Science*. 2000;290(5499):2144-
2148. doi:10.1126/science.290.5499.2144
37. Waldbauer JR, Rodrigue S, Coleman ML, Chisholm SW. Transcriptome and Proteome
Dynamics of a Light-Dark Synchronized Bacterial Cell Cycle. *PLOS ONE*.
2012;7(8):e43432. doi:10.1371/journal.pone.0043432
38. De Nisco NJ, Abo RP, Wu CM, Penterman J, Walker GC. Global analysis of cell cycle
gene expression of the legume symbiont Sinorhizobium meliloti. *PNAS*.
2014;111(9):3217-3224. doi:10.1073/pnas.1400421111
39. Bandekar AC, Subedi S, Ioerger TR, Sassetti CM. Cell-Cycle-Associated Expression
Patterns Predict Gene Function in Mycobacteria. *Current Biology*. 2020;30(20):3961-
3971.e6. doi:10.1016/j.cub.2020.07.070
40. Pountain A, Jiang P, Podkowik M, Shopsin B, Torres VJ, Yanai I. A quantitative model
for the transcriptional landscape of the bacterial cell cycle. Published online October 23,
2022:2022.10.22.513359. doi:10.1101/2022.10.22.513359
41. Teeseling MCF van, Thanbichler M. Generating asymmetry in a changing environment:
cell cycle regulation in dimorphic alphaproteobacteria. *Biological Chemistry*.
2020;401(12):1349-1363. doi:10.1515/hsz-2020-0235
42. Liberati NT, Urbach JM, Miyata S, et al. An ordered, nonredundant library of
Pseudomonas aeruginosa strain PA14 transposon insertion mutants. *Proceedings of the*
National Academy of Sciences. 2006;103(8):2833-2838. doi:10.1073/pnas.0511100103
43. Van P, Jiang W, Gottardo R, Finak G. ggCyto: next generation open-source visualization
software for cytometry. *Bioinformatics*. 2018;34(22):3951-3953.
doi:10.1093/bioinformatics/bty441
44. Bense S, Witte J, Preuße M, et al. Pseudomonas aeruginosa post-translational responses
to elevated c-di-GMP levels. *Molecular Microbiology*. 2022;117(5):1213-1226.
doi:10.1111/mmi.14902
45. Avraham R, Haseley N, Fan A, Bloom-Ackermann Z, Livny J, Hung DT. A highly
multiplexed and sensitive RNA-seq protocol for simultaneous analysis of host and
pathogen transcriptomes. *Nat Protoc*. 2016;11(8):1477-1491. doi:10.1038/nprot.2016.090
46. Freiherr von Boeselager R, Pfeifer E, Frunzke J. Cytometry meets next-generation
sequencing – RNA-Seq of sorted subpopulations reveals regional replication and iron-
triggered prophage induction in Corynebacterium glutamicum. *Sci Rep*. 2018;8(1):14856.
doi:10.1038/s41598-018-32997-9
47. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods*.
2012;9(4):357-359. doi:10.1038/nmeth.1923

48. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics*. 2014;30(7):923-930.
doi:10.1093/bioinformatics/btt656
49. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics*. 2010;26(1):139-140.
doi:10.1093/bioinformatics/btp616
50. Futschik ME, Carlisle B. Noise-robust soft clustering of gene expression time-course
data. *J Bioinform Comput Biol*. 2005;03(04):965-988. doi:10.1142/S0219720005001375

Figure Legends

Figure 1. Growth and Replication dynamics of *P. aeruginosa* in LB medium. (A) Optical density and cell numbers followed for 10 h of growth in LB medium. **(B)** Distribution of cell area as determined from EM micrographs. **(C)** Distribution of chromosome content revealed by flow cytometric analysis of SybrGreen fluorescence (One to four chromosome equivalents indicated by color). The lower panel shows representative distributions of fluorescence intensity for up to 7 h. R indicates replicating cells during coupled growth. **(D)** Representative EM micrographs of cells during uncoupled (3 h) and coupled (10 h) growth. Visible division planes are marked by a white arrow.

Figure 2. Transcriptome dynamics during growth in LB medium. (A) Expression dynamics of the eight clusters determined with mfuzz. Shown are the changes of the average expression in the according clusters during the course of a 10h-cultivation. **(B)** Significantly ($p < 0.05$) enriched KEGG-categories in the eight clusters. Size indicates the number of enriched genes in the category, color is according to p-value.

Figure 3. Global chromosomal gene expression changes between different growth phases. (A) Distribution of genes on the chromosomes that show the highest expression during uncoupled (clusters 1-3), transition (clusters 4-6) and coupled (clusters 7-8) growth phases. **(B)** Expression of genes located in the ori and ter proximal during uncouples (1-3 h), transition (4-5 h) and coupled (6-10 h) growth phases. **(C)** log_2 FCs between time points from different growth phases. Red lines show the fitted general additive models; orange lines show the models shifted up with the conserved region at the terminus set to log_2 FC of zero. Representative chromosome content indicative for the different growth phases is shown on the right.

Figure 4. Transcriptomes of replicating and non-replicating cells during coupled growth. (A) Sampling scheme for method evaluation. **(B)** Correlation between transcriptomes of differently treated RNAs. Data for two additional replicates are shown in Supplementary Figure S4A. **(C)** Differential expression between replicating (R) and non-replicating (C1, C2) cells. Number of significantly up- and down-regulated genes between fractions (dark red) are shown in the left and right corner at the bottom of each panel, respectively. **(D)** Chromosome-wide differential gene expression in replicating (R) versus non-replicating (C1) cells. Genes that change significantly in expression are marked in dark red. Operons discussed in the text are marked in yellow. The cell-division gene cluster is shown above the plot. The red line shows a fitted general additive model. Data for the comparisons R vs. C2 and C2 vs. C1 is shown in Supplementary Figure S4B.

Supplementary Figure Legends

Supplementary Figure S1. Flow cytometric determination of relative cell size and chromosome content during growth in LB medium. (A) Changes of the side scatter (SSC) indicates reductive cell division from 3 h to 7 h cultivation time. **(B)** Changes in the distribution of chromosome content for three biological replicates in the course of 10 h cultivation.

Supplementary Figure S2. Transcriptome dynamics during growth in LB medium. (A) Multidimensional scaling (MDS) plot of samples taken during 10 h cultivation. Note the different timing during the shift to coupled growth (6 h sample) for the two independent experiments. **(B)** Determination of ideal number of clusters based on the minimum centroid distance within the clusters. Increasing the number of clusters above 8 does not lead to further reduction of centroid distance. **(C)** Expression profiles of genes in the 8 clusters determined with the mfuzz-package. The number of genes within the cluster is shown below the cluster number. Cluster affiliation alongside expression data is also documented in Supplementary Table S1.

Supplementary Figure S3. Time-resolved chromosomal gene expression changes during growth in LB medium. log_2 fold changes between subsequent time points are shown. Red lines show the fitted general additive models.

Supplementary Figure S4. Transcriptomes of replicating and non-replicating cells during coupled growth. (A) Correlation between transcriptomes of differently treated RNAs (see Figure 4A). **(B)** Chromosome-wide differential gene expression in replicating pre-divisonal (C2) versus non-replicating (C1) and replicating (R) versus non-replicating cells. Genes that change significantly in expression are marked in dark red. The red line shows a fitted general additive model.